Polynomials that preserve nonnegative matrices

Benjamin J. Clark^{b,1}, Pietro Paparella^{a,2,*}

^aDivision of Engineering and Mathematics, University of Washington Bothell, Bothell, WA 98011-8246, USA ^bUniversity of Washington Bothell, Bothell, WA 98011-8246, USA

Abstract

In further pursuit of a solution to the celebrated nonnegative inverse eigenvalue problem, Loewy and London [Linear and Multilinear Algebra 6 (1978/79), no. 1, 83–90] posed the problem of characterizing all polynomials that preserve all nonnegative matrices of a fixed order. If \mathscr{P}_n denotes the set of all polynomials that preserve all *n*-by-*n* nonnegative matrices, then it is clear that polynomials with nonnegative coefficients belong to \mathscr{P}_n . However, it is known that \mathscr{P}_n contains polynomials with negative entries. In this work, novel results for \mathscr{P}_n with respect to the coefficients of the polynomials belonging to \mathscr{P}_n . Along the way, a generalization for the even-part and odd-part are given and shown to be equivalent to another construction that appeared in the literature. Implications for further research are discussed.

Keywords: polynomial, nonnegative matrix, nonnegative inverse eigenvalue problem, circulant matrix 2020 MSC: 15B48, 30C10

1. Introduction

The longstanding nonnegative inverse eigenvalue problem ("NIEP") is the problem of recognizing the spectra of nonnegative matrices. In particular, given a multi-set $\Lambda = \{\lambda_1, \ldots, \lambda_n\}$ of complex numbers, the NIEP asks for necessary and sufficient conditions such that Λ is the spectrum of an *n*-by-*n* entrywisenonnegative matrix A, in which case the multi-set Λ is called *realizable* and the matrix A is called a *realizing matrix* for Λ .

It is well-known that if Λ is realizable with realizing matrix A, then

$$s_k(\Lambda) := \sum_{i=1}^n \lambda_i^k = \operatorname{tr}(A^k) \ge 0, \ \forall k \in \mathbb{N},$$

Preprint submitted to Linear Algebra and its Applications

^{*}Corresponding author

Email addresses: thrin@uw.edu (Benjamin J. Clark), pietrop@uw.edu (Pietro Paparella) *URL:* http://faculty.washington.edu/pietrop/ (Pietro Paparella)

¹Supported by a 2018 University of Washington Bothell Founder's Fellow Award.

 $^{^2 \}mathrm{Supported}$ by a 2021 University of Washington Bothell Scholarship, Research, and Creative Practice Seed Grant.

$$s_k^m(\Lambda) \le n^{m-1} s_{km}(\Lambda), \forall k, m \in \mathbb{N}.$$
(1)

Condition (1) is called the *J-LL condition* and was established independently by Johnson [7] and by Loewy and London [8].

For a polynomial p, let $p(\Lambda) := \{p(\lambda_1), \dots, p(\lambda_n)\}$. If p is a polynomial that preserves the nonnegativity of all n-by-n nonnegative matrices, then

$$s_k(p(\Lambda)) = \sum_{i=1}^n p(\lambda_i)^k = \operatorname{tr}(p(A)^k) \ge 0, \ \forall k \in \mathbb{N},$$
(2)

and

$$s_k^m(p(\Lambda)) \le n^{m-1} s_{km}(p(\Lambda)), \forall k, m \in \mathbb{N}.$$
(3)

As noted by Loewy and London [8], determining whether the inequalities (2) and (3) are sufficient requires a characterization of

$$\mathscr{P}_n := \left\{ p \in \mathbb{C}[x] \mid p(A) \ge 0, \forall A \in \mathsf{M}_n(\mathbb{R}), A \ge 0 \right\}.$$

In particular, and for practical purposes, necessary and sufficient conditions are sought in terms of the coefficients of the polynomials. However, few results are known with respect to the coefficients of the polynomials in \mathscr{P}_n .

The characterization of \mathscr{P}_1 is known as the Pólya–Szegö theorem (see, e.g., Powers and Reznick [9, Proposition 2]), which asserts that $p \in \mathscr{P}_1$ if and only if

$$p(x) = \left(f_1(x)^2 + f_2(x)^2\right) + x\left(g_1(x)^2 + g_2(x)^2\right),$$

where $f_1, f_2, g_1, g_2 \in \mathbb{R}[x]$.

Bharali and Holtz [2] gave partial results for the set

$$\mathscr{F}_n := \{ f \text{ entire } | f(A) \ge 0, \forall A \in \mathsf{M}_n(\mathbb{R}), A \ge 0 \} \supset \mathscr{P}_n$$

and characterized entire functions that preserve certain structured nonnegative matrices, including upper-triangular matrices and circulant matrices.

In this work, novel results on the coefficients of polynomials in \mathscr{P}_n are presented. Along the way, a generalization for the even-part and odd-part is given and shown to be equivalent to a construction given by Balaich and Ondrus [1] for polynomials. We conclude with implications for further inquiry.

2. Notation and Background

The set of *m*-by-*n* matrices with entries from a field \mathbb{F} is denoted by $M_{m \times n}(\mathbb{F})$. If m = n, then $M_{m \times n}(\mathbb{F})$ is abbreviated to $M_n(\mathbb{F})$. The set of all *n*-by-1 column vectors is identified with the set of all ordered *n*-tuples with entries in \mathbb{F} and thus denoted by \mathbb{F}^n . The *n*-by-*n* identity matrix is denoted by $I = I_n$.

and

If $n \in \mathbb{N}$, n > 1, and $\lambda \in \mathbb{C}$, then $J_n(\lambda)$ denotes the Jordan block with eigenvalue λ , i.e.,

$$J_n(\lambda) = \begin{bmatrix} \lambda & 1 & & \\ & \lambda & 1 & & \\ & & \ddots & \ddots & \\ & & & \lambda & 1 \\ & & & & \lambda \end{bmatrix} \in \mathsf{M}_n(\mathbb{C}).$$

If $A \in \mathsf{M}_n(\mathbb{F})$, then a_{ij} denotes the (i, j)-entry of A. If $\mathbb{F} = \mathbb{R}$ and $a_{ij} \ge 0$, $1 \le i, j \le n$, then A is called *nonnegative* and this is denoted by $A \ge 0$.

Unless otherwise stated,

$$p(x) = \sum_{k=0}^{m} a_k x^k \in \mathbb{C}[x],$$

where $a_m \neq 0$. If n is a positive integer less than or equal to m, then the coefficients $a_0, a_1, \ldots, a_{n-1}$ are called the *first* n *terms of* p and the coefficients $a_{m-n+1}, \ldots, a_{m-1}, a_m$ are called the *last* n *terms of* p.

If $A \in M_n(\mathbb{F})$, then A is called a *circulant* or *circulant matrix* if there is a vector $v \in \mathbb{F}^n$, called the *reference vector of* A, such that $a_{ij} = v_{(j-i) \mod n+1}$. In such a case, we write $A = \operatorname{circ}(v_1, \ldots, v_n)$. The *n*-by-*n* circulant matrix with reference vector e_2 is denoted by $C = C_n$. Finally, note that if $q(x) = \sum_{k=0}^{n-1} v_{k+1} x^k$, then $A = q(C) = \sum_{k=0}^{n-1} v_{k+1} C^k$ [4, p. 68].

3. The Even and Odd Part of a Polynomial and a Generalization

In this section, we generalize the notion of the even and odd part of a polynomial and show that it is equivalent to a construction given by Balaich and Ondrus [1].

Recall that if $f : \mathbb{C} \longrightarrow \mathbb{C}$, then

$$f_e(x) := \frac{f(x) + f(-x)}{2}$$

is called the *even-part of* f and

$$f_o(x) := \frac{f(x) - f(-x)}{2}$$

is called the *odd-part* of f.

As is well-known, or otherwise easy to show,

$$p_e(x) = \sum_{k \equiv 0 \mod n} a_k x^k$$

and

$$p_o(x) = \sum_{k \equiv 1 \mod n} a_k x^k$$

The following construction generalizes the aforementioned functions.

Definition 3.1. Let $n \in \mathbb{N}$ and $r \in \{0, 1, \dots, n-1\}$. If

$$\mathcal{I}_{(m,n,r)} := \{ 0 \le k \le m \mid k \bmod n = r \},\$$

then the polynomial

$$p_{(r,n)}(x) := \sum_{k \in \mathcal{I}_{(m,n,r)}} a_k x^k,$$

is called the $r \mod n$ -part of p.

Observation 3.2. If $p \in \mathbb{C}[x]$, then $p(x) = \sum_{r=0}^{n-1} p_{(r,n)}(x)$.

Proof. Follows from the fact that if $r \neq \hat{r}$, then $\mathcal{I}_{(m,n,r)} \cap \mathcal{I}_{(m,n,\hat{r})} = \emptyset$ and the fact that

$$\bigcup_{r=0}^{n-1} \mathcal{I}_{(m,n,r)} = \{0, 1, \dots, m\}.$$

Balaich and Ondrus offered the following as a generalization to the even-part and odd-part of a function.

Definition 3.3 ([1, Definition 3]). If $f : \mathbb{C} \longrightarrow \mathbb{C}$, $r \in \{0, 1, ..., n-1\}$, and $\omega := \exp(2\pi i/n)$, then

$$f_{(r,\omega)}(z) := \frac{1}{n} \sum_{k=0}^{n-1} \omega^{-kr} f(\omega^k z)$$
(4)

is called the $r \mod n$ -part of f.

In the case of polynomials, the following result shows that the constructions yield the same function.

Theorem 3.4. If $p \in \mathbb{C}[x]$, then $p_{(r,n)}(x) = p_{(r,\omega)}(x)$.

Proof. Because

$$\sum_{k=0}^{n-1} \omega^{jk} = \begin{cases} n, & j \equiv 0 \mod n \\ 0, & j \not\equiv 0 \mod n \end{cases},$$

it follows that

$$p_{(r,\omega)}(x) = \frac{1}{n} \sum_{k=0}^{n-1} \omega^{-kr} p(\omega^k x)$$

= $\frac{1}{n} \sum_{k=0}^{n-1} \omega^{-kr} \sum_{j=0}^{m} a_j \omega^{jk} x^j$
= $\frac{1}{n} \sum_{k=0}^{n-1} \sum_{j=0}^{m} a_j x^j \omega^{k(j-r)}$
= $\frac{1}{n} \sum_{j=0}^{m} a_j x^j \sum_{k=0}^{n-1} \omega^{k(j-r)}$

$$\begin{split} &= \frac{1}{n} \left(\sum_{j \in \mathcal{I}_{(m,n,r)}} a_j x^j \sum_{k=0}^{n-1} \omega^{k(j-r)} + \sum_{j \notin \mathcal{I}_{(m,n,r)}} a_j x^j \sum_{k=0}^{n-1} \omega^{k(j-r)} \right) \\ &= \frac{1}{n} \left(\sum_{j \in \mathcal{I}_{(m,n,r)}} a_j x^j(n) + \sum_{j \notin \mathcal{I}_{(m,n,r)}} a_j x^j 0 \right) \\ &= \sum_{j \in \mathcal{I}_{(m,n,r)}} a_j x^j = p_{(r,n)}(x). \end{split}$$

4. Coefficients of polynomials in \mathscr{P}_n

Clark and Paparella recently showed that the coefficients of any polynomial belonging to \mathscr{P}_n must be real [3, Corollary 3.4]. As such, hereinafter it is assumed that the coefficients of p are real.

Bharali and Holtz [2, Theorem 6] established the following result for entire functions using block upper-triangular matrices. We offer a simpler proof using Jordan blocks that also applies to entire functions.

Theorem 4.1. If $p \in \mathscr{P}_n$, then $p, p^{(1)}, p^{(2)}, \ldots, p^{(n-1)} \in \mathscr{P}_1$.

Proof. The result follows by utilizing the well known fact [6, p. 386] that

$$p(J_n(t)) = \begin{cases} 1 & \cdots & k & \cdots & n \\ p(t) & \cdots & \frac{p^{(k-1)}(t)}{(k-1)!} & \cdots & \frac{p^{(n-1)}(t)}{(n-1)!} \\ \vdots & \ddots & \ddots & \vdots \\ p(t) & \frac{p(t)}{(k-1)!} \\ \vdots & & \ddots & \\ n & & & p(t) & \frac{p^{(k-1)}(t)}{(k-1)!} \\ \vdots & & & \ddots & \\ n & & & & p(t) \end{bmatrix}.$$

Corollary 4.2. If $p \in \mathscr{P}_n$ and $m \ge n-1$, then the first n terms of p are nonnegative; otherwise, if m < n-1, then all of its coefficients are nonnegative.

Proof. Follows from the fact that $a_k = p^{(k)}(0)/k! \ge 0$.

Observation 4.3. If $p, q \in \mathscr{P}_n$ and $\alpha \ge 0$, then $\alpha p, p+q$, pq, and $p \circ q \in \mathscr{P}_n$.

Proof. Follows from the fact that (p+q)(A) = p(A) + q(A) [5, Theorem 1.15], (pq)(A) = p(A)q(A) [5, Theorem 1.15], and $(p \circ q)(A) = p(q(A))$ [5, Theorem 1.17].

Since $C^n = I$ and C is invertible, it follows by the division algorithm that $C^k = C^r$, where $r = k \mod n$.

Proposition 4.4. If $p \in \mathbb{C}[x]$ and $t \in \mathbb{C}$, then

$$p(tC) = \sum_{r=0}^{n-1} p_{(r,n)}(t)C^r = \operatorname{circ}\left(p_{(0,n)}(t), p_{(1,n)}(t), \dots, p_{(n-1,n)}(t)\right).$$

Proof. By Observation 3.2 and the fact that (p+q)(A) = p(A) + q(A),

$$p(tC) = \sum_{r=0}^{n-1} p_{(r,n)}(tC) = \sum_{r=0}^{n-1} \sum_{k \in \mathcal{I}_r} a_k t^k C^k$$

=
$$\sum_{r=0}^{n-1} \sum_{k \in \mathcal{I}_r} a_k t^k C^r$$

=
$$\sum_{r=0}^{n-1} p_{(r,n)}(t) C^r$$

=
$$\operatorname{circ} \left(p_{(0,n)}(t), p_{(1,n)}(t), \dots, p_{(n-1,n)}(t) \right).$$

Corollary 4.5. If $p \in \mathscr{P}_n$, then $p_{(r,n)} \in \mathscr{P}_1$, $\forall r \in \{0, 1, \dots, n-1\}$.

Proof. If $p \in \mathscr{P}_n$ and $t \ge 0$, then

$$p(tC) = \operatorname{circ} \left(p_{(0,n)}(t), \dots, p_{(n-1,n)}(t) \right) \ge 0,$$

and the latter holds if and only if $p_{(r,n)} \in \mathscr{P}_1, \forall r \in \{0, 1, \dots, n-1\}$.

Corollary 4.6. If $p \in \mathscr{P}_n$, then

$$\sum_{k \in \mathcal{I}_{(m,n,r)}} a_k \ge 0, \ \forall r \in \{0, 1, \dots, n-1\}.$$

Proof. If $p \in \mathscr{P}_n$, then

$$\sum_{k \in \mathcal{I}_{(m,n,r)}} a_k = p_{(r,n)}(1) \ge 0, \ \forall r \in \{0, 1, \dots, n-1\}.$$

Remark 4.7. Since $a_r = p_{(r,n)}(0) \ge 0$, Corollary 4.5 yields another proof of Corollary 4.2, which can also be established by examining $p(J_n(0))$ [2, Proposition 2].

Lemma 4.8. If $n \in \mathbb{N}$, then $\mathscr{P}_{n+1} \subseteq \mathscr{P}_n$.

Proof. For completeness, we repeat the argument given by Bharali and Holtz [2, Lemma 1] for the containment $\mathscr{F}_{n+1} \subseteq \mathscr{F}_n$: let A be a nonnegative matrix of order n and let $p \in \mathscr{P}_{n+1}$. If $B := \operatorname{diag}(A, 0) \in \mathsf{M}_{n+1}(\mathbb{R})$, then $p(B) = \operatorname{diag}(p(A), 0)$ [5, Theorem 1.13(g)]. Since $p(B) \ge 0$, it follows that $p(A) \ge 0$, i.e., $p \in \mathscr{P}_n$.

Corollary 4.9. If $p \in \mathscr{P}_n$, then $p_{(r,\hat{n})} \in \mathscr{P}_1$ for every $\hat{n} \in \{1, \ldots, n\}$ and for every $r \in \{0, 1, \ldots, \hat{n} - 1\}$.

Proof. Immediate from Corollary 4.5 and Lemma 4.8.

Corollary 4.10. If $p \in \mathscr{P}_n$, then

$$\sum_{k \in \mathcal{I}_{(m,\hat{n},r)}} a_k \ge 0,$$

for every $\hat{n} \in \{1, ..., n\}$ and for every $r \in \{0, 1, ..., \hat{n} - 1\}$.

Proof. Immediate from Corollary 4.6 and Lemma 4.8.

Theorem 4.11. If $p \in \mathscr{P}_n$ and deg p > n, then the last n terms are nonnegative.

Proof. By Proposition 4.4, if $t \in \mathbb{R}$, then

$$p(tC) = \operatorname{circ}(p_{(0,n)}(t), \dots, p_{(n-1,n)}(t)) \ge 0.$$

For contradiction, suppose that $a_{m-n+k} < 0$, where $1 \le k \le n$. The *n* terms

 $m-n+1,\ldots,m-1,m$

form a complete residue system modulo n since they are consecutive; as such, if $r = (m - n + k) \mod n$, then a_{m-n+k} is the leading coefficient of $p_{(r,n)}$. Since $a_{m-n+k} < 0$, we may select t large enough such that $p_{(r,n)}(t) < 0$, a contradiction. Thus, the last n terms of p must be nonnegative.

Theorem 4.12. If $p \in \mathbb{C}[x]$ and deg p < 2n, then $p \in \mathscr{P}_n$ if and only if all of the coefficients are nonnegative.

Proof. Sufficiency is clear and necessity follows from Corollary 4.2 and Theorem 4.11. $\hfill \square$

5. Concluding Remarks

In addition to determining whether conditions presented previously are sufficient (or to disprove otherwise), we offer additional results and several other lines of inquiry.

Theorem 5.1. $\mathscr{P}_2 \subset \mathscr{P}_1$.

Proof. If $p(x) = x^2 - 4x + 4 = (x - 2)^2$, then $p(x) \in \mathscr{P}_1$ since $p(x) \ge 0, \forall x \in \mathbb{R}$. However, $p(x) \notin \mathscr{P}_2$ in view of Theorem 4.12.

Clark and Paparella [3, Theorem 5.6] recently showed that $\mathscr{P}_3 \subset \mathscr{P}_2$. Thus, we offer the following.

Conjecture 5.2. If $n \geq 3$, then $\mathscr{P}_{n+1} \subset \mathscr{P}_n$.

If V is a vector space over \mathbb{R} and U is a nonempty subset of V, then U is called a *convex cone* if $\alpha u + \beta v \in U$ for every $u, v \in U$ and $\alpha, \beta \geq 0$. A point x of a convex cone U is called an *extreme direction (or ray)* of U if, whenever $x = \alpha u + \beta v$, with $\alpha, \beta > 0$ and $u, v \in U$, then x = u = v. By Observation 4.3, \mathscr{P}_n is a convex cone of $\mathbb{C}[x]$ and $\{p_k(x) = x^k \in \mathbb{C}[x] \mid k \geq 0\}$ consists of extreme directions. As such, we pose the following line of inquiry.

Problem 5.3. Identify the extreme directions of \mathscr{P}_n .

Acknowledgements

We thank the anonymous referee for their careful review and suggestions that improved this work and the the University of Washington Bothell for generous funding that supported this work.

References

- M. Balaich and M. Ondrus. A generalization of even and odd functions. Involve, 4(1):91–102, 2011.
- [2] G. Bharali and O. Holtz. Functions preserving nonnegativity of matrices. SIAM J. Matrix Anal. Appl., 30(1):84–101, 2008.
- [3] B. J. Clark and P. Paparella. Polynomials that preserve nonnegative matrices of order two. https://arxiv.org/abs/2109.13365, 2021. Submitted.
- [4] P. J. Davis. Circulant matrices. A Wiley-Interscience Publication. John Wiley & Sons, New York-Chichester-Brisbane, 1979.
- [5] N. J. Higham. Functions of matrices. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2008. Theory and computation.
- [6] R. A. Horn and C. R. Johnson. *Topics in matrix analysis*. Cambridge University Press, Cambridge, 1994. Corrected reprint of the 1991 original.
- [7] C. R. Johnson. Row stochastic matrices similar to doubly stochastic matrices. Linear and Multilinear Algebra, 10(2):113–130, 1981.
- [8] R. Loewy and D. London. A note on an inverse problem for nonnegative matrices. *Linear and Multilinear Algebra*, 6(1):83–90, 1978/79.
- [9] V. Powers and B. Reznick. Polynomials that are positive on an interval. Trans. Amer. Math. Soc., 352(10):4677-4692, 2000.