

Qubit State Discrimination using Post-measurement Information

Donghoon Ha,¹ Jeong San Kim,^{1,*} and Younghun Kwon^{2,†}

¹*Department of Applied Mathematics and Institute of Natural Sciences,
Kyung Hee University, Yongin 17104, Republic of Korea*

²*Department of Applied Physics, Center for Bionano Intelligence Education and Research,
Hanyang University, Ansan 15588, Republic of Korea*

We consider the optimal discrimination of nonorthogonal qubit states with post-measurement information and provide an analytic structure of the optimal measurements. We also show that there is always a null optimal measurement when post-measurement information is given. Further, in discriminating four states using post-measurement information, we analytically provide the optimal probability of correct guessing and show that the uniqueness of optimal measurement is equivalent to the non-existence of non-null optimal measurement with post-measurement information.

I. INTRODUCTION

Whereas orthogonal quantum states can be perfectly discriminated in quantum physics, it is not generally true for nonorthogonal states[1–4]. For these reasons, various measurement strategies have been studied for optimal discrimination of nonorthogonal states, such as *minimum-error discrimination*(ME), unambiguous discrimination, and maximum-confidence discrimination[5–11]. ME is a discrimination scheme to minimize the average error probability without inconclusive results. Although a necessary and sufficient condition for realizing a minimum-error measurement in general cases is well known[12–15], the general solution for ME is not yet known except for ME of two states, symmetric states, and qubit states[5, 16–24]. In some cases, ME can be performed without the help of measurement, simply by guessing the state with the greatest prior probability is prepared[25].

When the post-measurement information about the prepared subensemble is available, some nonorthogonal states can be perfectly discriminated[26]. However, in general, nonorthogonal qubit states cannot be perfectly discriminated even with post-measurement information about the prepared subensemble. Therefore, for the case of qubit state, it is important to investigate *minimizing the average error probability with post-measurement information*(MEPI)[27–29]. Also, it is meaningful since MEPI is known to have a relation with the incompatibility of measurements[29–34].

MEPI problem can be understood in view of ME problem; a MEPI of quantum state ensemble consisting of m subensembles with n_1, \dots, n_m states can be translated into a ME of quantum state ensemble with $\prod_{b=1}^m n_b$ states by modifying the states and prior probabilities in the original ensemble[28]. This approach can be useful for characterizing MEPI of qubit states because useful properties and analytical results for ME of qubit states are already well known[18, 21–24].

In this paper, we analyze MEPI of nonorthogonal qubit states and provide an analytic structure of the optimal measurements based on the analysis of some ME problem. We first show that a null optimal measurement exists for any MEPI of qubit states. We also analytically provide a necessary and sufficient condition that pre-measurement information is strictly more favorable than post-measurement information when all subensembles have two states. Moreover, we characterize the optimal measurements by classifying MEPI into the two cases if it is possible or not without the help of measurement. In particular, for the case where the ensemble consists of two subensembles with two states and pre-measurement information is strictly more favorable than post-measurement information, we analytically provide the optimal probability of correct guessing. In this case, we further show that the uniqueness of optimal measurement is equivalent to the non-existence of non-null MEPI measurement.

This paper is organized as follows: In Sect. II, we review and analyze ME of qubit states. By applying the analysis for ME of qubit states to the ME problem associated with MEPI problem, we provide our results for MEPI of qubit states in Sect. III. In Sect. IV, we conclude our results.

*Electronic address: freddie1@khu.ac.kr

†Electronic address: yyhkwon@hanyang.ac.kr

II. MINIMUM-ERROR DISCRIMINATION OF QUBIT STATES

In two-level quantum systems (qubit), a state is expressed by a density operator on two-dimensional complex Hilbert space \mathcal{H} and a measurement with a finite outcome set $\Omega = \{1, \dots, n\}$ is described by a *positive operator valued measure* (POVM) \mathcal{M} , which is composed of n positive semidefinite operators M_1, \dots, M_n satisfying $\sum_{i \in \Omega} M_i = \mathbb{1}$. Here, $\mathbb{1}$ is the identity operator on \mathcal{H} . We say that $i \in \Omega$ is null(non-null) if $M_i \in \Omega$ is zero(non-zero). We also say that \mathcal{M} is *null* if it has at least one null outcome, otherwise *non-null*.

In this section, we consider ME of *qubit* state ensemble $\mathcal{E} = \{\eta_i, \rho_i\}_{i \in \Omega}$ in which the qubit state ρ_i is prepared with the probability η_i . We specify η_1 as the greatest prior probability to reduce the repetitive representation, that is,

$$\eta_1 \geq \eta_2, \dots, \eta_n. \quad (1)$$

A qubit state ρ_i can be described using the Bloch vector $\boldsymbol{\nu}_i$ in the three-dimensional real space \mathbb{R}^3 ,

$$\rho_i = \frac{1}{2}(\mathbb{1} + \boldsymbol{\nu}_i \cdot \boldsymbol{\sigma}), \quad i \in \Omega.$$

where $\boldsymbol{\sigma}$ is the Pauli matrices $(\sigma_X, \sigma_Y, \sigma_Z)$.

Given a qubit state ensemble $\mathcal{E} = \{\eta_i, \rho_i\}_{i \in \Omega}$ and distinct $i, j \in \Omega$, points in \mathbb{R}^3 , which has $|\eta_i - \eta_j|$ as distance difference from $\eta_i \boldsymbol{\nu}_i$ and $\eta_j \boldsymbol{\nu}_j$, form a hyperboloid of two sheets. The one hyperboloid sheet consists of points $\mathbf{v} \in \mathbb{R}^3$ satisfying $\varphi_i(\mathbf{v}) = \varphi_j(\mathbf{v})$, where

$$\varphi_i(\mathbf{v}) = \eta_i + \|\eta_i \boldsymbol{\nu}_i - \mathbf{v}\|, \quad i \in \Omega. \quad (2)$$

Here $\|\cdot\|$ is the Euclidean norm. The sheet divides \mathbb{R}^3 into two sets $\{\mathbf{v} \in \mathbb{R}^3 : \varphi_i(\mathbf{v}) \geq \varphi_j(\mathbf{v})\}$ and $\{\mathbf{v} \in \mathbb{R}^3 : \varphi_i(\mathbf{v}) < \varphi_j(\mathbf{v})\}$. We use the following definitions to express various conditions in ME of \mathcal{E} .

Definition 1. For each $S \subseteq \Omega$,

$$\mathcal{Z}_S := \{\mathbf{v} \in \mathcal{P}_S : \varphi_i(\mathbf{v}) = \varphi_j(\mathbf{v}) \quad \forall (i, j) \in S \times S, \\ \varphi_i(\mathbf{v}) \geq \varphi_j(\mathbf{v}) \quad \forall (i, j) \in S \times (\Omega - S)\}, \quad (3)$$

where \mathcal{P}_S is the relative interior of the convex hull of $\{\eta_i \boldsymbol{\nu}_i\}_{i \in S}$, that is,

$$\mathcal{P}_S = \{\sum_{i \in S} c_i \eta_i \boldsymbol{\nu}_i : c_i > 0 \quad \forall i \in S, \quad \sum_{i \in S} c_i = 1\}. \quad (4)$$

Also, we denote by \mathcal{Z} the union of all \mathcal{Z}_S , that is, $\mathcal{Z} = \bigcup_{S \subseteq \Omega} \mathcal{Z}_S$.

In ME of \mathcal{E} , we use a POVM $\mathcal{M} = \{M_i\}_{i \in \Omega}$ as a measurement such that the prepared state is guessed to be ρ_i for each $i \in \Omega$. Then, the maximal average probability of correctly guessing the given qubit state is

$$p_{\text{guess}} = \max_{\mathcal{M}} \sum_{i \in \Omega} \eta_i \text{Tr}[\rho_i M_i]. \quad (5)$$

ME is the task of finding optimal measurements that provides p_{guess} , called the *guessing probability*. We also note that the guessing probability in ME cannot be less than the greatest prior probability, that is,

$$p_{\text{guess}} \geq \eta_1. \quad (6)$$

In some cases, ME can be performed without the help of measurement[25]; the guessing probability can be obtained by taking the state with the greatest prior probability as the prepared state, that is, $p_{\text{guess}} = \eta_1$. Even for that case, nontrivial optimal measurements can possibly exist. The following proposition provides a necessary and sufficient condition for \mathcal{M} to be optimal when $p_{\text{guess}} = \eta_1$. The proof of Proposition 1 is given in Appendix A.

Proposition 1. For ME of qubit state ensemble \mathcal{E} ,

(a) $p_{\text{guess}} = \eta_1$ if and only if $\mathcal{Z}_{\{1\}}$ is not an empty set \emptyset or, equivalently,

$$\epsilon_i \geq \lambda_i \quad \forall i \in \Omega, \quad (7)$$

where

$$\epsilon_i = \eta_1 - \eta_i, \quad \lambda_i = \|\eta_1 \boldsymbol{\nu}_1 - \eta_i \boldsymbol{\nu}_i\|, \quad i \in \Omega. \quad (8)$$

(b) When $p_{\text{guess}} = \eta_1$, a POVM \mathcal{M} is optimal if and only if

$$\forall i \neq 1, M_i \propto \begin{cases} \mathbb{1} + \frac{\eta_i \boldsymbol{\nu}_i - \eta_1 \boldsymbol{\nu}_1}{\|\eta_i \boldsymbol{\nu}_i - \eta_1 \boldsymbol{\nu}_1\|} \cdot \boldsymbol{\sigma} & , \epsilon_i = \lambda_i, \\ 0 & , \epsilon_i > \lambda_i. \end{cases} \quad (9)$$

When $p_{\text{guess}} = \eta_1$, if $\epsilon_i > \lambda_i$ for all $i \neq 1$, the optimal measurement has only $1 \in \Omega$ as a non-null outcome, and M_1 becomes the identity operator. That is, the optimal measurement in this case is trivial and unique. However, if $\epsilon_i = \lambda_i$ for some $i \neq 1$, the optimal measurement can have a non-null outcome other than 1, which implies a non-trivial optimal measurement. Therefore, when $p_{\text{guess}} = \eta_1$, a nontrivial optimal measurement exists if and only if $\epsilon_i = \lambda_i$ for some $i \neq 1$.

Now, let us consider the qubit state ensemble \mathcal{E} in which a measurement is a necessary requirement for ME, that is, $p_{\text{guess}} > \eta_1$. The following proposition shows that finding $\mathbf{S} \subseteq \Omega$ with $\mathcal{Z}_{\mathbf{S}} \neq \emptyset$ and an element of \mathcal{Z} is directly related to obtaining optimal measurements, where \emptyset is the empty set. The proof of Proposition 2 is given in Appendix A.

Proposition 2. For ME of qubit state ensemble \mathcal{E} ,

(a) \mathcal{Z} is always a single-element set $\{\mathbf{v}\}$ and $p_{\text{guess}} = \varphi_i(\mathbf{v})$ for all i in \mathbf{S} with $\mathcal{Z}_{\mathbf{S}} \neq \emptyset$.

When $p_{\text{guess}} > \eta_1$,

(b) there is an optimal measurement having $\mathbf{S} \subseteq \Omega$ as the set of all non-null outcomes if and only if $\mathcal{Z}_{\mathbf{S}} \neq \emptyset$.

(c) Moreover, a POVM \mathcal{M} having \mathbf{S} as the set of all non-null outcomes is optimal if and only if

$$M_i \propto \mathbb{1} + \frac{\eta_i \boldsymbol{\nu}_i - \mathbf{v}}{\|\eta_i \boldsymbol{\nu}_i - \mathbf{v}\|} \cdot \boldsymbol{\sigma} \quad \forall i \in \mathbf{S}. \quad (10)$$

When $p_{\text{guess}} > \eta_1$, all optimal POVMs are characterized by the *single* element of \mathcal{Z} . Proposition 2 tells the following three facts. First, if $\mathcal{Z}_{\mathbf{S}}$ is empty, there is no optimal measurement having \mathbf{S} as the set of all non-null outcomes. This implies that all optimal measurements are null (that is, $M_i = 0$ for some $i \in \Omega$) if \mathcal{Z}_{Ω} is empty. Second, if $\mathcal{Z}_{\mathbf{S}}$ is non-empty and $\mathcal{Z} \subseteq \mathcal{P}_{\mathbf{S}'}$ for some $\mathbf{S}' \subseteq \mathbf{S}$, then $\mathcal{Z}_{\mathbf{S}'}$ is also non-empty. This implies that, if the *affine* dimension [35, 36] of $\{\eta_i \boldsymbol{\nu}_i\}_{i \in \Omega}$ is D , the guessing probability can be obtained by the detection of $D + 1$ qubit states [22]. Third, if $\mathcal{Z}_{\mathbf{S}}$ is non-empty and $\{\eta_i \boldsymbol{\nu}_i\}_{i \in \mathbf{S}}$ forms a simplex with affine dimension $|\mathbf{S}| - 1$, the optimal measurement having \mathbf{S} as the set of all non-null outcomes is unique.

Condition (10) is a necessary and sufficient condition for a POVM \mathcal{M} having \mathbf{S} as the set of all non-null outcomes to be optimal when $p_{\text{guess}} > \eta_1$. Therefore, when $p_{\text{guess}} > \eta_1$, all optimal measurements are obtained in the following three steps. The first step is to distinguish whether $\mathcal{Z}_{\mathbf{S}}$ is empty or nonempty for each \mathbf{S} with $|\mathbf{S}| \geq 2$. Note that, for $p_{\text{guess}} > \eta_1$, there exists no subset $\mathbf{S} \subseteq \Omega$ with $|\mathbf{S}| = 1$ and $\mathcal{Z}_{\mathbf{S}} \neq \emptyset$ because all optimal measurements have more than one non-null outcome. The second step is to find out what the single element of \mathcal{Z} is. The final step is to get a POVM \mathcal{M} that satisfies Condition (10).

III. MAIN RESULT: MEPI OF QUBIT STATES

In this section, we consider MEPI of *qubit* state ensemble,

$$\mathcal{E} = \bigcup_{b \in \mathbf{B}} \{\eta_{ib}, \rho_{ib}\}_{i \in \mathbf{A}_b}, \quad (11)$$

where

$$\begin{aligned} \mathbf{B} &= \{1, 2, \dots, m\}, \quad m \geq 2, \\ \mathbf{A}_b &= \{1, 2, \dots, n_b\}, \quad n_b \geq 2. \end{aligned} \quad (12)$$

The ensemble \mathcal{E} consists of m subensembles (we use the term “subensemble” regardless of the normalization of prior probability),

$$\mathcal{E}_b = \{\eta_{ib}, \rho_{ib}\}_{i \in \mathbf{A}_b}, \quad b \in \mathbf{B}. \quad (13)$$

Without loss of generality, we assume that

$$\eta_{1b} \geq \eta_{2b}, \dots, \eta_{n_b b} \quad \forall b \in \mathbf{B}. \quad (14)$$

The classical information $b \in \mathcal{B}$ of the prepared subensemble is provided after a measurement is performed. We use a POVM $\mathcal{M} = \{M_\omega\}_{\omega \in \Omega}$ to describe a measurement, where Ω is the Cartesian product of $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_m$, that is,

$$\Omega = \mathcal{A}_1 \times \mathcal{A}_2 \times \dots \times \mathcal{A}_m. \quad (15)$$

Each outcome $\omega = (\omega_1, \dots, \omega_m) \in \Omega$ means that the prepared state is guessed to be $\rho_{\omega_1 1}, \rho_{\omega_2 2}, \dots$, or $\rho_{\omega_m m}$ according to post-measurement information $b = 1, 2, \dots$, or m , respectively.

The maximal average probability of correctly guessing the prepared qubit state is

$$p_{\text{guess}}^{\text{post}} = \max_{\mathcal{M}} \sum_{b \in \mathcal{B}} \sum_{i \in \mathcal{A}_b} \sum_{\substack{\omega \in \Omega \\ \omega_b = i}} \eta_{ib} \text{Tr}[\rho_{ib} M_\omega] = \max_{\mathcal{M}} \sum_{\omega \in \Omega} \tilde{\eta}_\omega \text{Tr}[\tilde{\rho}_\omega M_\omega], \quad (16)$$

where $\tilde{\eta}_\omega$ and $\tilde{\rho}_\omega$ are positive numbers and density operators, respectively, such that

$$\tilde{\eta}_\omega = \sum_{b \in \mathcal{B}} \eta_{\omega_b b}, \quad \tilde{\rho}_\omega = \frac{\sum_{b \in \mathcal{B}} \eta_{\omega_b b} \rho_{\omega_b b}}{\sum_{b' \in \mathcal{B}} \eta_{\omega_b' b'}}, \quad \omega \in \Omega. \quad (17)$$

MEPI of \mathcal{E} is the task of finding optimal measurements that provides $p_{\text{guess}}^{\text{post}}$. From Eq. (16), we can see that $p_{\text{guess}}^{\text{post}}$ is the guessing probability of qubit state ensemble,

$$\tilde{\mathcal{E}} = \{\tilde{\eta}_\omega, \tilde{\rho}_\omega\}_{\omega \in \Omega}. \quad (18)$$

where $\{\tilde{\eta}_\omega\}_{\omega \in \Omega}$ is not normalized, that is, $\sum_{\omega \in \Omega} \tilde{\eta}_\omega > 1$. Therefore, MEPI of \mathcal{E} is equivalent to ME of $\tilde{\mathcal{E}}$; a POVM \mathcal{M} is optimal for MEPI of \mathcal{E} if and only if it is optimal for ME of $\tilde{\mathcal{E}}$ [28]. To distinguish between optimal measurements for ME and MEPI, we use ME and MEPI measurements, respectively. We also note that the assumption in (14) implies that $\tilde{\eta}_1$ is the greatest prior probability of $\tilde{\mathcal{E}}$ and a lower bound of $p_{\text{guess}}^{\text{post}}$, that is,

$$p_{\text{guess}}^{\text{post}} \geq \tilde{\eta}_1 \geq \tilde{\eta}_\omega \quad \forall \omega \in \Omega, \quad (19)$$

where

$$\mathbf{1} = (1, 1, \dots, 1). \quad (20)$$

A. Null MEPI measurement

Similar to ME, we use Bloch representation of qubit states as

$$\begin{aligned} \rho_{ib} &= \frac{1}{2}(\mathbb{1} + \boldsymbol{\nu}_{ib} \cdot \boldsymbol{\sigma}), \quad b \in \mathcal{B}, i \in \mathcal{A}_b, \\ \tilde{\rho}_\omega &= \frac{1}{2}(\mathbb{1} + \tilde{\boldsymbol{\nu}}_\omega \cdot \boldsymbol{\sigma}), \quad \omega \in \Omega. \end{aligned} \quad (21)$$

Then,

$$\sum_{b \in \mathcal{B}} \eta_{\omega_b b} \boldsymbol{\nu}_{\omega_b b} = \tilde{\eta}_\omega \tilde{\boldsymbol{\nu}}_\omega =: \tilde{\boldsymbol{\mu}}_\omega \quad \forall \omega \in \Omega. \quad (22)$$

When $m = n_1 = n_2 = 2$, the affine dimension D of $\{\tilde{\boldsymbol{\mu}}_\omega\}_{\omega \in \Omega}$ is less than three because

$$\begin{aligned} \tilde{\boldsymbol{\mu}}_{(2,1)} - \tilde{\boldsymbol{\mu}}_{(1,1)} &= \tilde{\boldsymbol{\mu}}_{(2,2)} - \tilde{\boldsymbol{\mu}}_{(1,2)} = \eta_{21} \boldsymbol{\nu}_{21} - \eta_{11} \boldsymbol{\nu}_{11}, \\ \tilde{\boldsymbol{\mu}}_{(1,2)} - \tilde{\boldsymbol{\mu}}_{(1,1)} &= \tilde{\boldsymbol{\mu}}_{(2,2)} - \tilde{\boldsymbol{\mu}}_{(2,1)} = \eta_{22} \boldsymbol{\nu}_{22} - \eta_{12} \boldsymbol{\nu}_{12}. \end{aligned} \quad (23)$$

Thus, in this case, ME of $\tilde{\mathcal{E}}$ is possible without detecting every state [22], and there exists a MEPI measurement of \mathcal{E} that is null (that is, $M_\omega = 0$ for some $\omega \in \Omega$).

Other than $m = n_1 = n_2 = 2$, the number of all outcomes, that is, $|\Omega| = \prod_{b \in \mathcal{B}} n_b$, is greater than four, and a null MEPI measurement of \mathcal{E} exists because, for any ME of more than four qubit states, there is a ME measurement that is null [18, 22, 37].

Corollary 1. *For any qubit state ensemble \mathcal{E} , a null MEPI measurement of \mathcal{E} exists.*

B. Upper bound of $p_{\text{guess}}^{\text{post}}$

When the classical information $b \in \mathbf{B}$ of the prepared subensemble is known prior to perform a measurement, the maximal average probability of correctly guessing the prepared qubit state, $p_{\text{guess}}^{\text{prior}}$, can be obtained by performing ME measurement of \mathcal{E}_b according to the pre-measurement information b . In other words,

$$p_{\text{guess}}^{\text{prior}} = \sum_{b \in \mathbf{B}} p_b^{\text{ME}}, \quad (24)$$

where p_b^{ME} is the guessing probability of \mathcal{E}_b , that is,

$$p_b^{\text{ME}} = \max_{\mathcal{M}_b} \sum_{i \in \mathbf{A}_b} \eta_{ib} \text{Tr}[\rho_{ib} M_{ib}], \quad b \in \mathbf{B}. \quad (25)$$

Here, \mathcal{M}_b is a POVM with n_b elements M_{ib} indicating the detection of ρ_{ib} .

Guessing the prepared qubit state using a POVM \mathcal{M} and post-measurement information is equivalent to guessing the prepared state by performing a POVM \mathcal{M}_b consisting of

$$M_{ib} = \sum_{\substack{\omega \in \Omega \\ \omega_b = i}} M_{\omega}, \quad i \in \mathbf{A}_b, \quad (26)$$

according to pre-measurement information $b \in \mathbf{B}$. Given an arbitrary POVM \mathcal{M}_b with n_b elements $M_{1b}, \dots, M_{n_b b}$ for each $b \in \mathbf{B}$, m POVMs $\mathcal{M}_1, \dots, \mathcal{M}_m$ are called *compatible* if there is a POVM \mathcal{M} satisfying Eq. (26) for all $b \in \mathbf{B}$; otherwise, they are called *incompatible*[30]. Thus, $p_{\text{guess}}^{\text{post}}$ is upper bounded as

$$p_{\text{guess}}^{\text{post}} \leq p_{\text{guess}}^{\text{prior}}, \quad (27)$$

where the equality holds if and only if there are m ME measurements of $\mathcal{E}_1, \dots, \mathcal{E}_m$ that are compatible[29].

Obviously, a POVM with the identity operator is compatible with any POVM; therefore, $p_{\text{guess}}^{\text{post}} = p_{\text{guess}}^{\text{prior}}$ if $p_b^{\text{ME}} = \eta_{1b}$ for some $b \in \mathbf{B}$. Moreover, POVMs with the same elements are compatible; thus, $p_{\text{guess}}^{\text{post}} = p_{\text{guess}}^{\text{prior}}$ if there are m optimal POVMs for MEs of $\mathcal{E}_1, \dots, \mathcal{E}_m$ that have the same elements. The following lemma provides a sufficient condition for $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$.

Lemma 1. Suppose that, for some $b, b' \in \mathbf{B}$ with $b \neq b'$, the ME measurements for \mathcal{E}_b and $\mathcal{E}_{b'}$ are unique and consist of rank-one elements $\{M_{ib}\}_{i \in \mathbf{A}_b}$, $\{M_{jb'}\}_{j \in \mathbf{A}_{b'}}$ such that $M_{ib} \not\propto M_{jb'}$ for all $i \in \mathbf{A}_b$ and all $j \in \mathbf{A}_{b'}$. Then, $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$.

Proof. Assume that $p_{\text{guess}}^{\text{post}} = p_{\text{guess}}^{\text{prior}}$ in which Eq. (26) holds for $b, b' \in \mathbf{B}$. Then, $\text{rank}(M_{ib}) = \text{rank}(M_{jb'}) = 1$ implies that $M_{\omega} \propto M_{ib}$ for all $\omega \in \Omega$ with $\omega_b = i$ and $M_{\omega} \propto M_{jb'}$ for all $\omega \in \Omega$ with $\omega_{b'} = j$. Thus, $M_{ib} \not\propto M_{jb'} \forall i, j$ means $M_{ib} = M_{jb'} = 0 \forall i, j$ which contradicts the POVM completeness. Therefore, $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. \square

In the case of $n_b = 2$ and $p_b^{\text{ME}} > \eta_{1b}$ for all $b \in \mathbf{B}$, from Helstrom bound[5] or Proposition 2, the optimal POVM elements for ME of \mathcal{E}_b are uniquely determined as follows:

$$M_{1b} = \frac{1}{2}(\mathbb{1} + \hat{\boldsymbol{\mu}}_{2b} \cdot \boldsymbol{\sigma}), \quad M_{2b} = \frac{1}{2}(\mathbb{1} - \hat{\boldsymbol{\mu}}_{2b} \cdot \boldsymbol{\sigma}), \quad b \in \mathbf{B}, \quad (28)$$

where

$$\hat{\boldsymbol{\mu}}_{ib} = \frac{\eta_{ib} \boldsymbol{\nu}_{ib} - \eta_{1b} \boldsymbol{\nu}_{1b}}{\|\eta_{ib} \boldsymbol{\nu}_{ib} - \eta_{1b} \boldsymbol{\nu}_{1b}\|}, \quad b \in \mathbf{B}. \quad (29)$$

Thus, if $\hat{\boldsymbol{\mu}}_{2b} \times \hat{\boldsymbol{\mu}}_{2b'} = \mathbf{0}$ for all $b, b' \in \mathbf{B}$, then $p_{\text{guess}}^{\text{post}} = p_{\text{guess}}^{\text{prior}}$ because m optimal POVMs for MEs of $\mathcal{E}_1, \dots, \mathcal{E}_m$ have the same elements; however, if $\hat{\boldsymbol{\mu}}_{2b} \times \hat{\boldsymbol{\mu}}_{2b'} \neq \mathbf{0}$ for some $b, b' \in \mathbf{B}$ with $b \neq b'$, then $p_{\text{guess}}^{\text{post}} > p_{\text{guess}}^{\text{prior}}$ from Lemma 1.

Corollary 2. When $n_b = 2$ for all $b \in \mathbf{B}$, $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$ if and only if $\epsilon_{2b} < \lambda_{2b}$ for all $b \in \mathbf{B}$ and $\hat{\boldsymbol{\mu}}_{2b} \times \hat{\boldsymbol{\mu}}_{2b'} \neq \mathbf{0}$ for some $b, b' \in \mathbf{B}$ with $b \neq b'$.

C. MEPI for $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$

From Inequalities (19) and (27), $p_{\text{guess}}^{\text{post}}$ has the following upper and lower bounds,

$$\tilde{\eta}_1 \leq p_{\text{guess}}^{\text{post}} \leq p_{\text{guess}}^{\text{prior}}. \quad (30)$$

Thus, if $p_b^{\text{ME}} = \eta_{1b}$ for all $b \in \mathcal{B}$, then $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$ in which MEPI of \mathcal{E} is possible without the help of measurement; the prepared state is guessed to be $\rho_{11}, \rho_{12}, \dots$, or ρ_{1m} according to post-measurement information $b = 1, 2, \dots$, or m , respectively.

From Proposition 1, we can see that $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$ if and only if

$$\tilde{\epsilon}_\omega \geq \tilde{\lambda}_\omega \quad \forall \omega \in \Omega, \quad (31)$$

where

$$\tilde{\epsilon}_\omega = \tilde{\eta}_1 - \tilde{\eta}_\omega, \quad \tilde{\lambda}_\omega = \|\tilde{\mu}_1 - \tilde{\mu}_\omega\|, \quad \omega \in \Omega. \quad (32)$$

Also, we can see that $p_b^{\text{ME}} = \eta_{1b}$ if and only if

$$\epsilon_{ib} \geq \lambda_{ib} \quad \forall i \in \mathcal{A}_b, \quad (33)$$

where

$$\epsilon_{ib} = \eta_{1b} - \eta_{ib}, \quad \lambda_{ib} = \|\eta_{1b}\nu_{1b} - \eta_{ib}\nu_{ib}\|, \quad b \in \mathcal{B}, i \in \mathcal{A}_b. \quad (34)$$

The following lemma shows that $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$ is equivalent to $p_b^{\text{ME}} = \eta_{1b}$ for all $b \in \mathcal{B}$.

Lemma 2. *For MEPI of qubit state ensemble \mathcal{E} , $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$ if and only if $\epsilon_{ib} \geq \lambda_{ib}$ holds for all b, i . When $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$, a POVM \mathcal{M} is optimal if and only if*

$$\forall \omega \neq \mathbf{1}, \quad M_\omega \propto \begin{cases} \mathbb{1} + \frac{\tilde{\mu}_\omega - \tilde{\mu}_1}{\|\tilde{\mu}_\omega - \tilde{\mu}_1\|} \cdot \boldsymbol{\sigma} & , \quad \tilde{\epsilon}_\omega = \tilde{\lambda}_\omega, \\ 0 & , \quad \tilde{\epsilon}_\omega > \tilde{\lambda}_\omega. \end{cases} \quad (35)$$

Proof. From the definitions of $\tilde{\eta}_\omega$ and $\tilde{\mu}_\omega$ in (17) and (22), we have

$$\tilde{\epsilon}_\omega = \sum_{b \in \mathcal{B}} \epsilon_{\omega_b b}, \quad \tilde{\lambda}_\omega = \sum_{b \in \mathcal{B}} \lambda_{\omega_b b} \quad \forall \omega \in \Omega. \quad (36)$$

Thus, if Inequality (31) holds for all $\omega \in \Omega$ or, equivalently, $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$, then Inequality (33) holds for all b, i because $\tilde{\epsilon}_\omega = \epsilon_{ib}$ and $\tilde{\lambda}_\omega = \lambda_{ib}$ for all $\omega \in \Omega$ such that $\omega_b = i$ for some $b \in \mathcal{B}$ and $\omega_{b'} = 1$ for all $b' \neq b$. Since the converse has already been proved, $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$ if and only if $\epsilon_{ib} \geq \lambda_{ib}$ for all b, i . In addition, directly from Proposition 1, we can see that Condition (35) is a necessary and sufficient condition for a POVM \mathcal{M} to be optimal when $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$. \square

For $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$ and $\omega \in \Omega$, if $\tilde{\epsilon}_\omega = \tilde{\lambda}_\omega$, then

$$\epsilon_{\omega_b b} = \lambda_{\omega_b b} \quad \forall b, \quad \tilde{\lambda}_\omega = \sum_{b \in \mathcal{B}} \lambda_{\omega_b b} \quad (37)$$

because

$$\sum_{b \in \mathcal{B}} \lambda_{\omega_b b} \leq \sum_{b \in \mathcal{B}} \epsilon_{\omega_b b} = \tilde{\epsilon}_\omega = \tilde{\lambda}_\omega \leq \sum_{b \in \mathcal{B}} \lambda_{\omega_b b}, \quad (38)$$

where the first and last inequalities follow from the inequalities in (33) and (36), respectively. Conversely, if Eq. (37) holds, $\tilde{\epsilon}_\omega = \tilde{\lambda}_\omega$ because

$$\tilde{\epsilon}_\omega = \sum_{b \in \mathcal{B}} \epsilon_{\omega_b b} = \sum_{b \in \mathcal{B}} \lambda_{\omega_b b} = \tilde{\lambda}_\omega. \quad (39)$$

Therefore, $\tilde{\epsilon}_\omega = \tilde{\lambda}_\omega$ is equivalent to Eq. (37) when $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$. The last condition of Eq. (37) is equivalent to that all unit vectors $\hat{\mu}_{\omega_b b}$ with $\omega_b \neq 1$ are the same, that is, for all $b \in \mathcal{B}$ with $\omega_b \neq 1$,

$$\hat{\mu}_{\omega_b b} = \frac{\tilde{\mu}_\omega - \tilde{\mu}_1}{\|\tilde{\mu}_\omega - \tilde{\mu}_1\|}, \quad (40)$$

where $\hat{\mu}_{ib}$ is defined in (29).




FIG. 1: In the case of $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, four vectors $\{\tilde{\mu}_\omega\}_{\omega \in \Omega}$ in \mathbb{R}^3 forms a parallelogram with nonempty interior \mathcal{P}_Ω (yellow in (a)). For $\mathcal{Z} \not\subseteq \mathcal{P}_\Omega$, the single element of \mathcal{Z} is in one of two edges $\mathcal{P}_{\{(1,1),(1,2)\}}$ and $\mathcal{P}_{\{(1,1),(2,1)\}}$ (red in (b)), but it cannot be in one of two edges $\mathcal{P}_{\{(1,2),(2,2)\}}$ and $\mathcal{P}_{\{(2,1),(2,2)\}}$ (blue in (b)).

Corollary 3. For MEPI of \mathcal{E} with $n_b = 2$ for all $b \in \mathcal{B}$ and $p_{\text{guess}}^{\text{post}} = \tilde{\eta}_1$, a POVM \mathcal{M} is optimal if and only if

$$\forall \omega \neq \mathbf{1}, M_\omega \propto \begin{cases} \mathbb{1} + \hat{\mu} \cdot \sigma, & \epsilon_{2b} = \lambda_{2b} \forall \omega_b \neq 1 \text{ and} \\ & \exists \hat{\mu} \in \mathbb{R}^3 \text{ such that } \hat{\mu} = \hat{\mu}_{2b} \forall \omega_b \neq 1, \\ 0 & , \text{ otherwise.} \end{cases} \quad (41)$$

D. MEPI for $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$

In order to consider the case of $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$, we redefine \mathcal{Z}_S of Definition 1 suitable for ME of $\tilde{\mathcal{E}}$.

Definition 2. For each $S \subseteq \Omega$,

$$\mathcal{Z}_S := \{v \in \mathcal{P}_S : \varphi_\omega(v) = \varphi_{\omega'}(v) \forall (\omega, \omega') \in S \times S, \\ \varphi_\omega(v) \geq \varphi_{\omega'}(v) \forall (\omega, \omega') \in S \times (\Omega - S)\}, \quad (42)$$

where

$$\varphi_\omega(v) = \tilde{\eta}_\omega + \|\tilde{\mu}_\omega - v\|, \quad \omega \in \Omega, \quad (43)$$

$$\mathcal{P}_S = \{\sum_{\omega \in S} c_\omega \tilde{\mu}_\omega : c_\omega > 0 \forall \omega \in S, \sum_{\omega \in S} c_\omega = 1\}.$$

We also use \mathcal{Z} to denote the union of all \mathcal{Z}_S , that is, $\mathcal{Z} = \bigcup_{S \subseteq \Omega} \mathcal{Z}_S$.

From Propositions 2, we can obtain the following lemma, showing that all MEPI measurements of \mathcal{E} are characterized by the single element of \mathcal{Z} when $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$.

Lemma 3. For MEPI of qubit state ensemble \mathcal{E} , \mathcal{Z} is always a single-element set $\{v\}$. If \mathcal{Z}_S is nonempty,

$$p_{\text{guess}}^{\text{post}} = \varphi_\omega(v) \quad \forall \omega \in S. \quad (44)$$

When $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$, there is an optimal measurement having $S \subseteq \Omega$ as the set of all non-null outcomes if and only if \mathcal{Z}_S is nonempty. Moreover, a POVM \mathcal{M} having S as the set of all non-null outcomes is optimal if and only if

$$M_\omega \propto \mathbb{1} + \frac{\tilde{\mu}_\omega - v}{\|\tilde{\mu}_\omega - v\|} \cdot \sigma \quad \forall \omega \in S. \quad (45)$$

For $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$, there is no subset $S \subseteq \Omega$ with $|S| = 1$ and $\mathcal{Z}_S \neq \emptyset$ because all MEPI measurements have more than one non-null outcome. Therefore, $\mathcal{Z} \not\subseteq \mathcal{P}_S$ for all $S \subseteq \Omega$ with $|S| = 1$.

For example, let us consider the case of $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. From Corollary 2, Lemma 2, and Eq. (23), $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$ and the polygon formed by $\{\tilde{\mu}_\omega\}_{\omega \in \Omega}$ is a parallelogram with nonempty interior \mathcal{P}_Ω illustrated in Fig. 1. The interior angle between two unit vector $\hat{\mu}_{21}$ and $\hat{\mu}_{22}$ defined in (29) is $\Phi \in (0, \pi)$ satisfying

$$\hat{\mu}_{21} \cdot \hat{\mu}_{22} = \cos \Phi. \quad (46)$$




FIG. 2: When $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, if $\mathcal{Z} \subseteq \mathcal{P}_\Omega$, the single element of \mathcal{Z} exists in one of the interiors of two line segments, $\mathcal{P}_{\{(1,1),(2,2)\}}$ (red in (a)) and $\mathcal{P}_{\{(1,2),(2,1)\}}$ (red in (b)), or in one of the interiors of four triangles, $\mathcal{P}_{\{(1,1),(1,2),(2,2)\}}$, $\mathcal{P}_{\{(1,2),(2,1),(2,2)\}}$ (green in (a)), $\mathcal{P}_{\{(1,1),(1,2),(2,1)\}}$ (green in (b)), and $\mathcal{P}_{\{(1,2),(2,1),(2,2)\}}$ (blue in (b)), but it cannot be an element of $\mathcal{Z}_{\{(1,2),(2,1),(2,2)\}}$.

Thus, we can classify this MEPI into two cases if the single element of \mathcal{Z} is in \mathcal{P}_Ω or not.

Before analyzing two cases $\mathcal{Z} \not\subseteq \mathcal{P}_\Omega$ and $\mathcal{Z} \subseteq \mathcal{P}_\Omega$, respectively, let us consider the uniqueness of MEPI measurement. When $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, Lemma 3 implies that, for each $S \subsetneq \Omega$ with $|S| \geq 2$, the MEPI measurement having S as the set of all non-null outcomes is unique because $\{\tilde{\mu}_\omega\}_{\omega \in S}$ forms a simplex with affine dimension $|S| - 1$. Since all convex combination of two different MEPI measurements are also MEPI measurements, if a non-null MEPI measurement does not exist (that is, $\mathcal{Z}_\Omega = \emptyset$), there cannot be more than one $S \subseteq \Omega$ satisfying $\mathcal{Z}_S \neq \emptyset$; thus, the MEPI measurement is unique. However, if a non-null MEPI measurements exists (that is, $\mathcal{Z}_\Omega \neq \emptyset$), the MEPI measurement is not unique because a null MEPI measurement also exists from Corollary 1.

Corollary 4. When $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, the MEPI measurement is unique if and only if a non-null MEPI measurement does not exist or, equivalently, $\mathcal{Z}_\Omega = \emptyset$.

Let us now analyze MEPI of two cases, $\mathcal{Z} \not\subseteq \mathcal{P}_\Omega$ and $\mathcal{Z} \subseteq \mathcal{P}_\Omega$. If $\mathcal{Z} \not\subseteq \mathcal{P}_\Omega$, then $\mathcal{Z} \subseteq \mathcal{P}_S$ for only one of the following four S 's:

$$\{(1,1),(1,2)\}, \{(1,1),(2,1)\}, \{(1,2),(2,2)\}, \{(2,1),(2,2)\}. \quad (47)$$

In other words, the single element of \mathcal{Z} is in one of the edges illustrated in Fig. 1. Thus, \mathcal{Z}_S is nonempty for only one S in Eq. (47). The following theorem provides a necessary and sufficient condition that \mathcal{Z}_S is nonempty for one S in Eq. (47) when $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. The proof of Theorem 1 is given in Appendix B.

Theorem 1. For MEPI of \mathcal{E} with $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, the followings are true:

$$\begin{aligned} \mathcal{Z}_{\{(1,1),(1,2)\}} \neq \emptyset &\Leftrightarrow \frac{\epsilon_{21} + \lambda_{21} \cos \Phi}{\lambda_{21} + \epsilon_{21}} \geq \frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \frac{\epsilon_{21} - \lambda_{21} \cos \Phi}{\lambda_{21} - \epsilon_{21}} \geq \frac{\lambda_{21} + \epsilon_{21}}{\lambda_{22} + \epsilon_{22}}, \\ \mathcal{Z}_{\{(1,1),(2,1)\}} \neq \emptyset &\Leftrightarrow \frac{\epsilon_{22} + \lambda_{22} \cos \Phi}{\lambda_{22} + \epsilon_{22}} \geq \frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \frac{\epsilon_{22} - \lambda_{22} \cos \Phi}{\lambda_{22} - \epsilon_{22}} \geq \frac{\lambda_{22} + \epsilon_{22}}{\lambda_{21} + \epsilon_{21}}, \\ \mathcal{Z}_{\{(2,1),(2,2)\}} &= \mathcal{Z}_{\{(1,2),(2,2)\}} = \emptyset, \end{aligned} \quad (48)$$

where ϵ_{ib} and λ_{ib} are defined in (34), and

$$p_{\text{guess}}^{\text{post}} = \begin{cases} \eta_{11} + \frac{1}{2}(\eta_{12} + \eta_{22} + \lambda_{22}) & , \mathcal{Z}_{\{(1,1),(1,2)\}} \neq \emptyset, \\ \eta_{12} + \frac{1}{2}(\eta_{11} + \eta_{21} + \lambda_{21}) & , \mathcal{Z}_{\{(1,1),(2,1)\}} \neq \emptyset. \end{cases} \quad (49)$$

When $\mathcal{Z}_{\{(1,1),(1,2)\}}$ is nonempty, $p_{\text{guess}}^{\text{prior}} - p_{\text{guess}}^{\text{post}} = \frac{1}{2}(\lambda_{21} - \epsilon_{21})$ which is a finite gap between the guessing probability and the greatest prior probability of \mathcal{E}_1 . In this case, $p_{\text{guess}}^{\text{post}}$ can be obtained by ME of \mathcal{E}_2 . More explicitly, we perform the ME measurement of \mathcal{E}_2 before obtaining the classical information b about the prepared subensemble. If $b = 1$, we ignore the result and guess the prepared state as ρ_{11} , whereas, if $b = 2$, we accept the measurement result as it is. Then, the average probability of correctly guessing the prepared state becomes $p_{\text{guess}}^{\text{post}}$. Similarly, MEPI of \mathcal{E} with $\mathcal{Z}_{\{(1,1),(2,1)\}} \neq \emptyset$ can be achieved with ME of \mathcal{E}_1 .

If $\mathcal{Z} \subseteq \mathcal{P}_\Omega$ equivalent to

$$\mathcal{Z}_{\{(1,1),(1,2)\}} = \mathcal{Z}_{\{(1,1),(2,1)\}} = \emptyset \quad (50)$$

from Theorem 1, then $\mathcal{Z} \subseteq \mathcal{P}_S$ for some S in the following seven S 's:

$$\Omega, \{(1, 1), (2, 2)\}, \{(1, 1), (1, 2), (2, 2)\}, \{(1, 1), (2, 1), (2, 2)\}, \\ \{(1, 2), (2, 1)\}, \{(1, 1), (1, 2), (2, 1)\}, \{(1, 2), (2, 1), (2, 2)\}. \quad (51)$$

In other words, the single element of \mathcal{Z} exists in one of the interiors of line segments or triangles illustrated in Fig. 2. Thus, \mathcal{Z}_S is nonempty for some S in (51). The following theorem provides a necessary and sufficient condition that $\mathcal{Z}_S \neq \emptyset$ for one S in (51) when $m = n_1 = n_2 = 2$, $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, and $\mathcal{Z} \subseteq \mathcal{P}_\Omega$. The proof of Theorem 2 is given in Appendix B.

Theorem 2. *For MEPI of \mathcal{E} with $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, the followings are true.*

(a) *We have*

$$\begin{aligned} \mathcal{Z}_{\{(1,1),(2,2)\}} \neq \emptyset &\Leftrightarrow \alpha \geq |\beta_+|/\gamma_+, \\ \mathcal{Z}_{\{(1,2),(2,1)\}} \neq \emptyset &\Leftrightarrow \alpha \leq -|\beta_-|/\gamma_-, \end{aligned} \quad (52)$$

and

$$p_{\text{guess}}^{\text{post}} = \begin{cases} \frac{1}{2}(1 + \gamma_+) & , \mathcal{Z}_{\{(1,1),(2,2)\}} \neq \emptyset, \\ \frac{1}{2}(1 + \gamma_-) & , \mathcal{Z}_{\{(1,2),(2,1)\}} \neq \emptyset, \end{cases} \quad (53)$$

where

$$\begin{aligned} \alpha &= \lambda_{21}\lambda_{22}\cos\Phi - \epsilon_{21}\epsilon_{22}, \\ \beta_{\pm} &= \epsilon_{21}\lambda_{22}^2 \mp \epsilon_{22}\lambda_{21}^2 \pm (\epsilon_{21} \mp \epsilon_{22})\lambda_{21}\lambda_{22}\cos\Phi, \\ \gamma_{\pm} &= \sqrt{\lambda_{21}^2 + \lambda_{22}^2 \pm 2\lambda_{21}\lambda_{22}\cos\Phi}. \end{aligned} \quad (54)$$

Here, ϵ_{ib} and λ_{ib} are defined in (34).

(b) *We have*

$$\begin{aligned} \mathcal{Z}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset &\Leftrightarrow \begin{cases} \frac{\epsilon_{21} + \lambda_{21}\cos\Phi}{\lambda_{21} + \epsilon_{21}} < \frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \\ \frac{\epsilon_{22} + \lambda_{22}\cos\Phi}{\lambda_{22} + \epsilon_{22}} < \frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \\ 0 \geq \alpha > -\beta_-/\gamma_-, \end{cases} \\ \mathcal{Z}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset &\Leftrightarrow \begin{cases} \frac{\epsilon_{21} - \lambda_{21}\cos\Phi}{\lambda_{21} - \epsilon_{21}} < \frac{\lambda_{21} + \epsilon_{21}}{\lambda_{22} + \epsilon_{22}}, \\ \frac{\epsilon_{22} + \lambda_{22}\cos\Phi}{\lambda_{22} + \epsilon_{22}} > -\frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \\ 0 \leq \alpha < -\beta_+/\gamma_+, \end{cases} \\ \mathcal{Z}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset &\Leftrightarrow \begin{cases} \frac{\epsilon_{22} - \lambda_{22}\cos\Phi}{\lambda_{22} - \epsilon_{22}} < \frac{\lambda_{22} + \epsilon_{22}}{\lambda_{21} + \epsilon_{21}}, \\ \frac{\epsilon_{21} + \lambda_{21}\cos\Phi}{\lambda_{21} + \epsilon_{21}} > -\frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \\ 0 \leq \alpha < \beta_+/\gamma_+, \end{cases} \\ \mathcal{Z}_{\{(1,2),(2,1),(2,2)\}} &= \emptyset, \end{aligned} \quad (55)$$

and

$$p_{\text{guess}}^{\text{post}} = \begin{cases} \tilde{\eta}_{(1,1)} + \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21}\cos(\Theta_- - \Xi_-) + \epsilon_{21}]} & , \mathcal{Z}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset, \\ \tilde{\eta}_{(1,2)} - \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21}\cos(\Theta_+ + \Xi_+) - \epsilon_{21}]} & , \mathcal{Z}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset, \\ \tilde{\eta}_{(2,1)} + \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21}\cos(\Theta_+ - \Xi_+) - \epsilon_{21}]} & , \mathcal{Z}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset, \end{cases} \quad (56)$$

where

$$\begin{aligned} \Theta_{\pm} &= \arccos \frac{\epsilon_{22}(\lambda_{21}^2 - \epsilon_{21}^2) \pm \epsilon_{21}(\lambda_{22}^2 - \epsilon_{22}^2)}{\sqrt{\lambda_{21}^2(\lambda_{22}^2 - \epsilon_{22}^2)^2 + \lambda_{22}^2(\lambda_{21}^2 - \epsilon_{21}^2)^2 \pm 2\lambda_{21}\lambda_{22}(\lambda_{21}^2 - \epsilon_{21}^2)(\lambda_{22}^2 - \epsilon_{22}^2)\cos\Phi}}, \\ \Xi_{\pm} &= \arccos \frac{\lambda_{21}(\lambda_{22}^2 - \epsilon_{22}^2) \pm \lambda_{22}(\lambda_{21}^2 - \epsilon_{21}^2)\cos\Phi}{\sqrt{\lambda_{21}^2(\lambda_{22}^2 - \epsilon_{22}^2)^2 + \lambda_{22}^2(\lambda_{21}^2 - \epsilon_{21}^2)^2 \pm 2\lambda_{21}\lambda_{22}(\lambda_{21}^2 - \epsilon_{21}^2)(\lambda_{22}^2 - \epsilon_{22}^2)\cos\Phi}}. \end{aligned} \quad (57)$$

(c) When $\mathcal{Z} \subseteq \mathcal{P}_\Omega$, a non-null MEPI measurement exists if and only if α in Eq. (54) is zero.

Finally, we mention the location of the single element of \mathcal{Z} characterizing all MEPI measurements for $p_{\text{guess}}^{\text{post}} > \tilde{\eta}_1$. When $\mathbf{z} \in \mathcal{Z}_S$ for some $S \subseteq \Omega$, the definition of \mathcal{Z}_S in Eq. (42) implies that two points \mathbf{z} and $\tilde{\boldsymbol{\mu}}_\omega$ are separated by $p_{\text{guess}}^{\text{post}} - \tilde{\eta}_\omega$ for each $\omega \in S$. Thus, the location of \mathbf{z} is easily determined by the value of $p_{\text{guess}}^{\text{post}}$ if the affine dimension of $\{\tilde{\boldsymbol{\mu}}_\omega\}_{\omega \in S}$ is one (that is, $\{\tilde{\boldsymbol{\mu}}_\omega\}_{\omega \in S}$ form a line segment). On the other hand, if the affine dimension of $\{\tilde{\boldsymbol{\mu}}_\omega\}_{\omega \in S}$ is two or three (that is, $\{\tilde{\boldsymbol{\mu}}_\omega\}_{\omega \in S}$ form a polygon or polyhedron), it is difficult to determine the location of \mathbf{z} only with the value of $p_{\text{guess}}^{\text{post}}$, and more information about \mathbf{z} is required. For example, Θ_\pm and Ξ_\pm in Eq. (57) are used to express the specific angles that determine \mathbf{z} when $m = n_1 = n_2 = 2$, $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, and $\mathcal{Z}_S \neq \emptyset$ for some $S \subseteq \Omega$ with $|S| = 3$. The details are given in Appendix B.

IV. CONCLUSION

We have analyzed MEPI of qubit state ensemble consisting of m subensembles with n_1, \dots, n_m states by considering the modified problem, that is, ME of $\prod_{b=1}^m n_b$ qubit states, and have characterized all optimal measurements. For the case where MEPI is impossible without the help of measurement, we have provided an analytic structure to characterize optimal measurements with fixed non-null outcomes. We have also shown that a null optimal measurement always exists for any MEPI of qubit states. In addition, we have analytically provided a necessary and sufficient condition that pre-measurement information is strictly more favorable than post-measurement information (that is, $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$) when $n_b = 2$ for all $b = 1, \dots, m$. For the case of $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, we have analytically provided the optimal probability of correct guessing. In this case, we have also shown that the uniqueness of optimal measurement is equivalent to the non-existence of non-null MEPI measurement. Whereas the existing results of MEPI only deal with some special cases of state ensembles consisting of equiprobable states or symmetric states [27–29], our results consider the general case of qubit state ensembles having four states and provide the analytic solutions.

Our results are closely related to the incompatibility of measurements [29–34]. Recently, it has been found that the gap between the two optimal probabilities of correct guessing with pre- and post-measurement information, $p_{\text{guess}}^{\text{prior}}$ and $p_{\text{guess}}^{\text{post}}$, is a witness of the incompatibility of measurements [31]. Moreover, the incompatibility robustness [32–34] of a given set of measurements $\{\mathcal{M}_b\}_{b \in \mathcal{B}}$ can be written by $p_{\text{guess}}^{\text{post}}$ as

$$I_R(\{\mathcal{M}_b\}_{b \in \mathcal{B}}) = \max_{\mathcal{E}} \frac{p_{\text{corr}}^{\text{prior}}(\mathcal{E}, \{\mathcal{M}_b\}_{b \in \mathcal{B}})}{p_{\text{guess}}^{\text{post}}(\mathcal{E})} - 1, \quad (58)$$

where the maximization is over all qubit state ensemble \mathcal{E} such as Eq. (11) and $p_{\text{corr}}^{\text{prior}}(\mathcal{E}, \{\mathcal{M}_b\}_{b \in \mathcal{B}})$ is the average probability of correct guessing that can be obtained by performing \mathcal{M}_b according to the pre-measurement information b . As quantum incompatibility is related with quantum resource theory [38] which is one of the most general framework in quantum information processing, it is an interesting future work to obtain an analytic form of Eq. (58).

Acknowledgements

We would like to thank Professor Joonwoo Bae for his contribution to the early stage of this work. We are grateful to Jaehee Shin and Jihwan Kim for useful comments. This work is supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF2018R1D1A1B07049420) and Institute of Information & communications Technology Planning & Evaluation (IITP) grant funded by the Korea government (MSIT) (No.2020001343, Artificial Intelligence Convergence Research Center (Hanyang University ERICA)). D.H. and J.S.K. acknowledge support from the National Research Foundation of Korea (NRF) grant funded by the Korea government (Ministry of Science and ICT) (NRF2020M3E4A1080088).

Appendix A: Proofs of Propositions

In the context of convex optimization [35], ME of \mathcal{E} is a convex optimization problem that maximizes $\sum_{i \in \Omega} \eta_i \text{Tr}[\rho_i M_i]$ subject to the following POVM constraints:

$$\begin{aligned} M_i &\succeq 0 \quad \forall i \in \Omega, \quad (\text{positive-semidefiniteness}) \\ \sum_{i \in \Omega} M_i &= \mathbb{1}. \quad (\text{completeness}) \end{aligned} \quad (A1)$$

The Lagrange dual problem is to minimize $\text{Tr}K$ subject to

$$\begin{aligned} W_i &\succeq 0 \quad \forall i \in \Omega, & (\text{positive-semidefiniteness}) \\ K &= \eta_i \rho_i + W_i \quad \forall i \in \Omega, & (\text{Lagrangian stability}) \end{aligned} \quad (\text{A2})$$

where W_i and K are Lagrange multipliers of $M_i \succeq 0$ and $\sum_{i \in \Omega} M_i = \mathbb{1}$, respectively[14, 24].

As the primal and dual problems have the same optimal value[14], primal and dual feasible variables are optimal if and only if they satisfy

$$\text{Tr}[M_i W_i] = 0 \quad \forall i \in \Omega. \quad (\text{complementary slackness}) \quad (\text{A3})$$

Therefore, ME of \mathcal{E} is equivalent to finding a set of primal and dual variables, $\{M_i\}_{i \in \Omega} \cup \{W_i\}_{i \in \Omega} \cup \{K\}$, that satisfies the so-called *Karush–Kuhn–Tucker* (KKT) condition consisting of Conditions (A1), (A2), and (A3)[24].

The primal and dual variables can be expressed as

$$M_i = p_i(\mathbb{1} + \mathbf{u}_i \cdot \boldsymbol{\sigma}), \quad W_i = \frac{r_i}{2}(\mathbb{1} + \mathbf{w}_i \cdot \boldsymbol{\sigma}), \quad K = \frac{1}{2}(s\mathbb{1} + \mathbf{v} \cdot \boldsymbol{\sigma}). \quad (\text{A4})$$

using $p_i, r_i, s \in \mathbb{R}$ and $\mathbf{u}_i, \mathbf{w}_i, \mathbf{v} \in \mathbb{R}^3$. From the expression of (A4), KKT condition can be expressed as follows:

$$\begin{aligned} (\text{P0}) \quad & p_i \geq 0, \quad \|\mathbf{u}_i\| \leq 1 \quad \forall i \in \Omega, \quad (\text{P1}) \quad \sum_{i \in \Omega} p_i = 1, \quad (\text{P2}) \quad \sum_{i \in \Omega} p_i \mathbf{u}_i = \mathbf{0}, \\ (\text{D0}) \quad & r_i \geq 0, \quad \|\mathbf{w}_i\| \leq 1 \quad \forall i \in \Omega, \quad (\text{D1}) \quad s = \eta_i + r_i \quad \forall i \in \Omega, \quad (\text{D2}) \quad \mathbf{v} = \eta_i \boldsymbol{\nu}_i + r_i \mathbf{w}_i \quad \forall i \in \Omega, \\ (\text{C0}) \quad & p_i r_i = 0 \quad \text{or} \quad \mathbf{u}_i \cdot \mathbf{w}_i = -1 \quad \forall i \in \Omega. \end{aligned} \quad (\text{A5})$$

Now, we prove Propositions 1 and 2 by using KKT condition (A5) with $\{p_i, \mathbf{u}_i\}_{i \in \Omega} \cup \{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$ as variables. We use the superscript $*$ to express the optimal variables. Note that s^*, \mathbf{v}^* , and r_i^* are always unique, whereas p_i^*, \mathbf{u}_i^* , and \mathbf{w}_i^* may not[24]. We also note that $s^* = p_{\text{guess}}$.

Proof of Proposition 1

Proof of (a). For the proof of Proposition 1(a)(\Rightarrow), assume $p_{\text{guess}} = \eta_1$. From (D1) and (D2) of (A5), this assumption implies $\eta_1 > \eta_i$ for all $i \neq 1$ because $\eta_j = \eta_1$ means $\boldsymbol{\nu}_j = \boldsymbol{\nu}_1$. Moreover,

$$r_i^* = \eta_1 - \eta_i, \quad \mathbf{w}_i^* = \frac{\eta_1 \boldsymbol{\nu}_1 - \eta_i \boldsymbol{\nu}_i}{\eta_1 - \eta_i} \quad \forall i \neq 1. \quad (\text{A6})$$

Thus,

$$\epsilon_i = \eta_1 - \eta_i \geq \|\mathbf{w}_i^*\|(\eta_1 - \eta_i) = \|\eta_1 \boldsymbol{\nu}_1 - \eta_i \boldsymbol{\nu}_i\| = \lambda_i \quad \forall i \in \Omega, \quad (\text{A7})$$

where the inequality is due to (D0) of (A5).

For the proof of Proposition 1(a)(\Leftarrow), suppose that $\epsilon_i \geq \lambda_i$ for all $i \in \Omega$. Since $\eta_j = \eta_1$ means $\boldsymbol{\nu}_j = \boldsymbol{\nu}_1$ by $\epsilon_j \geq \lambda_j$, it follows that $\eta_1 > \eta_i$ for all $i \neq 1$. Thus, $p_{\text{guess}} = \eta_1$ because (A5) holds for

$$\begin{aligned} p_i &= \delta_{1i}, & \mathbf{u}_i &= \mathbf{0}, \\ r_i &= \eta_1 - \eta_i, & \mathbf{w}_i &= \begin{cases} \mathbf{0} & \text{for } i = 1, \\ \frac{\eta_1 \boldsymbol{\nu}_1 - \eta_i \boldsymbol{\nu}_i}{\eta_1 - \eta_i} & \text{for } i \neq 1, \end{cases} \\ s &= \eta_1, & \mathbf{v} &= \eta_1 \boldsymbol{\nu}_1, \end{aligned} \quad (\text{A8})$$

where δ_{ij} is the Kronecker delta. This completes our proof of Proposition 1(a). We also note that $\epsilon_i \geq \lambda_i$ for all $i \in \Omega$ is equivalent to $\mathcal{Z}_{\{1\}} \neq \emptyset$ since $\mathcal{P}_{\{1\}}$ has only one element $\eta_1 \boldsymbol{\nu}_1$. \square

Proof of (b). For the proof of Proposition 1(b)(\Rightarrow), we assume that $p_{\text{guess}} = \eta_1$ and \mathcal{M} is an optimal POVM expressed by $\{p_i, \mathbf{u}_i\}_{i \in \Omega}$. Then, there is $\{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$ such that (A5) hold for $\{p_i, \mathbf{u}_i\}_{i \in \Omega} \cup \{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$.

Let us consider a non-null outcome $k \neq 1$. Since $s = p_{\text{guess}} = \eta_1$, it follows from (D1) and (D2) of (A5) that

$$r_k = \eta_1 - \eta_k, \quad r_k \mathbf{w}_k = \eta_1 \boldsymbol{\nu}_1 - \eta_k \boldsymbol{\nu}_k. \quad (\text{A9})$$

r_k is nonzero because $\eta_k = \eta_1$ means $\boldsymbol{\nu}_k = \boldsymbol{\nu}_1$ by $\epsilon_k \geq \lambda_k$, therefore,

$$\mathbf{w}_k = \frac{\eta_1 \boldsymbol{\nu}_1 - \eta_k \boldsymbol{\nu}_k}{\eta_1 - \eta_k}. \quad (\text{A10})$$

Since p_k and r_k are nonzero, it follows from (C0) of (A5) that \mathbf{u}_k and \mathbf{w}_k are unit vectors satisfying $\mathbf{u}_k \cdot \mathbf{w}_k = -1$. Thus,

$$\epsilon_k = \epsilon_k \|\mathbf{w}_k\| = \lambda_k \quad (\text{A11})$$

and

$$\mathbf{u}_k = \frac{\eta_k \boldsymbol{\nu}_k - \eta_1 \boldsymbol{\nu}_1}{\|\eta_k \boldsymbol{\nu}_k - \eta_1 \boldsymbol{\nu}_1\|}, \quad (\text{A12})$$

that is, Eq. (9) holds for $i = k$. We also note that Eq. (9) is obviously satisfied for every null outcome. Therefore, \mathcal{M} satisfies Eq. (9).

For the proof of Proposition 1(b)(\Leftarrow), suppose that $p_{\text{guess}} = \eta_1$ and \mathcal{M} is a POVM satisfying Eq. (9). Let $\{p_i, \mathbf{u}_i\}_{i \in \Omega}$ be a set of primal variables expressing \mathcal{M} . Also, let $\{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$ be a set of dual variables such that

$$\begin{aligned} r_i &= \eta_1 - \eta_i, \quad \mathbf{w}_i = \begin{cases} \mathbf{0} & \text{for } i = 1, \\ \frac{\eta_1 \boldsymbol{\nu}_1 - \eta_i \boldsymbol{\nu}_i}{\eta_1 - \eta_i} & \text{for } i \neq 1, \end{cases} \\ s &= \eta_1, \quad \mathbf{v} = \eta_1 \boldsymbol{\nu}_1. \end{aligned} \quad (\text{A13})$$

Then, (A5) holds for $\{p_i, \mathbf{u}_i\}_{i \in \Omega} \cup \{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$. Therefore, \mathcal{M} is optimal. \square

Proof of Proposition 2

Proof of (a), (b)(\Leftarrow), and (c)(\Leftarrow). Let us assume that S is a subset of Ω with $\mathbf{v} \in \mathcal{Z}_S$. The definition of \mathcal{Z}_S implies that there exists a real number s fulfilling $\varphi_i(\mathbf{v}) = s$ for all $i \in S$. We will show that $s = s^*$ and $\mathbf{v} = \mathbf{v}^*$. Since $s^* = p_{\text{guess}}$ and \mathbf{v}^* is unique, if $s = s^*$ and $\mathbf{v} = \mathbf{v}^*$, then $p_{\text{guess}} = \varphi_i(\mathbf{v})$ for all $i \in S$ and $\mathcal{Z}_S = \{\mathbf{v}^*\}$ which implies that \mathcal{Z} is a single-element set having only one element \mathbf{v} , that is, $\mathcal{Z} = \{\mathbf{v}\}$.

First, let us consider the case of $\eta_k = s$ for some $k \in \Omega$. Since $s \geq \varphi_i(\mathbf{v})$ for all $i \in \Omega$ from the definition of \mathcal{Z}_S and $\varphi_j(\mathbf{v}) = s$ for all $j \in S$, we have

$$\|\eta_k \boldsymbol{\nu}_k - \mathbf{v}\| = \varphi_k(\mathbf{v}) - \eta_k \leq s - \eta_k = 0, \quad (\text{A14})$$

which means $\mathbf{v} = \eta_k \boldsymbol{\nu}_k$. Thus, $k = 1$ because

$$\eta_k = s \geq \varphi_i(\eta_k \boldsymbol{\nu}_k) = \eta_i + \|\eta_i \boldsymbol{\nu}_i - \eta_k \boldsymbol{\nu}_k\| > \eta_i \quad \forall i \neq k. \quad (\text{A15})$$

(A5) holds for the primal and dual variables satisfying Eq. (A8); therefore, $s = s^*$ and $\mathbf{v} = \mathbf{v}^*$.

Now, let us consider the case of $\eta_i \neq s$ for all $i \in \Omega$. In this case, $\eta_i \boldsymbol{\nu}_i \neq \mathbf{v}$ for all $i \in S$ because otherwise

$$s = \varphi_j(\mathbf{v}) = \eta_j + \|\eta_j \boldsymbol{\nu}_j - \mathbf{v}\| = \eta_j \quad (\text{A16})$$

for some $j \in S$ with $\eta_j \boldsymbol{\nu}_j = \mathbf{v}$. Since $\mathbf{v} \in \mathcal{P}_S$, there exists $\{c_i\}_{i \in \Omega} \subseteq \mathbb{R}$ such that

$$c_i > 0 \quad \forall i \in S, \quad c_i = 0 \quad \forall i \notin S, \quad \sum_{i \in S} c_i = 1, \quad \sum_{i \in S} c_i (\eta_i \boldsymbol{\nu}_i) = \mathbf{v}. \quad (\text{A17})$$

Thus, $s = s^*$ and $\mathbf{v} = \mathbf{v}^*$ because (A5) holds for $\{s, \mathbf{v}\}$ along with $\{p_i, \mathbf{u}_i\}_{i \in \Omega} \cup \{r_i, \mathbf{w}_i\}_{i \in \Omega}$ satisfying

$$\begin{aligned} p_i &= \frac{c_i \|\eta_i \boldsymbol{\nu}_i - \mathbf{v}\|}{\sum_{j \in S} c_j \|\eta_j \boldsymbol{\nu}_j - \mathbf{v}\|}, & r_i &= s - \eta_i, \\ \mathbf{u}_i &= \begin{cases} \frac{\eta_i \boldsymbol{\nu}_i - \mathbf{v}}{\|\eta_i \boldsymbol{\nu}_i - \mathbf{v}\|}, & i \in S, \\ \mathbf{0}, & i \notin S, \end{cases} & \mathbf{w}_i &= \begin{cases} \frac{\mathbf{v} - \eta_i \boldsymbol{\nu}_i}{\|\mathbf{v} - \eta_i \boldsymbol{\nu}_i\|}, & i \in S, \\ \frac{\mathbf{v} - \eta_i \boldsymbol{\nu}_i}{s - \eta_i}, & i \notin S. \end{cases} \end{aligned} \quad (\text{A18})$$

This completes our proof of Proposition 2(a).

Moreover, the optimal POVM corresponding to $\{p_i, \mathbf{u}_i\}_{i \in \Omega}$ satisfying Eq. (A18) has S as the set of all non-null outcomes. Thus, Proposition 2(b)(\Leftarrow) is proved. We also note that every POVM \mathcal{M} fulfilling Condition (10) with $M_i = 0$ for all $i \notin S$ is optimal because (A5) holds for $\{s, \mathbf{v}\}$ along with $\{p_i, \mathbf{u}_i\}_{i \in \Omega}$ expressing \mathcal{M} and $\{r_i, \mathbf{w}_i\}_{i \in \Omega}$ satisfying (A18); therefore, Proposition 2(c)(\Leftarrow) is also proved. \square

Proof of (b)(\Rightarrow) and (c)(\Rightarrow). Suppose that $p_{\text{guess}} > \eta_1$ and \mathcal{M} is an optimal POVM for ME of \mathcal{E} having \mathbf{S} as the set of all non-null outcomes. We also assume that $\{p_i, \mathbf{u}_i\}_{i \in \Omega}$ is a set of primal variables expressing \mathcal{M} . Then, there exists a set of dual variables $\{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$ such that (A5) holds for $\{p_i, \mathbf{u}_i\}_{i \in \Omega} \cup \{r_i, \mathbf{w}_i\}_{i \in \Omega} \cup \{s, \mathbf{v}\}$. Since $p_{\text{guess}} > \eta_1$, it follows from (D1) of (A5) that r_i is nonzero for all $i \in \Omega$. Furthermore, (C0) of (A5) implies $\|\mathbf{u}_j\| = \|\mathbf{w}_j\| = 1$ and $\mathbf{u}_j \cdot \mathbf{w}_j = -1$ for each $j \in \mathbf{S}$ because both p_j and r_j are nonzero. Thus, from (D2) of (A5), we have

$$r_j = \|\eta_j \boldsymbol{\nu}_j - \mathbf{v}\|, \quad \mathbf{u}_j = \frac{\eta_j \boldsymbol{\nu}_j - \mathbf{v}}{\|\eta_j \boldsymbol{\nu}_j - \mathbf{v}\|} \quad \forall j \in \mathbf{S}. \quad (\text{A19})$$

The right equality in (A19) is equivalent to Condition (10); therefore, Proposition 2(c)(\Rightarrow) is proved.

Now, we show $\mathcal{Z}_{\mathbf{S}} \neq \emptyset$ to prove Proposition 2(b)(\Rightarrow). From (D0) and (D2) of (A5), we can see that

$$r_i \geq \|\mathbf{r}_i \mathbf{w}_i\| = \|\eta_i \boldsymbol{\nu}_i - \mathbf{v}\| \quad \forall i \in \Omega. \quad (\text{A20})$$

Thus,

$$\begin{aligned} \varphi_i(\mathbf{v}) &= q_i + \|\eta_i \boldsymbol{\nu}_i - \mathbf{v}\| = q_i + r_i = q_j + r_j \\ &\geq q_j + \|\eta_j \boldsymbol{\nu}_j - \mathbf{v}\| = \varphi_j(\mathbf{v}) \quad \forall i \in \mathbf{S}, j \in \Omega, \end{aligned} \quad (\text{A21})$$

where the second equality is due to Eq. (A19) and the third equality is from (D1) of (A5). Since the inequality in (A21) for $j \in \mathbf{S}$ becomes an equality by Eq. (A19), $\mathcal{Z}_{\mathbf{S}}$ is a nonempty set with \mathbf{v} . Note that $\mathbf{v} \in \mathcal{P}_{\mathbf{S}}$ because

$$\sum_{i \in \mathbf{S}} c_i (\eta_i \boldsymbol{\nu}_i) = \sum_{i \in \mathbf{S}} c_i (\mathbf{v} - r_i \mathbf{w}_i) = \mathbf{v} - \frac{\sum_{i \in \mathbf{S}} (p_i \mathbf{w}_i)}{\sum_{j \in \mathbf{S}} (p_j / r_j)} = \mathbf{v} + \frac{\sum_{i \in \mathbf{S}} (p_i \mathbf{u}_i)}{\sum_{j \in \mathbf{S}} (p_j / r_j)} = \mathbf{v}, \quad (\text{A22})$$

where

$$c_i = \frac{(p_i / r_i)}{\sum_{j \in \mathbf{S}} (p_j / r_j)}. \quad (\text{A23})$$

The first equality of (A22) is due to (D2) of (A5), the second equality is from $\sum_{i \in \mathbf{S}} c_i = 1$, and the third and last equality follow from (C0) and (P2), respectively. Therefore, Proposition 2(b)(\Rightarrow) is proved. \square

Appendix B: Proofs of Theorems

In this section, we prove Theorems 1 and 2 using $\mathcal{X}_{\mathbf{S}'}^{\mathbf{S}}$ and $\mathcal{Y}_{\mathbf{S}'}^{\mathbf{S}}$ ($\mathbf{S}' \subseteq \mathbf{S} \subseteq \Omega$) defined as

$$\begin{aligned} \mathcal{X}_{\mathbf{S}'}^{\mathbf{S}} &:= \{\mathbf{v} \in \mathcal{P}_{\mathbf{S}'} : \varphi_{\boldsymbol{\omega}}(\mathbf{v}) = \varphi_{\boldsymbol{\omega}'}(\mathbf{v}) \quad \forall (\boldsymbol{\omega}, \boldsymbol{\omega}') \in \mathbf{S}' \times \mathbf{S}', \\ &\quad \varphi_{\boldsymbol{\omega}}(\mathbf{v}) \geq \varphi_{\boldsymbol{\omega}'}(\mathbf{v}) \quad \forall (\boldsymbol{\omega}, \boldsymbol{\omega}') \in \mathbf{S}' \times (\mathbf{S} - \mathbf{S}')\}, \\ \mathcal{Y}_{\mathbf{S}'}^{\mathbf{S}} &:= \{\mathbf{v} \in \mathbb{R}^3 : \varphi_{\boldsymbol{\omega}}(\mathbf{v}) \geq \varphi_{\boldsymbol{\omega}'}(\mathbf{v}) \quad \forall (\boldsymbol{\omega}, \boldsymbol{\omega}') \in \mathbf{S}' \times (\Omega - \mathbf{S})\}, \end{aligned} \quad (\text{B1})$$

where we denote

$$\mathcal{X}_{\mathbf{S}} := \mathcal{X}_{\mathbf{S}}^{\mathbf{S}}, \quad \mathcal{Y}_{\mathbf{S}} := \mathcal{Y}_{\mathbf{S}}^{\mathbf{S}}. \quad (\text{B2})$$

Note that $\mathcal{X}_{\mathbf{S}'}^{\mathbf{S}} \cap \mathcal{Y}_{\mathbf{S}'}^{\mathbf{S}} = \mathcal{Z}_{\mathbf{S}'}$ for any $\mathbf{S}' \subseteq \mathbf{S} \subseteq \Omega$. We also note that $\bigcup_{\mathbf{S}' \subseteq \mathbf{S}} \mathcal{X}_{\mathbf{S}'}^{\mathbf{S}}$ is a single-element set for any $\mathbf{S} \subseteq \Omega$ in terms of ME of $\{\tilde{\eta}_{\boldsymbol{\omega}}, \tilde{\rho}_{\boldsymbol{\omega}}\}_{\boldsymbol{\omega} \in \mathbf{S}}$; thus, at least one of $\mathcal{X}_{\mathbf{S}'}^{\mathbf{S}}$ and $\mathcal{X}_{\mathbf{S}''}^{\mathbf{S}}$ is empty for any $\mathbf{S}', \mathbf{S}'' \subseteq \mathbf{S}$ with $\mathcal{P}_{\mathbf{S}'} \cap \mathcal{P}_{\mathbf{S}''} = \emptyset$. In particular, we use the following two lemmas when proving Theorem 2.

Lemma 4. *When $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, if \mathbf{S} is a proper subset of Ω with $\mathcal{X}_{\mathbf{S}} \neq \emptyset$ and \mathbf{S}' is a proper subset of \mathbf{S} satisfying $\mathbf{S} - \mathbf{S}' = \{\boldsymbol{\omega}\}$ and $\mathcal{X}_{\mathbf{S}'} = \{\mathbf{v}\}$, then*

$$\varphi_{\boldsymbol{\omega}'}(\mathbf{v}) < \varphi_{\boldsymbol{\omega}}(\mathbf{v}) \quad \forall \boldsymbol{\omega}' \in \mathbf{S}'. \quad (\text{B3})$$

Proof. $\mathcal{P}_{\mathbf{S}}$ and $\mathcal{P}_{\mathbf{S}'}$ are disjoint for any $\mathbf{S}' \subsetneq \mathbf{S} \subsetneq \Omega$ because $\{\tilde{\boldsymbol{\mu}}_{\boldsymbol{\omega}}\}_{\boldsymbol{\omega} \in \Omega}$ forms a parallelogram with nonempty interior when $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. Thus, if $\mathcal{X}_{\mathbf{S}}$ is nonempty for some $\mathbf{S} \subsetneq \Omega$, then $\mathcal{X}_{\mathbf{S}'}^{\mathbf{S}}$ is empty for all $\mathbf{S}' \subsetneq \mathbf{S}$. Now, we assume that \mathbf{S} is a proper subset of Ω with $\mathcal{X}_{\mathbf{S}} \neq \emptyset$ and \mathbf{S}' is a proper subset of \mathbf{S} satisfying $\mathbf{S} - \mathbf{S}' = \{\boldsymbol{\omega}\}$ and $\mathcal{X}_{\mathbf{S}'} = \{\mathbf{v}\}$. If $\varphi_{\boldsymbol{\omega}'}(\mathbf{v}) \geq \varphi_{\boldsymbol{\omega}}(\mathbf{v})$ for some $\boldsymbol{\omega}' \in \mathbf{S}'$, then $\mathcal{X}_{\mathbf{S}'}^{\mathbf{S}}$ becomes a nonempty set with \mathbf{v} as a element. Thus, $\mathcal{X}_{\mathbf{S}'}^{\mathbf{S}} = \emptyset$ implies (B3), which completes our proof. \square

Lemma 5. When $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$, \mathcal{X}_S is nonempty for $S \subsetneq \Omega$ if $\mathcal{X}_{S'} \neq \emptyset$ and $\mathcal{X}_{S'}^S = \emptyset$ for all $S' \subsetneq S$ with $|S - S'| = 1$.

Proof. For each $S'' \subsetneq S' \subsetneq S$ with $|S - S'| = 1$, $\mathcal{X}_{S''}^{S'}$ is empty because $\bigcup_{S'' \subsetneq S'} \mathcal{X}_{S''}^{S'}$ is a single-element set and $\mathcal{X}_{S'}$ is a nonempty set. Note that $\mathcal{P}_{S'}$ and $\mathcal{P}_{S''}$ are disjoint for any $S'' \subsetneq S' \subsetneq S$ since $\{\tilde{\mu}_\omega\}_{\omega \in \Omega}$ forms a parallelogram with nonempty interior when $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. For $S'' \subsetneq S' \subsetneq S$, emptiness of $\mathcal{X}_{S''}^{S'}$ implies emptiness of $\mathcal{X}_{S''}^S$, as $\mathcal{X}_{S''}^S \subseteq \mathcal{X}_{S''}^{S'}$; therefore, $\mathcal{X}_{S'}^S$ is empty for all $S' \subsetneq S$. Since $\bigcup_{S' \subsetneq S} \mathcal{X}_{S'}^S$ is a single-element set, \mathcal{X}_S is nonempty if $\mathcal{X}_{S'}^S$ is empty for all $S' \subsetneq S$. Thus, \mathcal{X}_S is nonempty. \square

Proof of Theorem 1

Proof. Suppose that $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. We can easily verify that

$$\begin{aligned} \mathcal{X}_{\{(1,1),(1,2)\}} &= \{\mathbf{v}_{11}\}, & \varphi_{(1,1)}(\mathbf{v}_{11}) + \varphi_{(1,2)}(\mathbf{v}_{11}) &= 2\eta_{11} + \eta_{12} + \eta_{22} + \lambda_{22}, \\ \mathcal{X}_{\{(1,1),(2,1)\}} &= \{\mathbf{v}_{12}\}, & \varphi_{(1,1)}(\mathbf{v}_{12}) + \varphi_{(2,1)}(\mathbf{v}_{12}) &= 2\eta_{12} + \eta_{11} + \eta_{21} + \lambda_{21}, \\ \mathcal{X}_{\{(2,1),(2,2)\}} &= \{\mathbf{v}_{21}\}, & \varphi_{(2,1)}(\mathbf{v}_{21}) + \varphi_{(2,2)}(\mathbf{v}_{21}) &= 2\eta_{21} + \eta_{12} + \eta_{22} + \lambda_{22}, \\ \mathcal{X}_{\{(1,2),(2,2)\}} &= \{\mathbf{v}_{22}\}, & \varphi_{(1,2)}(\mathbf{v}_{22}) + \varphi_{(2,2)}(\mathbf{v}_{22}) &= 2\eta_{22} + \eta_{11} + \eta_{21} + \lambda_{21}, \end{aligned} \quad (\text{B4})$$

where

$$\begin{aligned} \mathbf{v}_{11} &= \left(\frac{\lambda_{22} + \epsilon_{22}}{2\lambda_{22}} \right) \tilde{\mu}_{(1,1)} + \left(\frac{\lambda_{22} - \epsilon_{22}}{2\lambda_{22}} \right) \tilde{\mu}_{(1,2)}, & \mathbf{v}_{12} &= \left(\frac{\lambda_{21} + \epsilon_{21}}{2\lambda_{21}} \right) \tilde{\mu}_{(1,1)} + \left(\frac{\lambda_{21} - \epsilon_{21}}{2\lambda_{21}} \right) \tilde{\mu}_{(2,1)}, \\ \mathbf{v}_{21} &= \left(\frac{\lambda_{22} + \epsilon_{22}}{2\lambda_{22}} \right) \tilde{\mu}_{(2,1)} + \left(\frac{\lambda_{22} - \epsilon_{22}}{2\lambda_{22}} \right) \tilde{\mu}_{(2,2)}, & \mathbf{v}_{22} &= \left(\frac{\lambda_{21} + \epsilon_{21}}{2\lambda_{21}} \right) \tilde{\mu}_{(1,2)} + \left(\frac{\lambda_{21} - \epsilon_{21}}{2\lambda_{21}} \right) \tilde{\mu}_{(2,2)}. \end{aligned} \quad (\text{B5})$$

Since $\mathcal{X}_S \cap \mathcal{Y}_S = \mathcal{Z}_S$ for all $S \subseteq \Omega$, it follows from Eq. (B4) that

$$\begin{aligned} \mathcal{Z}_{\{(1,1),(1,2)\}} &\neq \emptyset \Leftrightarrow \mathbf{v}_{11} \in \mathcal{Y}_{\{(1,1),(1,2)\}}, \\ \mathcal{Z}_{\{(1,1),(2,1)\}} &\neq \emptyset \Leftrightarrow \mathbf{v}_{12} \in \mathcal{Y}_{\{(1,1),(2,1)\}}, \\ \mathcal{Z}_{\{(2,1),(2,2)\}} &\neq \emptyset \Leftrightarrow \mathbf{v}_{21} \in \mathcal{Y}_{\{(2,1),(2,2)\}}, \\ \mathcal{Z}_{\{(1,2),(2,2)\}} &\neq \emptyset \Leftrightarrow \mathbf{v}_{22} \in \mathcal{Y}_{\{(1,2),(2,2)\}}, \end{aligned} \quad (\text{B6})$$

and

$$p_{\text{guess}}^{\text{post}} = \begin{cases} \eta_{11} + \frac{1}{2}(\eta_{12} + \eta_{22} + \lambda_{22}), & \mathcal{Z}_{\{(1,1),(1,2)\}} \neq \emptyset, \\ \eta_{12} + \frac{1}{2}(\eta_{11} + \eta_{21} + \lambda_{21}), & \mathcal{Z}_{\{(1,1),(2,1)\}} \neq \emptyset, \\ \eta_{21} + \frac{1}{2}(\eta_{12} + \eta_{22} + \lambda_{22}), & \mathcal{Z}_{\{(2,1),(2,2)\}} \neq \emptyset, \\ \eta_{22} + \frac{1}{2}(\eta_{11} + \eta_{21} + \lambda_{21}), & \mathcal{Z}_{\{(1,2),(2,2)\}} \neq \emptyset. \end{cases} \quad (\text{B7})$$

From definition of \mathcal{Y}_S , we can see that

$$\begin{aligned} \mathbf{v}_{11} \in \mathcal{Y}_{\{(1,1),(1,2)\}} &\Leftrightarrow \varphi_{(1,1)}(\mathbf{v}_{11}) \geq \varphi_{(2,1)}(\mathbf{v}_{11}), & \varphi_{(1,2)}(\mathbf{v}_{11}) &\geq \varphi_{(2,2)}(\mathbf{v}_{11}), \\ \mathbf{v}_{12} \in \mathcal{Y}_{\{(1,1),(2,1)\}} &\Leftrightarrow \varphi_{(1,1)}(\mathbf{v}_{12}) \geq \varphi_{(1,2)}(\mathbf{v}_{12}), & \varphi_{(2,1)}(\mathbf{v}_{12}) &\geq \varphi_{(2,2)}(\mathbf{v}_{12}), \\ \mathbf{v}_{21} \in \mathcal{Y}_{\{(2,1),(2,2)\}} &\Leftrightarrow \varphi_{(1,1)}(\mathbf{v}_{21}) \leq \varphi_{(2,1)}(\mathbf{v}_{21}), & \varphi_{(1,2)}(\mathbf{v}_{21}) &\leq \varphi_{(2,2)}(\mathbf{v}_{21}), \\ \mathbf{v}_{22} \in \mathcal{Y}_{\{(1,2),(2,2)\}} &\Leftrightarrow \varphi_{(1,1)}(\mathbf{v}_{22}) \leq \varphi_{(1,2)}(\mathbf{v}_{22}), & \varphi_{(2,1)}(\mathbf{v}_{22}) &\leq \varphi_{(2,2)}(\mathbf{v}_{22}). \end{aligned} \quad (\text{B8})$$

It is straightforward to verify that

$$\begin{aligned}
\varphi_{(1,1)}(\mathbf{v}_{11}) - \varphi_{(2,1)}(\mathbf{v}_{11}) &= \frac{(\epsilon_{21} + \lambda_{21} \cos \Phi)(\lambda_{22} - \epsilon_{22}) - (\lambda_{21}^2 - \epsilon_{21}^2)}{\epsilon_{21} + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_{11}\| + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_{11}\|}, \\
\varphi_{(1,2)}(\mathbf{v}_{11}) - \varphi_{(2,2)}(\mathbf{v}_{11}) &= \frac{(\epsilon_{21} - \lambda_{21} \cos \Phi)(\lambda_{22} + \epsilon_{22}) - (\lambda_{21}^2 - \epsilon_{21}^2)}{\epsilon_{21} + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_{11}\| + \|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}_{11}\|}, \\
\varphi_{(1,1)}(\mathbf{v}_{12}) - \varphi_{(1,2)}(\mathbf{v}_{12}) &= \frac{(\epsilon_{22} + \lambda_{22} \cos \Phi)(\lambda_{21} - \epsilon_{21}) - (\lambda_{22}^2 - \epsilon_{22}^2)}{\epsilon_{22} + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_{12}\| + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_{12}\|}, \\
\varphi_{(2,1)}(\mathbf{v}_{12}) - \varphi_{(2,2)}(\mathbf{v}_{12}) &= \frac{(\epsilon_{22} - \lambda_{22} \cos \Phi)(\lambda_{21} + \epsilon_{21}) - (\lambda_{22}^2 - \epsilon_{22}^2)}{\epsilon_{22} + \|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}_{12}\| + \|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}_{12}\|}, \\
\varphi_{(1,1)}(\mathbf{v}_{21}) - \varphi_{(2,1)}(\mathbf{v}_{21}) &= \frac{(\epsilon_{21} + \lambda_{21} \cos \Phi)(\lambda_{22} - \epsilon_{22}) + (\lambda_{21}^2 - \epsilon_{21}^2)}{\|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}_{21}\| + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_{21}\| - \epsilon_{21}}, \\
\varphi_{(1,2)}(\mathbf{v}_{21}) - \varphi_{(2,2)}(\mathbf{v}_{21}) &= \frac{(\epsilon_{21} - \lambda_{21} \cos \Phi)(\lambda_{22} + \epsilon_{22}) + (\lambda_{21}^2 - \epsilon_{21}^2)}{\|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}_{21}\| + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_{21}\| - \epsilon_{21}}, \\
\varphi_{(1,1)}(\mathbf{v}_{22}) - \varphi_{(1,2)}(\mathbf{v}_{22}) &= \frac{(\epsilon_{22} + \lambda_{22} \cos \Phi)(\lambda_{21} - \epsilon_{21}) + (\lambda_{22}^2 - \epsilon_{22}^2)}{\|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_{22}\| + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_{22}\| - \epsilon_{22}}, \\
\varphi_{(2,1)}(\mathbf{v}_{22}) - \varphi_{(2,2)}(\mathbf{v}_{22}) &= \frac{(\epsilon_{22} - \lambda_{22} \cos \Phi)(\lambda_{21} + \epsilon_{21}) + (\lambda_{22}^2 - \epsilon_{22}^2)}{\|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}_{22}\| + \|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}_{22}\| - \epsilon_{22}}.
\end{aligned} \tag{B9}$$

Therefore,

$$\mathcal{Z}_{\{(1,1),(1,2)\}} \neq \emptyset \Leftrightarrow \frac{\epsilon_{21} + \lambda_{21} \cos \Phi}{\lambda_{21} + \epsilon_{21}} \geq \frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \quad \frac{\epsilon_{21} - \lambda_{21} \cos \Phi}{\lambda_{21} - \epsilon_{21}} \geq \frac{\lambda_{21} + \epsilon_{21}}{\lambda_{22} + \epsilon_{22}}, \tag{B10a}$$

$$\mathcal{Z}_{\{(1,1),(2,1)\}} \neq \emptyset \Leftrightarrow \frac{\epsilon_{22} + \lambda_{22} \cos \Phi}{\lambda_{22} + \epsilon_{22}} \geq \frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \quad \frac{\epsilon_{22} - \lambda_{22} \cos \Phi}{\lambda_{22} - \epsilon_{22}} \geq \frac{\lambda_{22} + \epsilon_{22}}{\lambda_{21} + \epsilon_{21}}, \tag{B10b}$$

$$\mathcal{Z}_{\{(2,1),(2,2)\}} \neq \emptyset \Leftrightarrow \frac{\epsilon_{21} + \lambda_{21} \cos \Phi}{\lambda_{21} + \epsilon_{21}} \leq -\frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \quad \frac{\epsilon_{21} - \lambda_{21} \cos \Phi}{\lambda_{21} - \epsilon_{21}} \leq -\frac{\lambda_{21} + \epsilon_{21}}{\lambda_{22} + \epsilon_{22}}, \tag{B10c}$$

$$\mathcal{Z}_{\{(1,2),(2,2)\}} \neq \emptyset \Leftrightarrow \frac{\epsilon_{22} + \lambda_{22} \cos \Phi}{\lambda_{22} + \epsilon_{22}} \leq -\frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \quad \frac{\epsilon_{22} - \lambda_{22} \cos \Phi}{\lambda_{22} - \epsilon_{22}} \leq -\frac{\lambda_{22} + \epsilon_{22}}{\lambda_{21} + \epsilon_{21}}. \tag{B10d}$$

Moreover, at least one of the two left-hand sides of Inequalities (B10c) is positive, but the two right-hand sides are both negative; thus, $\mathcal{Z}_{\{(2,1),(2,2)\}}$ is always empty. For similar reasons, $\mathcal{Z}_{\{(1,2),(2,2)\}}$ is always empty. \square

Proof of Theorem 2

Proof of (a). Suppose $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. Then, we can easily verify that

$$\begin{aligned}
\mathcal{X}_{\{(1,1),(2,2)\}} \neq \emptyset &\Rightarrow \mathcal{X}_{\{(1,1),(2,2)\}} = \{\mathbf{v}_+\}, \quad \varphi_{(1,1)}(\mathbf{v}_+) + \varphi_{(2,2)}(\mathbf{v}_+) = 1 + \gamma_+, \\
\mathcal{X}_{\{(1,2),(2,1)\}} \neq \emptyset &\Rightarrow \mathcal{X}_{\{(1,2),(2,1)\}} = \{\mathbf{v}_-\}, \quad \varphi_{(1,2)}(\mathbf{v}_-) + \varphi_{(2,1)}(\mathbf{v}_-) = 1 + \gamma_-,
\end{aligned} \tag{B11}$$

where

$$\begin{aligned}
\mathbf{v}_+ &= \left(\frac{\gamma_+ + \epsilon_{21} + \epsilon_{22}}{2\gamma_+} \right) \tilde{\boldsymbol{\mu}}_{(1,1)} + \left(\frac{\gamma_+ - \epsilon_{21} - \epsilon_{22}}{2\gamma_+} \right) \tilde{\boldsymbol{\mu}}_{(2,2)}, \\
\mathbf{v}_- &= \left(\frac{\gamma_- + \epsilon_{21} - \epsilon_{22}}{2\gamma_-} \right) \tilde{\boldsymbol{\mu}}_{(1,2)} + \left(\frac{\gamma_- - \epsilon_{21} + \epsilon_{22}}{2\gamma_-} \right) \tilde{\boldsymbol{\mu}}_{(2,1)}.
\end{aligned} \tag{B12}$$

Since $\mathcal{X}_S \cap \mathcal{Y}_S = \mathcal{Z}_S$ for all $S \subseteq \Omega$, Eq. (B11) implies

$$\begin{aligned}
\mathcal{Z}_{\{(1,1),(2,2)\}} \neq \emptyset &\Rightarrow \mathbf{v}_+ \in \mathcal{Y}_{\{(1,1),(2,2)\}}, \\
\mathcal{Z}_{\{(1,2),(2,1)\}} \neq \emptyset &\Rightarrow \mathbf{v}_- \in \mathcal{Y}_{\{(1,2),(2,1)\}},
\end{aligned} \tag{B13}$$

as well as the validity of Eq. (53).

From definition of \mathcal{Y}_S , we have

$$\begin{aligned}
\mathbf{v}_+ \in \mathcal{Y}_{\{(1,1),(2,2)\}} &\Leftrightarrow \varphi_{(1,1)}(\mathbf{v}_+) \geq \varphi_{(1,2)}(\mathbf{v}_+), \quad \varphi_{(1,1)}(\mathbf{v}_+) \geq \varphi_{(2,1)}(\mathbf{v}_+), \\
\mathbf{v}_- \in \mathcal{Y}_{\{(1,2),(2,1)\}} &\Leftrightarrow \varphi_{(1,1)}(\mathbf{v}_-) \leq \varphi_{(1,2)}(\mathbf{v}_-), \quad \varphi_{(2,2)}(\mathbf{v}_-) \leq \varphi_{(1,2)}(\mathbf{v}_-).
\end{aligned} \tag{B14}$$

It is straightforward to show that

$$\begin{aligned}
\varphi_{(1,1)}(\mathbf{v}_+) - \varphi_{(1,2)}(\mathbf{v}_+) &= \frac{\gamma_+ \alpha - \beta_+}{\gamma_+ (\epsilon_{21} + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_+\| + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_+\|)}, \\
\varphi_{(1,1)}(\mathbf{v}_+) - \varphi_{(2,1)}(\mathbf{v}_+) &= \frac{\gamma_+ \alpha + \beta_+}{\gamma_+ (\epsilon_{21} + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_+\| + \|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}_+\|)}, \\
\varphi_{(1,1)}(\mathbf{v}_-) - \varphi_{(1,2)}(\mathbf{v}_-) &= \frac{\gamma_- \alpha + \beta_-}{\gamma_- (\|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_-\| + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}_-\| - \epsilon_{22})}, \\
\varphi_{(1,1)}(\mathbf{v}_-) - \varphi_{(2,1)}(\mathbf{v}_-) &= \frac{\gamma_- \alpha - \beta_-}{\gamma_- (\epsilon_{21} + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}_-\| + \|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}_-\|)}.
\end{aligned} \tag{B15}$$

By applying Eq. (B15) to (B14) and (B14) to (B13), we can show that (52)(\Rightarrow) is true.

Now, we prove that (52)(\Leftarrow) is true. Because non-negativity and non-positivity of α imply $\gamma_+^2 \geq (\epsilon_{21} + \epsilon_{22})^2$ and $\gamma_-^2 \geq (\epsilon_{21} - \epsilon_{22})^2$, respectively,

$$\begin{aligned}\alpha \geq 0 &\Rightarrow \mathcal{X}_{\{(1,1),(2,2)\}} = \{\mathbf{v}_+\}, \\ \alpha \leq 0 &\Rightarrow \mathcal{X}_{\{(1,2),(2,1)\}} = \{\mathbf{v}_-\}.\end{aligned}\tag{B16}$$

Thus, both $\mathcal{X}_{\{(1,1),(2,2)\}}$ and $\mathcal{Y}_{\{(1,1),(2,2)\}}$ have \mathbf{v}_+ by Eq. (B15) and (B16) if $\alpha \geq |\beta_+|/\gamma_+$. Similarly, both $\mathcal{X}_{\{(1,2),(2,1)\}}$ and $\mathcal{Y}_{\{(1,2),(2,1)\}}$ have \mathbf{v}_- if $\alpha \leq -|\beta_-|/\gamma_-$. Therefore, (52)(\Leftarrow) is also proved. \square

Proof of (b). Assume $m = n_1 = n_2 = 2$ and $p_{\text{guess}}^{\text{post}} < p_{\text{guess}}^{\text{prior}}$. We can see from Lemma 4 and Eq. (B4) that

$$\begin{aligned}\mathcal{X}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset &\Rightarrow \varphi_{(1,1)}(\mathbf{v}_{11}) < \varphi_{(2,1)}(\mathbf{v}_{11}), \quad \varphi_{(1,1)}(\mathbf{v}_{12}) < \varphi_{(1,2)}(\mathbf{v}_{12}), \\ \mathcal{X}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset &\Rightarrow \varphi_{(1,2)}(\mathbf{v}_{11}) < \varphi_{(2,2)}(\mathbf{v}_{11}), \quad \varphi_{(1,2)}(\mathbf{v}_{22}) < \varphi_{(1,1)}(\mathbf{v}_{22}), \\ \mathcal{X}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \varphi_{(2,1)}(\mathbf{v}_{12}) < \varphi_{(2,2)}(\mathbf{v}_{12}), \quad \varphi_{(2,1)}(\mathbf{v}_{21}) < \varphi_{(1,1)}(\mathbf{v}_{21}), \\ \mathcal{X}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \varphi_{(2,2)}(\mathbf{v}_{21}) < \varphi_{(1,2)}(\mathbf{v}_{21}), \quad \varphi_{(2,2)}(\mathbf{v}_{22}) < \varphi_{(2,1)}(\mathbf{v}_{22}).\end{aligned}\tag{B17}$$

By applying Eq. (B9) to (B17), we have

$$\begin{aligned}\mathcal{X}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset &\Rightarrow \frac{\epsilon_{21} + \lambda_{21} \cos \Phi}{\lambda_{21} + \epsilon_{21}} < \frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \quad \frac{\epsilon_{22} + \lambda_{22} \cos \Phi}{\lambda_{22} + \epsilon_{22}} < \frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \\ \mathcal{X}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset &\Rightarrow \frac{\epsilon_{21} - \lambda_{21} \cos \Phi}{\lambda_{21} - \epsilon_{21}} < \frac{\lambda_{21} + \epsilon_{21}}{\lambda_{22} + \epsilon_{22}}, \quad \frac{\epsilon_{22} + \lambda_{22} \cos \Phi}{\lambda_{22} + \epsilon_{22}} > -\frac{\lambda_{22} - \epsilon_{22}}{\lambda_{21} - \epsilon_{21}}, \\ \mathcal{X}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \frac{\epsilon_{22} - \lambda_{22} \cos \Phi}{\lambda_{22} - \epsilon_{22}} < \frac{\lambda_{22} + \epsilon_{22}}{\lambda_{21} + \epsilon_{21}}, \quad \frac{\epsilon_{21} + \lambda_{21} \cos \Phi}{\lambda_{21} + \epsilon_{21}} > -\frac{\lambda_{21} - \epsilon_{21}}{\lambda_{22} - \epsilon_{22}}, \\ \mathcal{X}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \frac{\epsilon_{21} - \lambda_{21} \cos \Phi}{\lambda_{21} - \epsilon_{21}} > -\frac{\lambda_{21} + \epsilon_{21}}{\lambda_{22} + \epsilon_{22}}, \quad \frac{\epsilon_{22} - \lambda_{22} \cos \Phi}{\lambda_{22} - \epsilon_{22}} > -\frac{\lambda_{22} + \epsilon_{22}}{\lambda_{21} + \epsilon_{21}}.\end{aligned}\tag{B18}$$

Since the inequalities given in (B18) can also be expressed as

$$\begin{aligned}\frac{\epsilon_{22} \mp \lambda_{22} \cos \Phi}{\lambda_{22} \mp \epsilon_{22}} < \frac{\lambda_{22} \pm \epsilon_{22}}{\lambda_{21} \pm \epsilon_{21}} &\Leftrightarrow \Theta_{\pm} > \Xi_{\pm}, \\ \frac{\epsilon_{21} \mp \lambda_{21} \cos \Phi}{\lambda_{21} \mp \epsilon_{21}} < \frac{\lambda_{21} \pm \epsilon_{21}}{\lambda_{22} \pm \epsilon_{22}} &\Leftrightarrow \mp \Theta_{\pm} < \Phi \mp \Xi_{\pm} < \pi \mp \pi \pm \Theta_{\pm}, \\ \frac{\epsilon_{22} \pm \lambda_{22} \cos \Phi}{\lambda_{22} \pm \epsilon_{22}} > -\frac{\lambda_{22} \mp \epsilon_{22}}{\lambda_{21} \mp \epsilon_{21}} &\Leftrightarrow \Theta_{\pm} < \pi - \Xi_{\pm}, \\ \frac{\epsilon_{21} \pm \lambda_{21} \cos \Phi}{\lambda_{21} \pm \epsilon_{21}} > -\frac{\lambda_{21} \mp \epsilon_{21}}{\lambda_{22} \mp \epsilon_{22}} &\Leftrightarrow \mp \pi \pm \Theta_{\pm} < \Phi \mp \Xi_{\pm} < \pi \mp \Theta_{\pm},\end{aligned}\tag{B19}$$

a lengthy calculation show that

$$\begin{aligned}\mathbf{v} \in \mathcal{X}_{\{(1,1),(1,2),(2,1)\}} &\Rightarrow \begin{cases} \angle \mathbf{v} \tilde{\boldsymbol{\mu}}_{(1,1)} \tilde{\boldsymbol{\mu}}_{(2,1)} &= \Theta_- - \Xi_-, \\ \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}\| &= \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21} \cos(\Theta_- - \Xi_-) + \epsilon_{21}]}, \\ \varphi_{(1,1)}(\mathbf{v}) - \varphi_{(2,2)}(\mathbf{v}) &= \frac{-2\alpha}{\|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}\| + \|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}\| + \epsilon_{21} + \epsilon_{22}}, \end{cases} \\ \mathbf{v} \in \mathcal{X}_{\{(1,1),(1,2),(2,2)\}} &\Rightarrow \begin{cases} \angle \mathbf{v} \tilde{\boldsymbol{\mu}}_{(1,2)} \tilde{\boldsymbol{\mu}}_{(2,2)} &= \pi - \Theta_+ - \Xi_+, \\ \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}\| &= \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21} \cos(\pi - \Theta_+ - \Xi_+) + \epsilon_{21}]}, \\ \varphi_{(1,2)}(\mathbf{v}) - \varphi_{(2,1)}(\mathbf{v}) &= \frac{+2\alpha}{\|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}\| + \|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}\| + \epsilon_{21} - \epsilon_{22}}, \end{cases} \\ \mathbf{v} \in \mathcal{X}_{\{(1,1),(2,1),(2,2)\}} &\Rightarrow \begin{cases} \angle \mathbf{v} \tilde{\boldsymbol{\mu}}_{(2,1)} \tilde{\boldsymbol{\mu}}_{(1,1)} &= \Theta_+ - \Xi_+, \\ \|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}\| &= \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21} \cos(\Theta_+ - \Xi_+) - \epsilon_{21}]}, \\ \varphi_{(2,1)}(\mathbf{v}) - \varphi_{(1,2)}(\mathbf{v}) &= \frac{+2\alpha}{\|\tilde{\boldsymbol{\mu}}_{(2,1)} - \mathbf{v}\| + \|\tilde{\boldsymbol{\mu}}_{(1,2)} - \mathbf{v}\| - \epsilon_{21} + \epsilon_{22}}, \end{cases} \\ \mathbf{v} \in \mathcal{X}_{\{(1,2),(2,1),(2,2)\}} &\Rightarrow \begin{cases} \angle \mathbf{v} \tilde{\boldsymbol{\mu}}_{(2,2)} \tilde{\boldsymbol{\mu}}_{(1,2)} &= \pi - \Theta_- - \Xi_-, \\ \|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}\| &= \frac{\lambda_{21}^2 - \epsilon_{21}^2}{2[\lambda_{21} \cos(\pi - \Theta_- - \Xi_-) - \epsilon_{21}]}, \\ \varphi_{(2,2)}(\mathbf{v}) - \varphi_{(1,1)}(\mathbf{v}) &= \frac{-2\alpha}{\|\tilde{\boldsymbol{\mu}}_{(2,2)} - \mathbf{v}\| + \|\tilde{\boldsymbol{\mu}}_{(1,1)} - \mathbf{v}\| - \epsilon_{21} - \epsilon_{22}}.\end{cases}\end{aligned}\tag{B20}$$

Here, $\angle \mathbf{v} \tilde{\boldsymbol{\mu}}_{\omega} \tilde{\boldsymbol{\mu}}_{\omega'}$ is the internal angle between two line segments formed by $\{\mathbf{v}, \tilde{\boldsymbol{\mu}}_{\omega}\}$ and $\{\tilde{\boldsymbol{\mu}}_{\omega}, \tilde{\boldsymbol{\mu}}_{\omega'}\}$, respectively.

As $\mathcal{X}_S \cap \mathcal{Y}_S = \mathcal{Z}_S$ for all $S \subseteq \Omega$, Eq. (56) is true from Lemma 3 and Eq. (B20). Moreover, Eq. (B20) also implies

$$\begin{aligned}\mathcal{Z}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset &\Leftrightarrow \mathcal{X}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset, \quad \alpha \leq 0, \\ \mathcal{Z}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset &\Leftrightarrow \mathcal{X}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset, \quad \alpha \geq 0, \\ \mathcal{Z}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset &\Leftrightarrow \mathcal{X}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset, \quad \alpha \geq 0, \\ \mathcal{Z}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset &\Leftrightarrow \mathcal{X}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset, \quad \alpha \leq 0,\end{aligned}\tag{B21}$$

Now, (B16) and (B21) lead us to

$$\begin{aligned} \mathcal{Z}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset \text{ or } \mathcal{Z}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \mathcal{X}_{\{(1,2),(2,1)\}} = \{\mathbf{v}_-\}, \\ \mathcal{Z}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset \text{ or } \mathcal{Z}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \mathcal{X}_{\{(1,2),(2,1)\}} = \{\mathbf{v}_+\}. \end{aligned} \quad (\text{B22})$$

Thus, we can show from Eq. (B15), (B22), and Lemma 4 that

$$\begin{aligned} \mathcal{Z}_{\{(1,1),(1,2),(2,1)\}} \neq \emptyset &\Rightarrow \alpha > -\beta_-/\gamma_-, \\ \mathcal{Z}_{\{(1,1),(1,2),(2,2)\}} \neq \emptyset &\Rightarrow \alpha < -\beta_+/\gamma_+, \\ \mathcal{Z}_{\{(1,1),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \alpha < \beta_+/\gamma_+, \\ \mathcal{Z}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset &\Rightarrow \alpha > \beta_-/\gamma_-. \end{aligned} \quad (\text{B23})$$

Up to this point, (55)(\Rightarrow) for $S \neq \{(1,2), (2,1), (2,2)\}$ is proved by (B18), (B21), and (B23).

Since $\beta_-/\gamma_- < \alpha \leq 0$ is a necessary condition for $\mathcal{Z}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset$ from (B21) and (B23), $\mathcal{Z}_{\{(1,2),(2,1),(2,2)\}} \neq \emptyset$ requires non-positivity of α and β_- . However, both α and β_- cannot be non-positive because

$$\alpha(\epsilon_{21} + \epsilon_{22}) + \beta_- = \epsilon_{21}(\lambda_{22}^2 - \epsilon_{22}^2) + \epsilon_{22}(\lambda_{21}^2 - \epsilon_{21}^2) > 0. \quad (\text{B24})$$

Thus, $\mathcal{Z}_{\{(1,2),(2,1),(2,2)\}}$ is always empty.

Now, for the proof of (55)(\Leftarrow), let S be a proper subset of Ω satisfying the necessary condition of (55). Then, \mathcal{X}_S is nonempty for all $S' \subsetneq S$ with $|S'| = 2$ from (B4) and (B16). Thus, we can see from Lemma 5 and (B21) that \mathcal{Z}_S is nonempty if \mathcal{X}_S^S is empty for all $S' \subsetneq S$ with $|S'| = 2$. Using (B9) and (B15), we can easily show that $\mathcal{X}_S^S = \emptyset$ for all $S' \subsetneq S$ with $|S'| = 2$. Therefore, $\mathcal{Z}_S \neq \emptyset$. That is, (55)(\Leftarrow) is proved. \square

Proof of (c). Let us assume $\mathcal{Z} \subseteq \mathcal{P}_\Omega$. Since a null MEPI measurement exists from Corollary 1, $\mathcal{Z}_S \neq \emptyset$ for some S in (51) with $S \neq \Omega$. When \mathbf{v} is the single element in \mathcal{Z}_S , we can see from the definition of \mathcal{Z}_Ω that $\mathcal{Z}_\Omega \neq \emptyset$ if and only if $\varphi_\omega(\mathbf{v}) = \varphi_{\omega'}(\mathbf{v})$ for all $(\omega, \omega') \in S \times (\Omega - S)$. Thus, when $|S| = 3$, we can show from (B20) that $\mathcal{Z}_\Omega \neq \emptyset$ is equivalent to $\alpha = 0$.

When $|S| = 2$, we can verify from Eqs. (B11) and (B15) that $\mathcal{Z}_\Omega \neq \emptyset$ is equivalent to $\alpha = 0$. Note that $\alpha = 0$ means $\beta_+ = 0$ when $S = \{(1,1), (2,2)\}$ because $\alpha \geq |\beta_+|/\gamma_+$ is a necessary condition for $\mathcal{Z}_{\{(1,1),(2,2)\}} \neq \emptyset$. We also note that $\alpha = 0$ implies $\beta_- = 0$ when $S = \{(1,2), (2,1)\}$ because $\alpha \leq -|\beta_-|/\gamma_-$ is necessary for $\mathcal{Z}_{\{(1,2),(2,1)\}} \neq \emptyset$, therefore $\alpha \neq 0$ and $\mathcal{Z}_\Omega = \emptyset$ when $S = \{(1,2), (2,1)\}$ since both α and β_- cannot be zero by Inequality (B24). Thus, $\alpha = 0$ is a necessary and sufficient condition for $\mathcal{Z}_\Omega \neq \emptyset$ when $\mathcal{Z} \subseteq \mathcal{P}_\Omega$. This completes our proof of Theorem 2(c). \square

-
- [1] Chefles, A.: Quantum state discrimination. *Contemp. Phys.* **41**, 401 (2000)
 - [2] Barnett, S. M., Croke, S.: Quantum state discrimination. *Adv. Opt. Photon.* **1**, 238 (2009)
 - [3] Bergou, J. A.: Discrimination of quantum states. *J. Mod. Opt.* **57**, 160 (2010)
 - [4] Bae, J., Kwek, L.-C.: Quantum state discrimination and its applications. *J. Phys. A: Math. Theor.* **48**, 083001 (2015)
 - [5] Helstrom, C. W.: *Quantum Detection and Estimation Theory*. Academic Press, New York (1976)
 - [6] Ivanovic, I. D.: How to differentiate between non-orthogonal states. *Phys. Lett. A* **123**, 257 (1987)
 - [7] Dieks, D.: Overlap and distinguishability of quantum states. *Phys. Lett. A* **126**, 303 (1988)
 - [8] Peres, A.: How to differentiate between non-orthogonal states. *Phys. Lett. A* **128**, 19 (1988)
 - [9] Jaeger, G., Shimony, A.: Optimal distinction between two non-orthogonal quantum states. *Phys. Lett. A* **197**, 83 (1995)
 - [10] Ha, D., Kwon, Y.: Analysis of optimal unambiguous discrimination of three pure quantum states. *Phys. Rev. A* **91**, 062312 (2015)
 - [11] Croke, S., Andersson, E., Barnett, S. M., Gilson, C. R., Jeffers, J.: Maximum Confidence Quantum Measurements. *Phys. Rev. Lett.* **96**, 070401 (2006)
 - [12] Holevo, A. S.: *Probabilistic and Statistical Aspects of Quantum Theory*. North-Holland, Amsterdam (1979)
 - [13] Yuen, H. P., Kennedy, R. S., Lax, M.: Optimum Testing of Multiple Hypotheses in Quantum Detection Theory. *IEEE Trans. Inf. Theory* **21**, 125 (1975)
 - [14] Eldar, Y. C., Megretski, A., Verghese, G. C.: Designing Optimal Quantum Detectors Via Semidefinite Programming. *IEEE Trans. Inf. Theory*, **49**, 1007 (2003)
 - [15] Barnett, S. M., Croke, S.: On the conditions for discrimination between quantum states with minimum error. *J. Phys. A: Math. Theor.* **42**, 062001 (2009)
 - [16] Ban, M., Kurokawa, K., Momose, R., Hirota, O.: Optimum Measurements for Discrimination Among Symmetric Quantum States and Parameter Estimation. *Int. J. Theor. Phys.* **36**, 1269 (1997)
 - [17] Chou, C. L., Hsu, L. Y.: Minimum-error discrimination between symmetric mixed quantum states. *Phys. Rev. A* **68**, 042305 (2003)

- [18] Hunter, K.: Results in Optimal Discrimination. AIP Conf. Proc. **734**, 83 (2004)
- [19] Samsonov, B. F.: Minimum error discrimination problem for pure qubit states. Phys. Rev. A **80**, 052305 (2009)
- [20] Jafarizadeh, M. A., Mazhari, Y., Aali, M.: The minimum-error discrimination via Helstrom family of ensembles and convex optimization. Quantum Inf. Process. **10**, 155-176 (2011)
- [21] Ha, D., Kwon, Y.: Complete analysis for three-qubit mixed-state discrimination. Phys. Rev. A **87**, 062302 (2013)
- [22] Ha, D., Kwon, Y.: Discriminating N -qudit states using geometric structure. Phys. Rev. A **90**, 022330 (2014)
- [23] Bae, J., Hwang, W.-Y.: Minimum-error discrimination of qubit states: Methods, solutions, and properties. Phys. Rev. A **87**, 012334 (2013)
- [24] Bae, J.: Structure of minimum-error quantum state discrimination. New J. Phys. **15**, 073037 (2013)
- [25] Hunter, K.: Measurement does not always aid state discrimination. Phys. Rev. A **68**, 012306 (2003)
- [26] Akibue, S., Kato, G., Marumo, N.: Perfect discrimination of nonorthogonal quantum states with posterior classical partial information. Phys. Rev. A **99**, 020102(R) (2019)
- [27] Ballester, M. A., Wehner, S., Winter, A.: State Discrimination With Post-Measurement Information. IEEE Trans. Inf. Theory, **54**, 4183 (2008)
- [28] Gopal, D., Wehner, S.: Using postmeasurement information in state discrimination. Phys. Rev. A **82**, 022326 (2010)
- [29] Carmeli, C., Heinosaari, T., Toigo, A.: State discrimination with postmeasurement information and incompatibility of quantum measurements. Phys. Rev. A **98**, 012126 (2018)
- [30] Heinosaari, T., Miyadera, T., Ziman, M.: An invitation to quantum incompatibility. J. Phys. A: Math. Theor. **49**, 123001 (2016)
- [31] Carmeli, C., Heinosaari, T., Toigo, A.: Quantum Incompatibility Witnesses. Phys. Rev. Lett. **122**, 130402 (2019)
- [32] Skrzypczyk, P., Šupić, I., Cavalcanti, D.: All Sets of Incompatible Measurements give an Advantage in Quantum State Discrimination. Phys. Rev. Lett. **122**, 130403 (2019)
- [33] Uola, R., Kraft, T., Shang, J., Yu, X.-D., Gühne, O.: Quantifying Quantum Resources with Conic Programming. Phys. Rev. Lett. **122**, 130404 (2019)
- [34] Skrzypczyk, P., Linden, N.: Robustness of Measurement, Discrimination Games, and Accessible Information. Phys. Rev. Lett. **122**, 140403 (2019)
- [35] Boyd, S., Vandenberghe, L.: Convex Optimization. Cambridge University Press, UK (2004)
- [36] In \mathbb{R}^3 , a set X is *affine* if $px_0 + (1-p)x_1 \in X$ for any $x_0, x_1 \in X$ and $p \in \mathbb{R}$. If X is an affine set, there exist an element $x \in X$ and a subspace $V \subseteq \mathbb{R}^3$ such that $X = V + x$. Then, the dimension of X is defined as the dimension of V . Given a set X , the *affine dimension* is the dimension of the affine hull $\{px_0 + (1-p)x_1 \mid x_0, x_1 \in X \text{ and } p \in \mathbb{R}\}$. When D is the affine dimension of X , the affine hull is a point, line, plane, \mathbb{R}^3 according to $D = 0, 1, 2, 3$.
- [37] Davies, E. B.: Information and Quantum Measurement. IEEE Trans. Inf. Theory **24**, 596 (1978)
- [38] Chitambar, E., Gour, G.: Quantum resource theories. Rev. Mod. Phys. **91**, 025001 (2019)