

The groups and nilpotent Lie rings of order p^8 with maximal class

Seungjai Lee and Michael Vaughan-Lee

September 2021

Abstract

We classify the nilpotent Lie rings of order p^8 with maximal class for $p \geq 5$. This also provides a classification of the groups of order p^8 with maximal class for $p \geq 11$ via the Lazard correspondence.

1 Introduction

We give a classification of the nilpotent Lie rings of order p^8 ($p \geq 5$) which are of maximal class (i.e. nilpotent of class 7). The classification gives us the following theorem.

Theorem 1 *For $p \geq 5$ the number of nilpotent Lie rings of order p^8 which have maximal class is*

$$4p^3 + 7p^2 + 9p + 6 + (6p + 11) \gcd(p - 1, 3) + 4 \gcd(p - 1, 5) \\ + (p + 2) \gcd(p - 1, 7) + (p + 3) \gcd(p - 1, 8) + 2 \gcd(p - 1, 9) + \gcd(p - 1, 12).$$

By the Lazard correspondence between p -groups and nilpotent Lie rings, for $p \geq 11$ this formula also gives us the number of groups of order p^8 with maximal class.

We have constructed a database of the nilpotent Lie rings of order p^8 with maximal class which will be included in the next release of the **GAP** package LieP Ring [3]. The LieP Ring package can then be used to provide complete lists of the nilpotent Lie rings of order p^8 with maximal class for any given $p \geq 5$, and can also be used to provide complete lists of the groups of order p^8 with maximal class for any given $p \geq 11$. Both **GAP** [4] and **MAGMA** [1] have databases of the groups of order 2^8 and 3^8 , and it is easy to extract the groups of maximal class from these databases. For $p = 5, 7$ we can use the databases of groups of order p^7 to obtain a

list of the groups of maximal class, and then use the Descendants function in GAP or MAGMA to compute the groups of order p^8 with maximal class. (If P has order p^8 and maximal class then P is a “descendant” of the quotient P/Z where Z is the centre of P , and P/Z is a group of order p^7 of maximal class.)

2 Preliminaries

Let L be a nilpotent Lie ring of order p^n ($n \geq 3$) and class $n - 1$. Here “class” means nilpotency class, and not p -class, which is the length of lower p -central series of L . So L is a nilpotent Lie ring of maximal class. Let the lower central series of L be

$$L > L^2 > L^3 > \dots > L^{n-1} > L^n = \{0\},$$

where $L^2 = \langle ab \mid a, b \in L \rangle$, and where for $i > 2$, $L^i = \langle ab \mid a \in L^{i-1}, b \in L \rangle$. (We denote the Lie product of a and b by ab , rather than by $[a, b]$.) Then L/L^2 is elementary abelian of order p^2 , and for $2 \leq i < n$ the quotient L^i/L^{i+1} has order p . Note that if $n > 3$ then L/L^{n-1} has maximal class $n - 2$.

Lemma 2 $pL \leq L^{n-1}$.

Proof. The proof is by induction on n . Note that there is nothing to prove if $n = 3$. So assume that $n > 3$, and assume by induction that $pL \leq L^{n-2}$. Let L be generated by a, b , so that we can assume that $pa, pb \in L_{n-2}$. If $pa \notin L^{n-1}$ then a centralizes L^{n-2} . Similarly if $pb \notin L^{n-1}$ then b centralizes L^{n-2} , and if $pa + pb \notin L^{n-1}$ then $a + b$ centralizes L^{n-2} . Now if one or the other or both of pa, pb lie outside L^{n-1} then at least two of $pa, pb, pa + pb$ lie outside L^{n-1} , which implies that L is generated by elements which centralize L^{n-2} . Clearly this is impossible, and so $pa, pb \in L^{n-1}$. \square

Now let L be a nilpotent Lie ring of order p^8 and maximal class 7. Then L/L^7 is a nilpotent Lie ring of order p^7 with maximal class. By Lemma 2, L/L^7 has characteristic p . The GAP package LiePRing contains a database of the nilpotent Lie rings of order p^7 ($p \geq 5$). There are $p + 8$ Lie rings of maximal class and characteristic p in the database. These are as follows. (Here ba^3 denotes $baaa$, ba^4 denotes $baaaa$, and so on.)

$$\langle a, b \mid bab, ba^3b, pa, pb, \text{ class } 6 \rangle, \quad (7.623)$$

$$\langle a, b \mid bab, ba^3b - ba^5, pa, pb, \text{ class } 6 \rangle, \quad (7.627)$$

$$\langle a, b \mid bab - ba^5, ba^3b, pa, pb, \text{ class } 6 \rangle, \quad (7.633)$$

$$\langle a, b \mid bab - ba^4, ba^3b, pa, pb, \text{ class } 6 \rangle, \quad (7.641)$$

$$\langle a, b \mid bab - ba^4, ba^3b - ba^5, pa, pb, \text{ class } 6 \rangle, \quad (7.646)$$

$$\langle a, b \mid bab - ba^4 - ba^5, ba^3b, pa, pb, \text{ class } 6 \rangle, \quad (7.648)$$

$$\langle a, b \mid bab - ba^3, ba^3b - xba^5, pa, pb, \text{ class } 6 \rangle \quad (0 \leq x < p), \quad (7.650)$$

$$\langle a, b \mid bab - ba^3 - ba^5, ba^3b - ba^5, pa, pb, \text{ class } 6 \rangle, \quad (7.656)$$

$$\langle a, b \mid bab - ba^3 - wba^5, ba^3b - ba^5, pa, pb, \text{ class } 6 \rangle. \quad (7.657)$$

The numbering 7.623, 7.627, \dots gives the “LibraryName” of these Lie rings in the database. The parameter w in 7.657 is taken to be a (fixed) primitive element mod p . Note that p in these presentations can be replaced by 5 to give a complete list of the nilpotent Lie rings of order 5^7 of maximal class and characteristic 5, replaced by 7 to give a complete list of the nilpotent Lie rings of order 7^7 of maximal class and characteristic 7, and so on.

3 Computing descendants

We use the *Lie ring generation algorithm* as described in [5] and [7] to compute the descendants of order p^8 of the nilpotent Lie rings of order p^7 with maximal class. This algorithm is an analogue of the p -group generation algorithm described in [6]. The Lie ring generation algorithm makes use of the lower p -central series of a Lie ring L , which is defined in an analogous way to groups. We define the series

$$L = L_1 \geq L_2 \geq L_3 \geq \dots \geq L_c \geq \dots$$

by setting $L_1 = L$, $L_2 = L^2 + pL$, and for $c > 1$ we set $L_{c+1} = L_cL + pL_c$. (Here L_cL is $\langle ab \mid a \in L_c, b \in L \rangle$.) Note that we use superscripts to denote terms of the lower central series, and subscripts to denote terms of the lower p -central series. In the case of a nilpotent Lie ring of order p^n with maximal class the two series are identical. The ideal L_c consists of all linear combinations of terms of the form

$$a_1a_2 \dots a_c, pa_1a_2 \dots a_{c-1}, p^2a_1a_2 \dots a_{c-2}, \dots, p^{c-1}a_1.$$

We say that L has p -class c if $L_{c+1} = \{0\}$, $L_c \neq \{0\}$.

If L is a nilpotent Lie ring with finite order p^n for some prime p , then L_{c+1} will equal $\{0\}$ for some c . In fact if L is nilpotent of class k , and if the exponent of L as a finite abelian group is p^m then L has p -class c for some c with $k \leq c < k + m$.

If L and M are two finite nilpotent Lie rings with prime-power order, then L is a *descendant* of M if $L/L_c \cong M$ for some $c \geq 2$. If $L/L_c \cong M$ and L has p -class c (so that $L_c \neq \{0\}$, $L_{c+1} = \{0\}$) then L is an *immediate descendant* of M . Note that if L is a descendant of M then $L/L_2 \cong M/M_2$, so that L and M have the same generator number.

If M is a nilpotent d -generator Lie ring of order p^n , then we construct its p -covering ring \widehat{M} . This is the largest d -generator Lie ring \widehat{M} having a central elementary abelian ideal Z such that $\widehat{M}/Z \cong M$ and every immediate descendant

of M is isomorphic to \widehat{M}/T for some $T \leq Z$. However \widehat{M}/T is not an immediate descendant of M for every subring $T \leq Z$. If M has p -class c (so that $M_{c+1} = \{0\}$) then we define the *nucleus* of M to be \widehat{M}_{c+1} . Then \widehat{M}/T is an immediate descendant of M if and only if T is a proper subring of Z such that T supplements the nucleus \widehat{M}_{c+1} . It can happen that $\widehat{M}_{c+1} = \{0\}$, in which case M has no immediate descendants and is *terminal*.

Hence we obtain a complete list of the immediate descendants of M by calculating its p -covering ring \widehat{M} , and listing the proper subrings $T < Z$ such that $T + \widehat{M}_{c+1} = Z$. (These are the *allowable subrings* of Z .)

We now have a list of the immediate descendants of M , and we can easily restrict to those with a specified order. This list will usually contain redundancies, and we need to solve the isomorphism problem. This is done as follows. We compute the automorphism group of M and we extend each automorphism α of M to an automorphism α^* of \widehat{M} . (If M is generated by a_1, a_2, \dots, a_d then we choose preimages x_1, x_2, \dots, x_d in \widehat{M} for a_1, a_2, \dots, a_d , and preimages y_1, y_2, \dots, y_d in \widehat{M} for $a_1\alpha, a_2\alpha, \dots, a_d\alpha$. Then x_1, x_2, \dots, x_d generate \widehat{M} , and we define α^* by setting $x_i\alpha^* = y_i$ for $i = 1, 2, \dots, d$.) Then $Z\alpha^* = Z$, and the action of α^* on Z is uniquely determined by α . Two allowable subrings T_1, T_2 define isomorphic descendants $\widehat{M}/T_1, \widehat{M}/T_2$ if and only if $T_2\alpha^* = T_1$ for some automorphism α of M . We obtain a complete irredundant set of immediate descendants of M by choosing a set of representatives for the orbits of the allowable subrings of Z under this action of the automorphism group of M .

4 The nilpotent Lie rings of order p^8 with maximal class

In this section we give a complete list of presentations for the nilpotent Lie rings of order p^8 with maximal class ($p \geq 5$). Many of the presentations involve parameters w, x, y, z . These parameters take integer values in the range $0, 1, \dots, p-1$. The parameter w is always assumed to be a (fixed) primitive element mod p . Associated with the presentations are somewhat cryptic comments intended to describe when two sets of parameters give isomorphic Lie rings. For example four of the descendants of 7.623 have a single parameter x , together with the comment “ $x \neq 0$, $x \sim xa^6$ ”. Here (and for all these comments) a is assumed to range over all integers which are non-zero modulo p . So this comment is intended to mean that x can take any value in the range $1, 2, \dots, p-1$ and that if $0 < x, y < p$ then x and y give isomorphic Lie rings if and only if $x = ya^6 \pmod{p}$ for some integer a which is not divisible by p . Actually there is no reason to restrict x to the range $1, 2, \dots, p-1$ since x is the coefficient of an element of order p in the presentations of these Lie rings. So if $x = x' \pmod{p}$ then x and x' give identical Lie rings.

The simplest way to “solve” these conditions is to treat them as defining equivalence relations over $\text{GF}(p)$. Two non-zero elements in $\text{GF}(p)$ give isomorphic algebras if and only if they lie in the same coset of the subgroup $\{a^6 \mid a \in \text{GF}(p)^*\}$ of the multiplicative group $\text{GF}(p)^*$ of non-zero elements in $\text{GF}(p)$. This subgroup has order $\frac{p-1}{6}$ if $\gcd(p-1, 3) = 3$ and order $\frac{p-1}{2}$ if $\gcd(p-1, 3) = 1$. If we let u be a primitive element in $\text{GF}(p)$ then $1, u, u^2, u^3, u^4, u^5$ is a transversal for this subgroup when $\gcd(p-1, 3) = 3$, and $1, u$ is a transversal for the subgroup if $\gcd(p-1, 3) = 1$. So we obtain a complete and irredundant set of representatives for the isomorphism classes of these Lie rings as x ranges over $1, 2, \dots, p-1$ by taking $x = 1, w, w^2 \bmod p, w^3 \bmod p, w^4 \bmod p, w^5 \bmod p$ when $\gcd(p-1, 3) = 3$ and taking $x = 1, w$ when $\gcd(p-1, 3) = 1$.

The comments associated with the other Lie rings with a single parameter x are similar. One of the descendants of 7.627 has two parameters x, y with the comment “ $x, y \neq 0, [x, y] \sim [xa^7, ya^6]$ ”. We can solve this over $\text{GF}(p)$ by letting y range over a transversal for the subgroup $\{a^6 \mid a \in \text{GF}(p)^*\}$ of the group $\text{GF}(p)^*$. For any given y in this transversal the pair $[x, y]$ gives an isomorphic Lie ring to the pair $[x', y]$ if and only if $x' = xa$ for some $a \in \text{GF}(p)^*$ satisfying $a^6 = 1$. We can solve this equivalence relation on the values for x over $\text{GF}(p)$, and lift the a set of representatives for the equivalence classes to integers in the range $1, 2, \dots, p-1$.

All the Lie rings in the list below have nilpotency class 7, but we leave the class unspecified, to save space.

4.1 The descendants of 7.623

7.623 has $8 + 8 \gcd(p-1, 3) + 2 \gcd(p-1, 5) + \gcd(p-1, 8)$ descendants of order p^8 .

$$\langle a, b \mid bab, ba^3b, pa, pb, ba^5b \rangle$$

$$\langle a, b \mid bab, ba^3b, pa - ba^6, pb, ba^5b \rangle$$

$$\langle a, b \mid bab, ba^3b, pa, pb - xba^6, ba^5b \rangle \ (x \neq 0, x \sim xa^6)$$

$$\langle a, b \mid bab, ba^3b - ba^6, pa, pb, ba^5b \rangle$$

$$\langle a, b \mid bab, ba^3b - ba^6, pa - xba^6, pb, ba^5b \rangle \ (x \neq 0, x \sim xa^8)$$

$$\langle a, b \mid bab, ba^3b - ba^6, pa, pb - xba^6, ba^5b \rangle \ (x \neq 0, x \sim xa^6)$$

$$\langle a, b \mid bab - ba^6, ba^3b, pa, pb, ba^5b \rangle$$

$$\langle a, b \mid bab - ba^6, ba^3b, pa - xba^6, pb, ba^5b \rangle \ (x \neq 0, x \sim xa^{10})$$

$$\langle a, b \mid bab - ba^6, ba^3b, pa, pb - xba^6, ba^5b \rangle \ (x \neq 0, x \sim xa^6)$$

$$\langle a, b \mid bab, ba^3b, pa, pb, ba^6 \rangle$$

$$\langle a, b \mid bab, ba^3b, pa - ba^5b, pb, ba^6 \rangle$$

$$\begin{aligned}
&\langle a, b \mid bab, ba^3b, pa - wba^5b, pb, ba^6 \rangle \\
&\langle a, b \mid bab, ba^3b, pa, pb - ba^5b, ba^6 \rangle \\
&\langle a, b \mid bab, ba^3b, pa - xba^5b, pb - ba^5b, ba^6 \rangle \ (x \neq 0, x \sim xa^6)
\end{aligned}$$

4.2 Descendants of 7.627

7.627 has $p + 2 \gcd(p-1, 3) + \gcd(p-1, 7)$ descendants of order p^8 .

$$\begin{aligned}
&\langle a, b \mid bab, ba^3b - ba^5, pa, pb \rangle \\
&\langle a, b \mid bab, ba^3b - ba^5, pa - xba^6, pb \rangle \ (x \neq 0, x \sim xa^7) \\
&\langle a, b \mid bab, ba^3b - ba^5, pa, pb - xba^6 \rangle \ (x \neq 0, x \sim xa^6) \\
&\langle a, b \mid bab, ba^3b - ba^5, pa - xba^6, pb - yba^6 \rangle \ (x, y \neq 0, [x, y] \sim [xa^7, ya^6])
\end{aligned}$$

4.3 Descendants of 7.633

7.633 has $4p + (p+1) \gcd(p-1, 3) + 2 \gcd(p-1, 9) + \gcd(p-1, 12)$ descendants of order p^8 .

$$\begin{aligned}
&\langle a, b \mid bab - ba^5, ba^3b, pa, pb, ba^5b \rangle \\
&\langle a, b \mid bab - ba^5, ba^3b, pa, pb - xba^6, ba^5b \rangle \ (x \neq 0, x \sim xa^6) \\
&\langle a, b \mid bab - ba^5, ba^3b, pa - xba^6, pb, ba^5b \rangle \ (x \neq 0, x \sim xa^9) \\
&\langle a, b \mid bab - ba^5 - ba^6, ba^3b, pa, pb, ba^5b \rangle \\
&\langle a, b \mid bab - ba^5 - ba^6, ba^3b, pa - xba^6, pb, ba^5b \rangle \ (x \neq 0) \\
&\langle a, b \mid bab - ba^5 - ba^6, ba^3b, pa, pb - xba^6, ba^5b \rangle \ (x \neq 0) \\
&\langle a, b \mid bab - ba^5, ba^3b - ba^6, pa, pb, ba^5b \rangle \\
&\langle a, b \mid bab - ba^5, ba^3b - ba^6, pa - xba^6, pb, ba^5b \rangle \ (x \neq 0) \\
&\langle a, b \mid bab - ba^5, ba^3b - ba^6, pa, pb - xba^6, ba^5b \rangle \ (x \neq 0) \\
&\langle a, b \mid bab - ba^5, ba^3b, pa, pb, ba^6 \rangle \\
&\langle a, b \mid bab - ba^5, ba^3b, pa - xba^5b, pb, ba^6 \rangle \ (x \neq 0, x \sim xa^{12}) \\
&\langle a, b \mid bab - ba^5, ba^3b, pa, pb - xba^5b, ba^6 \rangle \ (x \neq 0, x \sim xa^9) \\
&\langle a, b \mid bab - ba^5, ba^3b, pa - xba^5b, pb - yba^5b, ba^6 \rangle \ (x, y \neq 0, [x, y] \sim [xa^{12}, ya^9])
\end{aligned}$$

4.4 Descendants of 7.641

7.641 has $5p - 3 + 2p \gcd(p - 1, 3) + 2 \gcd(p - 1, 5) + (p + 1) \gcd(p - 1, 8)$ descendants of order p^8 .

$$\begin{aligned}
& \langle a, b \mid bab - ba^4, ba^3b - xba^6, pa, pb, ba^5b \rangle \text{ (all } x) \\
& \langle a, b \mid bab - ba^4, ba^3b - xba^6, pa - yba^6, pb, ba^5b \rangle \text{ (} y \neq 0, [x, y] \sim [x, ya^8] \text{)} \\
& \langle a, b \mid bab - ba^4, ba^3b - xba^6, pa, pb - yba^6, ba^5b \rangle \text{ (} y \neq 0, [x, y] \sim [x, ya^6] \text{)} \\
& \langle a, b \mid bab - ba^4, ba^3b - ba^6, pa - xba^6, pb - yba^6, ba^5b \rangle \text{ (} x, y \neq 0, [x, y] \sim [xa^8, ya^6] \text{)} \\
& \langle a, b \mid bab - ba^4, ba^3b, pa, pb, ba^6 \rangle \\
& \langle a, b \mid bab - ba^4, ba^3b, pa - xba^5b, pb, ba^6 \rangle \text{ (} x \neq 0, x \sim xa^{10} \text{)} \\
& \langle a, b \mid bab - ba^4, ba^3b, pa, pb - xba^5b, ba^6 \rangle \text{ (} x \neq 0, x \sim xa^8 \text{)} \\
& \langle a, b \mid bab - ba^4, ba^3b, pa - xba^5b, pb - yba^5b, ba^6 \rangle \text{ (} x, y \neq 0, [x, y] \sim [xa^{10}, ya^8] \text{)}
\end{aligned}$$

4.5 Descendants of 7.646

7.646 has p^2 descendants of order p^8 .

$$\langle a, b \mid bab - ba^4, ba^3b - ba^5, pa - xba^6, pb - yba^6 \rangle \text{ (all } x, y)$$

4.6 Descendants of 7.648

7.648 has $4p^2 - 3p + 1$ descendants of order p^8 .

$$\begin{aligned}
& \langle a, b \mid bab - ba^4 - ba^5, ba^3b - xba^6, pa, pb, ba^5b \rangle \text{ (} x \neq 1 \text{)} \\
& \langle a, b \mid bab - ba^4 - ba^5, ba^3b - xba^6, pa - yba^6, pb, ba^5b \rangle \text{ (} x \neq 1, y \neq 0 \text{)} \\
& \langle a, b \mid bab - ba^4 - ba^5, ba^3b - xba^6, pa, pb - yba^6, ba^5b \rangle \text{ (} x \neq 1, y \neq 0 \text{)} \\
& \langle a, b \mid bab - ba^4 - ba^5, ba^3b - ba^6, pa - xba^6, pb - yba^6, ba^5b \rangle \text{ (all } x, y) \\
& \langle a, b \mid bab - ba^4 - ba^5, ba^3b, pa - xba^5b, pb - yba^5b, ba^6 \rangle \text{ (all } x, y)
\end{aligned}$$

4.7 Descendants of 7.650

7.650 is a family of p Lie rings, and between them they have

$$2p^3 + 3p^2 + p + 3p \gcd(p-1, 3) + (p+1) \gcd(p-1, 7) + \gcd(p-1, 8)$$

descendants of order p^8 .

$$\begin{aligned} &\langle a, b \mid bab - ba^3, ba^3b - ba^5, ba^5b, pa, pb \rangle \\ &\langle a, b \mid bab - ba^3, ba^3b - ba^5, ba^5b, pa - xba^6, pb \rangle (x \neq 0, x \sim xa^7) \\ &\langle a, b \mid bab - ba^3, ba^3b - ba^5, ba^5b, pa, pb - xba^6 \rangle (x \neq 0, x \sim xa^6) \\ &\langle a, b \mid bab - ba^3, ba^3b - ba^5, ba^5b, pa - xba^6, pb - yba^6 \rangle (x, y \neq 0, [x, y] \sim [xa^7, ya^6]) \\ &\langle a, b \mid bab - ba^3, ba^3b - ba^5 - ba^6, ba^5b, pa - xba^6, pb - yba^6 \rangle (\text{all } x, y) \\ &\langle a, b \mid bab - ba^3 - xba^6, ba^3b - ba^5, ba^5b, pa - yba^6, pb - zba^6 \rangle (x \neq 0, [x, y, z] \sim [xa^3, ya^7, za^6]) \\ &\langle a, b \mid bab - ba^3, ba^3b - xba^5, pa, pb, ba^5b \rangle (x \neq 1) \\ &\langle a, b \mid bab - ba^3, ba^3b - xba^5, pa - yba^6, pb, ba^5b \rangle (x \neq 1, y \neq 0, [x, y] \sim [x, ya^7]) \\ &\langle a, b \mid bab - ba^3, ba^3b - xba^5, pa, pb - yba^6, ba^5b \rangle (x \neq 1, y \neq 0, [x, y] \sim [x, ya^6]) \\ &\langle a, b \mid bab - ba^3, ba^3b - xba^5, pa - yba^6, pb - zba^6, ba^5b \rangle (x \neq 1, y, z \neq 0, [x, y, z] \sim [x, ya^7, za^6]) \\ &\langle a, b \mid bab - ba^3, ba^3b - xba^5 - ba^6, pa - yba^6, pb - zba^6, ba^5b \rangle (x \neq 1, \text{all } y, z) \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5, pa, pb, ba^6 \rangle \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5, pa - xba^5b, pb, ba^6 \rangle (x \neq 0, x \sim xa^8) \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5, pa, pb - xba^5b, ba^6 \rangle (x \neq 0, x \sim xa^7) \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5, pa - xba^5b, pb - yba^5b, ba^6 \rangle (x, y \neq 0, [x, y] \sim [xa^8, ya^7]) \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5 - ba^5b, pa - xba^5b, pb - yba^5b, ba^6 \rangle ([x, y] \sim [x, -y]) \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5 - wba^5b, pa - xba^5b, pb - yba^5b, ba^6 \rangle ([x, y] \sim [x, -y]) \\ &\langle a, b \mid bab - ba^3, ba^3b - 3ba^5 - xba^6, pa - yba^6, pb - zba^6, ba^5b - ba^6 \rangle (\text{all } x, y, z) \end{aligned}$$

4.8 Descendants of 7.656

7.656 has $p^3 - \frac{p^2-p}{2}$ descendants of order p^8 .

$$\begin{aligned} &\langle a, b \mid bab - ba^3 - ba^5, ba^3b - ba^5, ba^5b, pa - xba^6, pb - yba^6 \rangle ([x, y] \sim [-x, y]) \\ &\langle a, b \mid bab - ba^3 - ba^5 - xba^6, ba^3b - ba^5, ba^5b, pa - yba^6, pb - zba^6 \rangle (x \neq 0, [x, y, z] \sim [-x, y, z]) \\ &\langle a, b \mid bab - ba^3 - ba^5, ba^3b - ba^5 - xba^6, ba^5b, pa - yba^6, pb - zba^6 \rangle (x \neq 0, [x, y, z] \sim [-x, y, z]) \end{aligned}$$

4.9 Descendants of 7.657

7.657 has $p^3 - \frac{p^2-p}{2}$ descendants of order p^8 .

$$\langle a, b \mid bab - ba^3 - wba^5, ba^3b - ba^5, ba^5b, pa - xba^6, pb - yba^6 \rangle ([x, y] \sim [-x, y])$$

$$\langle a, b \mid bab - ba^3 - wba^5 - xba^6, ba^3b - ba^5, ba^5b, pa - yba^6, pb - zba^6 \rangle (x \neq 0, [x, y, z] \sim [-x, y, z])$$

$$\langle a, b \mid bab - ba^3 - wba^5, ba^3b - ba^5 - xba^6, ba^5b, pa - yba^6, pb - zba^6 \rangle (x \neq 0, [x, y, z] \sim [-x, y, z])$$

5 Acknowledgements

Our classification of the nilpotent Lie rings of order p^8 with maximal class ($p \geq 5$) is essentially a hand calculation, with some computer assistance with MAGMA [1]. We used Eamonn O'Brien's p -group generation algorithm [6] in MAGMA to compute the groups of order p^8 with maximal class for $11 \leq p \leq 43$, and confirmed that the number of groups in these cases agreed with the PORC formula given in Theorem 1. We also used Serena Cicalò and Willem de Graaf's implementation of the Lazard correspondence in their GAP package LieRing [2] to obtain the groups corresponding to the nilpotent Lie rings in our database, so that we could compare them with the groups provided by the p -group generation algorithm. We used Eamonn O'Brien's StandardPresentation function in MAGMA to prove that the two sets of groups are identical (up to isomorphism) for $11 \leq p \leq 43$.

The first author was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MEST), No. 2019R1A6A1A10073437.

References

- [1] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system I: The user language*, J. Symbolic Comput. **24** (1997), 235–265.
- [2] Serena Cicalò and Willem de Graaf, *LieRing – Computing with finitely presented Lie rings*, (2019), a GAP 4 package.
- [3] B. Eick and M. Vaughan-Lee, *LiePRing – Database and algorithms for Lie p -rings*, (2015), a GAP 4 package.
- [4] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.11*, (2020). Available from <http://www.gap-system.org>.
- [5] M.F. Newman, E.A. O'Brien, and M.R. Vaughan-Lee, *Groups and nilpotent Lie rings whose order is the sixth power of a prime*, J. Algebra **278** (2004), 383–401.

- [6] E.A. O'Brien, *The p -group generation algorithm*, J. Symbolic Comput. **9** (1990), 677–698.
- [7] E.A. O'Brien and M.R. Vaughan-Lee, *The groups with order p^7 for odd prime p* , J. Algebra **292** (2005), 243–358.