THE POSITIVE AND NEGATIVE DEFICIENCY INDICES OF FORMALLY SELF-ADJOINT DIFFERENCE EQUATIONS

GUOJING REN

ABSTRACT. This paper is concerned with formally self-adjoint difference equations and their positive and negative deficiency indices. It is shown that the order of any formally self-adjoint difference equation is even, and some characterizations of formally self-adjoint difference equations are established. Further, we show that the positive and negative deficiency indices are always equal, which implies the existence of the self-adjoint extensions of the minimal linear relations generated by the difference equations. This is an important and essential difference between formally self-adjoint difference equations and their corresponding differential equations in the spectral theory.

1. INTRODUCTION

Difference equations are usually regarded as the discretization of the corresponding differential equations. According to the existing results, most of the properties in spectral theory of difference equations coincide with those of the corresponding differential equations; only a few, but important, are different. It has been found that the maximal operator corresponding to a formally self-adjoint difference equation may be multi-valued, and the corresponding minimal operator may be non-densely defined [18, 20]. Therefore, the classical spectral theory for symmetric operators, i.e., densely defined and Hermitian single-valued operators, are not available in the studying of the spectral properties of difference equations in general. Due to this reason, some researchers focus on extending the spectral theory of linear operators to linear non-densely defined or multi-valued operators (which are called linear relations or linear subspaces), and many good results have been obtained (See [6, 7, 21, 22] and their references).

According to the generalized von Neumann theory and the GKN theory, a Hermitian linear relation, has a self-adjoint extension if and only if its positive and negative deficiency indices are equal, and in this case the domain of the self-adjoint

²⁰⁰⁰ Mathematics Subject Classification. 39A70, 47B39, 34B20.

Key words and phrases. Positive and negative deficiency indices, Hermitian operator, Formally self-adjoint, Difference equation, K-real.

This work is supported by the NSF of Shandong Province, P.R. China [grant numbers ZR2020MA012].

GUOJING REN

extension has a close relationship with the deficiency indices [6, 21]. So, the positive and negative deficiency indices of Hermitian linear relations play a key role in the study of spectral theory of linear relations.

Now, we briefly recall some important results about the deficiency indices of formally self-adjoint differential and difference equations, respectively.

The theory of positive and negative deficiency indices of Hermitian differential equations has been well developed. The canonical form of any formally self-adjoint differential expression τ of order m on some interval \mathcal{J} is given by

(1.1)
$$\tau y := \sum_{j=0}^{[m/2]} (-1)^j [a_j y^{(j)}]^{(j)} + i \sum_{k=0}^{[(m-1)/2]} \left[\left(b_k y^{(k+1)} \right)^{(k)} + \left(b_k y^{(k)} \right)^{(k+1)} \right],$$

where a_j and b_k are all real-valued functions belonging to $C^{\infty}(\mathcal{J})$, and $i = \sqrt{-1}$. The following equation

(1.2)
$$\tau y(x) = \lambda w(x)y(x), \quad x \in \mathcal{J}$$

is called the formally self-adjoint differential equation corresponding to τ [8, XIII,2], where $w \geq 0$ is called the weighted function, $\lambda \in \mathbb{C}$ is the spectral parameter. In the case that both endpoints of \mathcal{J} are singular, we can divided \mathcal{J} into two subintervals such that each subinterval has at least one regular endpoint. So we take $\mathcal{J} := [0, +\infty)$ without loss generality.

Since (1.1) is formally self-adjoint, the minimal linear relation generated by (1.2) in the corresponding Hilbert space is Hermitian, and consequently its deficiency index $d_{\lambda}(\tau)$ is constant when λ is in the upper half-plane and lower half-plane. Denoted $d_{\pm}(\tau) := d_{\pm i}(\tau)$, which are called the positive and negative deficiency indices of (1.2).

In addition, by $n_{\lambda}(\tau)$ denote the number of linearly independent solutions of (1.2) which satisfies

$$\int_{\mathcal{J}} w(x) |y(x)|^2 dx < +\infty.$$

and denote $n_{\pm}(\tau) := n_{\pm i}(\tau)$. The definiteness condition for (1.2) is given by:

 (A_1) there exists a bounded interval $\mathcal{J}_0 \subset \mathcal{J}$ such that for any $\lambda \in \mathbb{C}$ and for any non-trivial solution y of (1.2), the following always holds

$$\int_{\mathcal{J}_0} w(x) |y(x)|^2 dx > 0.$$

It is evident that (A_1) holds when w(t) > 0 on \mathcal{J} . Under the assumption (A_1) , it has been shown that $d_{\lambda}(\tau) = n_{\lambda}(\tau)$ for all $\lambda \in \mathbb{C}$. This equivalence does not hold without the assumption (A_1) [12, Proposition 2.19]). In the case that all the coefficients of τ are real-valued, i.e., all $b_k(t) \equiv 0$ on \mathcal{J} , it has been easily shown that $n_+(\tau) = n_-(\tau)$. This, together with (A_1) , implies $d_+(\tau) = d_-(\tau)$. However, the values of $d_{\pm}(\tau)$ may differ when τ has complexvalued coefficients. Mcleod [13] first gave an example of a fourth-order formally self-adjoint differential equation with $d_+(\tau) = 3$ and $d_-(\tau) = 2$. Later, Kogan and Rofe-Beketov [10, 11] showed that

$$|d_{+}(\tau) - d_{-}(\tau)| = 1$$

happens for any $m \geq 3$.

Many authors are interested in the positive and negative deficiency indices of formally self-adjoint difference equations, and have got many excellent results. A formally self-adjoint difference expression is necessary to be even, and it has the following form (see Theorem 3.3):

(1.3)
$$\mathcal{L}y := \sum_{j=0}^{n} F^{j}(A_{j}y) + \sum_{j=1}^{n} \overline{A}_{j}F^{-j}y,$$

where A_j , j = 1, ..., n, are complex-valued functions, and A_0 is real-valued on \mathcal{I} ; $A_n \neq 0$ on \mathcal{I} ; F is the forward shift operator, i.e., Fy(t) = y(t+1). The following equation

(1.4)
$$(\mathcal{L}y)(t) = \lambda w(t)y(t), \quad t \in \mathcal{I}$$

is called the formally self-adjoint equation generated by \mathcal{L} , where $w(t) \geq 0$ and $\lambda \in \mathbb{C}$. We take $\mathcal{I} := \{t\}_{t=0}^{+\infty}$ in the following. It is worth noting that the order of any formally self-adjoint difference equation is even. This is a difference between formally self-adjoint difference equations and differential equations.

In addition, \mathcal{L} defined by (1.3) can be rewritten as:

(1.5)
$$\mathcal{L}y = \sum_{j=0}^{n} (-1)^{j} \Delta^{j}(p_{j} \nabla^{j} y) + i \sum_{k=1}^{n} [(-1)^{k+1} \Delta^{k}(q_{k} y) + q_{k} \nabla^{k} y],$$

where Δ and ∇ are the forward and backward difference operators, respectively, i.e., $\Delta y(t) = y(t+1) - y(t)$ and $\nabla y(t) = y(t) - y(t-1)$; the coefficients p_j and q_k are all real-valued on \mathcal{I} . The coefficients between (1.3) and (1.5) have the following relationship:

$$A_{0}(t) = P_{0}(t), \quad A_{j}(t) = P_{j}(t) + iQ_{j}(t), \quad 1 \le j \le n,$$
$$P_{j}(t) = (-1)^{j} \sum_{s=j}^{n} \sum_{k=0}^{s-j} C_{s}^{k} C_{s}^{s-j-k} p_{s}(t+k), \quad 0 \le j \le n,$$
$$Q_{j}(t) = (-1)^{j+1} \sum_{k=j}^{n} C_{k}^{j} q_{k}(t), \quad 1 \le j \le n,$$

where $C_s^k = \frac{s!}{k!(s-k)!}$.

The definiteness condition for (1.4) is given by:

(A₂) there exists a bounded integer interval $\mathcal{I}_0 \subset \mathcal{I}$ such that for any $\lambda \in \mathbb{C}$ and for any non-trivial solution y of (1.4), the following always holds

$$\sum_{t\in\mathcal{I}_0} w(t)|y(t)|^2 > 0.$$

Similarly as the continuous case, the minimal linear relation generated by (1.4) is Hermitian and consequently its deficiency index $d_{\lambda}(\mathcal{L})$ is constant when λ is in the upper and lower half-planes [18]. Denote $d_{\pm}(\mathcal{L}) := d_{\pm i}(\mathcal{L})$. Under the assumption $(A_2), d_{\lambda}(\mathcal{L}) = n_{\lambda}(\mathcal{L})$ holds, where $n_{\lambda}(\mathcal{L})$ is the number of linearly independent solutions of (1.4), which satisfies

$$\sum_{t \in \mathcal{I}} w(t) |y(t)|^2 < +\infty.$$

Furthermore, $n \leq d_+(\mathcal{L}) = d_-(\mathcal{L}) \leq 2n$ holds and all the values in this range can be realized. The readers are referred to [2, 4, 5, 9, 14, 15, 23] for more details.

Up to now, all the existing results on the the positive and negative deficiency indices of formally self-adjoint difference equations coincide with those of their corresponding differential equations. So, one may think there would exist some special case of (1.4), corresponding to Mcleod's example mentioned above, satisfying $d_{+}(\mathcal{L}) \neq d_{-}(\mathcal{L})$. In this manuscript, we show that the positive and negative deficiency indices of any formally self-adjoint difference equations are equal (Theorem 4.1). This is a new and important difference in the spectral theory between the formally self-adjoint difference equations and their corresponding differential equations.

The rest of the paper is organized as follows. In Section 2, we introduce some basic concepts of the spectral theory in Hilbert space and give some sufficient and necessary conditions for an Hermitian operator to have equal positive and negative deficiency indices. In Section 3, we establish two characterizations of formally

self-adjoint difference expressions. In Section 4, we prove the positive and negative deficiency indices of any formally self-adjoint difference equations are equal (Theorem 4.1).

2. Fundamental results about linear relations and linear operators

In this section, we introduce some basic concepts of linear relations and linear operators, and show some results on the deficiency indices of Hermitian linear relations and linear operators and their self-adjoint extensions.

By \mathbb{C} , \mathbb{R} , and \mathbb{Z} denote the sets of all complex numbers, real numbers and integers, separately. By \bar{a} denote the conjugation of $a \in \mathbb{C}$, and $i := \sqrt{-1}$.

Let X be a complex Hilbert space with inner product $\langle \cdot, \cdot \rangle$. Let T_1 and T_2 be two linear relations (briefly, relation) in $X^2 := X \times X$. Denote

$$\mathcal{D}(T) := \{ x \in X : (x, f) \in T \text{ for some } f \in X \},\$$

$$\mathcal{R}(T) := \{ f \in X : (x, f) \in T \text{ for some } x \in X \},\$$

$$\mathcal{N}(T) := \{ x \in X : (x, 0) \in T \},\$$

$$T^* := \{ (x, f) \in X^2 : \langle x, g \rangle = \langle f, y \rangle \text{ for all } (y, g) \in T \},\$$

$$T(x) := \{ f \in X : (x, f) \in T \},\$$

$$T_1 + T_2 := \{ (x, f_1 + f_2) \in X^2 : (x, f_i) \in T_i, i = 1, 2 \}.$$

It is clear that $T(0) = \{0\}$ if and only if there exists a unique linear operator, denoted by $S_T : \mathcal{D}(T) \to X$, such that its graph $G(S_T) = T$. In this case, $S_T x = f$ for any $(x, f) \in T$.

In addition, T is said to be Hermitian if $T \subset T^*$; T is said to be symmetric if $T \subset T^*$ and $\mathcal{D}(T)$ is dense in X; T is said to be self-adjoint if $T = T^*$.

If T is a closed linear relation in X^2 , then T can be decomposed as [1]:

$$T = T_s \oplus T_{\infty},$$

where $T_{\infty} = \{(0, f) \in X^2 : f \in T(0)\}$ is called the pure multi-valued parts of T, and

$$T_s = T \ominus T_\infty$$

is called the operator part of T.

Lemma 2.1. [22, Proposition 2.1, 3.1] Let T be a closed Hermitian relation in X^2 . Then

$$T_{\infty} = T \cap (T(0))^2, \quad T_s = T \cap (T(0)^{\perp})^2,$$

and T_s is closed Hermitian in $(T(0)^{\perp})^2$. Further, if T is self-adjoint in X^2 , then T_s is self-adjoint in $(T(0)^{\perp})^2$.

Let T be a linear relation (operator) in $X^2(X)$. The subspace $\mathcal{R}(T - \lambda I))^{\perp}$ and the number $d_{\lambda}(T) := \dim \mathcal{R}(T - \lambda I)^{\perp}$ are called the deficiency space and deficiency index of T and λ , separately. By \overline{T} denote the closure of T. It can be easily verified that $d_{\lambda}(T) = d_{\lambda}(\overline{T})$ for all $\lambda \in \mathbb{C}$.

Let T be a linear relation in X^2 . The set

$$\Gamma(T) := \{\lambda \in \mathbb{C} : \exists c(\lambda) > 0 \text{ s.t. } \|f - \lambda x\| \ge c(\lambda) \|x\|, \forall (x, f) \in T\}$$

is called the regularity domain of T.

It has been shown by [21, Theorem 2.3] that the deficiency index $d_{\lambda}(T)$ is constant in each connected subset of $\Gamma(T)$. If T is Hermitian, then $d_{\lambda}(T)$ is constant in the upper and lower half-planes. So, we can denote $d_{\pm}(T) := d_{\pm i}(T)$ for an Hermitian linear relation T, and call $d_{\pm}(T)$ the positive and negative deficiency indices of T, respectively.

Lemma 2.2. [6, Theorem 15] Let T be a closed Hermitian relation in X^2 . Then T has self-adjoint extensions if and only if $d_+(T) = d_-(T)$.

Lemma 2.3. [22, Corollary 2.1] Let T be a closed Hermitian relation in X^2 . Then $d_{\pm}(T) = d_{\pm}(T_s)$, and consequently, T has a self-adjoint extension in X^2 if and only if T_s has a self-adjoint extension in $(T(0)^{\perp})^2$.

A relation T is said to be bounded from below (above) if there exists a number $c\in\mathbb{R}$ such that

$$\langle x, f \rangle \ge c \|x\|^2$$
, $(\langle x, f \rangle \le c \|x\|^2)$, $\forall (x, f) \in T$,

while such a constant c is called a lower (upper) bound of T.

Lemma 2.4. [3, Proposition 1.4.6] Let T be an Hermitian relation and be bounded from below with lower bound c. Then $\mathbb{C} \setminus [c, +\infty) \subset \Gamma(T)$, and the deficiency index $d_{\lambda}(T)$ is constant for all $\lambda \in \mathbb{C} \setminus [c, +\infty)$.

Combining Lemmas 2.2 and 2.4, one can get the following result.

Lemma 2.5. Let T be an Hermitian relation and be bounded from below. Then $d_+(T) = d_-(T)$ and T has self-adjoint extensions.

Next, we pay attention to the self-adjointness of an operator by using the concept of K-real. Let S be an operator on X.

Definition 2.6. [24, Section 8.1] Let X be a complex Hilbert space. A mapping K of X onto itself is called a conjugation if

- (1) $K(ax + by) = \bar{a}K(x) + \bar{b}K(y)$ for all $x, y \in X, a, b \in \mathbb{C}$.
- (2) $K^2 = I$.
- (3) $\langle Kx, Ky \rangle = \langle y, x \rangle$ for all $x, y \in X$.

An operator S on X is said to be a K-real if

- (1) $K\mathcal{D}(S) \subset \mathcal{D}(S)$
- (2) SKx = KSx for $x \in \mathcal{D}(S)$.

A sufficient condition for a symmetric operator to have equal positive and negative deficiency indices is given by [24, Theorem 8.9]. Since S is not required to be densely defined in the proof, the assertion is still true for Hermitian operators.

Lemma 2.7. [24, Theorem 8.9] Let X be a complex Hilbert space, and let K be a conjugation on X. If S is a K-real Hermitian operator on X, then $d_+(S) = d_-(S)$.

The following is a necessary condition for an operator S to be self-adjoint.

Theorem 2.8. Let X be a complex Hilbert space, and let S be a self-adjoint operator on X. Then there exists a conjugation K for which S is K-real.

To prove Theorem 2.8, we introduce several notations and a lemma first. The readers are referred to [24].

Let $\{\rho_{\alpha} : \alpha \in A\}$ be a family of right continuous non-decreasing function defined on \mathbb{R} . By $L^2(\mathbb{R}, \rho_{\alpha})$ denote the set of all the square integrable functions with respect to ρ_{α} on \mathbb{R} . It has been shown that $L^2(\mathbb{R}, \rho_{\alpha})$ is a Hilbert space with inner

$$\langle f,g \rangle_{\alpha} = \int_{\mathbb{R}} \bar{g}(t) f(t) d\rho_{\alpha}(t).$$

By $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_\alpha)$ denote the orthogonal sum of the spaces $L^2(\mathbb{R}, \rho_\alpha)$. Then $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_\alpha)$ is a Hilbert space with inner

$$\langle (x_{\alpha}), (y_{\alpha}) \rangle = \sum_{\alpha \in A} \langle x_{\alpha}, y_{\alpha} \rangle_{\alpha}, \quad (x_{\alpha}), (y_{\alpha}) \in \bigoplus_{\alpha \in A} L^{2}(\mathbb{R}, \rho_{\alpha}).$$

By K_0 denote the natural conjugation on $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_\alpha)$, i.e.,

(2.1)
$$K_0(x_\alpha) := (\bar{x}_\alpha), \quad (x_\alpha) \in \bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_\alpha).$$

The following result is a combination of Theorems 7.16-7.18 of [24].

Lemma 2.9. Let S be a self-adjoint operator on X. Then there exists exactly one spectral family E for which $T = \hat{E}(id)$. Moreover, for this spectral family E, there exists a family $\{\rho_{\alpha} : \alpha \in A\}$ of right continuous non-decreasing functions (the cardinality of A is at most the dimension of X) and a unitary operator $U : X \to$ $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_{\alpha})$ for which

$$(2.2) S = U^{-1} S_{\rm id} U,$$

where S_{id} denotes the maximal operator of multiplication by the function id on $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_{\alpha}).$

Now we give the proof of Theorem 2.8.

Proof. Let S be a self-adjoint operator on X. Then it follows from Lemma 2.9 that there exists a family $\{\rho_{\alpha} : \alpha \in A\}$ of right continuous non-decreasing functions (the cardinality of A is at most the dimension of X) and a unitary operator $U : X \to$ $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_{\alpha})$ such that (2.2) holds. Based on the proof of [24, Theorems 7.16], the unitary operator U can be constructed to be linear, i.e.,

$$U(ax + by) = aU(x) + bU(y), \quad x, y \in X, \ a, b \in \mathbb{C}.$$

Let

(2.3)
$$K = U^{-1} K_0 U,$$

where K_0 is the natural conjugation on $\bigoplus_{\alpha \in A} L^2(\mathbb{R}, \rho_\alpha)$ defined by (2.1). Then it follows that

(2.4)
$$K(ax+by) = \bar{a}K(x) + bK(y)$$

for all $x, y \in X$, $a, b \in \mathbb{C}$. Further, by the facts that K_0 is a conjugation and U is a unitary operator, it follows that

(2.5)
$$K^{2} = U^{-1}K_{0}UU^{-1}K_{0}U = I,$$
$$\langle Kx, Ky \rangle = \langle K_{0}Ux, K_{0}Uy \rangle = \langle Uy, Ux \rangle = \langle y, x \rangle, \quad x, y \in X.$$

These yield that K defined by (2.3) is a conjugation on X.

Further, it can be verified that S_{id} is K_0 -real, and $U\mathcal{D}(S) \subset \mathcal{D}(S_{id})$. Therefore it follows that $K\mathcal{D}(S) \subset \mathcal{D}(S)$, and

$$SKx = U^{-1}S_{\mathrm{id}}K_0Ux = U^{-1}K_0S_{\mathrm{id}}Ux = KSx, \quad x \in \mathcal{D}(S).$$

This implies that S is K-real. The proof is complete.

3. Characterization of formally self-adjoint difference expressions

In this section we pay attention to the characterization of formally self-adjoint difference expressions.

First, we introduce some notations and concepts. Let X be a vector space over \mathbb{C} . A mapping $s: X \times X \to \mathbb{C}$ is called a sesquilinear form on X if it follows

$$\begin{split} s[x,ay+bz] &= \bar{a}s[x,y] + \bar{b}s[x,z], \\ s[ay+bz,x] &= as[y,x] + bs[z,x], \end{split}$$

for all $x, y, z \in X$ and $a, b \in \mathbb{C}$. Let $\mathcal{I} = \{t\}_{t=0}^{+\infty}$. For some $n \in \mathbb{Z}$, denote

$$l(\mathcal{I}, n) = \{ x = \{ x(t) \}_{t=-n}^{+\infty} : \ x(t) \in \mathbb{C} \}$$

Let L be an arbitrary difference expression of order m defined on $l(\mathcal{I}, m)$:

$$(Ly)(t) := \sum_{j=0}^{k} A_j(t+j)y(t+j) + \sum_{j=1}^{s} A_{-j}(t)y(t-j),$$

where all $A_j(t)$, $-s \leq j \leq k$, are complex valued functions; k and s are non-negative integer satisfying k + s = m, and $A_k(t+k) \neq 0$, $A_{-s}(t) \neq 0$ for $t \in \mathcal{I}$. By F denote the forward shift operator, i.e.,

$$Fy(t) = y(t+1).$$

and define $F^{-1}y(t) = y(t-1), F^j = FF^{j-1}$. Thus L can be rewritten briefly as

(3.1)
$$Ly = \sum_{j=0}^{k} F^{j}(A_{j}y) + \sum_{j=1}^{s} A_{-j}F^{-j}y$$

Now, we give the definition of the formal adjoint of L and try to establish the characterization of its formal adjoint.

Definition 3.1. Let L be an *m*th-order difference expression. An *m*th-order difference expression, denoted by L^+ , is said to be a formal adjoint of L, if there exists a sesquilinear form $s[\cdot, \cdot]$ on $l(\mathcal{I}, m)$ such that

$$\overline{y}(t)(Lx)(t) - \overline{(L^+y)(t)}x(t) = \Delta s[x,y](t), \quad t \in \mathcal{I}.$$

Moreover, L is said to be formally self-adjoint if $L = L^+$.

For $n \ge 0$, let L_n be a difference expression defined as the following:

(3.2)
$$(L_n y)(t) := \sum_{j=0}^n F^j(B_j y)(t),$$

where all $B_j(t)$, $0 \le j \le n$, are complex-valued functions, $B_n(t) \ne 0$ on \mathcal{I} . We get the following result on the formal adjoint of L_n .

Lemma 3.2. Let L_n be an nth-order difference expression defined by (3.2). Then

(1) (Green Formula) For any $x, y \in l(\mathcal{I}, n), 0 \leq k \leq r$,

(3.3)
$$\sum_{t=k}^{r} \left[\bar{y}(t) L_n x(t) - \overline{(L_n^+ y(t))} x(t) \right] = s[x, y](r+1) - s[x, y](k),$$

where L_n^+ is given by

(3.4)
$$(L_n^+ y)(t) = \sum_{j=0}^n \overline{B}_j(t) F^{-j} y(t),$$

and s[x, y] is a sesquilinear on $l(\mathcal{I}, n)$ with the form

(3.5)
$$s[x,y](t) = \sum_{k=1}^{n} \sum_{j=0}^{k-1} \bar{y}(t-j-1)B_k(t+j)x(t+j).$$

(2)
$$L_n + L_n^+$$
 is a 2nth-order formally self-adjoint difference expression.

Proof. (1) Let L_n^+ and s[x, y] be defined as (3.4) and (3.5), respectively. It suffices to show

(3.6)
$$\bar{y}(t)L_nx(t) - \overline{(L_n^+y)(t)}x(t) = \Delta s[x,y](t), \quad t \in \mathcal{I}.$$

We will proof (3.6) by induction. For n = 1, it follows that

$$\bar{y}(t)(L_1x)(t) - \overline{(L_1^+y)(t)}x(t) = \bar{y}(t)B_1(t+1)x(t+1) - \bar{y}(t-1)B_1(t)x(t)$$
$$= \Delta[\bar{y}(t-1)B_1(t)x(t)].$$

This yields (3.6) with n = 1. Assume that (3.6) hold for n - 1. It can be easily verified

$$\bar{y}(t)B_n(t+n)x(t+n) - \bar{y}(t-n)B_n(t)x(t) = \Delta \sum_{j=0}^{n-1} [\bar{y}(t-j-1)B_n(t+j)x(t+j)].$$

Then it follows that

$$\begin{split} \bar{y}(t)(L_n x)(t) &- \overline{(L_n^+ y)(t)} x(t) \\ &= [\bar{y}(t)(L_{n-1} x)(t) - \overline{(L_{n-1}^+ y)(t)} x(t)] + \bar{y}(t) B_n(t+n) x(t+n) - \bar{y}(t-n) B_n(t) x(t) \\ &= \Delta \sum_{k=1}^{n-1} \sum_{j=0}^{k-1} \bar{y}(t-j-1) B_k(t+j) x(t+j) + \Delta \sum_{j=0}^{n-1} [\bar{y}(t-j-1) B_n(t+j) x(t+j)] \\ &= \Delta \sum_{k=1}^n \sum_{j=0}^{k-1} \bar{y}(t-j-1) B_k(t+j) x(t+j). \end{split}$$

Thus (3.6), and consequently, assertion (1) is proved.

(2) Assertion (1) yields that $(L_n^+)^+ = L_n$. So, $(L_n + L_n^+)^+ = L_n + L_n^+$, which implies that $L_n + L_n^+$ is formally self-adjoint. The proof is complete.

Based on the above discussion, we now establish a characterization of formally self-adjoint difference expressions.

Theorem 3.3. Let \mathcal{L} be a formally self-adjoint difference expression of order m. Then m is even, saying m = 2n, and \mathcal{L} has the the following form

(3.7)
$$\mathcal{L}y = \sum_{j=0}^{n} F^{j}(A_{j}y) + \sum_{j=1}^{n} \overline{A}_{j}F^{-j}y,$$

where A_j , j = 1, ..., n, are complex-valued functions, and A_0 is real-valued.

Proof. First, we shown \mathcal{L} defined by (3.7) is self-adjoint. Denote

$$L_n y = \sum_{j=1}^n F^j(A_j y) + \frac{1}{2} A_0 y.$$

Then one has that $\mathcal{L} = L_n + L_n^+$ by (1) of Lemma 3.2, and consequently, \mathcal{L} is formally self-adjoint by (2) of Lemma 3.2.

On the other hand, we proof \mathcal{L} has form (3.7) if \mathcal{L} is self-adjoint. Let L be any *m*th-order difference expression with the form of (3.1). Then it can be written as $L = L_k + L_s$, where

$$(L_k y)(t) = \sum_{j=0}^k A_j(t+j)y(t+j), \quad L_s y(t) = \sum_{j=1}^s A_{-j}(t)y(t-j).$$

It follows from (1) of Lemma 3.2 that

$$L^{+}y(t) = L_{k}^{+}y(t) + \hat{L}_{s}^{+}y(t) = \sum_{j=1}^{s} \overline{A}_{-j}(t+j)y(t+j) + \sum_{j=0}^{k} \overline{A}_{j}(t)y(t-j).$$

This yields that $L = L^+$ if and only if s = k, which implies that m is even, saying m = 2n; and $A_j(t) \equiv \overline{A}_{-j}(t)$ for j = 0, 1, ..., n, which implies that $A_0(t)$ is real-valued. The proof is complete.

Theorem 3.4. For any nth-order forward difference expression L_n , $L_nL_n^+$ is a formally self-adjoint difference expression of order 2n.

Proof. Let L_n be defined as (3.2). Then it can be verified that

$$(L_n L_n^+)y = L_n (L_n^+ y) = \sum_{j=0}^n F^j (D_j y) + \sum_{j=1}^n \overline{D}_j F^{-j} y,$$

where

(3.8)
$$D_j(t) = \sum_{k=j}^n B_k(t+k-j)\overline{B}_{k-j}(t+k-j), \quad j = 0, 1, \dots, n, \quad t \in \mathcal{I}.$$

It is clear $D_0(t) = \sum_{k=0}^n |B_k(t+k)|^2$ is real-valued on \mathcal{I} . So, $L_n L_n^+$ is formally self-adjoint by Theorem 3.3. The proof is complete.

Now we can give another characterization of formally self-adjoint difference expression \mathcal{L} by using L_n .

Theorem 3.5. For any formally self-adjoint difference expression \mathcal{L} with order 2n, there exist two difference expressions L_n and L_0 with the form of (3.3) such that

$$\mathcal{L} = L_n L_n^+ + L_0.$$

Proof. Let \mathcal{L} be formally self-adjoint difference expression defined as (3.7). By Theorem 3.4 and (3.8), it suffices to show there exists a set of complex-valued functions $B_j(t)$, $0 \leq j \leq n$, and a real-valued function C(t) such that

(3.9)
$$(L_n y)(t) = \sum_{j=0} F^j(B_j y)(t), \quad (L_0 y)(t) = C(t)y(t)$$

and

(3.10)
$$\sum_{\substack{k=j\\n}}^{n} B_k(t+k-j)\overline{B}_{k-j}(t+k-j) \equiv A_j(t), \quad j=1,2,\dots,n,$$

(3.11)
$$\sum_{k=0}^{n} B_k(t+k-j)\overline{B}_{k-j}(t+k-j) + C(t) \equiv A_0(t).$$

By taking $B_0(t) \equiv 1$ on \mathcal{I} , it can be easily verified that $B_n(t) = A_n(t)$ on \mathcal{I} yields (3.10) with j = n. In addition, by giving the following initial values

$$B_{1}(0), B_{2}(0), B_{2}(1), \dots B_{n-1}(0), B_{n-1}(1), \dots B_{n-1}(n-2),$$

$$C(t) = A_0(t) - \sum_{k=0}^n |B_k(t+k)|^2, \quad t \in \mathcal{I}.$$

Then (3.11) is satisfied. The proof is complete.

4. Proof of Theorem 1.1

Let $l(\mathcal{I}, n)$ be defined as that in Section 3, and \mathcal{L} be a formally self-adjoint difference expression of order 2n defined on $l(\mathcal{I}, n)$. Denote

$$l_w^2 := \left\{ x \in l(\mathcal{I}, n) : \sum_{t=0}^{+\infty} w(t) |x(t)|^2 < +\infty \right\}.$$

and

$$\langle x, y \rangle_w := \sum_{t=0}^{+\infty} \bar{y}(t) w(t) x(t), \quad \|x\|_w := \langle x, x \rangle_w^{1/2}.$$

For any $x, y \in l_w^2$, we say x = y if ||x|| = ||y||. Then l_w^2 is a Hilbert space with the inner product $\langle \cdot, \cdot \rangle_w$ (cf. [19, Lemma 2.5]).

Denote

$$l_{w,0}^{2}(\mathcal{I}) := \left\{ y \in l_{w}^{2}(\mathcal{I}) : y(t) = 0, \ -n \le t \le n-1 \text{ and } t \ge k \text{ for some } k \in \mathcal{I} \right\}.$$

and

$$T := \{ (x, f) \in (l_w^2(\mathcal{I}))^2 : (\mathcal{L}x)(t) = w(t)f(t) \text{ on } \mathcal{I} \}$$
$$T_0 := \{ (x, f) \in T : x \in l_{w,0}^2(\mathcal{I}) \}.$$

T, T_0 , and the closure of T_0 , denoted by \overline{T}_0 , are called the maximal, the preminimal, and the minimal linear relations corresponding to \mathcal{L} , separately. Thus, $d_{\lambda}(\mathcal{L})$, given in Section 1, just refers to $d_{\lambda}(\overline{T}_0)$, which is equal to $d_{\lambda}(T_0)$.

It follows from Theorem 3.5 there exist two difference expressions L_n and L_0 defined as (3.9) such that $\mathcal{L} = L_n L_n^+ + L_0$. Similarly, denote

$$H := \{ (x, f) \in (l_w^2(\mathcal{I}))^2 : (L_n L_n^+ x)(t) = w(t) f(t) \text{ on } \mathcal{I} \},\$$

$$H_0 := \{ (x, f) \in H : x \in l_{w,0}^2(\mathcal{I}) \}.$$

Lemma 4.1. Both T_0 and H_0 are Hermitian linear relations.

Proof. Since both \mathcal{L} and $L_n L_n^+$ are formally self-adjoint, it follows that both T_0 and H_0 are Hermitian linear relations by using (3.3).

Theorem 4.2. For any formally self-adjoint difference expression \mathcal{L} ,

(4.1)
$$d_+(\mathcal{L}) = d_-(\mathcal{L})$$

Proof. Based on the notations given above, (4.1) is equivalent to .

$$d_+(T_0) = d_-(T_0).$$

For any $(x, f) \in H_0$, it follows from (3.3) that

$$\langle x, f \rangle_w = \sum_{t \in \mathcal{I}} x(t) \overline{L_n L_n^+ x(t)} = \sum_{t \in \mathcal{I}} L_n^+ x(t) \overline{L_n^+ x(t)} \ge 0.$$

This yields that H_0 is Hermitian and bounded from below with lower bound 0. Hence, it follows from Lemma 2.5 that H_0 has self-adjoint extensions. Let H_1 be a self-adjoint extension of H_0 . By $H_{1,s}$ denote the operator part of H_1 . It follows from Lemma 2.1 that $H_{1,s}$ is self-adjoint in $(H_1^{\perp}(0))^2$, and there exists uniquely a self-adjoint operator, denoted by S_H in $H_1^{\perp}(0)$ such that $G(S_H) = H_{1,s}$. Then by Theorem 2.8, there exists a conjugate K on $H_1^{\perp}(0)$ such that S_H is K-real.

Corresponding to difference expression L_0 , we define

$$\mathcal{D}(S_0) = \{ x \in l_w^2(\mathcal{I}) \cap H_1^{\perp}(0) : \exists f \in l_w^2(\mathcal{I}) \ s.t. \ C(t)x(t) = w(t)f(t) \ \text{on } \mathcal{I} \}, \\ S_0 x = f.$$

It can be easily verified that S_0 is a well-defined operator on $l_w^2(\mathcal{I})$. Moreover, S_0 is K-real. In fact, for any $x \in \mathcal{D}(S_0) \cap H_1^{\perp}(0)$, there exists uniquely $f \in l_w^2(\mathcal{I})$ such that C(t)x(t) = w(t)f(t) on \mathcal{I} . Since both C(t) and w(t) are real-valued, it follows that

$$C(t)(Kx)(t) = w(t)(Kf)(t), \quad t \in \mathcal{I}.$$

This implies that $K\mathcal{D}(S_0) \subset \mathcal{D}(S_0)$ and $S_0Kx = Kf = KS_0x$. Thus S_0 is K-real, and consequently, $S_H + S_0$ is K-real. So, by Lemma 2.7 one has that

$$d_+(S_H + S_0) = d_-(S_H + S_0)$$

and consequently, $S_H + S_0$ has self-adjoint extensions by Lemma 2.2.

On the other hand, it is clear \overline{T}_0 is closed and Hermitian. Let $\overline{T}_{0,s}$ be the operator part of \overline{T}_0 and operator S_T be the operator which satisfies $G(S_T) = \overline{T}_{0,s}$. Then one has $S_T \subset S_H + S_0$. Thus, all the self-adjoint extensions of $S_H + S_0$ are still the self-adjoint extensions of S_T . Again by Lemma 2.2, one has $d_+(S_T) = d_-(S_T)$. This, together with $\mathcal{R}(\overline{T}_0) = \mathcal{R}(S_T)$, yields (4.1). The proof is complete.

References

- 1. R. Arens, Operational calculus of linear relations, Pac. J. Math., 11(1961), 9-23.
- F. V. Atkinson, Discrete and Continuous Boundary Problems, Academic Press, Inc., New York, 1964.
- J. Behrndt, S. Hassi, and H. Snoo, Boundary Value Problems, Weyl Functions, and Differential Operators, Monographs in Mathematiscs, 108, Birkhäuser, 2020.
- J. Chen, Y. Shi, The limit circle and limit point criteria for second-order linear difference equations, Comput. Math. Appl., 42(2001), 465-476.
- S. L. Clark, A spectral analysis for self-adjoint operators generated by a class of second order difference equations, J. Math. Anal. Appl., 197(1996), 267-285.
- E. A. Coddington, Extension theory of formally normal and symmetric linear relations, Mem. Amer. Math. Soc., 134(1973).
- 7. R. Cross, Multivalued Linear Operators, New York, Marcel Dekker, 1998.
- N. Dunford, J. T. Schwartz, Linear operators II: Spectral Theory of Self-adjoint Operators in Hilbert Space, New York, 1963.
- A. Jirari, Second-order Sturm-Liouville difference equations and orthogonal polynomials, Mem. Amer. Math. Soc., 542(1995), 1-136.
- V. I. Kogan, F.S. Rofe-Beketov, On the question of the deficiency indices of differential operators with complex coefficients, Proc. Royal Soc. Edin., 72(1973), 281-298.
- V. I. Kogan, F.S. Rofe-Beketov, On sqare-integrable solutions of symmetric systems of differential equations of arbitrary order, Proc. Royal Soc. Edin., 74(1974), 5-40.
- M. Lesch, M. Malamud, On the deficiency indices and self-asjointness of symmetric Hamiltonian systems, J. Differ. Equations 18(2003), 556-615.
- J. B. Mcleod, The number of integrable-square solutions of ordinary differential equations, Quart. J. Math. Oxford(2), 17(1966), 285-290.
- G. Ren, Y. Shi, The defect index of singular symmetric linear difference equations with real coefficients, Proc. Amer. Math. Soc., 138(2010), 2463-2475.
- G. Ren, Y. Shi, Defect indices and definiteness conditions for a class of discrete linear Hamiltonian systems, Appl. Math. Comp., 218(2011), 3414-3429.
- G. Ren, Defect indices of singular symmetric linear difference equations with complex coefficients, Adv. Differ. Equations 27(2012), 1-13.
- G. Ren, On the density of the minimal subspaces generated by discrete linear Hamiltonian systems, Appl. Math. Lett., 27 (2014), 1-5.
- G. Ren, Y. Shi, Self-adjoint extensions for discrete linear Hamiltonian systems, Linear Algebra Appl., 454 (2014), 1-48.
- Y. Shi, Weyl-Titchmarsh theory for a class of singular discrete linear Hamiltonian systems, Linear Algebra Appl., 416(2006), 452-519.
- Y. Shi and H. Sun, Self-adjoint extensions for second-order symmetric linear difference equations, Linear Algebra Appl. 434 (2011), no. 4, 903-930.
- Y. Shi, The Glazman-Krein-Naimark theory for Hermitian subspaces, J. Operat. Theor., 68(2012), 241-256.
- Y. Shi, C. Shao, and G.Ren, Spectral properties of self-adjoint subspaces, Linear Algebra Appl., 438 (2013) 191-218.
- H. Sun, Y. Shi, Limit-point and limit-circle criteria for singular second-order linear difference equations with complex coefficients, Comput. Math. Appl., 52 (2006), 539-554.
- 24. J. Weidmann, Linear Operators in Hilbert Spaces, Springer-Verlag, New York, 1980.

School of Mathematics and Quantitative Economics, Shandong University of Finance and Economics, 250014, P. R. China

Email address: gjren@sdufe.edu.cn