

LEFT NON-DEGENERATE SET-THEORETIC SOLUTIONS OF THE YANG-BAXTER EQUATION AND SEMITRUSSES

I. COLAZZO, E. JESPERS, A. VAN ANTWERPEN AND C. VERWIMP

ABSTRACT. To determine and analyze arbitrary left non-degenerate set-theoretic solutions of the Yang-Baxter equation (not necessarily bijective), we introduce an associative algebraic structure, called a YB-semitruss, that forms a subclass of the category of semitrusses as introduced by Brzeziński. Fundamental examples of YB-semitrusses are structure monoids of left non-degenerate set-theoretic solutions and (skew) left braces. Gateva-Ivanova and Van den Bergh introduced structure monoids and showed their importance (as well as that of the structure algebra) for studying involutive non-degenerate solutions. Skew left braces were introduced by Guarnieri, Vendramin and Rump to deal with bijective non-degenerate solutions. Hence, YB-semitrusses also yield a unified treatment of these different algebraic structures. The algebraic structure of YB-semitrusses is investigated and as a consequence it is proven, for example, that any finite left non-degenerate set-theoretic solution of the Yang-Baxter equation is right non-degenerate if and only if it is bijective. Furthermore, it is shown that some finite left non-degenerate solutions can be reduced to non-degenerate solutions of smaller size. The structure algebra of a finitely generated YB-semitruss is an algebra defined by homogeneous quadratic relations. We prove that it often is a left Noetherian algebra of finite Gelfand-Kirillov dimension that satisfies a polynomial identity, but in general it is not right Noetherian.

CONTENTS

1. Introduction	2
2. Left non-degenerate solutions and YB-semitrusses	5
2.1. Structure monoids as YB-semitrusses	14
2.2. Left cancellative YB-semitrusses	18
2.3. Idempotents in YB-semitrusses	21
2.4. Matched product of solutions and YB-semitrusses	23
3. YB-semitrusses with associated bijective solutions	25
4. Non-degenerate YB-semitrusses and the retract relation	28
5. Algebraic structure of YB-semitrusses	32
5.1. Additive structure of YB-semitruss and the structure algebra	32
5.2. YB-semitrusses with a left simple semigroup of endomorphisms and their solutions	37
Acknowledgement	39
References	39

2020 *Mathematics Subject Classification.* 16T25, 16W22, 20M25, 16S36, 16Y99.

Key words and phrases. Yang–Baxter equation, set-theoretic solution, structure semigroup, structure algebra, semitruss.

1. INTRODUCTION

Let V be a vector space over a field K . A linear map $R: V \otimes V \rightarrow V \otimes V$ is called a solution of the Yang–Baxter equation if

$$(R \otimes \text{id})(\text{id} \otimes R)(R \otimes \text{id}) = (\text{id} \otimes R)(R \otimes \text{id})(\text{id} \otimes R).$$

This equation originates from papers by Baxter [7] and Yang [64] on statistical physics and the search for solutions has attracted numerous studies both in mathematical physics and pure mathematics. As the study of arbitrary solutions is complex, Drinfeld in [27] initiated the investigations of set-theoretic solutions of the Yang–Baxter equation. These are solutions which are induced by a linear extension of a map $r: X \times X \rightarrow X \times X$, where X is a basis of V . In this case r satisfies

$$(r \times \text{id})(\text{id} \times r)(r \times \text{id}) = (\text{id} \times r)(r \times \text{id})(\text{id} \times r),$$

and one says that (X, r) is a set-theoretic solution of the Yang–Baxter equation (throughout the paper we will simply call this a solution). For any $x, y \in X$, we put $r(x, y) = (\lambda_x(y), \rho_y(x))$. The solution is said to be bijective if r is bijective. If furthermore $r^2 = \text{id}$ then it is said to be an involutive solution. A solution is said to be left non-degenerate (respectively right non-degenerate) if all maps λ_x (respectively all maps ρ_x) are bijective; a non-degenerate solution is one that is a left and right non-degenerate solution. Solutions of the Yang–Baxter equation also are of central importance for many mathematical topics, including quantum groups, Hopf algebras, Galois theory, group algebras, knot theory, radical rings, pre-Lie algebras, solvable groups and Garside groups (see for example [3, 16, 17, 23, 26, 28–30, 34, 46, 50, 56, 57, 59, 60]).

In recent years several fundamental new algebraic structures have been introduced to describe certain classes of solutions. Rump [55] introduced the notion of a (left) brace, that is a tuple $(B, +, \circ)$ with $(B, +)$ an abelian group and (B, \circ) a group so that $a \circ (b + c) = (a \circ b) - a + (a \circ c)$ holds for all $a, b, c \in B$ (see also [18]). If one assumes that $(B, +)$ is an arbitrary group then one obtains the notion of a skew left brace as introduced by Guarnieri and Vendramin [38]. It has been shown by Bachiller, Cedó and Jespers in [6] that all involutive non-degenerate solutions can be explicitly constructed from the description of all left braces, and later this has been extended by Bachiller in [5] to all bijective non-degenerate solutions from all skew left braces. This reduces the classification of all bijective non-degenerate solutions to classify all skew left braces.

The starting points of all these investigations is in the work of Gateva-Ivanova and Van den Bergh [34], Etingoff, Schedler and Soloviev [29], where a beautiful monoid and group theory translation was given of involutive non-degenerate solutions (X, r) , associating a monoid and group, denoted by $M(X, r)$ and $G(X, r)$ respectively. These are called the structure monoid and structure group. Recall that, for a solution (X, r) , the structure monoid $M(X, r)$ is the monoid (with operation written multiplicatively as \circ) generated by the elements of X subject to the relations $x \circ y = \lambda_x(y) \circ \rho_y(x)$. If one considers the group generated by X subject to these relations, considered as group relations, then one obtains the structure group $G(X, r)$. For a field K , the structure algebra is the K -algebra defined by these relations, and hence, since the relations are homogeneous, (quadratic), this is the monoid algebra $K[M(X, r)]$. It turns out [29, 41] that if (X, r) is an involutive non-degenerate solution then $M(X, r)$ is a regular submonoid of the holomorph of the free abelian monoid FaM_X of rank $|X|$ (with basis X) and $G(X, r)$ is the group of fractions of this cancellative monoid. The group $G(X, r)$ is a Bieberbach group and it is a regular subgroup of the holomorph of the free abelian group Fa_X (with basis X) of rank $|X|$. So, $M(X, r) = \{(a, \lambda_a) \mid a \in FaM_X\}$ is a submonoid of the natural

semidirect product $FaM_X \rtimes \text{Sym}(X)$, where $\lambda : FaM_X \rightarrow \text{Aut}(FaM_X) : a \mapsto \lambda_a$ is a map so that $\lambda_1 = \text{id}$ and λ_x is the mapping obtained from the defining solution (X, r) . Furthermore, as shown by Gateva-Ivanova and Van den Bergh [34], in this case, the structure algebra $K[M(X, r)]$ has a rich algebraic structure that resembles that of a polynomial algebra in finitely many commuting variables. In particular, if X is finite then it is left and right Noetherian domain that satisfies a polynomial identity (and it is a maximal order). For details and a survey we refer the reader to [42].

Lu, Yan and Zhu [52] and Soloviev [62] considered arbitrary bijective non-degenerate solutions and studied their structure groups. In contrast to the involutive case, the set X is not necessarily canonically embedded into the structure group $G(X, r)$ see [51, 61]. This problem can be avoided by working with the structure monoid and many of the above mentioned results have been extended to arbitrary bijective (left) non-degenerate solutions [39, 40]. Furthermore, it has been shown that the algebraic structure of the algebra $K[M(X, r)]$ determines crucial information of the solutions, for example, it is a domain (or equivalently it is a prime algebra) if and only if (X, r) is involutive. The structure algebra $R = K[M(X, r)] = \bigoplus_{n \in \mathbb{N}} R_n$ has a natural gradation and it is a locally finite connected graded algebra, i.e. $\dim_K(R_n) < \infty$ and $R_0 = K$. Moreover, it is generated by the elements of degree 1. In [32], Gateva-Ivanova posed several interesting questions on $\dim_K(R_2)$ for non-degenerate finite solutions (X, r) . This dimension is at most $\binom{|X|}{2}$ and this bound is reached precisely when (X, r) is involutive (see [39] and [34]). In [19] the minimal possible values of the dimension of R_2 are determined and one completely classifies non-degenerate finite solutions (X, r) for which these bounds are attained.

Rump introduced some other structures to study bijective (left) non-degenerate solutions (X, r) , such as cycle sets [54], q -cycle sets and q -braces [58]. Cycle sets and q -cycle sets are in a one-to-one correspondence with left non-degenerate involutive, respectively left non-degenerate solutions. Moreover, q -braces are groups equipped with two binary operations that give a q -cycle set structure. Such structures describe non-degenerate q -cycle sets, i.e. non-degenerate bijective solutions. Skew left braces are an intermediate step between left braces and q -braces.

Brzeziński in [8] initiated another notion, called a left semitruss, this to put the algebraic structures of skew left braces and structure monoids into a larger context, purely from an algebraic point of view, without making the link with solutions. A left semitruss is a tuple $(A, +, \circ, \lambda)$ such that $(A, +)$ and (A, \circ) are semigroups and $\lambda : A \rightarrow \text{Map}(A, A)$ is a function so that $a \circ (b + c) = (a \circ b) + \lambda_a(c)$, for all $a, b, c \in A$. A skew left brace $(A, +, \circ)$ is a left semitruss with $(A, +)$ and (A, \circ) groups and $\lambda_a(c) = -a + (a \circ c)$. Based on a fundamental result of Gateva-Ivanova and Majid [33], showing that any solution (X, r) can be lifted to a solution on the structure monoid $M(X, r)$, Cedó, Jespers and Verwimp showed in [20] that $M(X, r)$ is a left semitruss (by considering it as a regular submonoid of the structure monoid of an associated shelf solution) for any left non-degenerate solution (X, r) .

The aim of this paper is to study arbitrary left non-degenerate solutions via the study of certain, strongly related, associative structures. For this, we consider within the category of left semitrusses those that yield left non-degenerate solutions, and we therefore will call these objects YB-semitrusses. It turns out that such a YB-semitruss is a tuple $(A, +, \circ, \lambda, \sigma)$, with $(A, +, \circ, \lambda)$ a left semitruss, so that $a \circ b = a + \lambda_a(b)$ and $\sigma : A \rightarrow \text{Map}(A, A) : a \mapsto \sigma_a$ is a map that determines the addition: $a + b = b + \sigma_b(a)$. Furthermore λ is a semigroup homomorphism $(A, \circ) \rightarrow \text{Aut}(A, +)$, σ is a semigroup anti-homomorphism $(A, +) \rightarrow \text{End}(A, +)$,

and $\sigma_{\lambda_a(d)} \lambda_a(b) = \lambda_a \sigma_d(b)$, for all $a, b \in A$. The map λ is called the λ -map of the semitruss and the map σ is called the σ -map of the YB-semitruss.

We give a brief outline of the paper and mention parts of some of the main results. In Section 2, we set up the machinery of these YB-semitrusses and we show that this object determines all left non-degenerate solutions. In particular, in Corollary 2.14 we show the following result.

Theorem A. *Left non-degenerate solutions are restrictions of solutions associated to YB-semitrusses.*

Hence, the study of such solutions is being reduced to the study of YB-semitrusses. It turns out that a YB-semitruss is an epimorphic image of a structure semigroup $S(X, r) = M(X, r) \setminus \{1\}$ (considered as a YB-semitruss) of a left non-degenerate solution, where 1 is the identity of M . Further it is shown that all associative structures associated to left non-degenerate solutions that appear in the literature are special cases and correspond to YB-semitrusses with additional assumptions.

In Section 3, we characterize when the solution (A, r_A) associated to a YB-semitruss is bijective. In particular, we obtain the following result (Theorem 3.1).

Theorem B. *If (X, r) is a finite left non-degenerate solution, then r is bijective if and only if (X, r) is right non-degenerate.*

That the right non-degeneracy is a necessary condition already has been shown by Castelli, Catino and Stefanelli in [9]. Our proof also clarifies which data of the tuple defining a YB-semitruss controls specific properties of the associated solution.

In order to study solutions one would like techniques to reduce solutions to solutions of smaller size. For bijective non-degenerate solutions this for example has been successfully done via the retract of a solution (of a skew left brace); to do so one makes fundamentally use of the fact there are two group operations involved. In Section 4, we show that for non-degenerate solutions this can be done in general (Theorem 4.5 and Corollary 4.6).

Theorem C. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. Then $\mathcal{G}(A) = \{f(a) = (\sigma_a, \lambda_a, \rho_a) \mid a \in A\}$ is a YB-semitruss for the operations*

$$f(a) \circ f(b) = (\sigma_{a \circ b}, \lambda_{a \circ b}, \rho_{a \circ b}), \quad f(a) + f(b) = (\sigma_{a+b}, \lambda_{a+b}, \rho_{a+b}),$$

λ -map

$$\lambda_{f(a)} f(b) = (\sigma_{\lambda_a(b)}, \lambda_{\lambda_a(b)}, \rho_{\lambda_a(b)}) = f(\lambda_a(b)),$$

and σ -map

$$\sigma_{f(b)} f(a) = (\sigma_{\sigma_b(a)}, \lambda_{\sigma_b(a)}, \rho_{\sigma_b(a)}) = f(\sigma_b(a)).$$

The map $f : A \rightarrow \mathcal{G}(A) : a \mapsto f(a)$ is a YB-semitruss epimorphism and the associated solution of $\mathcal{G}(A)$ is non-degenerate.

Furthermore, $(\mathcal{G}(A), \circ)$ is a cancellative monoid (and thus also $(\mathcal{G}(A), +)$ is left cancellative) that satisfies the left and right Ore condition.

The YB-semitruss $\mathcal{G}(A)$ is thus “close” to being a left cancellative semi-brace as introduced by Catino, Colazzo and Stefanelli [10]. If this multiplicative semigroup is a group then it turns out that $\mathcal{G}(A)$ is a skew left brace and that the associated solution is bijective. In particular, it follows that if (X, r) is a non-degenerate solution so that for every $a \in M(X, r)$ there exists $b \in M(X, r)$ such that $\lambda_a \lambda_b = \text{id}$ and $\rho_b \rho_a = \text{id}$ (for example of all λ_a and ρ_b are of finite order) then r is bijective. This is another application of the machinery of YB-semitrusses.

Finally, in Section 5, we study the additive and multiplicative structure of a YB-semitruss A that is an epimorphic image of the structure monoid $M(X, r)$ of a finite left non-degenerate solution (X, r) . To do so we exploit the fact that in the finite semigroup $\mathcal{C}(A) := \{\sigma_a \mid a \in A\}$ every left ideal is a two-sided ideal. It

turns out that $(A, +)$ is a finite “module” over a subsemigroup that satisfies the ascending chain condition on left ideals and it has an ideal chain of which the Rees factors are finite unions of commutative semigroups. As a consequence we obtain the following results (Theorem 5.3 and Corollary 5.4).

Theorem D. *Let (X, r) be a finite left non-degenerate solution. Let K be a field. Then the algebra $K[(M(X, r), +)]$ is a left Noetherian PI-algebra of finite Gelfand-Kirillov dimension bounded by $|X|$.*

If, furthermore, the diagonal map $\mathfrak{q} : X \rightarrow X : x \mapsto \lambda_x^{-1}(x)$ is bijective then also the structure algebra $K[(M(X, r), \circ)]$ is a connected graded left Noetherian representable algebra.

Note that $K[M(X, r)]$ is a K -algebra defined by homogeneous quadratic relations. Such algebras have received a lot of interest in recent years in several contexts, such as arithmetical orders, classification of four dimensional noncommutative projective surfaces, Artin-Shelter regular algebras, regular languages and constructions of Noetherian algebras (see for example [22, 31, 42, 44, 65]). In case r is bijective, Theorem D has been proven in [39, 40]. It furthermore was shown that in this case $K[M(X, r)]$ is a prime algebra (or equivalently a domain) if and only if r is involutive. Hence showing that the algebraic structure of the algebra $K[M(X, r)]$ determines properties of the solution. For involutive finite solutions Gateva-Ivanova and Van den Bergh showed that homologically $K[M(X, r)]$ shares many properties with polynomial algebras in commuting variables. This rich algebraic structure of the algebra $K[M(X, r)]$ gives perspectives for developing and applying a variety of algebraic methods in the context of solutions of the Yang-Baxter equation.

With a YB-semitruss $(A, +, \circ, \lambda, \sigma)$ one associates another relevant solution, called the associated derived solution and which is denoted (A, s_A) . It is the associated solution of the YB-semitruss $(A, +, \circ, \iota, \sigma)$, where ι is the map $A \rightarrow \text{Aut}(A, +)$ defined by $\iota_a = \text{id}_A$. It is this structure that is fundamental in the proof of Theorem D.

We end the paper with describing the algebraic structure of YB-semitrusses for which $\mathcal{C}(A)$ is finite and has a principal ideal chain of length one, that is $\mathcal{C}(A)$ is a left simple semigroup. We show that their associated derived solutions are determined by finitely many bijective non-degenerate solutions. In particular we prove the following result (see Theorem 5.5).

Theorem E. *Let $A = (A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. Assume $\mathcal{C} = \mathcal{C}(A)$ is a finite left simple semigroup, let $E = E(\mathcal{C})$ be its set of idempotents and G_e the maximal subgroup of \mathcal{C} containing $e \in E$. Then, $A = \cup_{e \in E} A_e$, a disjoint union of left ideals $A_e = \sigma^{-1}(G_e)$, so that s_A restricts to a bijective non-degenerate solution on $G_e(A_e)$ and, for $a_e \in A_e$, $a_f \in A_f$, we have $s_A(a_e, a_f) = (a_f, \sigma_{f(a_f)}(f(a_e)))$, i.e. the derived solution is determined by bijective non-degenerate solutions.*

2. LEFT NON-DEGENERATE SOLUTIONS AND YB-SEMITRUSSES

In order to study the algebraic essence of several associative structures associated to solutions, Brzeziński in [8] introduced the notion of a left semitruss. A left semitruss is a quadruple $(A, +, \circ, \lambda)$ such that $(A, +)$ and (A, \circ) are non-empty semigroups and $\lambda : A \rightarrow \text{Map}(A, A) : a \mapsto \lambda_a$ is a function such that

$$a \circ (b + c) = (a \circ b) + \lambda_a(c), \quad (2.1)$$

for all $a, b, c \in A$. One calls $(A, +)$ the additive semigroup and (A, \circ) the multiplicative semigroup of the left semitruss and the relation (2.1) is called the left semitruss identity.

Since there is no natural construction to associate a (left non-degenerate) solution to an arbitrary semitruss, in [8, 25] special subclasses of left semi-trusses are considered that yield (left non-degenerate) solutions, but not every (left non-degenerate) solution can be obtained in this way. In order to resolve this for all left non-degenerate solutions, we define the following subclass. Strictly speaking these should be called left non-degenerate YB-left semitrusses, but for simplicity reasons we simply will call them YB-semitrusses. In Subsection 2.1 we will then show that they determine all left non-degenerate solutions.

Definition 2.1. A tuple $(A, +, \circ, \lambda, \sigma)$ is said to be a *YB-semitruss* if $(A, +, \circ, \lambda)$ is a left semitruss and $\sigma : A \rightarrow \text{Map}(A, A) : a \mapsto \sigma_a$ is map such that, for each $a, b, d \in A$, the following conditions are satisfied:

$$\lambda_a \in \text{Aut}(A, +) \text{ and } \lambda_a \lambda_b = \lambda_{a \circ b}, \quad (2.2)$$

$$a + \lambda_a(b) = a \circ b, \quad (2.3)$$

$$a + b = b + \sigma_b(a), \quad (2.4)$$

$$\sigma_a \in \text{End}(A, +) \text{ and } \sigma_{a+b} = \sigma_b \sigma_a, \quad (2.5)$$

$$\sigma_{\lambda_a(d)} \lambda_a(b) = \lambda_a \sigma_d(b). \quad (2.6)$$

The map $\lambda : (A, \circ) \rightarrow \text{Aut}(A, +) : a \mapsto \lambda_a$ is thus a semigroup homomorphism, it is called the λ -map of the semitruss and the map $\sigma : (A, +) \rightarrow \text{End}(A, +) : a \mapsto \sigma_a$ is a semigroup anti-homomorphism, called the σ -map of the YB-semitruss.

If $(A, +)$ is a semigroup and $\lambda : A \rightarrow \text{End}(A, +) : a \mapsto \lambda_a$ is a mapping so that $\lambda_a \lambda_b = \lambda_{a + \lambda_a(b)}$, for all $a, b \in A$, then (A, \circ) is a semigroup for the operation $a \circ b = a + \lambda_a(b)$ and A is a left semitruss. Clearly $(A, \circ) \rightarrow (A, +) \times \text{End}(A, +) : a \mapsto (a, \lambda_a)$ is a semigroup monomorphism. Of course one can transfer the addition on A to the image $\{(a, \lambda_a) \mid a \in A\}$, i.e. $(a, \lambda_a) + (b, \lambda_b) := (a + b, \lambda_{a+b})$. Hence, if $(A, +)$ is a semigroup then subsemigroups of the type $\{(a, \lambda_a) \mid a \in A\}$ of $(A, +) \times \text{End}(A, +)$ yield examples of left semitrusses and one has the stronger condition; $a \circ b = a + \lambda_a(b)$ (here we identify a with (a, λ_a)). Note that these have a similar flavour to regular subgroups of the holomorph of a group.

Examples of YB-semitrusses are regular subsemigroups of $(A, +) \times \text{Aut}(A, +)$ with $(A, +)$ an abelian semigroup (here each map $\sigma_a = \text{id}$). Specific examples of such semigroups are semigroups of I-type, studied by Gateva-Ivannova and Van den Bergh [34] and, more general, semigroups of IG-type, studied by Goffa, Jespers and Okniński in [36, 37]. Here $(A, +)$ is a free abelian semigroup, respectively an arbitrary cancellative abelian semigroup. It is shown in [34] that the former determine the involutive non-degenerate solutions; the latter have been studied in [36, 37] only from a pure algebraic point of view.

Now, we focus on the case in which the additive and the multiplicative structure of a YB-semitruss are monoids. Firstly, we prove that they share the same identity, say 1, then show that λ_1 and σ_1 are the identity map and $\lambda_a(1) = 1$ (for any $a \in A$). Later, we introduce the notion of unital YB-semitruss requiring in addition that $\sigma_a(1) = 1$ (for any $a \in A$). In Subsection 2.1 we show that structure monoids of solutions are examples of unital YB-semitruss.

Lemma 2.2. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss and assume $(A, +)$ and (A, \circ) are monoids. Then, condition (2.3) is equivalent with $(A, +)$ and (A, \circ) sharing the same identity, say 1. Moreover, $\lambda_1 = \sigma_1 = \text{id}$ and $\lambda_a(1) = 1$, for any $a \in A$.*

Proof. Assume that $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss such that $(A, +)$ and (A, \circ) are monoids. Let 0 denote the identity of $(A, +)$ and let 1 denote the identity of (A, \circ) . By (2.2), $\lambda_1 = \lambda_{1 \circ 1} = \lambda_1 \lambda_1$ and thus, because λ_1 is bijective, $\lambda_1 = \text{id}$. Hence, by (2.3), $0 = 1 \circ 0 = 1 + \lambda_1(0) = 1 + 0 = 1$. Conversely, assume that (A, \circ) and $(A, +)$

share the same identity, say 1. It then follows from the left semitruss identity that $a + \lambda_a(b) = a \circ 1 + \lambda_a(b) = a \circ (1 + b) = a \circ b$, as desired.

Moreover, for any $a \in A$, because of (2.4) we have $a = a + 1 = 1 + \sigma_1(a) = \sigma_1(a)$, i.e. $\sigma_1 = \text{id}$. Also, for any $a, b \in A$, we have $b + \lambda_a(1) = \lambda_a(\lambda_a^{-1}(b)) + \lambda_a(1) = \lambda_a(\lambda_a^{-1}(b) + 1) = \lambda_a(\lambda_a^{-1}(b)) = b$. This shows that $\lambda_a(1)$ is a right identity and thus $\lambda_a(1) = 1$. \square

Definition 2.3. A YB-semitruss $(A, +, \circ, \lambda, \sigma)$ is said to be *unital*, if $(A, +)$ and (A, \circ) are monoids and, for any $a \in A$,

$$\sigma_a(1) = 1, \quad (2.7)$$

where 1 denotes the identity in $(A, +)$ (and also in (A, \circ)).

Note that if $(A, +)$ is left cancellative monoid, with identity say 1, then (2.7) follows from the definition of YB-semitruss. Indeed $a + 1 = a = 1 + a \stackrel{(2.4)}{=} a + \sigma_a(1)$, and by left cancellativity, we get $1 = \sigma_a(1)$.

However, the above does not hold in general. Indeed, consider for example the multiplicative monoid $A = \{0, 1\}$. This can be made into a YB-semitruss by defining $(A, +) = (A, \circ)$ and $\lambda_1 = \lambda_0 = \text{id}_A$, $\sigma_1 = \text{id}$, and $\sigma_0(x) = 0$, for $x \in A$. Note that A is not a unital YB-semitruss since (2.7) does not holds in A because by definition $\sigma_0(1) = 0 \neq 1$.

Note that condition (2.4) implies, for all $a, b \in A$,

$$\sigma_a \sigma_b = \sigma_{\sigma_a(b)} \sigma_a. \quad (2.8)$$

Indeed, by (2.5) and (2.4), $\sigma_a \sigma_b = \sigma_{b+a} = \sigma_{a+\sigma_a(b)} = \sigma_{\sigma_a(b)} \sigma_a$.

Condition (2.3) links the multiplicative with the additive structure. Condition (2.4) implies that $A + b \subseteq b + A$, so right ideals in the semigroup $(A, +)$ are two-sided ideals. If, furthermore, each σ_b is bijective then $A + b = b + A$, i.e. every element of A is a normal element. As shown in [39] the latter is an essential property to describe the structure monoids of bijective non-degenerate solutions. Condition (2.5) can be avoided to associate a left non-degenerate solution to the considered YB-semitrusses (see Proposition 2.4). However, to deal with an arbitrary left non-degenerate solution we will show in Proposition 2.11 that, without loss of generality, we may assume that (2.5) holds.

Because of the equations (2.9), (2.10) and (2.11), equation (2.8) means that the map

$$s_A : A^2 \rightarrow A^2 : (a, b) \mapsto (b, \sigma_b(a)),$$

is a solution. We call it the *associated derived solution of the YB-semitruss* A (in the terminology of [49] one says that (A, \triangleleft) is a shelf, where $a \triangleleft b = \sigma_b(a)$, and equality (2.8) says that \triangleleft is self distributive, i.e. $(a \triangleleft b) \triangleleft d = (a \triangleleft d) \triangleleft (b \triangleleft d)$). Clearly s_A is bijective if and only if all σ_a are bijective, that is s_A is a non-degenerate bijective solution (or in the language of [45], (A, \triangleleft) is a rack). Note that s_A is the associated solution of the YB-semitruss $(A, +, +, \iota, \sigma)$, where ι is the map $A \rightarrow \text{Aut}(A, +)$ defined by $\iota_a = \text{id}_A$.

Recall (see for example [15]) that a mapping

$$r : X^2 \rightarrow X^2 : (x, y) \mapsto (\lambda_x(y), \rho_y(x)),$$

defines a solution if and only if, for all $a, b, c \in X$,

$$\lambda_a \lambda_b = \lambda_{\lambda_a(b)} \lambda_{\rho_b(a)}, \quad (2.9)$$

$$\lambda_{\rho_{\lambda_b(c)}(a)} \rho_c(b) = \rho_{\lambda_{\rho_b(a)}(c)} \lambda_a(b), \quad (2.10)$$

$$\rho_c \rho_b = \rho_{\rho_c(b)} \rho_{\lambda_b(c)}. \quad (2.11)$$

The meaning of the other requirements in Definition 2.1 are clarified in the following proposition. For this we introduce the so-called ρ -map of a left semitruss $(A, +, \circ, \lambda)$ for which there exists a map $\sigma : A \rightarrow \text{Map}(A, A)$ so that $a+b = b+\sigma_b(a)$, for all $a, b \in A$ (i.e. condition (2.4) holds) and $\lambda : (A, \circ) \rightarrow \text{Aut}(A, +) : a \mapsto \lambda_a$ is a homomorphism (for example A is a YB-semitruss). Then

$$\rho : A \rightarrow \text{Map}(A, A) : a \mapsto \rho_a,$$

with, for all $a, b \in A$,

$$\rho_b(a) = \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a),$$

is called its ρ -map.

We now prove a generalization of [20, Proposition 5.4]. The proof follows the same main idea as in [39, Proposition 2.2] and [62, Theorem 2.3].

Proposition 2.4. *Let $(A, +, \circ, \lambda)$ be a left semitruss and let $\sigma : A \rightarrow \text{Map}(A, A)$ be a map so that $a + b = b + \sigma_b(a)$, for all $a, b \in A$ (i.e. condition (2.4) holds). Assume that $\lambda : (A, \circ) \rightarrow \text{Aut}(A, +) : a \mapsto \lambda_a$ is a homomorphism and condition (2.3) is satisfied. Then*

$$r_A : A^2 \rightarrow A^2 : (a, b) \mapsto (\lambda_a(b), \rho_b(a)), \quad (2.12)$$

is a (left non-degenerate) solution if and only if $\sigma_{\lambda_a(b)} \lambda_a = \lambda_a \sigma_b$, for all $a, b \in A$ (i.e. condition (2.6) holds).

Furthermore, if $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss, then $r_A = \varphi^{-1} s_A \varphi$ where $\varphi : A^2 \rightarrow A^2 : (a, b) \mapsto (a, \lambda_a(b))$. In particular, r_A is bijective if and only if s_A is bijective, or equivalently s_A is (right) non-degenerate (i.e. all maps σ_a are bijective).

Proof. Let $J : A^3 \rightarrow A^3$ be the map defined by $J(a, b, d) = (a, \lambda_a(b), \lambda_a \lambda_b(d))$. Clearly J is bijective and $J^{-1}(a, b, d) = (a, \lambda_a^{-1}(b), \lambda_{\lambda_a^{-1}(b)}^{-1} \lambda_a^{-1}(d))$, for all $a, b, d \in A$.

We have

$$\begin{aligned} J^{-1}(s_A \times \text{id})J(a, b, d) &= J^{-1}(s_A \times \text{id})(a, \lambda_a(b), \lambda_a \lambda_b(d)) \\ &= J^{-1}(\lambda_a(b), \sigma_{\lambda_a(b)}(a), \lambda_a \lambda_b(d)) \\ &= (\lambda_a(b), \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a), \lambda_{\lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)}^{-1} \lambda_{\lambda_a(b)}^{-1} \lambda_a \lambda_b(d)) \\ &= (\lambda_a(b), \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a), \lambda_{\lambda_a(b) \circ \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)}^{-1} \lambda_{a \circ b}(d)) \\ &= (\lambda_a(b), \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a), \lambda_{a \circ b}^{-1} \lambda_{a \circ b}(d)) \\ &= (\lambda_a(b), \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a), d) \\ &= (\lambda_a(b), \rho_b(a), d), \end{aligned}$$

since $\lambda_a(b) \circ \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a) = \lambda_a(b) + \sigma_{\lambda_a(b)}(a) = a + \lambda_a(b) = a \circ b$. Hence, $J^{-1}(s_A \times \text{id})J = r_A \times \text{id}$. Moreover, if condition (2.6) holds then

$$\begin{aligned} J^{-1}(\text{id} \times s_A)J(a, b, d) &= J^{-1}(\text{id} \times s_A)(a, \lambda_a(b), \lambda_a \lambda_b(d)) \\ &= J^{-1}(a, \lambda_a \lambda_b(d), \sigma_{\lambda_a \lambda_b(d)} \lambda_a(b)) \\ &= (a, \lambda_a^{-1} \lambda_a \lambda_b(d), \lambda_{\lambda_a^{-1} \lambda_a \lambda_b(d)}^{-1} \lambda_a^{-1} \sigma_{\lambda_a \lambda_b(d)} \lambda_a(b)) \\ &= (a, \lambda_b(d), \lambda_{\lambda_b(d)}^{-1} \lambda_a^{-1} \lambda_a \sigma_{\lambda_b(d)}(b)) \\ &= (a, \lambda_b(d), \lambda_{\lambda_b(d)}^{-1} \sigma_{\lambda_b(d)}(b)) \\ &= (a, \lambda_b(d), \rho_d(b)), \end{aligned}$$

yields $J^{-1}(\text{id} \times s_A)J = \text{id} \times r_A$. Because (A, s_A) is a solution (as mentioned earlier the condition (2.6) is not needed for this) we get that (A, r_A) is a solution. This proves the necessity of condition (2.6).

Conversely, assume that (A, r_A) is a solution. As $\sigma_b(a) = \lambda_b \rho_{\lambda_a^{-1}(b)}(a)$ we get that

$$\begin{aligned}
\sigma_{\lambda_a(d)}(\lambda_a(b)) &= \lambda_{\lambda_a(d)} \rho_{\lambda_{\lambda_a^{-1}(b)}(d)} \lambda_a(b) \\
&= \lambda_{\lambda_a(d)} \rho_{\lambda_{\rho_b(a)} \lambda_b^{-1}(d)} \lambda_a(b) && \text{by (2.9)} \\
&= \lambda_{\lambda_a(d)} \lambda_{\rho_{\lambda_b \lambda_b^{-1}(d)}(a)} \rho_{\lambda_b^{-1}(d)}(b) && \text{by (2.10)} \\
&= \lambda_{\lambda_a(d)} \lambda_{\rho_d(a)} \rho_{\lambda_b^{-1}(d)}(b) \\
&= \lambda_a \lambda_d \rho_{\lambda_b^{-1}(d)}(b) && \text{by (2.9)} \\
&= \lambda_a \sigma_d(b).
\end{aligned}$$

Hence, $\lambda_a \sigma_d(b) = \sigma_{\lambda_a(d)}(\lambda_a(b))$.

To finish the proof it is enough to observe that $s_A = \varphi r_A \varphi^{-1}$. \square

The previous proposition states that a left non-degenerate solution r_A can be determined by a conjugate of the derived solution s_A , by a mapping say φ . However, φ is not an arbitrary permutation of A^2 . In the case of bijective solutions it has to be a so-called Drinfeld twist on solutions, see for example [35]. One can extend the latter definition to arbitrary left non-degenerate solutions and with the same strategy as in [35, Example 2.1] one proves that $r_A = \varphi^{-1} s_A \varphi$ for some permutation φ of A^2 so that (φ, Ψ, Φ) is a Drinfeld twist on the solution (A, r_A) .

For a YB-semitruss A we call r_A the *associated solution*. It is said to be the *trivial solution* if $r_A(a, b) = (b, a)$ for all $a, b \in A$. It turns out that if an associated solution is bijective then its inverse is the associated solution of a well-described associated YB-semitruss.

For a semigroup $(A, +)$ we denote by $(A, +^{op})$ its opposite semigroup, i.e. $a +^{op} b = b + a$, for $a, b \in A$.

Proposition 2.5. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss with σ_a bijective for any $a \in A$, that is, the associated solution r_A is bijective. Then,*

$$r_A^{-1}(a, b) = (\sigma_a^{-1} \lambda_a(b), \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)),$$

and $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$ is a YB-semitruss, with $\bar{\lambda}_a = \sigma_a^{-1} \lambda_a$ and $\bar{\sigma}_a = \sigma_a^{-1}$, and its associated solution is r_A^{-1} . We call $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$ the opposite YB-semitruss.

In particular, for a YB-semitruss $(A, +, \circ, \lambda, \sigma)$, the associated solution r_A is involutive if and only if the associated derived solution s_A is trivial (that is, all maps $\sigma_a = \text{id}$).

Proof. Put $r = r_A$. So, $r(a, b) = (\lambda_a(b), \lambda_{\lambda_a^{-1}(b)}^{-1} \sigma_{\lambda_a(b)}(a))$. Moreover, put $r'(a, b) = (\sigma_a^{-1} \lambda_a(b), \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a))$. Then

$$\begin{aligned}
rr'(a, b) &= r(\sigma_a^{-1} \lambda_a(b), \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)) \\
&= (\lambda_{\sigma_a^{-1} \lambda_a(b)} \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a), \lambda_{\lambda_{\sigma_a^{-1} \lambda_a(b)} \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)} \sigma_{\lambda_{\sigma_a^{-1} \lambda_a(b)} \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)} \sigma_a^{-1} \lambda_a(b)) \\
&= (a, \lambda_a^{-1} \sigma_a \sigma_a^{-1} \lambda_a(b)) = (a, b),
\end{aligned}$$

and

$$\begin{aligned}
r'r(a, b) &= r'(\lambda_a(b), \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)) \\
&= (\sigma_{\lambda_a(b)}^{-1} \lambda_{\lambda_a(b)} \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a), \lambda_{\sigma_{\lambda_a(b)}^{-1} \lambda_{\lambda_a(b)} \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)}^{-1} \lambda_a(b)) \\
&= (a, b).
\end{aligned}$$

So indeed $r' = r^{-1}$.

Now, assume that $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$, with $\bar{\lambda}_a = \sigma_a^{-1} \lambda_a$ and $\bar{\sigma}_a = \sigma_a^{-1}$, is a YB-semitruss. The associated solution to such a YB-semitruss is given by

$$\begin{aligned}
\bar{r}(a, b) &= (\bar{\lambda}_a(b), \bar{\lambda}_{\bar{\lambda}_a(b)}^{-1} \bar{\sigma}_{\bar{\lambda}_a(b)}(a)) \\
&= (\sigma_a^{-1} \lambda_a(b), \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1} \sigma_{\sigma_a^{-1} \lambda_a(b)} \sigma_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)) \\
&= (\sigma_a^{-1} \lambda_a(b), \lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)) \\
&= r^{-1}(a, b).
\end{aligned}$$

So it remains to prove that $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$ is a YB-semitruss. Clearly $\bar{\lambda}_a \in \text{Aut}(A, +^{op})$ for each $a \in A$. That $(A, \circ) \rightarrow \text{Aut}(A, +^{op}) : a \mapsto \bar{\lambda}_a$ is a homomorphism follows from the following:

$$\begin{aligned}
\bar{\lambda}_a \bar{\lambda}_b(d) &= \sigma_a^{-1} \lambda_a \sigma_b^{-1} \lambda_b(d) \\
&= \sigma_a^{-1} \sigma_{\lambda_a(b)}^{-1} \lambda_a \lambda_b(d) && \text{by (2.6)} \\
&= (\sigma_{\lambda_a(b)} \sigma_a)^{-1} \lambda_a \lambda_b(d) && \text{by (2.2)} \\
&= \sigma_{a+\lambda_a(b)}^{-1} \lambda_a \lambda_b(d) && \text{by (2.5)} \\
&= \sigma_{a \circ b}^{-1} \lambda_a \lambda_b(d) && \text{by (2.3)} \\
&= \bar{\lambda}_{a \circ b}(d).
\end{aligned}$$

Also, for each $a \in A$, it is clear that $\bar{\sigma}_a \in \text{Aut}(A, +^{op})$ and thus also that $(A, +^{op}) \rightarrow (\text{Map}(A, A), \circ) : a \mapsto \bar{\sigma}_a$ is semigroup anti-homomorphism.

Since $(A, +, \circ, \lambda, \sigma)$ satisfies (2.4), we have that $a + b = b + \sigma_b(a)$ for all $a, b \in A$. Hence, also $\sigma_b^{-1}(a) + b = b + a$, or equivalently, $b +^{op} \sigma_b^{-1}(a) = a +^{op} b$, i.e. (2.4) holds for $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$. Since $(A, +, \circ, \lambda, \sigma)$ satisfies equation (2.3), we obtain

$$\begin{aligned}
a +^{op} \bar{\lambda}_a(b) &= a +^{op} \sigma_a^{-1} \lambda_a(b) \\
&= a +^{op} \bar{\sigma}_a \lambda_a(b) \\
&= \lambda_a(b) +^{op} a && \text{(by (2.4) for } (A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})) \\
&= a + \lambda_a(b) \\
&= a \circ b.
\end{aligned}$$

So, (2.3) holds for $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$. Finally, (2.6) yields $\sigma_{\lambda_a(b)} \lambda_a = \lambda_a \sigma_b$ and thus $\lambda_a \sigma_b^{-1} = \sigma_{\lambda_a(b)}^{-1} \lambda_a$. Hence, $\sigma_a^{-1} \lambda_a \sigma_b^{-1} = \sigma_a^{-1} \sigma_{\lambda_a(b)}^{-1} \lambda_a$. Because of (2.8) we know that $\sigma_a^{-1} \sigma_{\lambda_a(b)}^{-1} = \sigma_{\sigma_a^{-1} \lambda_a(b)}^{-1} \sigma_a^{-1}$. Hence, $\bar{\lambda}_a \sigma_b^{-1} = \sigma_{\bar{\lambda}_a(b)}^{-1} \bar{\lambda}_a$. So, $(A, +^{op}, \circ, \bar{\lambda}, \bar{\sigma})$ is a YB-semitruss. \square

The λ and ρ maps provide relations in YB-semitrusses.

Proposition 2.6. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. Then for all $a, b \in A$ and for the λ -map and ρ -map we have*

$$a \circ b = \lambda_a(b) \circ \rho_b(a), \quad (2.13)$$

$$\lambda_b(a \circ d) = \lambda_b(a) \circ \lambda_{\rho_a(b)}(d), \quad (2.14)$$

$$\rho_b(d \circ a) = \rho_{\lambda_a(b)}(d) \circ \rho_b(a). \quad (2.15)$$

Proof. Let $a, b, d \in A$. Then,

$$\begin{aligned}\lambda_b(b) \circ \rho_b(a) &= \lambda_a(b) \circ \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a) \\ &= \lambda_a(b) + \sigma_{\lambda_a(b)}(a) = a + \lambda_a(b) = a \circ b.\end{aligned}$$

Hence, we get

$$\begin{aligned}\lambda_b(a \circ d) &= \lambda_b(a + \lambda_a(d)) = \lambda_b(a) + \lambda_b \lambda_a(d) \\ &= \lambda_b(a) + \lambda_{b \circ a}(d) = \lambda_b(a) + \lambda_{\lambda_b(a) \circ \rho_a(b)}(d) \\ &= \lambda_b(a) + \lambda_{\lambda_b(a)} \lambda_{\rho_a(b)}(d) \\ &= \lambda_b(a) \circ \lambda_{\rho_a(b)}(d).\end{aligned}$$

To prove (2.15) we first claim that

$$\rho_{\lambda_{\lambda_a^{-1}(b)}^{-1} \lambda_a^{-1}(d)} \lambda_a^{-1}(b) = \lambda_{\rho_{\lambda_a^{-1}(d)}^{-1}(a)}^{-1} \rho_{\lambda_b^{-1}(d)}(b). \quad (2.16)$$

Indeed by setting in (2.10) $c = \lambda_y^{-1}(t)$, $b = y$ and $a = z$, we get

$$\lambda_{\rho_{\lambda_y(\lambda_y^{-1}(t))}(z)} \rho_{\lambda_y^{-1}(t)}(y) = \rho_{\lambda_{\rho_y(z)} \lambda_y^{-1}(t)} \lambda_z(y).$$

With $t = \lambda_z^{-1}(u)$ and $y = \lambda_z^{-1}(v)$ and because of (2.9) this yields

$$\begin{aligned}\lambda_{\rho_{\lambda_z^{-1}(u)}(z)} \rho_{\lambda_z^{-1}(v)} \lambda_z^{-1}(u) \lambda_z^{-1}(v) &= \rho_{\lambda_{\rho_{\lambda_z^{-1}(v)}(z)} \lambda_z^{-1}(u)} \lambda_z \lambda_z^{-1}(v) \\ &= \rho_{\lambda_v^{-1}(u)}(v),\end{aligned}$$

as desired.

Moreover, we notice that

$$\sigma_a(b + d) = \lambda_a \rho_{\lambda_{b+d}^{-1}(a)}(b + d) = \lambda_a \rho_{\lambda_{b \circ \lambda_b^{-1}(d)}^{-1}(a)}(b \circ \lambda_b^{-1}(d)),$$

and

$$\begin{aligned}\sigma_a(b) + \sigma_a(d) &= \sigma_a(b) \circ \lambda_{\sigma_a(b)}^{-1}(\sigma_a(d)) \\ &= \lambda_a(\rho_{\lambda_b^{-1}(a)}(b)) \circ \lambda_{\lambda_a \rho_{\lambda_b^{-1}(a)}(b)}^{-1}(\lambda_a(\rho_{\lambda_d^{-1}(a)}(d))) \\ &= \lambda_a(\rho_{\lambda_b^{-1}(a)}(b)) \circ \lambda_{\rho_{\lambda_b^{-1}(a)}(b)(a)}(\lambda_{\rho_{\lambda_b^{-1}(a)}(b)}^{-1}(\rho_{\lambda_d^{-1}(a)}(d))) && \text{by (2.9)} \\ &= \lambda_a(\rho_{\lambda_b^{-1}(a)}(b)) \circ \lambda_{\rho_{\lambda_b^{-1}(a)}(b)(a)}(\rho_{\lambda_b^{-1}(d)} \lambda_b^{-1}(a)(\lambda_b^{-1}(d))) && \text{by (2.16)} \\ &= \lambda_a(\rho_{\lambda_b^{-1}(a)}(b) \circ \rho_{\lambda_{\lambda_b^{-1}(d)}^{-1}(a)} \lambda_b^{-1}(a)(\lambda_b^{-1}(d))) && \text{by (2.14)}.\end{aligned}$$

As σ_a is an additive homomorphism and λ_a is bijective, we thus obtain that

$$\rho_{\lambda_{b \circ \lambda_b^{-1}(d)}^{-1}(a)}(b \circ \lambda_b^{-1}(d)) = \rho_{\lambda_b^{-1}(a)}(b) \circ \rho_{\lambda_{\lambda_b^{-1}(d)}^{-1}(a)} \lambda_b^{-1}(a)(\lambda_b^{-1}(d)).$$

Put $e = \lambda_b^{-1}(d)$ and $x = \lambda_{b \circ e}^{-1}(a)$. Then the previous equality becomes, for arbitrary $x, b, e \in A$,

$$\rho_x(b \circ e) = \rho_{\lambda_b^{-1} \lambda_{b \circ e}(x)}(b) \circ \rho_x(e) = \rho_{\lambda_e(x)}(b) \circ \rho_x(e),$$

as desired. \square

The requirement that the σ -map is an anti-homomorphism implies that the ρ -map also is an anti-homomorphism.

Lemma 2.7. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss with ρ -map $\rho_b(a) = \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)$. Then $\rho : (A, \circ) \rightarrow \text{Map}(A, A) : a \mapsto \rho_a$ is an anti-homomorphism.*

Proof. Let $a, b \in A$. By (2.14) and because the σ -map is an anti-homomorphism, we get for all $x \in A$ that

$$\begin{aligned}
\rho_{b \circ a}(x) &= \lambda_{\lambda_x(b \circ a)}^{-1} \sigma_{\lambda_x(b \circ a)}(x) = \lambda_{\lambda_x(b) \circ \lambda_{\rho_b(x)}(a)}^{-1} \sigma_{\lambda_x(b + \lambda_b(a))}(x) \\
&= \lambda_{\lambda_{\rho_b(x)}(a)}^{-1} \lambda_{\lambda_x(b)}^{-1} \sigma_{\lambda_x(b) + \lambda_x \lambda_b(a)}(x) \\
&= \lambda_{\lambda_{\rho_b(x)}(a)}^{-1} \lambda_{\lambda_x(b)}^{-1} \sigma_{\lambda_x \lambda_b(a)} \sigma_{\lambda_x(b)}(x) \\
&= \lambda_{\lambda_{\rho_b(x)}(a)}^{-1} \sigma_{\lambda_{\lambda_x(b)}^{-1} \lambda_x \lambda_b(a)} \lambda_{\lambda_x(b)}^{-1} \sigma_{\lambda_x(b)}(x) \\
&= \lambda_{\lambda_{\rho_b(x)}(a)}^{-1} \sigma_{\lambda_{\rho_b(x)}(a)} \rho_b(x) \\
&= \rho_a \rho_b(x).
\end{aligned}$$

Hence, $\rho_{b \circ a}(x) = \rho_a \rho_b(x)$, as desired. \square

In fact, if we do not assume that a YB-semitruss $(A, +, \circ, \lambda, \sigma)$ satisfies $\sigma_{a+b} = \sigma_b \sigma_a$ for all $a, b \in A$, then we can prove that this condition follows from the ρ -map being an anti-homomorphism. Indeed as $\sigma_b(a) = \lambda_b \rho_{\lambda_x^{-1}(b)}(a)$ we get that, for any $a, b, x \in A$, using (2.9), (2.10) and that the ρ -map is an anti-homomorphism,

$$\begin{aligned}
\sigma_{a+b}(x) &= \lambda_{a+b} \rho_{\lambda_x^{-1}(a+b)}(x) \\
&= \lambda_a \lambda_{\lambda_x^{-1}(b)} \rho_{\lambda_x^{-1}(a) + \lambda_x^{-1}(b)}(x) \\
&= \lambda_b \lambda_{\rho_{\lambda_x^{-1}(b)}(a)} \rho_{\lambda_x^{-1}(a) \circ \lambda_{\lambda_x^{-1}(a)}^{-1}(\lambda_x^{-1}(b))}(x) \\
&= \lambda_b \lambda_{\rho_{\lambda_x^{-1}(b)}(a)} \rho_{\lambda_x^{-1}(a)}(\lambda_x^{-1}(b)) \rho_{\lambda_x^{-1}(a)}(x) \\
&= \lambda_b \lambda_{\rho_{\lambda_x^{-1}(b)}(a)} \rho_{\lambda_{\lambda_x^{-1}(a)}^{-1}(x)}(\lambda_x^{-1}(b)) \rho_{\lambda_x^{-1}(a)}(x) \\
&= \lambda_b \lambda_{\rho_{\lambda_{\rho_{\lambda_x^{-1}(a)}(x)}(\lambda_x^{-1}(a))}(\lambda_x^{-1}(b))} \rho_{\lambda_{\lambda_x^{-1}(a)}^{-1}(x)}(\lambda_x^{-1}(b)) (\rho_{\lambda_x^{-1}(a)}(x)) \\
&= \lambda_b \rho_{\lambda_{\rho_{\lambda_x^{-1}(a)}(x)}(a)} (\lambda_{\rho_{\lambda_x^{-1}(a)}^{-1}(x)} \lambda_x^{-1}(b)) \lambda_a (\rho_{\lambda_x^{-1}(a)}(x)) \\
&= \lambda_b \rho_{\lambda_{\lambda_a \rho_{\lambda_x^{-1}(a)}(x)}(a)} \lambda_a \lambda_x^{-1}(b) \lambda_a (\rho_{\lambda_x^{-1}(a)}(x)) \\
&= \lambda_b \rho_{\lambda_{\lambda_a \rho_{\lambda_x^{-1}(a)}(x)}(a)}(b) \lambda_a (\rho_{\lambda_x^{-1}(a)}(x)) \\
&= \sigma_b (\lambda_a \rho_{\lambda_x^{-1}(a)}(x)) \\
&= \sigma_b \sigma_a(x).
\end{aligned}$$

Hence, the map $\sigma : (A, +) \rightarrow \text{Map}(A, A) : a \mapsto \sigma_a$ is an anti-homomorphism.

Obviously, a homomorphism of YB-semitrusses should be a mapping that preserves the operations and is compatible with the λ -map and the σ -map.

Definition 2.8. A mapping $f : A \rightarrow B$ between YB-semitrusses $(A, +, \circ, \lambda, \sigma)$ and $(B, +', \circ', \lambda', \sigma')$ is said to be a *homomorphism* if

- (1) $f(a + b) = f(a) +' f(b)$, $f(a \circ b) = f(a) \circ' f(b)$,
- (2) $f(\lambda_a(b)) = \lambda'_{f(a)}(f(b))$ and $f(\sigma_b(a)) = \sigma'_{f(b)}(f(a))$, for all $a, b \in A$.

Note that a homomorphism f of YB-semitrusses also is compatible with the inverses of the λ -maps, i.e. $\lambda'_{f(a)}^{-1} f = f \lambda_a^{-1}$.

Lemma 2.9. *Let $f : A \rightarrow B$ be an epimorphism of YB-semitrusses $(A, +, \circ, \lambda, \sigma)$ and $(B, +', \circ', \lambda', \sigma')$. Then $\rho'_{f(a)} f(b) = f \rho_a(b)$ for all $a, b \in A$, i.e. the ρ -map is compatible with f .*

If, for all $a \in A$, ρ_a is surjective then for all $x \in B$, ρ'_x is surjective.

If, for all $a \in A$, ρ_a is bijective and the mapping $B \rightarrow B : f(b) \mapsto f\rho_a^{-1}(b)$ is well-defined, that is $f\rho_a^{-1}(b) = f\rho_{a'}^{-1}(b')$ for any $a, a', b, b' \in A$ such that $f(a) = f(a')$ and $f(b) = f(b')$, then all $(\rho')_{f(a)}$ are bijective and $(\rho')_{f(a)}^{-1}f = f\rho_a^{-1}$.

Proof. By definition we have that $\rho_b(a) = \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)$ and $\rho'_y(x) = (\lambda')_{\lambda'_x(y)}^{-1} \sigma'_{\lambda'_x(y)}(x)$, for all $a, b \in A, x, y \in B$. It follows that

$$\begin{aligned} \rho'_{f(b)}f(a) &= (\lambda')_{\lambda'_{f(a)}f(b)}^{-1} \sigma'_{\lambda'_{f(a)}f(b)} f(a) = (\lambda')_{f(\lambda_a(b))}^{-1} \sigma'_{f(\lambda_a(b))} f(a) \\ &= (\lambda')_{f(\lambda_a(b))}^{-1} f(\sigma_{\lambda_a(b)}(a)) = f(\lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)) \\ &= f\rho_b(a), \end{aligned}$$

for any $a, b \in A$. This proves the first part. Obviously this implies that $\rho'_{f(b)}$ is surjective if both f and ρ_b are surjective. Hence the second part also follows.

For the third part, assume ρ_a bijective, for any $a \in A$, and that the mapping $g : B \rightarrow B : f(b) \mapsto f\rho_a^{-1}(b)$ is well-defined. Then

$$g\rho'_{f(a)}f(b) = gf\rho_a(b) = f\rho_a^{-1}\rho_a(b) = f(b),$$

and

$$\rho'_{f(a)}gf(b) = \rho'_{f(a)}f\rho_a^{-1}(b) = f\rho_a\rho_a^{-1}(b) = f(b).$$

Therefore $\rho'_{f(a)}$ is bijective with inverse g . \square

Recall that a solution (X', r') is said to be an *epimorphic image* of a solution (X, r) if there exists a surjective map $f : X \rightarrow X'$ so that $(f \times f)r = r'(f \times f)$ on X^2 . If, moreover, f is bijective then one says that the solutions are isomorphic.

Proposition 2.10. *Let $(A, +, \circ, \lambda, \sigma)$ and $(A', +', \circ', \lambda', \sigma')$ be YB-semitrusses. Then the associated solution $r_{A'}$ is an epimorphic (respectively isomorphic) image of the associated solution r_A if and only if there exists a surjective (respectively bijective) map $f : A \rightarrow A'$ such that $f\lambda_a = \lambda'_{f(a)}f$ and $f\sigma_a = \sigma'_{f(a)}f$, for any $a \in A$ (that is condition (2) of Definition 2.8 holds).*

Proof. Assume that $r_{A'}$ is an epimorphic image of r_A . Then there exists a surjective map $f : A \rightarrow A'$ such that $(f \times f)r_A = r_{A'}(f \times f)$. It follows that

$$(f\lambda_a(b), f\lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)) = (\lambda'_{f(a)}f(b), (\lambda')_{\lambda'_{f(a)}f(b)}^{-1} \sigma'_{\lambda'_{f(a)}f(b)}(f(a))).$$

Hence, in particular $f\lambda_a(b) = \lambda'_{f(a)}f(b)$ for every $a, b \in A$. It yields $f\lambda_a = \lambda'_{f(a)}f$ and, since λ_a and $\lambda'_{f(a)}$ are bijective, $f\lambda_a^{-1} = (\lambda')_{f(a)}^{-1}f$. Hence, we get

$$\begin{aligned} (\lambda')_{\lambda'_{f(a)}f(b)}^{-1} \sigma'_{\lambda'_{f(a)}f(b)}(f(a)) &= f\lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a) \\ &= (\lambda')_{f\lambda_a(b)}^{-1} f\sigma_{\lambda_a(b)}(a) \\ &= (\lambda')_{\lambda'_{f(a)}f(b)}^{-1} f\sigma_{\lambda_a(b)}(a). \end{aligned}$$

It follows that $f\sigma_{\lambda_a(b)}(a) = \sigma'_{\lambda'_{f(a)}f(b)}(f(a)) = \sigma'_{f\lambda_a(b)}f(a)$, for every $a, b \in A$. Hence $f\sigma_b(a) = \sigma'_{f(b)}f(a)$.

Conversely, assume that $f\lambda_a = \lambda'_{f(a)}f$ and $f\sigma_a = \sigma'_{f(a)}f$, for any $a \in A$. Then

$$\begin{aligned} r_{A'}(f \times f)(a, b) &= (\lambda'_{f(a)}f(b), (\lambda')_{\lambda'_{f(a)}f(b)}^{-1} \sigma'_{\lambda'_{f(a)}f(b)} f(a)) \\ &= (f\lambda_a(b), (\lambda')_{f\lambda_a(b)}^{-1} \sigma'_{f\lambda_a(b)} f(a)) \\ &= (f\lambda_a(b), (\lambda')_{f\lambda_a(b)}^{-1} f\sigma_{\lambda_a(b)}(a)) \\ &= (f\lambda_a(b), f\lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)) \\ &= (f \times f)r_A(a, b). \end{aligned}$$

Hence the result follows. \square

So, isomorphic YB-semitrusses yield isomorphic associated solutions. The converse however is not true, take for example two non-isomorphic abelian groups of the same size $(G, +)$ and (H, \cdot) . Then $(G, +, +, \iota, \sigma)$ with $\iota_g = \sigma_g = \text{id}_G$ for all $g \in G$, and $(H, \cdot, \cdot, \iota, \sigma')$ with $\iota_h = \sigma'_h = \text{id}_H$ for all $h \in H$, are non-isomorphic YB-semitrusses with isomorphic associated solutions.

2.1. Structure monoids as YB-semitrusses.

In this section we show that all left non-degenerate solutions are determined by YB-semitrusses. We do this by putting a YB-semitruss structure on the structure monoid associated to a left non-degenerate solution, by specifically defining the σ -map using the derived solution. At the end of this section we show that isomorphic solutions have isomorphic (unital) structure YB-semitrusses and vice versa.

We begin by recalling results of Gateva-Ivanova and Majid [33] and Cedó, Jespers and Verwimp [20] using the same notation as in [20]. For completeness' sake we include all the necessary background. The notation $\langle X \rangle$ is used to denote the monoid generated by the set X .

Let (X, r) be a solution. As before, write $r(x, y) = (\lambda_x(y), \rho_y(x))$, for $x, y \in X$. The (left) derived monoid $A(X, r)$ of (X, r) is defined as the monoid with the following presentation

$$A = A(X, r) = \langle x \in X \mid x + \lambda_x(y) = \lambda_x(y) + \lambda_{\lambda_x(y)}\rho_y(x), x, y \in X \rangle,$$

and the structure monoid $M(X, r)$ of (X, r) is the monoid with presentation

$$M = M(X, r) = \langle x \in X \mid x \circ y = \lambda_x(y) \circ \rho_y(x), x, y \in X \rangle.$$

The subsemigroup $M \setminus \{1\}$ is called the structure semigroup of (X, r) and is denoted by $S(X, r)$. In [39, Proposition 2.2] it is shown that if (X, r) is left non-degenerate then

$$s : X^2 \rightarrow X^2 : (x, y) \mapsto (y, \sigma_y(x)),$$

also is a solution, called the derived solution of (X, r) , where

$$\sigma_y(x) = \lambda_y \rho_{\lambda_x^{-1}(y)}(x). \quad (2.17)$$

Note that in this case $x + y = y + \sigma_y(x)$ for all $x, y \in X$ (explaining the use of the notation σ_y in (2.17)), and furthermore, s is bijective if and only if r is bijective (or equivalently, s is right non-degenerate).

It was shown by Gateva-Ivanova and Majid [33, Theorem 3.6] that for an arbitrary solution (X, r) the maps λ and ρ can be extended uniquely to M (for simplicity we use the same notation for the extension) so that one obtains a monoid homomorphism $\lambda : (M, \circ) \rightarrow \text{Map}(M, M) : a \mapsto \lambda_a$, and a monoid anti-homomorphism $\rho : (M, \circ) \rightarrow \text{Map}(M, M) : a \mapsto \rho_a$. Furthermore, the map $r_M : M \times M \rightarrow M \times M$, defined by $r_M(a, b) = (\lambda_a(b), \rho_b(a))$, for all $a, b \in M$ is a solution. For $a, b \in M$ we have

$$a \circ b = \lambda_a(b) \circ \rho_b(a), \quad (2.18)$$

$$\rho_b(c \circ a) = \rho_{\lambda_a(b)}(c) \circ \rho_b(a), \quad (2.19)$$

$$\lambda_b(a \circ c) = \lambda_b(a) \circ \lambda_{\rho_a(b)}(c). \quad (2.20)$$

Also, r is left (respectively right) non-degenerate if and only if r_M is left (respectively right) non-degenerate; and r is bijective if and only if r_M is bijective. Note that the derived solution of (M, r_M) of a left non-degenerate solution is the solution

$$s_M : M^2 \rightarrow M^2 : (a, b) \mapsto (a, \sigma_b(a)),$$

where $\sigma_a(b) = \lambda_a \rho_{\lambda_b^{-1}(a)}(b)$. From the above we know that r_M is bijective if and only if s_M is bijective.

Actually one obtains more as shown by Cedó, Jespers and Verwimp in [20, Proposition 3.1]. Each λ induces a monoid homomorphism

$$\lambda' : (M, \circ) \rightarrow \text{End}(A, +) : a \mapsto \lambda'_a,$$

with $\lambda'_x(y) = \lambda_x(y)$, for $x, y \in X$.

In [20, Proposition 3.2] it is shown that there exists a 1-cocycle

$$\pi : M \rightarrow A,$$

with respect to the homomorphism λ' , such that $\pi(x) = x$ for all $x \in X$, and hence one obtains a homomorphism

$$f : M \rightarrow A \rtimes \text{Im}(\lambda') : m \mapsto (\pi(m), \lambda'_m).$$

Furthermore, if r is left non-degenerate then it follows from Proposition 3.6 and Proposition 3.7 in [20] that π is bijective and thus

$$M \cong \{(\pi(m), \lambda'_m) \mid m \in M\} \subseteq A \rtimes \text{Im}(\lambda').$$

For left non-degenerate solutions we will identify $M = M(X, r)$ with $A = A(X, r)$ via the mapping π and we thus denote λ' (respectively ρ') by λ (respectively ρ). Hence M also has an additive structure $(M, +)$ and we have $a \circ b = a + \lambda_a(b)$ and $a + b = b + \sigma_b(a)$ for all $a, b \in A$. Also,

$$\begin{aligned} \lambda_a \sigma_d(b) &= \lambda_a \lambda_d \rho_{\lambda_b^{-1}(d)}(b) \\ &= \lambda_{\lambda_a(d)} \lambda_{\rho_d(a)} \rho_{\lambda_b^{-1}(d)}(b) \\ &= \lambda_{\lambda_a(d)} \lambda_{\rho_{\lambda_b(\lambda_b^{-1}(d))}(a)} \rho_{\lambda_b^{-1}(d)}(b) \\ &= \lambda_{\lambda_a(d)} \rho_{\lambda_{\rho_b(a)}(\lambda_b^{-1}(d))} \lambda_a(b) \\ &= \lambda_{\lambda_a(d)} \rho_{\lambda_{\lambda_a(b)}^{-1}(\lambda_a(d))} \lambda_a(b) \\ &= \sigma_{\lambda_a(d)} \lambda_a(b). \end{aligned}$$

So equation (2.6) holds. Moreover, because of equation (2.19) one can show (with a reversed argument to that in the last part of the proof of Proposition 2.6) that each $\sigma_a \in \text{End}(M, +)$. Finally, since ρ is an anti-homomorphism we can prove that $\sigma_{a+b} = \sigma_b \sigma_a$, for all $a, b \in M(X, r)$.

To summarize we have shown the following.

Proposition 2.11. *Let (X, r) be a left non-degenerate solution. The structure monoid $M = M(X, r)$ is a unital YB-semitruss $(M, +, \circ, \lambda, \sigma)$ (with λ -map and σ -map as defined above) and with the structure semigroup $S(X, r)$ as a sub-YB-semitruss (not unital). The associated solution of the unital structure YB-semitruss M (respectively associated derived solution of the YB-semitruss M) (as defined in Proposition 2.4) is precisely the solution (M, r_M) (respectively (M, s_M)); hence justifying the terminology in Section 1).*

We refer to the YB-semitruss $(M(X, r), +, \circ, \lambda, \sigma)$ as the *unital structure YB-semitruss* associated to (X, r) and to the sub-YB-semitruss $S(X, r)$ as the *structure YB-semitruss* of (X, r) .

We note that for specific solutions (X, r) the monoid $M(X, r)$ possibly can be made into a YB-semitruss in several manners. The following example shows that one can consider another σ -map. This shows that given a set A with two operations $+, \circ$ and a λ -map, it is possible to have more than one σ -map that makes $(A, +, \circ, \lambda, \sigma)$ into a YB-semitruss.

Example 2.12. Let $X = \{1, 2\}$. Let $r : X \times X \rightarrow X \times X$ denote the left non-degenerate solution with $\lambda_1 = \lambda_2 = \rho_2 = \text{id}_X$ and $\rho_1 : X \rightarrow X : x \mapsto 1$. The associated left derived structure monoid is $A = A(X, r) = \langle X \mid 1+2 = 2+1 = 1+1 \rangle$

and the σ -map is such that σ_1 is the constant map onto 1, and $\sigma_2 = \text{id}$. Note that $M = M(X, r) = A$ and $r_A = r_M$. One can also consider another σ -map, denoted σ' , and defined by σ'_1 the constant mapping onto 2 and $\sigma'_2 = \text{id}$. Notice that the σ' -map yields a left non-degenerate solution $r' : X^2 \rightarrow X^2$ with λ -map (respectively ρ -map), denoted λ' (respectively ρ'), and defined by $\lambda'_1 = \lambda'_2 = \rho'_2 = \text{id}_X$ and ρ'_1 is the constant map onto 2. We get that $M(X, r') = A(X, r') = A = M(X, r)$. However, it easily is verified that the solutions r and r' are not isomorphic and thus also r_M and r'_M are not isomorphic.

Let \mathbb{N} denote the non-negative integers. Note that $M(X, r)$ has a natural strongly \mathbb{N} -gradation (its generators $x \in X$ are the elements of degree 1) in the following sense.

Definition 2.13. One says that a YB-semitruss $(A, +, \circ, \lambda, \sigma)$ is \mathbb{N} -graded (respectively *strongly \mathbb{N} -graded*) if $A = \bigcup_{n \in \mathbb{N}} A_n$, a disjoint union of subsets indexed by the non-negative integers, so that $A_n + A_m \subseteq A_{n+m}$ (respectively $A_n + A_m = A_{n+m}$) and all maps λ_a and σ_a are graded, i.e. degree preserving. In particular, if A is strongly \mathbb{N} -graded then the subsemigroup $A \setminus A_0$ is additively (and thus also multiplicatively) generated by its elements of degree 1. Note that the derived solution $r_A : A^2 \rightarrow A^2$ restricts to a solution $r_{A_1} : A_1^2 \rightarrow A_1^2$.

Corollary 2.14. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss and let $S(A, r_A)$ be the structure semigroup of its associated solution. Then the YB-semitruss A is an epimorphic image of the structure YB-semitruss $S(A, r_A)$. If A is a unital YB-semitruss then A is an epimorphic image of the unital structure YB-semitruss $M(A, r_A)$. Furthermore, if A is strongly \mathbb{N} -graded with $A_0 = \emptyset$ then A is a graded epimorphic image of the \mathbb{N} -graded structure YB-semitruss $S(A_1, r_{A_1})$. If, moreover, A is unital and $A_0 = \{1\}$ then A is a graded epimorphic image of the \mathbb{N} -graded unital structure YB-semitruss $M(A_1, r_{A_1})$.*

In particular, left non-degenerate solutions are restrictions of solutions associated to YB-semitrusses.

Proof. Of course there is a natural map of the generating set A of $S(A, r_A)$ to A . Because of the equations (2.18), (2.19) and (2.20) in $S(A, r_A)$ and the equations (2.13), (2.14) and (2.15) this map can be extended to a map $f : S(A, r_A) \rightarrow A$. Because of Lemma 2.7 and (2.17) it follows that f is an epimorphism of YB-semitrusses. In case A is unital, we have that $\lambda_1 = \text{id}$ in both YB-semitrusses and also $\lambda_a(1) = 1$ for all a . Hence we can extend f to $M = M(A, r_A)$. This proves the first part of the statement. The graded statements for $A_0 = \emptyset$ follow at once as A is generated by $A_1 \cup A_0$ and the natural map preserves degrees. \square

In the previous section it was shown that isomorphic YB-semitrusses yield isomorphic associated solutions. The converse is however not true. Yet, isomorphic left non-degenerate solutions have isomorphic (unital) structure YB-semitrusses and vice versa.

Proposition 2.15. *Let (X, r) and (Y, r') be left non-degenerate solutions. Then $(X, r) \cong (Y, r')$ if and only if the associated (unital) structure YB-semitrusses $S(X, r)$ and $S(Y, r')$ ($M(X, r)$ and $M(Y, r')$) are isomorphic as YB-semitrusses.*

Proof. Let (X, r) and (Y, r') be two isomorphic left non-degenerate solutions and denote $r(x, y) = (\lambda_x(y), \rho_y(x))$, for all $x, y \in X$, and $r'(x', y') = (\lambda'_{x'}(y'), \rho'_{y'}(x'))$, for all $x', y' \in Y$. Then, there exists a bijection $f : X \rightarrow Y$ such that $(f \times f)r = r'(f \times f)$, i.e. for any $x, y \in X$, $f(\lambda_x(y)) = \lambda'_{f(x)}(f(y))$ and $f(\rho_y(x)) = \rho'_{f(y)}(f(x))$. Defining the σ -map and σ' -map as (2.17) for $M(X, r)$ and $M(Y, r')$ respectively,

we get for any $x, y \in X$,

$$\begin{aligned}
f(\sigma_y(x)) &= f(\lambda_y(\rho_{\lambda_x^{-1}(y)}(x))) \\
&= \lambda'_f(y)(f(\rho_{\lambda_x^{-1}(y)}(x))) \\
&= \lambda'_f(y)(\rho'_{f(\lambda_x^{-1}(y))}(f(x))) \\
&= \lambda'_f(y)(\rho'_{\lambda_{f(x)}^{-1}(f(y))}(f(x))) \\
&= \sigma'_{f(y)}(f(x)).
\end{aligned}$$

We will prove that the associated unital structure YB-semitrusses $M = M(X, r)$ and $M' = M(Y, r')$ are isomorphic. The proof for the associated structure YB-semitrusses is similar. Extend f to M by defining $f(x \circ a) = f(x) \circ' f(a)$ for all $x \in X, a \in M$. Hence, $f(a \circ b) = f(a) \circ' f(b)$, for all $a, b \in M$. One can check that $f : M \rightarrow M'$ is a bijection. To prove that it is well-defined, let $x, y \in X$. Then,

$$\begin{aligned}
f(\lambda_x(y) \circ \rho_y(x)) &= f(\lambda_x(y)) \circ' f(\rho_y(x)) \\
&= \lambda'_{f(x)}(f(y)) \circ' \rho'_{f(y)}(f(x)) \\
&= f(x) \circ' f(y) \\
&= f(x \circ y).
\end{aligned}$$

By induction on the length of the elements in M , one can show that $f(\lambda_a(b)) = \lambda'_{f(a)}(f(b))$ and $f(\rho_b(a)) = \rho'_{f(b)}(f(a))$, for any $a, b \in M$. Finally, since $a + b = a \circ \lambda_a^{-1}(b)$ and $f(a) +' f(b) = f(a) \circ' \lambda_{f(a)}^{-1}(f(b))$ for all $a, b \in M$, we obtain

$$\begin{aligned}
f(a + b) &= f(a \circ \lambda_a^{-1}(b)) \\
&= f(a) \circ' f(\lambda_a^{-1}(b)) \\
&= f(a) \circ' \lambda_{f(a)}^{-1}(f(b)) \\
&= f(a) +' f(b).
\end{aligned}$$

Hence, by Definition 2.8, $(M(X, r), +, \circ, \lambda, \sigma) \cong (M(Y, r'), +', \circ', \lambda', \sigma')$ as YB-semitrusses.

Conversely, assume that $M = M(X, r)$ and $M' = M(Y, r')$ are isomorphic as YB-semitrusses, i.e. there exists a bijection $f : M \rightarrow M'$ such that $(f \times f)r_M = r_{M'}(f \times f)$, i.e. for any $a, b \in M$, $f(\lambda_a(b)) = \lambda'_{f(a)}(f(b))$ and $f(\rho_b(a)) = \rho'_{f(b)}(f(a))$. Hence, for any $x, y \in X$, $f(\lambda_x(y)) = \lambda'_{f(x)}(f(y))$ and $f(\rho_y(x)) = \rho'_{f(y)}(f(x))$. Furthermore, since the set X is the unique minimal set of generators of the monoid (M, \circ) and f is a monoid homomorphism, we get that $f(X) = Y$, the unique minimal set of generators of the monoid M' . Hence, $(X, r) \cong (Y, r')$. \square

Two structure monoids that are isomorphic as monoids do not necessarily imply isomorphic solutions. In [39, Example 1.1], two non-isomorphic bijective non-degenerate solutions are given with the same structure monoid. In case both solutions are involutive non-degenerate one has a satisfactory answer. This already has been pointed out in [39, page 6]. If one of the solutions is bijective while the other is still involutive, not necessarily non-degenerate, the result is still true.

Proposition 2.16. *Let (X, r) be an involutive solution. Suppose (Y, r') is a bijective solution such that $M(X, r) \cong M(Y, r')$ as monoids. Then, $(X, r) \cong (Y, r')$ as solutions.*

Proof. The monoid $M(X, r)$ has a unique minimal generating set X , which holds analogously for Y . Hence, a monoid isomorphism f between $M(X, r)$ and $M(Y, r')$ maps X to Y . Hence, this can be restricted to a map $f : X \rightarrow Y$. Moreover, considering that r is involutive, one obtains that at most two words of length 2

(in the alphabet X) become equal in the structure monoid of $M(X, r)$. Hence, the same can be said about words of length 2 (in alphabet Y) in $M(Y, r')$. In particular, r' is involutive. Let $x_1, x_2 \in X$. Then, either $x_1 \circ x_2$ can not be rewritten or $x_1 \circ x_2 = u \circ v$ with $r(x_1, x_2) = (u, v)$. In the former case, as f is a monoid isomorphism, it follows that $f(x_1 \circ x_2) = f(x_1) \circ f(x_2)$ can also not be rewritten, implying that $\lambda_{f(x_1)}(f(x_2)) = f(x_1)$ and $\rho_{f(x_2)}(f(x_1)) = f(x_2)$. In the latter case, $f(x_1) \circ f(x_2) = f(u) \circ f(v)$ are the two only ways to write this element. Considering the defining relation of $M(Y, s)$, it follows that $\lambda_{f(x_1)}(f(x_2)) = f(u)$ and $\rho_{f(x_2)}(f(x_1)) = f(v)$. In conclusion, in both cases one obtains that $f\lambda_x = \lambda_{f(x)}f$ and $f\rho_x = \rho_{f(x)}f$ for all $x \in X$, which shows that f is an isomorphism of solutions. \square

The following example shows that the assumption that (Y, r') is bijective is essential. Let $X = \{1, 2\}$ and (X, r) the trivial solution, i.e. $r(x, y) = (y, x)$, for $x, y \in X$. Its structure monoid is $M(X, r) = \langle X \mid 1 \circ 2 = 2 \circ 1 \rangle$, the free abelian monoid of rank 2. On X one can also have a degenerate idempotent solution (X, r') with $r'(x, y) = (\lambda_x(y), \rho_y(x))$ and $\lambda_1 : x \mapsto 1, \lambda_2 = \rho_1 = \text{id}$ and $\rho_2 : x \mapsto 2$, i.e. $r'(2, 1) = (1, 2)$ and the other pairs are fixed under r' . Then, $M(X, r') = \langle X \mid 2 \circ 1 = 1 \circ 2 \rangle = M(X, r)$. However, $(X, r) \not\cong (X, r')$. Note that the example (X, r') is not left non-degenerate and that, because of Theorem 3.1 no counterexamples exists in case (Y, r') is finite non-degenerate.

We end this section with a class of examples that yield left non-degenerate idempotent solutions.

Example 2.17. If (X, r) is an idempotent solution, i.e. $r^2 = r$, then $\lambda_{\lambda_x(y)}(\rho_y(x)) = \lambda_x(y)$ and $\rho_{\rho_y(x)}(\lambda_x(y)) = \rho_y(x)$. If furthermore the solution is left non-degenerate, the associated σ -map on the unital structure YB-semitruss $M(X, r)$ satisfies $\sigma_b(a) = \lambda_b(\rho_{\lambda_a^{-1}(b)}(a)) = \lambda_{\lambda_a(t)}(\rho_t(a)) = \lambda_a(t) = \lambda_a(\lambda_a^{-1}(b)) = b$, with $t = \lambda_a^{-1}(b)$. Conversely, assume $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss with $\sigma_b(a) = b$ for all $a, b \in A$, then one easily verifies that $(r_A)^2 = r_A$, i.e. the solution (A, r_A) is idempotent. Hence, idempotent left non-degenerate solutions correspond with YB-semitrusses $(A, +, \circ, \lambda, \sigma)$ with σ_b the constant mapping onto b , for every $b \in A$.

2.2. Left cancellative YB-semitrusses.

In this section we show that the known algebraic associative structures that determine left non-degenerate solutions, and that are mentioned in the introduction, are special classes of YB-semitrusses.

Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. Because $a \circ b = a + \lambda_a(b)$ for all $a, b \in A$, it easily is verified that $(A, +)$ being left cancellative (i.e. $a + b = a + d$ implies $b = d$) is equivalent to (A, \circ) being left cancellative. So if $(A, +)$ is a left cancellative semigroup, then the mappings σ_b are uniquely determined as $a + b = b + \sigma_b(a)$ for any $a, b \in A$ (in the non-cancellative case this is not necessarily true, see Example 2.12). Furthermore, since (by Proposition 2.6) $a \circ b = \lambda_a(b) \circ \lambda_{\lambda_a^{-1}(b)}^{-1} \sigma_{\lambda_a(b)}(a)$, there exists a unique $x \in A$ so that $a \circ b = \lambda_a(b) \circ x$. It turns out that several of the requirements to be a YB-semitruss are then redundant.

Proposition 2.18. *Let $(A, +, \circ, \lambda)$ be a left semitruss with $(A, +)$ left cancellative (and thus also (A, \circ) is left cancellative) and λ_a bijective, for any $a \in A$ such that for all $a, b \in A$, the equation*

$$a \circ b = a + \lambda_a(b),$$

holds and there exists an $x \in A$ such that

$$a \circ b = \lambda_a(b) \circ x.$$

We put $\rho_b(a) = x$. Then there exists a unique semigroup anti-morphism $\sigma : (A, +) \rightarrow \text{End}(A, +) : a \mapsto \sigma_a$ such that $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss. For $a, b \in A$ we have $\sigma_b(a) = \lambda_b \rho_{\lambda_a^{-1}(b)}(a)$.

Proof. By [8, Proposition 2.1], we have that $\lambda : (A, \circ) \rightarrow \text{Aut}(A, +) : a \mapsto \lambda_a$ is a semigroup morphism. By the previous and the assumptions, for any $a, b \in A$, there is a unique element $x \in A$ such that $a \circ b = \lambda_a(b) \circ x$. We denote x by $\rho_b(a)$. Now, set $\sigma_b(a) = \lambda_b \rho_{\lambda_a^{-1}(b)}(a)$. Then

$$\begin{aligned} b + \sigma_b(a) &= b + \lambda_b \rho_{\lambda_a^{-1}(b)}(a) = b \circ \rho_{\lambda_a^{-1}(b)}(a) = \lambda_a \lambda_a^{-1}(b) \circ \rho_{\lambda_a^{-1}(b)}(a) \\ &= a \circ \lambda_a^{-1}(b) = a + \lambda_a \lambda_a^{-1}(b) = a + b. \end{aligned}$$

Hence, (2.4) holds. Moreover,

$$\begin{aligned} \lambda_a(b) + \sigma_{\lambda_a(b)} \lambda_a(d) &= \lambda_a(d) + \lambda_a(b) = \lambda_a(d + b) = \lambda_a(b + \sigma_b(d)) \\ &= \lambda_a(b) + \lambda_a \sigma_b(d), \end{aligned}$$

and, since $(A, +)$ is left cancellative, we get $\sigma_{\lambda_a(b)} \lambda_a(d) = \lambda_a \sigma_b(d)$, i.e. condition (2.6) is satisfied. Furthermore, note that

$$d + \sigma_d(a + b) = a + (b + d) = a + d + \sigma_d(b) = d + \sigma_d(a) + \sigma_d(b),$$

i.e. $\sigma_d(a + b) = \sigma_d(a) + \sigma_d(b)$ and

$$b + a + \sigma_a \sigma_b(d) = b + \sigma_b(d) + a = d + (b + a) = b + a + \sigma_{b+a}(d),$$

i.e. $\sigma_a \sigma_b = \sigma_{b+a}$. Hence $\sigma : (A, +) \rightarrow \text{End}(A, +)$ is an anti-morphism, i.e. (2.5) holds. Finally, since (2.3) holds by assumption, $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss. \square

We call a YB-semitruss $(A, +, \circ, \lambda, \sigma)$ with $(A, +)$ left cancellative (and so also (A, \circ) left cancellative) a *left cancellative YB-semitruss*.

Example 2.19. Let (X, r) be a left non-degenerate solution and let η be the left cancellative congruence on $(M(X, r), +)$. In [20, Lemma 5.8, Corollary 5.9], it is shown that the left cancellative image $\bar{M} = M/\eta$ of $(M(X, r), +)$ has a left semitruss structure, denoted by $(\bar{M}, +, \circ, \bar{\lambda})$, with a unique $\bar{\sigma}$ -map such that $a + \bar{\lambda}_a(b) = a \circ b$ and $a + b = b + \bar{\sigma}_b(a)$. We claim that $(\bar{M}, +, \circ, \bar{\lambda}, \bar{\sigma})$ is a left cancellative YB-semitruss. Indeed $(\bar{M}, +)$ and (\bar{M}, \circ) are left cancellative and

$$a \circ b = a + \bar{\lambda}_a(b) = \bar{\lambda}_a(b) + \bar{\sigma}_{\bar{\lambda}_a(b)}(a) = \bar{\lambda}_a(b) \circ \bar{\lambda}_{\bar{\lambda}_a(b)}^{-1} \bar{\sigma}_{\bar{\lambda}_a(b)}(a),$$

i.e. $(\bar{M}, +, \circ, \bar{\lambda})$ is a left semitruss satisfying the assumptions of Proposition 2.18. Therefore $(\bar{M}, +, \circ, \bar{\lambda}, \bar{\sigma})$ is a left cancellative YB-semitruss.

Example 2.20. As defined by Catino, Colazzo and Stefanelli in [10], a *left cancellative semi-brace* is a triple $(A, +, \circ)$ such that (A, \circ) is a group, $(A, +)$ is a left cancellative semigroup and the condition $a \circ (b + d) = a \circ b + a \circ (\bar{a} + d)$ holds for any $a, b, d \in A$, where \bar{a} denotes the inverse of a in (A, \circ) . Any left cancellative semi-brace is a YB-semitruss with (A, \circ) a group, $\lambda_a(b) = a \circ (\bar{a} + b)$ and $\sigma_b(a) = \lambda_b(\bar{b} \circ (a + b)) = \lambda_b(\bar{b}) + a + b$ for all $a, b \in A$. Conversely, a YB-semitruss with (A, \circ) a group is a left cancellative semi-brace.

Proof. Let $(A, +, \circ)$ be a left cancellative semi-brace. Let 1 denote the identity of (A, \circ) . Then $1 + 1 = 1 \circ (1 + 1) = 1 \circ 1 + 1 \circ (\bar{1} + 1) = 1 + 1 + 1$ and thus the left cancellativity of $(A, +)$ yields $1 = 1 + 1$. So, for any $a \in A$ we obtain that $1 + a = 1 + 1 + a$ and thus $a = 1 + a$, i.e. 1 is a left identity of $(A, +)$. Put $\lambda_a(b) = a \circ (\bar{a} + b)$, by [10, Proposition 3] the map $\lambda : (A, \circ) \rightarrow \text{Aut}(A, +) : a \mapsto \lambda_a$ is a group homomorphism and $a \circ (b + d) = a \circ b + \lambda_a(d)$. In addition, $a + \lambda_a(b) = a \circ 1 + \lambda_a(b) = a \circ (1 + b) = a \circ b$. Because (A, \circ) is a group, for any $a, b \in A$ there

exists a unique $x \in A$ such that $a \circ b = \lambda_a(b) \circ x$. Hence, from Proposition 2.18 we obtain that indeed $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss.

Conversely, let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss with (A, \circ) a group. Then, $(A, +)$ is a left cancellative semigroup. Moreover, (2.3) yields $\lambda_a^{-1}(b) = \bar{a} \circ (a + b)$ and, since $\lambda_a^{-1} = \lambda_{\bar{a}}$, we get $\lambda_a(b) = a \circ (\bar{a} + b)$. Hence, $a \circ (b + d) = a \circ b + \lambda_a(d) = a \circ b + a \circ (\bar{a} + d)$. Therefore, $(A, +, \circ)$ is a left cancellative semi-brace. \square

Example 2.21. As defined by Guarnieri and Vendramin in [38], a *skew left brace* is a triple $(A, +, \circ)$, with $(A, +)$ and (A, \circ) groups, so that $a \circ (b + d) = a \circ b - a + a \circ d$, for all $a, b, d \in A$ (here $-a$ denotes the inverse of a in the additive group). If, furthermore, $(A, +)$ is abelian, then this is called a *left brace*, as introduced by Rump in [55] (see also [18]). Skew left braces are (unital) YB-semitrusses $(A, +, \circ, \lambda, \sigma)$ where both $(A, +)$ and (A, \circ) are groups. In this case $\lambda_a(b) = -a + a \circ b$ and $\sigma_b(a) = -b + a + b$. Furthermore, skew left braces are left cancellative semi-braces for which also ρ_a is bijective, for any $a \in A$.

Proof. Note that if $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss with $(A, +)$ and (A, \circ) groups, they share the same identity, say 1, and by Example 2.20, $(A, +, \circ)$ it is a left cancellative semi-brace. Then $\lambda_a(b) = a \circ (\bar{a} + b) = -a + (a \circ 1) + (a \circ (\bar{a} + b)) = -a + a \circ (1 + b) = -a + (a \circ b)$ and $\sigma_b(a) = \lambda_b(\bar{b}) + a + b = -b + (b \circ \bar{b}) + a + b = -b + a + b$. Hence $(A, +, \circ)$ is a skew left brace. The previous equalities also show that a skew left brace is a left cancellative semi-brace with $(A, +)$ also a group.

The last statement of the example has been proven in [10, p. 167]. For completeness' sake we include a short proof. So, let $(A, +, \circ)$ be a left cancellative semi-brace and assume that ρ_a is bijective for any $a \in B$. We know that $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss and thus $\lambda : (A, \circ) \rightarrow \text{Aut}(A, +)$ is a monoid homomorphism and, by Lemma 2.7, $\rho : (A, \circ) \rightarrow \text{Map}(A, A) : b \mapsto \rho_b$ is a monoid anti-homomorphism. In particular, since λ_a and ρ_a are bijective for any $a \in A$ we get $\lambda_1 = \text{id}$ and $\rho_1 = \text{id}$ where 1 denotes the identity of (A, \circ) . Moreover, by Proposition 2.18 we get $\rho_b(a) = \overline{(\lambda_a(b))} \circ a \circ b = \overline{a \circ (\bar{a} + b)} \circ a \circ b = \overline{(\bar{a} + b)} \circ \bar{a} \circ a \circ b = \overline{(\bar{a} + b)} \circ b$, which yields

$$a + b = b \circ \overline{\rho_b(\bar{a})}. \quad (2.21)$$

It follows that 1 also is an identity for $(A, +)$. Indeed, for any $a \in A$, we have $1 + a = 1 + \lambda_1(a) = 1 \circ a = a$ and $a + 1 \stackrel{(2.21)}{=} 1 \circ \rho_1(\bar{a}) = \bar{a} = a$. Furthermore, put $b = \lambda_a(\bar{a})$. Then $a + b = a + \lambda_a(\bar{a}) = a \circ \bar{a} = 1$. Since $\overline{\rho_{\bar{a}}(a)} = \overline{(\bar{a} + \bar{a})} \circ \bar{a} = a \circ (\bar{a} + \bar{a}) = \lambda_a(\bar{a}) = b$ we also get that $b + a \stackrel{(2.21)}{=} a \circ \rho_a(\bar{b}) = a \circ \rho_a \rho_{\bar{a}}(\bar{a}) = a \circ \rho_{\bar{a} \circ a}(\bar{a}) = a \circ \rho_1(\bar{a}) = a \circ \bar{a} = 1$. Therefore, $(A, +)$ is a group and thus $(A, +, \circ)$ is skew left brace and the result follows. \square

Note that if $(B, +, \circ)$ is a skew left brace, the opposite YB-semitruss as defined in Proposition 2.5 is a skew left brace and the definition coincides with the definition of an opposite skew left brace given in [48, Proposition 3.1].

In order to deal with degenerate and non-bijective solutions the notion of a left semi-brace has been introduced in [43]. The definition is as in Example 2.20 but without the restriction that $(A, +)$ has to be left cancellative. In general this is not a YB-semitruss as, for example, $A + b$ is not necessarily contained in $b + A$. It is easy to see that a left semi-brace is a YB-semitruss if and only if it is a left cancellative semi-brace.

Catino, Colazzo and Stefanelli in [13] and Catino, Mazzotta and Stefanelli in [14] introduced a generalization of left semi-braces. In general these do not yield left non-degenerate solutions.

We conclude this section with an example of a left cancellative YB-semitruss that is finite but is not a left cancellative semi-brace.

Example 2.22. Let $A = \{1, 2, 3, 4\}$. Define for any $a, b \in A$, $a + b = b$, $\sigma_b(a) = b$ and $a \circ b = \lambda_a(b)$, where $\lambda_1 = \lambda_2 = \text{id}_A$ and $\lambda_3 = \lambda_4 = (13)(24)$. Then $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss with $(A, +)$ left cancellative and ρ -map given by $\rho_1 = \rho_3 : t \mapsto 1$ and $\rho_2 = \rho_4 : t \mapsto 2$. In particular $1 \circ 1 = 2 \circ 1$ and thus (A, \circ) is not a group.

2.3. Idempotents in YB-semitrusses.

For a semigroup $(S, *)$ we denote by $E_*(S)$ its subset of idempotents. In case $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss we have two such subsets $E_+(A)$ and $E_\circ(A)$. It has been shown in [10] that if A is a left cancellative semi-brace then $E_+(A)$ is a subsemigroup of $(A, +)$ and, furthermore $A = A + 1 + E_+(A)$, where $\{1\} = E_\circ(A)$, and $A + 1$ is a maximal subgroup of $(A, +)$ and it is a sub-YB-semitruss that is skew left brace. We will prove that for an arbitrary YB-semitruss $E_+(A)$ is subsemigroup of $(A, +)$ and also $E_\circ(A)$ is a subsemigroup of (A, \circ) . In the terminology of [1], $(A, +)$ and (A, \circ) are E-semigroups and they thus both have a so-called unitary cover. Actually we will show that $E_+(A)$ is a sub-YB-semitrusses and $E_\circ(A)$ is a sub-semitruss of A .

Proposition 2.23. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. The following properties hold.*

- (1) $\lambda_e = \text{id}$ for $e \in E_\circ(A)$.
- (2) $\sigma_e^2 = \sigma_e$ for $e \in E_+(A)$.
- (3) $E_\circ(A) \subseteq E_+(A)$.
- (4) $E_+(A)$ is a sub-YB-semitruss of A .
- (5) $E_\circ(A)$ is a sub-semitruss of A .
- (6) *If the associated solution (A, r_A) is bijective, then $E_+(A)$ and $E_\circ(A)$ are commutative semigroups.*

Proof. (1) Follows from the fact that λ_e is bijective and $\lambda_e^2 = \lambda_{e \circ e} = \lambda_e$. (2) Follows from the fact that the σ -map is a anti-homomorphism with domain $(A, +)$. To prove (3), let $e \in E_\circ(A)$ then $e = e \circ e = e + \lambda_e(e) = e + e$ and thus $e \in E_+(A)$.

(4) Let $e, f \in E_+(A)$. Then,

$$(e + f) + (e + f) \stackrel{(2.4)}{=} e + f + f + \sigma_f(e) = e + f + \sigma_f(e) \stackrel{(2.4)}{=} e + e + f = e + f,$$

i.e. $e + f \in E_+(A)$. Hence $E_+(A)$ is a subsemigroup. Since $e \circ f = e + \lambda_e(f)$ and because $\lambda_e(f) \in E_+(A)$, as $\lambda_e \in \text{End}(A, +)$, we get that $e \circ f \in E_+(A)$. So, $E_+(A)$ is a subsemigroup of (A, \circ) . Since $E_+(A)$ is λ - and σ -invariant it follows that $E_+(A)$ is a sub-YB-semitruss of A .

(5) Let $e, f \in E_\circ(A)$. Then,

$$\begin{aligned} (e \circ f) \circ (e \circ f) &= (e \circ f) + \lambda_{e \circ f}(e \circ f) = e + f + \lambda_e \lambda_f(e + f) \\ &\stackrel{(1)}{=} e + f + e + f \stackrel{(4)}{=} e + f \stackrel{(1)}{=} e \circ f, \end{aligned}$$

i.e. $e \circ f \in E_\circ(A)$. Hence, also $e + f = e \circ \lambda_e^{-1}(f) \stackrel{(1)}{=} e \circ f \in E_\circ(A)$. Moreover, $\lambda_e(f) \stackrel{(1)}{=} f \in E_\circ(A)$. It follows that $E_\circ(A)$ is a sub-semitruss.

(6) Assume r_A is bijective, i.e. σ_x is bijective for any $x \in A$. Hence, by part (2) it follows that $\sigma_e = \text{id}$ for any $e \in E_+(A)$. Therefore, $e + f \stackrel{(2.4)}{=} f + \sigma_f(e) = f + e$. Hence, $E_+(A)$ is commutative. In case $e, f \in E_\circ(A)$ then we get that $e \circ f = e + f = f + e = f \circ e$. So also $E_\circ(A)$ is commutative. \square

Recall that if S is a right simple semigroup, i.e. S is the only right ideal, or a left cancellative semigroup then any idempotent is a left identity. For both classes

of semigroups we obtain additional information. We begin with the class of left cancellative semigroups.

Proposition 2.24. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. If $e \in E_+(A)$ is a left identity of $(A, +)$, then $\sigma_e(a) = a + e$. In particular, if $(A, +, \circ, \lambda, \sigma)$ is left cancellative, then $E_\circ(A)$ is a sub-YB-semitruss.*

Proof. Let $e \in E_+(A)$ be a left identity of $(A, +)$ and $a \in A$. It follows that

$$\sigma_e(a) = e + \sigma_e(a) \stackrel{(2.4)}{=} a + e.$$

Hence, if $(A, +)$ is left cancellative then, for any $e, f \in E_\circ(A)$, we get that $\sigma_e(f) = f + e = e \in E_\circ(A)$. So $E_\circ(A)$ is λ - and σ -invariant. Hence, by Proposition 2.23.(5), $E_\circ(A)$ is a sub-YB-semitruss. \square

Proposition 2.25. *Let $(A, +, \circ, \lambda, \sigma)$ be a left cancellative YB-semitruss. If the associated solution (A, r_A) is bijective then*

- (1) $|E_+(A)| \leq 1$, and
- (2) $(A, +)$ is right cancellative.

In particular, in this case, if furthermore A is finite, then $(A, +)$ is a group.

Proof. (1) Let $e, f \in E_+(A)$. Since $(A, +)$ is left cancellative, e and f also are left identities. Moreover, since r_A is bijective, we have that σ_e is bijective and thus $\sigma_e = \text{id}$. It follows that $e = f + e = e + \sigma_e(f) = e + f = f$.

(2) Let $a, b, d \in A$. Assume that $a + b = d + b$. Equation (2.4) yields $b + \sigma_b(a) = b + \sigma_b(d)$. Since $(A, +)$ is left cancellative, it follows that $\sigma_b(a) = \sigma_b(d)$. Hence, since r_A is bijective, and thus σ_b bijective, $a = d$. Therefore $(A, +)$ is right cancellative. \square

Proposition 2.26. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. If $(A, +)$ is a group, then (A, \circ) is a group.*

Proof. First note that (2.3) and $(A, +)$ a group yield that (A, \circ) is left cancellative. Let 0 denote the identity of $(A, +)$. We claim that $0 \in E_\circ(A)$ so in particular, 0 is a left identity. First note that $\lambda_0(0) = \lambda_0(0 + 0) = \lambda_0(0) + \lambda_0(0)$, i.e. $\lambda_0(0) = 0$. It yields $0 \circ 0 \stackrel{(2.3)}{=} 0 + \lambda_0(0) = 0$ and thus indeed $0 \in E_\circ(A)$. Similarly, for $a \in A$ we have $\lambda_a(0) = \lambda_a(0 + 0) = \lambda_a(0) + \lambda_a(0)$, i.e. $\lambda_a(0) = 0$. It follows that $a \circ 0 = a + \lambda_a(0) = a + 0 = a$. Hence 0 is also a right identity and (A, \circ) is a monoid. Now set $b = \lambda_a^{-1}(-a)$, then we get $a \circ b = a \circ \lambda_a^{-1}(-a) \stackrel{(2.3)}{=} a + (-a) = 0$. Therefore, any $a \in A$ has a right inverse in (A, \circ) . It follows that (A, \circ) is a group. \square

Corollary 2.27. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. If $(A, +)$ is a group, then $(A, +, \circ)$ is a skew left brace and the associated solution (A, r_A) is bijective non-degenerate.*

Proof. By Proposition 2.26, (A, \circ) is a group. Moreover, by Example 2.21, a YB-semitruss with $(A, +)$ and (A, \circ) groups is a skew left brace. \square

Recall that an inverse semigroup is a regular semigroup S (i.e. for any $a \in S$ there exists $b \in S$ so that $aba = a$) such that any two idempotents of S commute [24, Theorem 1.17].

Lemma 2.28. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. If $a \in A$ is regular in (A, \circ) then a is regular in $(A, +)$.*

Proof. Let $x \in A$ such that $a \circ x \circ a = a$. It follows

$$\begin{aligned}
a &= a \circ x \circ a = a + \lambda_a(x + \lambda_x(a)) = a + \lambda_a(x) + \lambda_a \lambda_x(a) \\
&= a + \lambda_a(x) + \lambda_a \lambda_x \lambda_a \lambda_a^{-1}(a) \\
&= a + \lambda_a(x) + \lambda_{a \circ x \circ a} \lambda_a^{-1}(a) \\
&= a + \lambda_a(x) + \lambda_a \lambda_a^{-1}(a) && \text{(since } a \circ x \circ a = a \text{)} \\
&= a + \lambda_a(x) + a.
\end{aligned}$$

Hence $a + \lambda_a(x) + a = a$ and a is regular in $(A, +)$. \square

Proposition 2.29. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. If $E_+(A) = E_\circ(A)$ and (A, \circ) is an inverse semigroup, then $(A, +)$ is an inverse semigroup.*

Proof. Recall that $e_1 \circ e_2 = e_1 + e_2$ when $e_1 \in E_\circ(A)$. Hence, if (A, \circ) is an inverse semigroup then, for $e_1, e_2 \in E_\circ(A)$, we have $e_2 + e_1 = e_2 \circ e_1 = e_1 \circ e_2 = e_1 + e_2$. Hence $E_+(A)$ is a commutative semigroup and thus, by Lemma 2.28, the result follows. \square

2.4. Matched product of solutions and YB-semitrusses.

In this section we construct from two YB-semitrusses a new YB-semitruss. In particular we give examples of YB-semitrusses that are not as in the examples in Subsection 2.2. We do this by defining a matched product of YB-semitrusses, a construction technique that generalizes the matched product of left cancellative semi-braces [10], of skew left braces [61] and braces [4]. Note that if B_1 and B_2 are left cancellative semi-braces (or skew left braces or braces), then the multiplicative group of the matched product of B_1 and B_2 is the Zappa-Szép product of the groups (B_1, \circ) and (B_2, \circ) [11], i.e. (B_1, \circ) and (B_2, \circ) form a matched pair of groups [63]. In the same manner, the multiplicative structure of the matched product of two unital YB-semitrusses forms a matched pair of monoids. Matched pairs of monoids were successfully used by Gateva-Ivanova and Majid in [33] to construct new solutions. Moreover, we prove that the solution associated to the matched product of two YB-semitrusses is indeed the matched product of the solutions associated to those two YB-semitrusses.

Definition 2.30. Let $(A_1, +, \circ, \lambda^{(1)}, \sigma^{(1)})$, $(A_2, +, \circ, \lambda^{(2)}, \sigma^{(2)})$ be YB-semitrusses, $\alpha : (A_2, \circ) \rightarrow \text{Aut}(A_1, +)$, $\beta : (A_1, \circ) \rightarrow \text{Aut}(A_2, +)$ semigroups morphisms such that the following hold

$$\lambda_a^{(1)} \alpha_{\beta_a^{-1}(u)} = \alpha_u \lambda_{\alpha_u^{-1}(a)}^{(1)}, \quad \lambda_u^{(2)} \beta_{\alpha_u^{-1}(a)} = \beta_a \lambda_{\beta_a^{-1}(u)}^{(2)}, \quad (2.22)$$

$$\alpha_u \sigma_a^{(1)} = \sigma_{\alpha_u(a)}^{(1)} \alpha_u, \quad \beta_a \sigma_u^{(2)} = \sigma_{\beta_a(u)}^{(2)} \beta_a, \quad (2.23)$$

for all $a \in A_1$, $u \in A_2$. The quadruple $(A_1, A_2, \alpha, \beta)$ is called a *matched product system of YB-semitrusses*.

In order to show that the matched product of a matched product system of YB-semitrusses is again a YB-semitruss we need the following lemma.

Lemma 2.31. *Let $(A, +)$ be a semigroup and $\lambda : A \rightarrow \text{End}(A, +) : a \mapsto \lambda_a$. Define $a \circ b = a + \lambda_a(b)$, for any $a, b \in A$. If $\lambda_{a+\lambda_a(b)} = \lambda_a \lambda_b$ for any $a, b \in A$, then $(A, +, \circ, \lambda)$ is a left semitruss.*

Proof. First we show that (A, \circ) is associative and thus a semigroup. For $a, b, d \in A$ we have

$$\begin{aligned} (a \circ b) \circ d &= (a + \lambda_a(b)) \circ d = a + \lambda_a(b) + \lambda_{a+\lambda_a(b)}(d) \\ &= a + \lambda_a(b) + \lambda_a \lambda_b(d) = a + \lambda_a(b + \lambda_b(d)) \\ &= a \circ (b \circ d). \end{aligned}$$

Hence, (A, \circ) is a semigroup. Furthermore,

$$a \circ (b + d) = a + \lambda_a(b + d) = (a + \lambda_a(b)) + \lambda_a(d) = a \circ b + \lambda_a(d).$$

Hence, $(A, +, \circ, \lambda)$ is a left semitruss. \square

Theorem 2.32. *Let $(A_1, A_2, \alpha, \beta)$ a matched product system of YB-semitrusses. Define, on the cartesian product $A_1 \times A_2$, the following*

$$(a, u) + (b, v) = (a + b, u + v), \quad (2.24)$$

$$(a, u) \circ (b, v) = (\alpha_u(\alpha_u^{-1}(a) \circ b), \beta_a(\beta_a^{-1}(u) \circ v)), \quad (2.25)$$

$$\lambda_{(a,u)}(b, v) = (\lambda_a^{(1)} \alpha_{\beta_a^{-1}(u)}(b), \lambda_u^{(2)} \beta_{\alpha_u^{-1}(a)}(v)), \quad (2.26)$$

$$\sigma_{(a,u)}(b, v) = (\sigma_a(b), \sigma_u(v)). \quad (2.27)$$

Then $(A_1 \times A_2, +, \circ, \lambda, \sigma)$ is a YB-semitruss, called the matched product of A_1 and A_2 (via α and β) and denoted by $A_1 \bowtie A_2$.

Proof. Let $a, b, d \in A_1$ and $u, v, w \in A_2$. Denote by $\bar{a} = \alpha_u^{-1}(a)$, $\bar{u} = \beta_a^{-1}(u)$, $\bar{b} = \alpha_v^{-1}(b)$, $\bar{v} = \beta_b^{-1}(v)$. First note that by (2.22), (2.3) and (2.24) we get

$$\begin{aligned} (a, u) + \lambda_{(a,u)}(b, v) &= (a + \lambda_a^{(1)} \alpha_{\bar{u}}(b), u + \lambda_u^{(2)} \beta_{\bar{a}}(v)) = (a + \alpha_u \lambda_{\bar{a}}^{(1)}(b), u + \beta_a \lambda_{\bar{u}}^{(2)}(v)) \\ &= (\alpha_u(\alpha_u^{-1}(a) + \lambda_{\bar{a}}^{(1)}(b)), \beta_a(\beta_a^{-1}(u) + \lambda_{\bar{u}}^{(2)}(v))) \\ &= (\alpha_u(\bar{a} \circ b), \beta_a(\bar{u} \circ v)) = (a, u) \circ (b, v). \end{aligned}$$

Hence, $A_1 \times A_2$ satisfies condition (2.3). Moreover, it is a routine computation to verify that $\lambda_{(a,u)}(b, v) + \lambda_{(a,u)}(d, w) = \lambda_{(a,u)}((b, v) + (d, w))$, i.e. $\lambda : A_1 \times A_2 \rightarrow \text{End}(A_1 \times A_2, +)$. Furthermore, note that $\alpha_u(\bar{a} \circ b) = a \circ \alpha_{\bar{u}}(b)$ and $\beta_{\alpha_{\bar{u}}(b)}^{-1}(\bar{u} \circ v) = \beta_{\alpha_{\bar{u}}(b)}^{-1}(\bar{u}) \circ \beta_b^{-1}(v)$, that yields

$$\begin{aligned} \lambda_{\alpha_u(\bar{a} \circ b)}^{(1)} \alpha_{\beta_{\alpha_u(\bar{a} \circ b)}^{-1}(\bar{u} \circ v)} \beta_a(\bar{u} \circ v)(d) &= \lambda_{a \circ \alpha_{\bar{u}}(b)}^{(1)} \alpha_{\beta_{a \circ \alpha_{\bar{u}}(b)}^{-1}(\bar{u} \circ v)} \beta_a(\bar{u} \circ v)(d) \\ &= \lambda_a^{(1)} \lambda_{\alpha_{\bar{u}}(b)}^{(1)} \alpha_{\beta_{\alpha_{\bar{u}}(b)}^{-1}(\bar{u} \circ v)}(d) \\ &= \lambda_a^{(1)} \lambda_{\alpha_{\bar{u}}(b)}^{(1)} \alpha_{\beta_{\alpha_{\bar{u}}(b)}^{-1}(\bar{u}) \circ \beta_b^{-1}(v)}(d) \\ &= \lambda_a^{(1)} \lambda_{\alpha_{\bar{u}}(b)}^{(1)} \alpha_{\beta_{\alpha_{\bar{u}}(b)}^{-1}(\bar{u})} \alpha_{\beta_b^{-1}(v)}(d) \\ &= \lambda_a^{(1)} \alpha_{\bar{u}} \lambda_{\alpha_{\bar{u}}^{-1} \alpha_{\bar{u}}(b)}^{(1)} \alpha_{\bar{v}}(d) \\ &= \lambda_a^{(1)} \alpha_{\bar{u}} \lambda_b^{(1)} \alpha_{\bar{v}}(d), \end{aligned}$$

In the same way, we get $\lambda_{\beta_a(\bar{u} \circ v)}^{(2)} \beta_{\alpha_{\beta_a(\bar{u} \circ v)}^{-1}(a \circ \alpha_{\bar{u}}(b))} \alpha_u(a \circ \alpha_{\bar{u}}(b))(w) = \lambda_u^{(2)} \beta_{\bar{a}} \lambda_{\bar{v}}^{(2)} \beta_{\bar{b}}(w)$. It follows $\lambda_{(a,u) \circ (b,v)}(d, w) = \lambda_{(\alpha_u(\bar{a} \circ b), \beta_a(\bar{u} \circ v))}(d, w) = \lambda_{(a,u)} \lambda_{(b,v)}(d, w)$. Hence, Lemma 2.31 yields that $(A_1 \times A_2, +, \circ, \lambda)$ is a left semitruss. Moreover, by (2.27) and (2.4), we obtain that $A_1 \times A_2$ satisfies (2.4). In addition, condition (2.5) and (2.6) for $A_1 \times A_2$ follow from the respective conditions in A_1 and A_2 and from (2.27) and (2.23). Since $\lambda_{(a,u)}$ is bijective with inverse given by $\lambda_{(a,u)}^{-1}(b, v) = (\alpha_{\beta_a^{-1}(u)}^{-1} \lambda_a^{(1)-1}(b), \beta_{\alpha_u^{-1}(a)}^{-1} \lambda_u^{(2)-1}(v))$, we conclude that $(A_1 \times A_2, +, \circ, \lambda, \sigma)$ is a YB-semitruss. \square

Via the matched product we now give an example of a YB-semitruss that is not a left cancellative semi-brace, and where the additive and multiplicative operations are different.

Example 2.33. Let $A = \{1, \dots, n\}$ and $B = C_n = \langle \xi \rangle$ the cyclic group of n -elements. Let $a + b = a \circ b = b$, $\lambda_a = \text{id}_A$ and $\sigma_a(b) = a$, for any $a, b \in A$, then $(A, +, \circ, \lambda, \sigma)$ is a YB-semitruss, and let B the trivial skew left brace with $(B, +) = (B, \circ) = C_n$, i.e. a YB-semitruss with $\lambda_x = \sigma_x = \text{id}_B$. Consider the natural action of C_n over A : $\alpha : B \rightarrow A$ defined by $\alpha(\xi^i)(j) = (1 \ 2 \ \dots \ n)^i(j)$. Note that since $\text{Aut}(A, +) = \text{Sym}(A)$ we have that $\alpha : (B, \circ) \rightarrow \text{Aut}(A, +)$. It is easy to check that setting $\beta_a = \text{id}$ for any $a \in A$ we get that (A, B, α, β) is a matched product system of YB-semitrusses.

As in the semi-braces' case we establish a connection between the matched product of YB-semitrusses and the matched product of their solutions.

For completeness' sake we recall the definition of matched product of solutions given in [11]. Let (S, r_S) and (T, r_T) be solutions, $\alpha : T \rightarrow \text{Sym}(S)$ and $\beta : S \rightarrow \text{Sym}(T)$ maps. Put $\alpha_u = \alpha(u)$, for every $u \in T$, and $\beta_a = \beta(a)$, for every $a \in S$, then the quadruple $(r_S, r_T, \alpha, \beta)$ is said to be a *matched product system of solutions* if the following conditions hold

$$\begin{aligned} \alpha_u \alpha_v &= \alpha_{\lambda_u(v)} \alpha_{\rho_v(u)}, & \text{(s1)} & & \beta_a \beta_b &= \beta_{\lambda_a(b)} \beta_{\rho_b(a)}, & \text{(s2)} \\ \rho_{\alpha_u^{-1}(b)} \alpha_{\beta_a(u)}^{-1}(a) &= \alpha_{\beta_{\rho_b(a)} \beta_b^{-1}(u)}^{-1} \rho_b(a), & \text{(s3)} & & \rho_{\beta_a^{-1}(v)} \beta_{\alpha_u(a)}^{-1}(u) &= \beta_{\alpha_{\rho_v(u)} \alpha_v^{-1}(a)}^{-1} \rho_v(u), & \text{(s4)} \\ \lambda_a \alpha_{\beta_a^{-1}(u)} &= \alpha_u \lambda_{\alpha_u^{-1}(a)}, & \text{(s5)} & & \lambda_u \beta_{\alpha_u^{-1}(a)} &= \beta_a \lambda_{\beta_a^{-1}(u)}, & \text{(s6)} \end{aligned}$$

for all $a, b \in S$ and $u, v \in T$.

As shown in [11, Theorem 1], any matched product system of solutions determines a new solution on the set $S \times T$. Specifically, if $(r_S, r_T, \alpha, \beta)$ is a matched product system of solutions, then the map $r : S \times T \times S \times T \rightarrow S \times T \times S \times T$ defined by

$$r((a, u), (b, v)) = \left((\alpha_u \lambda_{\bar{a}}(b), \beta_a \lambda_{\bar{u}}(v)), \left(\alpha_{\bar{U}}^{-1} \rho_{\alpha_{\bar{a}}(b)}(a), \beta_{\bar{A}}^{-1} \rho_{\beta_{\bar{a}}(v)}(u) \right) \right),$$

where

$$\bar{a} = \alpha_u^{-1}(a), \quad \bar{u} = \beta_a^{-1}(u), \quad A = \alpha_u \lambda_{\bar{a}}(b), \quad U = \beta_a \lambda_{\bar{u}}(v), \quad \bar{A} = \alpha_{\bar{U}}^{-1}(A), \quad \bar{U} = \beta_{\bar{A}}^{-1}(U),$$

for all $(a, u), (b, v) \in S \times T$, is a solution. This solution is called the *matched product of the solutions* r_S and r_T (via α and β) and it is denoted by $r_S \bowtie r_T$.

It is a routine computation to prove the following result (see for example [11, Theorem 9] [12, Theorem 6] in the context of (left cancellative) semi-braces).

Proposition 2.34. *Let $(A_1, A_2, \alpha, \beta)$ a matched product system of YB-semitrusses. Then $r_{A_1 \bowtie A_2} = r_{A_1} \bowtie r_{A_2}$.*

3. YB-SEMITRUSSES WITH ASSOCIATED BIJECTIVE SOLUTIONS

The aim of this section is to prove the following result.

Theorem 3.1. *If (X, r) is a finite left non-degenerate solution, then r is bijective if and only if (X, r) is right non-degenerate.*

Castelli, Catino and Stefanelli in [9] have proven the necessity of the condition. Their proof is based on the notion of q -cycle sets introduced by Rump in [58]. We will prove the sufficiency as well and we reprove the necessity in terms of YB-semitrusses.

Recall that the solution (A, r_A) associated to a YB-semitruss A is always left non-degenerate. If the solution also is right non-degenerate, we call the YB-semitruss a *non-degenerate YB-semitruss*. An example is the structure YB-semitruss $S(X, r)$ associated to a non-degenerate solution (X, r) .

Let

$$\mathfrak{q} : A \rightarrow A : a \mapsto \lambda_a^{-1}(a),$$

it is called the *diagonal map* on a YB-semitruss A .

In [58, Corollary 2 of Proposition 8], Rump proved that for any bijective solution (X, r) , \mathfrak{q} is bijective if and only if (X, r) is non-degenerate. In the following lemma we show that one does not need the bijectiveness of r_A to prove that \mathfrak{q} is injective. So for any \mathbb{N} -graded non-degenerate YB-semitruss with all homogeneous components A_n finite, we get that \mathfrak{q} is bijective.

Lemma 3.2. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. Then the diagonal map \mathfrak{q} is injective.*

Proof. Let $a, d \in A$ and assume that $\mathfrak{q}(a) = \mathfrak{q}(d)$. Put say $t = \mathfrak{q}(a)$. It follows that

$$\begin{aligned} \rho_t(a) &= \lambda_{\lambda_a^{-1}(t)}^{-1} \sigma_{\lambda_a(t)}(a) = \lambda_{\lambda_a \lambda_a^{-1}(a)}^{-1} \sigma_{\lambda_a(t)}(a) \\ &= \lambda_a^{-1} \sigma_{\lambda_a(t)}(a) = \sigma_{\lambda_a^{-1} \lambda_a(t)} \lambda_a^{-1}(a) = \sigma_t(t). \end{aligned}$$

Similarly, $\rho_t(d) = \sigma_t(t)$. Hence, $\rho_t(a) = \rho_t(d)$. Since, by assumption, ρ_t is bijective, it follows that $a = d$. So, \mathfrak{q} is injective. \square

Let X be a set with $|X| \geq 2$. Consider the idempotent solution on X given by $r(x, y) = (y, y)$ (see Example 2.17). Clearly r is neither bijective nor right non-degenerate. However, the solution is left non-degenerate (as each $\lambda_x = \text{id}_X$) and \mathfrak{q} is bijective (since $\mathfrak{q}(x) = \lambda_x^{-1}(x) = x$, for any $x \in X$). This shows that for an arbitrary solution (X, r) one can have \mathfrak{q} bijective even if r is neither bijective nor right non-degenerate; and thus the bijectiveness assumption is essential in Rump's result.

Lemma 3.3. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. If the diagonal map \mathfrak{q} is bijective, then r_A is bijective.*

Proof. By Proposition 2.4, r_A is bijective if and only if σ_a is bijective for any $a \in A$. From (2.6) and (2.9) we get that, for $a, b \in A$,

$$\rho_b(a) = \lambda_{\lambda_a^{-1}(b)}^{-1} \sigma_{\lambda_a(b)}(a) = \lambda_{\lambda_a^{-1}(b)}^{-1} \lambda_a \sigma_b \lambda_a^{-1}(a) = \lambda_{\rho_b(a)} \lambda_b^{-1} \sigma_b \mathfrak{q}(a).$$

Since, by assumption, ρ_b is bijective it yields $\lambda_a \lambda_b^{-1} \sigma_b \mathfrak{q} \rho_b^{-1}(a) = a$ and thus

$$\mathfrak{q}(a) = \lambda_b^{-1} \sigma_b \mathfrak{q} \rho_b^{-1}(a).$$

Now, assume \mathfrak{q} is bijective. Then

$$\sigma_b(a) = \lambda_b \mathfrak{q} \rho_b \mathfrak{q}^{-1}(a),$$

i.e. σ_b is bijective. So r_A is bijective. \square

To prove the main result of this section we need one more lemma.

Lemma 3.4. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss with σ_a bijective for any $a \in A$ and let \mathfrak{q} be the diagonal map. For any $a, b \in A$, it holds*

$$\lambda_b^{-1} \mathfrak{q}(a) = \mathfrak{q}(\lambda_{\sigma_a^{-1} \lambda_a(b)}^{-1}(a)).$$

Proof. By (2.9), we get

$$\lambda_{\rho_{\lambda_a^{-1}(b)}^{-1}(a)}^{-1} \lambda_b^{-1} = \lambda_{\rho_{\lambda_a^{-1}(b)}^{-1}(a)}^{-1} \lambda_{\lambda_a(\lambda_a^{-1}(b))}^{-1} = \lambda_{\lambda_a^{-1}(b)}^{-1} \lambda_a^{-1}.$$

Next, put $a = \sigma_b^{-1} \lambda_b(x)$, to obtain

$$\lambda_{\rho_{\lambda_a^{-1} \lambda_b(x)}^{-1}(\sigma_b^{-1} \lambda_b(x))}^{-1} \lambda_b^{-1}(b) = \lambda_{\lambda_{\sigma_b^{-1} \lambda_b(x)}^{-1}(b)}^{-1} \lambda_{\sigma_b^{-1} \lambda_b(x)}^{-1}(b). \quad (3.1)$$

By definition, $\rho_b(a) = \lambda_{\lambda_a(b)}^{-1} \sigma_{\lambda_a(b)}(a)$, and thus

$$\rho_{\lambda_{\sigma_b^{-1} \lambda_b(x)}^{-1}(b)}(\sigma_b^{-1} \lambda_b(x)) = \lambda_b^{-1} \sigma_b(\sigma_b^{-1} \lambda_b(x)) = x.$$

Hence, (3.1) is equivalent to

$$\lambda_x^{-1} \mathbf{q}(b) = \mathbf{q}(\lambda_{\sigma_b^{-1} \lambda_b(x)}^{-1}(b)),$$

as desired. \square

The sufficiency of the next proposition was proven by Castelli, Catino and Stefanelli in [9]. We translated their proof in the language of YB-semitrusses and added it for completeness' sake.

Proposition 3.5. *Let $(A, +, \circ, \lambda, \sigma)$ be an \mathbb{N} -graded YB-semitruss with all homogeneous components A_n finite and with (A, r_A) its associated left non-degenerate solution. Then, (A, r_A) is right non-degenerate if and only if r_A is bijective.*

Proof. Assume first that the solution (A, r_A) is non-degenerate, i.e. $(A, +, \circ, \lambda, \sigma)$ is a non-degenerate YB-semitruss. By Lemma 3.2 the diagonal map \mathbf{q} is injective. As \mathbf{q} is degree preserving and A is an \mathbb{N} -graded YB-semitruss with all homogeneous components A_n finite, it follows that \mathbf{q} is bijective. By Lemma 3.3, we conclude that r_A is bijective.

To prove the converse, assume that (A, r_A) is bijective (i.e. σ_a is bijective for all $a \in A$). First, we prove that the diagonal map \mathbf{q} is bijective. Let $n \in \mathbb{N}$. Without loss of generality, we focus on the restrictions of λ, ρ, σ to A_n for a fixed $n \in \mathbb{N}$. Since A_n is finite, there exists $m \in \mathbb{N}$ such that $(\lambda_z^{-1})^m = \text{id}_{A_n}$ for any $z \in A_n$. Now, if $m = 1$, then $\mathbf{q} = \text{id}_{A_n}$ and in particular \mathbf{q} is bijective. If $m \geq 2$, then by Lemma 3.4 for $a \in A_n$,

$$\begin{aligned} a &= (\lambda_a^{-1})^m(a) = (\lambda_a^{-1})^{m-2} \lambda_a^{-1} \mathbf{q}(a) \\ &= (\lambda_a^{-1})^{m-2} \mathbf{q} \lambda_{\sigma_a^{-1} \lambda_a(a)}^{-1}(a) \\ &= (\lambda_a^{-1})^{m-3} \lambda_a^{-1} \mathbf{q} \lambda_{\sigma_a^{-1} \lambda_a(a)}^{-1}(a) \\ &= (\lambda_a^{-1})^{m-3} \mathbf{q} \lambda_{\sigma_{\lambda_{\sigma_a^{-1} \lambda_a(a)}^{-1}(a)}^{-1} \lambda_{\sigma_a^{-1} \lambda_a(a)}(a)}^{-1} \lambda_{\sigma_a^{-1} \lambda_a(a)}^{-1}(a), \end{aligned}$$

and so on. So there exists $t \in A_n$ such that $a = \mathbf{q}(t)$. Therefore \mathbf{q} is surjective and since A_n is finite, \mathbf{q} is bijective. Now we will prove that (A, r_A) is right non-degenerate. As the ρ -map is degree preserving, it is enough to prove that ρ_a is injective, for all $a \in A$. Assume $\rho_a(b) = \rho_a(d)$, i.e. $\lambda_{\lambda_b(a)}^{-1} \sigma_{\lambda_b(a)}(b) = \lambda_{\lambda_d(a)}^{-1} \sigma_{\lambda_d(a)}(d)$. Therefore (2.6) yields

$$\lambda_{\lambda_b(a)}^{-1} \lambda_b \sigma_a \mathbf{q}(b) = \lambda_{\lambda_b(a)}^{-1} \lambda_b \sigma_a \lambda_b^{-1}(b) = \lambda_{\lambda_d(a)}^{-1} \lambda_d \sigma_a \lambda_d^{-1}(d) = \lambda_{\lambda_d(a)}^{-1} \lambda_d \sigma_a \mathbf{q}(d).$$

Hence, since $\lambda_{\lambda_b(a)}^{-1} \lambda_b = \lambda_{\rho_a(b)} \lambda_a^{-1} = \lambda_{\rho_a(d)} \lambda_a^{-1} = \lambda_{\lambda_d(a)}^{-1} \lambda_d$ we get $\sigma_a \mathbf{q}(b) = \sigma_a \mathbf{q}(d)$. As σ_a is bijective, this implies $\mathbf{q}(b) = \mathbf{q}(d)$, i.e. $b = d$ and thus ρ_a is injective. \square

We are now in a position to prove the main result of this section.

Proof of Theorem 3.1. Let (X, r) be a finite left non-degenerate solution. From Proposition 2.11, we know that we can associate to (X, r) its unital structure YB-semitruss $M = (M(X, r), +, \circ, \lambda, \sigma)$ and a left non-degenerate solution (M, r_M) . Furthermore, M is an \mathbb{N} -graded YB-semitruss with all homogeneous components finite. From Proposition 3.5 it follows that (M, r_M) is right non-degenerate if and only if r_M is bijective. Since (M, r_M) being right non-degenerate is equivalent to (X, r) being right non-degenerate, and r_M being bijective is equivalent to r being bijective, we get that (X, r) is right non-degenerate if and only if r is bijective. \square

In Lemma 3.3 it was shown that the bijectiveness of \mathfrak{q} is an important property to prove that r is bijective. In [20, Lemma 4.4] the bijectiveness of \mathfrak{q} is proven in case the solution is irretractable. The following proposition slightly extends this result and the proof follows the same lines.

Proposition 3.6. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. Assume the following property holds for all $a, b \in A$: $(\lambda_a, \rho_a) = (\lambda_b, \rho_b)$ implies $a = b$. Then the diagonal map \mathfrak{q} is bijective and its inverse is given by $a \mapsto \rho_a^{-1}(a)$.*

Proof. By (2.9),

$$\lambda_{\lambda_{\rho_a^{-1}(a)}(a)}\lambda_a = \lambda_{\lambda_{\rho_a^{-1}(a)}(a)}\lambda_{\rho_a(\rho_a^{-1}(a))} = \lambda_{\rho_a^{-1}(a)}\lambda_a.$$

Because λ_a is bijective, we thus get

$$\lambda_{\lambda_{\rho_a^{-1}(a)}(a)} = \lambda_{\rho_a^{-1}(a)}. \quad (3.2)$$

Furthermore, by (2.11),

$$\rho_a \rho_{\lambda_{\rho_a^{-1}(a)}(a)} = \rho_{\rho_a(\rho_a^{-1}(a))} \rho_{\lambda_{\rho_a^{-1}(a)}(a)} = \rho_a \rho_{\rho_a^{-1}(a)}.$$

Thus, since, by assumption, ρ_a is bijective,

$$\rho_{\lambda_{\rho_a^{-1}(a)}(a)} = \rho_{\rho_a^{-1}(a)}. \quad (3.3)$$

By (3.2), (3.3) and the assumption we get $\lambda_{\rho_a^{-1}(a)}(a) = \rho_a^{-1}(a)$, and therefore $a = \lambda_{\rho_a^{-1}(a)}^{-1}(\rho_a^{-1}(a)) = \mathfrak{q}(\rho_a^{-1}(a))$. It follows that the diagonal map \mathfrak{q} is bijective and its inverse is given by $a \mapsto \rho_a^{-1}(a)$. \square

4. NON-DEGENERATE YB-SEMITRUSSES AND THE RETRACT RELATION

In the previous section we showed that non-degenerate solutions on a finite set are bijective. In this section we prove this for a much wider class of non-degenerate solutions. To do so we introduce, in analogy with skew left braces (see for example [29, 51]), i.e. for non-degenerate bijective solutions (see also [21] for various equivalent definitions), the retract of a non-degenerate YB-semitruss.

Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. Consider the following set

$$\mathcal{G}(A) = \{f(a) := (\sigma_a, \lambda_a, \rho_a) \mid a \in A\}.$$

Lemma 4.1. *Let $a, a', b, b' \in A$.*

- (1) *If $f(a) = f(a')$ then $\sigma_{b \circ a} = \sigma_{b \circ a'}$, $\lambda_{a \circ b} = \lambda_{a' \circ b}$, $\rho_{b \circ a} = \rho_{b \circ a'}$, and $\lambda_{a+b} = \lambda_{a'+b}$, $\rho_{a+b} = \rho_{a'+b}$, $\sigma_{a+b} = \sigma_{a'+b}$.*
- (2) *if $f(b) = f(b')$ then $\lambda_{a \circ b} = \lambda_{a \circ b'}$, $\rho_{b \circ a} = \rho_{b' \circ a}$, $\sigma_{b \circ a} = \sigma_{b' \circ a}$, and $\lambda_{a+b} = \lambda_{a+b'}$, $\rho_{a+b} = \rho_{a+b'}$, $\sigma_{a+b} = \sigma_{a+b'}$.*

Proof. (1) We have that $\lambda_{a \circ b} = \lambda_a \lambda_b = \lambda_{a'} \lambda_b = \lambda_{a' \circ b}$ and similarly $\rho_{b \circ a} = \rho_{b \circ a'}$. Further, $\sigma_{\lambda_b(a)} = \lambda_b \sigma_a \lambda_b^{-1} = \lambda_b \sigma_{a'} \lambda_b^{-1} = \sigma_{\lambda_b(a')}$ yields

$$\sigma_{b \circ a} = \sigma_{b + \lambda_b(a)} = \sigma_{\lambda_b(a)} \sigma_b = \sigma_{\lambda_b(a')} \sigma_b = \sigma_{b \circ a'}.$$

Now, $\lambda_{a+b} = \lambda_{a \circ \lambda_a^{-1}(b)} = \lambda_a \lambda_{\lambda_a^{-1}(b)} = \lambda_{a'} \lambda_{\lambda_a^{-1}(b)} = \lambda_{a'+b}$, $\rho_{a+b} = \rho_{a'+b}$ and $\sigma_{a+b} = \sigma_b \sigma_a = \sigma_b \sigma_{a'} = \sigma_{a'+b}$.

- (2) We have $\lambda_{a \circ b} = \lambda_{a \circ b'}$, $\rho_{b \circ a} = \rho_{b' \circ a}$. Furthermore,

$$\sigma_{b \circ a} = \sigma_{b + \lambda_b(a)} = \sigma_{\lambda_b(a)} \sigma_b = \sigma_{\lambda_b(a')} \sigma_{b'} = \sigma_{b' \circ a}.$$

Moreover, $a + b = b + \sigma_b(a) = b \circ \lambda_b \sigma_b(a)$, then $\lambda_{a+b} = \lambda_{b \circ \lambda_b \sigma_b(a)} = \lambda_b \lambda_{\lambda_b \sigma_b(a)} = \lambda_{b'} \lambda_{\lambda_{b'} \sigma_{b'}(a)} = \lambda_{a+b'}$, similarly $\rho_{a+b} = \rho_{a+b'}$. Finally, $\sigma_{a+b} = \sigma_b \sigma_a = \sigma_{b'} \sigma_a = \sigma_{a+b'}$. \square

As a consequence of Lemma 4.1 we have the following (well-defined) additive and multiplicative operations on $\mathcal{G}(A)$:

$$f(a) + f(b) := (\sigma_{a+b}, \lambda_{a+b}, \rho_{a+b}) = f(a + b),$$

and

$$f(a) \circ f(b) := (\sigma_{a \circ b}, \lambda_{a \circ b}, \rho_{a \circ b}) = f(a \circ b).$$

Hence,

$$f : A \rightarrow \mathcal{G}(A) : a \mapsto f(a) = (\sigma_a, \lambda_a, \rho_a),$$

is a semigroup homomorphism for both the additive and multiplicative operations.

We will show that $\mathcal{G}(A)$ is a YB-semitruss provided A is non-degenerate. First we introduce a λ -map on $\mathcal{G}(A)$.

Lemma 4.2. *The mapping $\lambda : \mathcal{G}(A) \rightarrow \text{Map}(\mathcal{G}(A), \mathcal{G}(A)) : a \mapsto \lambda_{f(a)}$ given by $\lambda_{f(a)}(f(b)) = f(\lambda_a(b))$, is well-defined. Furthermore, each $\lambda_{f(a)}$ is surjective.*

Proof. To prove that the map λ is well-defined, we first note that if $f(a) = f(a')$ then $\lambda_{f(a)} = \lambda_{f(a')}$. Next assume $f(b) = f(b')$. Then $\lambda_{f(a)}(f(b)) = f(\lambda_a(b))$ and $\lambda_{f(a)}(f(b')) = f(\lambda_a(b'))$.

By (2.9), $\lambda_{\lambda_a(b)} = \lambda_a \lambda_b \lambda_{\rho_b(a)}^{-1} = \lambda_a \lambda_{b'} \lambda_{\rho_{b'}(a)}^{-1} = \lambda_{\lambda_a(b')}$. By (2.6) $\sigma_{\lambda_a(b)} = \lambda_a \sigma_b \lambda_a^{-1} = \lambda_a \sigma_{b'} \lambda_a^{-1} = \sigma_{\lambda_a(b')}$ and by (2.6) and (2.9), for all $x \in A$,

$$\begin{aligned} \rho_{\lambda_a(b)}(x) &= \lambda_{\lambda_x \lambda_a(b)}^{-1} \sigma_{\lambda_x \lambda_a(b)}(x) \\ &= \lambda_{\lambda_{x \circ a}(b)}^{-1} \lambda_x \lambda_a \sigma_b \lambda_a^{-1} \lambda_x^{-1}(x) \\ &= \lambda_{\rho_b(x \circ a)} \lambda_b^{-1} \lambda_{x \circ a}^{-1} \lambda_x \lambda_a \sigma_b \lambda_a^{-1} \lambda_x^{-1}(x) \\ &= \lambda_{\rho_{b'}(x \circ a)} \lambda_{b'}^{-1} \lambda_{x \circ a}^{-1} \lambda_x \lambda_a \sigma_{b'} \lambda_a^{-1} \lambda_x^{-1}(x) \\ &= \rho_{\lambda_a(b')}(x). \end{aligned}$$

This shows that $\lambda_{f(a)}(f(b)) = \lambda_{f(a)}(f(b'))$ and thus the map λ is well-defined.

To prove that the map $\lambda_{f(a)}$ is surjective, let $f(d) \in \text{Im}(f)$. Then there exists $b \in A$ such that $\lambda_a(b) = d$. Hence, $\lambda_{f(a)}(f(b)) = (\sigma_{\lambda_a(b)}, \lambda_{\lambda_a(b)}, \rho_{\lambda_a(b)}) = (\sigma_d, \lambda_d, \rho_d) = f(d)$, as desired. \square

Next we introduce a σ -map in $\mathcal{G}(A)$ as follows

$$\sigma : \mathcal{G}(A) \rightarrow \text{Map}(\mathcal{G}(A), \mathcal{G}(A)) : f(b) \mapsto \sigma_{f(b)},$$

with

$$\sigma_{f(b)}(f(a)) = f(\sigma_b(a)).$$

The well-definedness of this map will follow from Lemma 4.4, but first we need another proposition.

Proposition 4.3. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. Let $a, b \in A$. If $\lambda_a = \lambda_b$ and $\rho_a = \rho_b$ then $\sigma_a = \sigma_b$. Furthermore, if $\lambda_a = \rho_a = \text{id}$ then $\sigma_a = \text{id}$.*

Proof. First note that, for $x, y, z \in A$, because of Proposition 2.4 and (2.10),

$$\begin{aligned} \lambda_{\rho_{\lambda_y \lambda_x^{-1}(z)} \rho_y^{-1}(x)} \rho_{\lambda_x^{-1}(z)}(y) &= \rho_{\lambda_{\rho_y \rho_y^{-1}(x)} \lambda_x^{-1}(z)} \lambda_{\rho_y^{-1}(x)}(y) \\ &= \rho_z \lambda_{\rho_y^{-1}(x)}(y), \end{aligned}$$

yields

$$\rho_{\lambda_x^{-1}(z)}(y) = \lambda_{\rho_{\lambda_y \lambda_x^{-1}(z)} \rho_y^{-1}(x)}^{-1} \rho_z \lambda_{\rho_y^{-1}(x)}(y). \quad (4.1)$$

Let $a, b \in A$ and assume $\lambda_a = \lambda_b$ and $\rho_a = \rho_b$. For any $x \in A$, using (4.1) it follows that

$$\begin{aligned}\rho_{\lambda_x^{-1}(a)}(x) &= \lambda_{\rho_{\lambda_x \lambda_x^{-1}(a)} \rho_x^{-1}(x)}^{-1} \rho_a \lambda_{\rho_x^{-1}(x)}(x) \\ &= \lambda_{\rho_a \rho_x^{-1}(x)}^{-1} \rho_a \lambda_{\rho_x^{-1}(x)}(x) \\ &= \lambda_{\rho_b \rho_x^{-1}(x)}^{-1} \rho_b \lambda_{\rho_x^{-1}(x)}(x) \\ &= \rho_{\lambda_x^{-1}(b)}(x).\end{aligned}$$

So,

$$\sigma_a(x) = \lambda_a \rho_{\lambda_x^{-1}(a)}(x) = \lambda_b \rho_{\lambda_x^{-1}(b)}(x) = \sigma_b(x). \quad (4.2)$$

This proves the first part of the statement.

To prove the second part, assume $\lambda_a = \rho_a = \text{id}$. By (4.1) we get that $\rho_{\lambda_x^{-1}(a)}(x) = x$. Then, as in (4.2), $\sigma_a(x) = \lambda_a \rho_{\lambda_x^{-1}(a)}(x) = x$, as desired. \square

Lemma 4.4. *Let $(A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. Let $a, a', x \in A$.*

- (1) *If $\lambda_a = \lambda_{a'}$ then $\lambda_{\rho_x(a)} = \lambda_{\rho_x(a')}$.*
- (2) *if $\lambda_a = \lambda_{a'}$ and $\rho_a = \rho_{a'}$ then $\lambda_{\lambda_x(a)} = \lambda_{\lambda_x(a')}$.*

If, furthermore, A is non-degenerate then $\lambda_a = \lambda_{a'}$ and $\rho_a = \rho_{a'}$ implies

- (3) *$\lambda_{\lambda_x^{-1}(a)} = \lambda_{\lambda_x^{-1}(a')}$ and $\rho_{\lambda_x^{-1}(a)} = \rho_{\lambda_x^{-1}(a')}$.*
- (4) *$\rho_{\rho_x(a)} = \rho_{\rho_x(a')}$ and $\rho_{\lambda_x(a)} = \rho_{\lambda_x(a')}$.*
- (5) *$\lambda_{\sigma_x(a)} = \lambda_{\sigma_x(a')}$ and $\rho_{\sigma_x(a)} = \rho_{\sigma_x(a')}$.*
- (6) *$\sigma_{\sigma_x(a)} = \sigma_{\sigma_x(a')}$.*

Proof. (1) Since $\lambda_{\lambda_a(x)} \lambda_{\rho_x(a)} = \lambda_a \lambda_x = \lambda_{a'} \lambda_x = \lambda_{\lambda_{a'}(x)} \lambda_{\rho_x(a')} = \lambda_{\lambda_a(x)} \lambda_{\rho_x(a')}$ and, because $\lambda_{\lambda_a(x)}$ is bijective, we get that $\lambda_{\rho_x(a)} = \lambda_{\rho_x(a')}$.

(2) Since $\lambda_{\lambda_x(a)} \lambda_{\rho_a(x)} = \lambda_x \lambda_a = \lambda_x \lambda_{a'} = \lambda_{\lambda_x(a')} \lambda_{\rho_{a'}(x)} = \lambda_{\lambda_x(a')} \lambda_{\rho_a(x)}$ and, because $\lambda_{\rho_a(x)}$ is bijective, we get that $\lambda_{\lambda_x(a)} = \lambda_{\lambda_x(a')}$.

Assume now that A is non-degenerate.

(3) First we claim that $\lambda_x \lambda_{\lambda_x^{-1}(a)} = \lambda_a \lambda_{\rho_{\lambda_x^{-1}(a)}}(x)$. Indeed, by (2.9),

$$\lambda_x \lambda_{\lambda_x^{-1}(a)} = \lambda_{\lambda_x(\lambda_x^{-1}(a))} \lambda_{\rho_{\lambda_x^{-1}(a)}}(x) = \lambda_a \lambda_{\rho_{\lambda_x^{-1}(a)}}(x). \quad (4.3)$$

Using (2.10), we also get

$$\begin{aligned}\rho_{\lambda_x^{-1}(a)}(x) &= \lambda_{\rho_{\lambda_x(\lambda_x^{-1}(a))} \rho_x^{-1}(x)}^{-1} \rho_a \lambda_{\rho_x(\rho_x^{-1}(x))} \lambda_x^{-1}(a) \lambda_{\rho_x^{-1}(x)}(x) \\ &= \lambda_{\rho_a \rho_x^{-1}(x)}^{-1} \rho_a \lambda_{\lambda_x^{-1}(a)} \lambda_{\rho_x^{-1}(x)}(x) \\ &= \lambda_{\rho_a \rho_x^{-1}(x)}^{-1} \rho_a \lambda_{\rho_x^{-1}(x)}(x),\end{aligned}$$

so as $\rho_a = \rho_{a'}$ this implies

$$\begin{aligned}\rho_{\lambda_x^{-1}(a)}(x) &= \lambda_{\rho_a \rho_x^{-1}(x)}^{-1} \rho_a \lambda_{\rho_x^{-1}(x)}(x) \\ &= \lambda_{\rho_{a'} \rho_x^{-1}(x)}^{-1} \rho_{a'} \lambda_{\rho_x^{-1}(x)}(x) \\ &= \rho_{\lambda_x^{-1}(a')}(x).\end{aligned}$$

Hence,

$$\lambda_x \lambda_{\lambda_x^{-1}(a)} \stackrel{(4.3)}{=} \lambda_a \lambda_{\rho_{\lambda_x^{-1}(a)}}(x) = \lambda_a \lambda_{\rho_{\lambda_x^{-1}(a')}}(x) = \lambda_{a'} \lambda_{\rho_{\lambda_x^{-1}(a')}}(x) \stackrel{(4.3)}{=} \lambda_x \lambda_{\lambda_x^{-1}(a')}.$$

Since λ_x is bijective, we conclude that $\lambda_{\lambda_x^{-1}(a)} = \lambda_{\lambda_x^{-1}(a')}$. Similarly, one can prove that $\rho_{\lambda_x^{-1}(a)} = \rho_{\lambda_x^{-1}(a')}$.

(4) Because $\rho_{\rho_x(a)} \rho_{\lambda_a(x)} = \rho_x \rho_a = \rho_x \rho_{a'} = \rho_{\rho_x(a')} \rho_{\lambda_{a'}(x)} = \rho_{\rho_x(a')} \rho_{\lambda_a(x)}$ and, because of the assumption that $\rho_{\lambda_a(x)}$ is bijective, we get that $\rho_{\rho_x(a)} = \rho_{\rho_x(a')}$.

Because $\rho_{\rho_a(x)}\rho_{\lambda_x(a)} = \rho_a\rho_x = \rho_{a'}\rho_x = \rho_{\rho_{a'}(x)}\rho_{\lambda_x(a')} = \rho_{\rho_a(x)}\rho_{\lambda_x(a')}$ and, because of the assumption that $\rho_{\rho_a(x)}$ is bijective, we get that $\rho_{\lambda_x(a)} = \rho_{\lambda_x(a')}$.

(5) Recall that $\sigma_x(a) = \lambda_x\rho_{\lambda_x^{-1}(x)}(a)$. Hence, $\lambda_{\sigma_x(a)} = \lambda_{\lambda_x\rho_{\lambda_x^{-1}(x)}(a)}$. Substitute $e = \lambda_x^{-1}(x) = \lambda_{a'}^{-1}(x)$ then we get $\lambda_{\sigma_x(a)} = \lambda_{\lambda_{\lambda_a(e)}\rho_e(a)}$. Hence statement (5) is equivalent with $\lambda_{\lambda_{\lambda_a(e)}\rho_e(a)} = \lambda_{\lambda_{\lambda_{a'}(e)}\rho_e(a')} = \lambda_{\lambda_{\lambda_a(e)}\rho_e(a')}$ for all $e \in A$. Now, from part (1) we know that $\lambda_{\rho_e(a)} = \lambda_{\rho_e(a')}$ and from part (4) we also get that $\rho_{\rho_e(a)} = \rho_{\rho_e(a')}$. Therefore, by part (2), $\lambda_{\lambda_{\lambda_a(e)}\rho_e(a)} = \lambda_{\lambda_{\lambda_a(e)}\rho_e(a')}$, as desired.

Repeat the steps in the previous, but replace part (2) by part (4), i.e. if ρ_x is bijective and $\lambda_a = \lambda_{a'}$, $\rho_a = \rho_{a'}$ then $\rho_{\lambda_x(a)} = \rho_{\lambda_x(a')}$.

(6) Note that because of part (5) and Proposition 4.3 we immediately obtain that $\sigma_{\sigma_b(a)} = \sigma_{\sigma_b(a')}$. \square

Theorem 4.5. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. Then $\mathcal{G}(A) = \{f(a) = (\sigma_a, \lambda_a, \rho_a) \mid a \in A\}$ is a YB-semitruss for the operations*

$$f(a) \circ f(b) = (\sigma_{a \circ b}, \lambda_{a \circ b}, \rho_{a \circ b}), \quad f(a) + f(b) = (\sigma_{a+b}, \lambda_{a+b}, \rho_{a+b}),$$

λ -map

$$\lambda_{f(a)}f(b) = (\sigma_{\lambda_a(b)}, \lambda_{\lambda_a(b)}, \rho_{\lambda_a(b)}) = f(\lambda_a(b)).$$

and σ -map

$$\sigma_{f(b)}f(a) = (\sigma_{\sigma_b(a)}, \lambda_{\sigma_b(a)}, \rho_{\sigma_b(a)}) = f(\sigma_b(a)).$$

The map $f : A \rightarrow \mathcal{G}(A) : a \mapsto f(a)$ is a YB-semitruss epimorphism and the associated solution of $\mathcal{G}(A)$ is non-degenerate.

Note that because of Proposition 4.3 the natural mapping $\mathcal{G}(A) \rightarrow \{(\lambda_a, \rho_a) \mid a \in A\}$ is bijective and thus the latter can be considered as a YB-semitruss.

Proof. That the mentioned operations are well-defined has been proven in the beginning of this section in Lemma 4.1. That the λ -map is well-defined follows from Lemma 4.2 and that the σ -map is well-defined follows from Lemma 4.4.

Note that, for each $a \in A$ we know from Lemma 4.2 that $\lambda_{f(a)}$ is surjective. To prove it also is injective assume that $\lambda_{f(a)}(f(b)) = \lambda_{f(a)}(f(b'))$. Then $\lambda_{\lambda_a(b)} = \lambda_{\lambda_a(b')}$ and $\rho_{\lambda_a(b)} = \rho_{\lambda_a(b')}$ and thus it follows from Lemma 4.4.(2) that $\lambda_b = \lambda_{\lambda_a^{-1}\lambda_a(b)} = \lambda_{\lambda_a^{-1}\lambda_a(b')} = \lambda_{b'}$. Similarly, we get that $\rho_b = \rho_{b'}$. From Proposition 4.3 it follows that also $\sigma_b = \sigma_{b'}$. Hence, $f(b) = f(b')$, as desired.

It is now straightforward to verify that all required conditions for $\mathcal{G}(A)$ to be a YB-semitruss are satisfied (in particular, $\lambda : \mathcal{G}(A) \rightarrow \text{Aut}(\mathcal{G}(A), +)$) and that $f : A \rightarrow \mathcal{G}(A)$ is an epimorphism of YB-semitrusses.

It remains to show that the associated solution of $\mathcal{G}(A)$ is right non-degenerate. We prove this via Lemma 2.9. Hence, because of Proposition 4.3, we need to prove the following: if $f(a) = f(a')$, $f(b) = f(b')$ then $\lambda_{\rho_a^{-1}(b)} = \lambda_{\rho_{a'}^{-1}(b')}$ and $\rho_{\rho_a^{-1}(b)} = \rho_{\rho_{a'}^{-1}(b')}$. Clearly, without loss of generality, we may assume that $a = a'$. Hence we need to prove Lemma 4.4.(3) for ρ and λ interchanged. For this notice that if $r : X \times X \rightarrow X \times X : (x, y) \mapsto (\lambda_x(y), \rho_y(x))$ is a solution then so is $\tau r \tau : X^2 \times X^2 : (x, y) \mapsto (\rho_x(y), \lambda_y(x))$, where $\tau(x, y) = (y, x)$. Indeed, $r_{12}r_{23}r_{12} = r_{23}r_{12}r_{23}$ if and only if $(\tau r_{12}\tau)(\tau r_{23}\tau)(\tau r_{12}\tau) = (\tau r_{23}\tau)(\tau r_{12}\tau)(\tau r_{23}\tau)$. Hence we indeed may interchange λ and ρ and the result follows. \square

Corollary 4.6. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. Then $(\mathcal{G}(A), +, \circ, \lambda, \sigma)$ is a YB-semitruss with $(\mathcal{G}(A), \circ)$ a cancellative monoid (and thus also $(\mathcal{G}(A), +)$ is left cancellative) that satisfies the left and right Ore condition.*

Proof. To prove left cancellativity, suppose $f(a) \circ f(b) = f(a) \circ f(b')$. Then $\lambda_a\lambda_b = \lambda_a\lambda_{b'}$ and $\rho_a\rho_b = \rho_a\rho_{b'}$. As λ_a and ρ_a are bijective we obtain that $\lambda_b = \lambda_{b'}$ and $\rho_b = \rho_{b'}$. Because of Proposition 4.3 we then also have $\sigma_b = \sigma_{b'}$. Hence $f(b) =$

$f(b')$, as desired. So both semigroups $(\mathcal{G}(A), \circ)$ and $(\mathcal{G}(A), +)$ are left cancellative. Similarly one proves that $(\mathcal{G}(A), \circ)$ is right cancellative. Indeed, suppose $f(a) \circ f(b) = f(a') \circ f(b)$. Then, $\lambda_a \lambda_b = \lambda_{a'} \lambda_b$ and $\rho_a \rho_b = \rho_{a'} \rho_b$. As λ_b and ρ_b are bijective it follows that $\lambda_a = \lambda_{a'}$ and $\rho_a = \rho_{a'}$. Hence, also $\sigma_a = \sigma_{a'}$ and thus $f(a) = f(a')$, as desired.

Since $\mathcal{G}(A)$ is a non-degenerate YB-semitruss, we obtain from Proposition 2.6 that $f(a) \circ f(b) = \lambda_{f(a)}(f(b)) \circ \rho_{f(b)}(f(a))$, where ρ denotes the ρ -map for $\mathcal{G}(A)$. As each $\lambda_{f(a)}$ and $\rho_{f(b)}$ is bijective, the left and right condition follow at once. \square

We are now in a position to extend Proposition 3.5 to a much larger class of solutions. For this, in the non-degenerate case, we consider $\{(\lambda_a, \rho_a) \mid a \in A\}$ as a subsemigroup of the direct product of $(\text{Sym}(A), \circ)$ and its opposite group $(\text{Sym}(A), \circ^{op})$, where \circ is the composition of functions.

Corollary 4.7. *Let $(A, +, \circ, \lambda, \sigma)$ be a non-degenerate YB-semitruss. If for every $a \in A$, there exists $b \in A$ such that $\lambda_a \lambda_b = \text{id}$ and $\rho_a \rho_b = \text{id}$ (for example if all λ_a and ρ_a are of finite order), then the associated solution r_A is bijective and $\mathcal{G}(A)$ is a skew left brace.*

Proof. Note that it follows from Proposition 4.3 that if $\{(\lambda_a, \rho_a) \mid a \in A\}$ is a group then also $(\mathcal{G}(A), \circ)$ is a group. Indeed, if $a \in A$ then, by assumption, $(\lambda_a, \rho_a)(\lambda_b, \rho_b) = (\text{id}, \text{id})$ for some $b \in A$. Hence $\lambda_{a \circ b} = \rho_{a \circ b} = \text{id}$. So, by Proposition 4.3, $\sigma_{a \circ b} = \text{id}$. Hence $f(a) \circ f(b) = f(a \circ b) = \text{id}$. It follows that $(\mathcal{G}(A), \circ)$ is indeed a group. In particular, all maps σ_a are bijective and thus, by Proposition 2.4, r_A is bijective.

The last part of the result now follows at once from Example 2.21 and Corollary 4.6. \square

Corollary 4.8. *Let (X, r) be a non-degenerate solution. If the subsemigroup $\langle (\lambda_x, \rho_x) \mid x \in X \rangle$ of the group $(\text{Sym}(X), \circ) \times (\text{Sym}(X), \circ^{op})$ is a group itself then r is bijective.*

Proof. As before, the structure semigroup $S(X, r)$ is a non-degenerate YB-semitruss and, because of the assumptions, $\mathcal{G}(S(X, r))$ is a group. Hence, by Corollary 4.7, $r_{S(X, r)}$ is bijective. Since this map, restricted to X^2 is r , we obtain that r is bijective. \square

5. ALGEBRAIC STRUCTURE OF YB-SEMITRUSSES

5.1. Additive structure of YB-semitruss and the structure algebra.

In this section we investigate the algebra $K[(A, \circ)]$ of a YB-semitruss $(A, +, \circ, \lambda, \sigma)$. Our main result states that this algebra is left Noetherian and satisfies a polynomial identity provided A is a unital strongly \mathbb{N} -graded YB-semitruss with A_1 finite, $A_0 = \{1\}$ and the diagonal map is bijective. As such an algebra is an epimorphic image of the structure algebra $K[M(X, r)]$, with (X, r) a left non-degenerate finite solution, it is sufficient to deal with the unital structure YB-semitruss $M(X, r)$. To prove this result we first determine the algebraic structure of the derived monoid $A(X, r)$.

Put

$$A = A(X, r) = \langle x \in X \mid x + y = y + \sigma_y(x) \rangle,$$

the additive monoid of the unital structure YB-semitruss $M(X, r)$. Recall from Lemma 2.2 that for a unital YB-semitruss we have $0 = 1$, where 0 and 1 are the identities of the additive and the multiplicative monoid respectively. For any $a, b \in A$,

$$a + b = b + \sigma_b(a), \tag{5.1}$$

where $\sigma : (A, +) \rightarrow \text{End}(A, +) : a \mapsto \sigma_a$, is a monoid anti-homomorphism. In particular, each right ideal $b + A$ of $(A, +)$ is a two-sided ideal of $(A, +)$. One says that $a \in A$ is *divisible* by b if $a = e + b + d$ for some $e, d \in A$. If $e = 0$ then one says that a is *left divisible* by b . Because of (5.1), a is divisible by b if and only if a is left divisible by b .

Consider the subsemigroup

$$\mathcal{C} = \mathcal{C}(A) := \{\sigma_a \mid a \in A\} = \langle \sigma_x \mid x \in X \rangle,$$

of $\text{End}(A, +)$. Furthermore, for $a, b \in A$,

$$\sigma_a \sigma_b = \sigma_{b+a} = \sigma_{a+\sigma_a(b)} = \sigma_{\sigma_a(b)} \sigma_a, \quad (5.2)$$

and thus

$$\sigma_a \mathcal{C} \subseteq \mathcal{C} \sigma_a. \quad (5.3)$$

So, every left ideal of the semigroup \mathcal{C} is a two-sided ideal. If \mathcal{C} is finite (for example if X is finite), there exists a positive integer, say v , so that

$$\sigma_x^v = \sigma_{vx} = \sigma_{vx}^2 \text{ is an idempotent,}$$

for each $x \in X$.

From now on $X = \{x_1, \dots, x_n\}$ is a finite set and thus any element of $A = A(X, r)$ is divisible by only finitely many elements. Let $a \in A = \langle x_1, \dots, x_n \rangle$ and let m_1 be the maximal non-negative integer so that a is left divisible by $m_1 x_1$, that is $a = m_1 x_1 + b$ and $b \in A$ can not be written as $x_1 + d$ for some $d \in A$. Repeat this argument on b and consider the maximal non-negative integer m_2 so that $b = m_2 x_2 + d$, for some $d \in A$. After at most n steps we get that

$$a = m_1 x_1 + m_2 x_2 + \dots + m_n x_n,$$

for some non-negative integers m_i . Notice that $vx_i + vx_j = vx_j + \sigma_{vx_j}(vx_i) = vx_j + v \sigma_{vx_j}(x_i)$ and $\sigma_{vx_j}(x_i) \in X$. Hence,

$$B(v) = \{m_1 vx_1 + \dots + m_n vx_n \mid m_1, \dots, m_n \geq 0\},$$

is a submonoid of $(A, +)$. Furthermore

$$A = B(v) + F(v), \text{ with } F(v) = \{m_1 x_1 + \dots + m_n x_n \mid v > m_1, \dots, m_n \geq 0\}. \quad (5.4)$$

Thus A is a finite module over the finitely generated submonoid $B(v)$. For $1 \leq i \leq n$, put

$$y_i = vx_i \quad \text{and} \quad X(v) = \{y_1, \dots, y_n\}.$$

The solution $s : X^2 \rightarrow X^2 : (x, y) \mapsto (y, \sigma_y(x))$ induces a solution $s_{X(v)} : X(v)^2 \rightarrow X(v)^2 : (y_i, y_j) \mapsto (y_j, \sigma_{y_j}(y_i))$. Clearly,

$$B(v) = A(X(v), s_{X(v)}) = \langle y_1, \dots, y_n \rangle.$$

Put

$$\sigma_j = \sigma_{y_j}, \quad \text{for } 1 \leq j \leq n.$$

Because of (5.2) and since each element σ_y , for $y \in Y$ is idempotent, any non-identity element of $\mathcal{C}(B(v))$ can be written as $\sigma_{j_1} \dots \sigma_{j_k}$ for some j_1, \dots, j_k and some $k \leq n$. If v is clear from the context we simply write $B(v)$ as B .

Let $\sigma_{j_1} \sigma_{j_2} \dots \sigma_{j_k} \in \mathcal{C}(B)$ with $J = \{j_1, \dots, j_k\}$ a subset of $\{1, \dots, n\}$. Then, for any $j_l \in J$, we have

$$\begin{aligned} \sigma_{j_1} \dots \sigma_{j_k} &= \sigma_{\sigma_{j_l}(y_{j_{l+1}})} \sigma_{j_l} \sigma_{j_{l+2}} \dots \sigma_{j_k} \\ &= \sigma_{\sigma_{j_l}(y_{j_{l+1}})} \sigma_{\sigma_{j_l}(y_{j_{l+2}})} \dots \sigma_{\sigma_{j_l}(y_{j_k})} \sigma_{j_l}. \end{aligned}$$

Because σ_{j_l} and σ_{j_k} are idempotent we get

$$\begin{aligned} \sigma_{j_l} \cdots \sigma_{j_k} \sigma_{j_l} &= \sigma_{\sigma_{j_l}(y_{j_{l+1}})} \sigma_{\sigma_{j_l}(y_{j_{l+2}})} \cdots \sigma_{\sigma_{j_l}(y_{j_k})} \sigma_{j_l} \sigma_{j_l} \\ &= \sigma_{\sigma_{j_l}(y_{j_{l+1}})} \sigma_{\sigma_{j_l}(y_{j_{l+2}})} \cdots \sigma_{\sigma_{j_l}(y_{j_k})} \sigma_{j_l} \\ &= \sigma_{j_l} \cdots \sigma_{j_k} \end{aligned} \tag{5.5}$$

Hence it follows from (5.5) that each $\sigma_{j_1} \cdots \sigma_{j_k}$ is an idempotent and we have shown the following property.

Lemma 5.1. *The semigroup $\mathcal{C}(B(v)) = \{\sigma_{j_1} \sigma_{j_2} \cdots \sigma_{j_k} \mid 1 \leq j_1, \dots, j_k \leq n\} \cup \{\text{id}\} = \{\sigma_{j_1} \sigma_{j_2} \cdots \sigma_{j_k} \mid 1 \leq j_1, \dots, j_k \leq n, \text{ all distinct}\} \cup \{\text{id}\}$ is a band, i.e. a semigroup consisting of idempotents.*

For $1 \leq k \leq n$ and $\kappa \in \text{Sym}_n$, the symmetric group of degree n , put

$$\begin{aligned} t_k &= y_1 + \cdots + y_k \in A, \\ \kappa(t_k) &= y_{\kappa(1)} + \cdots + y_{\kappa(k)} \in A, \\ B_{\kappa(t_k)} &= \langle y_{\kappa(i)} \mid 1 \leq i \leq k \rangle, \end{aligned}$$

and put

$$T = \{\kappa(t_k) \mid 1 \leq k \leq n, \kappa \in \text{Sym}_n\}.$$

Let $0 \neq w \in B$. Assume $w = w' + y_k$ for some $k \leq n$. Let m be the maximal non-negative integer so that w' is left divisible by my_k . Then $w = my_k + w'' + y_k$ for some $w'' \in B$ that is not left divisible by y_k . If $0 \neq w''$ then repeat the argument on w'' and after at most n steps we get that

$$w = w_t + t,$$

for some $t \in T$ and $w_t \in B_t$. Since $\sigma_w = \sigma_t \sigma_{w_t}$, it follows from (5.5) that

$$\sigma_w = \sigma_t.$$

Hence, for $a, b \in B_t$

$$a + b + t = a + t + \sigma_t(b) = a + t + \sigma_{a+t}(b) = b + a + t, \tag{5.6}$$

in particular

$$a + t + b + t = b + t + a + t,$$

that is, each $B_t + t$ is an abelian semigroup. With the notation introduced above we thus obtain the following property.

Lemma 5.2. *Let (X, r) be a finite left non-degenerate solution and $A = A(X, r)$. Then $A = B(v) + F(v)$ and $B(v) = \{0\} \cup \bigcup_{t \in T} (B(v)_t + t)$, a finite union of abelian subsemigroups.*

We also have that for $\kappa \in \text{Sym}_n$, $1 \leq k \leq n$ and $1 \leq l \leq k$

$$\kappa(t_k) + y_{\kappa(l)} = y_{\eta(l')} + \eta(t_k) \tag{5.7}$$

for some $\eta \in \text{Sym}_n$ and $1 \leq l' \leq k$.

To prove this it is sufficient to deal with the case that $\kappa = \text{id}$. Hence, we need to show that for $1 \leq l \leq k \leq n$ one can rewrite $t_k + y_l = y_1 + y_2 + \cdots + y_k + y_l$ as $y_{\eta(l')} + \eta(t_k)$ for some $\eta \in \text{Sym}_n$ and $1 \leq l' \leq k$. If $l = 1$ then this is obvious. Assume $k > l > 1$. Then

$$t_k + y_l = y_l + \cdots + y_k + y_l + \sigma_{y_l + \cdots + y_k + y_l}(y_1 + \cdots + y_{l-1}).$$

Hence, since σ is an antihomomorphism and because of (5.5),

$$\begin{aligned} t_k + y_l &= y_l + y_{l+1} + \cdots + y_k + y_l + \sigma_{y_{l+1} + \cdots + y_k + y_l}(y_1 + \cdots + y_{l-1}) \\ &= y_l + y_1 + \cdots + y_{l-1} + y_{l+1} + \cdots + y_k + y_l \\ &= y_l + \eta(t_k) \\ &= y_{\eta(l')} + \eta(t_k), \end{aligned}$$

for some η and $l' = k$, as desired. It remains to deal with the case $k = l$. In this case

$$t_k + y_k = y_k + y_k + \sigma_{y_k + y_k}(y_1 + \cdots + y_{k-1}),$$

and thus, because σ_{y_k} is an idempotent and σ is an anti-homomorphism,

$$\begin{aligned} t_k + y_k &= y_k + y_k + \sigma_{y_k}(y_1 + \cdots + y_{k-1}) \\ &= y_k + y_1 + \cdots + y_{k-1} + y_k \\ &= y_k + t_k, \end{aligned}$$

again as desired.

Let K be a field. Each $K[B_t + t]$ is a left $K[B_t]$ -module for the module action $K[B_t] \times K[B_t + t] \rightarrow K[B_t + t]$ defined by the linear extension of $(a, d+t) \mapsto a+d+t$, where $a, d \in B_t$ and $t \in T$. Because of (5.6), $K[B_t + t]$ is a cyclic left module over the finitely generated commutative K -algebra $K[B_t]^{ab} = K[B_t]/[K[B_t], K[B_t]]$, i.e. the abelianization of the algebra $K[B_t]$. Hence the commutative (non-unital) algebra $K[B_t + t]$ is a Noetherian left $K[B_t]$ -module.

Obviously, for $\kappa \in \text{Sym}_n$, we have that $B_{\kappa(t_n)} + \kappa(t_n) = B + \kappa(t_n)$ is a left ideal of B . Hence, $K[B_{\kappa(t_n)} + \kappa(t_n)]$ is a left ideal of $K[B]$ and thus it is a Noetherian left $K[B]$ -module and, as ring, it is commutative. Put

$$R_n = \sum_{\kappa \in \text{Sym}_n} K[B_{\kappa(t_n)} + \kappa(t_n)].$$

Clearly this is a Noetherian left $K[B]$ -module. Furthermore, from (5.7) it follows that R_n is an ideal of $K[B]$. As a ring it is a sum of left ideals that are PI-algebras (actually each of the left ideals is commutative as a ring) and thus, by a result of Rowen [53] (or more general a result of Kępczyk [47], which states that a ring which is a sum of two PI rings is again a PI ring), R_n is a PI-algebra.

Next consider the K -algebra $K[B]/R_n$. For any $\kappa \in \text{Sym}_n$, it easily is verified that $K[B_{\kappa(t_{n-1})} + \kappa(t_{n-1})]/R_n$ is a left $K[B]/R_n$ -submodule of $K[B]/R_n$. With a similar reasoning as above, we obtain that this is a Noetherian cyclic left module over $K[B_{\kappa(t_{n-1})}]$ and thus $K[B_{\kappa(t_{n-1})} + \kappa(t_{n-1})]/R_n$ is a Noetherian left module over $K[B]/R_n$. Furthermore, as a ring it is commutative. Put

$$R_{n-1} = \sum_{\kappa \in \text{Sym}_n} K[B_{\kappa(t_{n-1})} + \kappa(t_{n-1})].$$

By (5.7), $R_{n-1} + R_n/R_n$ is an ideal in $K[B]/R_n$ and it is a Noetherian left $K[B]/R_n$ -module. Consequently $R_{n-1} + R_n$ is a Noetherian left $K[B]$ -module. One can repeat this process in an obvious manner making use of the sets

$$R_k = \sum_{\kappa \in \text{Sym}_n} K[B_{\kappa(t_k)} + \kappa(t_k)],$$

where $1 \leq k \leq n$. Put $R_0 = K$. It follows, in particular the algebra $K[B] = R_0 + R_1 + \cdots + R_n$ is left Noetherian. Since a K -algebra V is PI if it has an ideal I so that both algebras I and V/I are PI, we obtain that $K[B]$ also is a PI-algebra. By (5.4), $K[A] = \sum_{f \in F} K[B] + f$. So the K -algebra $K[A]$ is a finite module over the subalgebra $K[B]$ and thus $K[A]$ also is a left Noetherian PI algebra (actually it is a Noetherian left $K[B]$ -module). We thus have proven the following result.

Theorem 5.3. *Let (X, r) be a finite left non-degenerate solution and put $X = \{x_1, \dots, x_n\}$. Let v be a positive integer so that σ_x^v is an idempotent endomorphism for each $x \in X$. Then, with the notation introduced above, the derived monoid $A = A(X, r)$ satisfies the following properties.*

- (1) $A(X, r) = B(v) + F(v)$, i.e. $A(X, r)$ is a finite module over $B(v) = A(Y, s_Y) = \langle y_1 = vx_1, \dots, y_n = vx_n \rangle$.
- (2) $\mathcal{C}(B(v))$ is a band;
- (3) for $1 \leq k \leq n$, $t_k = y_1 + \dots + y_k \in B$ and $\kappa \in \text{Sym}_n$, each $B_{\kappa(t_k)} + \kappa(t_k)$ is a commutative semigroup, where $B_{\kappa(t_k)} = \langle y_{\kappa(i)} \mid 1 \leq i \leq k \rangle$.
- (4) for $1 \leq k \leq n$ and $B_k = \bigcup_{\kappa \in \text{Sym}_n} B_{\kappa(t_k)} + \kappa(t_k)$,

$$B_n \subseteq B_n \cup B_{n-1} \subseteq \dots \subseteq B_n \cup \dots \cup B_2 \cup B_1 \subseteq B,$$

is an ideal chain of $B(v)$,

- (5) each Rees factor semigroup of the ideal chain is a union of left ideals

$$(B_n \cup B_{n-1} \cup \dots \cup B_{i-1} \cup B_{\kappa(t_i)} + \kappa(t_i)) / (B_n \cup B_{n-1} \cup \dots \cup B_{i-1}),$$

with $\kappa \in \text{Sym}_n$.

- (6) A satisfies the ascending chain condition on left ideals.
- (7) for any field K , each factor of the ideal chain

$$\{0\} \subseteq K[B_n] \subseteq K[B_n] + K[B_{n-1}] \subseteq \dots \subseteq K[B_n] + \dots + K[B_1] \subseteq K[B(v)],$$

of $K[B(v)]$ is a Noetherian left $K[B(v)]$ -module that is a finite sum of commutative rings.

In particular, $K[A]$ is a Noetherian left $K[B(v)]$ -module, and $K[A]$ is a left Noetherian PI-algebra of finite Gelfand-Kirillov dimension bounded by n .

In general the algebra $K[A]$ is not right Noetherian. Indeed, again as in Example 2.17, consider solutions (X, r) with $r(x, y) = (y, y)$ for $x, y \in X$. So r is idempotent and left non-degenerate. For simplicity take $X = \{x, y\}$. To avoid confusion of the operations in the structure algebra, we will write $A = A(X, r)$ multiplicatively, i.e. $A = A(X, r) = \langle x, y \mid xy = yy, yx = xx \rangle = \{x^n, y^n \mid n \geq 0\}$. In $K[A]$ we have $(x^n - y^n)a = 0$ for any $a \in A$. Hence $\sum_{n>0} K(x^n - y^n)$ is a right ideal of $K[A]$ that obviously is not finitely generated as a right ideal.

As an application of Theorem 5.3 we claim that the structure algebra $K[M(X, r)]$ is left Noetherian in case (X, r) is a finite left non-degenerate solution, provided the diagonal map \mathfrak{q} is surjective (i.e. $X \rightarrow X : x \mapsto \lambda_x^{-1}(x)$ is surjective). With a standard length and induction argument one can easily show that the latter is equivalent with the diagonal map $\mathfrak{q} : M(X, r) \rightarrow M(X, r) : a \mapsto \lambda_a^{-1}(a)$ being surjective.

From Section 2.1, we know that $M(X, r)$ is a submonoid of the semidirect product $A(X, r) \rtimes \text{Im } \lambda$ and $\text{Im } \lambda = \{\lambda_a \mid a \in M(X, r)\}$ is a finite group. From Theorem 5.3 we get that the algebra $K[A(X, r) \rtimes \text{Im } \lambda]$ is a Noetherian left $K[B(v)]$ -module. Hence, to prove that $K[M(X, r)]$ is left Noetherian it is sufficient to show that one can choose the positive integer v in the statement of Theorem 5.3 so that $B(v) \subseteq M(X, r)$, i.e. for all $x \in X$ we also may assume that $\lambda_{vx} = \text{id}$, and thus $M(X, r)$ is a finite module over $B(v)$. This is what will be shown in the following result.

Also recall that an algebra over a field is said to be *representable* if it can be embedded into a matrix algebra over some field. A well-known result of Ananin [2] states that any finitely generated left Noetherian PI-algebra over a field is representable. Conversely, any representable algebra is a PI-algebra.

Corollary 5.4. *Let (X, r) be a finite left non-degenerate solution and let K be a field. If the diagonal map $\mathfrak{q} : X \rightarrow X$ is bijective, then there exists a positive integer v so that σ_x^v is an idempotent endomorphism for each $x \in X$ and $B(v) \subseteq M(X, r)$.*

Consequently, $M(X, r) = B(v) + F$, for some finite subset F of $M(X, r)$ and the structure algebra $K[M(X, r)]$ is a connected graded left Noetherian representable algebra. In particular, so is the algebra $K[(A, \circ)]$ for any unital strongly \mathbb{N} -graded YB-semitruss $(A, +, \circ, \lambda, \sigma)$ with A_1 finite, $A_0 = \{1\}$ and with bijective diagonal map.

Proof. From Theorem 5.3 we know that $A = A(X, r)$ is a finite module over the left Noetherian ring $K[B]$, where $B = B(v)$ and v is any positive integer so that each σ_x^v is an idempotent, for $x \in X$. Without loss of generality, replacing v by a multiple if needed, we also may assume that $\lambda_a^v = \text{id}$, for each $a \in M(X, r)$.

Now, for $x \in X$, define $x^{(1)} = x$ and recursively $x^{(i+1)} = \mathfrak{q}(x^{(i)}) = \lambda_{x^{(i)}}^{-1}(x^{(i)})$. Since, by assumption, \mathfrak{q} is a bijective map and thus it is of finite order, i.e. $\mathfrak{q}^m = \text{id}$, for some positive integer m . So $x^{(m+1)} = x^{(1)} = x$. Since for any positive integer k we have that $kx = x^{(1)} \circ x^{(2)} \circ \dots \circ x^{(k)}$, we obtain that $vmx = (x^{(1)} \circ x^{(2)} \circ \dots \circ x^{(m)})^v$ and thus $\lambda_{vmx} = (\lambda_x \lambda_{x^{(1)}} \dots \lambda_{x^{(m)}})^v = \text{id}$. So $B(mv) \subseteq M(X, r)$, as desired.

Because of Corollary 2.14, for any strongly \mathbb{N} -graded unital YB-semitruss $(A, +, \circ, \lambda, \sigma)$ with bijective diagonal map and with A_1 finite and $A_0 = \{1\}$, we have that the algebra $K[(A, \circ)]$ is a graded epimorphic image of the structure algebra $K[M(A_1, r_{A_1})]$ for the left non-degenerate finite solution (A_1, r_{A_1}) with bijective diagonal map. Hence, the last statement of the result follows from the first one. \square

In case (X, r) is a finite left non-degenerate bijective solution (so a finite non-degenerate bijective solution, by Theorem 3.1), then each element of $A(X, r)$ is a normal element, as each σ_a is bijective, and thus $\mathcal{C}(A)$ is a group with id as the only idempotent. Hence, for an appropriate choice of v we get that $B(v) \subseteq (Z(A) \cap M(X, r))$. It follows that $w \circ B(v) = B(v) \circ w$ for each element $w \in M(X, r)$. Hence one obtains Theorem 4 in [40] (see also [39]); in particular $K[M(X, r)]$ is a module-finite normal extension of the commutative affine subalgebra $K[B(v)]$ and thus $K[M(X, r)]$ is a left and right Noetherian PI-algebra. Note that the bijective left non-degenerate assumption implies that the diagonal map is bijective, because of Theorem 3.1 and Lemma 3.2.

5.2. YB-semitrusses with a left simple semigroup of endomorphisms and their solutions.

Let $A = (A, +, \circ, \lambda, \sigma)$ be a YB-semitruss and assume $\mathcal{C} = \{\sigma_a \mid a \in A\}$ is a finite subsemigroup of $\text{End}(A, +)$. We know from (5.3) that every left ideal of \mathcal{C} is a two-sided ideal. It is then well known that this semigroup has an ideal chain

$$C_0 \subseteq C_1 \cdots \subseteq C_{n-1} \subseteq C_n = \mathcal{C},$$

with $C_0 = \{\theta\}$ or C_0 the empty set, and all factors C_i/C_{i-1} are either a null semigroup or completely (θ) -simple. In the latter case we know that such a semigroup is of the form $\mathcal{M}^0(G, k, l, P)$, with G a maximal subgroup of \mathcal{C} and P a regular sandwich matrix. Since every element of such a semigroup has an idempotent as left identity, it is easy to see that $R \cup C_{i-1}$, where R is a ‘‘column’’ of this semigroup, is a left ideal of \mathcal{C} . But since this must be a two-sided ideal, we get that $l = 1$.

So the principal factors that are completely (θ) -simple (or possibly simple in case of C_1) are of the form $\mathcal{M}^0(G, k, 1, P)$, and after normalising, we also may assume that each entry of the row matrix P is the identity of G . So, this semigroup is a finite union of finite groups and the idempotents form a subsemigroup so that $ef = e$ for any two non-zero idempotents.

In this subsection we will deal with the case that \mathcal{C} is left simple (for example if $(A, +)$ is a right simple semigroup), i.e. $\mathcal{C} = C_1$ and $C_0 = \emptyset$ and show that $(A, +)$ is determined by skew left braces (actually racks).

So, from now on, assume \mathcal{C} is a finite left simple semigroup and thus

$$\mathcal{C} = G_1 \cup \dots \cup G_k,$$

a disjoint union of finite groups, say with identity $1_1, \dots, 1_k$ respectively. We have

$$G_i G_j \subseteq G_i \quad \text{and} \quad 1_i 1_j = 1_i. \quad (5.8)$$

Consider the anti-homomorphism $\sigma : (A, +) \rightarrow \text{End}(A, +)$ and define the following left ideals of $(A, +)$, for $1 \leq i \leq n$,

$$A_i = \sigma^{-1}(G_i).$$

For $a \in A_j$ and $b \in A_i$, since $\sigma_b \sigma_a = \sigma_{\sigma_b(a)} \sigma_b$, and thus, by (5.8), we get that $\sigma_{\sigma_b(a)} \in G_i$, i.e. $\{\sigma_b(a) \mid \sigma_b \in G_i, a \in A\} \subseteq A_i$. Because $1_i \sigma_b = \sigma_b$ the set $\{\sigma_b(a) \mid \sigma_b \in G_i, a \in A\}$ is a subsemigroup of $(A, +)$. We will denote it as $G_i(A)$. For a subset T of A we denote by $G_i(T)$ the set $\{\sigma_b(t) \mid \sigma_b \in G_i, t \in T\}$. Note that $G_i(A_i) \subseteq G_i(A) = G_i 1_i(A) \subseteq G_i(A_i)$ and thus

$$G_i(A) = G_i(A_i).$$

We also have that

$$a + b = b + \sigma_b(a) = \sigma_b(a) + \sigma_{\sigma_b(a)}(b). \quad (5.9)$$

Hence it follows that $a + b \in G_i(A_i)$ and that each $G_i(A_i)$ is a left ideal of the semigroup $(A, +)$. Furthermore, we get the following restrictions of the derived solution of A ,

$$\sigma_{j,i} : A_j \times A_i \rightarrow A_i \times A_i : (a, b) \mapsto (b, \sigma_b(a)).$$

In particular, each

$$\sigma_{i,i} : A_i \times A_i \rightarrow A_i \times A_i : (a, b) \mapsto (b, \sigma_b(a)),$$

is a solution. Clearly this map restricts to solutions

$$\tilde{\sigma}_{i,i} : G_i(A_i) \times G_i(A_i) \rightarrow G_i(A_i) \times G_i(A_i).$$

As \mathcal{C} is finite, there exists a positive integer, say v , so that if $b \in A_i$ then $\sigma_b^v = 1_i$. Hence, it follows that

$$\sigma_b = 1_i \sigma_b = \sigma_b^{v+1} = \sigma_{\sigma_b^v(b)} \sigma_b^v = \sigma_{1_i(b)} 1_i = \sigma_{1_i(b)}.$$

So

$$\sigma_{j,i} : A_j \times A_i \rightarrow A_i \times A_i : (a, b) \mapsto (b, \sigma_{1_i(b)}(a)).$$

Now for $a \in A_i$ we have that $\sigma_a(G_i(A_i)) \subseteq G_i(A_i)$ and 1_i acts as the identity on $G_i(A_i)$. As G_i is a group we get that each map $\sigma_a : G_i(A_i) \rightarrow G_i(A_i)$ is bijective (and of finite order, as \mathcal{C} is finite). Hence the restriction

$$\tilde{\sigma}_{i,i} : G_i(A_i) \times G_i(A_i) \rightarrow G_i(A_i) \times G_i(A_i),$$

is a bijective non-degenerate solution and thus $G_i(A_i)$ is a normalising semigroup, i.e. $G_i(A_i) + a = a + G_i(A_i)$ for all $a \in A_i$.

Furthermore, for $a \in A_i$ and $b \in A$, we have that $\sigma_a(b) = \sigma_a(1_i(b))$. Hence, the solution $\sigma_{i,i}$ on A_i is determined by its bijective non-degenerate subsolution on $G_i(A_i)$ and the projections 1_i .

To conclude, we have proved the first part of the following result.

Theorem 5.5. *Let $A = (A, +, \circ, \lambda, \sigma)$ be a YB-semitruss. Assume \mathcal{C} is a finite left simple semigroup, let $E = E(\mathcal{C})$ be the subsemigroup consisting of the idempotents and G_e the maximal subgroup of \mathcal{C} containing $e \in E$. Then, $A = \cup_{e \in E} A_e$, a disjoint union of left ideals $A_e = \sigma^{-1}(G_e)$, so that the derived solution s_A associated to A restricts to a bijective non-degenerate solution on $G_e(A_e)$, say s_{A_e} and for $a_e \in A_e$, $a_f \in A_f$ we have $s_A(a_e, a_f) = s_{A_f}(f(a_e), a_f) = (a_f, \sigma_{f(a_f)}(f(a_e)))$, i.e. the derived solution is determined by bijective non-degenerate solutions.*

If, furthermore, A is strongly \mathbb{N} -graded with A_1 finite if $A_0 = \emptyset$ then, for each $e \in E(\mathcal{C})$, $A_e \setminus G_e(A_e)$ is finite and $G_e(A_e)$ also is strongly \mathbb{N} -graded.

Proof. It only remains to prove the second part of the statement. So, assume A is strongly \mathbb{N} -graded with A_1 finite if $A_0 = \emptyset$. Let $e \in E$. Because of (5.9) each element of $A_e \setminus (A_0 \cup A_1)$ belongs to $G_e(A_e)$. Clearly A_e is strongly \mathbb{N} -graded and hence so is $G_e(A_e)$. Hence the result follows. \square

ACKNOWLEDGEMENT

The first named author is supported by the Engineering and Physical Sciences Research Council [grant number EP/V005995/1]. The second author was supported in part by Onderzoeksraad of Vrije Universiteit Brussel and Fonds voor Wetenschappelijk Onderzoek (Flanders), grant G016117. The third author is supported by Fonds voor Wetenschappelijk Onderzoek (Flanders), via an FWO post-doctoral fellowship, grant 12ZG221N. The fourth author is supported by Fonds voor Wetenschappelijk Onderzoek (Flanders), via an FWO Aspirant-fellowship, grant 11C9421N.

REFERENCES

- [1] J. Almeida, J.-E. Pin, and P. Weil. Semigroups whose idempotents form a subsemigroup. *Math. Proc. Cambridge Philos. Soc.*, 111(2):241–253, 1992.
- [2] A. Z. Anan'in. An intriguing story about representable algebras. In *Ring theory 1989 (Ramat Gan and Jerusalem, 1988/1989)*, volume 1 of *Israel Math. Conf. Proc.*, pages 31–38. Weizmann, Jerusalem, 1989.
- [3] I. Angiono, C. Galindo, and L. Vendramin. Hopf braces and Yang-Baxter operators. *Proc. Amer. Math. Soc.*, 145(5):1981–1995, 2017.
- [4] D. Bachiller. Extensions, matched products, and simple braces. *J. Pure Appl. Algebra*, 222(7):1670–1691, 2018.
- [5] D. Bachiller. Solutions of the Yang-Baxter equation associated to skew left braces, with applications to racks. *J. Knot Theory Ramifications*, 27(8):1850055, 36, 2018.
- [6] D. Bachiller, F. Cedó, and E. Jespers. Solutions of the Yang-Baxter equation associated with a left brace. *J. Algebra*, 463:80–102, 2016.
- [7] R. J. Baxter. Partition function of the eight-vertex lattice model. *Ann. Physics*, 70(1):193–228, 1972.
- [8] T. Brzeziński. Towards semi-trusses. *Rev. Roumaine Math. Pures Appl.*, 63(2):75–89, 2018.
- [9] M. Castelli, F. Catino, and P. Stefanelli. Left non-degenerate set-theoretic solutions of the yang-baxter equation and dynamical extensions of q-cycle sets. *Journal of Algebra and Its Applications*, 2021, doi:10.1142/S0219498822501547.
- [10] F. Catino, I. Colazzo, and P. Stefanelli. Semi-braces and the Yang-Baxter equation. *J. Algebra*, 483:163–187, 2017.
- [11] F. Catino, I. Colazzo, and P. Stefanelli. The matched product of set-theoretical solutions of the Yang-Baxter equation. *J. Pure Appl. Algebra*, 224(3):1173–1194, 2020.
- [12] F. Catino, I. Colazzo, and P. Stefanelli. The matched product of the solutions to the Yang-Baxter equation of finite order. *Mediterr. J. Math.*, 17(2):Paper No. 58, 22, 2020.
- [13] F. Catino, I. Colazzo, and P. Stefanelli. Set-theoretic solutions to the Yang-Baxter equation and generalized semi-braces. *Forum Mathematicum*, 2021.
- [14] F. Catino, M. Mazzotta, and P. Stefanelli. Inverse semi-braces and the Yang-Baxter equation. *J. Algebra*, 573:576–619, 2021.
- [15] F. Cedó. Left braces: solutions of the Yang-Baxter equation. *Adv. Group Theory Appl.*, 5:33–90, 2018.

- [16] F. Cedó, E. Jespers, and A. del Río. Involutive Yang-Baxter groups. *Trans. Amer. Math. Soc.*, 362(5):2541–2558, 2010.
- [17] F. Cedó, E. Jespers, and J. Okniński. Retractability of set theoretic solutions of the Yang-Baxter equation. *Adv. Math.*, 224(6):2472–2484, 2010.
- [18] F. Cedó, E. Jespers, and J. Okniński. Braces and the Yang–Baxter equation. *Commun. Math. Phys.*, 327:101–116, 2014.
- [19] F. Cedó, E. Jespers, and J. Okniński. Set-theoretic solutions of the Yang-Baxter equation, associated quadratic algebras and the minimality condition. *Rev. Mat. Complut.*, 34(1):99–129, 2021.
- [20] F. Cedó, E. Jespers, and C. Verwimp. Structure monoids of set-theoretic solutions of the Yang-Baxter equation. *Publ. Mat.*, 65:499–528, 2021.
- [21] F. Cedó, E. Jespers, Lukasz Kubat, A. Van Antwerpen, and C. Verwimp. On various types of nilpotency of the structure monoid and group of a set-theoretic solution of the Yang–Baxter equation. *Arxiv:2011.01724*, 2020.
- [22] F. Cedó and J. Okniński. Gröbner bases for quadratic algebras of skew type. *Proc. Edinb. Math. Soc. (2)*, 55(2):387–401, 2012.
- [23] F. Chouraqui. Garside groups and Yang-Baxter equation. *Comm. Algebra*, 38(12):4441–4460, 2010.
- [24] A. H. Clifford and G. B. Preston. *The algebraic theory of semigroups. Vol. I.* Mathematical Surveys, No. 7. American Mathematical Society, Providence, R.I., 1961.
- [25] I. Colazzo and A. Van Antwerpen. The algebraic structure of left semi-trusses. *J. Pure Appl. Algebra*, 225(2):Paper No. 106467, 15, 2021.
- [26] P. Dehornoy. Set-theoretic solutions of the Yang-Baxter equation, RC-calculus, and Garside germs. *Adv. Math.*, 282:93–127, 2015.
- [27] V. G. Drinfel'd. On some unsolved problems in quantum group theory. In *Quantum groups (Leningrad, 1990)*, volume 1510 of *Lecture Notes in Math.*, pages 1–8. Springer, Berlin, 1992.
- [28] P. Etingof and S. Gelaki. A method of construction of finite-dimensional triangular semisimple Hopf algebras. *Math. Res. Lett.*, 5(4):551–561, 1998.
- [29] P. Etingof, T. Schedler, and A. Soloviev. Set-theoretical solutions to the quantum Yang-Baxter equation. *Duke Math. J.*, 100(2):169–209, 1999.
- [30] S. C. Featherstonhaugh, A. Caranti, and L. N. Childs. Abelian Hopf Galois structures on prime-power Galois field extensions. *Trans. Amer. Math. Soc.*, 364(7):3675–3684, 2012.
- [31] T. Gateva-Ivanova. Quadratic algebras, Yang-Baxter equation, and Artin-Schelter regularity. *Adv. Math.*, 230(4-6):2152–2175, 2012.
- [32] T. Gateva-Ivanova. A combinatorial approach to noninvolutive set-theoretic solutions of the Yang-Baxter equation. *Publ. Mat.*, 65:747–808, 2021.
- [33] T. Gateva-Ivanova and S. Majid. Matched pairs approach to set theoretic solutions of the Yang-Baxter equation. *J. Algebra*, 319(4):1462–1529, 2008.
- [34] T. Gateva-Ivanova and M. Van den Bergh. Semigroups of I -type. *J. Algebra*, 206(1):97–112, 1998.
- [35] A. Ghobadi. Drinfeld twists on skew braces. *Arxiv:2105.03286v1*, 2021.
- [36] I. Goffa and E. Jespers. Monoids of IG-type and maximal orders. *J. Algebra*, 308(1):44–62, 2007.
- [37] I. Goffa, E. Jespers, and J. Okniński. Primes of height one and a class of Noetherian finitely presented algebras. *Internat. J. Algebra Comput.*, 17(7):1465–1491, 2007.
- [38] L. Guarnieri and L. Vendramin. Skew braces and the Yang–Baxter equation. *Math. Comp.*, 86(307):2519–2534, 2017.
- [39] E. Jespers, L. Kubat, and A. Van Antwerpen. The structure monoid and algebra of a non-degenerate set-theoretic solution of the Yang-Baxter equation. *Trans. Amer. Math. Soc.*, 372(10):7191–7223, 2019.
- [40] E. Jespers, L. Kubat, and A. Van Antwerpen. Corrigendum and addendum to “The structure monoid and algebra of a non-degenerate set-theoretic solution of the Yang–Baxter equation”. *Trans. Amer. Math. Soc.*, 373(6):4517–4521, 2020.
- [41] E. Jespers and J. Okniński. Monoids and groups of I -type. *Algebr. Represent. Theory*, 8(5):709–729, 2005.
- [42] E. Jespers and J. Okniński. *Noetherian semigroup algebras*, volume 7 of *Algebra and Applications*. Springer, Dordrecht, 2007.
- [43] E. Jespers and A. Van Antwerpen. Left semi-braces and solutions of the Yang-Baxter equation. *Forum Math.*, 31(1):241–263, 2019.
- [44] E. Jespers and M. Van Campenhout. Finitely generated algebras defined by homogeneous quadratic monomial relations and their underlying monoids II. *J. Algebra*, 492:524–546, 2017.

- [45] S. Kamada. Knot invariants derived from quandles and racks. In *Invariants of knots and 3-manifolds (Kyoto, 2001)*, volume 4 of *Geom. Topol. Monogr.*, pages 103–117. Geom. Topol. Publ., Coventry, 2002.
- [46] C. Kassel. *Quantum groups*, volume 155 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [47] M. Kępczyk. A ring which is a sum of two *PI* subrings is always a *PI* ring. *Israel J. Math.*, 221(1):481–487, 2017.
- [48] A. Koch and P. J. Truman. Opposite skew left braces and applications. *J. Algebra*, 546:218–235, 2020.
- [49] V. Lebed. Applications of self-distributivity to Yang-Baxter operators and their cohomology. *J. Knot Theory Ramifications*, 27(11):1843012, 20, 2018.
- [50] V. Lebed and L. Vendramin. Homology of left non-degenerate set-theoretic solutions to the Yang-Baxter equation. *Adv. Math.*, 304:1219–1261, 2017.
- [51] V. Lebed and L. Vendramin. On structure groups of set-theoretic solutions to the Yang-Baxter equation. *Proc. Edinb. Math. Soc. (2)*, 62(3):683–717, 2019.
- [52] J.-H. Lu, M. Yan, and Y.-C. Zhu. On the set-theoretical Yang-Baxter equation. *Duke Math. J.*, 104(1):1–18, 2000.
- [53] L. H. Rowen. General polynomial identities. II. *J. Algebra*, 38(2):380–392, 1976.
- [54] W. Rump. A decomposition theorem for square-free unitary solutions of the quantum Yang-Baxter equation. *Adv. Math.*, 193(1):40–55, 2005.
- [55] W. Rump. Braces, radical rings, and the quantum Yang-Baxter equation. *J. Algebra*, 307(1):153–170, 2007.
- [56] W. Rump. Generalized radical rings, unknotted biquandles, and quantum groups. *Colloq. Math.*, 109(1):85–100, 2007.
- [57] W. Rump. The brace of a classical group. *Note Mat.*, 34(1):115–144, 2014.
- [58] W. Rump. A covering theory for non-involutive set-theoretic solutions to the Yang-Baxter equation. *J. Algebra*, 520:136–170, 2019.
- [59] A. Smoktunowicz. Algebraic approach to rump’s results on relations between braces and pre-lie algebras. *Arxiv:2007.09403*, 2020.
- [60] A. Smoktunowicz. A new formula for lazarid’s correspondence for finite braces and pre-lie algebras. *Arxiv:2011.07611*, 2020.
- [61] A. Smoktunowicz and L. Vendramin. On skew braces (with an appendix by N. Byott and L. Vendramin). *J. Comb. Algebra*, 2(1):47–86, 2018.
- [62] A. Soloviev. Non-unitary set-theoretical solutions to the quantum Yang-Baxter equation. *Math. Res. Lett.*, 7(5-6):577–596, 2000.
- [63] M. Takeuchi. Survey on matched pairs of groups—an elementary approach to the ESS-LYZ theory. In *Noncommutative geometry and quantum groups (Warsaw, 2001)*, volume 61 of *Banach Center Publ.*, pages 305–331. Polish Acad. Sci. Inst. Math., Warsaw, 2003.
- [64] C. N. Yang. Some exact results for the many-body problem in one dimension with repulsive delta-function interaction. *Phys. Rev. Lett.*, 19(23):1312–1315, 1967.
- [65] A. Yekutieli and J. J. Zhang. Homological transcendence degree. *Proc. London Math. Soc. (3)*, 93(1):105–137, 2006.

(I. Colazzo (ORCID: 0000-0002-2713-0409)) DEPARTMENT OF MATHEMATICS, COLLEGE OF ENGINEERING, MATHEMATICS AND PHYSICAL SCIENCES, UNIVERSITY OF EXETER, EXETER EX4 4QF, UK

Email address: I.Colazzo@exeter.ac.uk

(E. Jespers (ORCID: 0000-0002-2695-7949), A. Van Antwerpen (ORCID: 0000-0001-7619-6298), and C. Verwimp (ORCID: 0000-0003-3128-4854)) DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL

Email address: eric.jespers@vub.be

Email address: arne.van.antwerpen@vub.be

Email address: charlotte.verwimp@vub.be