

Dual Fields of Massive/Massless Gravitons in IR/UV Completions*

Ashkbiz Danehkar^{†,||}, Hassan Alshal^{‡,§,**}, and Thomas L. Curtright^{¶,††}

[†]*Department of Astronomy, University of Michigan, Ann Arbor, MI 48109, USA*

[‡]*Department of Chemistry and Physics, Lincoln University, PA 19352, USA*

[§]*Department of Physics, Cairo University, Giza 12613, Egypt*

[¶]*Department of Physics, University of Miami, Coral Gables, FL 33146, USA*

^{||}*danehkar@umich.edu*

^{**}*halshal@sci.cu.edu.eg*

^{††}*curtright@miami.edu*

Received 21 May 2021

Accepted 10 September 2021

In the holographic picture, the BEH mechanism in d -dimensional Yang-Mills theories is conjectured to provide a Higgs-like mechanism for gravity in $d + 1$ dimensions, resulting in massive (or massless) gravitons in IR (or UV) completions. Accordingly, one could imagine dual (magnetic-type) fields of massive gravitons in the IR (low-energy) limit that are coupled to the curl of their own energy-momentum, as well as to the rotation of matter fields on large scales. This hypothesis, which might solve cosmological issues currently ascribed to dark matter and dark energy, needs to be examined by the future LISA mission using observations of gravitational waves emitted from extragalactic sources.

Keywords: massive gravity; dual fields; gravitation; gravitational waves

PACS numbers: 04.50.-h, 95.30.Sf, 04.50.Kd

According to the holographic principle,¹ a theory of gravity could be related to a gauge field theory in one less dimension. This leads to the AdS _{$d+1$} /CFT _{d} correspondence,² also known as gauge/gravity duality. The holographic correspondence between d -dimensional Yang-Mills theories and gravitational fields in $d + 1$ dimensions thereby hints at a geometric counterpart of the BEH mechanism.³ In the context of the standard model of particle physics, the electroweak W and Z bosons acquire mass through the BEH mechanism,⁴ which is analogous to the behavior of photons inside superconductors.⁵ While photons are massless in empty space, they are apparently massive inside superconductors at very low temperatures, and hence described by the Maxwell-Proca equations. From the holographic principle, it follows that a Higgs-like mechanism analogue to that in Yang-Mills theories could

*This essay received an Honorable Mention in the 2021 Essay Competition of the Gravity Research Foundation.

be expected for gravitational theories in one higher dimension. This could give a mass to gravitons in the IR completion on a large scale, resulting in massive gravity (see Ref. 6 for a general review of massive gravity in various circumstances).

In the theory of general relativity, the graviton is taken to be massless, with tidal and frame-dragging fields encoded in the Weyl conformal tensor $C_{\mu\alpha\nu\beta}$.⁷ The decomposed parts of the Weyl tensor, $E_{\mu\nu} \equiv C_{\mu\alpha\nu\beta}u^\alpha u^\beta$ and $H_{\mu\nu} \equiv -\frac{1}{2}\epsilon_{\mu\gamma\alpha\beta}C^{\alpha\beta}{}_{\nu\lambda}u^\gamma u^\lambda$ (where $\epsilon_{\mu\gamma\alpha\beta}$ is the Levi-Civita symbol), are analogous to the electric and magnetic fields in Maxwell theory, thereby suggesting gravitational electric-magnetic duality (see Ref. 8 for a review of this idea). By analogy with the four-potential vector A_μ in Maxwell theory (spin 1), it is standard lore to anticipate a 2nd-rank symmetric tensor $h_{\mu\nu}$ in gravitational theories (spin 2). According to the electric-magnetic invariance, the electromagnetic potential vector A_μ is dual to a $(d-3)$ -form potential field $\tilde{A}_{\mu_1\dots\mu_{d-3}}$ in d dimensions, while it is dual to itself in $d=4$.

The extension of the electric-magnetic duality principle to d -dimensional gravitational theories implies that the symmetric tensor $h_{\mu\nu}$ of a massless graviton is dual to a mixed symmetry $(d-3, 1)$ tensor $\tilde{h}_{[\mu_1\dots\mu_{d-3}]\nu}$ of rank $d-2$ (also called Curtright fields),^{9,10} whereas the $h_{\mu\nu}$ of a massive graviton is dual to a mixed symmetry $(d-2, 1)$ tensor $\tilde{h}_{[\mu_1\dots\mu_{d-2}]\nu}$ of rank $d-1$.¹¹⁻¹⁴ In $d=4$, the massless field $h_{\mu\nu}$ is dual to itself, whereas the massive field $h_{\mu\nu}$ is dual to a mixed symmetry $(2, 1)$ tensor $\tilde{h}_{[\mu\nu]\lambda}$. The number of propagating helicity modes of $h_{\mu\nu}$ and its dual field is $d(d-3)/2$ for massless gravitons and $(d+1)(d-2)/2$ for massive gravitons. In $d=4$, a massless graviton propagates with 2 helicity modes, whereas a massive graviton propagates with 5 helicity modes.

We know that the (gravito-)magnetic part of the Weyl tensor of massless gravitons can be geometrically interpreted as frame-dragging vortexes around spinning massive objects.¹⁵ Similarly, by comparison with $\tilde{A}_{\mu_1\dots\mu_{d-3}}$, one could speculate about magnetic-like effects introduced by $\tilde{h}_{[\mu_1\dots\mu_{d-3}]\nu}$. This stems from the fact that the *geometric* Lanczos tensor $H_{[\mu\nu]\alpha}$ generating the Weyl tensor $C_{\mu\nu\alpha\beta}$ ¹⁶ has gauge symmetries similar to those of the Curtright fields, which can be used to predict frame-dragging effects for the Curtright fields in a way same as *gravitomagnetism* in general relativity.

The Lagrangian density for dual fields of massless gravitons in d dimensions can be expressed as $\mathcal{L}_{\text{massless}} = \mathcal{F}_\mu{}^\nu \mathcal{F}_\nu{}^\mu$, where $\mathcal{F}_\mu{}^\nu$ is the curl of dual fields defined as:^{9,17}

$$\mathcal{F}_\mu{}^\nu \equiv \frac{1}{(d-3)!} \epsilon^{\lambda_1\dots\lambda_{d-2}\nu} \partial_{\mu_1} \tilde{h}_{[\lambda_2\dots\lambda_{d-2}]\mu}. \quad (1)$$

For dual fields of massive gravity, the Lagrangian density is as follows:¹¹⁻¹⁴

$$\begin{aligned} \mathcal{L}_{\text{massive}} = & \mathcal{K}_\mu{}^\nu \mathcal{K}_\nu{}^\mu + \frac{(-1)^d}{(d-2)!} m^2 \left(\tilde{h}_{[\lambda_1\dots\lambda_{d-2}]\mu} \tilde{h}^{[\lambda_1\dots\lambda_{d-2}]\mu} \right. \\ & \left. - (d-2) \tilde{h}_{[\lambda_1\dots\lambda_{d-3}]\nu}{}^\nu \tilde{h}^{[\lambda_1\dots\lambda_{d-3}]\rho}{}_\rho \right), \quad (2) \end{aligned}$$

where m is the graviton mass, and $\mathcal{K}_\mu{}^\nu$ is the curl of dual fields of massive gravity:

$$\mathcal{K}_\mu{}^\nu \equiv \frac{1}{(d-2)!} \epsilon^{\lambda_1 \dots \lambda_{d-1} \nu} \partial_{\mu_1} \tilde{h}_{[\lambda_2 \dots \lambda_{d-1}] \mu}. \quad (3)$$

The Lagrangian density (2) is analogue to the Proca action for massive photons, $\mathcal{L}_{\text{Proca}} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} m^2 A_\mu A^\mu$.

The dual fields of massive gravitons can be coupled to the curl of any conserved, energy-momentum tensor:^{11,13}

$$(\square + m^2) \tilde{h}_{[\lambda_1 \dots \lambda_{d-2}] \nu} = \kappa P_{\lambda_1 \dots \lambda_{d-2} \nu}{}^{\alpha\beta\gamma} \partial_\alpha (\mathcal{T}_{\beta\gamma} + \mathfrak{t}_{\beta\gamma}), \quad (4)$$

$$P_{\lambda_1 \dots \lambda_{d-2} \nu}{}^{\alpha\beta\gamma} \equiv (d-3) \epsilon_{\lambda_1 \dots \lambda_{d-2}}{}^{\alpha\beta} \eta_\nu{}^\gamma + (d-1) \epsilon_{(\lambda_1 \dots \lambda_{d-2}}{}^{\alpha\beta} \eta_{\nu)}{}^\gamma, \quad (5)$$

where $P_{\lambda_1 \dots \lambda_{d-2} \nu}{}^{\alpha\beta\gamma}$ is a Young symmetrizer, $\eta_{ab} = \text{diag}(-1, +1, +1, +1)$ the metric tensor, $\square \equiv \partial_\mu \partial^\mu$ the d'Alembert operator, κ a gravitational constant in d -dimensional spacetime, $\mathcal{T}_{\alpha\beta}$ the energy-momentum tensor of arbitrary matter fields, and $\mathfrak{t}_{\alpha\beta}$ the energy-momentum tensor of dual fields of massive gravitons (see Ref. 13). The field equation (4) is analogous with $(\square + m^2) A_\mu = J_\mu$ in Maxwell-Proca theory.

From Eq. (4), it follows the on-shell equation (see Eq. 35 in Ref. 12) for the dual field strength $\mathcal{K}_\mu{}^\nu$:^{12,13}

$$(\square + m^2) \mathcal{K}_\mu{}^\nu = \kappa(1-d) \square (\mathcal{T}_\mu{}^\nu + \mathfrak{t}_\mu{}^\nu) + \kappa (\delta_\mu{}^\nu \square - \partial_\mu \partial^\nu) (\mathcal{T} + \mathfrak{t}), \quad (6)$$

where $\mathcal{T} \equiv \mathcal{T}_\lambda{}^\lambda$ and $\mathfrak{t} \equiv \mathfrak{t}_\lambda{}^\lambda$, and $\delta_\mu{}^\nu \equiv \eta_{\mu\lambda} \eta^{\lambda\nu}$ is the Kronecker delta. The field equation (6) is similar to the conventional one in massive gravity, $(\square + m^2) h_{\mu\nu} = \kappa \mathcal{T}_{\mu\nu}$, apart from its constant with an opposite sign ($d > 3$) and the manifestly conserved trace term. As shown in Ref. 12, Eq. (6) leads to the Ogievetsky-Polubarinov theory,¹⁸ implying that the Curtright field is the exact dual of the Ogievetsky-Polubarinov model for massive gravitons.

According to Eqs. (4) and (6), the curl of massive gravitons in d -dimensional spacetime is associated with dual (magnetic-type) fields described by a $(d-1)$ -rank tensor of Young symmetry type $(D-2, 1)$, which are coupled to the curl of their own energy-momentum, as well as the rotation of matter fields on large scales. In this picture, dual fields of massive gravity that strongly emerge in the IR completion could have major implications for some phenomena on large scales. Moreover, as graviton is massless in the UV limit, the theory of general relativity still remains valid within the solar system and high energy domains.

It is argued that duality breaks asymptotic flatness of the Schwarzschild solution, and consequently fills the empty space surrounding isolated celestial objects, because of the extra constant that appears in the g_{tt} , which is responsible for creating the Barriola-Vilenkin monopole solution.¹⁹ In massive dual gravity, we have a constant, with the negative sign (see also Eq. 6), that plays the similar role in creating the dual graviton. The constant appears after expanding the exponent term in the static spherically symmetric solution of the h_{00} in the interaction theory of massive dual gravity (see Eqs. 39–40 in Ref. 12). Thus, if gravitoelectromagnetism prescribed

by general relativity was a good candidate as a Machian theory, then the massive gravity with its dual field could be too.

Gravitons that acquire a mass in the IR completion could potentially offer some solutions to the current issues in cosmology, namely flat rotation curves of *disk* galaxies and accelerating expansion of the universe since about 7 billion years ago. Gravitational forces mediated by massive gravitons are weakened at distances larger than the graviton's Compton wavelength $\lambda_g = \hbar/(m_g c^2)$ according to $\sim \frac{1}{r} \exp(-r/\lambda_g)$, by contrast, dual (magnetic-type) fields produced by the rotation of matter fields and massive gravitons turn to be stronger than gravitomagnetic fields associated with massless gravitons. Massive gravity yields dual fields associated with the curl of arbitrary conserved energy-momentum that gravitationally confine stars rotating with galactic disks of spiral galaxies. Moreover, dual fields of massive gravity may lead to the cosmic acceleration at late times if graviton is massless in the UV, but massive in the IR. This may naturally explain why the acceleration of the cosmic expansion occurs at late times on large scales when intergalactic space predominantly got much cooler. In particular, dual fields of massive gravity on a large scale may contribute to some preferential orientations in spiral galaxies in filaments, similar to those reported in the literature.²⁰

Gravitational waves (GW) in general relativity ($d = 4$) propagate at the speed of light with two tensor polarization modes: + and \times . In contrast, the speed of gravitational waves in massive gravity could be frequency-dependent, slightly below the speed of light. Apart from the frequency-dependent speed, massive gravity in $d = 4$ also generates five polarization modes: two tensors, two vectors, and one scalar (see the review by Ref. 21). If graviton remains massless in the UV regime, gravitational waves detected from nearby sources within our Galaxy should perfectly agree with general relativity. However, graviton may enter its massive mode as it propagates on intergalactic scales in the IR limit. In this IR/UV scenario, that makes massive/massless gravitons, GW emitted from a binary merger in a nearby galaxy could potentially carry three additional polarization modes, with a slight time delay between the GW and its electromagnetic counterpart. The presence of additional GW polarization modes can be investigated using 3-arm space-borne GW detectors such as the Laser Interferometer Space Antenna (LISA; planned for 2034), whereas 2-arm GW detectors similar to the Laser Interferometer Gravitational-Wave Observatory (LIGO) are inadequate for this analysis. Future space-borne GW detectors will decide whether gravitons propagate in the massive mode in the IR regime.

Vector bosons in Yang-Mills theories can acquire a rest mass through the BEH mechanism by ingesting the Higgs scalar field (spin 0). However, the Higgs field is supposed to provide the BEH mechanism for vector bosons (spin 1). As argued by Ref. 22, the holographic correspondence maps a 4-dimensional Yang-Mills theory onto a 5-dimensional theory of gravity that could also be projected onto a massive graviton in 4 dimensions. In this scenario, four massless scalar fields can assemble a Higgs-like mechanism for gravity in $d = 4$, by which a massive graviton and a

massive scalar boson survive,^{22,23} while the other three scalar fields are absorbed into the massive graviton. This geometric counterpart of the Higgs boson should be detectable in our future high-energy particle physics experiments if the nature also supports the Higgs-like mechanism for gravity.

References

1. G. 't Hooft, *Conf. Proc. C* **930308**, 284 (1993), arXiv:gr-qc/9310026; L. Susskind, *J. Math. Phys.* **36**, 6377 (1995).
2. J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998); J. Maldacena, *Int. J. Theor. Phys.* **38**, 1113 (1999); E. Witten, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
3. M. Porrati, *JHEP* **2002**, 058 (2002); M. Porrati, *Mod. Phys. Lett. A* **18**, 1793 (2003); O. Aharony, A. B. Clark, and A. Karch, *Phys. Rev. D* **74**, 086006 (2006).
4. F. Englert and R. Brout, *Phys. Rev. Lett.* **13**, 321 (1964); P. W. Higgs, *Phys. Rev. Lett.* **13**, 508 (1964); G. S. Guralnik, C. R. Hagen, and T. W. Kibble, *Phys. Rev. Lett.* **13**, 585 (1964).
5. A. S. Goldhaber and M. M. Nieto, *Rev. Mod. Phys.* **82**, 939 (2010); F. Wilczek, *Central Eur. J. Phys.* **10**, 1021 (2012).
6. K. Hinterbichler, *Rev. Mod. Phys.* **84**, 671 (2012); C. de Rham, *Living Rev. Rel.* **17**, 7 (2014).
7. R. Maartens and B. A. Bassett, *Class. Quant. Grav.* **15**, 705 (1998); G. F. R. Ellis, *Gen. Rel. Grav.* **41**, 581 (2009); A. Danehkar, *Mod. Phys. Lett. A* **24**, 3113 (2009).
8. A. Danehkar, *Front. Phys.* **6**, 146 (2019).
9. T. Curtright, *Phys. Lett. B* **165**, 304 (1985).
10. C. M. Hull, *Nucl. Phys. B* **583**, 237 (2000); C. M. Hull, *JHEP* **2001**, 027 (2001); P. West, *Class. Quant. Grav.* **18**, 4443 (2001).
11. T. L. Curtright and P. G. O. Freund, *Nucl. Phys. B* **172**, 413 (1980); T. L. Curtright, *Nucl. Phys. B* **948**, 114784 (2019).
12. T. L. Curtright and H. Alshal, *Nucl. Phys. B* **948**, 114777 (2019).
13. H. Alshal and T. L. Curtright, *JHEP* **2019**, 63 (2019); H. Alshal, *Aspects of Massive Dual Gravity*, Ph.D. thesis, University of Miami, Florida (2020).
14. B. González, A. Khoudeir, R. Montemayor, et al., *JHEP* **2008**, 058 (2008).
15. R. Owen, J. Brink, Y. Chen, et al., *Phys. Rev. Lett.* **106**, 151101 (2011); D. A. Nichols, R. Owen, F. Zhang, et al., *Phys. Rev. D* **84**, 124014 (2011); D. A. Nichols, A. Zimmerman, Y. Chen, et al., *Phys. Rev. D* **86**, 104028 (2012); A. Danehkar, *Int. J. Mod. Phys. D* **29**, 2043001 (2020).
16. C. Lanczos, *Rev. Mod. Phys.* **34**, 379 (1962); H. Takeno, *Tensor N.S.* **15**, 103 (1964).
17. N. Boulanger and O. Hohm, *Phys. Rev. D* **78**, 064027 (2008).
18. V. I. Ogievetsky and I. V. Polubarinov, *Annals Phys.* **35**, 167 (1965).

6 REFERENCES

19. N. Dadhich, *Gen. Rel. Grav.* **32**, 1009 (2000); N. Dadhich, *arXiv e-prints* , gr-qc/9902066 (1999).
20. I. Trujillo, C. Carretero, and S. G. Patiri, *Astrophys. J. Lett.* **640**, L111 (2006); J. Lee and P. Erdogdu, *Astrophys. J.* **671**, 1248 (2007); E. Tempel and N. I. Libeskind, *Astrophys. J. Lett.* **775**, L42 (2013); Y. Zhang, X. Yang, H. Wang, et al., *Astrophys. J.* **779**, 160 (2013).
21. J. M. Ezquiaga and M. Zumalacárregui, *Front. Astron. Space Sci.* **5**, 44 (2018).
22. G. 't Hooft, *Preprint* , arXiv:0708.3184 (2007), report No. ITP-UU-07-44, SPIN-07-32.
23. Z. Kakushadze, *Int. J. Mod. Phys. A* **23**, 1581 (2008); A. H. Chamseddine and V. Mukhanov, *JHEP* **2010**, 11 (2010).