

Improved DHOST Genesis

Mian Zhu^{1,2,3,*} and Yunlong Zheng^{3,4,5,6,†}

¹*Department of Physics, The Hong Kong University of Science and Technology,
Clear Water Bay, Hong Kong S.A.R., P.R.China*

²*HKUST Jockey Club Institute for Advanced Study,
The Hong Kong University of Science and Technology,
Clear Water Bay, Hong Kong S.A.R., P.R.China*

³*Department of Astronomy, School of Physical Sciences,
University of Science and Technology of China, Hefei, Anhui 230026, China*

⁴*CAS Key Laboratory for Researches in Galaxies and Cosmology,
University of Science and Technology of China, Hefei, Anhui 230026, China*

⁵*School of Astronomy and Space Science, University of Science and Technology of China, Hefei, Anhui 230026, China*

⁶*ICRANet, Piazza della Repubblica 10, I-65122 Pescara, Italy*

We improve the DHOST Genesis proposed in [1], such that the near scale invariant scalar power spectrum can be generated from the model itself, without invoking extra mechanism like a string gas. Besides, the superluminality problem of scalar perturbation plagued in [1] can be rescued by choosing proper DHOST action.

I. INTRODUCTION

Currently, Inflation [2–5] is the standard paradigm of the very early universe. Alternative scenarios, like bounce cosmology [6–16], emergent universe scenario [17, 18] and string gas cosmology [19–26], are also of people’s interests. It is believed that certain alternative scenarios can not only explain the formation of the Large Scale Structure of our universe as good as inflation, but also evade the initial spacetime singularity which plagues in inflationary cosmology [27, 28] (see [29, 30] for reviews on alternative cosmologies).

In this paper we focus on the emergent universe scenario, in which the universe is emergent from a quasi-Minkowskian spacetime. Such a scenario requires a violation of Null Energy Condition (NEC). One possible way to realize the NEC violation is introducing Horndeski/Generalized Galileon theory [31–34], which results in the Galileon Genesis model [35, 36] (see also earlier in [37]). The physics of Galileon Genesis is studied in [38–42] and extended in [43–52].

The Galileon Genesis faces several challenges. Firstly, there are no-go theorems stating that no classically stable Genesis solution exists within the framework of Horndeski/Galileon theory [53–56]¹. One possible ways out is to consider models which are strongly coupled in the asymptotic past/future [48, 59], at the cost of potentially triggering the strong-coupling problem [60, 61]. Besides, it is not very clear how such a Genesis scenario can be connected to the standard cosmology².

To evade the possible puzzles, we recently propose an emergent universe model from the DHOST cosmology [1]. We modified the original action of Galileon Genesis such that the universe can gracefully exit from the quasi-Minkowskian state to a radiation dominated phase. Moreover, we introduce a DHOST coupling to cure the gradient instability. However, in our model the scale invariant scalar power spectrum cannot be generated within the model. To be consistent with astrophysical observations, we introduce a quasi-static Hagedorn phase from string gas cosmology, and hence the model itself is incomplete.

We complete the previous model by a slight deformation of the action. In Galileon Genesis, the behavior of Hubble parameter $H \propto (-t)^{-2\alpha-1}$ determines the scalar spectrum index as $n_s = 5 - 2\alpha$ [37, 47, 48, 63, 64]. We improve the action of our previous model [1] accordingly, and show that the improved Genesis model can produce a scale invariant power spectrum. Hence our improved model can serve as a possible candidate for the early universe, without invoking additional mechanisms.

The paper is organized as follows. In section II, we briefly introduce the improved model and the genesis solution. The background dynamics is analyzed in section III, where we verify that our model can gracefully exit to a radiation dominated universe. We work out the evolution of the scalar perturbation in section IV, and show that the scale invariance can be obtained. The tensor perturbation is discussed in section V. Finally, we conclude our work in section VI.

Throughout the paper, we take the sign of the metric as $(+, -, -, -)$. The canonical kinetic term of the scalar field π is defined as $X \equiv \frac{1}{2} \nabla_\mu \pi \nabla^\mu \pi$. The d’Alembert operator is $\square \equiv \nabla_\mu \nabla^\mu$. The reduced Planck mass $M_p^2 \equiv \frac{1}{8\pi G}$ is set to be 1.

* mzhuan@connect.ust.hk

† Corresponding author: zhyunl@ustc.edu.cn

¹ It is proved in the framework of effective field theory [49, 57, 58] that, a fully stable NEC violation can be realized only in theories beyond Horndeski.

² In [62], a model of inflation preceded by (generalized) Galilean Genesis is proposed.

II. IMPROVED DHOST GENESIS

A. Action

The action of the model can be written as

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2} + \mathcal{L}_{H2} + \mathcal{L}_{H3} + \mathcal{L}_D \right), \quad (1)$$

where \mathcal{L}_{H2} and \mathcal{L}_{H3} come from the first two classes of Horndeski/Galileon theory [32, 34], and \mathcal{L}_D represents the type ⁽²⁾ $N - II$ DHOST Lagrangian [65–67]³:

$$\mathcal{L}_{H2} = K(\pi, X) = -g_1(\pi)X + g_2(\pi)X^2, \quad (2)$$

$$\mathcal{L}_{H3} = G(X)\square X, \quad G(X) = \gamma X^2, \quad (3)$$

$$\begin{aligned} \mathcal{L}_D = & \frac{R}{2}h - \frac{h}{4X} [\pi_{\mu\nu}\pi^{\mu\nu} - (\square\pi)^2] \\ & + \frac{h - 2Xh_X}{4X^2} [\pi_{\mu}\pi^{\mu\rho}\pi_{\rho\nu}\pi^{\nu} - (\square\pi)\pi^{\mu}\pi_{\mu\nu}\pi^{\nu}]. \end{aligned} \quad (4)$$

In equation (1), $R/2$ is the standard Einstein-Hilbert action. $g_1(\pi)$, $g_2(\pi)$ are function of the scalar field π , while $h \equiv h(X)$ is a function of the canonical kinetic term X only. Their detailed expressions are taken to be

$$g_1(\pi) = \frac{3}{2}f^2e^{4\pi} \frac{1 + e^{2\pi}}{1 + e^{m\pi}}, \quad g_2(\pi) = e^{2\pi} \frac{1 + e^{2\pi}}{1 + e^{4\pi}}, \quad (5)$$

$$h(X) = d_1X + d_2X^2, \quad (6)$$

in which $m > 6$ is a fixed parameter which determines the behavior when the universe exiting the quasi-Minkowskian state. As we will show in next sections, the value of m doesn't change the scale dependence, so we will simply take $m = 7$ through out this paper.

From equation (5), we have

$$\lim_{\pi \rightarrow -\infty} g_1(\pi) = \frac{3}{2}f^2e^{4\pi}, \quad \lim_{\pi \rightarrow -\infty} g_2(\pi) = e^{2\pi}; \quad (7)$$

$$\lim_{\pi \rightarrow \infty} g_1(\pi) = 0, \quad \lim_{\pi \rightarrow \infty} g_2(\pi) = 1. \quad (8)$$

Applying the FLRW background to the action (1), we recover the Friedmann equations

$$3H^2 = \rho_{\pi}, \quad -2\dot{H} = \rho_{\pi} + p_{\pi}, \quad (9)$$

where H is the Hubble parameter, ρ_{π} and p_{π} are the density and pressure of the matter field π , expressed by

$$\rho_{\pi} = -\frac{1}{2}g_1\dot{\pi}^2 + \frac{3}{4}g_2\dot{\pi}^4 + 3H\gamma\dot{\pi}^5, \quad (10)$$

$$p_{\pi} = -\frac{1}{2}g_1\dot{\pi}^2 + \frac{1}{4}g_2\dot{\pi}^4 - \gamma\dot{\pi}^4\ddot{\pi}. \quad (11)$$

We see from equation (10) and (11) that the DHOST term (4) doesn't enter into the background dynamics. This fact is already found in [1, 74] as long as the DHOST coupling is non-trivial only during the NEC violation phase. Moreover, observational signals from scalar perturbation, such as the scalar spectra index and tensor-to-scalar ratio, will not be affected by DHOST action in bounce cosmology [75], and in section IV, we will similarly find that the scalar spectra index is independent of the DHOST action.

B. The Genesis Solution

We assume the initial condition of scalar field π at far past to be $\pi \ll -1$. From equation (7), we see the first Friedmann's equation becomes

$$3H^2 = -\frac{3}{4}f^2e^{4\pi}\dot{\pi}^2 + \frac{3}{4}e^{2\pi}\dot{\pi}^4 + 3H\gamma\dot{\pi}^5, \quad (12)$$

and there is a quasi-Minkowskian solution with $H \approx 0$ as long as

$$\dot{\pi}^2 e^{-2\pi} = f^2 \rightarrow \pi = \ln \left[\frac{1}{f(-t)} \right]. \quad (13)$$

Substituting equation (13) into equation (11), the pressure becomes

$$p_{\pi} = -\frac{1}{(-t)^6} \left(\frac{1}{2f^2} + \gamma \right), \quad (14)$$

which is of order $\mathcal{O}(t^{-6})$ and larger than that of the density. So the universe will deviate from the quasi-Minkowskian configuration by the negative and non-vanishing pressure.

C. Model Parameters

To illustrate the dynamics of the universe and numerically verify the approximations used in next sections, we will use the following parameters through out this paper:

$$f = 0.8, \quad \gamma = 1, \quad d_1 = 0, \quad d_2 = 6. \quad (15)$$

As we shall see in section IV B, the sound speed of scalar perturbation in pre-emergent period is $c_{s0}^2 = (4f^2\gamma - 1)/3$. To avoid the gradient instability and superluminality problem in this period, we have $0 < c_{s0}^2 < 1$ and the model parameters are constrained by

$$\frac{1}{4} < \gamma f^2 < 1. \quad (16)$$

Finally, we will show in section IV A that the DHOST coupling chosen in equation (15) can evade the gradient instability and superluminality problem of scalar perturbation in the whole cosmological evolution.

³ More explicitly, the action (4) belongs to the GLPV theory [68–70], which propagates only one scalar degree of freedom [71–73] and hence included by the DHOST theory.




FIG. 1. Dynamics of background geometry a , H and scalar field π , $\dot{\pi}$ as a function of cosmic time t with the special parameter set (15).

III. BACKGROUND EVOLUTION

In this section we briefly discuss the background dynamics of our model. As shown in section II B, at far past $t \rightarrow -\infty$, the model permits a quasi-Minkowskian solution described by equation (13). Substituting the solution (13) into the second Friedmann equation (9) and use $p_\pi \gg \rho_\pi$, we have

$$-2\dot{H} \simeq p_\pi = -\frac{1}{(-t)^6} \left(\frac{1}{2f^2} + \gamma \right), \quad (17)$$

so that

$$H = \frac{h_0}{(-t)^5}, \quad h_0 \equiv \frac{1}{10} \left(\frac{1}{2f^2} + \gamma \right). \quad (18)$$

When the universe evolves towards the emergent event $t = 0$, the approximated equations (13), (18) become invalid, and the universe will deviate from the quasi-Minkowskian state. We define a typical time scale t_E , so that the approximations (13) and (18) is valid when $t < t_E < 0$. For the specific set of parameters (15), t_E is of order $\mathcal{O}(1)$. After the emergent event $t = 0$, the scalar field π become positive and the function $g_1(\pi)$, $g_2(\pi)$ quickly turns to the form (8). The pressure and density then becomes

$$\rho_\pi = \frac{3}{4}\dot{\pi}^4 + 3H\gamma\dot{\pi}^5, \quad p_\pi = \frac{1}{4}\dot{\pi}^4 - \gamma\dot{\pi}^4\ddot{\pi}, \quad (19)$$

and there is an approximated solution

$$a \propto t^{\frac{1}{2}}, \quad \dot{\pi} \propto t^{-\frac{1}{2}}, \quad (20)$$

which represents a radiation dominated universe.

As shown above, the universe starts from a quasi-Minkowskian state, then quit the status around the emergent event $t = 0$, and finally transits to a radiation dominated universe, thus connected to the standard Cosmology. Our model then provides a graceful exit mechanism




FIG. 2. Dynamics of z_s^2 and c_s^2 with respect to cosmic time t . The model parameters are set by (15).

to the regular Galileon Genesis scenario. We illustrate the cosmological evolution numerically in figure 1.

IV. SCALAR PERTURBATION

A. Dynamic equation and the stability issue

In this section, we study the scalar perturbation of the current model. Applying the previous result developed in [1, 74], the scalar perturbation at linear level is expressed by

$$S_{2,s} = \int d\tau d^3x \frac{z_s^2}{2} [\zeta'^2 - c_s^2 (\partial_i \zeta)^2], \quad (21)$$

where ζ is the curvature perturbation and

$$\frac{z_s^2}{2a^2} = 3 + 2 \frac{\dot{\pi}^2(-g_1 + 3g_2\dot{\pi}^2) + 18H\gamma\dot{\pi}^5 - 6H^2}{(\gamma\dot{\pi}^5 - 2H)^2}, \quad (22)$$

$$\left(-\frac{z_s^2}{2a^2} \right) c_s^2 = 1 + h + \frac{2}{a} \frac{d}{dt} \frac{a(h_X\dot{\pi}^2 - h - 1)}{2H - \gamma\dot{\pi}^5}. \quad (23)$$

The dynamical equation of the scalar perturbation is the standard Mukhanov-Sasaki(MS) equation [76–78]:

$$v_k'' + \left(k^2 c_s^2 - \frac{z_s''}{z_s} \right) v_k = 0, \quad (24)$$

where $v_k \equiv z_s \zeta$ is the MS variable, and a prime denotes differentiation with respect to the conformal time $d\eta = dt/a$.

To obtain a healthy model, we shall avoid the gradient instability and the existence of the ghost. This requires

$$z_s^2 > 0, \quad c_s^2 > 0. \quad (25)$$

We numerically illustrate the dynamics of z_s^2 and c_s^2 in figure 2 with parameters being (15). The gradient and ghost instabilities are absent. Moreover, the scalar sound speed does not exceed the speed of light, i.e. $c_s^2 < 1$, and hence the superluminality problem plagued in [1] is resolved.

B. General expression for the scalar power spectrum

In the pre-emergent period $t < 0$, the universe is almost static and $d\eta = dt/a \simeq dt$. By properly setting the integration constant, we can interchangeably use η and t in this region. Moreover, the DHOST coupling $h \approx 0$, along with the asymptotic behavior (7) and (18), the parameters reduce to

$$z_s^2 = \frac{600f^2}{(1 - 8f^2\gamma)^2} (-t)^4, \quad c_s^2 = \frac{4f^2\gamma - 1}{3} \equiv c_{s0}^2. \quad (26)$$

In equation (26), only the highest order of $(-t)$ are kept. The MS equation (24) becomes

$$v_k'' + \left(k^2 c_{s0}^2 - \frac{2}{\eta^2} \right) v_k = 0, \quad (27)$$

whose general solution is

$$v_k(\eta) \propto \sqrt{k\eta} \left[c_1(k) J_{\frac{3}{2}}(kc_{s0}\eta) + c_2(k) Y_{\frac{3}{2}}(kc_{s0}\eta) \right], \quad (28)$$

where $J_a(z)$ and $Y_a(z)$ are the Bessel functions.

Before proceeding, we mention that in generalized Galilean Genesis, the curvature perturbation ζ grows even on superhorizon scales when the second Horndeski term $\mathcal{L}_{H3} \propto X^\alpha \square \pi$ satisfies $\alpha > 1/2$ [35, 48]. Our model corresponds to the $\alpha = 2$ case, so we don't need to determine the time for each wavemode to exit the horizon during the pre-emergent period. However, after the emergent event $t = 0$, the universe becomes radiation dominated and the scalar perturbation contains a constant mode and a decreasing mode. Hence the dominated mode which is superhorizon at $t > 0$ will be the constant one, and the scalar perturbation will then remains invariant, just as the case in standard cosmology.

Now, as we argued in [1], the wavemodes of observational interests will all become superhorizon in the pre-emergent period $t < 0$. So we should evaluate the dynamics of scalar perturbation from past infinity to the critical time scale $\eta_E = t_E$, where the emergent solution breaks down. After $t = t_E$, the universe quickly turns to a radiation dominated period, and the scalar perturbation becomes conserved.

Hence, the scalar power spectrum can be evaluated by

$$P_\zeta(k) = \frac{k^3}{2\pi^2} \frac{|v_k(\eta_E)|^2}{z_s(\eta_E)^2}. \quad (29)$$

C. Vacuum initial condition

In this section we impose the vacuum initial condition and work out the scalar spectra index. Firstly, the argument of the Bessel function $kc_{s0}\eta_E \ll 1$, so we can expand the Bessel function with small argument approximation:

$$J_a(z) \sim \frac{1}{\Gamma(a+1)} \left(\frac{z}{2}\right)^a, \quad Y_a(z) \sim -\frac{\Gamma(a)}{\pi} \left(\frac{z}{2}\right)^a, \quad (30)$$




FIG. 3. Dynamics of c_T^2 with respect to cosmic time t . The model parameters are set by (15).

and the Bessel Y function will dominate at $\eta = \eta_E$. The MS variable is approximately

$$v_k(\eta_H) \propto c_2(k) \sqrt{k\eta_H} (k\eta_H)^{-\frac{3}{2}} = c_2(k) (k\eta_H)^{-1}. \quad (31)$$

Now we determine the $c_2(k)$ term from the vacuum initial condition. In the far past, the linearized action (21) is of a harmonic oscillator form with a nearly constant sound speed c_{s0} . So we can impose the quantum vacuum initial condition

$$v_k(\eta) \rightarrow \frac{e^{-ic_{s0}k(\eta-\eta_0)}}{\sqrt{2k}}, \quad (32)$$

where η_0 is an integration constant representing the initial phase. Comparing equation (32) and (28), and with the help of large argument expansion of Bessel function, we find

$$c_2(k) \propto k^{-1/2}. \quad (33)$$

Substituting equation (33) and (31) into the expression of scalar spectrum (29), we find

$$P_\zeta \propto k^3 \times |k^{-\frac{3}{2}}|^2 = k^0, \quad (34)$$

and the scale invariant scalar power spectrum is acquired.

V. TENSOR PERTURBATION

The tensor perturbation at the linearized level is [1]

$$S_{2,T} = \int d\eta d^3x \frac{a^2}{8} [\gamma_{ij}^{\prime 2} - (1+h)\gamma_{ij}^2], \quad (35)$$

in which γ_{ij} represents the tensor perturbation.

Since the coefficient of the kinetic term $\gamma_{ij}^{\prime 2}$ is always positive, there is no ghost model of tensor perturbation. We numerically plot the sound speed of tensor perturbation $c_T^2 = 1 + h$ in figure 3, and show that there is no gradient instability as well.

The tensor perturbation contains two modes. In our case the two polarization modes propagate at the same

speed [79], so we can collectively denote them as γ , and the MS equation becomes

$$\nu_k'' + \left(c_T^2 k^2 - \frac{a''}{a} \right) \nu_k = 0, \quad (36)$$

where $\nu_k \equiv 1/2a\gamma_k$ is the MS variable for tensor perturbation.

In the radiation dominated period, $a \propto t^{1/2}$, so the conformal time $\eta = \int dt/a \propto t^{1/2}$. We then have $a \propto \eta$ and a''/a vanishes. Hence the tensor perturbation propagates as a plane wave, and the tensor spectrum index is invariant in this period. Similarly, in the pre-emergent period the background is quasi-Minkowskian, and the same conclusion applied. One may also see the fact by noticing that, $H \propto (-t)^{-5}$ when $t < t_E$ and hence $a''/a \propto (-t)^{-6}$, which is negligible compared to $c_T^2 k^2$ in the MS equation (36). Finally, the duration of the transition period is small (of order $\mathcal{O}(10^0)$ in our set of parameters). Since there is no divergent part in MS equation (36) during this period, the contribution from this period to tensor perturbation will be suppressed by its short duration, so the tensor spectrum index is also invariant. We then conclude that, the tensor spectrum index we observed is just that in the beginning of the universe. Substituting the vacuum initial condition $\nu_k \propto k^{-1/2}$ ⁴ into the expression for tensor spectrum $P_t \propto k^3 |\nu_k|^2$, we see that $n_t = \frac{d \ln P_t}{d \ln k} = 2$, which is consistent with previous studies [35, 47].

VI. CONCLUSION AND OUTLOOK

In this paper we improve the DHOST Genesis model [1] to get the scale invariance scalar power spectrum. Compared to the original Galileon Genesis model [35], the improved model consists three new ingredient. The deformed Horndeski term \mathcal{L}_{H2} provides the realization for a graceful exit from emergent universe to standard cosmology; the \mathcal{L}_{H3} term determines the scalar spectrum in-

dex, resulting in the scale invariance; The DHOST term \mathcal{L}_D modifies the sound speed of scalar perturbation c_s^2 near the emergent event $t = 0$ without altering the background evolution, hence solves the gradient instability. The emergent universe scenario presented here is then healthy and can meet the basic observational constraint without other mechanism invoked.

There are still plenty of rooms for us to improve the current model. Firstly, most of the emergent universe models (including the one here) predict a blue tensor spectrum with $n_t = 2$ (counterexamples can be found in, say, [48, 80]), so updating the current model to allow more possibility for n_t would be worth investigating. Secondly, it is interesting to develop genesis models with alternative mechanism (like the curvaton mechanism [35, 39]) to realize the scale invariance, and see whether they can be distinguished by more observational data. Thirdly, it is worthy to develop the model to make the sound speed c_T^2 are less than $c^2 = 1$, hence evading the superluminality issue. Finally, a detailed phase space analyze can help to decide whether there is fine-tuning problem in our model.

ACKNOWLEDGMENTS

We are grateful to Yifu Cai, Robert Brandenberger, Yong Cai, Yi Wang, Damien Easson, Zhibang Yao, Xian Gao, Emmanuel Saridakis, Misao Sasaki and Dong-Gang Wang for stimulating discussions. This work is supported in part by the NSFC (Nos. 11847239, 11421303), by the Fundamental Research Funds for Central Universities, by the China Scholarship Council (CSC No.202006345019), and by GRF Grant 16304418 from the Research Grants Council of Hong Kong. YZ would like to thank the ICRANet for their hospitality during his visit. All numeric are operated on the computer clusters *LINDA* & *JUDY* in the particle cosmology group at USTC.

-
- [1] Amara Ilyas, Mian Zhu, Yunlong Zheng, and Yi-Fu Cai. Emergent Universe and Genesis from the DHOST Cosmology. *JHEP*, 01:141, 2021.
- [2] Alan H. Guth. The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems. *Phys. Rev.*, D23:347–356, 1981. [Adv. Ser. Astrophys. Cosmol.3,139(1987)].
- [3] Andrei D. Linde. A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems. *Phys. Lett.*, 108B:389–393, 1982. [Adv. Ser. Astrophys. Cosmol.3,149(1987)].

- [4] Andreas Albrecht and Paul J. Steinhardt. Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking. *Phys. Rev. Lett.*, 48:1220–1223, 1982. [Adv. Ser. Astrophys. Cosmol.3,158(1987)].
- [5] S. W. Hawking and I. G. Moss. Supercooled Phase Transitions in the Very Early Universe. *Phys. Lett.*, 110B:35–38, 1982. [Adv. Ser. Astrophys. Cosmol.3,154(1987)].
- [6] Robert H. Brandenberger. A Nonsingular universe. In *International School of Astrophysics, 'D. Chalonge': 2nd Course: Current Topics in Astrofundamental Physics*, 9 1992.
- [7] Robert H. Brandenberger, Viatcheslav F. Mukhanov, and A. Sornborger. A Cosmological theory without singularities. *Phys. Rev. D*, 48:1629–1642, 1993.
- [8] Yi-Fu Cai, Tao-tao Qiu, Robert Brandenberger, and Xinmin Zhang. A Nonsingular Cosmology with a Scale-

⁴ The expression is analog to that for scalar perturbation.

- Invariant Spectrum of Cosmological Perturbations from Lee-Wick Theory. *Phys. Rev. D*, 80:023511, 2009.
- [9] Yi-Fu Cai, Damien A. Easson, and Robert Brandenberger. Towards a Nonsingular Bouncing Cosmology. *JCAP*, 1208:020, 2012.
- [10] Daisuke Yoshida, Jerome Quintin, Masahide Yamaguchi, and Robert H. Brandenberger. Cosmological perturbations and stability of nonsingular cosmologies with limiting curvature. *Phys. Rev. D*, 96(4):043502, 2017.
- [11] M. Novello and S. E. Perez Bergliaffa. Bouncing Cosmologies. *Phys. Rept.*, 463:127–213, 2008.
- [12] Jean-Luc Lehners. Ekpyrotic and Cyclic Cosmology. *Phys. Rept.*, 465:223–263, 2008.
- [13] Yi-Fu Cai. Exploring Bouncing Cosmologies with Cosmological Surveys. *Sci. China Phys. Mech. Astron.*, 57:1414–1430, 2014.
- [14] D. Battefeld and Patrick Peter. A Critical Review of Classical Bouncing Cosmologies. *Phys. Rept.*, 571:1–66, 2015.
- [15] Robert Brandenberger and Patrick Peter. Bouncing Cosmologies: Progress and Problems. *Found. Phys.*, 47(6):797–850, 2017.
- [16] Yi-Fu Cai, Antonino Marciano, Dong-Gang Wang, and Edward Wilson-Ewing. Bouncing cosmologies with dark matter and dark energy. *Universe*, 3(1):1, 2016.
- [17] George F. R. Ellis and Roy Maartens. The emergent universe: Inflationary cosmology with no singularity. *Class. Quant. Grav.*, 21:223–232, 2004.
- [18] George F. R. Ellis, Jeff Murugan, and Christos G. Tsagas. The Emergent universe: An Explicit construction. *Class. Quant. Grav.*, 21(1):233–250, 2004.
- [19] Robert H. Brandenberger and C. Vafa. Superstrings in the Early Universe. *Nucl. Phys.*, B316:391–410, 1989.
- [20] Ali Nayeri, Robert H. Brandenberger, and Cumrun Vafa. Producing a scale-invariant spectrum of perturbations in a Hagedorn phase of string cosmology. *Phys. Rev. Lett.*, 97:021302, 2006.
- [21] Robert H. Brandenberger, Ali Nayeri, Subodh P. Patil, and Cumrun Vafa. Tensor Modes from a Primordial Hagedorn Phase of String Cosmology. *Phys. Rev. Lett.*, 98:231302, 2007.
- [22] Robert H. Brandenberger, Ali Nayeri, Subodh P. Patil, and Cumrun Vafa. String gas cosmology and structure formation. *Int. J. Mod. Phys. A*, 22:3621–3642, 2007.
- [23] Minxi He, Junyu Liu, Shiyun Lu, Siyi Zhou, Yi-Fu Cai, Yi Wang, and Robert Brandenberger. Differentiating G-inflation from String Gas Cosmology using the Effective Field Theory Approach. *JCAP*, 12:040, 2016.
- [24] Thorsten Battefeld and Scott Watson. String gas cosmology. *Rev. Mod. Phys.*, 78:435–454, 2006.
- [25] Robert H. Brandenberger. String Gas Cosmology: Progress and Problems. *Class. Quant. Grav.*, 28:204005, 2011.
- [26] Robert H. Brandenberger. String Gas Cosmology after Planck. *Class. Quant. Grav.*, 32(23):234002, 2015.
- [27] Arvind Borde and Alexander Vilenkin. Eternal inflation and the initial singularity. *Phys. Rev. Lett.*, 72:3305–3309, 1994.
- [28] Arvind Borde, Alan H. Guth, and Alexander Vilenkin. Inflationary space-times are incomplete in past directions. *Phys. Rev. Lett.*, 90:151301, 2003.
- [29] Robert H. Brandenberger. Alternatives to the inflationary paradigm of structure formation. *Int. J. Mod. Phys. Conf. Ser.*, 01:67–79, 2011.
- [30] Robert H. Brandenberger. Introduction to Early Universe Cosmology. *PoS*, ICFI2010:001, 2010.
- [31] Alberto Nicolis, Riccardo Rattazzi, and Enrico Trincherini. The Galileon as a local modification of gravity. *Phys. Rev.*, D79:064036, 2009.
- [32] C. Deffayet, Xian Gao, D. A. Steer, and G. Zahariade. From k-essence to generalised Galileons. *Phys. Rev. D*, 84:064039, 2011.
- [33] Tsutomu Kobayashi, Masahide Yamaguchi, and Jun’ichi Yokoyama. Generalized G-inflation: Inflation with the most general second-order field equations. *Prog. Theor. Phys.*, 126:511–529, 2011.
- [34] Gregory Walter Horndeski. Second-order scalar-tensor field equations in a four-dimensional space. *Int. J. Theor. Phys.*, 10:363–384, 1974.
- [35] Paolo Creminelli, Alberto Nicolis, and Enrico Trincherini. Galilean Genesis: An Alternative to inflation. *JCAP*, 1011:021, 2010.
- [36] Paolo Creminelli, Markus A. Luty, Alberto Nicolis, and Leonardo Senatore. Starting the Universe: Stable Violation of the Null Energy Condition and Non-standard Cosmologies. *JHEP*, 12:080, 2006.
- [37] Yun-Song Piao and E. Zhou. Nearly scale invariant spectrum of adiabatic fluctuations may be from a very slowly expanding phase of the universe. *Phys. Rev. D*, 68:083515, 2003.
- [38] Laurence Perreault L’vasseur, Robert Brandenberger, and Anne-Christine Davis. Defrosting in an Emergent Galileon Cosmology. *Phys. Rev.*, D84:103512, 2011.
- [39] Yi Wang and Robert Brandenberger. Scale-Invariant Fluctuations from Galilean Genesis. *JCAP*, 10:021, 2012.
- [40] Damien A. Easson, Ignacy Sawicki, and Alexander Vikman. When Matter Matters. *JCAP*, 1307:014, 2013.
- [41] V. A. Rubakov. Consistent NEC-violation: towards creating a universe in the laboratory. *Phys. Rev. D*, 88:044015, 2013.
- [42] Benjamin Elder, Austin Joyce, and Justin Khoury. From Satisfying to Violating the Null Energy Condition. *Phys. Rev. D*, 89(4):044027, 2014.
- [43] Paolo Creminelli, Kurt Hinterbichler, Justin Khoury, Alberto Nicolis, and Enrico Trincherini. Subluminal Galilean Genesis. *JHEP*, 02:006, 2013.
- [44] Kurt Hinterbichler, Austin Joyce, Justin Khoury, and Godfrey E. J. Miller. DBI Realizations of the Pseudo-Conformal Universe and Galilean Genesis Scenarios. *JCAP*, 12:030, 2012.
- [45] Kurt Hinterbichler, Austin Joyce, Justin Khoury, and Godfrey E. J. Miller. Dirac-Born-Infeld Genesis: An Improved Violation of the Null Energy Condition. *Phys. Rev. Lett.*, 110(24):241303, 2013.
- [46] David Pirtskhalava, Luca Santoni, Enrico Trincherini, and Patipan Uttayarat. Inflation from Minkowski Space. *JHEP*, 12:151, 2014.
- [47] Sakine Nishi and Tsutomu Kobayashi. Generalized Galilean Genesis. *JCAP*, 03:057, 2015.
- [48] Sakine Nishi and Tsutomu Kobayashi. Scale-invariant perturbations from null-energy-condition violation: A new variant of Galilean genesis. *Phys. Rev. D*, 95(6):064001, 2017.
- [49] Yong Cai, Hai-Guang Li, Taotao Qiu, and Yun-Song Piao. The Effective Field Theory of nonsingular cosmology: II. *Eur. Phys. J. C*, 77(6):369, 2017.
- [50] S. Mironov, V. Rubakov, and V. Volkova. Genesis with general relativity asymptotics in beyond Horndeski the-

- ory. *Phys. Rev. D*, 100(8):083521, 2019.
- [51] Yulia Ageeva, Pavel Petrov, and Valery Rubakov. Horndeski genesis: consistency of classical theory. *JHEP*, 12:107, 2020.
- [52] Yong Cai and Yun-Song Piao. Intermittent null energy condition violations during inflation and primordial gravitational waves. *Phys. Rev. D*, 103(8):083521, 2021.
- [53] M. Libanov, S. Mironov, and V. Rubakov. Generalized Galileons: instabilities of bouncing and Genesis cosmologies and modified Genesis. *JCAP*, 08:037, 2016.
- [54] Tsutomu Kobayashi. Generic instabilities of nonsingular cosmologies in Horndeski theory: A no-go theorem. *Phys. Rev. D*, 94(4):043511, 2016.
- [55] R. Kolevatorov and S. Mironov. Cosmological bounces and Lorentzian wormholes in Galileon theories with an extra scalar field. *Phys. Rev. D*, 94(12):123516, 2016.
- [56] Shingo Akama and Tsutomu Kobayashi. Generalized multi-Galileons, covariantized new terms, and the no-go theorem for nonsingular cosmologies. *Phys. Rev. D*, 95(6):064011, 2017.
- [57] Yong Cai, Youping Wan, Hai-Guang Li, Taotao Qiu, and Yun-Song Piao. The Effective Field Theory of nonsingular cosmology. *JHEP*, 01:090, 2017.
- [58] Yong Cai and Yun-Song Piao. A covariant Lagrangian for stable nonsingular bounce. *JHEP*, 09:027, 2017.
- [59] Anna Ijjas and Paul J. Steinhardt. Fully stable cosmological solutions with a non-singular classical bounce. *Phys. Lett. B*, 764:289–294, 2017.
- [60] Y. A. Ageeva, O. A. Evseev, O. I. Melichev, and V. A. Rubakov. Horndeski Genesis: strong coupling and absence thereof. *EPJ Web Conf.*, 191:07010, 2018.
- [61] Y. Ageeva, O. Evseev, O. Melichev, and V. Rubakov. Toward evading the strong coupling problem in Horndeski genesis. *Phys. Rev. D*, 102(2):023519, 2020.
- [62] Tsutomu Kobayashi, Masahide Yamaguchi, and Jun'ichi Yokoyama. Galilean Creation of the Inflationary Universe. *JCAP*, 07:017, 2015.
- [63] Zhi-Guo Liu, Jun Zhang, and Yun-Song Piao. A Galileon Design of Slow Expansion. *Phys. Rev. D*, 84:063508, 2011.
- [64] Zhi-Guo Liu and Yun-Song Piao. A Galileon Design of Slow Expansion: Emergent universe. *Phys. Lett. B*, 718:734–739, 2013.
- [65] David Langlois and Karim Noui. Hamiltonian analysis of higher derivative scalar-tensor theories. *JCAP*, 07:016, 2016.
- [66] Jibril Ben Achour, Marco Crisostomi, Kazuya Koyama, David Langlois, Karim Noui, and Gianmassimo Tasinato. Degenerate higher order scalar-tensor theories beyond horndeski up to cubic order. *JHEP*, 12:100, 2016.
- [67] David Langlois, Michele Mancarella, Karim Noui, and Filippo Vernizzi. Effective Description of Higher-Order Scalar-Tensor Theories. *JCAP*, 05:033, 2017.
- [68] Jerome Gleyzes, David Langlois, Federico Piazza, and Filippo Vernizzi. Essential Building Blocks of Dark Energy. *JCAP*, 08:025, 2013.
- [69] Jérôme Gleyzes, David Langlois, Federico Piazza, and Filippo Vernizzi. Healthy theories beyond Horndeski. *Phys. Rev. Lett.*, 114(21):211101, 2015.
- [70] Jérôme Gleyzes, David Langlois, Federico Piazza, and Filippo Vernizzi. Exploring gravitational theories beyond Horndeski. *JCAP*, 02:018, 2015.
- [71] Chunshan Lin, Shinji Mukohyama, Ryo Namba, and Rio Saitou. Hamiltonian structure of scalar-tensor theories beyond Horndeski. *JCAP*, 10:071, 2014.
- [72] Xian Gao. Hamiltonian analysis of spatially covariant gravity. *Phys. Rev. D*, 90:104033, 2014.
- [73] Cedric Deffayet, Gilles Esposito-Farese, and Daniele A. Steer. Counting the degrees of freedom of generalized Galileons. *Phys. Rev. D*, 92:084013, 2015.
- [74] Amara Ilyas, Mian Zhu, Yunlong Zheng, Yi-Fu Cai, and Emmanuel N. Saridakis. DHOST Bounce. *JCAP*, 09:002, 2020.
- [75] Mian Zhu, Amara Ilyas, Yunlong Zheng, Yi-Fu Cai, and Emmanuel N. Saridakis. Scalar and Tensor Perturbations in DHOST Bounce Cosmology. 8 2021.
- [76] Misao Sasaki. Gauge Invariant Scalar Perturbations in the New Inflationary Universe. *Prog. Theor. Phys.*, 70:394, 1983.
- [77] Hideo Kodama and Misao Sasaki. Cosmological Perturbation Theory. *Prog. Theor. Phys. Suppl.*, 78:1–166, 1984.
- [78] Viatcheslav F. Mukhanov. Quantum Theory of Gauge Invariant Cosmological Perturbations. *Sov. Phys. JETP*, 67:1297–1302, 1988.
- [79] Xian Gao and Xun-Yang Hong. Propagation of gravitational waves in a cosmological background. *Phys. Rev. D*, 101(6):064057, 2020.
- [80] Yong Cai and Yun-Song Piao. The slow expansion with nonminimal derivative coupling and its conformal dual. *JHEP*, 03:134, 2016.