

WELL-POSEDNESS OF PATH-DEPENDENT SEMILINEAR PARABOLIC MASTER EQUATIONS

SHANJIAN TANG AND HUILIN ZHANG

ABSTRACT. Master equations are partial differential equations for measure-dependent unknowns, and are introduced to describe asymptotic equilibrium of large scale mean-field interacting systems, especially in games and control theory. In this paper we introduce new semilinear master equations whose unknowns are functionals of both paths and path measures. They include state-dependent master equations, path-dependent partial differential equations (PPDEs), history information dependent master equations and time inconsistent (e.g. time-delayed) equations, which naturally arise in stochastic control theory and games. We give a classical solution to the master equation by introducing a new notation called strong vertical derivative (SVD) for path-dependent functionals, inspired by Dupire's vertical derivative [20], and applying stochastic forward-backward system argument. Moreover, we consider a general non-smooth case with a functional mollifying method.

CONTENTS

1. Introduction	2
2. Basic setup and Itô calculus for functionals of path and path-measure	4
2.1. The canonical setup	4
2.2. Strong vertical derivatives with respect to path and path-measure	5
2.3. Itô-Dupire formula	12
3. Differentiability of solutions for path-dependent mean-field BSDEs	16
3.1. First-order differentiability	20
3.2. Second-order differentiability	34
4. Solutions of semilinear path-dependent master equations	38
4.1. The decoupling field and its regularity	38
4.2. Classical solutions of path-dependent master equations	42
4.3. Some typical cases	45
4.4. The general case via functional mollifying	47
5. Appendix	50
5.1. Proof of Lemma 3.4	50
5.2. An extension of [41, Theorem 4.5] without assumption of local Lipschitz continuity in time variable	53
References	54

2010 *Mathematics Subject Classification.* 60G22, 60H10, 34C29.

Key words and phrases. path-dependent, master equation, mean-field games, Itô-Dupire formula.

1. INTRODUCTION

Denote by $\mathbb{C}_{T,d}$ the space of continuous functions on $[0, T]$ with values in \mathbb{R}^d and by \mathcal{P}_2^C the totality of probability measures on $\mathbb{C}_{T,d}$ with finite second order moments. Given deterministic functions (b_1, σ_1) and (b_2, σ_2) on $[0, T]$ with values in $\mathbb{R}^d \times \mathbb{R}^{d \times d}$, we study the following path-dependent parabolic master equation,

$$(1) \quad \begin{cases} \partial_t u(t, \omega, \mu) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \omega, \mu) \sigma_1(t) \sigma_1(t)^T] + \partial_\omega u(t, \omega, \mu) b_1(t) \\ \quad + \frac{1}{2} \text{Tr} [\int_{\mathbb{C}_{T,d}} \partial_{\tilde{\omega}} \partial_\mu u(t, \omega, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) \sigma_2(t) \sigma_2(t)^T] + \int_{\mathbb{C}_{T,d}} \partial_\mu u(t, \omega, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) b_2(t) \\ \quad + f(t, \omega, u(t, \omega, \mu), \sigma_1(t) \partial_\omega u(t, \omega, \mu), \mu, \mathcal{L}_{u(t, W^\mu, \mu)}) = 0, \\ u(T, \omega, \mu) = \Phi(\omega, \mu), \quad (t, \omega, \mu) \in [0, T] \times \mathbb{C}_{T,d} \times \mathcal{P}_2^C. \end{cases}$$

Here, (functional) derivatives ∂_ω and ∂_μ are taken in the spirit of Dupire and Lions (see the subsequent definitions (12) and (29)), respectively, and W^μ represents the canonical processes on $\mathbb{C}_{T,d}$ under μ . Master equations (also called Hamiltonian-Jacobi-Bellman (HJB) equations in a Wasserstein space when concerned with control problems) naturally arise from mean-field games. The mean-field game is introduced independently by Lasry and Lions [35] and Huang, Malhamé and Caines [31] to the study of Nash equilibrium for systems consist of a large number of ‘‘agents’’. The classical mean-field theory, popular in statistical physics (e.g. McKean-Vlasov and Boltzmann models), quantum mechanics and quantum chemistry (e.g. Hartree-Fock type model), was developed in the last century to study systems with large size of particles (see Kac [33], McKean [37], Sznitman [45, 46, 47, 48], Bossy [3] and references therein). In the last decade, the mean-field theory has been widely studied and applied from theoretical areas including stochastic differential games, partial differential equations (PDEs) and stochastic control, etc., to practical areas such as engineering, economics and social science, just to mention a few publications and references therein: [36], [15], [7], [11], [12], [27]. The master equation is a PDE involving measure variables on a Wasserstein space of infinite dimension and introduced by Lions in his lecture [36] for differential games. It is a powerful analytic tool for the study of large systems in physics and games with a mean-field interaction (see [14], [15]), and includes interesting particular cases such as HJB and Fokker-Planck (FP) equations in dynamical systems, stochastic control and mathematical finance of mean-field type. Results are available in various frameworks: Bensoussan et al. [5] consider the regular case when measure variables are restricted on those measures of square integrable density functions, Cardaliaguet [15] gives a viscosity solution for first-order HJB equations on a Wasserstein space, Gomes and Saude [30] survey well-posedness of HJB-FP equations for reduced mean-field games, Buckdahn et al. [9] and Chassagneux et al. [10] study classical solutions for second order master equations through stochastic differential equations (SDEs) and forward backward stochastic differential equations (FBSDEs) respectively, Carmona and Delarue [13] consider the mean-field games and corresponding master equation with common noise, Cardaliaguet et al. [14] give an analytic approach for master equations, Pham and Wei [42] study the dynamic programming principle for Bellman master equation, etc. However, all these works consider the state-dependent case, where (ω, μ) in Equation (1) take values in $\mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^k)$. Here, $\mathcal{P}_2(\mathbb{R}^k)$ is the set of probability measures on \mathbb{R}^k with finite second order moments. In practice, many problems could be non-Markovian or path-dependent: to mention a few, optional pricing for exotic options (e.g. Asian, chooser, lookback and barrier options [20], [19], [32], [26]), stochastic game theory and stochastic control with delayed information ([2], [28], [44], [51], [49]), rough volatility [29], [6], etc.

Dupire [20] introduces a functional Itô formula to incorporate the calculus of path-dependent functionals, which is subsequently developed by Cont-Fournié [16, 17] and references therein (on the other hand, see another approach to path-dependent problems of Flandoli and Zanco [25] by lifting the primal problem into a functional one in Banach spaces). In contrast to the classical functional approach to the path-dependent stochastic analysis (see Ahn [1]), Dupire’s approach is featured by the finite dimensional *vertical derivative* (see the following definition (12)), and is admitted to solve non-Markovian problems, in particular the one proposed by Peng in his ICM 2010 lecture [40] that whether non-Markovian FBSDEs can be connected with path-dependent PDEs (PPDEs). Concerning the well-posedness of PPDEs, Peng and Wang [41] consider smooth solutions of parabolic PPDEs; Ekren et al. [21, 22, 23] study the viscosity solution of quasilinear and fully nonlinear PPDEs; Cosso et al. [18] treat PPDEs as the Hilbert space valued equations and build the viscosity solution; Peng and Song [43] introduce a new path derivative and build Sobolev solutions for corresponding parabolic fully-nonlinear PPDEs via G -BSDEs [39]; Wu and Zhang [50] solve a master equation with solutions in a form of $V(t, \mu)$, $\mu \in \mathcal{P}_2^C$. However, the path-dependent master equation with solutions in the general form of $u(t, \omega, \mu)$, $(\omega, \mu) \in \mathbb{C}_{T,d} \times \mathcal{P}_2^C$ still remains to be studied.

In this article, we study the classical solution of path-dependent master equation (1). In contrast to the state-dependent case [10], smooth solution of equation (1) by FBSDEs meets with new issues. The first comes from the very definition of vertical derivatives with respect to paths and measures on $\mathbb{C}_{T,d}$ (see identities (12) and (29) for details). Dupire’s vertical derivative [20] depends on the cut-off time for paths. In the same spirit, the derivative with respect to measures on the path space (see also [50]), also depends on the cut-off time. Note that in this case two different flows are involved in the probabilistic representation of the path-dependent field u . Application of the flow property of solutions for SDEs involves vertical derivatives with respect to the path and the path measure at two different times, while only one cut-off time is available in the field u . Indeed, one of them should describe the smoothness of u on “past” information and is not defined in the sense of Dupire, which is only available for non-anticipative (or adapted) functionals and describe the smoothness on the “present” information. Secondly, the existence of the derivative with respect to measure in Lions’ sense usually requires the separability of the measurable space, which is the space of càdlàg functions here in view of Dupire’s vertical derivative. However, FBSDEs are more consistent with the uniform norm instead of Skorokhod norm, which leaves us without the general existence result for derivatives. To handle the first issue, we propose a new notation called “strong vertical derivative” (SVD) (see Definition 2.1) which is derived from the vertical derivative of Dupire and restricts functionals to be smooth before the cut-off time. By definition, non-anticipative functionals with SVDs also have vertical derivatives. Moreover, the assumption of SVDs is general enough to include all interesting continuously vertical differentiable functionals (see Example 2.2). On the other hand, the SVD can be viewed as a pathwise definition for the Malliavin derivative (see e.g. [38]) on the canonical space (see subsequent Remark 2.3 for details). Based on the differentiability with respect to paths and path measures, we build the Itô formula and the partial Itô formula. Then we consider the non-Markovian FBSDE

$$(2) \quad \begin{cases} dX(r) = b_1(r)dr + \sigma_1(r)dB(r), \\ dX'(r) = b_2(r)dr + \sigma_2(r)dB(r), \\ dY(r) = -f(X_r, Y(r), Z(r), \mathcal{L}_{X'_r}, \mathcal{L}_{Y_{\eta_t(r)}})dr + Z(r)dB(r), \quad r \geq t, \\ X_t = \gamma_t, \quad X'_t = \eta_t, \quad Y(T) = \Phi(X_T, \mathcal{L}_{X'_T}), \end{cases}$$

where η is a continuous process with law μ , and Y^{η_t} solves the associated mean-field FBSDE

$$(3) \quad \mathcal{Y}(s) = \Phi(X_T^{\eta_t}, \mathcal{L}_{X_T^{\eta_t}}) + \int_s^T f(X_r^{\eta_t}, \mathcal{Y}(r), \mathcal{Z}(r), \mathcal{L}_{X_r^{\eta_t}}, \mathcal{L}_{\mathcal{Y}(r)}) dr - \int_s^T \mathcal{Z}(r) dB(r), \quad s \in [t, T],$$

with $X_r^{\eta_t} := X_r|_{\gamma=\eta}$ for $r \in [t, T]$. Here we denote by $\omega(t)$ the value of path ω at time t and by $\omega_t(\cdot) := \omega(t \wedge \cdot)$ the path up to time t . Assuming that the functional generator f and terminal value Φ have smooth SVDs, we prove that the solution of corresponding FBSDE also has smooth SVDs. Moreover, we construct all strong vertical derivatives of $u(t, \omega, \mu)$ via FBSDE, and give the required regularity to apply our Itô formula (see Theorem 2.15 and Corollary 2.16)—which will benefit both numerical and theoretical approximation of equilibrium by finite systems (see [24], [34]). Furthermore, we also address some non-smooth case and connect it with viscosity solutions via a functional mollifying argument, which is illustrated with typical examples of the time-delayed case.

In summary, our main contribution is three-fold. Firstly, we propose the general form of path-dependent master equation (1) and give the well-posedness. Secondly, we introduce the notation of strong vertical differentiability and build Itô and partial Itô formulas in this framework which are fundamental tools in the study of path-dependent mean-field problems. Thirdly, the argument of functional smoothing also seems new in the path-dependent framework.

The rest of the article is organized as follows. In Section 2, we introduce notations of SVD with respect to paths and measures on path space, and build in the framework functional Itô calculus incorporating paths and path measures. In Section 3, we show the differentiability and regularity of associated FBSDE solutions. In Section 4, we prove the existence and uniqueness of smooth solutions for path-dependent parabolic master equation. Moreover, we extend our result to more general cases by a functional mollifying argument.

Acknowledgements. This work is supported by NSF of China (Grant Numbers 12031009, 11901104) and Laboratory of Mathematics for Nonlinear Science, Fudan University, Shanghai, China.

2. BASIC SETUP AND ITÔ CALCULUS FOR FUNCTIONALS OF PATH AND PATH-MEASURE

2.1. The canonical setup. For any fixed $T > 0$, we denote by $\mathbb{C}_{T,d} = C([0, T], \mathbb{R}^d)$ the canonical space and equip it with the supreme norm $\|\cdot\|_{[0,T]}$. W is the canonical process and $\{\mathcal{F}_t^W\}_{0 \leq t \leq T}$ is the natural filtration. For any $(t, \omega) \in [0, T] \times \mathbb{C}_{T,d}$, ω_t is the cut-off path, meaning that $\omega_t \in \mathbb{C}_{T,d}$ such that

$$(4) \quad \omega_t(r) = \omega(r)1_{[0,t)}(r) + \omega(t)1_{[t,T]}(r), \quad r \in [0, T];$$

and $\omega(t)$ is the state of ω at time t . Let \mathcal{P}_2^C be the set of probability measures on $(\mathbb{C}_{T,d}, \mathcal{F}_T^W)$ with finite second order moments, i.e. $\mu \in \mathcal{P}_2^C$ iff $\|\mu\|^2 := \mathbb{E}^\mu[\|W\|_{[0,T]}^2] < \infty$. For $\mu \in \mathcal{P}_2^C$, $\mu_t \in \mathcal{P}_2^C$ is the distribution of stopped process W_t under μ . For any $\mu, \nu \in \mathcal{P}_2^C$, we define the following classical 2-Wasserstein distance

$$(5) \quad W_2(\mu, \nu) = \inf_{P \in \mathcal{P}(\mu, \nu)} \left(\int_{\mathbb{C}_{T,d} \times \mathbb{C}_{T,d}} \|u - v\|_{[0,T]}^2 dP(u, v) \right)^{\frac{1}{2}},$$

where $\mathcal{P}(\mu, \nu)$ is the set of all probability measures on $(\mathbb{C}_{T,d} \times \mathbb{C}_{T,d}, \mathcal{F}_T^W \times \mathcal{F}_T^W)$ with marginal measures μ and ν . To introduce functional derivative in the space of paths, we consider the space of càdlàg paths $\mathbb{D}_{T,d} := D([0, T], \mathbb{R}^d)$, which can be equipped with the uniform topology $\|\cdot\|_{[0,T]}$, or the Skorohod topology

$$(6) \quad d(\omega, \omega') := \inf_{\lambda \in \Lambda_{[0,T]}} \sup_{t \in [0,T]} (|t - \lambda(t)| + |\omega(t) - \omega'(t)|),$$

where $\Lambda_{[0,T]}$ is the set of all strictly increasing continuous mappings on $[0, T]$ with $\lambda(0) = 0$ and $\lambda(T) = T$. In the following, we equip $\mathbb{D}_{T,d}$ with the uniform topology unless stated otherwise. With the space $\mathbb{C}_{T,d}$ being replaced with $\mathbb{D}_{T,d}$, notations such as \mathcal{P}_2^D and $W_2(\mu, \nu)$ are self-explained.

Suppose that (Ω, \mathcal{F}, P) is an atomless probability space supporting a d -dimensional Brownian motion B , and $\{\mathcal{F}_t\}_{t \in [0, T]}$ is the natural augmented filtration. For any $t \in [0, T]$ and $r \in [t, T]$, we define \mathcal{F}_r^t as the σ -algebra generated by $\{B(s) - B(t); t \leq s \leq r\}$ and completed under P . For any (stopped up to time t) process X_t , we denote by \mathcal{L}_{X_t} the law of the process X_t and $\mathcal{L}_{X(t)}$ the law of the random variable $X(t)$. In the following, we use notation \mathbb{M}_2^C (\mathbb{M}_2^D , resp.) as the collection of measurable continuous processes (càdlàg processes, resp.) with laws in \mathcal{P}_2^C (\mathcal{P}_2^D , resp.). Since for any $\mu \in \mathcal{P}_2^D$, we can always find an atomless probability space $(\Omega, \mathcal{F}_t, P)$ such that there exists a càdlàg process η on this probability space with law μ , we will always suppose for any $\mu \in \mathcal{P}_2^D$, (Ω, \mathcal{F}, P) is rich enough to support a càdlàg process η such that $\mathcal{L}_\eta = \mu$. Moreover, for any progressively measurable process X and random variable ξ on (Ω, \mathcal{F}, P) , we define the following norms if they are finite: for any $t \in [0, T]$, $p \in \mathbb{N}^+$,

$$(7) \quad \|X\|_{\mathbb{S}^p, [t, T]}^p := \mathbb{E}^P[\|X\|_{[t, T]}^p], \quad \|X\|_{\mathbb{H}^p, [t, T]}^p := \mathbb{E}^P[(\int_t^T |X(r)|^2 dr)^{\frac{p}{2}}], \quad \|\xi\|_{L^p}^p := \mathbb{E}^P[|\xi|^p].$$

We write $\mathbb{S}^p([t, T], \mathbb{R}^k)$, $\mathbb{H}^p([t, T], \mathbb{R}^k)$ and $L^p(\mathcal{F}_T, \mathbb{R}^k)$ for spaces of progressively measurable processes on $[t, T]$ and random variables with values in \mathbb{R}^k and finite corresponding norms. Denote by $C^n(\mathbb{R}^m, \mathbb{R}^k)$ ($C_b^n(\mathbb{R}^m, \mathbb{R}^k)$, resp.) the space of (bounded, resp.) continuous functions from \mathbb{R}^m to \mathbb{R}^k with (bounded, resp.) continuous derivatives up to order n . Usually, we omit \mathbb{R}^k in $\mathbb{S}^p([t, T], \mathbb{R}^k)$, $\mathbb{H}^p([t, T], \mathbb{R}^k)$, $L^p(\mathcal{F}_T, \mathbb{R}^k)$, $C(\mathbb{R}^m, \mathbb{R}^k)$ when $k = 1$, and also omit the time interval $[t, T]$ if no confusion raised. Moreover, for $(Y, Z) \in \mathbb{S}^p([t, T], \mathbb{R}^m) \times \mathbb{H}^p([t, T], \mathbb{R}^n)$, we write

$$(8) \quad \|(Y, Z)\|_{\mathbb{S}^p \times \mathbb{H}^p} := (\|Y\|_{\mathbb{S}^p}^p + \|Z\|_{\mathbb{H}^p}^p)^{\frac{1}{p}}.$$

2.2. Strong vertical derivatives with respect to path and path-measure. Denote by $\hat{\mathbb{D}}_{T,d}$ the product space $[0, T] \times \mathbb{D}_{T,d} \times \mathcal{P}_2^D$ and by \mathcal{D} the space of functionals on $\hat{\mathbb{D}}_{T,d}$. A functional $f \in \mathcal{D}$ is said to be *non-anticipative* if for any (t, ω, μ) , $f(t, \omega, \mu) = f(t, \omega_t, \mu_t)$. For non-anticipative $f \in \mathcal{D}$, we call f continuous on $\hat{\mathbb{D}}_{T,d}$ and write $f \in \mathcal{C}(\hat{\mathbb{D}}_{T,d})$ if f is continuous in the product space $[0, T] \times \mathbb{D}_{T,d} \times \mathcal{P}_2^D$ equipped with the premetric:

$$(9) \quad d_p((t, \omega, \mu), (t', \omega', \mu')) := |t - t'| + \|\omega_t - \omega_{t'}\| + W_2(\mu_t, \mu_{t'}).$$

For any non-anticipative $f \in \mathcal{D}$, the horizontal derivative is defined as

$$(10) \quad \partial_t f(t, \omega, \mu) := \lim_{h \rightarrow 0^+} \frac{1}{h} [f(t+h, \omega_t, \mu_t) - f(t, \omega_t, \mu_t)], \quad \forall (t, \omega, \mu) \in \hat{\mathbb{D}}_{T,d}.$$

For any $(t, x) \in [0, T] \times \mathbb{R}^d$, define $\omega^{t,x} \in \mathbb{D}_{T,d}$ by

$$(11) \quad \omega^{t,x} := \omega + x1_{[t, T]}.$$

For any fixed $(t, \mu) \in [0, T] \times \mathcal{P}_2^D$, $f(t, \cdot, \mu) : \mathbb{D}_{T,d} \rightarrow \mathbb{R}$ is called *vertically differentiable* at (t, ω) (or ω_t for short), if $f(t, \omega^{t,x}, \mu)$ is differentiable at $x = 0$, i.e. there exists $\partial_\omega f(t, \omega, \mu) \in \mathbb{R}^d$ such that

$$(12) \quad f(t, \omega + x1_{[t,T]}, \mu) = f(t, \omega, \mu) + \partial_\omega f(t, \omega, \mu)x + o(|x|), \quad \forall x \in \mathbb{R}^d,$$

and $\partial_\omega f(t, \omega, \mu)$ is then called the vertical derivative. Now we extend the vertical derivative from non-anticipative functionals taken at cut-off time to path functionals at any time before the cut-off time.

Definition 2.1. *Suppose that $f : [0, T] \times \mathbb{D}_{T,d} \rightarrow \mathbb{R}$. For any $\tau \leq t$, we call f strongly vertically differentiable at (τ, t, ω) (or ω_τ for short), if there exists $\partial_{\omega_\tau} f(t, \omega) \in \mathbb{R}^d$ such that*

$$(13) \quad f(t, \omega + x1_{[\tau,T]}) = f(t, \omega) + \partial_{\omega_\tau} f(t, \omega)x + o(|x|), \quad \forall x \in \mathbb{R}^d.$$

In this case, $\partial_{\omega_\tau} f(t, \omega)$ is called the strong vertical derivative of f at (τ, t, ω) . Moreover, if f is strongly vertically differentiable at (τ, t, ω) for any $\tau \leq t$, we call f strongly vertically differentiable at (t, ω) (or ω_t for short).

Clearly, f is strongly vertical differentiable at ω_t if and only if the mapping $x \rightarrow f(t, \omega^{\tau,x})$ is differentiable at $x = 0$ for any $\tau \leq t$. In particular, if f is non-anticipative and strongly vertically differentiable, f is vertically differentiable and its vertical derivative at (t, x) agrees with its strong vertical derivative at (t, t, ω) . For the SVD $\partial_{\omega_\tau} f(t, \omega)$, we can further define its SVDs in the same spirit: for any $\tau' \leq t$, define $\partial_{\omega_{\tau'}} \partial_{\omega_\tau} f(t, \omega)$ as the SVD of $\partial_{\omega_\tau} f(t, \omega)$ at (τ', t, ω) . In the following, we only need to consider the case $\tau' = \tau$. We call f has continuous SVDs or $\partial_{\omega_\tau} f(t, \omega)$ is continuous if $\partial_{\omega_\tau} f$ is continuous with respect to the metric: for any (τ, t, ω) and (τ', t', ω') with $\tau \leq t$, $\tau' \leq t'$,

$$(14) \quad d_{sp}((\tau, t, \omega), (\tau', t', \omega')) := |\tau - \tau'| + |t - t'| + \|\omega_t - \omega'_{t'}\|.$$

Here are examples of strongly vertically differentiable functionals.

Example 2.2. *Let $f : [0, T] \times \mathbb{D}_{T,d} \rightarrow \mathbb{R}$ and $(t, \omega) \in [0, T] \times \mathbb{D}_{T,d}$.*

(i) *If $f(t, \omega) = F(t, \omega(t))$ for a function $F \in C^{1,k}([0, T] \times \mathbb{R}^d)$, then we have that for any $\tau_1, \tau_2, \dots, \tau_j \in [0, t]$, $j \leq k$,*

$$(15) \quad \partial_t f(t, \omega) = \partial_t F(t, \omega(t)), \quad \partial_{\omega_{\tau_j}} \cdots \partial_{\omega_{\tau_1}} f(t, \omega) = D_x^j F(t, \omega(t)),$$

and thus f has continuous strong vertical derivatives up to order k .

(ii) *Suppose that $f(t, \omega) = \int_0^t F(r, \omega(r))dr$ with $F \in C^{1,k}([0, T] \times \mathbb{R}^d)$. Then for any $\tau_1, \tau_2, \dots, \tau_j \in [0, t]$, $j \leq k$,*

$$(16) \quad \partial_t f(t, \omega) = F(t, \omega(t)), \quad \partial_{\omega_{\tau_j}} \cdots \partial_{\omega_{\tau_1}} f(t, \omega) = \int_\tau^t D_x^j F(r, \omega(r))dr,$$

with $\tau = \max_{1 \leq i \leq j} \{\tau_i\}$. Thus f has continuous SVDs up to order k .

(iii) *For a partition $0 = t_0 < t_1 < \dots < t_n = T$, and a continuously differentiable function $F : \underbrace{\mathbb{R}^d \times \mathbb{R}^d \times \dots \times \mathbb{R}^d}_n \rightarrow \mathbb{R}$, let*

$$(17) \quad f(T, \omega) := F(\omega(t_1), \omega(t_2) - \omega(t_1), \dots, \omega(T) - \omega(t_{n-1})).$$

Then f is strongly vertically differentiable at (T, ω) with

$$(18) \quad \partial_{\omega_t} f(T, \omega) = \sum_{j=1}^n \partial_{x_j} F(\omega(t_1), \omega(t_2) - \omega(t_1), \dots, \omega(T) - \omega(t_{n-1})) 1_{(t_{j-1}, t_j]}(t), \quad t > 0.$$

(iv) For fixed $t_0 \in (0, T)$ and $F \in C^1(\mathbb{R}^d)$, define $f(T, \omega) := F(\omega(t_0))$. Thus f has SVDs

$$(19) \quad \partial_{\omega_t} f(T, \omega) = D_x F(\omega(t_0)) 1_{[0, t_0]}(t),$$

which may not be continuous in $t \in [0, T]$. However, consider

$$(20) \quad f_\varepsilon(T, \omega) := \int_0^T \rho_\varepsilon(t_0 - s) F(\omega(s)) ds, \quad \omega \in \mathbb{D}_{T,d}, \quad \varepsilon > 0,$$

with ρ_ε a standard mollifier on \mathbb{R} . Then, for any $\omega \in \mathbb{C}_{T,d}$, $\lim_{\varepsilon \rightarrow 0} f_\varepsilon(T, \omega) = f(T, \omega)$. Moreover, according to (ii), $f_\varepsilon(T, \omega)$ has continuous SVDs. Therefore, we can approximate path-dependent master equations with non-smooth driven and terminal functionals by those with smooth ones, see Example 4.11 for further details.

(v) For a given partition of $[0, T] : 0 = t_0 < t_1 < \dots < t_n = T$ and smooth functions $\{f_i\}_{i=0}^{n-1}$ on \mathbb{R}^d , consider

$$(21) \quad f(t, \omega) := \sum_{i=0}^{n-1} f_i(\omega(t_i)) 1_{[t_i, t_{i+1})}(t).$$

Then f is strongly vertically differentiable at ω_t with

$$(22) \quad \partial_{\omega_\tau} f(t, \omega) = \sum_{i=0}^{n-1} Df_i(\omega(t_i)) 1_{[t_i, t_{i+1})}(t) 1_{[0, t_i]}(\tau), \quad \forall \tau \leq t.$$

Remark 2.3. The relation between vertical derivative and Malliavin derivative is considered in [17], where an equivalence is built through martingale representation in both frameworks (see [17, Theorem 6.1]). However, according to (iii) of Example 2.2, there is a direct equivalence between SVDs and Malliavin derivatives. Recall that W is the canonical process, and then $f(W) := F(W(t_1), W(t_2) - W(t_1), \dots, W(T) - W(t_{n-1}))$ gives a cylindrical random variable under the Wiener measure. Then its Malliavin derivative $\mathcal{D}_r f(W)$ agrees with its SVD at (r, T, W) and then the SVD can be viewed as a pathwise definition of Malliavin derivative without involving any probability measure. Furthermore, we can consider SVDs with respect to driven signals for integrals and equations by restricting the domain of definition. Denote by $\mathbb{C}_{T,d}^1$ the subspace of $\mathbb{C}_{T,d}$ with continuous derivatives. Consider functional $g : [0, T] \times \mathbb{C}_{T,d}^1 \rightarrow \mathbb{R}$ given by

$$(23) \quad g(t, \omega) := \int_0^t G(\omega(r)) d\omega(r), \quad (t, \omega) \in [0, T] \times \mathbb{C}_{T,d}^1,$$

with $G \in C_b^1(\mathbb{R}^d, \mathbb{R}^d)$. Then, we have

$$(24) \quad \partial_{\omega_\tau} g(t, \omega) = G(\omega(\tau)) + \int_\tau^t DG(\omega(r)) d\omega(r), \quad \tau \in [0, t].$$

Similarly, consider $\phi : [0, T] \times \mathbb{C}_{T,d}^1 \rightarrow \mathbb{R}$ given by $\phi(t, \omega) := x(t)$ with

$$(25) \quad x(t) = \int_0^t H(x(r)) d\omega(r),$$

for a given function $H \in C_b^2(\mathbb{R}, \mathbb{R}^d)$. Then by a nontrivial argument (note that ordinary differential equations (ODEs) are not continuous with respect to the driven signal under the uniform norm), we have that for any $(t, \omega) \in [0, T] \times \mathbb{C}_{T,d}^1$, ϕ is strongly vertically differentiable at any (t, ω) and the derivative $\partial_{\omega_\tau} \phi(t, \omega)$ at (τ, t, ω) solves the following linear ODE,

$$(26) \quad \partial_{\omega_\tau} x(t) = H(x(\tau)) + \int_\tau^t \partial_{\omega_\tau} x(r) H'(x(r)) d\omega(r), \quad t \geq \tau.$$

The following lemma follows from Definition 2.1 directly.

Lemma 2.4. *Suppose that $f : [0, T] \times \mathbb{D}_{T,d} \rightarrow \mathbb{R}$ is strongly vertically differentiable, and uniformly Lipschitz continuous in ω :*

$$(27) \quad |f(t, \omega) - f(t, \omega')| \leq C \|\omega_t - \omega'_t\|, \quad \forall (t, \omega, \omega') \in [0, T] \times \mathbb{D}_{T,d} \times \mathbb{D}_{T,d}.$$

Then we have $|\partial_{\omega_\tau} f(t, \omega)| \leq C$ for any $(t, \omega) \in [0, T] \times \mathbb{D}_{T,d}$ and $\tau \leq t$.

For a non-anticipative functional $f \in \mathcal{D}$, consider its lift $\mathbf{f} : [0, T] \times \mathbb{D}_{T,d} \times \mathbb{M}_2^D \rightarrow \mathbb{R}$,

$$(28) \quad \mathbf{f}(t, \omega, \eta) := f(t, \omega, \mathcal{L}_\eta).$$

In the spirit of Lions [36] (also see [50] for derivative with respect to measure on the path space), we call f Fréchet (vertically) differentiable at (t, μ) (or μ_t for short), if for any fixed ω , \mathbf{f} is Fréchet (vertically) differentiable at (t, η) (or η_t for short) with $\mathcal{L}_\eta = \mu$ in the following sense: there exists $D_\eta \mathbf{f}(t, \omega, \eta) \in L_P^2(\mathcal{F}_t, \mathbb{R}^d)$ such that for any $\xi \in L_P^2(\mathcal{F}_t, \mathbb{R}^d)$,

$$(29) \quad \mathbf{f}(t, \omega, \eta + \xi 1_{[t, T]}) = \mathbf{f}(t, \omega, \eta) + \mathbb{E}^P[D_\eta \mathbf{f}(t, \omega, \eta) \xi] + o(\|\xi\|_{L^2}).$$

In particular, it means that the following Gâteaux derivative exists

$$(30) \quad \lim_{h \rightarrow 0} \frac{1}{h} [\mathbf{f}(t, \omega, \eta + h \xi 1_{[t, T]}) - \mathbf{f}(t, \omega, \eta)] = \mathbb{E}^P[D_\eta \mathbf{f}(t, \omega, \eta) \xi].$$

Moreover, if there exists a non-anticipative jointly measurable functional $\partial_\mu f : \hat{\mathbb{D}}_{T,d} \times \mathbb{D}_{T,d} \rightarrow \mathbb{R}$, such that

$$(31) \quad D_\eta \mathbf{f}(t, \omega, \eta) = \partial_\mu f(t, \omega, \mu, \eta), \quad P\text{-a.s.},$$

we call f vertically differentiable at (t, μ) and $\partial_\mu f(t, \omega, \mu, \tilde{\omega})$ the vertical derivative of $f(t, \omega, \cdot)$ at (t, μ) (or μ_t).

Remark 2.5. *Here we give a crucial remark about the validity for notations of Fréchet and Gâteaux differentiability. Denote by \mathbf{f} the lift of $f \in \mathcal{D}$. For any $\xi \in L_P^2(\mathcal{F}_t, \mathbb{R}^d)$, let $F(t, \omega, \eta, \xi) := \mathbf{f}(t, \omega, \eta + \xi 1_{[t, T]})$. Then \mathbf{f} is Fréchet differentiable at (t, η) in the above sense is equivalent to that $F(t, \omega, \eta, \xi)$ is Fréchet differentiable at $\xi = 0$ in the classical sense. Similar argument for Gâteaux differentiability also holds.*

Remark 2.6. *Concerning the existence of the derivative functional $\partial_\mu f$. If the lift $\mathbf{f}(t, \omega, \eta)$ of $f(t, \omega, \mu)$ is Fréchet differentiable at η_t , and the derivative $D_\eta \mathbf{f}(t, \omega, \eta)$ is continuous in the sense that $D_\eta \mathbf{f}(t, \omega, \eta^n) \xrightarrow{L^2} D_\eta \mathbf{f}(t, \omega, \eta)$ as $\eta^n \xrightarrow{L^2} \eta$ under the Skorohod topology (6), then according to [50, Theorem 2.2], $\partial_\mu f$ exists in the sense of (31). However, to build smooth solutions for (1),*

we need our Itô formula (Theorem 2.15 and Corollary 2.16) to be applicable for the larger class of functionals, which only need to be continuous with respect to the uniform topology. Luckily, we can construct the derivative directly by corresponding FBSDEs.

For the uniqueness of $\partial_\mu f(t, \omega, \mu, \cdot)$, in view of identity (31), we see that it is unique μ -a.s. in $\mathbb{D}_{T,d}$. Then for any $\mu \in \mathcal{P}_2^D$ such that $\text{supp}(\mu) = \mathbb{D}_{T,d}$, if $\partial_\mu f(t, \omega, \mu, \tilde{\omega})$ is continuous in $\tilde{\omega} \in \mathbb{D}_{T,d}$, $\partial_\mu f(t, \omega, \mu, \cdot)$ is unique on $\mathbb{D}_{T,d}$. Moreover, suppose that $\partial_\mu f(t, \omega, \cdot, \cdot)$ is jointly continuous on $\mathcal{P}_2^D \times \mathbb{D}_{T,d}$. Then for any $\mu_0 \in \mathcal{P}_2^D$, $\partial_\mu f(t, \omega, \mu_0, \cdot)$ is unique on $\mathbb{D}_{T,d}$. Indeed, choose any $\eta \in \mathbb{M}_2^D$ with $\mathcal{L}_\eta = \mu_0 \in \mathcal{P}_2^D$, and any $\eta' \in (\mathbb{M}_2^D)'$, which is independent of η , such that $\text{supp}(\mathcal{L}_{\eta'}) = \mathbb{D}_{T,d}$. Then for any $\varepsilon > 0$, the functional $\partial_\mu f(t, \omega, \mathcal{L}_{\eta+\varepsilon\eta'}, \cdot)$ is unique on $\mathbb{D}_{T,d}$. It follows by continuity of $\partial_\mu f(t, \omega, \cdot, \cdot)$ that $\partial_\mu f(t, \omega, \mu_0, \tilde{\omega})$ is unique as the limit of $\partial_\mu f(t, \omega, \mathcal{L}_{\eta+\varepsilon\eta'}, \tilde{\omega})$ as ε goes to zero. In conclusion, we have the following lemma.

Lemma 2.7. *Suppose that for any fixed $(t, \omega) \in [0, T] \times \mathbb{D}_{T,d}$, the functional derivative $\partial_\mu f(t, \omega, \cdot, \cdot)$ is jointly continuous in $\mathcal{P}_2^D \times \mathbb{D}_{T,d}$. Then for any $(t, \omega, \mu) \in \mathbb{D}_{T,d}$, $\partial_\mu f(t, \omega, \mu, \cdot)$ is unique on $\mathbb{D}_{T,d}$.*

Remark 2.8. *The definition of vertical derivative given by (29) and (30) has natural extension for Banach space valued functionals. For any $t \in [0, T]$, suppose that $f(t, \omega, \mu)$ takes values in a (stochastic) Banach space E_t (e.g. $\mathbb{S}^2([t, T])$, $\mathbb{H}^2([t, T])$, $L^2(\mathcal{F}_t)$). Indeed, $f(t, \omega, \mu)$ has the natural lift $\mathbf{f}(t, \omega, \eta) \in E_t$ with $\mathcal{L}_\eta = \mu$. If the mapping from $L^2(\mathcal{F}_t)$ to E_t*

$$\mathbf{f}(t, \omega, \eta + \cdot 1_{[t, T]}) : L^2(\mathcal{F}_t) \longrightarrow E_t \\ \xi \qquad \qquad \qquad \mathbf{f}(t, \omega, \eta + \xi 1_{[t, T]})$$

is Fréchet (vertical) differentiable with derivative $D_\eta \mathbf{f}(t, \omega, \eta) \in L(L^2(\mathcal{F}_t), E_t)$ at $\xi = 0$, we call $f(t, \omega, \cdot)$ Fréchet (vertically) differentiable at μ_t . Moreover, if there exists a jointly measurable functional $U : \mathbb{D}_{T,d} \times \mathbb{D}_{T,d} \rightarrow E_t$ such that for any $\xi \in L^2(\mathcal{F}_t)$, $D_\eta \mathbf{f}(t, \omega, \eta)(\xi) = \mathbb{E}^P[U(t, \omega, \mu, \eta)\xi]$, we call $\partial_\mu f(t, \omega, \mu, \cdot) := U(t, \omega, \mu, \cdot)$ the vertical derivative of $f(t, \omega, \cdot)$ at μ_t .

Now we introduce SVDs with respect to path-measure.

Definition 2.9. *For any $\tau, t \in [0, T]$ with $\tau \leq t$ and $\mu \in \mathcal{P}_2^D$, we call a non-anticipative functional $f : [0, T] \times \mathcal{P}_2^D \rightarrow \mathbb{R}$ Fréchet (strongly vertically) differentiable at (τ, t, μ) if its lift $\mathbf{f}(t, \eta)$ with $\mathcal{L}_\eta = \mu$ is Fréchet (strongly vertically) differentiable: there exists $D_{\eta_\tau} \mathbf{f}(t, \eta) \in L_P^2(\mathcal{F}_t, \mathbb{R}^d)$ such that for any $\xi \in L_P^2(\mathcal{F}_\tau, \mathbb{R}^d)$,*

$$(32) \quad \mathbf{f}(t, \eta + \xi 1_{[\tau, T]}) = \mathbf{f}(t, \eta) + \mathbb{E}^P[D_{\eta_\tau} \mathbf{f}(t, \eta)\xi] + o(\|\xi\|_{L^2}).$$

In particular, it means that the following Gâteaux derivative exists,

$$(33) \quad \lim_{h \rightarrow 0} \frac{1}{h} [\mathbf{f}(t, \eta + h\xi 1_{[\tau, T]}) - \mathbf{f}(t, \eta)] = \mathbb{E}^P[D_{\eta_\tau} \mathbf{f}(t, \eta)\xi].$$

We call f strongly vertically differentiable at (t, μ) or μ_t , if it is Fréchet differentiable at (τ, t, μ) for any $\tau \leq t$, and moreover, there exists a jointly measurable non-anticipative functional $\partial_{\mu_\tau} f : [0, T] \times \mathcal{P}_2^D \times \mathbb{D}_{T,d} \rightarrow \mathbb{R}^d$ such that

$$(34) \quad D_{\eta_\tau} \mathbf{f}(t, \eta) = \partial_{\mu_\tau} f(t, \mu, \eta), \quad P\text{-a.s.}$$

$\partial_{\mu_\tau} f(t, \mu, \cdot)$ is then called the strong vertical derivative of $f(t, \cdot)$ at (τ, t, μ) .

Remark 2.10. *For the existence and uniqueness of the SVD at μ_τ , we have similar results as Remark 2.6 and Lemma 2.7 for vertical derivatives. In particular, if for any $t \in [0, T]$, $\partial_{\mu_\tau} f(t, \cdot, \cdot)$ is jointly continuous on $\mathcal{P}_2^D \times \mathbb{D}_{T,d}$, then the SVD is unique. Moreover, we can extend SVDs in path-measure to the (stochastic) Banach framework as Remark 2.8.*

Given strongly vertically differentiable $f : [0, T] \times \mathcal{P}_2^D \rightarrow \mathbb{R}$, for any $(t, \mu, \tilde{\omega}) \in [0, T] \times \mathcal{P}_2^D \times \mathbb{D}_{T,d}$ and $\tau \leq t$, we can further consider SVDs of $\partial_{\mu_\tau} f$ with respect to μ_t and $\tilde{\omega}_t$: for any $\tau' \leq t$, consider $\partial_{\tilde{\omega}_{\tau'}} \partial_{\mu_{\tau'}} f(t, \mu, \tilde{\omega})$ as the SVD of $\partial_{\mu_\tau} f(t, \mu, \tilde{\omega})$ at $(\tau', t, \tilde{\omega})$; $\partial_{\mu_{\tau'}} \partial_{\mu_\tau} f(t, \mu, \tilde{\omega}, \tilde{\omega}')$ as the SVD of $\partial_{\mu_\tau} f(t, \mu, \tilde{\omega})$ at (τ', t, μ) . In the subsequent sections, we only need to consider the case $\tau' = \tau$ and the second order derivative $\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(t, \mu, \tilde{\omega})$. Moreover, we call f has continuous SVDs or $\partial_{\mu_\tau} f(t, \mu, \tilde{\omega})$ is continuous if $\partial_{\mu_\tau} f$ is continuous with respect to the following premetric: for any $(t, \mu, \tilde{\omega})$ and $(t', \mu', \tilde{\omega}')$ with $\tau \leq t, \tau' \leq t'$,

$$(35) \quad d_{sp}((\tau, t, \mu, \tilde{\omega}), (\tau', t', \mu', \tilde{\omega}')) := |\tau - \tau'| + |t - t'| + W_2(\mu_t, \mu_{t'}) + \|\tilde{\omega}_t - \tilde{\omega}'_{t'}\|.$$

f is said to have continuous SVDs in path-measure up to order 2, if both $\partial_{\mu_\tau} f$ and $\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f$ are continuous with respect to the above topology.

Example 2.11. Here we consider $f : [0, T] \times \mathcal{P}_2^D \rightarrow \mathbb{R}$ and $(t, \mu) \in [0, T] \times \mathcal{P}_2^D$.

- (i) Suppose that $F \in C^{1,2}([0, T] \times \mathbb{R}^d)$ with $|D_x^2 F|$ being uniformly bounded, and $f(t, \mu) := \mathbb{E}^\mu[F(t, W(t))]$. Then we have that

$$\begin{aligned} \partial_t f(t, \mu) &= \mathbb{E}^\mu[\partial_t F(t, W(t))], & \partial_{\mu_\tau} f(t, \mu, \tilde{\omega}) &= D_x F(t, \tilde{\omega}(t)), \\ \text{and } \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(t, \mu, \tilde{\omega}) &= D_x^2 F(t, \tilde{\omega}(t)), & \forall \tau \in [0, t]. \end{aligned}$$

Thus f has continuous SVDs up to order 2.

- (ii) Let F as defined in (i) and $f(t, \mu) := \mathbb{E}^\mu[\int_0^t F(r, W(r))dr]$. Then for any $\tau \in [0, t]$,

$$\begin{aligned} \partial_t f(t, \mu) &= \mathbb{E}^\mu[F(t, W(t))], & \partial_{\mu_\tau} f(t, \mu, \tilde{\omega}) &= \int_\tau^t D_x F(r, \tilde{\omega}(r))dr, \\ \text{and } \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(t, \mu, \tilde{\omega}) &= \int_\tau^t D_x^2 F(r, \tilde{\omega}(r))dr. \end{aligned}$$

Therefore, the functional f also has continuous SVDs up to order 2.

- (iii) Let $F \in C^1(\mathbb{R}^d)$ such that $|DF(x)| \leq C(1 + |x|)$ for some $C \geq 0$. For fixed $t_0 \in (0, T)$, consider $\Phi(T, \mu) := \mathbb{E}^\mu[F(W(t_0))]$. Then the SVD at μ_t is $\partial_{\mu_t} \Phi(T, \mu, \tilde{\omega}) := D_x F(\tilde{\omega}(t_0))1_{[0, t_0]}(t)$, which may not be continuous in $t \in [0, T]$. However, for any $\mu \in \mathcal{P}_2^D$, consider

$$(36) \quad \Phi_\varepsilon(T, \mu) := \mathbb{E}^\mu\left[\int_0^T \rho_\varepsilon(t_0 - s)F(W(s))ds\right],$$

with ρ_ε being a standard mollifier on \mathbb{R} . Then by applying the dominated convergence theorem, we have

$$(37) \quad \lim_{\varepsilon \rightarrow 0} \Phi_\varepsilon(T, \mu) = \Phi(T, \mu), \quad \forall \mu \in \mathcal{P}_2^C.$$

Moreover, according to (ii), $\Phi_\varepsilon(T, \mu)$ has continuous SVDs. Therefore, we may approximate functionals with non-smooth SVDs by smooth ones. See Example 4.11 for further application in path-dependent master equations.

Example 2.12. We consider non-anticipative functionals $f \in \mathcal{D}$ by combining Example 2.2 and Example 2.11. For simplicity take $d = 1$. Suppose that $F \in C_b^{1,2}([0, T] \times \mathbb{R}^5)$ and $f_1, f_2, f_3, f_5 \in$

$C_b^2(\mathbb{R})$. $f_4 \in C_b^2(\mathbb{R}^2)$. Consider the following functional

$$f(t, \omega, \mu) := F\left(t, \omega(t), \int_0^t f_1(\omega(r))dr, \mathbb{E}^\mu[f_2(W(t))], \mathbb{E}^\mu\left[\int_0^t f_3(W(r))dr\right], \right. \\ \left. \mathbb{E}^\mu[f_4(W(t), \int_0^t f_5(W(r))dr)\right], \quad \forall (t, \omega, \mu) \in \hat{\mathbb{D}}_{T,d}.$$

Then we check that f has continuous horizontal derivatives and twice continuous SVDs in ω_t and μ_t . Indeed, for any $\tau \leq t$,

$$\partial_t f(t, \omega, \mu) = \partial_t F(t, x) + \partial_{x_2} F(t, x) f_1(\omega(t)) + \partial_{x_4} F(t, x) \mathbb{E}^\mu[f_3(W(t))] \\ + \partial_{x_5} F(t, x) \mathbb{E}^\mu[\partial_{y_2} f_4(Y) f_5(W(t))],$$

$$\partial_{\omega_\tau} f(t, \omega, \mu) = \partial_{x_1} F(t, x) + \partial_{x_2} F(t, x) \int_\tau^t f_1'(\omega(r))dr,$$

$$\partial_{\omega_\tau}^2 f(t, \omega, \mu) = \partial_{x_1}^2 F(t, x) + \partial_{x_2}^2 F(t, x) \left(\int_\tau^t f_1'(\omega(r))dr \right)^2 + \partial_{x_2} F(t, x) \int_\tau^t f_1^{(2)}(\omega(r))dr,$$

$$\partial_{\mu_\tau} f(t, \omega, \mu, \tilde{\omega}) = \partial_{x_3} F(t, x) f_2'(\tilde{\omega}(t)) + \partial_{x_4} F(t, x) \int_\tau^t f_3'(\tilde{\omega}(r))dr \\ + \partial_{x_5} F(t, x) \left[\partial_{y_1} f_4(\tilde{y}) + \partial_{y_2} f_4(\tilde{y}) \int_\tau^t f_5'(\tilde{\omega}(r))dr \right], \quad \text{and}$$

$$\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(t, \omega, \mu, \tilde{\omega}) = \partial_{x_3} F(t, x) f_2^{(2)}(\tilde{\omega}(t)) + \partial_{x_4} F(t, x) \int_\tau^t f_3^{(2)}(\tilde{\omega}(r))dr + \partial_{x_5} F(t, x) \left[\partial_{y_1}^2 f_4(\tilde{y}), \right. \\ \left. + 2\partial_{y_2} \partial_{y_1} f_4(\tilde{y}) \int_\tau^t f_5'(\tilde{\omega}(r))dr + \partial_{y_2}^2 f_4(\tilde{y}) \left(\int_\tau^t f_5'(\tilde{\omega}(r))dr \right)^2 \right],$$

where

$$(t, x) = \left(t, \omega(t), \int_0^t f_1(\omega(r))dr, \mathbb{E}^\mu[f_2(W(t))], \mathbb{E}^\mu \left[\int_0^t f_3(W(r))dr \right], \mathbb{E}^\mu \left[f_4(W(t), \int_0^t f_5(W(r))dr) \right] \right),$$

$$Y = \left(W(t), \int_0^t f_5(W(r))dr \right), \quad \text{and} \quad \tilde{y} = \left(\tilde{\omega}(t), \int_0^t f_5(\tilde{\omega}(r))dr \right).$$

In the following, for any $f \in \mathcal{D}$, we use generic notations $(\partial_\omega f, \partial_\omega^2 f)$ ($(\partial_{\omega_\tau} f, \partial_{\omega_\tau}^2 f)$, resp.) to denote the vertical derivative (SVD, resp.) in path, and $(\partial_\mu f, \partial_{\tilde{\omega}} \partial_\mu f)$ ($(\partial_{\mu_\tau} f, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f)$, resp.) to denote the vertical derivative (SVD, resp.) in measure if there is no confusion.

Definition 2.13. Denote by $\mathcal{C}(\hat{\mathbb{D}}_{T,d})$ (or \mathcal{C} when there is no confusion), the subspace of \mathcal{D} which consists of all non-anticipative and continuous functionals. Furthermore,

(i) $\mathcal{C}_s^{1,1,1}$ ($\mathcal{C}_s^{1,1,1}$, resp.) is the subset of \mathcal{C} whose element is continuously horizontally differentiable, (strongly, resp.) vertically differentiable w.r.t. both path and measure, with all derivatives being continuous;

(ii) $\mathcal{C}_s^{1,2,1}$ ($\mathcal{C}_s^{1,2,1}$, resp.) is the subset of $\mathcal{C}^{1,1,1}$ ($\mathcal{C}_s^{1,1,1}$, resp.) whose element's derivative $\partial_\omega f(t, \cdot, \mu, \tilde{\omega})$ ($\partial_{\omega_\tau} f(t, \cdot, \mu, \tilde{\omega})$, $\tau \leq t$, resp.), $(t, \omega, \mu, \tilde{\omega}) \in \hat{\mathbb{D}}_{T,d} \times \mathbb{D}_{T,d}$, is further vertically differentiable (strongly vertically differentiable at (τ, t, ω) , resp.), with all derivatives being continuous;

(iii) $\mathcal{C}^{1,2,1,1}$ ($\mathcal{C}_s^{1,2,1,1}$, resp.) is the subset of $\mathcal{C}^{1,2,1}$ ($\mathcal{C}_s^{1,2,1}$, resp.) whose element's derivative functional $\partial_\mu f(t, \omega, \mu, \cdot)$ ($\partial_{\mu_\tau} f(t, \omega, \mu, \cdot)$, $\tau \leq t$, resp.), $(t, \omega, \mu, \tilde{\omega}) \in \hat{\mathbb{D}}_{T,d} \times \mathbb{D}_{T,d}$, is further vertically differentiable (strongly vertically differentiable at $(\tau, t, \tilde{\omega})$, resp.), with all derivatives being continuous.

Moreover, denote by $\mathcal{C}_p^{1,1,1}$ the subset of $\mathcal{C}^{1,1,1}$ such that the functional and all its first order derivatives have at most polynomial growth in the path variable: there exists $k \in \mathbb{Z}^+$, such that for $\phi = f, \partial_t f, \partial_\omega f$, $\psi = \partial_\mu f$ and any $K > 0$,

$$(38) \quad \begin{aligned} |\phi(t, \omega, \mu)| &\leq C_K(1 + \|\omega_t\|^k), & |\psi(t, \omega, \mu, \tilde{\omega})| &\leq C_K(1 + \|\omega_t\|^k + \|\tilde{\omega}_t\|^k), \\ \forall(t, \omega, \mu, \tilde{\omega}) &\in \hat{\mathbb{D}}_{T,d} \times \mathbb{D}_{T,d} \text{ such that } \|\mu_t\| &\leq K, \end{aligned}$$

for a constant C_K depending only on K . Notations such as \mathcal{C}_p , $\mathcal{C}_{s,p}^{1,1,1}$, $\mathcal{C}^{0,1,1}$ and $\mathcal{C}_p^{1,2,1,1}$ are defined similarly.

Remark 2.14. Assume that $f \in \mathcal{D}$ is non-anticipative and has a state-dependent structure: $f(t, \omega, \mu) = \tilde{f}(t, \omega(t), \mu(t))$ for some function \tilde{f} defined on $[0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$. Then the horizontal differentiability and strongly vertical differentiability of f is reduced to the differentiability of \tilde{f} on $[0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d)$. Moreover,

$$(39) \quad \partial_t f(t, \omega, \mu) = \partial_t \tilde{f}(t, \omega(t), \mu(t)), \quad \partial_\omega f(t, \omega, \mu) = D_x f(t, \omega(t), \mu(t)), \quad \text{and}$$

$$(40) \quad \partial_{\mu_\tau} f(t, \omega, \mu, \tilde{\omega}) = \partial_\nu \tilde{f}(t, \omega(t), \mu(t), \tilde{\omega}(t)), \quad \forall(t, \omega, \mu) \in [0, T] \times \mathbb{D}_{T,d} \times \mathcal{P}_2^D, \quad \tau \leq t,$$

where $\partial_\nu \tilde{f}$ is the Lions' derivative (see e.g. [36]).

2.3. Itô-Dupire formula. Suppose that (a, b) is a bounded progressively measurable process on (Ω, \mathcal{F}, P) with values in $\mathbb{R}^m \times \mathbb{R}^{m \times d}$. For any $(t, \gamma) \in [0, T] \times \mathbb{D}_{T,d}$, X is the solution of SDE

$$(41) \quad \begin{cases} dX(r) = a(r)dr + b(r)dB(r), \\ X_t = \gamma_t, \quad r \geq t. \end{cases}$$

$(\Omega', \mathcal{F}', P')$ is an atomless probability space with a k -dimensional Brownian motion B' and (c, d) is a bounded progressively measurable process on $(\Omega', \mathcal{F}', P')$ with values in $\mathbb{R}^n \times \mathbb{R}^{n \times k}$. Given $\eta \in (\mathbb{M}_2^D)'$, let X' defined by SDE

$$(42) \quad \begin{cases} dX'(r) = c(r)dr + d(r)dB'(r), \\ X'_t = \eta_t, \quad r \geq t. \end{cases}$$

Moreover, let $(\tilde{X}', \tilde{c}, \tilde{d}, \tilde{B}', \tilde{\eta})$ be an independent copy of (X', c, d, B', η) , which means that $(\tilde{X}', \tilde{c}, \tilde{d}, \tilde{B}', \tilde{\eta})$ is defined in an independent probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ from (Ω, \mathcal{F}, P) and $(\Omega', \mathcal{F}', P')$, and it has the same law as (X', c, d, B', η) . Then we have the following Itô-Dupire formula.

Theorem 2.15. For any fixed $(t, \gamma, \eta) \in [0, T] \times \mathbb{D}_{T,d} \times (\mathbb{M}_2^D)'$, X and X' are diffusion processes defined by (41) and (42) respectively. Suppose that $f \in \mathcal{C}_p^{1,2,1,1}(\hat{\mathbb{D}}_{T,d})$, and then we have

$$\begin{aligned}
& f(s, X, \mathcal{L}_{X'}) - f(t, \gamma, \mathcal{L}_\eta) \\
&= \int_t^s \partial_r f(r, X, \mathcal{L}_{X'}) dr + \int_t^s \partial_\omega f(r, X, \mathcal{L}_{X'}) dX(r) \\
(43) \quad &+ \frac{1}{2} \int_t^s \text{Tr} [\partial_\omega^2 f(r, X_r, \mathcal{L}_{X'}) d\langle X \rangle(r)] + \mathbb{E}^{\tilde{P}'} \left[\int_t^s \partial_\mu f(r, X, \mathcal{L}_{X'}, \tilde{X}') d\tilde{X}'(r) \right] \\
&+ \frac{1}{2} \mathbb{E}^{\tilde{P}'} \int_t^s \text{Tr} [\partial_{\tilde{\omega}} \partial_\mu f(r, X, \mathcal{L}_{X'}, \tilde{X}') \tilde{d}(r) \tilde{d}(r)^T] dr, \quad \forall s \geq t.
\end{aligned}$$

Proof. Without loss of generality, assume $d = k = m = n = 1$ and $s = T$. Since both sides of identity (43) depend on (X', c, d, η) through its law, we assume that $(\Omega', \mathcal{F}', P')$ is independent from (Ω, \mathcal{F}, P) for simplicity of notations. Consider the following discretization of X and X' : for any $n \geq 1$, take $t = t_0 < t_1 < \dots < t_n = T$ as any partition of $[0, T]$ with vanishing modulus δ_n . Define càdlàg processes X^n, X'^n with $X_t^n = \gamma_t, X_t'^n = \eta_t$ by

$$\begin{aligned}
X^n(r) &:= \sum_{i=0}^{n-1} X(t_i) 1_{[t_i, t_{i+1})}(r) + X(T) 1_{\{T\}}(r), \\
X'^n(r) &:= \sum_{i=0}^{n-1} X'(t_i) 1_{[t_i, t_{i+1})}(r) + X'(T) 1_{\{T\}}(r), \quad r \geq t.
\end{aligned}$$

Since (a, b, c, d) is bounded, we see that for any $r \in [0, T]$,

$$(44) \quad \mathbb{E} \|X^n\|_{\mathbb{S}^p}^p \leq \mathbb{E} \|X\|_{\mathbb{S}^p}^p < \infty, \quad \lim_{n \rightarrow \infty} \|X_{t_i}^n - X_r\| = 0, \quad P\text{-a.s.},$$

$$(45) \quad \|\mathcal{L}_{X'^n}\|^2 = \mathbb{E} \|X'^n\|_{\mathbb{S}^2}^2 \leq \mathbb{E} \|X'\|_{\mathbb{S}^2}^2 < \infty, \quad \lim_{n \rightarrow \infty} \|X_{t_i}^n - X'_r\| = 0, \quad P'\text{-a.s.},$$

where i above satisfies $r \in [t_i, t_{i+1})$. It follows from (45) that

$$(46) \quad \lim_{n \rightarrow \infty} W_2(\mathcal{L}_{X'^n}, \mathcal{L}_{X'_r}) = 0.$$

Then we have

$$\begin{aligned}
& f(T, X_T^n, \mathcal{L}_{X'^n}) - f(t, \gamma_t, \mathcal{L}_{\eta_t}) \\
&= \sum_{i=0}^{n-1} [f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{(X'^n)_{t_{i+1}}}) - f(t_i, X_{t_i}^n, \mathcal{L}_{(X'^n)_{t_i}})] \\
(47) \quad &= \sum_{i=0}^{n-1} \left[(f(t_{i+1}, X_{t_i}^n, \mathcal{L}_{X'^n}) - f(t_i, X_{t_i}^n, \mathcal{L}_{X'^n})) + (f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X'^n}) \right. \\
&\quad \left. - f(t_{i+1}, X_{t_i}^n, \mathcal{L}_{X'^n})) + (f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X'^n}) - f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X'^n})) \right].
\end{aligned}$$

Since

$$\begin{aligned}
(48) \quad f(t_{i+1}, X_{t_i}^n, \mathcal{L}_{X'^n}) - f(t_i, X_{t_i}^n, \mathcal{L}_{X'^n}) &= \int_{t_i}^{t_{i+1}} \partial_r f(r, X_{t_i}^n, \mathcal{L}_{X'^n}) dr \\
&= \int_t^T \partial_r f(r, X_{t_i}^n, \mathcal{L}_{X'^n}) 1_{[t_i, t_{i+1})}(r) dr,
\end{aligned}$$

in view of inequalities (44) and (46), applying the dominated convergence theorem and passing to the limit for a subsequence, we have

$$(49) \quad \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \left(f(t_{i+1}, X_{t_i}^n, \mathcal{L}_{X_{t_i}^{\prime n}}) - f(t_i, X_{t_i}^n, \mathcal{L}_{X_{t_i}^{\prime n}}) \right) = \int_t^T \partial_r f(r, X, \mathcal{L}_{X'}) dr, \quad P\text{-a.s.}$$

For the second term on the right hand side of (47), since $f \in \mathcal{C}_p^{1,2,1,1}$, we have that $\phi_i(\theta) := f(t_{i+1}, X_{t_i}^n + \theta 1_{[t_{i+1}, T]}, \mathcal{L}_{X_{t_i}^{\prime n}})$ is twice continuously differentiable in θ , and moreover,

$$(50) \quad \phi_i'(\theta) = \partial_\omega f(t_{i+1}, X_{t_i}^n + \theta 1_{[t_{i+1}, T]}, \mathcal{L}_{X_{t_i}^{\prime n}}), \quad \phi_i''(\theta) = \partial_\omega^2 f(t_{i+1}, X_{t_i}^n + \theta 1_{[t_{i+1}, T]}, \mathcal{L}_{X_{t_i}^{\prime n}}).$$

In the following, we will write $X_i := X(t_i) \equiv X^n(t_i)$ and $\delta X_i := X_{i+1} - X_i$. Similar notations such as X_i' are self-explained. Note that $X_{t_{i+1}}^n = X_{t_i}^n + (X_{i+1} - X_i) 1_{[t_{i+1}, T]}$. Using the Itô formula to $\phi(X(r) - X_i)$ on $r \in [t_i, t_{i+1}]$, we have

$$(51) \quad \begin{aligned} & f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_{i+1}}^{\prime n}}) - f(t_{i+1}, X_{t_i}^n, \mathcal{L}_{X_{t_i}^{\prime n}}) \\ &= \int_{t_i}^{t_{i+1}} \partial_\omega f(t_{i+1}, X_{t_i}^n + (X(r) - X_i) 1_{[t_{i+1}, T]}, \mathcal{L}_{X_{t_i}^{\prime n}}) dX(r) \\ & \quad + \frac{1}{2} \int_{t_i}^{t_{i+1}} \partial_\omega^2 f(t_{i+1}, X_{t_i}^n + (X(r) - X_i) 1_{[t_{i+1}, T]}, \mathcal{L}_{X_{t_i}^{\prime n}}) d\langle X \rangle(r). \end{aligned}$$

Since $\|X_{t_i}^n + (X(r) - X_i) 1_{[t_{i+1}, T]} - X_r\| \rightarrow 0$, P -a.s. for any $r \in [t_i, t_{i+1})$, according to inequality (51), passing to the limit in a subsequence, we have

$$(52) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \left(f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_{i+1}}^{\prime n}}) - f(t_{i+1}, X_{t_i}^n, \mathcal{L}_{X_{t_i}^{\prime n}}) \right) \\ &= \int_t^T \partial_\omega f(r, X, \mathcal{L}_{X'}) dX(r) + \frac{1}{2} \int_t^T \partial_\omega^2 f(r, X, \mathcal{L}_{X'}) d\langle X \rangle(r), \quad P\text{-a.s.} \end{aligned}$$

For the last term in the decomposition (47), we have

$$\begin{aligned} & f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_{i+1}}^{\prime n}}) - f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^{\prime n}}) \\ &= \int_0^1 \mathbb{E}' \left[\partial_\mu f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^{\prime n} + \theta(\delta X_i') 1_{[t_{i+1}, T]}}), X_{t_i}^n + \theta(\delta X_i') 1_{[t_{i+1}, T]} \right] (\delta X_i') d\theta \\ &= \int_0^1 \mathbb{E}' \left[\partial_\mu f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^{\prime n} + \theta(\delta X_i') 1_{[t_{i+1}, T]}}), X_{t_i}^n \right] (\delta X_i') d\theta \\ & \quad + \int_0^1 \int_0^1 \mathbb{E}' \left[\partial_\omega \partial_\mu f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^{\prime n} + \theta(\delta X_i') 1_{[t_{i+1}, T]}}), X_{t_i}^n + \lambda \theta(\delta X_i') 1_{[t_{i+1}, T]} \right] \theta (\delta X_i')^2 d\theta d\lambda. \end{aligned}$$

Since $\|X_{t_i}^n + \theta(\delta X_i') 1_{[t_{i+1}, T]} - X_r'\| \rightarrow 0$, P' -a.s. for any $r \in [0, T]$ with $r \in [t_i, t_{i+1}]$, we have

$$\lim_{n \rightarrow \infty} W_2(\mathcal{L}_{X_{t_i}^{\prime n} + \theta(\delta X_i') 1_{[t_{i+1}, T]}}, \mathcal{L}_{X_r'}) = 0.$$

In view of (44), (46) and the dominated convergence theorem, we have

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \int_0^1 \left[\partial_\mu f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^n + \theta(\delta X_i') 1_{[t_{i+1}, T]}}}, X_{t_i}^n)(\delta X_i') \right] d\theta \\ &= \int_t^T \partial_\mu f(r, X^{\gamma_t}, \mathcal{L}_{X'}, X') dX'(r), \quad P \times P' \text{-a.s.} \end{aligned}$$

Then, according to Fubini's theorem, we have

$$(53) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \mathbb{E}' \int_0^1 \left[\partial_\mu f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^n + \theta(\delta X_i') 1_{[t_{i+1}, T]}}}, X_{t_i}^n)(\delta X_i') \right] d\theta \\ &= \mathbb{E}' \left[\int_t^T \partial_\mu f(r, X, \mathcal{L}_{X'}, X') dX'(r) \right], \quad P \text{-a.s.} \end{aligned}$$

By a similar argument as above, we have

$$(54) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_0^1 \int_0^1 \mathbb{E}' \left[\partial_{\tilde{\omega}} \partial_\mu f(t_{i+1}, X_{t_{i+1}}^n, \mathcal{L}_{X_{t_i}^n + \theta(\delta X_i') 1_{[t_{i+1}, T]}}}, X_{t_i}^n + \lambda \theta(\delta X_i') 1_{[t_{i+1}, T]}) \theta(\delta X_i')^2 \right] d\theta d\lambda \\ &= \mathbb{E}' \left[\int_t^T \partial_{\tilde{\omega}} \partial_\mu f(r, X, \mathcal{L}_{X'}, X') dr \right], \quad P \text{-a.s.} \end{aligned}$$

In view of (49), (52), (53) and (54), taking $n \rightarrow \infty$ in (47), we obtain the desired identity. \square

Note that $(\omega_s)_\tau = \omega_s$ and $(\mu_s)_\tau = \mu_s$ for any $\tau \geq s$. In particular, if the non-anticipative functional f is strongly vertically differentiable, we have the following partial Itô-Dupire formula.

Corollary 2.16. *Suppose that (X, X') is defined as in Theorem 2.15, and $f \in \mathcal{C}_{s,p}^{0,2,1,1}(\hat{\mathbb{D}}_{T,d})$. Then we have that for any $t \leq s \leq v \leq T$,*

$$\begin{aligned} & f(v, X_s, \mathcal{L}_{X'_s}) - f(v, \gamma_t, \mathcal{L}_{\eta_t}) \\ &= \int_t^s \partial_{\omega_r} f(v, X_r, \mathcal{L}_{X'_r}) dX(r) + \frac{1}{2} \int_t^s \text{Tr} [\partial_{\omega_r}^2 f(v, X_r, \mathcal{L}_{X'_r}) d\langle X \rangle(r)] \\ &+ \mathbb{E}^{\tilde{P}'} \left[\int_t^s \partial_{\mu_r} f(v, X_r, \mathcal{L}_{X'_r}, \tilde{X}'_r) d\tilde{X}'(r) \right] + \frac{1}{2} \mathbb{E}^{\tilde{P}'} \int_t^s \text{Tr} [\partial_{\tilde{\omega}_r} \partial_{\mu_r} f(v, X_r, \mathcal{L}_{X'_r}, \tilde{X}'_r) \tilde{d}(r) \tilde{d}(r)^T] dr. \end{aligned}$$

Proof. Without loss of generality, assume $v = T$. For any $r \in [t, s]$, let

$$(54) \quad \tilde{f}(r, \omega, \mu) := f(T, \omega_r, \mu_r).$$

Obviously, \tilde{f} is non-anticipative, and moreover, we have that for any $h \geq 0$,

$$\tilde{f}(r+h, \omega_r, \mu_r) = f(T, (\omega_r)_{r+h}, (\mu_r)_{r+h}) = f(T, \omega_r, \mu_r) = \tilde{f}(r, \omega_r, \mu_r),$$

which implies $\partial_r \tilde{f}(r, \omega_r, \mu_r) = 0$. Furthermore, it follows by definitions of vertical derivatives and strongly vertical derivatives that

$$\begin{aligned} \partial_\omega \tilde{f}(r, \omega, \mu) &= \partial_{\omega_r} f(T, \omega_r, \mu_r), & \partial_\omega^2 \tilde{f}(r, \omega, \mu) &= \partial_{\omega_r}^2 f(T, \omega_r, \mu_r), \\ \partial_\mu \tilde{f}(r, \omega, \mu, \tilde{\omega}) &= \partial_{\mu_r} f(T, \omega_r, \mu_r, \tilde{\omega}), & \text{and } \partial_{\tilde{\omega}} \partial_\mu \tilde{f}(r, \omega, \mu, \tilde{\omega}) &= \partial_{\tilde{\omega}_r} \partial_{\mu_r} f(T, \omega_r, \mu_r, \tilde{\omega}). \end{aligned}$$

Applying Theorem 2.15 to $\tilde{f}(r, X, \mathcal{L}_{X'})$ on $r \in [t, s]$, and we obtain the desired formula. \square

3. DIFFERENTIABILITY OF SOLUTIONS FOR PATH-DEPENDENT MEAN-FIELD BSDEs

In the following, for any process (X, Y, Z) on the probability space (Ω, \mathcal{F}, P) , we denote by $(\tilde{X}, \tilde{Y}, \tilde{Z})$ an independent copy of (X, Y, Z) , which means that $(\tilde{X}, \tilde{Y}, \tilde{Z})$ is defined in an independent probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ and has the same law as (X, Y, Z) . Recall that B is a d -dimensional Brownian motion on (Ω, \mathcal{F}, P) . The following linear mean-field BSDEs and estimates are frequently used in subsequent discussions. Note that for a classical linear BSDE, the generator has a linear growth in Y , which implies the global well-posedness. Here in the mean-field case, things are different since the expected evolutionary equation is an ODE. For simplicity, we only address the one-dimensional case. Similar assertions in this section are still true in the multi-dimensional case.

Lemma 3.1. *Let $\xi \in L^2(\mathcal{F}_T)$ and $t \in [0, T)$. Suppose that $(\alpha, \beta) \in \mathbb{H}^2([t, T], \mathbb{R} \times \mathbb{R}^d)$ is bounded, $c \in \mathbb{H}^2([t, T], \mathbb{R}^k)$, and h is a real valued progressively measurable process such that $\int_t^T |h(r)| dr \in L^2(\mathcal{F}_T)$. For any $(r, x) \in [t, T] \times \mathbb{R}^k$, $g(\cdot, x) \in \mathbb{H}^2([t, T])$ and $g(r, \cdot)$ is uniformly Lipschitz continuous:*

$$\sup_{r \in [t, T]} |g(r, x) - g(r, y)| \leq L|x - y|, \quad \forall y \in \mathbb{R}^k, \quad P\text{-a.s.}$$

for a constant L . Then, the following linear mean-field BSDE

$$(55) \quad Y(s) = \xi + \int_s^T \left(\alpha(r)Y(r) + \beta(r)Z(r) + \tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{Y}(r)] + h(r) \right) dr - \int_s^T Z(r)dB(r), \quad s \in [t, T]$$

with (\tilde{c}, \tilde{Y}) being an independent copy of (c, Y) , has a unique solution $(Y, Z) \in \mathbb{S}^2([t, T]) \times \mathbb{H}^2([t, T], \mathbb{R}^d)$. Moreover, we have

$$(56) \quad \|(Y, Z)\|_{\mathbb{S}^2 \times \mathbb{H}^2}^2 \leq C(\|\xi\|_{L^2}^2 + \|\int_t^T |h(r)| dr\|_{L^2}^2) e^{C(\|c\|_{\mathbb{H}^2} + \|g(\cdot, 0)\|_{\mathbb{H}^2})}$$

for a constant C depending on the bound of α, β and L . In particular, if g is uniformly bounded, we have

$$(57) \quad \|(Y, Z)\|_{\mathbb{S}^2 \times \mathbb{H}^2}^2 \leq C(\|\xi\|_{L^2}^2 + \|\int_t^T |h(r)| dr\|_{L^2}^2).$$

Remark 3.2. *Since neither $g(t, x)$ nor $g(r, c(r))$ is bounded or uniformly integrable for any $c(r) \in \mathbb{H}^2([t, T], \mathbb{R}^k)$, the well-posedness of the mean-field BSDE is not an immediate consequence of existing works such as [8].*

Proof. For any $y \in \mathbb{H}^2$, consider the following classical linear BSDE

$$(58) \quad Y(s) = \xi + \int_s^T \left(\alpha(r)Y(r) + \beta(r)Z(r) + \tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{y}(r)] + h(r) \right) dr - \int_s^T Z(r)dB(r),$$

where (\tilde{c}, \tilde{y}) is an independent copy of (c, y) . To prove that it is well-posed on $[t, T]$, we only need to show

$$(59) \quad \mathbb{E} \left[\int_t^T \left| \tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{y}(r)] \right| dr \right]^2 < \infty.$$

Indeed, by the uniformly Lipschitz continuity of g , we have

$$\begin{aligned}
(60) \quad & \mathbb{E} \left| \int_t^T \tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{y}(r)] dr \right|^2 \leq C \mathbb{E} \left[\int_t^T |\tilde{\mathbb{E}}[g(r, 0)\tilde{y}(r)] dr + \tilde{\mathbb{E}}[\tilde{c}(r)\tilde{y}(r)] dr \right]^2 \\
& \leq C \mathbb{E} \left[\int_t^T |g(r, 0)\tilde{\mathbb{E}}[\tilde{y}(r)]| dr \right]^2 + C \left[\int_t^T \tilde{\mathbb{E}}[|\tilde{c}(r)\tilde{y}(r)|] dr \right]^2 \\
& \leq C \left[(\mathbb{E} \int_t^T |g(r, 0)|^2 dr) (\int_t^T [\tilde{\mathbb{E}}|\tilde{y}(r)|]^2 dr) + (\int_t^T \tilde{\mathbb{E}}|\tilde{c}(r)|^2 dr) (\int_t^T \tilde{\mathbb{E}}|\tilde{y}(r)|^2 dr) \right] \\
& \leq C \left[\|g(\cdot, 0)\|_{\mathbb{H}^2}^2 \|y\|_{\mathbb{H}^2}^2 + \|c\|_{\mathbb{H}^2}^2 \|y\|_{\mathbb{H}^2}^2 \right] \leq C (\|g(\cdot, 0)\|_{\mathbb{H}^2}^2 + \|c\|_{\mathbb{H}^2}^2) \|y\|_{\mathbb{H}^2}^2,
\end{aligned}$$

where we apply the Hölder inequality to integrals in forms of \int_t^T and $\int_t^T \tilde{\mathbb{E}}$ respectively in the third inequality above. Then for any $y \in \mathbb{H}^2$, there exists a unique solution $(Y, Z) \in \mathbb{H}^2 \times \mathbb{H}^2$ of BSDE (58). The mapping $y \rightarrow Y$, denoted by Φ , is a transformation on \mathbb{H}^2 , and will be shown to be a contraction under the following equivalent norm on \mathbb{H}^2

$$(61) \quad \|Y\|^2 := \mathbb{E} \int_t^T e^{As - \int_s^T (\|g(r, 0)\|_{L^2}^2 + \|c(r)\|_{L^2}^2) dr} |Y(s)|^2 ds$$

with A being a constant to be determined later. Take any $y^{(1)}, y^{(2)} \in \mathbb{H}^2$ and denote the corresponding solutions of classical BSDE (58) by $(Y^{(1)}, Z^{(1)}), (Y^{(2)}, Z^{(2)})$. Set $(\Delta Y, \Delta Z) := (Y^{(1)} - Y^{(2)}, Z^{(1)} - Z^{(2)})$, $\Delta y := y^{(1)} - y^{(2)}$, and $f(r) := \|g(r, 0)\|_{L^2}^2 + \|c(r)\|_{L^2}^2$. Apply Itô's formula to $e^{As - \int_s^T f(r) dr} |\Delta Y(s)|^2$ on $s \in [t, T]$, and we have

$$\begin{aligned}
& -e^{At - \int_t^T f(r) dr} |\Delta Y(t)|^2 = \int_t^T (A + f(s)) e^{As - \int_s^T f(r) dr} |\Delta Y(s)|^2 ds \\
& \quad + 2 \int_t^T e^{As - \int_s^T f(r) dr} \Delta Y d(\Delta Y) + \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Z|^2 ds.
\end{aligned}$$

Therefore,

$$\begin{aligned}
& e^{At - \int_t^T f(r) dr} |\Delta Y(t)|^2 + \int_t^T (A + f(s)) e^{As - \int_s^T f(r) dr} |\Delta Y(s)|^2 ds + \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Z(s)|^2 ds \\
& = 2 \int_t^T e^{As - \int_s^T f(r) dr} \Delta Y [\alpha \Delta Y + \beta \Delta Z + \tilde{\mathbb{E}}[g(s, \tilde{c}(s))\Delta \tilde{y}]] ds - 2 \int_t^T e^{As - \int_s^T f(r) dr} \Delta Y \Delta Z dW(s) \\
& \leq C \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Y|^2 ds + C \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Y|^2 ds + \frac{1}{2} \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Z|^2 ds \\
& \quad + 2 \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Y| \|g(s, 0)\| \|\Delta \tilde{y}\|_{L^2} ds + 2 \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Y| \|c\|_{L^2} \|\Delta \tilde{y}\|_{L^2} ds \\
& \quad - 2 \int_t^T e^{As - \int_s^T f(r) dr} \Delta Y \Delta Z dB(s) \\
& \leq C \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Y|^2 ds + \frac{1}{2} \int_t^T e^{As - \int_s^T f(r) dr} |\Delta Z|^2 ds - 2 \int_t^T e^{As - \int_s^T f(r) dr} \Delta Y \Delta Z dB(s) \\
& \quad + 2 \int_t^T e^{As - \int_s^T f(r) dr} (|\Delta Y| \|g(s, 0)\| \|\Delta \tilde{y}\|_{L^2}) ds + \int_t^T e^{As - \int_s^T f(r) dr} (|\Delta Y|^2 \|c\|_{L^2}^2 + \|\Delta \tilde{y}\|_{L^2}^2) ds.
\end{aligned}$$

Taking expectation on both sides of the above inequality, we have

$$\begin{aligned}
& \int_t^T (A - C + f(s)) e^{As - \int_s^T f(r) dr} \|\Delta Y(s)\|_{L^2}^2 ds \\
& \leq 2 \int_t^T e^{As - \int_s^T f(r) dr} \mathbb{E}[\|\Delta Y\|g(s, 0)\|] \|\Delta \tilde{y}\|_{L^2} ds + \int_t^T e^{As - \int_s^T f(r) dr} (\|\Delta Y\|_{L^2}^2 \|c\|_{L^2}^2 + \|\Delta \tilde{y}\|_{L^2}^2) ds \\
& \leq \int_t^T e^{As - \int_s^T f(r) dr} \left[\left(\mathbb{E}[\|\Delta Y\|g(s, 0)\|] \right)^2 + \|\Delta \tilde{y}\|_{L^2}^2 \right] ds + \int_t^T e^{As - \int_s^T f(r) dr} (\|\Delta Y\|_{L^2}^2 \|c\|_{L^2}^2 + \|\Delta \tilde{y}\|_{L^2}^2) ds \\
& \leq \int_t^T e^{As - \int_s^T f(r) dr} \left(\|g(s, 0)\|_{L^2}^2 + \|c\|_{L^2}^2 \right) \|\Delta Y\|_{L^2}^2 ds + \int_t^T e^{As - \int_s^T f(r) dr} \|\Delta \tilde{y}\|_{L^2}^2 ds.
\end{aligned}$$

Therefore, choosing a sufficiently large number A such that $A - C > 1$, we obtain a contraction and then the well-posedness of (55).

Now BSDE (55) can be written as the following classical BSDE

$$(62) \quad Y(s) = \xi + \int_s^T \left(\alpha(r)Y(r) + \beta(r)Z(r) + h'(r) \right) dr - \int_s^T Z(r) dB(r),$$

with $h'(r) = \tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{Y}(r)] + h(r)$. Thus it is standard that

$$\begin{aligned}
(63) \quad & \|Y\|_{\mathbb{S}^2}^2 + \|Z\|_{\mathbb{H}^2}^2 \leq C(\|\xi\|_{L^2}^2 + \|\int_t^T |h'(r)| dr\|_{L^2}^2) \\
& \leq C(\|\xi\|_{L^2}^2 + \|\int_t^T |h(r)| dr\|_{L^2}^2 + \|\int_t^T |\tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{Y}(r)]| dr\|_{L^2}^2).
\end{aligned}$$

Furthermore, similar to the proof of inequality (60), we have

$$(64) \quad \|\int_t^T |\tilde{\mathbb{E}}[g(r, \tilde{c}(r))\tilde{Y}(r)]| dr\|_{L^2}^2 \leq C[\int_t^T \|g(r, 0)\|_{L^2}^2 \|Y\|_{\mathbb{S}^2, [t, r]}^2 dr + \int_t^T \|c(r)\|_{L^2}^2 \|Y\|_{\mathbb{S}^2, [t, r]}^2 dr].$$

Then, using Gronwall's inequality, we obtain the desired estimate (56). \square

We consider the following path-dependent master equation

$$(65) \quad \begin{cases} \partial_t u(t, \gamma, \mu) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \gamma, \mu) \sigma_1(t) \sigma_1(t)^T] + \partial_\omega u(t, \gamma, \mu) b_1(t) \\ \quad + \frac{1}{2} \text{Tr} [\mathbb{E}^P [\partial_\omega \partial_\mu u(t, \gamma, \mu, \eta)] \sigma_2(t) \sigma_2(t)^T] + \mathbb{E}^P [\partial_\mu u(t, \gamma, \mu, \eta)] b_2(t) \\ \quad + f(t, \gamma, u(t, \gamma, \mu), \sigma_1(t) \partial_\omega u(t, \gamma, \mu), \mu, \mathcal{L}_{u(t, \gamma, \mu)}) = 0, \\ u(T, \gamma, \mu) = \Phi(\gamma_T, \mu_T), \quad (t, \gamma, \mu) \in [0, T] \times \mathbb{C}_{T, d} \times \mathcal{P}_2^C, \end{cases}$$

where $(b_1, \sigma_1, b_2, \sigma_2)$ are continuous functions and $\eta \in \mathbb{M}_2^C$ with law μ . For simplicity, we take $(b_1, \sigma_1) = (b_2, \sigma_2) = (0, I)$ and refer to Remark 4.9 for the above form. In the following, we write $f(\omega_t, \mu_t) := f(t, \omega, \mu)$ for simplicity since f is non-anticipative. Moreover, for any $\gamma, \omega \in \mathbb{D}_{T, d}$, define $\omega^{\gamma t} \in \mathbb{D}_{T, d}$ as the following

$$(66) \quad \omega^{\gamma t}(\cdot) := \gamma_t(\cdot) + (\omega(\cdot) - \omega(t))1_{[t, T]}(\cdot).$$

To give a classical solution through FBSDEs, for any $(t, \eta) \in [0, T] \times \mathbb{M}_2^D$, we denote by (Y^{η_t}, Z^{η_t}) the solution of the following path-dependent mean-field BSDE

$$(67) \quad Y(s) = \Phi(B_T^{\eta_t}, \mathcal{L}_{B_T^{\eta_t}}) + \int_s^T f(B_r^{\eta_t}, Y(r), Z(r), \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y(r)}) dr - \int_s^T Z(r) dB(r), \quad s \in [t, T].$$

On the other hand, for any $\gamma \in \mathbb{D}_{T,d}$, let $(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t})$ solve the associated path-dependent BSDE

$$(68) \quad \mathcal{Y}(s) = \Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\gamma_t}}) + \int_s^T f(B_r^{\gamma_t}, \mathcal{Y}(r), \mathcal{Z}(r), \mathcal{L}_{B_r^{\gamma_t}}, \mathcal{L}_{Y^{\eta_t}(r)}) dr - \int_s^T \mathcal{Z}(r) dB(r), \quad s \in [t, T].$$

Definition 3.3. We write $\Phi \in \mathcal{C}_T(\hat{\mathbb{D}}_{T,d})$ (or \mathcal{C}_T if no confusion raised) if $\Phi : \mathbb{D}_{T,d} \times \mathcal{P}_2^D \rightarrow \mathbb{R}$ is continuous on $\mathbb{D}_{T,d} \times \mathcal{P}_2^D$. Furthermore, we write

(i) $\Phi \in \mathcal{C}_{T,lip}$ if it is uniformly Lipschitz continuous on $\mathbb{D}_{T,d} \times \mathcal{P}_2^D$:

$$|\Phi(\omega_T, \mu_T) - \Phi(\omega'_T, \mu'_T)| \leq C(\|\omega_T - \omega'_T\| + W_2(\mu_T, \mu'_T)), \quad \forall (\omega, \mu), (\omega', \mu') \in \mathbb{D}_{T,d} \times \mathcal{P}_2^D,$$

for some constant C ;

(ii) $\Phi \in \mathcal{C}_{T,lip}^{1,1}$ if $\Phi \in \mathcal{C}_{T,lip}$ and Φ is continuously strongly vertically differentiable in path and measure. Furthermore, for any $\tau \in [0, T)$, SVDs $\partial_{\omega_\tau} \Phi$ and $\partial_{\mu_\tau} \Phi$ are uniformly Lipschitz continuous in $(\omega, \mu) \in \mathbb{D}_{T,d} \times \mathcal{P}_2^D$ and $(\omega, \mu, \tilde{\omega}) \in \mathbb{D}_{T,d} \times \mathcal{P}_2^D \times \mathbb{D}_{T,d}$, respectively;

(iii) $\Phi \in \mathcal{C}_{T,lip}^{2,1,1}$ if $\Phi \in \mathcal{C}_{T,lip}^{1,1}$ and for any $(\tau, \omega, \mu, \tilde{\omega}) \in \hat{\mathbb{D}}_{T,d} \times \mathbb{D}_{T,d}$, its SVDs $\partial_{\omega_\tau} \Phi(\cdot, \mu_T)$ and $\partial_{\mu_\tau} \Phi(\omega_T, \mu_T, \cdot)$ are continuously strongly vertically differentiable at (τ, T, ω) and $(\tau, T, \tilde{\omega})$ respectively. Moreover, all second-order derivatives are uniformly Lipschitz continuous.

To obtain the well-posedness and estimates of BSDEs (67) and (68), we assume that

(H0) (i) The functional $\Phi \in \mathcal{C}_{T,lip}(\hat{\mathbb{D}}_{T,d})$; (ii) f is a non-anticipative continuous function on $[0, T] \times \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^D \times \mathcal{P}_2(\mathbb{R})$, and for any $(t, \omega, \mu) \in [0, T] \times \mathbb{D}_{T,d} \times \mathcal{P}_2^D$, $f(t, \omega, \cdot, \cdot, \mu, \cdot)$ is continuously differentiable on $\mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R})$. Moreover, for any $t \in [0, T]$, $f(t, \cdot, \cdot, \cdot, \cdot, \cdot)$ and $\partial_\nu f(t, \cdot, \cdot, \cdot, \cdot, \cdot)$ are uniformly Lipschitz continuous.

Note that under Assumption **(H0)**, the functional

$$(69) \quad \hat{f}(r, y, z, \nu) := f(B_r^{\eta_t}, y, z, \mathcal{L}_{B_r^{\eta_t}}, \nu), \quad (r, y, z, \nu) \in [t, T] \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}),$$

is uniformly Lipschitz continuous in $(y, z) \in \mathbb{R} \times \mathbb{R}^d$. According to [11, Theorem 4.23], BSDE (67) is well posed with $(Y^{\eta_t}, Z^{\eta_t}, \mathcal{L}_{Y^{\eta_t}}) \in \mathbb{S}^2 \times \mathbb{H}^2 \times \mathcal{P}_2(\mathbb{R})$. Then (68) is a well-defined classical BSDE with $(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t}) \in \mathbb{S}^p \times \mathbb{H}^p$ for any $p \geq 1$. In the following BSDEs, we write $\Theta_r^{\eta_t} := (B_r^{\eta_t}, Y^{\eta_t}(r), Z^{\eta_t}(r))$, $\Theta_r^{\gamma_t, \eta_t} := (B_r^{\gamma_t}, Y^{\gamma_t, \eta_t}(r), Z^{\gamma_t, \eta_t}(r))$, $\mathcal{L}_{\Theta_r^{\eta_t}} := (\mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(r)})$ and $(Y, Z) := (Y(t), Z(t))$ if no confusion is raised. Then we have the following basic estimates for BSDEs (67) and (68).

Lemma 3.4. Assume that (Φ, f) satisfies **(H0)**. For any $K > 0$ and $(\gamma, \eta), (\gamma', \eta') \in \mathbb{D}_{T,d} \times \mathbb{M}_2^D$ such that $\|\mathcal{L}_{\eta_t}\|, \|\mathcal{L}_{\eta'_t}\| \leq K$, we have

$$(70) \quad \|(Y^{\eta_t}, Z^{\eta_t})\|_{\mathbb{S}^2 \times \mathbb{H}^2} \leq C(1 + \|\eta_t\|_{\mathbb{S}^2}),$$

$$(71) \quad \|(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \leq C_p(1 + \|\gamma_t\| + \|\eta_t\|_{\mathbb{S}^2}), \quad \forall p \geq 1,$$

$$(72) \quad \|(Y^{\eta_t} - Y^{\eta'_t}, Z^{\eta_t} - Z^{\eta'_t})\|_{\mathbb{S}^2 \times \mathbb{H}^2} \leq C_K \|\eta_t - \eta'_t\|_{\mathbb{S}^2}, \quad \text{and}$$

$$(73) \quad \|(Y^{\gamma_t, \eta_t} - Y^{\gamma'_t, \eta'_t}, Z^{\gamma_t, \eta_t} - Z^{\gamma'_t, \eta'_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \leq C_{K,p}(\|\gamma_t - \gamma'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t})), \quad \forall p \geq 1,$$

where (C, C_p) does not depend on (γ, η) , and $(C_K, C_{K,p})$ does not depend on (γ, γ') .

Remark 3.5. According to inequality (73), $(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t})$ and $(Y^{\gamma_t, \eta'_t}, Z^{\gamma_t, \eta'_t})$ are indistinguishable if $\mathcal{L}_{\eta_t} = \mathcal{L}_{\eta'_t}$, which implies the following definition is well-posed

$$(74) \quad (Y^{\gamma_t, \mathcal{L}_{\eta_t}}, Z^{\gamma_t, \mathcal{L}_{\eta_t}}) := (Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t}).$$

The proof of Lemma 3.4 is rather standard with an application of Lemma 3.1, and is left in the appendix.

3.1. First-order differentiability. For any $(\gamma, \eta) \in [0, T] \times \mathbb{M}_2^D$, in the following, we consider the first order differentiability of $Y^{\gamma_t, \eta_t} = Y^{\gamma_t, \mathcal{L}_{\eta_t}}$ with respect to γ_t and \mathcal{L}_{η_t} . For the differentiability in γ_t , let

$$(75) \quad \begin{aligned} \hat{f}(\omega_s, y, z) &:= f(\omega_s, y, z, \mathcal{L}_{B_s^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(s)}), \\ \hat{\Phi}(\omega_T) &:= \Phi(\omega_T, \mathcal{L}_{B_T^{\eta_t}}), \quad \forall (s, \omega, y, z) \in [t, T] \times \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d, \end{aligned}$$

and then the solution $Y^{\gamma_t, \eta_t}(s)$ to equation (68) solves the following path-dependent BSDE

$$(76) \quad \hat{Y}(s) = \hat{\Phi}(B_T^{\eta_t}) + \int_s^T \hat{f}(B_r^{\eta_t}, \hat{Y}(r), \hat{Z}(r)) dr - \int_s^T \hat{Z}(r) dB(r).$$

Define $\hat{u}_{\eta_t}(t, \gamma) := Y^{\gamma_t, \eta_t}(t)$. If f and Φ are regular enough, according to [41, Theorem 4.5], $\hat{u}_{\eta_t}(t, \gamma)$ is twice vertically differentiable at (t, γ) , and moreover for any $s \geq t$,

$$(77) \quad \hat{u}_{\eta_t}(s, B^{\gamma_t}) = Y^{\gamma_t, \eta_t}(s), \quad \partial_{\gamma_t} \hat{u}_{\eta_t}(s, B^{\gamma_t}) = Z^{\gamma_t, \eta_t}(s).$$

Furthermore, $\hat{u}_{\eta_t}(t, \gamma)$ is the unique solution to the following semilinear PPDE

$$(78) \quad \begin{cases} \partial_t \hat{u}_{\eta_t}(t, \gamma) + \frac{1}{2} \text{Tr} [\partial_{\omega}^2 \hat{u}_{\eta_t}(t, \gamma)] + \hat{f}(\gamma_t, \hat{u}_{\eta_t}(t, \gamma), \partial_{\omega} \hat{u}_{\eta_t}(t, \gamma)) = 0, \\ \hat{u}_{\eta_t}(T, \gamma) = \hat{\Phi}(\gamma), \quad (t, \gamma) \in [0, T] \times \mathbb{C}_{T,d}. \end{cases}$$

In the following, we denote by $\partial_{(t, \omega, y, z, \mu, \nu, \omega_{\tau}, \mu_{\tau})} f$ the derivative vector

$$(\partial_t f, \partial_{\omega} f, \partial_y f, \partial_z f, \partial_{\mu} f, \partial_{\nu} f, \partial_{\omega_{\tau}} f, \partial_{\mu_{\tau}} f).$$

In this subsection, we assume that

- (H1)** (i) The functional $\Phi \in \mathcal{C}_{T, \text{lip}}^{1,1}(\hat{\mathbb{D}}_{T,d})$; (ii) f is a non-anticipative continuous function on $[0, T] \times \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^D \times \mathcal{P}_2(\mathbb{R})$, and for any $(t, \omega, \mu) \in [0, T] \times \mathbb{D}_{T,d} \times \mathcal{P}_2^D$, $f(t, \omega, \cdot, \cdot, \mu, \cdot)$ is differentiable on $\mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R})$ with bounded derivatives. For any $(y, z, \nu) \in \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R})$, $f(t, \omega, y, z, \cdot, \nu)$ is strongly vertically differentiable at μ_t and $f(t, \cdot, y, z, \mu, \nu)$ is strongly vertically differentiable at ω_t . Moreover, $\partial_{(y, z, \nu, \omega_{\tau}, \mu_{\tau})} f$ is continuous, and for any $\tau \leq t$, $(I, \partial_{(y, z, \nu, \omega_{\tau}, \mu_{\tau})}) f(t, \cdot)$ is uniformly Lipschitz continuous.

Remark 3.6. Assume that $\Phi : \mathbb{D}_{T,d} \rightarrow \mathbb{R}$ is twice continuously strongly vertically differentiable and satisfies the following locally Lipschitz continuous condition: for any $t \in [0, T]$ and $\phi = \Phi, \partial_{\omega_t} \Phi, \partial_{\omega_t}^2 \Phi$,

$$(79) \quad |\phi(\omega_T) - \phi(\omega'_T)| \leq C(1 + \|\omega_T\|^k + \|\omega'_T\|^k) \|\omega_T - \omega'_T\|, \quad \forall (\omega, \omega') \in \mathbb{D}_{T,d}^2.$$

Then, the main result [41, Theorem 4.5] is still true. For the reader's convenience, the proof is sketched in the appendix, using our partial Itô-Dupire formula.

Lemma 3.7. *Let (f, Φ) satisfy Assumption **(H1)**. Then $(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t})$ is almost surely vertically differentiable at (t, γ) . The derivative $(\partial_{\omega_t} Y^{\gamma_t, \eta_t}, \partial_{\omega_t} Z^{\gamma_t, \eta_t}) \in \mathbb{S}^p([t, T], \mathbb{R}^d) \times \mathbb{H}^p([t, T], \mathbb{R}^{d \times d})$, for any $p \geq 1$, is the unique solution to BSDE*

$$(80) \quad \begin{aligned} \mathcal{Y}(s) &= \partial_{\omega_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}) + \int_s^T \partial_{\omega_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr + \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Y}(r) dr \\ &+ \int_s^T \mathcal{Z}(r) \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr - \int_s^T \mathcal{Z}(r) dB(r), \quad s \in [t, T]. \end{aligned}$$

Furthermore, since $(\partial_{\omega_t} Y^{\gamma_t, \eta_t}, \partial_{\omega_t} Z^{\gamma_t, \eta_t})$ is independent of \mathcal{F}_t , we have that for any $K > 0$, there are positive constants C_p and $C_{K,p}$ such that

$$(81) \quad \begin{aligned} &\|(\partial_{\omega_t} Y^{\gamma_t, \eta_t}, \partial_{\omega_t} Z^{\gamma_t, \eta_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} < C_p, \\ &\|(\partial_{\omega_t} Y^{\gamma_t, \eta_t} - \partial_{\omega_t} Y^{\gamma'_t, \eta'_t}, \partial_{\omega_t} Z^{\gamma_t, \eta_t} - \partial_{\omega_t} Z^{\gamma'_t, \eta'_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} < C_{K,p} (\|\gamma_t - \gamma'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t})), \\ &\forall (\gamma, \eta), (\gamma', \eta') \in \mathbb{D}_{T,d} \times \mathbb{M}_2^D \text{ such that } \|\mathcal{L}_{\eta_t}\|, \|\mathcal{L}_{\eta'_t}\| \leq K. \end{aligned}$$

Proof. According to the preceding remark and [41, Lemma 3.8], we have the first two assertions. In view of the standard estimate for linear BSDEs, Lipschitz continuity of (Φ, f) and Lemma 2.4, we have

$$\begin{aligned} &\|\partial_{\omega_t} Y^{\gamma_t, \eta_t}\|_{\mathbb{S}^p} + \|\partial_{\omega_t} Z^{\gamma_t, \eta_t}\|_{\mathbb{H}^p} \\ &\leq C (\|\partial_{\omega_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}})\|_{L^p} + \|\int_t^T |\partial_{\omega_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})| dr\|_{L^p}), \end{aligned}$$

and thus the first inequality of (81). The last inequality is proved in a similar way to Lemma 3.4. \square

Furthermore, we have

Proposition 3.8. *Let (f, Φ) satisfy Assumption **(H1)**. Then for any $\tau \leq t$, $(Y^{\gamma_t, \eta_t}(s), Z^{\gamma_t, \eta_t}(s))$ is strongly vertically differentiable at (τ, t, γ) . Moreover, the derivative $(\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}, \partial_{\omega_\tau} Z^{\gamma_t, \eta_t}) \in \mathbb{S}^p([t, T], \mathbb{R}^d) \times \mathbb{H}^p([t, T], \mathbb{R}^{d \times d})$, $\forall p \geq 1$, is the unique solution to BSDE*

$$(82) \quad \begin{aligned} \mathcal{Y}(s) &= \partial_{\omega_\tau} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}) + \int_s^T \partial_{\omega_\tau} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr + \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Y}(r) dr \\ &+ \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Z}(r) dr - \int_s^T \mathcal{Z}(r) dB(r), \quad s \in [t, T]. \end{aligned}$$

Furthermore, since $(\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}, \partial_{\omega_\tau} Z^{\gamma_t, \eta_t})$ is independent of \mathcal{F}_t , we have that for any $K > 0$,

$$(83) \quad \begin{aligned} &\|(\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}, \partial_{\omega_\tau} Z^{\gamma_t, \eta_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} < C_p, \\ &\|(\partial_{\omega_\tau} Y^{\gamma_t, \eta_t} - \partial_{\omega_\tau} Y^{\gamma'_t, \eta'_t}, \partial_{\omega_\tau} Z^{\gamma_t, \eta_t} - \partial_{\omega_\tau} Z^{\gamma'_t, \eta'_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} < C_{K,p} (\|\gamma_t - \gamma'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t})), \\ &\forall (\gamma, \eta), (\gamma', \eta') \in \mathbb{D}_{T,d} \times \mathbb{M}_2^D \text{ such that } \|\mathcal{L}_{\eta_t}\|, \|\mathcal{L}_{\eta'_t}\| \leq K, \end{aligned}$$

for some positive constants C_p and $C_{K,p}$.

Proof. In view of Assumption **(H1)** and Lemma 3.1, we see that equation (82) has a unique solution $(\partial_{\omega_\tau} Y, \partial_{\omega_\tau} Z) \in \mathbb{S}^p \times \mathbb{H}^p$, $\forall p \geq 1$. Here, we consider the one-dimensional case for simplicity. For any

$h > 0$, recall that $\gamma^{\tau, h} = \gamma + h1_{[r, T]}$. Set

$$(84) \quad \begin{aligned} \gamma' &:= \gamma^{\tau, h}, \quad \Delta_h Y := \frac{1}{h}(Y' - Y) := \frac{1}{h}(Y^{\gamma_t^{\tau, h}, \eta_t} - Y^{\gamma_t, \eta_t}), \quad \text{and} \\ \Delta_h Z &:= \frac{1}{h}(Z' - Z) := \frac{1}{h}(Z^{\gamma_t^{\tau, h}, \eta_t} - Z^{\gamma_t, \eta_t}). \end{aligned}$$

Then we know that $(\Delta_h Y, \Delta_h Z)$ solves the following BSDE

$$\begin{aligned} \Delta_h Y(s) &= \frac{1}{h}(\Phi' - \Phi) + \frac{1}{h} \int_s^T [f(\Theta_r^{\gamma', \eta}, \mathcal{L}_{\Theta_r^{\eta}}) - f(\Theta_r^{\gamma, \eta}, \mathcal{L}_{\Theta_r^{\eta}})] dr - \int_s^T \Delta_h Z(r) dB(r) \\ &=: \Delta_h \Phi + \int_s^T (a_r \Delta_h Y(r) + b_r \Delta_h Z(r) + \Delta_h f) dr - \int_s^T \Delta_h Z(r) dB(r), \end{aligned}$$

where

$$\begin{aligned} \Phi' &:= \Phi(B^{\gamma'}, \mathcal{L}_{B^{\eta}}), \quad \Phi := \Phi(B^{\gamma}, \mathcal{L}_{B^{\eta}}), \quad \Delta_h \Phi := \int_0^1 \partial_{\gamma_{\tau}} \Phi(B^{\gamma^{\tau, h\theta}}, \mathcal{L}_{B^{\eta}}) d\theta, \\ a_r &:= \int_0^1 \partial_y f(B_r^{\gamma'}, Y + \theta(Y' - Y), Z', \mathcal{L}_{\Theta_r^{\eta}}) d\theta, \quad b_r := \int_0^1 \partial_z f(B_r^{\gamma'}, Y, Z + \theta(Z' - Z), \mathcal{L}_{\Theta_r^{\eta}}) d\theta, \\ \text{and } \Delta_h f &:= \frac{1}{h} f(B_r^{\omega}, Y, Z, \mathcal{L}_{B_r^{\eta}}, \mathcal{L}_Y) \Big|_{\omega=\gamma}^{\omega=\gamma'} = \int_0^1 \partial_{\omega_{\tau}} f(B^{\gamma^{\tau, h\theta}}, Y, Z, \mathcal{L}_{B_r^{\eta}}, \mathcal{L}_Y) d\theta. \end{aligned}$$

Then $(\delta Y, \delta Z) := (\Delta_h Y - \partial_{\omega_{\tau}} Y, \Delta_h Z - \partial_{\omega_{\tau}} Z)$ satisfies BSDE

$$\begin{aligned} \delta Y(s) &= (\Delta_h \Phi - \partial_{\omega_{\tau}} \Phi) + \int_s^T (a_r \delta Y + b_r \delta Z + (\Delta_h f - \partial_{\omega_{\tau}} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta}}))) dr \\ &\quad + \int_s^T [(a_r - \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta}})) \partial_{\omega_{\tau}} Y + (b_r - \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta}})) \partial_{\omega_{\tau}} Z] dr - \int_s^T \delta Z dB(r). \end{aligned}$$

According to standard estimate for BSDEs (or Lemma 3.1 for $p = 2$) and Lemma 3.4, we have

$$\begin{aligned} \|\delta Y\|_{\mathbb{S}^p}^p + \|\delta Z\|_{\mathbb{H}^p}^p &\leq C \|\Delta_h \Phi - \partial_{\omega_{\tau}} \Phi\|_{L^p}^p + \left\| \int_t^T |\Delta_h f - \partial_{\omega_{\tau}} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta}})| dr \right\|_{L^p}^p + O(|h|) \\ &\leq O(|h|), \end{aligned}$$

and thus the strongly vertical differentiability. \square

To show the differentiability of Y^{γ_t, η_t} with respect to η_t , we follow a similar argument as in the state-dependent case for SDEs made in [9]. Firstly we show that Y^{γ_t, η_t} is Gâteaux differentiable in η_t in the sense of (30) and Remark 2.8. To this end, we need to prove that for any $\xi \in L^2(\mathcal{F}_t, \mathbb{R}^d)$ and $\eta_t^{\lambda \xi} := \eta_t + \lambda \xi 1_{[t, T]}$, $\lambda > 0$, the following limit exits in $\mathbb{S}^2([t, T], \mathbb{R}^d)$,

$$(85) \quad \partial_{\eta} Y^{\gamma_t, \eta_t}(\xi) := \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (Y^{\gamma_t, \eta_t^{\lambda \xi}} - Y^{\gamma_t, \eta_t}).$$

Then we show that $\partial_{\eta} Y^{\gamma_t, \eta_t}(\cdot) : L^2(\mathcal{F}_t, \mathbb{R}^d) \rightarrow \mathbb{S}^2([t, T], \mathbb{R}^d)$ is a bounded linear operator, and moreover, it is continuous in the following sense: for any $\zeta \in L^2(\mathcal{F}_t, \mathbb{R}^d)$, $\partial_{\eta} Y^{\gamma_t, \eta_t + \zeta 1_{[t, T]}}$ converges to $\partial_{\eta} Y^{\gamma_t, \eta_t}$ in the sense of operators as ζ goes to zero. In view of Remark 2.5, we see that Y^{γ_t, η_t} is Fréchet (vertically) differentiable in the sense of (29) and Remark 2.8.

To this end, consider the following linear BSDE

$$\begin{aligned}
(86) \quad \mathcal{Y}^{\gamma_t, \eta_t, \xi}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}_r^{\tilde{\eta}_t}) \tilde{\xi}] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Y}^{\gamma_t, \eta_t, \xi}(r) dr + \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Z}^{\gamma_t, \eta_t, \xi}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})(\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi} + \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}})(r)] dr \\
&- \int_s^T \mathcal{Z}^{\gamma_t, \eta_t, \xi}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

Here, $(\tilde{B}, \tilde{\eta}, \tilde{\xi}, \tilde{Y}^{\tilde{\eta}}, \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}}, \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}})$ is an independent copy of $(B, \eta, \xi, Y^{\eta_t}, \partial_{\omega_t} Y^{\gamma_t, \mathcal{L}_{\eta_t}} |_{\gamma=\eta}, \mathcal{Y}^{\eta_t, \xi})$, and $\mathcal{Y}^{\eta_t, \xi}$ satisfies the following linear mean-field BSDE

$$\begin{aligned}
(87) \quad \mathcal{Y}^{\eta_t, \xi}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}_r^{\tilde{\eta}_t}) \tilde{\xi}] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Y}^{\eta_t, \xi}(r) dr + \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \mathcal{Z}^{\eta_t, \xi}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})(\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi} + \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}})(r)] dr \\
&- \int_s^T \mathcal{Z}^{\eta_t, \xi}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

Lemma 3.9. *For any $\xi \in L^2(\mathcal{F}_t, \mathbb{R}^d)$, there exists a unique solution $(\mathcal{Y}^{\eta_t, \xi}, \mathcal{Z}^{\eta_t, \xi}) \in \mathbb{S}^2([t, T]) \times \mathbb{H}^2([t, T], \mathbb{R}^d)$ for BSDE (87). Moreover, $(\mathcal{Y}^{\eta_t, \xi}, \mathcal{Z}^{\eta_t, \xi})$ is linear in ξ , and we have*

$$(88) \quad \|(\mathcal{Y}^{\eta_t, \xi}, \mathcal{Z}^{\eta_t, \xi})\|_{\mathbb{S}^2 \times \mathbb{H}^2} \leq C \|\xi\|_{L^2}$$

for some constant C .

Proof. By Lipschitz continuity of $(\partial_{\mu_t} \Phi, \partial_{\mu_t} f)$, we have

$$\tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B_T^{\eta_t}, \mathcal{L}_{B_T^{\eta_t}}, \tilde{B}_T^{\tilde{\eta}_t}) \tilde{\xi}] \in L^2(\mathcal{F}_T), \quad \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}_r^{\tilde{\eta}_t}) \tilde{\xi}] \in L^2(\mathcal{F}_r).$$

Since f is uniformly Lipschitz continuous in (y, z) , $\partial_{(y, z)} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})$ is uniformly bounded. Set $g(r, x) := \partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, x)$. In view of Lemma 3.4 and Assumption **(H1)**, we see that $g(\cdot, 0) \in \mathbb{H}^2$. Then by Lemma 3.1, to show the well-posedness of linear mean-field BSDE (87), we only need to check the following

$$\int_t^T \left| \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})(\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi})] \right| dr \in L^2(\mathcal{F}_T).$$

Let

$$(89) \quad F_2(t, x, y, z, \mu, \nu) := \tilde{\mathbb{E}}[\partial_\nu f(t, x, y, z, \mu, \nu, \tilde{Y}^{\tilde{\eta}_t}(r))(\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}}(r) \tilde{\xi})].$$

Then by Lipschitz continuity of $\partial_\nu f$ and Lemma 3.7, we have

$$\begin{aligned}
& F_2(t, x, y, z, \mu, \nu) \\
&= \tilde{\mathbb{E}} \left[\tilde{\mathbb{E}}_{\tilde{\mathcal{F}}_t} [\partial_\nu f(t, x, y, z, \mu, \nu, \tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}(r)) (\partial_{\omega_t} \tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}(r))] \Big|_{\gamma_t = \tilde{\eta}_t} \tilde{\xi} \right] \\
&\leq C \tilde{\mathbb{E}} \left[\tilde{\mathbb{E}}_{\tilde{\mathcal{F}}_t} [|\tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}| |\partial_{\omega_t} \tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}(r)|] \Big|_{\gamma_t = \tilde{\eta}_t} \tilde{\xi} \right] + \partial_\nu f(t, x, y, z, \mu, \nu, 0) \tilde{\mathbb{E}} \left[\tilde{\mathbb{E}}_{\tilde{\mathcal{F}}_t} [|\tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}(r)|] \Big|_{\gamma_t = \tilde{\eta}_t} \tilde{\xi} \right] \\
&\leq C \tilde{\mathbb{E}} \left[(\tilde{\mathbb{E}}_{\tilde{\mathcal{F}}_t} [|\tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}|^2]^{\frac{1}{2}}) (\tilde{\mathbb{E}}_{\tilde{\mathcal{F}}_t} [|\partial_{\omega_t} \tilde{Y}^{\gamma_t, \mathcal{L}_{\eta_t}}(r)|^2]^{\frac{1}{2}}) \Big|_{\gamma_t = \tilde{\eta}_t} \tilde{\xi} \right] + C \partial_\nu f(t, x, y, z, \mu, \nu, 0) \\
&\leq C \tilde{\mathbb{E}} \left[\tilde{\mathbb{E}}_{\tilde{\mathcal{F}}_t} [(1 + \|\gamma_t\|)] \Big|_{\gamma_t = \tilde{\eta}_t} \tilde{\xi} \right] + C \partial_\nu f(t, x, y, z, \mu, \nu, 0) \\
&\leq C + C \partial_\nu f(t, x, y, z, \mu, \nu, 0),
\end{aligned}$$

where we applied Lemmas 3.4 and 3.7 in the second last inequality. Then according to Lemma 3.4, we have $\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, 0) \in \mathbb{H}^2$, and thus the well-posedness of (87). For inequality (88), similar to the proof of Lemma 3.4, we have

$$\begin{aligned}
& \|\mathcal{Y}^{\eta_t, \xi}\|_{\mathbb{S}^2}^2 + \|\mathcal{Z}^{\eta_t, \xi}\|_{\mathbb{H}^2}^2 \\
&\leq C \mathbb{E} \left[|\tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}]|^2 + \int_s^T |\tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}]|^2 dr \right. \\
&\quad \left. + \int_s^T |\tilde{\mathbb{E}} \partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) (\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi})|^2 dr \right] \\
&\leq C \left((\tilde{\mathbb{E}} \|\tilde{B}^{\tilde{\eta}_t}\| |\tilde{\xi}|)^2 + \|\xi\|_{L^2}^2 \mathbb{E} |\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, 0)|^2 \right. \\
&\quad \left. + \|\xi\|_{L^2}^2 \mathbb{E} \int_s^T |\partial_{\mu_t} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, 0)|^2 dr + \mathbb{E} \int_s^T |\tilde{\mathbb{E}}[\tilde{Y}^{\tilde{\eta}_t} \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi}]|^2 dr \right. \\
&\quad \left. + \|\xi\|_{L^2}^2 \mathbb{E} \int_s^T |\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, 0)|^2 dr \right) \leq C \|\xi\|_{L^2}^2.
\end{aligned}$$

□

Since BSDE (87) is well-posed, we see that BSDE (86) is also well-posed. In conclusion, we have

Corollary 3.10. *There exists a unique solution $(\mathcal{Y}^{\gamma_t, \eta_t, \xi}, \mathcal{Z}^{\gamma_t, \eta_t, \xi}) \in \mathbb{S}^2([t, T]) \times \mathbb{H}^2([t, T], \mathbb{R}^d)$ to BSDE (86). Moreover,*

$$(90) \quad (\mathcal{Y}^{\eta_t, \xi}, \mathcal{Z}^{\eta_t, \xi}) = (\mathcal{Y}^{\gamma_t, \eta_t, \xi}, \mathcal{Z}^{\gamma_t, \eta_t, \xi})|_{\gamma = \eta}.$$

Lemma 3.11. *The map $\xi \rightarrow \mathcal{Y}^{\gamma_t, \eta_t, \xi}$ is a bounded linear operator from $L^2(\mathcal{F}_t, \mathbb{R}^d)$ to $\mathbb{S}^2([t, T])$. Moreover, it is the Gâteaux derivative of Y^{γ_t, η_t} with respect to η_t in the following sense*

$$(91) \quad \mathcal{Y}^{\gamma_t, \eta_t, \xi} = \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (Y^{\gamma_t, \eta_t^{\lambda \xi}} - Y^{\gamma_t, \eta_t}) \quad \text{strongly in } \mathbb{S}^2([t, T]).$$

In particular, $\mathcal{Y}^{\gamma_t, \eta_t, \xi}(s)$ is the Gâteaux derivative of $Y^{\gamma_t, \eta_t}(s)$ in the sense of (30).

Proof. Since $\mathcal{Y}^{\eta_t, \xi}$ is linear in ξ , we see that $(\mathcal{Y}^{\gamma_t, \eta_t, \xi}, \mathcal{Z}^{\gamma_t, \eta_t, \xi})$ is also linear in ξ . Moreover, we have the following estimate

$$(92) \quad \|(\mathcal{Y}^{\gamma_t, \eta_t, \xi}, \mathcal{Z}^{\gamma_t, \eta_t, \xi})\|_{\mathbb{S}^2 \times \mathbb{H}^2} \leq C \|\xi\|_{L^2}.$$

Therefore, we have the first assertion.

In the following, we omit the fixed subscript t and write $(Y, Z) := (Y(r), Z(r))$ if no confusion raised. Besides, the constant C may change from line to line. Set

$$(93) \quad \begin{aligned} \Delta_\lambda Y &:= \frac{1}{\lambda}(Y^{\gamma, \eta^{\lambda\xi}} - Y^{\gamma, \eta}), \quad \Delta_\lambda Z := \frac{1}{\lambda}(Z^{\gamma, \eta^{\lambda\xi}} - Z^{\gamma, \eta}), \quad \text{and} \\ \Delta_\lambda \Phi &:= \frac{1}{\lambda}[\Phi(B_T^\gamma, \mathcal{L}_{B_T^{\eta^{\lambda\xi}}}) - \Phi(B_T^\gamma, \mathcal{L}_{B_T^\eta})]. \end{aligned}$$

Then according to Lemma 3.4, we have

$$(94) \quad \|\Delta_\lambda Y\|_{\mathbb{S}^2} + \|\Delta_\lambda Z\|_{\mathbb{H}^2} \leq C \frac{1}{\lambda} \|\eta_t^{\lambda\xi} - \eta_t\|_{\mathbb{S}^2} \leq C \|\xi\|_{L^2}.$$

In view of BSDE (68), we see that $(\Delta_\lambda Y, \Delta_\lambda Z)$ satisfies the following linear mean-field BSDE

$$(95) \quad \begin{aligned} \Delta_\lambda Y &= \Delta_\lambda \Phi + \int_s^T \left[\alpha(r) \Delta_\lambda Y + \beta(r) \Delta_\lambda Z + \tilde{\mathbb{E}}[\tilde{g}(r) \frac{1}{\lambda} (\tilde{Y}^{\tilde{\eta}^{\lambda\xi}} - \tilde{Y}^{\tilde{\eta}})] + \Delta_\lambda f \right] dr \\ &\quad - \int_s^T \Delta_\lambda Z dB(r), \end{aligned}$$

where

$$\begin{aligned} \alpha(r) &:= \int_0^1 \partial_y f(B_r^\gamma, Y^{\gamma, \eta} + \theta(Y^{\gamma, \eta^{\lambda\xi}} - Y^{\gamma, \eta}), Z^{\gamma, \eta^{\lambda\xi}}, \mathcal{L}_{\Theta_r^{\eta^{\lambda\xi}}}) d\theta, \\ \beta(r) &:= \int_0^1 \partial_z f(B_r^\gamma, Y^{\gamma, \eta}, Z^{\gamma, \eta} + \theta(Z^{\gamma, \eta^{\lambda\xi}} - Z^{\gamma, \eta}), \mathcal{L}_{\Theta_r^{\eta^{\lambda\xi}}}) d\theta, \\ \tilde{g}(r) &:= \int_0^1 \partial_\nu f(\Theta^{\gamma, \eta}, \mathcal{L}_{B^{\eta^{\lambda\xi}}}, \mathcal{L}_{Y^{\eta} + \theta(Y^{\eta^{\lambda\xi}} - Y^\eta)}, \tilde{Y}^{\tilde{\eta}} + \theta(\tilde{Y}^{\tilde{\eta}^{\lambda\xi}} - \tilde{Y}^{\tilde{\eta}})) d\theta, \quad \text{and} \\ \Delta_\lambda f(r) &:= \frac{1}{\lambda} [f(\Theta^{\gamma, \eta}, \mathcal{L}_{B^{\eta^{\lambda\xi}}}, \mathcal{L}_{Y^\eta}) - f(\Theta^{\gamma, \eta}, \mathcal{L}_{B^\eta}, \mathcal{L}_{Y^\eta})]. \end{aligned}$$

According to estimate (72) in Lemma 3.4, we have

$$(96) \quad \|\Delta_\lambda \tilde{Y}^{\tilde{\eta}}\|_{\mathbb{S}^2} := \left\| \frac{1}{\lambda} (\tilde{Y}^{\tilde{\eta}^{\lambda\xi}} - \tilde{Y}^{\tilde{\eta}}) \right\|_{\mathbb{S}^2} \leq C \frac{1}{\lambda} \|\eta_t^{\lambda\xi} - \eta_t\|_{\mathbb{S}^2} \leq C \|\xi\|_{L^2}.$$

Then, in view of Assumption **(H1)**, we have

$$(97) \quad \left\| \int_t^T \tilde{\mathbb{E}}[\tilde{g}(r) \Delta_\lambda \tilde{Y}^{\tilde{\eta}}] dr \right\|_{L^2} + \left\| \int_t^T \Delta_\lambda f dr \right\|_{L^2} \leq C \|\xi\|_{L^2}.$$

Thus BSDE (95) has a unique solution $(\Delta_\lambda Y, \Delta_\lambda Z)$, and therefore, $(\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}, \Delta_\lambda Z - \mathcal{Z}^{\gamma_t, \eta_t, \xi})$ is the unique solution of the following BSDE

$$\begin{aligned} Y(s) &= (\Delta_\lambda \Phi - \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}]) + \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) Y dr \\ &\quad + \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) Z(r) dr + \int_s^T (\Delta_\lambda f - \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}]) dr \\ &\quad + \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) (\Delta_\lambda \tilde{Y}^{\tilde{\eta}} - \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi} - \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}})] dr \\ &\quad + \int_t^T R_1(r) dr - \int_s^T Z dB(r) \end{aligned}$$

with

$$\begin{aligned} R_1(r) &:= (\alpha(r) - \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})) \Delta_\lambda Y + (\beta(r) - \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})) \Delta_\lambda Z \\ &\quad + \tilde{\mathbb{E}} \left[(\tilde{g}(r) - \partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})) \Delta_\lambda \tilde{Y}^{\tilde{\eta}} \right]. \end{aligned}$$

Since $\partial_{(y,z)} f$ is bounded, from the standard estimate for solutions of BSDEs, we have

$$(98) \quad \|\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}\|_{\mathbb{S}^2}^2 \leq C(\|A_1\|_{L^2}^2 + \|A_2\|_{L^2}^2 + \|A_3\|_{L^2}^2 + \|A_4\|_{L^2}^2)$$

with

$$\begin{aligned} A_1 &:= \Delta_\lambda \Phi - \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}], \quad A_2 := \int_t^T |R_1(r)| dr, \\ A_3 &:= \int_t^T \left| (\Delta_\lambda f - \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}]) \right| dr, \quad \text{and} \\ A_4 &:= \int_t^T \left| \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) (\Delta_\lambda \tilde{Y}^{\tilde{\eta}} - \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi} - \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}})](r) \right| dr. \end{aligned}$$

For A_1 , according to the Lipschitz continuity of $\partial_{\mu_t} \Phi$, we have

$$\begin{aligned} (99) \quad \mathbb{E}|A_1|^2 &= \mathbb{E} \left| \int_0^1 \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^\gamma, \mathcal{L}_{B^{\eta+\theta(B^{\eta^{\lambda\xi}} - B^\eta)}, \tilde{B}^{\tilde{\eta}} + \theta(\tilde{B}^{\tilde{\eta}^{\lambda\xi}} - \tilde{B}^{\tilde{\eta}})) \tilde{\xi} \right. \\ &\quad \left. - \partial_{\mu_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}] d\theta \right|^2 \\ &\leq C \left([\bar{E} \|\bar{B}^{\tilde{\eta}^{\lambda\xi}} - \bar{B}^{\tilde{\eta}}\|^2]^{\frac{1}{2}} \|\xi\|_{L^2} + \tilde{\mathbb{E}}[\|\tilde{B}^{\tilde{\eta}^{\lambda\xi}} - \tilde{B}^{\tilde{\eta}}\| \|\tilde{\xi}\|] \right)^2 \leq C\lambda^2 \|\xi\|_{L^2}^4, \end{aligned}$$

for a constant C independent of γ and η .

Term A_2 is estimated as follows:

$$\begin{aligned} (100) \quad |A_2|^2 &\leq C \left| \int_t^T (\alpha(r) - \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})) \Delta_\lambda Y dr \right|^2 \\ &\quad + C \left| \int_t^T (\beta(r) - \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})) \Delta_\lambda Z dr \right|^2 \\ &\quad + C \left| \int_t^T \tilde{\mathbb{E}}[(\tilde{g}(r) - \partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})) \Delta_\lambda \tilde{Y}^{\tilde{\eta}}] dr \right|^2. \end{aligned}$$

For the first two terms on the right hand side of the above inequality, by the Lipschitz continuity of $\partial_{(y,z)}f$ and inequality (94), we obtain

$$\mathbb{E} \left| \int_t^T (\alpha(r) - \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})) \Delta_\lambda Y dr \right|^2 + \left| \int_t^T (\beta(r) - \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}})) \Delta_\lambda Z dr \right|^2 \leq C\lambda^2 \|\xi\|_{L^2}^4.$$

For the third term, we claim that

$$(101) \quad \mathbb{E} \left| \int_t^T \tilde{\mathbb{E}} \left[(\tilde{g}(r) - \partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})) \Delta_\lambda \tilde{Y}^{\tilde{\eta}} \right] dr \right|^2 \leq C\lambda^2 \|\xi\|_{L^2}^4,$$

with C depending only on $\|\eta_t\|_{\mathbb{S}^2}$, and therefore we have

$$(102) \quad \mathbb{E}|A_2|^2 \leq C\lambda^2 \|\xi\|_{L^2}^4,$$

in view of (100) and above estimates. Indeed, by the Hölder inequality and estimate (96), we have

$$\begin{aligned} \mathbb{E} \left| \int_t^T \tilde{\mathbb{E}}[(\tilde{g}(r) - \partial_\nu f) \Delta_\lambda \tilde{Y}^{\tilde{\eta}}] dr \right|^2 &\leq \mathbb{E} \left[\int_t^T \tilde{\mathbb{E}}|\tilde{g} - \partial_\nu f|^2 dr \right] \int_t^T \tilde{\mathbb{E}}|\Delta_\lambda \tilde{Y}^{\tilde{\eta}}|^2 dr \\ &\leq C\|\xi\|_{L^2}^2 \mathbb{E} \int_t^T \tilde{\mathbb{E}} \left| \int_0^1 (\partial_\nu f(\Theta_r^{\gamma, \eta}, \mathcal{L}_{B^{\eta\lambda\xi}}, \mathcal{L}_{Y^{\eta+\theta(Y^{\eta\lambda\xi} - Y^\eta)}, \tilde{Y}^{\tilde{\eta}} + \theta(\tilde{Y}^{\tilde{\eta}\lambda\xi} - \tilde{Y}^{\tilde{\eta}})) \right. \\ &\quad \left. - \partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})) d\theta \right|^2 dr \\ &\leq C\|\xi\|_{L^2}^2 (\|\tilde{Y}^{\tilde{\eta}\lambda\xi} - \tilde{Y}^{\tilde{\eta}}\|_{\mathbb{S}^2} + \|B^{\eta\lambda\xi} - B^\eta\|_{\mathbb{S}^2})^2 \leq C\lambda^2 \|\xi\|_{L^2}^4. \end{aligned}$$

For A_3 , from Lipschitz continuity of $\partial_{\mu_t} f$ in $(\mu, \nu, \tilde{\omega})$, we have

$$(103) \quad \begin{aligned} \mathbb{E}|A_3|^2 &= \mathbb{E} \left| \int_t^T \int_0^1 \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{B^{\eta+\theta(\eta\lambda\xi - \eta)}}, \mathcal{L}_{Y^\eta}, \tilde{B}^{\tilde{\eta}+\theta(\tilde{\eta}\lambda\xi - \tilde{\eta})}) \right. \\ &\quad \left. - \partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t})] \tilde{\xi} d\theta dr \right|^2 \\ &\leq C \left| \tilde{\mathbb{E}}[(\|\eta_t^{\lambda\xi} - \eta_t\|_{\mathbb{S}^2} + \|\tilde{\eta}^{\lambda\xi} - \tilde{\eta}\|) |\tilde{\xi}|] \right|^2 \\ &\leq C \left| \tilde{\mathbb{E}}[\lambda\|\xi\|_{L^2} \tilde{\xi} + \lambda|\tilde{\xi}|^2] \right|^2 \leq C\lambda^2 \|\xi\|_{L^2}^4. \end{aligned}$$

We now estimate A_4 . Since

$$(104) \quad \Delta_\lambda \tilde{Y}^{\tilde{\eta}} - \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t} \tilde{\xi}} - \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}} = A_{41} + A_{42}$$

with

$$(105) \quad \begin{aligned} A_{41} &:= \left[\frac{1}{\lambda} (\tilde{Y}^{\tilde{\eta}\lambda\xi, \mathcal{L}_{\tilde{\eta}\lambda\xi}} - \tilde{Y}^{\tilde{\eta}, \mathcal{L}_{\tilde{\eta}\lambda\xi}}) - \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t} \tilde{\xi}} \right], \quad \text{and} \\ A_{42} &:= \left[\frac{1}{\lambda} (\tilde{Y}^{\tilde{\eta}, \mathcal{L}_{\tilde{\eta}\lambda\xi}} - \tilde{Y}^{\tilde{\eta}, \mathcal{L}_{\tilde{\eta}}}) - \tilde{\mathcal{Y}}^{\tilde{\eta}_t, \tilde{\xi}} \right], \end{aligned}$$

then, from boundedness of $\partial_\nu f$, we have

$$(106) \quad \begin{aligned} \mathbb{E}|A_4|^2 &= \mathbb{E} \left| \int_t^T \tilde{\mathbb{E}} \left[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) (A_{41}(r) + A_{42}(r)) \right] dr \right|^2 \\ &\leq C \left(\left| \int_t^T \tilde{\mathbb{E}}[A_{41}] dr \right|^2 + \left| \int_t^T \tilde{\mathbb{E}}[A_{42}] dr \right|^2 \right). \end{aligned}$$

From Lemma 3.7, we have

$$\begin{aligned}
(107) \quad \left| \int_t^T \tilde{\mathbb{E}}[A_{41}] dr \right|^2 &\leq C \int_t^T \left[\tilde{\mathbb{E}} \int_0^1 |(\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}^{\lambda\theta\tilde{\xi}}, \mathcal{L}_{\eta^{\lambda\xi}} - \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}, \mathcal{L}_{\eta}}) \tilde{\xi}| d\theta \right]^2 dr \\
&\leq C \|\xi\|_{L^2}^2 \int_t^T \int_0^1 \tilde{\mathbb{E}} |\partial_{\omega_t} \tilde{Y}^{\tilde{\eta}^{\lambda\theta\tilde{\xi}}, \mathcal{L}_{\eta^{\lambda\xi}} - \partial_{\omega_t} \tilde{Y}^{\tilde{\eta}, \mathcal{L}_{\eta}}|^2 d\theta dr \\
&\leq C \|\xi\|_{L^2}^2 \int_t^T \int_0^1 \tilde{\mathbb{E}} (\|\tilde{\eta}_t^{\lambda\theta\tilde{\xi}} - \tilde{\eta}_t\|)^2 d\theta dr \leq C \lambda^2 \|\xi\|_{L^2}^4
\end{aligned}$$

for a constant C only depending on $\|\eta_t\|_{\mathbb{S}^2}$. Since

$$(108) \quad \left| \int_t^T \tilde{\mathbb{E}}[A_{42}] dr \right|^2 \leq \int_t^T \tilde{\mathbb{E}} |A_{42}|^2 dr \leq C \sup_{\gamma_t} \int_t^T \tilde{\mathbb{E}} |\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}|^2 dr,$$

for a constant C independent of (γ, η) , we have

$$(109) \quad \mathbb{E} |A_4|^2 \leq C \left(\lambda^2 \|\xi\|_{L^2}^4 + \sup_{\gamma_t} \int_t^T \tilde{\mathbb{E}} |\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}|^2 dr \right).$$

Finally, in view of inequalities (99), (102), (103), (109) and (98), we have

$$\|\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}\|_{\mathbb{S}^2}^2 \leq C (\lambda^2 \|\xi\|_{L^2}^4 + \sup_{\gamma_t} \int_t^T \|\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}\|_{\mathbb{S}^2}^2 dr),$$

where C only depends on $\|\eta_t\|_{\mathbb{S}^2}$. Then, using Gronwall's inequality, we have

$$(110) \quad \|\Delta_\lambda Y - \mathcal{Y}^{\gamma_t, \eta_t, \xi}\|_{\mathbb{S}^2}^2 \leq C \lambda^2 \|\xi\|_{L^2}^4 \rightarrow 0, \quad \text{as } \lambda \rightarrow 0.$$

□

In view of our Assumption **(H1)**, the solution $(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t})$ of BSDE (68) is indeed strongly vertically differentiable in η_t . According to Definition 2.9, for any $\tau \leq t$ and $\xi \in L^2(\mathcal{F}_\tau, \mathbb{R}^d)$, let $\eta_t^{\tau, \lambda\xi} := \eta_t + \lambda\xi 1_{[\tau, T]}$. Similar as the vertical differentiable case, we firstly need to show the following limit exists in $\mathbb{S}^2([t, T])$,

$$(111) \quad \partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi} := \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (Y^{\gamma_t, \eta_t^{\tau, \lambda\xi}} - Y^{\gamma_t, \eta_t}).$$

Indeed, $\partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}$ is the unique solution of the following BSDE

$$\begin{aligned}
(112) \quad \partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_\tau} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_\tau} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{\tilde{\eta}_t}) \tilde{\xi}] dr \\
&\quad + \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}(r) dr \\
&\quad + \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t})(\partial_{\omega_\tau} \tilde{Y}^{\tilde{\eta}_t, \mathcal{L}_{\eta_t}} \tilde{\xi} + \partial_{\eta_\tau} \tilde{Y}^{\tilde{\eta}_t, \tilde{\xi}})(r)] dr \\
&\quad + \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\eta_\tau} Z^{\gamma_t, \eta_t, \xi}(r) dr \\
&\quad - \int_s^T \partial_{\eta_\tau} Z^{\gamma_t, \eta_t, \xi}(r) dB(r), \quad s \in [t, T],
\end{aligned}$$

where $\partial_{\eta_\tau} Y^{\eta_t, \xi}$ solves the following mean-field BSDE

$$\begin{aligned}
(113) \quad \partial_{\eta_\tau} Y^{\eta_t, \xi}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_\tau} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\eta_t}) \tilde{\xi}] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_\tau} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{\eta_t}) \tilde{\xi}] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\eta_\tau} Y^{\eta_t, \xi}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\eta_t})(\partial_{\omega_\tau} \tilde{Y}^{\eta_t, \mathcal{L}_{\eta_t}} \tilde{\xi} + \partial_{\eta_\tau} \tilde{Y}^{\eta_t, \xi})(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\eta_\tau} Z^{\eta_t, \xi}(r) dr - \int_s^T \partial_{\eta_\tau} Z^{\eta_t, \xi}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

According to Assumption **(H1)**, we see that BSDEs (113) and (112) are well-posed. Moreover, following a similar argument as in Lemma 3.11, for the Gâteaux strong vertical differentiability, we have

Lemma 3.12. $\partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \cdot}$ is a bounded linear operator from $L^2(\mathcal{F}_\tau, \mathbb{R}^d)$ to $\mathbb{S}^2([t, T])$. Moreover, $\partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}$ is the Gâteaux strong vertical derivative of Y^{γ_t, η_t} at (τ, t, η) :

$$(114) \quad \partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi} = \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (Y^{\gamma_t, \eta_t^{\tau, \lambda \xi}} - Y^{\gamma_t, \eta_t}), \quad \text{strongly in } \mathbb{S}^2([t, T]).$$

In particular, $\partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \cdot}(s)$ is the Gâteaux derivative of $Y^{\gamma_t, \eta_t}(s)$ at (τ, t, η) in the sense of (33).

To give an explicit representation of the vertical derivative $Y^{\gamma_t, \mathcal{L}_{\eta_t}}(\cdot)$ with respect to \mathcal{L}_{η_t} in view of (31), we need to find out a measurable random field $U^{\gamma_t, \mathcal{L}_{\eta_t}}(\cdot) : \mathbb{D}_{T, d} \rightarrow \mathbb{S}^2([t, T], \mathbb{R}^d)$, such that for any $s \geq t$ and $\xi \in L^2(\mathcal{F}_t, \mathbb{R}^d)$,

$$(115) \quad \mathcal{Y}^{\gamma_t, \eta_t, \xi}(s) = \bar{E}[U^{\gamma_t, \mathcal{L}_{\eta_t}}(\bar{\eta}_t)(s) \bar{\xi}],$$

where $(\bar{\eta}, \bar{\xi})$ is an independent copy of (η, ξ) . If (115) holds and moreover we show that Y^{γ_t, η_t} is Fréchet differentiable with respect to η_t in the sense of (29) and Remark 2.8, we have that

$$(116) \quad \partial_{\mu_t} Y^{\gamma_t, \mathcal{L}_{\eta_t}}(x_t) := U^{\gamma_t, \mathcal{L}_{\eta_t}}(x_t), \quad \forall x \in \mathbb{D}_{T, d},$$

is the vertical derivative of $Y^{\gamma_t, \mathcal{L}_{\eta_t}}$ at \mathcal{L}_{η_t} . Here and in the following, we write ∂_μ instead of $\partial_{\mathcal{L}_\eta}$. In view of (86) and (115), we formally deduce that $(U^{\gamma_t, \mathcal{L}_{\eta_t}}(x_t), V^{\gamma_t, \mathcal{L}_{\eta_t}}(x_t))$ solves the following BSDE

$$\begin{aligned}
(117) \quad U^{\gamma_t, \eta_t, x_t}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) U^{\gamma_t, \eta_t, x_t}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\omega_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\eta_t}) \tilde{U}^{\eta_t, x_t}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) V^{\gamma_t, \eta_t, x_t}(r) dr - \int_s^T V^{\gamma_t, \eta_t, x_t}(r) dB(r), \quad s \in [t, T],
\end{aligned}$$

where U^{η_t, x_t} solves the associated mean-field BSDE:

$$\begin{aligned}
(118) \quad U^{\eta_t, x_t}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) U^{\eta_t, x_t}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\omega_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \tilde{U}^{\tilde{\eta}_t, x_t}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) V^{\eta_t, x_t}(r) dr - \int_s^T V^{\eta_t, x_t}(r) dB(r).
\end{aligned}$$

According to Lemma 3.1, we see that mean-field BSDE (118) is well posed with $(U^{\eta_t, x_t}, V^{\eta_t, x_t}) \in \mathbb{S}^2([t, T], \mathbb{R}^d) \times \mathbb{H}^2([t, T], \mathbb{R}^{d \times d})$. Then BSDE (117) also has a unique solution $(U^{\gamma_t, \eta_t, x_t}, V^{\gamma_t, \eta_t, x_t}) \in \mathbb{S}^2([t, T], \mathbb{R}^d) \times \mathbb{H}^2([t, T], \mathbb{R}^{d \times d})$. Moreover, according to the uniqueness of solutions for BSDEs (118), we see $U^{\eta_t, x_t} = U^{\gamma_t, \eta_t, x_t}|_{\gamma=\eta}$. Concerning the regularity of $U^{\gamma_t, \eta_t, x_t}$ and U^{η_t, x_t} with respect to (γ, η, x) , we have

Lemma 3.13. *For any $x, x', \gamma, \gamma' \in \mathbb{D}_{T, d}$, and $\eta, \eta' \in \mathbb{M}_2^D$, we have*

$$(119) \quad \|U^{\eta_t, x_t} - U^{\eta'_t, x'_t}\|_{\mathbb{S}^2} \leq C(\|\eta_t - \eta'_t\|_{\mathbb{S}^2} + \|x_t - x'_t\|), \quad \text{and}$$

$$(120) \quad \|U^{\gamma_t, \eta_t, x_t} - U^{\gamma'_t, \eta'_t, x'_t}\|_{\mathbb{S}^2} \leq C(\|\gamma_t - \gamma'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t}) + \|x_t - x'_t\|),$$

with C only depending on $\|\eta_t\|_{\mathbb{S}^2} + \|\eta'_t\|_{\mathbb{S}^2}$.

Remark 3.14. *Similar to Lemma 3.4, according to estimate (120), $U^{\gamma_t, \mathcal{L}_{\eta_t}, x_t} := U^{\gamma_t, \eta_t, x_t}$ is well-defined.*

Proof. In the following we omit the subscript t and write $(U, V, Y, Z) := (U(r), V(r), Y(r), Z(r))$. Moreover, we only show the proof for (119) since (120) follows by (119) and similar argument. Denote

$$\begin{aligned}
(\Delta U, \Delta V) &:= (U^{\eta, x} - U^{\eta', x'}, V^{\eta, x} - V^{\eta', x'}), \\
\Delta \partial_\mu \Phi &:= \partial_{\mu_t} \Phi(B^\eta, \mathcal{L}_{B^\eta}, B^x) - \partial_{\mu_t} \Phi(B^{\eta'}, \mathcal{L}_{B^{\eta'}}, B^{x'}), \\
\Delta \partial_\mu f &:= \partial_{\mu_t} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{B}^{x_t}) - \partial_{\mu_t} f(\Theta_r^{\eta'}, \mathcal{L}_{\Theta_r^{\eta'}}, \tilde{B}^{x'_t}), \\
\Delta \partial_\nu f^{(1)} &:= \partial_\nu f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) - \partial_\nu f(\Theta_r^{\eta'}, \mathcal{L}_{\Theta_r^{\eta'}}, \tilde{Y}^{x'_t, \mathcal{L}_{\eta'_t}}), \\
\Delta \partial_\nu f^{(2)} &:= \partial_\nu f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{Y}^{\tilde{\eta}_t}) - \partial_\nu f(\Theta_r^{\eta'}, \mathcal{L}_{\Theta_r^{\eta'}}, \tilde{Y}^{\tilde{\eta}'_t}), \\
\Delta \partial_{(y, z)} f &:= \partial_{(y, z)} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) - \partial_{(y, z)} f(\Theta_r^{\eta'}, \mathcal{L}_{\Theta_r^{\eta'}}), \quad \text{and} \\
\Delta \partial_\omega \tilde{Y} &:= \partial_{\omega_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}} - \partial_{\omega_t} \tilde{Y}^{x'_t, \mathcal{L}_{\eta'_t}},
\end{aligned}$$

and we see that $(\Delta U, \Delta V)$ is the unique solution of BSDE

$$\begin{aligned}
(121) \quad \Delta U(s) &= \Delta \partial_\mu \Phi + \int_s^T \tilde{\mathbb{E}}[\Delta \partial_\mu f] dr + \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{Y}^{\tilde{\eta}}) \Delta \tilde{U}] dr \\
&+ \int_s^T (\partial_y f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) \Delta U + \partial_z f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) \Delta V) dr - \int_s^T \Delta V dB(r) \\
&+ \int_s^T \tilde{\mathbb{E}}[(\Delta \partial_\nu f^{(1)}) \partial_\omega \tilde{Y}^{x', \tilde{\eta}'} + (\Delta \partial_\nu f^{(2)}) \tilde{U}^{\tilde{\eta}', x'}] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{Y}^{x, \mathcal{L}_\eta}) \Delta \partial_\omega \tilde{Y}^{x, \mathcal{L}_\eta}] dr \\
&+ \int_s^T \left((\Delta \partial_y f) U^{\eta', x'} + (\Delta \partial_z f) V^{\eta', x'} \right) dr.
\end{aligned}$$

By Lipschitz continuity of $(\partial_{(\mu, \nu, y, z)} f, \partial_\mu \Phi)$, and boundedness of $\partial_{(y, z)} f$, we see that

$$\begin{aligned}
(122) \quad &\|\Delta \partial_\mu \Phi\|_{L^2}^2 + \left\| \int_t^T \tilde{\mathbb{E}}[\Delta \partial_\mu f] dr \right\|_{L^2}^2 + \mathbb{E} \left[\int_t^T \tilde{\mathbb{E}}(|\Delta \partial_\nu f^{(1)}|^2 + |\Delta \partial_\nu f^{(2)}|^2) dr \right] \\
&+ \mathbb{E} \left[\int_t^T |\Delta \partial_y f|^2 dr \right] + \mathbb{E} \left[\int_t^T |\Delta \partial_z f|^2 dr \right] \\
&\leq C(\|\eta_t - \eta'_t\|_{\mathbb{S}^2}^2 + \|x_t - x'_t\|^2).
\end{aligned}$$

Moreover, since $\partial_\omega \tilde{Y}^{x', \tilde{\eta}'}, \tilde{U}^{\eta', x'}, \tilde{Y}^{x, \mathcal{L}_\eta}, \tilde{Y}^{\tilde{\eta}} \in \mathbb{S}^2$, and $V^{\eta', x'} \in \mathbb{H}^2$, from the above estimate and the Cauchy-Schwartz inequality, we obtain

$$\begin{aligned}
(123) \quad &\left\| \int_s^T \tilde{\mathbb{E}} \left[(\Delta \partial_\nu f^{(1)}) \partial_\omega \tilde{Y}^{x', \tilde{\eta}'} + (\Delta \partial_\nu f^{(2)}) \tilde{U}^{\eta', x'} \right] dr \right\|_{L^2} \\
&+ \left\| \int_s^T \left((\Delta \partial_y f) U^{\eta', x'} + (\Delta \partial_z f) V^{\eta', x'} \right) dr \right\|_{L^2} \\
&\leq C(\|\eta_t - \eta'_t\|_{\mathbb{S}^2} + \|x_t - x'_t\|).
\end{aligned}$$

According to estimates given in Lemma 3.7 and boundedness of $\partial_\nu f$, we check that

$$(124) \quad \left\| \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{Y}^{x, \mathcal{L}_\eta}) \Delta \partial_\omega \tilde{Y}^{x, \mathcal{L}_\eta}] dr \right\|_{L^2} \leq C(\|\eta_t - \eta'_t\|_{\mathbb{S}^2} + \|x_t - x'_t\|).$$

Then in view of Lemma 3.1, inequalities (122), (123) and (124), we obtain the desired inequality (119). \square

Concerning the SVD $\partial_{\mu_\tau} Y^{\gamma_t, \eta_t, \cdot}$ of $Y^{\gamma_t, \mathcal{L}_{\eta_t}}$ at $(\tau, t, \mathcal{L}_\eta)$, $\tau \leq t$, in view of Definition 2.9 and BSDE (112), we deduce that it is the unique solution of the following BSDE: for any $x \in \mathbb{D}_{T,d}$,

$$\begin{aligned}
(125) \quad \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_\tau} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_\tau} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\mu_\tau} \tilde{Y}^{\tilde{\eta}_t, x_t}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\omega_\tau} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t}(r) dr \\
&- \int_s^T \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t}(r) dB(r), \quad s \in [t, T],
\end{aligned}$$

where $\partial_{\mu_\tau} Y^{\eta_t, x_t}$ solves the mean-field BSDE below

$$\begin{aligned}
(126) \quad \partial_{\mu_\tau} Y^{\eta_t, x_t}(s) &= \tilde{\mathbb{E}}[\partial_{\mu_\tau} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_\tau} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\mu_\tau} Y^{\eta_t, x_t}(r) dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\omega_\tau} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\mu_\tau} \tilde{Y}^{\tilde{\eta}_t, x_t}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\mu_\tau} Z^{\eta_t, x_t}(r) dr - \int_s^T \partial_{\mu_\tau} Z^{\eta_t, x_t}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

Thanks to Lemma 3.1 again, mean-field BSDE (126) has a unique solution $(\partial_{\mu_\tau} Y^{\eta_t, x_t}, \partial_{\mu_\tau} Z^{\eta_t, x_t}) \in \mathbb{S}^2 \times \mathbb{H}^2$. Then the well-posedness of equation (125) follows similarly. Moreover we have that if $\tau = t$,

$$(127) \quad \partial_{\mu_t} Y^{\gamma_t, \eta_t, x_t} = U^{\gamma_t, \eta_t, x_t}, \quad \partial_{\mu_t} Y^{\eta_t, x_t} = U^{\eta_t, x_t},$$

and $\partial_{\mu_\tau} Y^{\eta_t, x_t} = \partial_{\mu_\tau} Y^{\gamma_t, \mathcal{L}_{\eta_t}, x_t}|_{\gamma=\eta}$. Thus the following lemma follows similarly as that of Lemma 3.13.

Lemma 3.15. *For any $x, x', \gamma, \gamma' \in \mathbb{D}_{T,d}$, and $\eta, \eta' \in \mathbb{M}_2^D$, we have*

$$(128) \quad \|\partial_{\mu_\tau} Y^{\eta_t, x_t} - \partial_{\mu_\tau} Y^{\eta'_t, x'_t}\|_{\mathbb{S}^2} \leq C(\|\eta_t - \eta'_t\|_{\mathbb{S}^2} + \|x_t - x'_t\|), \quad \text{and}$$

$$(129) \quad \|\partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t} - \partial_{\mu_\tau} Y^{\gamma'_t, \eta'_t, x'_t}\|_{\mathbb{S}^2} \leq C(\|\gamma_t - \gamma'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t}) + \|x_t - x'_t\|),$$

with C only depending on $\|\eta_t\|_{\mathbb{S}^2} + \|\eta'_t\|_{\mathbb{S}^2}$.

Recall that $\mathcal{Y}^{\gamma_t, \eta_t, \xi}$ is the solution of BSDE (86) and $U^{\gamma_t, \eta_t, x_t}$ solves equation (117). The following lemma implies that $U^{\gamma_t, \mathcal{L}_{\eta_t}, \cdot} := U^{\gamma_t, \eta_t, \cdot}$ is the derivative of $Y^{\gamma_t, \mathcal{L}_{\eta_t}}$ with respect to \mathcal{L}_{η_t} .

Lemma 3.16. For any $\xi \in L^2(\mathcal{F}_t, \mathbb{R}^d)$, we have

$$(130) \quad \mathcal{Y}^{\gamma_t, \eta_t, \xi}(s) = \bar{E}[U^{\gamma_t, \mathcal{L}_{\eta_t}, \bar{\eta}_t}(s) \bar{\xi}],$$

where $(\bar{\eta}, \bar{\xi})$ is an independent copy of (η, ξ) .

Proof. Substitute $\bar{\eta}_t$ for x_t in equation (118) and multiply the equation by $\bar{\xi}$. Then we take the expectation \bar{E} on both sides of the relation, and obtain

$$(131) \quad \begin{aligned} \bar{E}[U^{\eta_t, \bar{\eta}_t}(s) \bar{\xi}] &= \bar{E} \left[\tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\bar{\eta}_t})] \bar{\xi} \right] + \int_s^T \bar{E} \left[\tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{B}^{\bar{\eta}_t})] \bar{\xi} \right] dr \\ &\quad + \int_s^T \partial_y f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) \bar{E}[U^{\eta_t, \bar{\eta}_t}(r) \bar{\xi}] dr \\ &\quad + \int_s^T \bar{E}[\tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\bar{\eta}_t, \mathcal{L}_{\eta_t}}) \partial_{\omega_t} \tilde{Y}^{\bar{\eta}_t, \mathcal{L}_{\eta_t}}(r)] \bar{\xi}] dr \\ &\quad + \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\bar{\eta}_t}) \bar{E}[\tilde{U}^{\bar{\eta}_t, \bar{\eta}_t}(r) \bar{\xi}]] dr \\ &\quad + \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \bar{E}[V^{\eta_t, \bar{\eta}_t}(r) \bar{\xi}] dr - \int_s^T \bar{E}[V^{\eta_t, \bar{\eta}_t}(r) \bar{\xi}] dB(r). \end{aligned}$$

Since random vectors $(\bar{\eta}, \bar{\xi})$, $(\tilde{B}, \tilde{Y}, \tilde{U})$, (B, Y, Z, η) are mutually independent, and $(B^{\gamma_t}, Y^{\gamma_t, \mathcal{L}_{\eta_t}}, \partial_{x_t} Y^{\gamma_t, \mathcal{L}_{\eta_t}})$ is independent of \mathcal{F}_t , we have

$$\begin{aligned} \bar{E} \left[\tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\bar{\eta}_t})] \bar{\xi} \right] &= \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\bar{\eta}_t}) \bar{\xi}], \\ \bar{E} \left[\tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{B}^{\bar{\eta}_t})] \bar{\xi} \right] &= \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{B}^{\bar{\eta}_t}) \bar{\xi}], \quad \text{and} \\ \bar{E} \left[\tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\bar{\eta}_t, \mathcal{L}_{\eta_t}}) \partial_{x_t} \tilde{Y}^{\bar{\eta}_t, \mathcal{L}_{\eta_t}}(r)] \bar{\xi} \right] &= \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\bar{\eta}_t}) \partial_{x_t} \tilde{Y}^{\bar{\eta}_t, \mathcal{L}_{\eta_t}}(r) \bar{\xi}]. \end{aligned}$$

Then, identity (131) is equivalent to

$$\begin{aligned} \bar{E}[U^{\eta_t, \bar{\eta}_t}(s) \bar{\xi}] &= \tilde{\mathbb{E}}[\partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{\bar{\eta}_t}) \bar{\xi}] + \int_s^T \tilde{\mathbb{E}}[\partial_{\mu_t} f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}, \tilde{B}^{\bar{\eta}_t}) \bar{\xi}] dr \\ &\quad + \int_s^T \partial_y f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) \bar{E}[U^{\eta_t, \bar{\eta}_t}(r) \bar{\xi}] dr - \int_s^T \bar{E}[V^{\eta_t, \bar{\eta}_t}(r) \bar{\xi}] dB(r) \\ &\quad + \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\bar{\eta}_t}) \partial_{\omega_t} \tilde{Y}^{\bar{\eta}_t, \mathcal{L}_{\eta_t}}(r) \bar{\xi}] dr \\ &\quad + \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\bar{\eta}_t}) \bar{E}[\tilde{U}^{\bar{\eta}_t, \bar{\eta}_t}(r) \bar{\xi}]] dr + \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \bar{E}[V^{\eta_t, \bar{\eta}_t}(r) \bar{\xi}] dr, \end{aligned}$$

and therefore $(\bar{E}[U^{\eta_t, \bar{\eta}_t}(s) \bar{\xi}], \bar{E}[V^{\eta_t, \bar{\eta}_t}(s) \bar{\xi}])$ satisfies BSDE (87). In view of the uniqueness of solutions of BSDE (87), we see $\mathcal{Y}^{\eta_t, \xi} = \bar{E}[U^{\eta_t, \bar{\eta}_t}(s) \bar{\xi}]$. Then identity (130) follows from a similar argument. \square

Finally we have main results of this subsection.

Theorem 3.17. *Suppose that (Φ, f) satisfies Assumption **(H1)**. For any $(t, \gamma, \eta) \in [0, T] \times \mathbb{D}_{T,d} \times \mathbb{M}_2^D$, Y^{γ_t, η_t} is Fréchet differentiable with respect to η_t in the sense of (29) and Remark 2.8. Moreover, the Fréchet derivative $D_{\eta_t} Y^{\gamma_t, \eta_t}$ has the following representation: for any $\xi \in L^2(\mathcal{F}_t, \mathbb{R}^d)$,*

$$(132) \quad D_{\eta_t} Y^{\gamma_t, \eta_t}(s)(\xi) = \mathcal{Y}^{\gamma_t, \eta_t, \xi}(s) = \bar{E}[U^{\gamma_t, \mathcal{L}_{\eta_t}, \bar{\eta}_t}(s)\bar{\xi}],$$

where $\mathcal{Y}^{\gamma_t, \eta_t, \xi}$ is the solution of BSDE (86) and $U^{\gamma_t, \mathcal{L}_{\eta_t}, x_t}$, $x \in \mathbb{D}_{T,d}$, is the solution of BSDE (117). In particular, $U^{\gamma_t, \mathcal{L}_{\eta_t}, \cdot}$ is the vertical derivative of $Y^{\gamma_t, \mathcal{L}_{\eta_t}}$ at \mathcal{L}_{η_t} in the sense of (31) and Remark 2.8.

Proof. According to inequality (92) and argument therein, we see that $\mathcal{Y}^{\gamma_t, \eta_t, \cdot}$ is a bounded linear operator from $L^2(\mathcal{F}_t, \mathbb{R}^d)$ to $\mathbb{S}^2([t, T])$. Moreover, in view of Lemma 3.11, for any $\xi \in L^2(\mathcal{F}_t, \mathbb{R}^d)$, $\mathcal{Y}^{\gamma_t, \eta_t, \xi}$ is the Gâteaux derivative of Y^{γ_t, η_t} with respect to η_t . To show $\mathcal{Y}^{\gamma_t, \eta_t, \cdot}$ is the Fréchet derivative of Y^{γ_t, η_t} , it suffices to prove that $\mathcal{Y}^{\gamma_t, \eta_t, \cdot}$ is continuous in $\eta_t \in \mathbb{S}^2([0, t])$ as a linear bounded operator from $L^2(\mathcal{F}_t)$ to $\mathbb{S}^2([t, T])$. Indeed, due to the representation (130) and estimate (120), we have that for any $\eta, \eta' \in \mathbb{M}_2^D$,

$$\|\mathcal{Y}^{\gamma_t, \eta_t, \xi} - \mathcal{Y}^{\gamma_t, \eta'_t, \xi}\|_{\mathbb{S}^2}^2 = \mathbb{E}\|\bar{E}[U^{\gamma_t, \mathcal{L}_{\eta_t}, \bar{\eta}_t}(s)\bar{\xi}] - \bar{E}[U^{\gamma_t, \mathcal{L}_{\eta'_t}, \bar{\eta}'_t}(s)\bar{\xi}]\|^2 \leq C\|\bar{\xi}\|_{L^2}^2\|\bar{\eta}_t - \bar{\eta}'_t\|_{\mathbb{S}^2}^2.$$

Thus we have the following estimate and complete our proof

$$\|\mathcal{Y}^{\gamma_t, \eta_t, \cdot} - \mathcal{Y}^{\gamma_t, \eta'_t, \cdot}\|_{L(L^2(\mathcal{F}_t), \mathbb{S}^2)} \leq C\|\bar{\eta}_t - \bar{\eta}'_t\|_{\mathbb{S}^2}^2.$$

□

For the strong vertical differentiability of Y^{γ_t, η_t} at $(\tau, t, \mathcal{L}_\eta)$, similar as the proof of Lemma 3.16, we have that for any $\xi \in L^2(\mathcal{F}_\tau, \mathbb{R}^d)$,

$$(133) \quad \partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}(s) = \bar{E}[\partial_{\mu_\tau} Y^{\gamma_t, \mathcal{L}_{\eta_t}, \bar{\eta}_t}(s)\bar{\xi}].$$

Moreover, the following proposition implies that $\partial_{\mu_\tau} Y^{\gamma_t, \mathcal{L}_{\eta_t}, \cdot}$ is the SVD of Y^{γ_t, η_t} at $(\tau, t, \mathcal{L}_\eta)$, the proof of which follows from Lemma 3.12, Lemma 3.15 and identity (133).

Proposition 3.18. *For any $(t, \gamma, \eta) \in [0, T] \times \mathbb{D}_{T,d} \times \mathbb{M}_2^D$ and $\tau \in [0, t]$, Y^{γ_t, η_t} is Fréchet differentiable with respect to η_τ in the sense of (32) and Remark 2.10. Moreover, the Fréchet derivative $D_{\eta_\tau} Y^{\gamma_t, \eta_t}$ has the representation: for any $\xi \in L^2(\mathcal{F}_\tau, \mathbb{R}^d)$*

$$(134) \quad D_{\eta_\tau} Y^{\gamma_t, \eta_t}(s)(\xi) = \partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}(s) = \bar{E}[\partial_{\mu_\tau} Y^{\gamma_t, \mathcal{L}_{\eta_t}, \bar{\eta}_t}(s)\bar{\xi}],$$

where $\partial_{\eta_\tau} Y^{\gamma_t, \eta_t, \xi}$ is the solution of BSDE (112) and $\partial_{\mu_\tau} Y^{\gamma_t, \mathcal{L}_{\eta_t}, x_t}$, $x \in \mathbb{D}_{T,d}$, is the solution of BSDE (125).

3.2. Second-order differentiability. Now we consider the second order differentiability of Y^{γ_t, η_t} . To this end, we assume that (Φ, f) satisfies the following assumption.

- (H2)** (i) $\Phi \in \mathcal{C}_{T, \text{lip}}^{2,1,1}(\hat{\mathbb{D}}_{T,d})$; (ii) $f : [0, T] \times \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^D \times \mathcal{P}_2(\mathbb{R}) \rightarrow \mathbb{R}$ satisfies Assumption **(H1)**(ii). Moreover, for any $(t, \omega, y, z, \mu, \nu) \in [0, T] \times \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^D \times \mathcal{P}_2(\mathbb{R})$, $(\partial_y f(t, \omega, \cdot, \cdot, \mu, \nu), \partial_z f(t, \omega, \cdot, \cdot, \mu, \nu))$ is differentiable on $\mathbb{R} \times \mathbb{R}^d$; $(\partial_y f(t, \cdot, y, z, \mu, \nu), \partial_z f(t, \cdot, y, z, \mu, \nu))$ is strongly vertically differentiable at (t, ω) ; for any $\tau \leq t$, $\partial_{\omega_\tau} f(t, \cdot, y, z, \mu, \nu)$ is differentiable at (τ, t, ω) ; $\partial_\nu f(t, \omega, y, z, \mu, \nu, \cdot)$ is differentiable on \mathbb{R} ; for any $\tilde{\omega} \in \mathbb{D}_{T,d}$, $\partial_{\mu_\tau} f(t, \omega, y, z, \mu, \nu, \cdot)$ is differentiable at $(\tau, t, \tilde{\omega})$. All second order derivatives are continuous and $(\partial_y^2, \partial_y \partial_z, \partial_z^2, \partial_{\omega_\tau}^2, \partial_{\tilde{y}} \partial_\nu, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau}) f(t, \cdot)$ is uniformly Lipschitz continuous.

Under Assumption **(H2)**, according to Remark 3.6, we see that $Y^{\gamma_t, \eta_t}(s)$ is almost surely twice vertically differentiable at (t, γ) . Moreover, the second order vertical derivative $\partial_{\omega_t}^2 Y^{\gamma_t, \eta_t}$ satisfies the following linear BSDE

$$\begin{aligned}
& \partial_{\omega_t}^2 Y^{\gamma_t, \eta_t}(s) \\
&= \partial_{\omega_t}^2 \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}) + \int_s^T \partial_{\omega_t}^2 f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr \\
&+ \int_s^T [\partial_y^2 f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) (\partial_{\omega_t} Y^{\gamma_t, \eta_t})^2 + \partial_z^2 f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) (\partial_{\omega_t} Z^{\gamma_t, \eta_t})^2] dr \\
(135) \quad &+ 2 \int_s^T [\partial_{\omega_t} \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\omega_t} Y^{\gamma_t, \eta_t} + \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\omega_t} Y^{\gamma_t, \eta_t} \partial_{\omega_t} Z^{\gamma_t, \eta_t}] dr \\
&+ 2 \int_s^T \partial_{\omega_t} \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\omega_t} Z^{\gamma_t, \eta_t} dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\omega_t}^2 Y^{\gamma_t, \eta_t}(r) dr + \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\omega_t}^2 Z^{\gamma_t, \eta_t}(r) dr \\
&- \int_s^T \partial_{\omega_t}^2 Z^{\gamma_t, \eta_t}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

For the rest of this section, we consider the case of $d = 1$ for sake of simplicity. By boundedness of $\partial_{(y,z)} f$ and Lipschitz continuity of $\partial_{(\omega_t, y, z)} f$, we have similar estimates to those of Lemma 3.7.

Lemma 3.19. *Suppose that (f, Φ) satisfies Assumption **(H2)**, $\partial_{\omega_t} Y^{\gamma_t, \eta_t}$ is almost surely vertically differentiable at γ_t , and the derivative satisfies BSDE (135). Moreover, for any $K > 0$, there exist constants $C_p, C_{K,p}$ such that*

$$(136) \quad \|(\partial_{\omega_t}^2 Y^{\gamma_t, \eta_t}, \partial_{\omega_t}^2 Z^{\gamma_t, \eta_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \leq C_p, \quad \text{and}$$

$$\begin{aligned}
(137) \quad & \|(\partial_{\omega_t}^2 Y^{\gamma_t, \eta_t} - \partial_{\omega_t}^2 Y^{\gamma'_t, \eta'_t}, \partial_{\omega_t}^2 Z^{\gamma_t, \eta_t} - \partial_{\omega_t}^2 Z^{\gamma'_t, \eta'_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \leq C_{K,p} (\|\gamma_t - \gamma'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t})), \\
& \forall (\gamma, \eta), (\gamma', \eta') \in \mathbb{D}_{T,d} \times \mathbb{M}_2^D \text{ such that } \|\mathcal{L}_{\eta_t}\|, \|\mathcal{L}_{\eta'_t}\| \leq K.
\end{aligned}$$

According to Proposition 3.8, Y^{γ_t, η_t} is strongly vertically differentiable at (t, γ) , and the derivative $\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}$ at (τ, t, γ) solves the linear BSDE (82). Similarly, we see that $\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}$ is strongly vertically differentiable at (τ, t, γ) , and moreover, the derivative $\partial_{\omega_\tau}^2 Y^{\gamma_t, \eta_t}$ is the unique solution of BSDE (135) with the derivatives $(\partial_{\omega_t}, \partial_{\omega_t}^2)$ being replaced with $(\partial_{\omega_\tau}, \partial_{\omega_\tau}^2)$. Furthermore, we have estimates (136) and (137) on the solution $(\partial_{\omega_\tau}^2 Y^{\gamma_t, \eta_t}, \partial_{\omega_\tau}^2 Z^{\gamma_t, \eta_t})$, with the derivatives $(\partial_{\omega_t}, \partial_{\omega_t}^2)$ in the left hand sides being replaced with $(\partial_{\omega_\tau}, \partial_{\omega_\tau}^2)$. To apply Theorem 2.15, it remains to study the differentiability of $\partial_{\mu_t} Y^{\gamma_t, \eta_t}(x_t) = U^{\gamma_t, \eta_t, x_t}$ with respect to $x_t \in \mathbb{D}_{T,d}$. Since $U^{\gamma_t, \eta_t, x_t}$ is the unique solution of BSDE (117), by formally taking vertical derivative at (t, x) , we obtain the following

linear BSDE

$$\begin{aligned}
& \partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t}(s) \\
&= \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_t} \partial_{\mu_t} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_t} \partial_{\mu_t} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t}(r) dr \\
(138) \quad &+ \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{y}} \partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}} (\partial_{\omega_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}})^T(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_t}^2 \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\tilde{\omega}_t} \tilde{U}^{\tilde{\eta}_t, x_t}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\tilde{\omega}_t} V^{\gamma_t, \eta_t, x_t}(r) dr - \int_s^T \partial_{\tilde{\omega}_t} V^{\gamma_t, \eta_t, x_t}(r) dB(r), \quad s \in [t, T],
\end{aligned}$$

where $\partial_{\tilde{\omega}_t} U^{\eta_t, x_t}$ solves a mean-field linear BSDE

$$\begin{aligned}
& \partial_{\tilde{\omega}_t} U^{\eta_t, x_t}(s) \\
&= \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_t} \partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_t} \partial_{\mu_t} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\tilde{\omega}_t} U^{\eta_t, x_t}(r) dr \\
(139) \quad &+ \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{y}} \partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}} (\partial_{\omega_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}})^T(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_t}^2 \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\tilde{\omega}_t} \tilde{U}^{\tilde{\eta}_t, x_t}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\tilde{\omega}_t} V^{\eta_t, x_t}(r) dr - \int_s^T \partial_{\tilde{\omega}_t} V^{\eta_t, x_t}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

Lemma 3.20. *There exist unique solutions $(\partial_{\tilde{\omega}_t} U^{\eta_t, x_t}, \partial_{\tilde{\omega}_t} V^{\eta_t, x_t}) \in \mathbb{S}^2([t, T]) \times \mathbb{H}^2([t, T])$ and $(\partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t}, \partial_{\tilde{\omega}_t} V^{\gamma_t, \eta_t, x_t}) \in \mathbb{S}^p([t, T]) \times \mathbb{H}^p([t, T])$ of equations (139) and (138), respectively. Moreover, $\partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t}$ is the vertical derivative of $U^{\gamma_t, \eta_t, x_t}$ at (t, x) , and for any $K > 0$,*

$$\begin{aligned}
& \|(\partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t}, \partial_{\tilde{\omega}_t} V^{\gamma_t, \eta_t, x_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \leq C_p, \\
(140) \quad & \|(\partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t} - \partial_{\tilde{\omega}_t} U^{\gamma'_t, \eta'_t, x'_t}, \partial_{\tilde{\omega}_t} V^{\gamma_t, \eta_t, x_t} - \partial_{\tilde{\omega}_t} V^{\gamma'_t, \eta'_t, x'_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \\
& \leq C_{K,p} (\|x_t - x'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t}) + \|\gamma_t - \gamma'_t\|), \\
& \quad \forall (\gamma, \eta, x), (\gamma', \eta', x') \in \mathbb{D}_{T,d} \times \mathbb{M}_{2,D}^D \times \mathbb{D}_{T,d} \text{ such that } \|\mathcal{L}_{\eta_t}\|, \|\mathcal{L}_{\eta'_t}\| \leq K,
\end{aligned}$$

with some constants C_p and $C_{K,p}$.

Proof. To show the well-posedness of (139), according to Lemma 3.1, it remains to check the following terms belong to $L^2(\mathcal{F}_T)$,

$$\begin{aligned} & \left| \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_t} \partial_{\mu_t} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] \right|, \int_t^T \left| \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_t} \partial_{\mu_t} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] \right| dr, \\ & \int_t^T \left| \tilde{\mathbb{E}}[\partial_{\nu} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_t}^2 \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] \right| dr, \quad \text{and} \\ & \int_t^T \left| \tilde{\mathbb{E}}[\partial_{\tilde{y}} \partial_{\nu} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}} (\partial_{\tilde{\omega}_t} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r))^T] \right|, \end{aligned}$$

which follows easily by the boundedness of $(\partial_{\tilde{\omega}_t} \partial_{\mu_t} \Phi, \partial_{\tilde{\omega}_t} \partial_{\mu_t} f, \partial_{\nu} f, \partial_{\tilde{y}} \partial_{\nu} f)$, Lemmas 3.7 and 3.19. Moreover, we see

$$(141) \quad \|\partial_{\tilde{\omega}_t} U^{\eta_t, x_t}\|_{\mathbb{S}^2} + \|\partial_{\tilde{\omega}_t} V^{\eta_t, x_t}\|_{\mathbb{H}^2} \leq C_p, \quad \text{and}$$

$$(142) \quad \|\partial_{\tilde{\omega}_t} U^{\eta_t, x_t} - \partial_{\tilde{\omega}_t} U^{\eta'_t, x'_t}\|_{\mathbb{S}^2} + \|\partial_{\tilde{\omega}_t} V^{\eta_t, x_t} - \partial_{\tilde{\omega}_t} V^{\eta'_t, x'_t}\|_{\mathbb{H}^2} \leq C_{K,p}(\|x_t - x'_t\| + \|\eta_t - \eta'_t\|_{\mathbb{S}^2}).$$

Concerning the well-posedness of (138), since $(\partial_{\tilde{\omega}_t} U^{\eta_t, x_t}, \partial_{\tilde{\omega}_t} V^{\eta_t, x_t}) \in \mathbb{S}^2 \times \mathbb{H}^2$, we have

$$\int_t^T \left| \tilde{\mathbb{E}}[\partial_{\nu} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\tilde{\omega}_t} \tilde{U}^{\tilde{\eta}_t, x_t}(r)] \right| dr \in L^p(\mathcal{F}_T),$$

and therefore, there exists a unique solution $(\partial_{\tilde{\omega}_t} U^{\gamma_t, \eta_t, x_t}, \partial_{\tilde{\omega}_t} V^{\gamma_t, \eta_t, x_t}) \in \mathbb{S}^p \times \mathbb{H}^p$. In view of the boundedness of $(\partial_{\tilde{\omega}_t} \partial_{\mu_t} \Phi, \partial_{\tilde{\omega}_t} \partial_{\mu_t} f, \partial_{\tilde{y}} \partial_{\nu} f, \partial_{\tilde{y}} f, \partial_z f)$ and standard estimate for BSDEs, we have our desired estimates. \square

Recall that $\partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}$ is the solution of BSDE (125). To prove the strong vertical differentiability of $\partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}$ at (τ, t, x) , we consider BSDE

$$\begin{aligned} & \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}(s) \\ &= \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\gamma_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}, \tilde{B}^{x_t})] dr \\ &+ \int_s^T \partial_{\tilde{y}} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}) \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}(r) dr \\ &+ \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{y}} \partial_{\nu} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_\tau} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}} (\partial_{\tilde{\omega}_\tau} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r))^T] dr \\ (143) \quad &+ \int_s^T \tilde{\mathbb{E}}[\partial_{\nu} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_\tau}^2 \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\ &+ \int_s^T \tilde{\mathbb{E}}[\partial_{\nu} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} \tilde{Y}^{\tilde{\eta}_t, x_t}(r)] dr \\ &+ \int_s^T \partial_z f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\gamma_t, \eta_t}}) \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t}(r) dr \\ &- \int_s^T \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t}(r) dB(r), \quad s \in [t, T], \end{aligned}$$

where $\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\eta_t, x_t}$ solves the following mean-field linear BSDE

$$\begin{aligned}
& \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\eta_t, x_t}(s) \\
&= \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} \Phi(B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}, \tilde{B}^{x_t})] + \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{B}^{x_t})] dr \\
&+ \int_s^T \partial_y f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\eta_t, x_t}(r) dr \\
(144) \quad &+ \int_s^T \tilde{\mathbb{E}}[\partial_{\tilde{y}} \partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_\tau} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}} (\partial_{\tilde{\omega}_\tau} \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r))^T] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}) \partial_{\tilde{\omega}_\tau}^2 \tilde{Y}^{x_t, \mathcal{L}_{\eta_t}}(r)] dr \\
&+ \int_s^T \tilde{\mathbb{E}}[\partial_\nu f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}, \tilde{Y}^{\tilde{\eta}_t}) \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} \tilde{Y}^{\tilde{\eta}_t, x_t}(r)] dr \\
&+ \int_s^T \partial_z f(\Theta_r^{\eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\eta_t, x_t}(r) dr - \int_s^T \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\eta_t, x_t}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

Then we have the following lemma via a similar proof of Lemma 3.20.

Lemma 3.21. *There exists a unique solution $(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t})$ of BSDE (143). Moreover, $\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}$ is the SVD of $\partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}$ at (τ, t, x) , and for any $K > 0$,*

$$\begin{aligned}
(145) \quad & \|(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t}, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \leq C_p, \\
& \|(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \eta_t, x_t} - \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma'_t, \eta'_t, x'_t}, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\gamma_t, \eta_t, x_t} - \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Z^{\gamma'_t, \eta'_t, x'_t})\|_{\mathbb{S}^p \times \mathbb{H}^p} \\
& \leq C_{K,p} (\|x_t - x'_t\| + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t}) + \|\gamma_t - \gamma'_t\|), \\
& \quad \forall (\gamma, \eta, x), (\gamma', \eta', x') \in \mathbb{D}_{T,d} \times \mathbb{M}_2^D \times \mathbb{D}_{T,d} \text{ such that } \|\mathcal{L}_{\eta_t}\|, \|\mathcal{L}_{\eta'_t}\| \leq K,
\end{aligned}$$

with some constants C_p and $C_{K,p}$.

4. SOLUTIONS OF SEMILINEAR PATH-DEPENDENT MASTER EQUATIONS

4.1. The decoupling field and its regularity. For any $(t, \gamma, \mu) \in \hat{\mathbb{D}}_{T,d}$, recall that Y^{γ_t, η_t} solves the path-dependent BSDE

$$\begin{aligned}
(146) \quad & Y^{\gamma_t, \eta_t}(s) = \Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\gamma_t}}) + \int_s^T f(B_r^{\gamma_t}, Y^{\gamma_t, \eta_t}(r), Z^{\gamma_t, \eta_t}(r), \mathcal{L}_{B_r^{\gamma_t}}, \mathcal{L}_{Y^{\eta_t}(r)}) dr \\
& - \int_s^T Z^{\gamma_t, \eta_t}(r) dB(r), \quad s \in [t, T],
\end{aligned}$$

where $Y^{\eta_t} = Y^{\gamma_t, \eta_t}|_{\gamma=\eta}$ is the unique solution of the mean-field BSDE

$$\begin{aligned}
(147) \quad & Y^{\eta_t}(s) = \Phi(B_T^{\eta_t}, \mathcal{L}_{B_T^{\eta_t}}) + \int_s^T f(B_r^{\eta_t}, Y^{\eta_t}(r), Z^{\eta_t}(r), \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(r)}) dr \\
& - \int_s^T Z^{\eta_t}(r) dB(r), \quad s \in [t, T].
\end{aligned}$$

For any $(t, \gamma, \mu) \in \hat{\mathbb{D}}_{T,d}$ with $\mu = \mathcal{L}_\eta$, define the decoupling field

$$(148) \quad u(t, \gamma, \mu) := Y^{\gamma_t, \mathcal{L}_{\eta_t}}(t).$$

In view of classical BSDE theory, we see that $u \in \mathcal{D}$ and it is non-anticipative. On the other hand, for any $v \geq t$, $(\omega, y, z) \in \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d$, denote

$$(149) \quad \hat{\Phi}_{\mu_t}(\omega_T) := \Phi(\omega_T, \mathcal{L}_{B_T^{\eta_t}}), \quad \hat{f}_{\mu_t}(v, \omega, y, z) := f(v, \omega, y, z, \mathcal{L}_{B_v^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(v)}).$$

Then $Y^{\gamma v, \mathcal{L}^{\eta_t}}$ is the unique solution of the following (path-dependent) BSDE

$$(150) \quad Y^{\gamma v, \eta_t}(s) = \hat{\Phi}_{\mu_t}(B_T^{\gamma v}) + \int_s^T \hat{f}_{\mu_t}(r, B_r^{\gamma v}, Y^{\gamma v, \eta_t}(r), Z^{\gamma v, \eta_t}(r)) dr - \int_s^T Z^{\gamma v, \eta_t}(r) dB(r), \quad s \geq v.$$

According to Remark 3.6 and [41, Theorem 3.9], there exists a non-anticipative mapping $\hat{u}_{\mu_t} : [v, T] \times \mathbb{D}_{T,d} \rightarrow \mathbb{R}$, such that for any $s \geq v$,

$$(151) \quad \hat{u}_{\mu_t}(s, B^{\gamma v}) = Y^{\gamma v, \mu_t}(s), \quad \partial_{\gamma v} \hat{u}_{\mu_t}(s, B^{\gamma v}) = Z^{\gamma v, \mu_t}(s).$$

Moreover, \hat{u}_{μ_t} is the classical solution of the following semilinear PPDE

$$(152) \quad \begin{cases} \partial_v \hat{u}_{\mu_t}(v, \gamma) + \frac{1}{2} \text{Tr} [\partial_{\omega_v}^2 \hat{u}_{\mu_t}(v, \gamma)] + \hat{f}_{\mu_t}(v, \gamma, \hat{u}_{\mu_t}(v, \gamma), \partial_{\omega_v} \hat{u}_{\mu_t}(v, \gamma)) = 0, \\ \hat{u}_{\mu_t}(T, \gamma) = \hat{\Phi}(\gamma), \quad v \geq t. \end{cases}$$

Indeed, denote $\hat{\eta} := B^{\eta_t}$, and we have

$$(153) \quad \hat{u}_{\mu_t}(v, \gamma) = Y^{\gamma v, \hat{\eta}_v}(v).$$

Concerning the relation among $Y^{\gamma t, \eta_t}$, $u(t, \gamma, \mu)$ and $\hat{u}_{\mu_t}(v, \gamma)$, we have

Proposition 4.1. *For any $(t, \gamma, \mu) \in \hat{\mathbb{D}}_{T,d}$ and $s \geq t$,*

$$(154) \quad u(t, \gamma, \mu) = \hat{u}_{\mu_t}(t, \gamma),$$

$$(155) \quad u(s, \omega_s^{\gamma t}, \mathcal{L}_{B_s^{\eta_t}}) = \hat{u}_{\mu_t}(s, \omega_s^{\gamma t}), \quad \forall \omega \in \mathbb{C}_{T,d}, \quad \text{and}$$

$$(156) \quad u(s, B_s^{\eta_t}, \mathcal{L}_{B_s^{\eta_t}}) = Y^{\eta_t}(s).$$

Proof. The first identity follows immediately from (148) and (151). By the uniqueness of BSDE (147), we see that for any $t \leq v \leq s$,

$$(157) \quad (B_s^{B_v^{\eta_t}}, Y^{B_v^{\eta_t}}(s)) = (B_s^{\eta_t}, Y^{\eta_t}(s)),$$

and in particular

$$(158) \quad \mathcal{L}_{(B_s^{B_v^{\eta_t}}, Y^{B_v^{\eta_t}}(s))} = \mathcal{L}_{(B_s^{\eta_t}, Y^{\eta_t}(s))}.$$

Then in view of the uniqueness of solutions of BSDEs (146) and (151), we have

$$(159) \quad \hat{u}_{\mu_t}(s, B_s^{\gamma t}) = \hat{u}_{\mathcal{L}_{B_v^{\eta_t}}}(s, B_s^{\gamma t}).$$

In particular if $v = s$, $\hat{u}_{\mu_t}(s, B_s^{\gamma t}) = \hat{u}_{\mathcal{L}_{B_s^{\eta_t}}}(s, B_s^{\gamma t})$. On the other hand, by relation (154), we have

$$(160) \quad \hat{u}_{\mathcal{L}_{B_s^{\eta_t}}}(s, B_s^{\gamma t}) = u(s, B_s^{\gamma t}, \mathcal{L}_{B_s^{\eta_t}}),$$

and thus (155) in view of the continuity of $u(t, \omega, \mu)$ in $\omega \in \mathbb{C}_{T,d}$ and the support theorem for diffusion processes. Finally, since

$$(161) \quad Y^{\eta_t}(s) = Y^{\gamma t, \eta_t}(s)|_{\gamma=\eta} = \hat{u}_{\mu_t}(s, B_s^{\gamma t})|_{\gamma=\eta} = \hat{u}_{\mu_t}(s, B_s^{\eta_t}),$$

identity (156) follows from (155). □

Proposition 4.2. *Suppose that (f, Φ) satisfies Assumption (H2). The decoupling field u given by (148) belongs to $\mathcal{C}_{s,p}^{0,2,1,1}(\hat{\mathbb{D}}_{T,d})$.*

Proof. According to Lemma 3.4, $u(t, \gamma, \mu) = Y^{\gamma_t, \eta_t}(t)$ satisfies the polynomial growth condition in the sense of (38). To prove that $u \in \mathcal{C}_p$, we only need to show its continuity in $(t, \gamma, \mu) \in \hat{\mathbb{D}}_{T,d}$. For any $(t, \gamma, \mu), (t', \gamma', \mu') \in \hat{\mathbb{D}}_{T,d}$, without loss of generality, assume $t \geq t'$. We have

$$\begin{aligned}
(162) \quad & |u(t, \gamma, \mu) - u(t', \gamma', \mu')| = |Y^{\gamma_t, \eta_t}(t) - Y^{\gamma'_{t'}, \eta'_{t'}}(t')| \\
& \leq \mathbb{E}|Y^{\gamma_t, \eta_t}(t) - Y^{\gamma'_{t'}, \eta'_{t'}}(t)| + \mathbb{E}|Y^{\gamma'_{t'}, \eta'_{t'}}(t) - Y^{\gamma'_{t'}, \eta'_{t'}}(t')| \\
& \leq \mathbb{E}|Y^{\gamma_t, \eta_t}(t) - Y^{\gamma'_{t'}, \eta'_{t'}}(t)| + \mathbb{E} \left| \int_{t'}^t f(\Theta_r^{\gamma'_{t'}, \eta'_{t'}}, \mathcal{L}_{\Theta_r^{\eta'_{t'}}}) dr - \int_{t'}^t Z'(r) dB(r) \right| \\
& \leq \mathbb{E}|Y^{\gamma_t, \eta_t}(t) - Y^{\gamma'_{t'}, \eta'_{t'}}(t)| + C(1 + \|\gamma_t\| + \|\eta_t\|_{\mathbb{S}^2})(t - t')^{\frac{1}{2}}.
\end{aligned}$$

It remains to prove $\mathbb{E}|Y^{\gamma_t, \eta_t}(t) - Y^{\gamma'_{t'}, \eta'_{t'}}(t)| \rightarrow 0$ as $(t, \gamma, \mu) \rightarrow (t', \gamma', \mu')$. Set $Y' := Y^{\gamma'_{t'}, \eta'_{t'}}$, $Y := Y^{\gamma_t, \eta_t}$, and $(\delta Y, \delta Z) := (Y - Y', Z - Z')$, and omit subscripts t and t' . Then $(\delta Y, \delta Z)$ is the unique solution of BSDE

$$\begin{aligned}
\delta Y(s) &= \Phi(B^\gamma, \mathcal{L}_{B^\gamma}) - \Phi(B^{\gamma'}, \mathcal{L}_{B^{\gamma'}}) + \int_s^T [f(\Theta_r^{\gamma, \eta}, \mathcal{L}_{\Theta_r^\eta}) - f(\Theta_r^{\gamma', \eta'}, \mathcal{L}_{\Theta_r^{\eta'}})] dr - \int_s^T \delta Z(r) dB(r) \\
&=: \delta \Phi + \int_s^T \left(a_r \delta Y(r) + b_r \delta Z(r) + \tilde{\mathbb{E}}[\tilde{c}_r \delta \tilde{Y}(r)] + \delta h_r \right) dr - \int_s^T \delta Z(r) dB(r),
\end{aligned}$$

where

$$\begin{aligned}
a_r &:= \int_0^1 \partial_y f(B_r^\gamma, Y' + \theta(Y - Y'), Z, \mathcal{L}_{\Theta_r^\eta}) d\theta, \\
b_r &:= \int_0^1 \partial_z f(B_r^\gamma, Y', Z + \theta(Z - Z'), \mathcal{L}_{\Theta_r^\eta}) d\theta, \\
\tilde{c}_r &:= \int_0^1 \partial_\nu f(B_r^\gamma, Y', Z', \mathcal{L}_{B_r^\eta}, \mathcal{L}_{Y^{\eta'} + \theta(Y^\eta - Y^{\eta'})}, \tilde{Y}^{\eta'} + \theta(\tilde{Y}^\eta - \tilde{Y}^{\eta'})) d\theta, \quad \text{and} \\
\delta h_r &:= f(B_r^\gamma, Y', Z', \mathcal{L}_{B_r^\eta}, \mathcal{L}_{Y^{\eta'}}) - f(B_r^{\gamma'}, Y', Z', \mathcal{L}_{B_r^{\eta'}}, \mathcal{L}_{Y^{\eta'}}).
\end{aligned}$$

Using Lemma 3.1 to the above BSDE, we have

$$\begin{aligned}
\|(\delta Y, \delta Z)\|_{\mathbb{S}^2 \times \mathbb{H}^2}^2 &\leq C(\|\delta \Phi\|_{L^2}^2 + \|\int_t^T |\delta h_r| dr\|_{L^2}^2) \\
&\leq C(\|B^\gamma - B^{\gamma'}\|_{L^2}^2 + W_2(\mathcal{L}_{B^\eta}, \mathcal{L}_{B^{\eta'}})^2) \\
&\leq C(\|\gamma_t - \gamma'_{t'}\|^2 + W_2(\mu_t, \mu'_{t'})^2 + (t - t')),
\end{aligned}$$

where C depends on $\|\mu_t\| + \|\mu'_{t'}\|$, $\|\gamma_t\|$ and $\|\gamma'_{t'}\|$, and thus the continuity of u in view of (162).

Since $u(t, \gamma, \mu) = Y^{\gamma_t, \eta_t}(t) = \hat{u}_{\mu_t}(t, \gamma)$, according to Proposition 3.8, we see that for any $\tau \leq t$, $u(t, \gamma, \mu)$ is twice strongly vertically differentiable at (τ, t, γ) , and moreover, $\partial_{\omega_\tau} u(t, \gamma, \mu) = \partial_{\omega_\tau} Y^{\gamma_t, \eta_t}(t)$ satisfies the polynomial growth condition. To show $u(t, \gamma, \mu) \in \mathcal{C}_{s,p}^{0,1,0}$, we only need to prove that $\partial_{\gamma_\tau} u(t, \gamma, \mu)$ is continuous at any (τ, t, γ, μ) . Indeed, for any (τ, t, γ, μ) and $(\tau', t', \gamma', \mu')$ with $\tau \leq t$, $\tau' \leq t'$, denote solutions of equation (82) corresponding to parameters (τ, t, γ, μ) and $(\tau', t', \gamma', \mu')$ by

$$(163) \quad (\mathcal{Y}, \mathcal{Z}) := (\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}, \partial_{\omega_\tau} Z^{\gamma_t, \eta_t}), \quad (\mathcal{Y}', \mathcal{Z}') := (\partial_{\omega_{\tau'}} Y^{\gamma'_{t'}, \eta'_{t'}}, \partial_{\omega_{\tau'}} Z^{\gamma'_{t'}, \eta'_{t'}}).$$

Without loss of generality, let $t \geq t'$. By inserting the term $\mathcal{Y}'(t)$ and applying Proposition 3.8, we have

$$\begin{aligned} & |\partial_{\omega_\tau} u(t, \gamma, \mu) - \partial_{\omega_{\tau'}} u(t', \gamma', \mu')| \\ &= |\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}(t) - \partial_{\omega_{\tau'}} Y^{\gamma_{t'}, \eta_{t'}}(t')| \\ &\leq \mathbb{E} |\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}(t) - \partial_{\omega_{\tau'}} Y^{\gamma_{t'}, \eta_{t'}}(t)| + \mathbb{E} |\partial_{\omega_{\tau'}} Y^{\gamma_{t'}, \eta_{t'}}(t) - \partial_{\omega_{\tau'}} Y^{\gamma_{t'}, \eta_{t'}}(t')| \\ &\leq \mathbb{E} |\partial_{\omega_\tau} Y^{\gamma_t, \eta_t}(t) - \partial_{\omega_{\tau'}} Y^{\gamma_{t'}, \eta_{t'}}(t)| + C(t - t')^{\frac{1}{2}}. \end{aligned}$$

Set $(\delta\mathcal{Y}, \delta\mathcal{Z}) := (\mathcal{Y} - \mathcal{Y}', \mathcal{Z} - \mathcal{Z}')$. We see that $(\delta\mathcal{Y}, \delta\mathcal{Z})$ is the unique solution of the following BSDE

$$\begin{aligned} \delta\mathcal{Y}(s) &= [\partial_{\omega_\tau} \Phi - \partial_{\omega_{\tau'}} \Phi'] + \int_s^T [\partial_{\omega_\tau} f - \partial_{\omega_{\tau'}} f'] dr + \int_s^T \partial_y f \delta\mathcal{Y}(r) dr \\ &+ \int_s^T (\partial_y f - \partial_y f') \mathcal{Y}'(r) dr + \int_s^T (\partial_z f)^T \delta\mathcal{Z}(r) dr + \int_s^T (\partial_z f - \partial_z f')^T \mathcal{Z}'(r) dr \\ &- \int_s^T \delta\mathcal{Z} dB(r), \end{aligned}$$

where

$$\begin{aligned} \partial_{\omega_\tau} \Phi &:= \partial_{\omega_\tau} \Phi(B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}), \quad \partial_{\omega_{\tau'}} \Phi' := \partial_{\omega_{\tau'}} \Phi(B^{\gamma_{t'}}, \mathcal{L}_{B^{\eta_{t'}}}), \quad \partial_{(\omega_\tau, y, z)} f := \partial_{(\omega_\tau, y, z)} f(\Theta^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta^{\eta_t}}), \\ \text{and } \partial_{(\omega_{\tau'}, y, z)} f' &:= \partial_{(\omega_{\tau'}, y, z)} f(\Theta^{\gamma_{t'}, \eta_{t'}}, \mathcal{L}_{\Theta^{\eta_{t'}}}). \end{aligned}$$

In view of estimates in Lemmas 3.1 and 3.7, using Cauchy inequality, we have

$$\begin{aligned} |\mathcal{Y}(t) - \mathcal{Y}'(t)|^2 &\leq \left[\|\partial_{\omega_\tau} \Phi - \partial_{\omega_{\tau'}} \Phi'\|_{L^2}^2 + \left\| \int_t^T [\partial_{\omega_\tau} f - \partial_{\omega_{\tau'}} f'] dr \right\|_{L^2}^2 \right. \\ &\quad \left. + \left\| \int_t^T [\partial_y f - \partial_y f'] \mathcal{Y}' dr \right\|_{L^2}^2 + \left\| \int_t^T [\partial_z f - \partial_z f'] \mathcal{Z}' dr \right\|_{L^2}^2 \right] \\ &\leq \left[\|\partial_{\omega_\tau} \Phi - \partial_{\omega_{\tau'}} \Phi'\|_{L^2}^2 + \mathbb{E} \left[\int_t^T |\partial_{\omega_\tau} f - \partial_{\omega_{\tau'}} f'|^2 dr \right] \right. \\ &\quad \left. + \mathbb{E} \left[\int_t^T |\partial_y f - \partial_y f'|^4 dr \right] + \mathbb{E} \left[\int_t^T |\partial_z f - \partial_z f'|^4 dr \right] \right]. \end{aligned}$$

Then the desired continuity follows from that of $(\partial_{\omega_\tau} \Phi, \partial_{(\omega_\tau, y, z)} f)$ and the bounded convergence theorem. Similarly, we have $\partial_{\omega_\tau} u \in \mathcal{C}_{s,p}^{0,1,0}$ and therefore $u \in \mathcal{C}_{s,p}^{0,2,0}$.

For the differentiability with respect to the measure variable, according to Lemmas 3.12 and 3.21, we have that for any $x \in \mathbb{D}_{T,d}$,

$$(164) \quad \partial_{\mu_\tau} u(t, \gamma, \mu, x) = \partial_{\mu_\tau} Y^{\gamma_t, \mu_t, x_t}(t), \quad \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} u(t, \gamma, \mu, x) = \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \mu_t, x_t}(t).$$

Here, $\partial_{\mu_\tau} Y^{\gamma_t, \mu_t, x_t}$ solves BSDE (125) and $\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} Y^{\gamma_t, \mu_t, x_t}$ solves BSDE (143). Following a similar argument as above, we see that $u(t, \gamma, \mu) \in \mathcal{C}_{s,p}^{0,2,1,1}$. \square

4.2. Classical solutions of path-dependent master equations. In this subsection we consider well-posedness of the path-dependent semilinear parabolic master equation

$$(165) \quad \begin{cases} \partial_t u(t, \gamma, \mu) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \gamma, \mu)] + \frac{1}{2} \text{Tr} \left[\int_{\mathbb{C}_{T,d}} \partial_{\tilde{\omega}} \partial_\mu u(t, \gamma, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) \right] \\ \quad + f(t, \gamma, u(t, \gamma, \mu), \partial_\omega u(t, \gamma, \mu), \mu, \mathcal{L}_{u(t, W^\mu, \mu)}) = 0, \\ u(T, \gamma, \mu) = \Phi(\gamma_T, \mu_T), \quad (t, \gamma, \mu) \in [0, T] \times \mathbb{C}_{T,d} \times \mathcal{P}_2^C, \end{cases}$$

where W^μ is the canonical process under μ . In applications, (γ, μ) takes values in $\mathbb{C}_{T,d} \times \mathcal{P}_2^C$. Thus we need to give a description of equation (165) restricted on $\mathbb{C}_{T,d} \times \mathcal{P}_2^C$. Denote by $\hat{\mathbb{C}}_{T,d}$ the product space $[0, T] \times \mathbb{C}_{T,d} \times \mathcal{P}_2^C$, and for a $n \times n$ matrix A , we write $\text{Sym}(A) := \frac{1}{2}(A + A^T)$. For any $f \in \mathcal{D}$, we write $(\partial_{\omega_\tau}, \partial_{\tilde{\omega}_\tau}^2, \partial_{\mu_\tau}, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau})f := (\partial_{\omega_\tau} f, \partial_{\tilde{\omega}_\tau}^2 f, \partial_{\mu_\tau} f, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f)$ if the right hand side exists.

Definition 4.3. Denote by $\mathcal{C}_{s,p}^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$ the set of functionals $f : \hat{\mathbb{C}}_{T,d} \rightarrow \mathbb{R}$ such that there exists an extension $F \in \mathcal{C}_{s,p}^{1,2,1,1}(\hat{\mathbb{D}}_{T,d})$ with $f = F$ on $\hat{\mathbb{C}}_{T,d}$. In this case, for any $(t, \omega, \mu, \tilde{\omega}) \in \hat{\mathbb{C}}_{T,d} \times \mathbb{C}_{T,d}$ and $\tau \leq t$, we write

$$(166) \quad \begin{aligned} \partial_t f(t, \omega, \mu) &:= \partial_t F(t, \omega, \mu), \quad (\partial_{\omega_\tau}, \partial_{\tilde{\omega}_\tau}^2) f(t, \omega_\tau, \mu) := (\partial_{\omega_\tau}, \partial_{\tilde{\omega}_\tau}^2) F(t, \omega_\tau, \mu), \\ \text{and} \quad (\partial_{\mu_\tau}, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau}) f(t, \omega, \mu_\tau, \tilde{\omega}_\tau) &:= (\partial_{\mu_\tau}, \partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau}) F(t, \omega, \mu_\tau, \tilde{\omega}_\tau). \end{aligned}$$

Notations such as $\mathcal{C}^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$, $\mathcal{C}_p^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$ and $\mathcal{C}_{s,p}^{0,2,1,1}(\hat{\mathbb{C}}_{T,d})$ are defined in a similar way.

In view of Itô-Dupire formulas given in Theorem 2.15 and Corollary 2.16, we have

Corollary 4.4. For any $(t, \gamma, \mu, \eta) \in \hat{\mathbb{C}}_{T,d} \times \mathbb{M}_2^C$, X and X' are diffusion processes given by (41) and (42) respectively.

(i) Suppose that $f \in \mathcal{C}_s^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$ ($\mathcal{C}^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$, resp.). For any $(\tau, \tilde{\omega}) \in [0, t] \times \mathbb{C}_{T,d}$, derivatives

$$\partial_t f(t, \omega, \mu), \quad (\partial_{\omega_\tau}, \partial_{\tilde{\omega}_\tau}^2) f(t, \omega_\tau, \mu), \quad \partial_{\mu_\tau} f(t, \omega, \mu_\tau, \tilde{\omega}_\tau), \quad \text{and} \quad \text{Sym}(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(t, \omega, \mu_\tau, \tilde{\omega}_\tau)),$$

($\partial_t f$, $(\partial_\omega, \partial_\omega^2) f$, $\partial_\mu f$, and $\text{Sym}(\partial_{\tilde{\omega}} \partial_\mu f)$, resp.) defined as in (166) do not depend on the choice of the extended functional.

(ii) Suppose that $f \in \mathcal{C}_p^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$. For any $s \geq t$, we have

$$(167) \quad \begin{aligned} & f(s, X, \mathcal{L}_{X'}) - f(t, \gamma, \mathcal{L}_\eta) \\ &= \int_t^s \partial_r f(r, X, \mathcal{L}_{X'}) dr + \int_t^s \partial_\omega f(r, X, \mathcal{L}_{X'}) dX(r) \\ & \quad + \frac{1}{2} \int_t^s \text{Tr} [\partial_\omega^2 f(r, X, \mathcal{L}_{X'}) d\langle X \rangle(r)] + \mathbb{E}^{\tilde{P}'} \left[\int_t^s \partial_\mu f(r, X, \mathcal{L}_{X'}, \tilde{X}') d\tilde{X}'(r) \right] \\ & \quad + \frac{1}{2} \mathbb{E}^{\tilde{P}'} \int_t^s \text{Tr} [\partial_{\tilde{\omega}} \partial_\mu f(r, X, \mathcal{L}_{X'}, \tilde{X}') \tilde{d}(r) \tilde{d}(r)^T] dr. \end{aligned}$$

(iii) Suppose that $f \in \mathcal{C}_{s,p}^{0,2,1,1}(\hat{\mathbb{C}}_{T,d})$. For any $t \leq s \leq s'$, we have the partial Itô-Dupire formula

$$\begin{aligned}
& f(s', X_s, \mathcal{L}_{X'_s}) - f(s', \gamma_t, \mathcal{L}_{\eta_t}) \\
&= \int_t^s \partial_{\omega_r} f(s', X_r, \mathcal{L}_{X'_r}) dX(r) \\
(168) \quad &+ \frac{1}{2} \int_t^s \text{Tr} [\partial_{\omega_r}^2 f(s', X_r, \mathcal{L}_{X'_r}) d\langle X \rangle(r)] + \mathbb{E}^{\tilde{P}'} \left[\int_t^s \partial_{\mu_r} f(s', X_r, \mathcal{L}_{X'_r}, \tilde{X}'_r) d\tilde{X}'(r) \right] \\
&+ \frac{1}{2} \mathbb{E}^{\tilde{P}'} \int_t^s \text{Tr} [\partial_{\tilde{\omega}_r} \partial_{\mu_r} f(s', X_r, \mathcal{L}_{X'_r}, \tilde{X}'_r) \tilde{d}(r) \tilde{d}(r)^T] dr.
\end{aligned}$$

Proof. Since for any $n \times n$ matrix A and symmetric $n \times n$ matrix B , $\text{Tr}[AB]$ depends only on $\text{Sym}(A)$, Assertions (ii) and (iii) follow directly from Assertion (i), Theorem 2.15 and Corollary 2.16.

To end the proof, we only need to show (i). Indeed, the uniqueness of $\partial_t f(t, \gamma, \mu)$ follows from its definition. For the uniqueness of $(\partial_{\gamma_\tau}, \partial_{\gamma_\tau}^2) f(t, \gamma_\tau, \mu)$, without loss of generality, assume $\tau = t$. Otherwise consider the non-anticipative path-dependent function $\tilde{f}_{\mu_t}(\tau, \gamma) := f(t, \gamma_\tau, \mu)$ instead of $f(t, \gamma_\tau, \mu)$. For any $(\gamma, \mu) \in \mathbb{C}_{T,d} \times \mathcal{P}_2^C$, take $c(\cdot) = d(\cdot) = 0$ in equation (42) for X' and $a = 0$ in equation (41) for X . For any extension F of f , applying Itô formula (43) to $F(s, X, \mathcal{L}_{\eta_t})$ on $s \in [t, T]$, we have

$$\begin{aligned}
(169) \quad f(T, X, \mathcal{L}_{\eta_t}) - f(t, \gamma_t, \mathcal{L}_{\eta_t}) &= \int_t^T \partial_r f(r, X, \mathcal{L}_{\eta_t}) dr + \int_t^T \partial_{\omega_r} F(r, X, \mathcal{L}_{\eta_t}) b(r) dB(r) \\
&+ \frac{1}{2} \int_t^T \text{Tr} [\partial_{\omega_r}^2 F(r, X, \mathcal{L}_{\eta_t}) b(r) b(r)^T] dr.
\end{aligned}$$

In view of identity (169) and the Doob-Meyer theorem for semimartingales, we obtain the uniqueness of $\partial_{\omega_t} F(t, \gamma_t, \mu_t)$ and $\text{Sym}(\partial_{\omega_t}^2 F(t, \gamma_t, \mu_t))$. For the uniqueness of $\partial_{\mu_\tau} f$ and $\text{Sym}(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f)$, again we assume $\tau = t$. Otherwise consider $\tilde{f}_{\omega_t}(\tau, \mu) = f(t, \omega, \mu_\tau)$, and then by definition

$$\partial_{\mu_\tau} f(t, \omega, \mu_\tau, \tilde{\omega}_\tau) = \partial_{\mu_\tau} \tilde{f}_{\omega_t}(\tau, \mu, \tilde{\omega}), \quad \text{Sym}(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} f(t, \omega, \mu_\tau, \tilde{\omega}_\tau)) = \text{Sym}(\partial_{\tilde{\omega}_\tau} \partial_{\mu_\tau} \tilde{f}_{\omega_t}(\tau, \mu, \tilde{\omega})).$$

Then the uniqueness of $\partial_{\mu} f$ and $\text{Sym}(\partial_{\tilde{\omega}} \partial_{\mu} f)$ follows from [50, Theorem 2.9]. \square

Remark 4.5. *The uniqueness of $\partial_{\tilde{\omega}} \partial_{\mu} f$ can be proved via a similar argument as above from the uniqueness of $\partial_{\mu} f$ under a stronger assumption on the regularity of f . However, our Itô-Dupire formulas and analysis below only depend on $\text{Sym}(\partial_{\tilde{\omega}} \partial_{\mu} f)$. Indeed, equation (165) also only depends on $\text{Sym}(\partial_{\tilde{\omega}} \partial_{\mu} u)$ instead of $\partial_{\tilde{\omega}} \partial_{\mu} u$.*

For (f, Φ) in equation (165), we assume

- (A2) $\Phi : \mathbb{C}_{T,d} \times \mathcal{P}_2^C \rightarrow \mathbb{R}$ such that there exists Φ' satisfying **(H2)**(i) and $\Phi = \Phi'$ on $\mathbb{C}_{T,d} \times \mathcal{P}_2^C$.
 $f : [0, T] \times \mathbb{C}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^C \times \mathcal{P}_2(\mathbb{R}) \rightarrow \mathbb{R}$ such that there exists f' satisfying **(H2)**(ii) and $f = f'$ on $[0, T] \times \mathbb{C}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^C \times \mathcal{P}_2(\mathbb{R})$.

A functional $u \in \mathcal{C}_p^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$ is called a classical solution to equation (165) if it satisfies equation (165). The following theorem states the uniqueness of solutions for master equation (165).

Theorem 4.6. *Suppose that u^1 and u^2 are two classical solutions to the path-dependent master equation (165). Then $u^1 = u^2$.*

Proof. Apply Itô formula (167) to $u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}})$ on $r \in [t, s]$, $i = 1, 2$, and we obtain that

$$\begin{aligned} du_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}) &= \partial_r u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}) dr + \partial_\omega u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}) dB(r) + \frac{1}{2} \text{Tr}[\partial_\omega^2 u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}})] dr \\ &\quad + \frac{1}{2} \tilde{\mathbb{E}} \left[\text{Tr}[\partial_{\tilde{\omega}_r} \partial_{\mu_r} u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}, \tilde{B}^{\eta_t})] dr \right]. \end{aligned}$$

In view of equation (165), we have

$$\begin{aligned} du_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}) &= -f(r, B^{\gamma_t}, u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}), \partial_\omega u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}), \mathcal{L}_{\tilde{B}^{\eta_t}}, \mathcal{L}_{u_i(r, B^{\eta_t}, \mathcal{L}_{\tilde{B}^{\eta_t}})}) dr \\ &\quad + \partial_\omega u_i(r, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}) dB(r). \end{aligned}$$

Then processes $(Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t})$ and (Y^{η_t}, Z^{η_t}) given by

$$\begin{aligned} (Y^{\gamma_t, \eta_t}(s), Z^{\gamma_t, \eta_t}(s)) &:= (u_i(s, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}), \partial_\omega u_i(s, B^{\gamma_t}, \mathcal{L}_{\tilde{B}^{\eta_t}})), \quad \text{and} \\ (Y^{\eta_t}(s), Z^{\eta_t}(s)) &:= (u_i(s, B^{\eta_t}, \mathcal{L}_{\tilde{B}^{\eta_t}}), \partial_\omega u_i(s, B^{\eta_t}, \mathcal{L}_{\tilde{B}^{\eta_t}})), \quad s \geq t, \end{aligned}$$

define solutions to equations (146) and (147), respectively. From the uniqueness of solutions for BSDEs (146) and (147), our conclusion follows. \square

In view of the comparison theorem for BSDEs, we have the following comparison theorem for one dimensional path-dependent master equation (165).

Corollary 4.7. *Suppose that (f_i, Φ_i) , $i = 1, 2$, satisfy the following conditions: for any $(t, \omega, y, z, \mu, \nu) \in [0, T] \times \mathbb{C}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^C \times \mathcal{P}_2(\mathbb{R})$,*

$$(i) \ f_1(t, \omega, y, z, \mu, \nu) \leq f_2(t, \omega, y, z, \mu, \nu); \quad (ii) \ \Phi_1(\omega_T, \mu_T) \leq \Phi_2(\omega_T, \mu_T).$$

If $u_i \in \mathcal{C}_p^{1,2,1,1}(\hat{\mathbb{C}}_{T,d})$ is a solution to equation (165) with $(f, \Phi) := (f_i, \Phi_i)$ for $i = 1, 2$, then

$$(170) \quad u_1(t, \gamma, \mu) \leq u_2(t, \gamma, \mu), \quad \forall (t, \gamma, \mu) \in \hat{\mathbb{C}}_{T,d}.$$

Now we show that there is a classical solution to (165) via FBSDEs.

Theorem 4.8. *Suppose that (f, Φ) satisfies Assumption **(A2)** and u is given by (148). Then u restricted on $\hat{\mathbb{C}}_{T,d}$ is a classical solution of (165).*

Proof. In view of Proposition 4.2, we have $u \in \mathcal{C}_{s,p}^{0,2,1,1}(\hat{\mathbb{C}}_{T,d})$. For any $(t, \gamma, \mu) \in \hat{\mathbb{C}}_{T,d}$ and $h > 0$,

$$(171) \quad \begin{aligned} &u(t+h, \gamma_t, \mu_t) - u(t, \gamma, \mu) \\ &= u(t+h, \gamma_t, \mu_t) - \mathbb{E}[u(t+h, B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}})] + \mathbb{E}[u(t+h, B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}})] - u(t, \gamma, \mu). \end{aligned}$$

Applying partial Itô formula (168) to $u(t+h, B_r^{\gamma_t}, \mathcal{L}_{B_r^{\eta_t}})$ on $r \in [t, t+h]$, we have

$$(172) \quad \begin{aligned} &u(t+h, \gamma_t, \mu_t) - u(t+h, B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}) \\ &= - \int_t^{t+h} \partial_{\omega_r} u(t+h, B_r^{\gamma_t}, \mathcal{L}_{B_r^{\eta_t}}) dB(r) - \frac{1}{2} \int_t^{t+h} \text{Tr}[\partial_{\omega_r}^2 u(t+h, B_r^{\gamma_t}, \mathcal{L}_{B_r^{\eta_t}})] dr \\ &\quad - \frac{1}{2} \tilde{\mathbb{E}} \int_t^{t+h} \text{Tr}[\partial_{\tilde{\omega}_r} \partial_{\mu_r} u(t+h, B_r^{\gamma_t}, \mathcal{L}_{B_r^{\eta_t}}, \tilde{B}_r^{\eta_t})] dr. \end{aligned}$$

On the other hand, in view of identities (155), (151), and BSDE (146)

$$(173) \quad \begin{aligned} &u(t+h, B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}) - u(t, \gamma, \mu) = Y^{\gamma_t, \eta_t}(t+h) - Y^{\gamma_t, \eta_t}(t) \\ &= - \int_t^{t+h} f(r, B^{\gamma_t}, Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t}, \mathcal{L}_{B^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}}) dr + \int_t^{t+h} Z^{\gamma_t, \eta_t}(r) dB(r). \end{aligned}$$

Putting (172) and (173) to (171), and taking expectation \mathbb{E} , we obtain

$$(174) \quad \begin{aligned} u(t+h, \gamma_t, \mu_t) - u(t, \gamma, \mu) &= -\frac{1}{2} \int_t^{t+h} \mathbb{E} [\text{Tr}[\partial_{\omega_r}^2 u(t+h, B_r^{\gamma_t}, \mathcal{L}_{B_r^{\eta_t}})]] dr \\ &\quad - \frac{1}{2} \int_t^{t+h} \mathbb{E} [\tilde{\mathbb{E}} \text{Tr}[\partial_{\tilde{\omega}_r} \partial_{\mu_r} u(t+h, B_r^{\gamma_t}, \mathcal{L}_{B_r^{\eta_t}}, \tilde{B}_r^{\tilde{\eta}_t})]] dr \\ &\quad - \int_t^{t+h} \mathbb{E} [f(r, B^{\gamma_t}, Y^{\gamma_t, \eta_t}, Z^{\gamma_t, \eta_t}, \mathcal{L}_{B^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}})] dr. \end{aligned}$$

Moreover, in view of (151), (155) and (156), we have

$$(175) \quad Y^{\gamma_t, \eta_t}(r) = u(r, B^{\gamma_t}, \mathcal{L}_{B^{\eta_t}}), \quad Z^{\gamma_t, \eta_t}(r) = \partial_{\gamma_t} u(r, B^{\omega_t}, \mathcal{L}_{B^{\eta_t}}),$$

$$(176) \quad \text{and} \quad Y^{\eta_t}(r) = u(r, B^{\eta_t}, \mathcal{L}_{B^{\eta_t}}).$$

Then dividing both sides of (174) by h and taking $h \rightarrow 0^+$, according to the dominated convergence theorem and Proposition 4.2, we obtain

$$\begin{aligned} \partial_t u(t, \gamma, \mu) &= -\frac{1}{2} \text{Tr} [\partial_{\omega}^2 u(t, \gamma, \mu)] - \frac{1}{2} \text{Tr} [\mathbb{E}^\mu [\partial_{\tilde{\omega}} \partial_{\mu} u(t, \gamma, \mu, W)]] \\ &\quad - f(t, \gamma, u(t, \gamma, \mu), \partial_{\omega} u(t, \gamma, \mu), \mu, \mathcal{L}_{u(t, B^{\eta_t}, \mu)}). \end{aligned}$$

□

Remark 4.9. Substitute $(X^{\gamma_t}, X^{\eta_t})$ for $(B^{\gamma_t}, B^{\eta_t})$ in BSDEs (146), where

$$(177) \quad \begin{cases} dX^{\gamma_t}(r) = b_1(r)dr + \sigma_1(r)dB(r), \\ dX^{\eta_t}(r) = b_2(r)dr + \sigma_2(r)dB(r), \\ X_t = \gamma_t, \quad X'_t = \eta_t, \quad r \in [t, T], \end{cases}$$

with $(b_1, \sigma_1, b_2, \sigma_2) : [0, T] \rightarrow (\mathbb{R}^d \times \mathbb{R}^{d \times d})^2$ being continuous functions. Following a similar argument, we deduce that under Assumption **(A2)**, $u(t, \gamma, \mu) := Y^{\gamma_t, \eta_t}(t)$ is the unique classical solution to the following path-dependent master equation

$$(178) \quad \begin{cases} \partial_t u(t, \gamma, \mu) + \frac{1}{2} \text{Tr} [\partial_{\omega}^2 u(t, \gamma, \mu) \sigma_1(t) \sigma_1(t)^T] + \partial_{\omega} u(t, \gamma, \mu) b_1(t) \\ \quad + \frac{1}{2} \text{Tr} [\mathbb{E}^\mu [\partial_{\tilde{\omega}} \partial_{\mu} u(t, \gamma, \mu, W) \sigma_2(t) \sigma_2(t)^T] + \mathbb{E}^\mu [\partial_{\mu} u(t, \gamma, \mu, W) b_2(t)] \\ \quad + f(t, \gamma, u(t, \gamma, \mu), \sigma_1(t) \partial_{\omega} u(t, \gamma, \mu), \mu, \mathcal{L}_{u(t, W^\mu, \mu)}) = 0, \\ u(T, \gamma, \mu) = \Phi(\gamma_T, \mu_T), \quad (t, \gamma, \mu) \in \hat{\mathbb{C}}_{T,d}. \end{cases}$$

Moreover, our probabilistic approach works in the multiple dimensional case when $u(t, \gamma, \mu) \in \mathbb{R}^k$ with $k \geq 1$.

4.3. Some typical cases. In view of Remark 2.14, the path-dependent master equation (165) involves many interesting special cases. In the following we list some typical ones, where we always assume that (f, Φ) satisfies Assumption **(A2)** and $(t, \omega, y, z, \mu, \nu) \in [0, T] \times \mathbb{C}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^C \times \mathcal{P}_2(\mathbb{R})$.

(i) **The state-dependent master equation.** Suppose that (f, Φ) has a state-dependent form:

$$(179) \quad f(t, \omega, y, z, \mu, \nu) = F(t, \omega(t), y, z, \mu(t), \nu),$$

$$(180) \quad \Phi(T, \omega, \mu) = G(\omega(T), \mu(T)),$$

for functionals $F : [0, T] \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{P}_2(\mathbb{R}) \rightarrow \mathbb{R}$ and $G : \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$. In this case, the differentiability of (f, Φ) is equivalent to that of (F, G) in its corresponding domain, and master equation (165) has the form

$$(181) \quad \begin{cases} \partial_t u(t, \gamma, \mu) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \gamma, \mu)] + \frac{1}{2} \text{Tr} \left[\int_{\mathbb{C}_{T,d}} \partial_{\tilde{\omega}} \partial_\mu u(t, \gamma, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) \right] \\ \quad + F(t, \gamma(t), u(t, \gamma, \mu), \partial_\omega u(t, \gamma, \mu), \mu(t), \mathcal{L}_{u(t, W^\mu, \mu)}) = 0, \\ u(T, \gamma, \mu) = G(\gamma(T), \mu(T)), \quad (t, \gamma, \mu) \in \hat{\mathcal{C}}_{T,d}. \end{cases}$$

Since the corresponding FBSDE is Markovian, we see that $u(t, \gamma, \mu) = U(t, \gamma(t), \mu(t))$ for a smooth (indeed $\mathcal{C}^{1,2,1,1}$ in view of Definition 2.13 with obvious adjustment) functional $U : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ thanks to Remark 2.14. Then we obtain the well-posedness of the (state-dependent) master equation

$$(182) \quad \begin{cases} \partial_t U(t, a, \lambda) + \frac{1}{2} \text{Tr} [\partial_a^2 U(t, a, \lambda)] + \frac{1}{2} \int_{\mathbb{R}^d} \partial_{\tilde{a}} \partial_\lambda U(t, a, \lambda, \tilde{a}) \lambda(d\tilde{a}) \\ \quad + F(t, a, U(t, a, \lambda), \partial_a U(t, a, \lambda), \lambda, \mathcal{L}_{u(t, \xi, \lambda)}) = 0, \\ U(T, a, \lambda) = G(a, \lambda), \quad (t, a, \lambda) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d), \end{cases}$$

which is considered in [9, 10]. Here ξ is a random variable on \mathbb{R}^d with law λ .

(ii) **The PPDE.** Suppose that (f, Φ) does not depend on measures:

$$(183) \quad f(t, \omega, y, z, \mu, \nu) = H(t, \omega, y, z),$$

$$(184) \quad \Phi(T, \omega, \mu) = I(\omega_T),$$

with $H : [0, T] \times \mathbb{C}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $I : \mathbb{C}_{T,d} \rightarrow \mathbb{R}$. Then, Equation (165) becomes the following PPDE

$$(185) \quad \begin{cases} \partial_t u(t, \gamma) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \gamma)] + H(t, \gamma, u(t, \gamma), \partial_\omega u(t, \gamma)) = 0, \\ u(T, \gamma) = I(\gamma_T), \quad (t, \gamma) \in [0, T] \times \mathbb{C}_{T,d}. \end{cases}$$

Then, Theorem 4.8 includes as a special case the well-posedness of PPDEs shown in [41, Theorem 4.5] under a stronger assumption in view of the integrability of BSDE (147).

(iii) **The measure-dependent master equation.** Suppose that (f, Φ) does not depend on the path/state variable and has the following structure

$$(186) \quad f(t, \omega, y, z, \mu, \nu) = J(t, y, \mu),$$

$$(187) \quad \Phi(T, \omega, \mu) = K(\mu_T),$$

where $J : [0, T] \times \mathbb{R} \times \mathcal{P}_2^C \rightarrow \mathbb{R}$ and $K : \mathcal{P}_2^C \rightarrow \mathbb{R}$. In this case our path-dependent master equation is reduced to

$$(188) \quad \begin{cases} \partial_t u(t, \mu) + \frac{1}{2} \text{Tr} \left[\int_{\mathbb{C}_{T,d}} \partial_{\tilde{\omega}} \partial_\mu u(t, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) \right] + J(t, u(t, \mu), \mu) = 0, \\ u(T, \mu) = K(\mu_T), \quad (t, \mu) \in [0, T] \times \mathcal{P}_2^C. \end{cases}$$

Such master equation is introduced in [50] for a closed-loop control problem with control being the form of $\alpha_t = \alpha(t, \mathcal{L}_{X_t})$.

(iv) **Path-state mixed cases.** Suppose that (f, Φ) has the following form

$$(189) \quad f(t, \omega, y, z, \mu, \nu) = L(t, \omega, y, z, \mu(t)),$$

$$(190) \quad \Phi(T, \omega, \mu) = M(\omega_T, \mu(T)),$$

where $L : [0, T] \times \mathbb{C}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $M : \mathbb{C}_{T,d} \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$. Then we have a unique smooth solution for the following master equation

$$(191) \quad \begin{cases} \partial_t u(t, \gamma, \mu) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \gamma, \mu)] + \frac{1}{2} \text{Tr} \left[\int_{\mathbb{C}_{T,d}} \partial_{\tilde{\omega}} \partial_\mu u(t, \gamma, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) \right] \\ \quad + L(t, \gamma, u(t, \gamma, \mu), \partial_\omega u(t, \gamma, \mu), \mu(t)) = 0, \\ u(T, \gamma, \mu) = M(\gamma_T, \mu(T)), \quad (t, \gamma, \mu) \in \hat{\mathbb{C}}_{T,d}. \end{cases}$$

In view of the corresponding FBSDE (146), we see that $u(t, \gamma, \mu) = U(t, \gamma, \mu(t))$ for a functional $U : [0, T] \times \mathbb{C}_{T,d} \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$. Since $u \in \mathcal{C}^{1,2,1,1}$, U is the unique classical solution to the master equation

$$(192) \quad \begin{cases} \partial_t U(t, \gamma, \lambda) + \frac{1}{2} \text{Tr} [\partial_\omega^2 U(t, \gamma, \lambda)] + \frac{1}{2} \text{Tr} \left[\int_{\mathbb{R}^d} \partial_{\tilde{a}} \partial_\lambda U(t, \gamma, \lambda, \tilde{a}) \lambda(d\tilde{a}) \right] \\ \quad + L(t, \gamma, U(t, \gamma, \lambda), \partial_\omega U(t, \gamma, \lambda), \mu(t)) = 0, \\ U(T, \gamma, \lambda) = M(\gamma_T, \lambda), \quad (t, \gamma, \lambda) \in [0, T] \times \mathbb{C}_{T,d} \times \mathcal{P}_2(\mathbb{R}^d). \end{cases}$$

On the other hand, if (f, Φ) has the the following structure

$$(193) \quad f(t, \omega, y, z, \mu, \nu) = N(t, \omega(t), y, z, \mu, \nu) \quad \text{and}$$

$$(194) \quad \Phi(T, \omega, \mu) = P(\omega(T), \mu_T)$$

for some functionals $N : [0, T] \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathcal{P}_2^C \times \mathcal{P}_2(\mathbb{R}) \rightarrow \mathbb{R}$ and $P : \mathbb{R}^d \times \mathcal{P}_2^C \rightarrow \mathbb{R}$, then $u(t, \omega, \mu) = V(t, \omega(t), \mu_t)$ for a functional $V : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2^C \rightarrow \mathbb{R}$. Then, V is the unique smooth solution to the master equation

$$(195) \quad \begin{cases} \partial_t V(t, a, \mu) + \frac{1}{2} \text{Tr} [\partial_a^2 V(t, a, \mu)] + \frac{1}{2} \text{Tr} \left[\int_{\mathbb{C}_{T,d}} \partial_{\tilde{\omega}} \partial_\mu V(t, x, \mu, \tilde{\omega}) \mu(d\tilde{\omega}) \right] \\ \quad + N(t, a, V(t, a, \mu), \partial_a V(t, a, \mu), \mu, \mathcal{L}_{V(t, W^\mu(t), \mu)}) = 0, \\ V(T, a, \mu) = P(a, \mu_T), \quad (t, a, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}_2^C. \end{cases}$$

4.4. The general case via functional mollifying. To our best knowledge, there is no viscosity solution theory corresponding to the path-dependent master equation in the general form of (165) yet (see our future work). However, if the decoupling field only depends on γ_t or μ_t , viscosity solutions are introduced in [21] and [50] respectively. Moreover, we can check that the (degenerate) smooth solution introduced here is a viscosity solution. On the other hand, Examples 2.2 and 2.11 enlighten that we can “smoothen” the equation such that it adapts a smooth solution which converges to the unique limit. Though in the path-dependent or functional case, there is no general way to smoothen the terminal functional or the generator functional yet, we consider the following kind of inhomogeneous path-dependent master equation

$$(196) \quad \begin{cases} \partial_t u(t, \omega, \mu) + \frac{1}{2} \text{Tr} [\partial_\omega^2 u(t, \omega, \mu)] + \frac{1}{2} \text{Tr} [\partial_{\omega'} \partial_\mu u(t, \omega, \mu, \omega')] = f(t, \omega, \mu), \\ u(T, \omega, \mu) = \Phi(T, \omega, \mu), \quad (t, \omega, \mu) \in \hat{\mathbb{C}}_{T,d}. \end{cases}$$

In this subsection, we assume that (f, Φ) satisfies

- (A3)** (i) there is a sequence of functionals $\{(f_\varepsilon, \Phi_\varepsilon)\}_{\varepsilon \geq 0}$ which satisfies Assumption **(A2)** and the following polynomial growth condition: for any $K > 0$,

$$(197) \quad \begin{aligned} |f_\varepsilon(t, \omega, \mu)| &\leq C_K(1 + \|\omega_t\|^p), & |\Phi_\varepsilon(\omega_T, \mu_T)| &\leq C_K(1 + \|\omega_T\|^p), \\ \forall (t, \omega, \mu) &\in \hat{\mathbb{C}}_{T,d}, \text{ such that } \|\mu\| &\leq K, \end{aligned}$$

with constants $p \in \mathbb{N}^+$ and $C_K \in \mathbb{R}^+$.

- (ii) $(f(t, \omega, \mu), \Phi(T, \omega, \mu)) = \lim_{\varepsilon \rightarrow 0} (f_\varepsilon(t, \omega, \mu), \Phi_\varepsilon(T, \omega, \mu)), \forall (t, \omega, \mu) \in \hat{\mathbb{C}}_{T,d}$.

According to Theorem 4.8, the solution of (196) driven by $(f_\varepsilon, \Phi_\varepsilon)$ satisfies

$$(198) \quad u_\varepsilon(t, \omega, \mu) = Y_\varepsilon^{\omega_t, \mu_t}(t),$$

where $Y_\varepsilon^{\omega_t, \mu_t}$ is the unique solution of the following FBSDE

$$(199) \quad Y_\varepsilon(s) = \Phi_\varepsilon(B_T^{\omega_t}, \mathcal{L}_{B_T^{\eta_t}}) + \int_s^T f_\varepsilon(r, B_r^{\omega_t}, \mathcal{L}_{B_r^{\eta_t}}) dr - \int_s^T Z_\varepsilon(r) dB(r),$$

with $\mathcal{L}_{\eta_t} = \mu_t$. In this case, by taking the conditional expectation, we obtain

$$(200) \quad u_\varepsilon(t, \omega, \mu) = Y_\varepsilon(t) = \mathbb{E}_{\mathcal{F}_t}[\Phi_\varepsilon(B_T^{\omega_t}, \mathcal{L}_{B_T^{\eta_t}}) + \int_t^T f_\varepsilon(r, B_r^{\omega_t}, \mathcal{L}_{B_r^{\eta_t}}) dr].$$

Moreover, according to the dominated convergence theorem, u_ε converges to the following limit as ε goes to null,

$$(201) \quad u(t, \omega, \mu) := \mathbb{E}_{\mathcal{F}_t}[\Phi(B_T^{\omega_t}, \mathcal{L}_{B_T^{\eta_t}}) + \int_t^T f(r, B_r^{\omega_t}, \mathcal{L}_{B_r^{\eta_t}}) dr].$$

Then $u(t, \omega, \mu)$ formally defines a unique solution of equation (196) and we call u a Sobolev solution. In conclusion, we show

Theorem 4.10. *Suppose (f, Φ) satisfies Assumption **(A3)**. Then there exists a unique Sobolev solution to equation (196) on $\hat{\mathbb{C}}_{T,d}$.*

In the following, we consider path-dependent master equations with time delayed functionals.

Example 4.11. (i) For any $t_0 \in (0, T)$ and $F \in C_b^3(\mathbb{R}^d)$, consider PPDE

$$(202) \quad \begin{cases} \partial_t u(t, \omega) + \frac{1}{2} \text{Tr}[\partial_\omega^2 u(t, \omega)] = 0, \\ u(T, \omega) = F(\omega(t_0)), \quad (t, \omega) \in [0, T] \times \mathbb{C}_{T,d}. \end{cases}$$

Let $\Phi_\varepsilon(T, \omega) := \int_0^T \rho_\varepsilon(t_0 - s) F(\omega(s)) ds$, where $\rho_\varepsilon \in C_c^\infty$ is a positive standard modifier on \mathbb{R} . According to Example 2.2 (iv), Φ_ε satisfies Assumption **(A2)**, and it is easy to check that Φ_ε satisfies condition (197). Moreover, $\lim_{\varepsilon \rightarrow 0} \Phi_\varepsilon(T, \omega) = F(\omega(t_0))$, for any $\omega \in \mathbb{C}_{T,d}$. Then according to Theorem 4.10, there exists a unique Sobolev solution of this PPDE. In particular, if $F(x) = ax$ for some $a \in \mathbb{R}^d$, by solvability of the corresponding BSDE with terminal value $\Phi_\varepsilon(T, B_T)$, we have

$$(203) \quad u_\varepsilon(t, \omega) = \int_0^t \rho_\varepsilon(t_0 - s) a \omega(s) ds + a \omega(t) \int_t^T \rho_\varepsilon(t_0 - s) ds.$$

Then, it follows that $u_\varepsilon(t, \omega) \rightarrow a\omega(t \wedge t_0)$, as $\varepsilon \rightarrow 0$. Indeed, $u(t, \omega) := a\omega(t \wedge t_0)$ is the unique viscosity solution in the sense of [21].

(ii) Similarly, for any $F \in C_b^3(\mathbb{R}^d)$ and $G \in C_b^3(\mathbb{R}^d, \mathbb{R}^d)$, consider the following kind of (degenerate) master equation

$$(204) \quad \begin{cases} \partial_t u(t, \mu) + \frac{1}{2} \text{Tr}[\partial_{\omega'} \partial_\mu u(t, \mu, \omega')] = 0, \\ u(T, \mu) = F(\mathbb{E}^\mu[G(W(t_0))]), \quad (t, \mu) \in [0, T] \times \mathcal{P}_2^C. \end{cases}$$

Let $\Phi_\varepsilon(T, \mu) := \int_0^T \rho_\varepsilon(t_0 - s) F(\mathbb{E}^\mu[G(W(s))]) ds$. In view of Example 2.11 (iii) and basic estimates, Φ_ε satisfies Assumption **(A2)** and condition (197). Moreover, for any $\mu \in \mathcal{P}_2^C$, $\lim_{\varepsilon \rightarrow 0} \Phi_\varepsilon(T, \mu) = \Phi(T, \mu)$. Then the master equation has a unique solution given by (201). In particular if $\Phi(T, \mu) = \mathbb{E}^\mu[aW(t_0)]$, equation (204) with terminal value Φ_ε has a unique non-anticipative classical solution,

$$(205) \quad u_\varepsilon(t, \mu) = \mathbb{E}^\mu\left[\int_0^t \rho_\varepsilon(t_0 - s) aW(s) ds\right] + \mathbb{E}^\mu[aW(t)] \int_t^T \rho_\varepsilon(t_0 - s) ds.$$

Furthermore, for any $(t, \mu) \in [0, T] \times \mathcal{P}_2^C$, $u_\varepsilon(t, \mu) \rightarrow \mathbb{E}^\mu[aW(t \wedge t_0)]$ as $\varepsilon \rightarrow 0$, and $u(t, \mu) := \mathbb{E}^\mu[aW(t \wedge t_0)]$ is the unique viscosity solution of equation (202) in the sense of [50].

(iii) Generally, suppose that f satisfies **(A2)** and $H \in C_b^3(\mathbb{R}^d \times \mathbb{R}^d)$. For any $t_1, t_2 \in (0, T)$, consider

$$(206) \quad \begin{cases} \partial_t u(t, \omega, \mu) + \frac{1}{2} \partial_\omega^2 u(t, \omega, \mu) + \frac{1}{2} \partial_{\omega'} \partial_\mu u(t, \omega, \mu, \omega') = f(t, \omega, \mu), \\ u(T, \omega, \mu) = H(\omega(t_1), \mathbb{E}^\mu[W(t_2)]). \end{cases}$$

Let $\Phi_\varepsilon(T, \omega, \mu) := \int_0^T \int_0^T H(\omega(r_1), \mathbb{E}^\mu[W(r_2)]) \rho_\varepsilon(t_1 - r_1) \rho_\varepsilon(t_2 - r_2) dr_1 dr_2$. According to Theorem 4.10, there exists a unique solution u_ε of equation (206) with terminal Φ_ε . Moreover, u_ε converges to a unique solution u of (206) with terminal $H(\omega(t_1), \mathbb{E}^\mu[W(t_2)])$. In particular, if $H(T, \omega, \mu) = a\omega(t_1) + \mathbb{E}^\mu[bW(t_2)]$,

$$(207) \quad u(t, \omega, \mu) = a\omega(t \wedge t_1) + \mathbb{E}^\mu[bW(t \wedge t_2)], \quad (t, \omega, \mu) \in \hat{\mathcal{C}}_{T,d}.$$

(iv) For any fixed $t_0 \in (0, T)$ and $g \in C_b^3(\mathbb{R}^d \times \mathbb{R}^d)$, consider

$$(208) \quad \begin{cases} \partial_t u(t, \omega, \mu) + \frac{1}{2} \text{Tr}[\partial_\omega^2 u(t, \omega, \mu) + \partial_{\omega'} \partial_\mu u(t, \omega, \mu, \omega')] = g(\omega(t_0), \mathbb{E}^\mu(W(t_0))) 1_{[t_0, T]}(t), \\ u(T, \mu) = H(\omega(t_1), \mathbb{E}^\mu[W(t_2)]), \quad (t, \omega, \mu) \in \hat{\mathcal{C}}_{T,d}, \end{cases}$$

with $H(\omega(t_1), \mathbb{E}^\mu[W(t_2)])$ defined as in (iii). Let

$$(209) \quad f_\varepsilon(t, \omega, \mu) := \int_0^t \int_0^t g(\omega(r_1), \mathbb{E}^\mu[W(r_2)]) \rho_\varepsilon(t_0 - r_1) \rho_\varepsilon(t_0 - r_2) dr_1 dr_2.$$

We check that f_ε satisfies **(A2)** and condition (197). Furthermore,

$$\lim_{\varepsilon \rightarrow 0} f_\varepsilon(t, \omega, \mu) = g(\omega(t_0), \mathbb{E}^\mu(W(t_0))) 1_{[t_0, T]}(t), \quad \forall (t, \omega, \mu) \in \hat{\mathcal{C}}_{T,d}.$$

Then our Theorem 4.10 applies, and there exists a unique Sobolev solution to equation (208) driven by functional f without continuity.

5. APPENDIX

5.1. **Proof of Lemma 3.4.** We omit the proof of inequality (70) since its proof is easiest and is similar to that of (71) for $p = 2$.

Now suppose that (70) is true, and we show inequality (72) first. In what follows, we write $(\mathcal{Y}, \mathcal{Z}) := (Y^{\eta_t}, Z^{\eta_t})$ and notations such as $(\mathcal{Y}', \mathcal{Z}')$ and (Φ, Φ') are defined in a similar way. Set $(\delta\mathcal{Y}, \delta\mathcal{Z}) := (\mathcal{Y} - \mathcal{Y}', \mathcal{Z} - \mathcal{Z}')$. We see that $(\delta\mathcal{Y}, \delta\mathcal{Z})$ solves the following linearized BSDE

$$(210) \quad \begin{aligned} \delta\mathcal{Y}(s) &= \Phi - \Phi' + \int_s^T [f(\Theta_r^\eta, \mathcal{L}_{\Theta_r^\eta}) - f(\Theta_r^{\eta'}, \mathcal{L}_{\Theta_r^{\eta'}})] dr - \int_s^T \delta\mathcal{Z}(r) dB(r) \\ &=: \delta\Phi + \int_s^T \left(a_r \delta\mathcal{Y}(r) + b_r \delta\mathcal{Z}(r) + \tilde{\mathbb{E}}[\tilde{c}_r \delta\tilde{\mathcal{Y}}(r)] + \delta h_r \right) dr - \int_s^T \delta\mathcal{Z}(r) dB(r), \end{aligned}$$

where

$$\begin{aligned} a_r &:= \int_0^1 \partial_y f(B_r^{\eta_t}, \mathcal{Y}' + \theta(\mathcal{Y} - \mathcal{Y}'), \mathcal{Z}, \mathcal{L}_{\Theta_r^\eta}) d\theta, \\ b_r &:= \int_0^1 \partial_z f(B_r^{\eta_t}, \mathcal{Y}', \mathcal{Z}' + \theta(\mathcal{Z} - \mathcal{Z}'), \mathcal{L}_{\Theta_r^{\eta_t}}) d\theta, \\ \tilde{c}_r &:= \int_0^1 \partial_\nu f(B_r^{\eta_t}, \mathcal{Y}', \mathcal{Z}', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{\mathcal{Y}' + \theta(\mathcal{Y} - \mathcal{Y}')} , \tilde{\mathcal{Y}}' + \theta(\tilde{\mathcal{Y}} - \tilde{\mathcal{Y}}')) d\theta, \quad \text{and} \\ \delta h_r &:= f(B_r^{\eta_t}, \mathcal{Y}', \mathcal{Z}', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{\tilde{\mathcal{Y}}'}) - f(B_r^{\eta_t}, \mathcal{Y}', \mathcal{Z}', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{\tilde{\mathcal{Y}}}). \end{aligned}$$

Let

$$F(\Theta_t^\eta, y_1, y_2) := \int_0^1 \partial_\nu f(B_r^{\eta_t}, \mathcal{Y}', \mathcal{Z}', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{\mathcal{Y}' + \theta(\mathcal{Y} - \mathcal{Y}')} , y_1 + \theta y_2) d\theta.$$

Then, F is uniformly Lipschitz continuous in (y_1, y_2) in view of Assumption **(H0)**. On the other hand, since $\mathcal{Y}', \mathcal{Z}' \in \mathbb{H}^2$, we deduce that $F(\Theta_t^\eta, 0, 0) \in \mathbb{H}^2$, and moreover, we have

$$(211) \quad \|\mathcal{Y}\|_{\mathbb{H}^2} + \|\mathcal{Y}'\|_{\mathbb{H}^2} + \|F(\Theta_t^\eta, 0, 0)\|_{\mathbb{H}^2} \leq C(1 + \|\eta_t\|_{\mathbb{S}^2} + \|\eta_t'\|_{\mathbb{S}^2}).$$

Then applying estimates of Lemma 3.1 to BSDE (210), we have

$$(212) \quad \|\delta\mathcal{Y}\|_{\mathbb{S}^2}^2 + \|\delta\mathcal{Z}\|_{\mathbb{H}^2}^2 \leq C(\|\Phi - \Phi'\|_{L^2}^2 + \|\delta h\|_{\mathbb{H}^2}^2) e^{C(\|F(\Theta_t^\eta, 0, 0)\|_{\mathbb{H}^2} + \|\mathcal{Y}, \mathcal{Y}'\|_{\mathbb{H}^2})}.$$

Furthermore, using the Lipschitz continuity of Φ and f , we have

$$(213) \quad \|\Phi - \Phi'\|_{L^2} + \|\delta h\|_{\mathbb{H}^2} \leq C\|\eta_t - \eta_t'\|_{\mathbb{S}^2},$$

and thus the desired estimate (72) in view of inequality (212).

Now we show inequalities (71). In what follows, we omit the superscript (γ_t, η_t) for simplicity. Without loss of generality, we assume $p = 2q$, $q \in \mathbb{Z}^+$. Otherwise, we replace $|Y|$ with $(|Y|^2 + \varepsilon)^{\frac{1}{2}}$ in the following argument and then take the limit $\varepsilon \rightarrow 0$. Applying Itô's formula to $|Y|^p$ on $[s, T]$, we have

$$(214) \quad \begin{aligned} |Y(s)|^p + \frac{1}{2}p(p-1) \int_s^T |Y|^{p-2}|Z|^2 dr \\ = |\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\gamma_t}})|^p + p \int_s^T |Y(r)|^{p-1} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr - p \int_s^T |Y(r)|^{p-1} Z(r) dB(r). \end{aligned}$$

Since Φ and f are Lipschitz continuous, we have

$$\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}}), \int_t^T f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(r)}) dr \in L^q(\mathcal{F}_T), \quad \forall q \geq 1.$$

Then, using standard estimates of BSDEs, we obtain $\|Y\|_{\mathbb{S}^q} + \|Z\|_{\mathbb{H}^q} < \infty$. Taking the expectation on both sides of identity (214), we have

$$(215) \quad \mathbb{E}[|Y(s)|^p + \frac{1}{2}p(p-1) \int_s^T |Y|^{p-2}|Z|^2 dr] \leq \mathbb{E}[|\Phi|^p] + p \int_s^T |Y(r)|^{p-1} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr.$$

Applying Young's inequality to the last integral, we have

$$(216) \quad \begin{aligned} & \int_s^T |Y(r)|^{p-1} f(\Theta_r^{\gamma_t, \eta_t}, \mathcal{L}_{\Theta_r^{\eta_t}}) dr \\ & \leq \int_s^T \left[|Y|^{p-1} f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0) + C|Y|^p + C|Y|^{p-1} [\tilde{\mathbb{E}}[\tilde{Y}^{\tilde{\eta}_t}(r)]]^{\frac{1}{2}} + C|Y|^{p-1}|Z| \right] dr \\ & \leq (C_p + \frac{C}{\varepsilon}) \int_s^T |Y|^p dr + \frac{1}{p} \int_s^T |f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr \\ & \quad + \varepsilon C \int_s^T |Y|^{p-2}|Z|^2 dr + \frac{C}{2p} \int_s^T \|\tilde{Y}^{\tilde{\eta}_t}(r)\|_{L^2}^p dr, \quad \forall \varepsilon > 0. \end{aligned}$$

Then by choosing a small enough ε such that $\frac{1}{2}p(p-1) - \varepsilon C > 0$, we obtain

$$(217) \quad \begin{aligned} & \mathbb{E}[|Y(s)|^p + C_p \int_s^T |Y(r)|^{p-2}|Z(r)|^2 dr] \\ & \leq \mathbb{E}[|\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p] + C_p \int_s^T [|f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr + \int_s^T |Y|^p dr] \\ & \quad + \int_s^T \|\tilde{Y}^{\tilde{\eta}_t}(r)\|_{L^2}^p dr. \end{aligned}$$

Apply Gronwall's inequality to (217), and we obtain

$$(218) \quad \begin{aligned} & \mathbb{E}[|Y(s)|^p + C_p \int_s^T |Y(r)|^{p-2}|Z(r)|^2 dr] \\ & \leq C_p \mathbb{E}[|\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p] + \int_s^T |f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr + \int_s^T \|\tilde{Y}^{\tilde{\eta}_t}(r)\|_{L^2}^p dr. \end{aligned}$$

Then in view of inequalities (216) and (214), choosing ε sufficiently small, we have

$$(219) \quad \begin{aligned} |Y(s)|^p & \leq |\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p + C_p \left[\int_s^T |f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr \right. \\ & \quad \left. + \int_s^T |Y(r)|^p dr + \int_s^T \|\tilde{Y}^{\tilde{\eta}_t}(r)\|_{L^2}^p dr \right] - p \int_s^T |Y|^{p-1} Z(r) dB(r). \end{aligned}$$

Applying Burkholder-Davis-Gundy inequality to the right hand side of inequality (219), we obtain

$$\begin{aligned}
\mathbb{E}[\sup_{s \in [t, T]} |Y(s)|^p] &\leq \mathbb{E}\left[|\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p + C_p \int_t^T |f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr \right. \\
&\quad \left. + C_p \int_s^T \|Y^{\eta_t}(r)\|_{L_2}^p dr\right] + C_p \mathbb{E}\left[\int_t^T |Y(r)|^p dr + \left(\int_t^T |Y|^{2p-2} |Z|^2 dr\right)^{\frac{1}{2}}\right] \\
(220) \quad &\leq \mathbb{E}\left[|\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p + C_p \int_t^T |f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr + C_p \int_s^T \|Y^{\eta_t}(r)\|_{L_2}^p dr\right] \\
&\quad + C_p \mathbb{E}\left[\int_t^T |Y(r)|^p dr + \varepsilon \mathbb{E}[\sup_{s \in [t, T]} |Y(s)|^p] + \frac{1}{4\varepsilon} \mathbb{E}\left[\int_t^T |Y|^{p-2} |Z|^2 dr\right]\right].
\end{aligned}$$

Then in view of (218), (70) and (220), choosing ε small again, we have

$$\begin{aligned}
\mathbb{E}[\sup_{s \in [t, T]} |Y(s)|^p] &\leq C_p \mathbb{E}\left[|\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p + \int_t^T |f(B_r^{\gamma_t}, 0, 0, \mathcal{L}_{B_r^{\eta_t}}, \delta_0)|^p dr + C_p \int_s^T \|Y^{\eta_t}(r)\|_{L_2}^p dr\right] \\
(221) \quad &\leq C_p (1 + \mathbb{E}[\|B^{\gamma_t}\|^p] + \|\|B^{\eta_t}\|\|_{L_2}^p + \|\|\eta_t\|\|_{L_2}^p) \\
&\leq C_p \left(1 + \mathbb{E}(\|B\| + \|\gamma_t\|)^p + [\mathbb{E}(\|B\| + \|\eta_t\|)^2]^{\frac{p}{2}} + \|\|\eta_t\|\|_{L_2}^p\right) \\
&\leq C_p (1 + \|\gamma_t\|^p + \|\|\eta_t\|\|_{L_2}^p).
\end{aligned}$$

Let $\tilde{f}(Y, Z) := f(B_r^{\gamma_t}, Y, Z, \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(r)})$. Following a standard argument of BSDEs, we have

$$\begin{aligned}
\mathbb{E}[\int_t^T |Z|^2 dr]^{\frac{p}{2}} &\leq C_p \mathbb{E}\left[|\Phi(B_T^{\gamma_t}, \mathcal{L}_{B_T^{\eta_t}})|^p + \int_t^T |\tilde{f}(0, 0)|^p dr + C_p \int_s^T \|Y(r)\|_{L_2}^p dr + \sup_{s \in [t, T]} |Y(s)|^p\right] \\
&\leq C_p (1 + \|\gamma_t\|^p + \|\|\eta_t\|\|_{L_2}^p),
\end{aligned}$$

and thus (71).

It remains to prove (73). Note that $(\delta Y, \delta Z) := (Y - Y', Z - Z')$ solves the following linearized BSDE

$$\begin{aligned}
(222) \quad \delta Y(s) &= \Phi - \Phi' + \int_s^T [f(\Theta_r, \mathcal{L}_{\Theta_r^{\eta_t}}) - f(\Theta'_r, \mathcal{L}_{\Theta'_r^{\eta_t}})] dr - \int_s^T \delta Z(r) dB(r) \\
&=: \delta \Phi + \int_s^T (\alpha_r \delta Y(r) + \beta_r \delta Z(r) + \delta h_r + \delta f_r) dr - \int_s^T \delta Z(r) dB(r),
\end{aligned}$$

where

$$\begin{aligned}
\alpha_r &:= \int_0^1 \partial_y f(B_r^{\gamma_t}, Y' + \theta(Y - Y'), Z, \mathcal{L}_{\Theta_r}) d\theta, \\
\beta_r &:= \int_0^1 \partial_z f(B_r^{\gamma_t}, Y', Z + \theta(Z - Z'), \mathcal{L}_{\Theta_r}) d\theta, \quad \delta \tilde{Y}^{\eta_t} := \tilde{Y}^{\eta_t} - \tilde{Y}'^{\eta_t}, \\
\delta h_r &:= f(B_r^{\gamma_t}, Y', Z', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(r)}) - f(B_r^{\gamma_t}, Y', Z', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y'^{\eta_t}(r)}), \quad \text{and} \\
\delta f_r &:= f(B_r^{\gamma_t}, Y', Z', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y^{\eta_t}(r)}) - f(B_r^{\gamma_t}, Y', Z', \mathcal{L}_{B_r^{\eta_t}}, \mathcal{L}_{Y'^{\eta_t}(r)}).
\end{aligned}$$

Using the Lipschitz continuity of f in (y, z) and standard estimate for linear BSDEs (see e.g. [4]), we have

$$(223) \quad \mathbb{E} \left[\sup_{s \in [t, T]} |\delta Y(s)|^p + \left(\int_t^T |\delta Z(s)|^2 ds \right)^{\frac{p}{2}} \right] \leq C_p \left(\mathbb{E} |\delta \Phi|^p + \mathbb{E} \left| \int_t^T (\delta h_r + \delta f_r) dr \right|^p \right).$$

Then, in view of estimate (72), we have

$$(224) \quad \mathbb{E} |\delta \Phi|^p + \mathbb{E} \left| \int_t^T \delta h_r dr \right|^p + \mathbb{E} \left| \int_t^T \delta f_r dr \right|^p \leq C_K (\|\gamma_t - \gamma'_t\|^p + W_2(\mathcal{L}_{\eta_t}, \mathcal{L}_{\eta'_t})^p),$$

and thus the desired estimate (73).

5.2. An extension of [41, Theorem 4.5] without assumption of local Lipschitz continuity in time variable.

Lemma 5.1. *Let non-anticipative functional $f : [0, T] \times \mathbb{D}_{T,d} \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ lie in $\mathcal{C}_s^{0,2,2,2}$. Assume that for any $t \in [0, T]$ and $\tau \leq t$, f and all its derivatives are locally Lipschitz continuous on $\mathbb{D}_{T,d}$: for $\phi(t, \cdot) = (I, \partial_{\omega_\tau}, \partial_{\omega_\tau}^2) f(t, \cdot, 0, 0)$,*

$$(225) \quad |\phi(t, \omega) - \phi(t, \omega')| \leq C(1 + \|\omega_t\|^k + \|\omega'_t\|^k)(\|\omega_t - \omega'_t\|), \quad \forall (\omega, \omega') \in \mathbb{D}_{T,d}^2,$$

for some constant C and integer k . Moreover, suppose that the first order derivatives in y and z , as well as their first order derivatives w.r.t. ω_τ , y and z are uniformly bounded. If Φ satisfies (79), there is a unique classical solution of the following PPDE

$$(226) \quad \begin{cases} \partial_t u(t, \gamma) + \frac{1}{2} \text{Tr} [\partial_{\omega}^2 u(t, \gamma)] + f(t, \gamma, u(t, \gamma), \partial_{\omega} u(t, \gamma)) = 0, \\ u(T, \gamma) = \Phi(\gamma_T), \quad (t, \gamma) \in [0, T] \times \mathbb{C}_{T,d}. \end{cases}$$

Proof. The uniqueness is a consequence of that of the following non-Markovian BSDE

$$(227) \quad Y^{\gamma_t}(s) = \Phi(B_T^{\gamma_t}) + \int_s^T f(r, B^{\gamma_t}, Y^{\gamma_t}(r), Z^{\gamma_t}(r)) dr - \int_s^T Z^{\gamma_t}(r) dB(r).$$

We now sketch the proof of the existence. Set

$$(228) \quad u(t, \gamma) := Y^{\gamma_t}(t).$$

Via a similar argument as in the proof of [41, Theorems 3.9 and 3.10], we have $u \in \mathcal{C}_{s,p}^{0,2}$ and moreover,

$$(229) \quad u(s, B^{\gamma_t}) = Y^{\gamma_t}(s), \quad \partial_{\omega_t} u(s, B^{\gamma_t}) = Z^{\gamma_t}(s), \quad s \geq t.$$

Then, applying the partial Itô formula (168), we have that for any $\delta > 0$,

$$\begin{aligned} & u(t + \delta, \gamma_t) - u(t, \gamma_t) \\ &= u(t + \delta, \gamma_t) - \mathbb{E}[u(t + \delta, B^{\gamma_t})] + \mathbb{E}[u(t + \delta, B^{\gamma_t})] - u(t, \gamma_t) \\ &= \mathbb{E} \left[- \int_t^{t+\delta} \partial_{\omega_r} u(t + \delta, B_r^{\gamma_t}) dB(r) - \frac{1}{2} \int_t^{t+\delta} \text{Tr}[\partial_{\omega_r}^2 u(t + \delta, B_r^{\gamma_t})] dr \right] \\ & \quad + \mathbb{E} \left[- \int_t^{t+\delta} f(r, B^{\gamma_t}, Y^{\gamma_t}(r), Z^{\gamma_t}(r)) dr \right]. \end{aligned}$$

Dividing both sides of the above identity by δ and taking $\delta \rightarrow 0$, we complete the proof. \square

REFERENCES

- [1] H. Ahn, Semimartingale integral representation *Ann. Probab.*, 25(2): 997-1010, 1997.
- [2] A. Bensoussan, *Stochastic Control of Partially Observable Systems*, Cambridge University Press, Cambridge, UK, 1992.
- [3] M. Bossy, Some stochastic particle methods for nonlinear parabolic PDEs, in GRIP-Research Group on Particle Interaction. *ESAIM Proceedings*, 15, 18-57, MR2441320, 2005.
- [4] P. Briand, B. Delyon, Y. Hu, E. Pardoux and L. Stoica. L^p -solutions of backward stochastic differential equations, *Stoch. Proc. App.* 108:109-129, 2003.
- [5] A. Bensoussan, J. Frehse, and P. Yam, The Master Equation in Mean Field Theory, <http://arxiv.org/abs/1404.4150>, 2014.
- [6] C. Bayer, P. Friz and J. Gatheral, Pricing under rough volatility, *Quantitative Finance*, 16, 887-904, 2016.
- [7] A. Bensoussan, J. Frehse, and P. Yam, *Mean Field Games and Mean Field Type Control Theory*, Springer Briefs in Mathematics, Springer, New York, 2013.
- [8] Rainer Buckdahn, Juan Li and Shige Peng, Mean-field backward stochastic differential equations and related partial differential equations, *Stoch. Proc. App.* 119:3133-3154, 2009.
- [9] Rainer Buckdahn, Juan Li, Shige Peng and Catherine Rainer, Mean-field stochastic differential equations and associated PDEs, *Ann. Probab.* 45(2):824-878, 2017.
- [10] J.-F. Chassagneux, D. Crisan and F. Delarue, A probabilistic approach to classical solutions of the master equation for large population equilibria, Preprint. Available at arXiv:1411.3009, 2015.
- [11] R. Carmona and F. Dularue, Probabilistic Theory of Mean Field Games with Applications. I: Mean Field FBSDEs, Control and Games. *Probability Theory and Stochastic Modelling* 83. Springer, Cham. MR3752669, 2018.
- [12] R. Carmona and F. Dularue, Probabilistic Theory of Mean Field Games with Applications. I: Mean Field FBSDEs, Control and Games. *Probability Theory and Stochastic Modelling* 84. Springer, Cham. MR3753660, 2018.
- [13] R. Carmona and F. Dularue, The master equation for large population equilibria, In *Stochastic Analysis and Applications*, Springer Proc. Math. Stat. 100, 77-128. Springer, Cham, 2014.
- [14] P. Cardaliaguet, F. Delarue, J.-M. Lasry and P.-L. Lions, The Master Equation and the Convergence Problem in Mean Field Games, *Annals of Mathematics Studies* 201. Princeton Univ. Press, Princeton, 2019.
- [15] P. Cardaliaguet, Notes on mean field games. Notes from P.L. Lions' lectures at the Collège de France, <https://www.ceremade.dauphine.fr/~cardalia/MFG100629.pdf>, 2012.
- [16] Rama Cont and David Fournié, Change of variable formula for non-anticipative functionals on path space, *Journal of Functional Analysis* 259:1043-1072, 2010.
- [17] Rama Cont and David Fournié, Functional Itô calculus and stochastic integral representation of martingales, *Ann. Probab.* 41:109-133, 2013.
- [18] A. Cosso, S. Federico, F. Gozzi, M. Rosestolato and N. Touzi, Path-dependent equations and viscosity solutions in infinite dimension, *Ann. Probab.* 46(1):126-174, 2018.
- [19] A. Conze and Viswanathan, Path-dependent options: the case of lookback options, *J. Finance* 46(5): 1893-1907, 1991.
- [20] B. Dupire, Functional Itô calculus, Bloomberg Portfolio Research Paper No. 2009-04-FRONTIERS, Available at SSRN: <https://ssrn.com/abstract=1435551>, 2009.
- [21] Ibrahim Ekren, Christian Keller, Nizar Touzi and Jianfeng Zhang, On viscosity solutions of path dependent pdes, *Ann. Probab.* 42(1):204-236, 2014.
- [22] I. Ekren, N. Touzi and J. Zhang, Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part I, *Ann. Probab.*, 44, 1212-1253, 2016.
- [23] I. Ekren, N. Touzi and J. Zhang, Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part II, *Ann. Probab.*, 44, 2507-2553, 2016.
- [24] M. Fischer, On the connection between symmetric N-player games and mean field games, *Ann. Appl. Probab.*, 27(2), 757-810, 2017.
- [25] F. Flandoli and G. Zanco, An infinite-dimensional approach to path-dependent Kolmogorov equations, *Ann. Probab.* 44(4):2643-2693, 2016.
- [26] E. Haug, *The complete guide to option pricing formulas*. Vol. 2., New York: McGraw-Hill, 2007.
- [27] O. Gueant, J-M Lasry and P-L Lions, Mean field games and applications, Paris-Princeton lectures on mathematical finance, 2010.

- [28] F. Gozzi and C. Marinelli, Stochastic optimal control of delay equations arising in advertising models, in Stochastic Partial Differential Equations and Applications VII, G. D. Prato and L. Tubaro, eds., Chapman & Hall/CRC, Press, Boca Raton, FL, 133-148, 2006.
- [29] J. Gatheral, T. Jaisson and M. Rosenbaum, Volatility is rough, *Quantitative Finance*, 18(6): 933-949, 2018.
- [30] D. A. Gomes and J. Saúde, Mean field games models - a brief survey, *Dynamic Games and Applications* volume 4, 110-154, 2014.
- [31] M. Huang, R. P. Malhamé and P. E. Caines, Large population stochastic dynamic games: Closed-loop McKean-Vlasov systems and Nash certainty equivalence principle, *Commun. Inf. Syst.* 6:221-251, 2006.
- [32] T. Ichiba and S. M. Mousavi, Option pricing with delayed information, preprint available at arXiv:1707.01600, 2017.
- [33] M. Kac, Foundation of kinetic theory, *Proc. Third Berkeley Symp. on Math. Stat. and Probab.* 3: 171-197, Univ. of Calif. Press, 1956.
- [34] D. Lacker, A general characterization of the mean field limit for stochastic differential games, *Probab. Theory Relat. Fields*, 165, 581-648, 2016.
- [35] J.-M. Lasry and P.-L. Lions, Mean field games, *Jpn. J. Math.* 2:229-260, 2007.
- [36] P. L. Lions, Cours au Collège de France, Available at www.college-de-france.fr.
- [37] H.P. McKean, Propagation of chaos for a class of non linear parabolic equations, in *Lecture Series in Differential Equations*, 7: 41-57, 1967.
- [38] D. Nualart, The Malliavin calculus and related topics, *Probability and its Applications (New-York)*, Springer-Verlag, Berlin, second ed., 2006.
- [39] S. Peng, Nonlinear Expectations and Stochastic Calculus under Uncertainty-with Robust CLT and G-Brownian Motion, *Probability Theory and Stochastic Modelling*, Volume 95, Springer, Berlin, 2019.
- [40] S. Peng, Backward stochastic differential equation, nonlinear expectation and their applications, In *Proceedings of International Congress of Mathematicians*, Volume I 393-432, Hindustan Book Agency, New Delhi, MR2827899, 2010.
- [41] S. Peng and F. Wang, BSDE, path-dependent PDE and nonlinear Feynman-Kac formula, *Science China Math.* 59(1), 19-36, 2016.
- [42] H. Pham and X. Wei, Bellman equation and viscosity solutions for mean-field stochastic control problem, *ESAIM Control Optim. Calc. Var.*, 24, 437-461, 2018.
- [43] S. Peng and Y. Song, G -expectation weighted Sobolev spaces, backward SDE and path dependent PDE, *J. Math. Soc. Japan*, 67(4): 1725-1757, 2015.
- [44] Y. F. Saporito, Stochastic Control and Differential Games with Path-Dependent Influence of Controls on Dynamics and Running Cost *SIAM J. Control Optim.*, 57(2), 1312-1327, 2019.
- [45] A.S. Sznitman, Nonlinear reflecting diffusion process, and the propagation of chaos and fluctuations associated, *J.F.A.* 56, 311-336, 1984.
- [46] A.S. Sznitman, A fluctuation result for nonlinear diffusions, in *Infinite dimensional Analysis and Stochastic Processes*, (S. Albeverio, ed.), Pitman, 145-160, 1985.
- [47] A.S. Sznitman, A propagation of chaos result for Burgers's equation, *Probability Theory and Related Fields* 71,581-613, 1986.
- [48] A.S. Sznitman, Topics in propagation of chaos, *Ecole d'ite de Probabilites de Saint-Flour XIX - 1989*, *Led. Notes in Math.*, 1464, Springer, 1991.
- [49] Y. F. Saporito, and J. Zhang, Stochastic control with delayed information and related nonlinear master equation, *SIAM J. Control Optim.*, 57, 693-717, 2019.
- [50] Cong Wu and Jianfeng Zhang, Viscosity solutions to parabolic master equations and McKean-Vlasov SDEs with closed-loop controls, *Ann. App. Prob.* 30(2), 936-986, 2020.
- [51] J. Zhang, *Backward Stochastic Differential Equations—From Linear to Fully Nonlinear Theory*, Springer, New York, 2017.

S. TANG, SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, HANDAN ROAD 220, 200433, SHANGHAI, PRC.

Email address: sjtang@fudan.edu.cn

H. ZHANG, RESEARCH CENTER FOR MATHEMATICS AND INTERDISCIPLINARY SCIENCES, SHANDONG UNIVERSITY, BINHAI ROAD 72, 266237, QINGDAO, PRC.

Email address: huilinzhang@sdu.edu.cn