arXiv:2109.06001v1 [nlin.SI] 13 Sep 2021

Meshy soliton structures for (2+1)-dimensional integrable systems and interactions

Shoufeng Shen, Guofang Wang, Yongyang Jin *

Department of Applied Mathematics, Zhejiang University of Technology, Hangzhou 310023, China

Abstract:

In this letter, we construct new meshy soliton structures by using two concrete (2+1)-dimensional integrable systems. The explicit expressions based on corresponding Cole-Hopf type transformations are obtained. Constraint equation $f_t + \sum_{j=1}^{N} h_j(y) f_{jx} = 0$ shows that these meshy soliton structures can be linear or parabolic. Interaction between meshy soliton structure and Lump structure are also revealed.

Keywords: Meshy soliton structure; integrable system; Cole-Hopf transformation; Lump

PACS numbers:

1 Introduction

The concept of soliton introduced by Zabusky and Kruskal, is originally used to describe the collision of two localized KdV solitary waves in FPU problem. Nowadays, solitons are receiving much attention in many natural sciences such as biology, chemistry, mathematics, communication, and especially in almost all branches of physics like condense matter physics, quantum field theory, plasma physics, fluid mechanics and nonlinear optics. Usually, one considers that the solitons are the basic excitations of integrable systems while the chaos and fractals are the basic behaviors of non-integrable systems. Thus, one of the most important research fields in soliton theory is to seek for exact solutions of integrable systems and use these solutions to simulate various natural science phenomena [1–6].

Recently, many new (2+1)-dimensional soliton structures such as dromions, lumps, ring solitons, breathers, instantons, compactons have been obtained. These studies are restricted in the single-valued situations. For more complicated cases, multivalued-functions also have been used to construct folded solitary waves and foldons. For example, the dromion structures which decay exponentially in all directions, can be obtained by two non-perpendicular line solutions for

* Corresponding Author

the Kadomtsev-Petviashvilli (KP) equation, while for the Davey-Stewartson (DS) and Nizhnik-Novikov-Veselov (NNV) equations, the dromion solutions are constructed by two perpendicular line solutions.

In this letter, we construct new soliton structures, which are called the meshy soliton structures by using two concrete (2+1)-dimensional integrable systems. It should be emphasized that the meshy soliton structure was first named and studied by Prof. Yong Chen. The obtained results may be able to simulate and explain the water wave phenomenon in Internet figure 1, which has happened on a sea surface in France.



Figure 1: Meshy wave on a sea surface in France.

2 Meshy soliton structures and interactions

We focus on the following coupled nonlinear system [7, 8],

$$\begin{cases} u_t - 2uu_x - v_{xx} = 0, \\ v_{yt} - u_{xxy} - 2uv_{xy} - 2u_x v_y = 0, \end{cases} \tag{1}$$

which deserves the name as a coupled Burgers system because by setting $u = v$ and $x = y$, it is reduced to the standard Burgers equation. Substituting

$$\begin{cases} u = \frac{f_x}{f} + h(y), \\ v = u + \alpha, \end{cases} \tag{2}$$

into the above system, we have the following linear equation,

$$f_t - 2h(y)f_x - f_{xx} = 0. \tag{3}$$

Here $f \equiv f(x, y, t)$ and α is an arbitrary constant. Obviously, Eq.(2) is a Cole-Hopf type transformation which transform the original coupled Burgers system (1) to the heat conduction type equation (3). Moreover, another integrable coupled system of the modified KdV equation and the

potential Boiti-Leon-Manna-Pempinelli equation (mKdV-pBMLP) is given in Refs.[7, 8] as

$$\begin{cases} u_t + 3u^2u_x + u_{xxx} + \frac{3}{2}(uv_x)_x + \frac{3}{2}\partial_y^{-1}(uv_y)_{xx} = 0, \\ v_{yt} + \frac{3}{2}(v_xv_y)_x + v_{xxxy} + 3(u^2v_y)_x + \frac{3}{2}(uu_y)_{xx} + \frac{3}{2}(uu_{xy})_x = 0. \end{cases} \tag{4}$$

By means of Eq.(2), we have the following linear equation

$$f_t + 3h^2(y)f_x + 3h(y)f_{xx} + f_{xxx} = 0. \tag{5}$$

Thus we can construct new meshy soliton structures for the physical quantity

$$U = \gamma(\ln f)_{xy} = \gamma\frac{f_{xy}f - f_xf_y}{f^2}, \tag{6}$$

via suitable selections of function f in Eq.(3) and Eq.(5). Here γ is a constant, just for the sake of computer simulation and drawing. In this letter, we can take the value of γ to be -1.

If we set $f = 1 + \sum_{j=1}^{N} c_j e^{a_jx+b_jy+a_j^2t+d_j}$ in Eq.(3) with $h(y) = 0$, where a_j, b_j, c_j, d_j are appropriate constants, then we have dromion structure or multiple solitoff structure. Here we call a half-straight line soliton structure as a solitoff. These conclusions have been presented in many literatures. Thanks to the arbitrariness of the constants b_j, we can construct new Meshy soliton structure displayed in Figures 2-7 by setting

$$\begin{aligned} f_1 = 1 &+ e^{-4x+y+16t} + e^{4x-y+16t-30} + e^{-12x+3y+144t-50} + e^{x-4y+t} + e^{-3x-3y+9t} \\ &+ e^{5x-5y+25t-30} + e^{-11x-y+121t-50} + e^{-x+4y+t-30} + e^{-5x+5y+25t-30} \\ &+ e^{3x+3y+9t-60} + e^{-13x+7y+169t-80} + e^{3x-12y+9t-50} + e^{-x-11y+t-50} \\ &+ e^{7x-13y+49t-80} + e^{-9x-9y+81t-100}. \end{aligned} \tag{7}$$

From Figures 2-4 to Figures 5-7, we can see that the shape of meshy soliton structure is most regular as the time variable goes to 0 and becoming a parallelogram type soliton structure. Theoretically, we give the interpretation of water wave simulation in Figure 1.




Figure 2: Meshy kink-soliton structure of the field function $-u$ in Eq.(2) by using f_1 at $t = -1$.




Figure 3: Meshy soliton structure of U by using f_1 at $t = -1$.




Figure 4: Density plot of U by using f_1 at $t = -1$.




Figure 5: Meshy kink-soliton structure of the field function $-u$ in Eq.(2) by using f_1 at $t = 0$




Figure 6: Meshy soliton structure of U by using f_1 at $t = 0$.




Figure 7: Density plot of U by using f_1 at $t = 0$.

To consider furthermore, we write another kind of meshy soliton structure by using the following functional expression (8), which is composed of linear solitons and parabolic solitons.

$$\begin{aligned} f_2 = 1 &+ e^{x-4y+t} + e^{-x+4y+t-30} + e^{3x-12y+9t-50} + e^{-4x+y^2+16t} + e^{-3x-4y+y^2+9t} \\ &+ e^{-5x+4y+y^2+25t-30} + e^{-x-12y+y^2+t-50} + e^{4x-y^2+16t-30} + e^{5x-4y-y^2+25t-30} \\ &+ e^{3x+4y-y^2+9t-60} + e^{7x-12y-y^2+49t-80} + e^{-12x+3y^2+144t-50} + e^{-11x-4y+3y^2+121t-50} \\ &+ e^{-13x+4y+3y^2+169t-80} + e^{-9x-12y+3y^2+81t-100}. \end{aligned} \tag{8}$$

For the sake of readability and simplicity, we present only two figures to illustrate this soliton structure (see Figures 8-9).




Figure 8: Meshy soliton structure of U by using f_2 at $t = 0$




Figure 9: Density plot of U by using f_2 at $t = 0$

Finally, we construct the interaction behavior between meshy soliton structure and Lump structure by using the following functional expression (9),

$$f_3 = f_1 + x^2 + 2t + y^2 - \frac{39}{20}. \tag{9}$$

Figures 9-10 show the interaction behavior at $t = 0$, which has parallelogram type soliton-Lump structure. For $f_4 = f_2 + x^2 + 2t + y^2 - \frac{39}{20}$, we have an almost similar result.




Figure 10: Interaction behavior between meshy soliton structure and Lump structure of U by using f_3 at $t = 0$.




Figure 11: Density plot of U by using f_3 at $t = 0$.

Remark 1. *The choice of function $f \equiv f(x, y, t)$ depends on the constraint equations* (3) *or* (5). *In fact, starting from the following generalized constraint equation*

$$f_t + \sum_{j=1}^{N} h_j(y) f_{jx} = 0 \tag{10}$$

and corresponding Cole-Hopf type transformation, we can construct some new integrable systems from the inverse problem point of view. We can also consider differential-difference equations and nonlocal constraint equations such as $f_t + h(y) f_x(-x) + f_{xxx} = 0$.

Remark 2. *We can also study meshy peakon structure and meshy loop-soliton structure.*

3 Conclusions

We considered the (2+1)-dimensional coupled Burgers and mKdV-pBLMP system, which are reduced to linear constraint equations (3) and (5). Although the construction of (2+1)-dimensional soliton structures is more difficult than that of (1+1)-dimensional soliton structures [9, 10], new meshy soliton structures represented by Figures 2-8 for the physical quantity U are obtained, which can be linear or parabolic. These results may be used to simulate the water wave phenomenon in Figure 1, which has happened on a sea surface in France. In Figures 9-10, interaction between meshy soliton structure and Lump structure are also revealed.

Acknowledgments

The authors thank Prof. Yong Chen of East China Normal University for helpful discussions. The work was supported by the National Natural Science Foundation of China (11771395).

References

[1] E. Infeld, A. Senatorski, A.A. Skorupski, Decay of Kadomtsev-Petviashvili solitons, Phys. Rew. Lett. 72 (1994) 1345-1347.

[2] V.N. Serkin, A. Hasegawa, Novel soliton solutions of the nonlinear Schrödinger equation model, Phys. Rew. Lett. 85 (2000) 4502-4505.

[3] F. Lu, Q. Lin, W.H. Knox, Govind P. Agrawal, Vector soliton fission, Phys. Rew. Lett. 93 (2004) 183901(4pp).

[4] Jiefang Zhang, Pin Han, New multisoliton solutions of the(2+1)-dimensional dispersive long wave equations, Commun. Nonl. Sci. Nume. Simu. 6 (2001) 178-182.

[5] A. Maccari, Non-resonant interacting water waves in 2+1 dimensions, Chaos, Solitons and Fractals 14 (2002) 105-116.

[6] Song Wang, Xiaoyan Tang, Senyue Lou, Soliton fission and fusion: Burgers equation and Sharma-Tasso-Olver equation, Chaos, Solitons and Fractals 21 (2004) 231-239.

[7] Jianyong Wang, Zufeng Liang, Xiaoyan Tang, Infinitely many generalized symmetries and Painlevé analysis of a (2+1)-dimensional Burgers system, Phys. Scr. 89 (2014) 025201 (6pp).

[8] A.M. Wazwaz, Multiple kink solutions for two coupled integrable (2 + 1)-dimensional systems, Appl. Math. Let. 58 (2016) 1-6.

[9] Shoufeng Shen, Yongyang Jin, Jun Zhang, Bäcklund Transformations and solutions of some generalized nonlinear evolution equations, , Rep. Math. Phys. 73 (2014) 225-279.

[10] Gaizhu Qu, Xiaorui Hu, Zhengwu Miao, Shoufeng Shen, Mengmeng Wang, Soliton molecules and abundant interaction solutions of a general high-order Burgers equation, Results in Physics 23 (2021) 104052(7pp).