

THE POSITIVE ENERGY CONJECTURE FOR A CLASS OF AHM METRICS ON $\mathbb{R}^2 \times \mathbb{T}^{n-2}$

ZHUOBIN LIANG[†] AND XIAO ZHANG^b

ABSTRACT. We prove the positive energy conjecture for a class of asymptotically Horowitz-Myers (AHM) metrics on $\mathbb{R}^2 \times \mathbb{T}^{n-2}$. This generalizes the previous results of Barzegar-Chruściel-Hörzinger-Maliborski-Nguyen [1] as well as the authors [5].

1. INTRODUCTION

In [4], Horowitz-Myers constructed so-called AdS solitons with toroidal topology

$$-r^2 dt^2 + \frac{1}{\frac{r^2}{\ell^2} \left(1 - \frac{\check{r}_0^n}{r^n}\right)} dr^2 + \frac{r^2}{\ell^2} \left(1 - \frac{\check{r}_0^n}{r^n}\right) d\xi^2 + r^2 \sum_{i=3}^n (d\phi^i)^2$$

which are globally static vacuum $(n+1)$ -dimensional spacetime with cosmological constant

$$\Lambda = -\frac{n(n-1)}{2\ell^2}.$$

For convenience, we assume $\ell = 1$ throughout the paper. These metrics are geodesically complete when

$$r \in [\check{r}_0, \infty), \quad \xi \in [0, \beta_0], \quad \phi^i \in [0, \lambda_i]$$

for constants $\check{r}_0 > 0$, $\lambda_i > 0$, $i = 3, \dots, n$ and

$$\beta_0 = \frac{4\pi}{n\check{r}_0}. \quad (1.1)$$

The induced Riemannian metrics g_{HM} on the constant time slices

$$g_{\text{HM}} = \frac{1}{r^2 \left(1 - \frac{\check{r}_0^n}{r^n}\right)} dr^2 + r^2 \left(1 - \frac{\check{r}_0^n}{r^n}\right) d\xi^2 + r^2 \sum_{i=3}^n (d\phi^i)^2 \quad (1.2)$$

are asymptotically locally hyperbolic (ALH) and are referred as Horowitz-Myers metrics. Horowitz-Myers also verified that the Hawking-Horowitz mass of g_{HM} is negative and conjectured that, among all metrics which are asymptotic to g_{HM} with the same period β_0 for ξ and with the scalar curvature

$$R \geq -n(n-1), \quad (1.3)$$

g_{HM} is the unique metric with the least Hawking-Horowitz mass [4].

Throughout the paper, we misuse the notation and refer a function f type $O(\frac{1}{r^m})$ if for *all* $k \geq 0$, its k th partial derivative with respect to r satisfies

$$\partial_r^k f \sim \frac{1}{r^{m+k}}, \quad r \rightarrow \infty. \quad (1.4)$$

Furthermore, a tensor is referred as type $O(\frac{1}{r^m})$ if its components belong to $O(\frac{1}{r^m})$ in coordinates $\{r, \xi, \phi^i\}$.

As both Schoen-Yau's method and Witten's method are hard to apply in this situation, there is less progress towards proof of the conjecture. In 2020, Barzegar-Chruściel-Hörzinger-Maliborski-Nguyen studied the following asymptotically Horowitz-Myers metrics on $\mathbb{R}^2 \times \mathbb{T}^{n-2}$

$$g_1 = e^{2u(r)} dr^2 + e^{2v(r)} d\xi^2 + e^{2w(r)} \sum_{i=3}^n (d\phi^i)^2, \quad (1.5)$$

where u, v, w are functions of r which satisfy

$$\begin{aligned} u(r) &= -\ln r - \frac{1}{2} \ln \left(1 - \frac{\check{r}_0^n}{r^n} \right) + \frac{u_n}{r^n} + O\left(\frac{1}{r^{n+1}}\right), \\ v(r) &= \ln r + \frac{1}{2} \ln \left(1 - \frac{\check{r}_0^n}{r^n} \right) + \frac{v_n}{r^n} + O\left(\frac{1}{r^{n+1}}\right), \\ w(r) &= \ln r + \frac{w_n}{r^n} + O\left(\frac{1}{r^{n+1}}\right), \end{aligned} \quad (1.6)$$

for constants u_n, v_n and w_n . The metric g_1 is asymptotic to g_{HM} up to order $O(\frac{1}{r^n})$. In [1], they verified Horowitz-Myers conjecture for g_1 if its scalar curvature satisfies (1.3) with $\ell = 1$.

Let a be certain constant and r_+ be the largest positive root of the equation

$$1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} = 0$$

where constant $r_0 > 0$. Denote

$$\beta = \frac{4\pi}{r_+ \left(n - 1 + \frac{r_0^n}{r_+^n} \right)}. \quad (1.7)$$

In [5], we construct metrics of Horowitz-Myers type with the negative constant scalar curvature

$$\dot{g} = \frac{1}{r^2 \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right)} dr^2 + r^2 \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right) d\xi^2 + r^2 \sum_{i=3}^n (d\phi^i)^2, \quad (1.8)$$

where

$$r \in [r_+, \infty), \quad \xi \in [0, \beta], \quad \phi^i \in [0, \lambda_i].$$

This is a geodesically complete metric on $\mathbb{R}^2 \times \mathbb{T}^{n-2}$ of constant scalar curvature $-n(n-1)$. And it is asymptotic to g_{HM} up to order $O(\frac{1}{r^{n-1}})$ provided $\beta_0 = \beta$. As \dot{g} decays weaker than g_1 , we can not apply their positive energy

theorem to \hat{g} directly. However, we can still verify that the Horowitz-Myers conjecture holds for \hat{g} [5].

Now we study the following more general perturbation of Horowitz-Myers metrics on $\mathbb{R}^2 \times \mathbb{T}^{n-2}$

$$g = e^{2u(r)} dr^2 + e^{2v(r, \xi)} d\xi^2 + r^2 \left(\sum_{i=3}^n (d\phi^i)^2 + \hat{w}(r, \xi, \phi^i) \right), \quad (1.9)$$

where

$$\begin{aligned} u(r) &= -\ln r - \frac{1}{2} \ln \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right) + \hat{u}(r), \\ v(r, \xi) &= \ln r + \frac{1}{2} \ln \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right) + \hat{v}(r, \xi), \\ e^{\hat{u}(r)} &= 1 + \frac{u_{n-1}}{r^{n-1}} + \frac{u_n}{r^n} + O\left(\frac{1}{r^{n+1}}\right), \\ e^{\hat{v}(r, \xi)} &= 1 + \frac{v_{n-1}(\xi)}{r^{n-1}} + \frac{v_n(\xi)}{r^n} + O\left(\frac{1}{r^{n+1}}\right), \\ \hat{w}(r, \xi, \phi^i) &= \frac{2w_{n-1}(\xi, \phi^i)}{r^{n-1}} + \frac{2w_n(\xi, \phi^i)}{r^n} + O\left(\frac{1}{r^{n+1}}\right), \end{aligned} \quad (1.10)$$

and u_{n-1} , u_n are real constants, $v_{n-1}(\xi)$, $v_n(\xi)$ are smooth functions of ξ , $w_{n-1}(\xi, \phi^i)$ and $w_n(\xi, \phi^i)$ are 2-tensors on \mathbb{T}^{n-2} which depend on ξ, ϕ^i . The restriction of g on \mathbb{T}^{n-2} is denoted as

$$\gamma = r^2 \left(\sum_{i=3}^n (d\phi^i)^2 + \hat{w}(r, \xi, \phi^i) \right).$$

In this paper, we show that the Horowitz-Myers conjecture is true for the above AHM metrics.

Theorem 1.1. *Let $n \geq 3$ and let g be the smooth metric (1.9) on $\mathbb{R}^2 \times \mathbb{T}^{n-2}$, which satisfies (1.10). Suppose the scalar curvature $R(g)$ of g satisfies*

$$R(g) + n(n-1) \geq 0, \quad [R(g) + n(n-1)]r \in L^1(\mathbb{R}^2 \times \mathbb{T}^{n-2}, g), \quad (1.11)$$

and suppose

$$\beta_0 = \beta. \quad (1.12)$$

If $n \geq 4$, we assume in addition that the scalar curvature $R(\gamma)$ of γ satisfies

$$\int_{\mathbb{T}^{n-2}} R(\gamma) d\phi^3 \cdots d\phi^n \leq 0 \quad (1.13)$$

for all r and ξ . Then the total energy

$$E(g) - E(g_{HM}) \geq 0.$$

The equality holds if and only if g is isometric to g_{HM} .

Remark 1.1. *The assumption that (1.4) holds for all $k \geq 0$ is only for the sake of simplicity of notations appeared in the paper. Indeed, the proof of Theorem 1.1 needs only $k = 0, 1, 2$.*

Remark 1.2. Condition (1.13) is trivial when $n = 3$, since in this case, γ is a one dimensional metric with $R(\gamma) \equiv 0$. If $n = 4$, then the Gauss-Bonnet theorem states that

$$\int_{\mathbb{T}^2} R(\gamma) dV_\gamma = 0,$$

which can not implies (1.13).

Remark 1.3. In [1], the components of γ are assumed to be independent of ϕ^i (e.g. (1.5)). In this case, γ is a flat metric on \mathbb{T}^{n-2} and (1.13) holds for all n .

The paper is organized as follows. In Section 2, we show that g can be written as an asymptotically Poincaré-Einstein (APE) metric. Then the total energy can be computed in terms of the definitions of Wang [6] and Chruściel-Herzlich [2]. In Section 3, we prove the Horowitz-Myers conjecture for g .

2. APE METRICS

In this section, we show that g is actually an APE metric. The boundary-at-infinity $\partial_\infty M$ of $\mathbb{R}^2 \times \mathbb{T}^{n-2}$ is \mathbb{T}^{n-1} . Let x be a special defining function for $\partial_\infty M$. That is, on a collar neighborhood of $\partial_\infty M$,

$$(1) \ x \geq 0 \text{ with } \{x = 0\} = \partial_\infty M,$$

$$(2) \ |dx|_{x^2g} \equiv 1.$$

On this collar neighborhood, the metric g can be written in the form

$$g = \frac{1}{x^2}(dx^2 \oplus h_x), \quad \lim_{r \rightarrow \infty} xr = 1$$

where h_x are the induced metrics of x^2g on the constant x slices,

$$\frac{1}{x}dx = -\frac{e^{\hat{u}(r)}}{r\sqrt{1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n}}}dr.$$

It follows that

$$\begin{aligned} \ln x &= -\int_{r_+}^r \frac{e^{\hat{u}(s)}}{s\sqrt{1 + \frac{a}{s^{n-1}} - \frac{r_0^n}{s^n}}}ds + C \\ &= -\ln r - \int_{r_+}^r \left(\frac{u_{n-1} - \frac{1}{2}a}{s^n} + \frac{u_n + \frac{1}{2}r_0^n}{s^{n+1}} + O(s^{-n-2}) \right) ds \\ &\quad + \ln r_+ + C, \end{aligned}$$

where

$$C = -\ln r_+ + \int_{r_+}^{\infty} \frac{1}{s} \left(\frac{e^{\hat{u}(s)}}{\sqrt{1 + \frac{a}{s^{n-1}} - \frac{r_0^n}{s^n}}} - 1 \right) ds.$$

Therefore, we obtain

$$r = x^{-1} + \frac{u_{n-1} - \frac{1}{2}a}{n-1}x^{n-2} + \frac{u_n + \frac{1}{2}r_0^n}{n}x^{n-1} + O(x^n).$$

Putting this into (1.9) and applying the asymptotic conditions (1.10), we obtain

$$\begin{aligned} g = \frac{1}{x^2} & \left\{ dx^2 + d\xi^2 + \sum_{i=3}^n (d\phi^i)^2 \right. \\ & + \frac{x^{n-1}}{n-1} \left[((n-2)a + 2u_{n-1} + 2(n-1)v_{n-1})d\xi^2 \right. \\ & + (2u_{n-1} - a) \sum_{i=3}^n (d\phi^i)^2 + 2(n-1)w_{n-1} \Big] \\ & + \frac{x^n}{n} \left[(-(n-1)r_0^n + 2u_n + 2nv_n)d\xi^2 \right. \\ & + (r_0^n + 2u_n) \sum_{i=3}^n (d\phi^i)^2 + 2nw_n \Big] + O(x^{n+1}) \Big\}. \end{aligned} \quad (2.1)$$

Denote

$$\begin{aligned} h_0 &= d\xi^2 + \sum_{i=3}^n (d\phi^i)^2, \\ \theta &= \left((n-2)a + 2u_{n-1} + 2(n-1)v_{n-1} \right) d\xi^2 \\ &\quad + (2u_{n-1} - a) \sum_{i=3}^n (d\phi^i)^2 + 2(n-1)w_{n-1}, \\ \kappa &= \left(-(n-1)r_0^n + 2u_n + 2nv_n \right) d\xi^2 \\ &\quad + (r_0^n + 2u_n) \sum_{i=3}^n (d\phi^i)^2 + 2nw_n. \end{aligned}$$

The conformal infinity of g is defined as the conformal class $[h_0]$ on $\partial_\infty M$ (cf. [3]). The mass aspect function, the total energy are defined as

$$\text{tr}_{h_0} \kappa = -r_0^n + 2(n-1)u_n + 2nv_n + 2n\text{tr}_{h_0} w_n, \quad (2.2)$$

$$E(g) = \int_{\partial_\infty M} \text{tr}_{h_0} \kappa dV_{h_0} \quad (2.3)$$

respectively [6, 2]. When $u_n = v_n = w_n = 0$, we obtain

$$E(g_{\text{HM}}) = -\check{r}_0^n \text{Vol}(\partial_\infty M, h_0).$$

Therefore

$$E(g) - E(g_{\text{HM}}) = \int_{\partial_\infty M} \left[\check{r}_0^n - r_0^n + 2(n-1)u_n + 2nv_n + 2n\text{tr}_{h_0} w_n \right] dV_{h_0}. \quad (2.4)$$

Now we express the scalar curvature $R(g)$ in some appropriate form. Denote

$$W^r = \frac{1}{2} \text{tr}_\gamma (\partial_r \gamma) = \frac{1}{2} (\partial_r \gamma_{ij}) \gamma^{ij}$$

and let \mathring{W}^r be the trace-less part of $\partial_r \gamma$. It is straightforward that

$$\partial_r \gamma = \frac{2W^r}{n-2} \gamma + \mathring{W}^r. \quad (2.5)$$

By the asymptotic condition of γ , we obtain

$$W^r = \frac{n-2}{r} - \frac{(n-1)\text{tr}_{h_0} w_{n-1}}{r^n} - \frac{n\text{tr}_{h_0} w_n}{r^{n+1}} + O(r^{-n-2}). \quad (2.6)$$

Similarly, denote

$$W^\xi = \frac{1}{2} \text{tr}_\gamma (\partial_\xi \gamma) = \frac{1}{2} (\partial_\xi \gamma_{ij}) \gamma^{ij}$$

and let \mathring{W}^ξ be the trace-less part of $\partial_\xi \gamma$. We can also get

$$\partial_\xi \gamma = \frac{2W^\xi}{n-2} \gamma + \mathring{W}^\xi. \quad (2.7)$$

Set

$$\hat{W}^r = W^r - \frac{n-2}{r}. \quad (2.8)$$

Lemma 2.1. *With the above notations, the scalar curvature $R(g)$ of g is*

$$\begin{aligned} R(g) = & -2e^{-u-v} \partial_r \left[e^{v-u} (\partial_r v + W^r) \right] - 2e^{-v} \partial_\xi (e^{-v} W^\xi) \\ & + R(\gamma) - 2e^{-2u} \left[\frac{n-1}{2(n-2)} (W^r)^2 + \frac{1}{8} |\mathring{W}^r|_\gamma^2 \right] \\ & - 2e^{-2v} \left[\frac{n-1}{2(n-2)} (W^\xi)^2 + \frac{1}{8} |\mathring{W}^\xi|_\gamma^2 \right]. \end{aligned} \quad (2.9)$$

Proof. For the following metrics

$$N(s)^2 ds^2 + \eta, \quad \eta = \eta_{ab}(s, y^1, \dots, y^m) dy^a dy^b,$$

it is well-known that their scalar curvature is

$$R(\eta) - N^{-1} (N^{-1} \text{tr}_\eta \eta')' - \frac{1}{4N^2} |\eta'|_\eta^2 - \frac{1}{4N^2} (\text{tr}_\eta \eta')^2, \quad (2.10)$$

where $R(\eta)$ is the scalar curvature of η , $'$ is the partial derivative with respect to s ,

$$\eta' = \eta'_{ab} dy^a dy^b.$$

Now we apply formula (2.10) to the following part of g

$$\rho = e^{2v} d\xi^2 + \gamma,$$

and obtain

$$\begin{aligned} R(\rho) &= R(\gamma) - e^{-v} \partial_\xi (e^{-v} \text{tr}_\gamma \partial_\xi \gamma) - \frac{1}{4} e^{-2v} |\partial_\xi \gamma|_\gamma^2 - \frac{1}{4} e^{-2v} (\text{tr}_\gamma \partial_\xi \gamma)^2 \\ &= R(\gamma) - 2e^{-v} \partial_\xi (e^{-v} W^\xi) - 2e^{-2v} \left[\frac{n-1}{2(n-2)} (W^\xi)^2 + \frac{1}{8} |\mathring{W}^\xi|_\gamma^2 \right]. \end{aligned}$$

Next we apply (2.10) to the metric

$$g = e^{2u} dr^2 + \rho$$

and obtain

$$R(g) = R(\rho) - e^{-u} \partial_r (e^{-u} \text{tr}_\rho \partial_r \rho) - \frac{1}{4} e^{-2u} |\partial_r \rho|_\rho^2 - \frac{1}{4} e^{-2u} (\text{tr}_\rho \partial_r \rho)^2.$$

Since

$$\begin{aligned} \partial_r \rho &= 2(\partial_r v) e^{2v} d\xi^2 + \partial_r \gamma, \\ \text{tr}_\rho \partial_r \rho &= 2(\partial_r v) + \text{tr}_\gamma \partial_r \gamma = 2(\partial_r v) + 2W^r, \\ |\partial_r \rho|_\rho^2 &= 4(\partial_r v)^2 + |\partial_r \gamma|_\gamma^2 = 4(\partial_r v)^2 + \frac{4}{n-2} (W^r)^2 + |\mathring{W}^r|_\gamma^2. \end{aligned}$$

we obtain

$$\begin{aligned} R(g) &= R(\rho) - 2e^{-u} \partial_r [e^{-u} (\partial_r v + W^r)] \\ &\quad - e^{-2u} \left[2(\partial_r v)^2 + \frac{n-1}{n-2} (W^r)^2 + 2\partial_r v W^r + \frac{1}{4} |\mathring{W}^r|_\gamma^2 \right] \\ &= R(\rho) - 2e^{-u-v} \partial_r [e^{v-u} (\partial_r v + W^r)] \\ &\quad - 2e^{-2u} \left[\frac{n-1}{2(n-2)} (W^r)^2 + \frac{1}{8} |\mathring{W}^r|_\gamma^2 \right]. \end{aligned}$$

Therefore the lemma follows. Q.E.D.

Proposition 2.1. *For the metric (1.9), the condition*

$$[R(g) + n(n-1)]r \in L^1(\mathbb{R}^2 \times \mathbb{T}^{n-2}, g)$$

holds if and only if

$$\text{tr}_{h_0} \theta = 0 \iff u_{n-1} + v_{n-1} + \text{tr}_{h_0} w_{n-1} = 0. \quad (2.11)$$

In this case,

$$R + n(n-1) = O\left(\frac{1}{r^{n+1}}\right). \quad (2.12)$$

Proof. Note if all \hat{u} , \hat{v} , \hat{W}^r , W^ξ , \mathring{W}^r and \mathring{W}^ξ vanish, then g reduces to

$$e^{-2f} dr^2 + e^{2f} d\xi^2 + r^2 \sum_{i,j=3}^n \gamma_{ij}(\phi^3, \dots, \phi^n) d\phi^i d\phi^j,$$

where

$$f = \ln r + \frac{1}{2} \ln \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right).$$

By (2.9), the scalar curvature of this metric is

$$-n(n-1) + R(\gamma).$$

Putting (1.10) and (2.8) into (2.9), we obtain

$$\begin{aligned} R(g) = & -n(n-1)e^{-2\hat{u}} + 2e^{2f-2\hat{u}} \left[-\partial_r(\partial_r \hat{v} + \hat{W}^r) \right. \\ & - \partial_r(\hat{v} - \hat{u} + 2f + (n-1)\ln r) \cdot (\partial_r \hat{v} + \hat{W}^r) \\ & + \left(\frac{1}{r} - \partial_r f \right) \partial_r \hat{v} + \left(\partial_r f + \frac{n-2}{r} \right) \partial_r \hat{u} \\ & \left. - \frac{(n-1)}{2(n-2)} (\hat{W}^r)^2 - \frac{1}{8} |\mathring{W}^r|_\gamma^2 \right] \\ & - 2e^{-v} \partial_\xi (e^{-v} W^\xi) + R(\gamma) \\ & - 2e^{-2v} \left[\frac{n-1}{2(n-2)} (W^\xi)^2 + \frac{1}{8} |\mathring{W}^\xi|_\gamma^2 \right] \\ = & -n(n-1) + (\text{tr}_{h_0} \theta) r^{1-n} + O(r^{-n-1}). \end{aligned}$$

Here we use that

$$R(\gamma) = O(r^{-n-1}),$$

which could be confirmed by direct computation. Since

$$\sqrt{\det g} = O(r^{n-2}),$$

and

$$\text{tr}_{h_0} \theta = 2(n-1)(v_{n-1} + u_{n-1} + \text{tr}_{h_0} w_{n-1}),$$

so the lemma follows. Q.E.D.

By (2.1), we have

$$g = x^{-2} \bar{g}, \quad \bar{g} = dx^2 + d\xi^2 + \sum_{i=3}^n (d\phi^i)^2 + \frac{x^{n-1}}{n-1} \theta + O(x^n).$$

We denote $\phi^0 = x$ and $\phi^1 = \xi$. Let a, b and c be indices ranging from 1 to n . Under the coordinates $\{\phi^0, \phi^1, \dots, \phi^n\}$, it holds

$$\bar{g}_{00} = 1, \quad \bar{g}_{0a} = 0, \quad \bar{g}_{ab} = \delta_{ab} + \frac{x^{n-1}}{n-1} \theta_{ab} + O(x^n).$$

It is direct to show that the Christoffel symbols of \bar{g} satisfy

$$\begin{aligned}\Gamma_{ab}^c &= O(x^{n-1}), \\ \Gamma_{ab}^0 &= -\frac{1}{2}x^{n-2}\theta_{ab} + O(x^{n-1}), \\ \Gamma_{a0}^b &= \Gamma_{0a}^b = \frac{1}{2}x^{n-2}\theta_{ab} + O(x^{2n-3}),\end{aligned}$$

and the others vanish. So the Ricci curvature of \bar{g} satisfies

$$\begin{aligned}\bar{R}_{00} &= -\frac{1}{2}(n-2)x^{n-3}\text{tr}_{h_0}\theta + O(x^{2n-4}), \\ \bar{R}_{0a} &= O(x^{n-2}), \\ \bar{R}_{ab} &= O(x^{n-2}).\end{aligned}$$

Denote by Ric the Ricci curvature of g . Let

$$\Omega = Ric + (n-1)g.$$

By the formulas of conformal transformation of the Ricci and scalar curvatures, we finally obtain

$$\begin{aligned}\Omega(\partial_x, \partial_x) &= -\frac{1}{2}(n-3)x^{n-3}\text{tr}_{h_0}\theta + O(x^{n-2}), \\ \Omega(\partial_a, \partial_b) &= \frac{1}{2}x^{n-3}(\text{tr}_{h_0}\theta)\delta_{ab} + O(x^{n-2}), \\ \Omega(\partial_x, \partial_a) &= O(x^{n-2}).\end{aligned}$$

Recall Definition 2.1 in [7] that the metric g is asymptotically Poincaré-Einstein (APE) if

$$\lim_{x \rightarrow 0} |\Omega|_g = O(x^n).$$

By Lemma 2.2 [7], g is APE if and only if $\text{tr}_{h_0}\theta = 0$. (This can also be seen from the above computation.) Therefore we have

Proposition 2.2. *For the metric (1.9), the condition*

$$[R(g) + n(n-1)]r \in L^1(\mathbb{R}^2 \times \mathbb{T}^{n-2}, g)$$

holds if and only if g is APE.

Finally, we study the consequence of smoothness of g at $r = r_+$.

Proposition 2.3. *If the metric g is smooth at $r = r_+$, then*

$$\hat{v}(r_+, \xi) = \hat{u}(r_+) \quad \text{for all } \xi. \quad (2.13)$$

Proof. We follow the proof of Theorem II.1 in [5] to construct a regular coordinate system around $r = r_+$. Set

$$A(r) = r^2 \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right)$$

for $r > 0$. Differentiating A at $r = r_+$ and using (1.7), we find

$$A'(r_+) = \frac{4\pi}{\beta} > 0.$$

This implies that A has a smooth inverse, which we denote by B , round r_+ . Clearly $B(0) = r_+$. Now we can introduce new coordinates (ρ, Ξ) given by

$$\rho = A^{\frac{1}{2}}(r), \quad \Xi = \frac{2\pi\xi}{\beta}$$

for $r_+ < r < r_+ + \varepsilon$, $0 \leq \Xi < 2\pi$, with some $\varepsilon > 0$. So (ρ, Ξ) can be viewed as the standard polar coordinates around the origin in \mathbb{R}^2 . Set

$$z^1 = \rho \cos \Xi, \quad z^2 = \rho \sin \Xi.$$

Obviously, $\{r = r_+\} = \{z^1 = z^2 = 0\}$. It is shown in [5] that $(z^1, z^2, \phi^3, \dots, \phi^n)$ is a regular coordinate system around $r = r_+$ for \dot{g} . Since g is smooth, the components in this coordinate system are smooth around $(z^1, z^2) = (0, 0)$. For our purpose, we only rewrite the following part

$$g_{(0)} := e^{2\hat{u}(r)} A^{-1}(r) dr^2 + e^{2\hat{v}(r, \xi)} A(r) d\xi^2$$

of g in the new coordinate system. Since

$$A^{-1}(r) dr^2 = \frac{4}{A'(r)^2} d\rho^2, \quad A(r) d\xi^2 = \frac{4\rho^2}{A'(r_+)^2} d\Xi^2,$$

we have

$$g_{(0)} = \frac{4e^{2\hat{u}(r)}}{A'(r)^2} d\rho^2 + \frac{4e^{2\hat{v}(r, \xi)}}{A'(r_+)^2} \rho^2 d\Xi^2 = g_{(\rho)} + g_{(\Xi)} + g_{\mathbb{R}^2}$$

where

$$\begin{aligned} g_{(\rho)} &= \left[\frac{4e^{2\hat{u}(r)}}{A'(r)^2} - \frac{4e^{2\hat{u}(r_+)}}{A'(r_+)^2} \right] d\rho^2, \\ g_{(\Xi)} &= \frac{4e^{2\hat{u}(r_+)}}{A'(r_+)^2} \left[e^{2(\hat{v}(r, \xi) - \hat{u}(r_+))} - 1 \right] \rho^2 d\Xi^2, \\ g_{\mathbb{R}^2} &= \frac{4e^{2\hat{u}(r_+)}}{A'(r_+)^2} (d\rho^2 + \rho^2 d\Xi^2). \end{aligned}$$

Since $\frac{4e^{2\hat{u}(r)}}{A'(r)^2}$ is smooth near $r = r_+$, there exists a smooth function $\varphi(r)$ so that

$$\frac{4e^{2\hat{u}(r)}}{A'(r)^2} - \frac{4e^{2\hat{u}(r_+)}}{A'(r_+)^2} = (r - r_+) \varphi(r)$$

for r near r_+ . Note that

$$(r - r_+) \varphi(r) = [B(\rho^2) - B(0)](\varphi \circ B)(\rho^2).$$

Since B is smooth near 0, there also exists a smooth function $\psi(\rho)$ such that

$$B(\rho^2) - B(0) = \rho^2 \psi(\rho^2).$$

So

$$\begin{aligned} g_{(\rho)} &= (\varphi \circ B)(\rho^2)\psi(\rho^2)\rho^2 d\rho^2 \\ &= ((\varphi \circ B) \cdot \psi)((z^1)^2 + (z^2)^2)(z^1 dz^1 + z^2 dz^2)^2. \end{aligned}$$

This implies that $g_{(\rho)}$ is smooth around $(z^1, z^2) = (0, 0)$. As a result, $g_{(\Xi)}$ is also smooth around $(z^1, z^2) = (0, 0)$. However,

$$\begin{aligned} \rho^2 d\Xi^2 &= (dz^1)^2 + (dz^2)^2 - d\rho^2 \\ &= (dz^1)^2 + (dz^2)^2 - \frac{1}{\rho^2}(z^1 dz^1 + z^2 dz^2)^2 \end{aligned}$$

is singular at $(z^1, z^2) = (0, 0)$. So it must hold that

$$\lim_{r \rightarrow r_+} \hat{v}(r, \xi) = \hat{u}(r_+)$$

for all ξ . This proves (2.13).

Q.E.D.

3. PROOF OF THE THEOREM

In this section, we adopt the idea in [1] to prove Theorem 1.1. We will replace r by a new coordinate function \tilde{r} to change $e^{2u(r)} dr^2$ in (1.9) to $\tilde{r}^{-2} \left(1 - \frac{\tilde{r}_0^n}{\tilde{r}^n}\right)^{-1} d\tilde{r}^2$.

Lemma 3.1. *There exist a positive number \tilde{r}_0 and a smooth increasing function $r \mapsto \tilde{r}(r)$ from $[r_+, \infty)$ to $[\tilde{r}_0, \infty)$, with $\tilde{r}(r_+) = \tilde{r}_0$, such that*

$$\frac{1}{\tilde{r} \sqrt{1 - \frac{\tilde{r}_0^n}{\tilde{r}^n}}} d\tilde{r} = \frac{1}{r \sqrt{1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n}}} e^{\hat{u}(r)} dr, \quad (3.1)$$

and

$$\tilde{r}(r) = r + \frac{a - 2u_{n-1}}{2(n-1)} r^{2-n} + \frac{\tilde{r}_0^n - r_0^n - 2u_n}{2n} r^{1-n} + O(r^{-n}). \quad (3.2)$$

Proof. Define

$$F(r) = \int_1^r \frac{1}{s \sqrt{1 - \frac{1}{s^n}}} ds, \quad r \geq 1.$$

For large r , we have

$$F(r) = F_0 + \ln r - \frac{1}{2n} r^{-n} + O(r^{-n-1}), \quad (3.3)$$

where

$$F_0 = \int_1^\infty \frac{1}{s} \left(\frac{1}{\sqrt{1 - \frac{1}{s^n}}} - 1 \right) ds.$$

Clearly, F is a smooth increasing function. Define $\tilde{r}(r)$ on $[r_+, \infty)$ by

$$F\left(\frac{\tilde{r}(r)}{\tilde{r}_0}\right) = \int_{r_+}^r \frac{e^{\hat{u}(s)}}{s \sqrt{1 + \frac{a}{s^{n-1}} - \frac{r_0^n}{s^n}}} ds. \quad (3.4)$$

Here \tilde{r}_0 is a positive number to be specified later. The integral on the right hand side has the following asymptotic expansion as $r \rightarrow \infty$:

$$C + \ln r + \frac{a - 2u_{n-1}}{2(n-1)} r^{1-n} - \frac{2u_n + r_0^n}{2n} r^{-n} + O(r^{-n-1}), \quad (3.5)$$

where

$$\begin{aligned} C &= \lim_{r \rightarrow \infty} \left(\int_{r_+}^r \frac{1}{s \sqrt{1 + \frac{a}{s^{n-1}} - \frac{r_0^n}{s^n}}} ds - \ln r \right) \\ &= -\ln r_+ + \int_{r_+}^{\infty} \frac{1}{s} \left(\frac{1}{\sqrt{1 + \frac{a}{s^{n-1}} - \frac{r_0^n}{s^n}}} - 1 \right) ds. \end{aligned}$$

Putting (3.3) and (3.5) into (3.4), we obtain

$$\begin{aligned} F_0 - \ln \tilde{r}_0 + \ln \tilde{r} - \frac{\tilde{r}_0^n}{2n} \tilde{r}^{-n} + O(\tilde{r}^{-n-1}) \\ = C + \ln r + \frac{a - 2u_{n-1}}{2(n-1)} r^{1-n} - \frac{2u_n + r_0^n}{2n} r^{-n} + O(r^{-n-1}). \end{aligned}$$

Now we set

$$\tilde{r}_0 = e^{F_0 - C}.$$

The above equation implies the asymptotic expansion (3.2). The equation (3.1) follows directly by the definition of \tilde{r} . Q.E.D.

With the function \tilde{r} , we rewrite the metric

$$g = e^{2\tilde{u}(\tilde{r})} d\tilde{r}^2 + e^{2\tilde{v}(\tilde{r}, \xi)} d\xi^2 + \tilde{r}^2 \left(\sum_{i=3}^n (d\phi^i)^2 + \hat{w}(\tilde{r}, \xi, \phi^i) \right), \quad (3.6)$$

where

$$\begin{aligned} \tilde{u}(\tilde{r}) &= -\ln \tilde{r} - \frac{1}{2} \ln \left(1 - \frac{\tilde{r}_0^n}{\tilde{r}^n} \right), \\ \tilde{v}(\tilde{r}, \xi) &= -\tilde{u}(\tilde{r}) + \hat{v}(\tilde{r}, \xi), \\ e^{\hat{v}(\tilde{r}, \xi)} &= 1 + \frac{\tilde{v}_{n-1}(\xi)}{\tilde{r}^{n-1}} + \frac{\tilde{v}_n(\xi)}{\tilde{r}^n} + O\left(\frac{1}{\tilde{r}^{n+1}}\right), \\ \hat{w}(\tilde{r}, \xi, \phi^i) &= \frac{2\tilde{w}_{n-1}}{\tilde{r}^{n-1}} + \frac{2\tilde{w}_n}{\tilde{r}^n} + O\left(\frac{1}{\tilde{r}^{n+1}}\right). \end{aligned} \quad (3.7)$$

By comparing (1.9) and (3.6), we have

$$\hat{\hat{v}} = \hat{v} + \hat{u} - \ln \frac{d\tilde{r}}{dr} \quad (3.8)$$

and

$$\hat{w} = \frac{r^2}{\tilde{r}^2} \hat{w} + \left(\frac{r^2}{\tilde{r}^2} - 1 \right) \sum_{i=3}^n (d\phi^i)^2. \quad (3.9)$$

In particular, we obtain

$$\begin{aligned} \tilde{v}_{n-1} &= v_{n-1} + \frac{(n-2)a + 2u_{n-1}}{2(n-1)}, \\ \tilde{v}_n &= v_n + \frac{(n-1)(\tilde{r}_0^n - r_0^n) + 2u_n}{2n}, \\ \tilde{w}_{n-1} &= w_{n-1} + \frac{2u_{n-1} - a}{2(n-1)} \sum_{i=3}^n (d\phi^i)^2, \\ \tilde{w}_n &= w_n + \frac{-(\tilde{r}_0^n - r_0^n) + 2u_n}{2n} \sum_{i=3}^n (d\phi^i)^2. \end{aligned} \quad (3.10)$$

As a result, (2.11) becomes

$$\tilde{v}_{n-1} + \text{tr}_{h_0} \tilde{w}_{n-1} = 0, \quad (3.11)$$

and (2.4) becomes

$$E(g) - E(g_{\text{HM}}) = (\tilde{r}_0^n - r_0^n) \text{Vol}(\partial_\infty M, h_0) + 2n \int_{\partial_\infty M} (\tilde{v}_n + \text{tr}_{h_0} \tilde{w}_n) dV_{h_0}. \quad (3.12)$$

Lemma 3.2. *Under the above condition, it holds that*

$$e^{\hat{v}(\tilde{r}_0, \xi)} = \frac{\tilde{r}_0}{\tilde{r}_0} \quad (3.13)$$

for all ξ .

Proof. By (3.8) and (2.13), we have

$$e^{\hat{v}(\tilde{r}_0, \xi)} = \frac{e^{2\hat{u}(r_+)}}{\frac{d\tilde{r}}{dr}(r_+)}.$$

Since (3.1) gives that

$$\left(\frac{d\tilde{r}}{dr} \right)^2 = \frac{\tilde{r}^2 \left(1 - \frac{\tilde{r}_0^n}{\tilde{r}^n} \right)}{r^2 \left(1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} \right)} e^{2\hat{u}(r)},$$

it follows

$$\left(\frac{d\tilde{r}}{dr} \right)^2 (r_+) = \frac{\tilde{r}_0^2}{r_+^2} e^{2\hat{u}(r_+)} \lim_{r \rightarrow r_+} \frac{1 - \frac{\tilde{r}_0^n}{\tilde{r}^n}}{1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n}} = \frac{n\tilde{r}_0 e^{2\hat{u}(r_+)} \frac{d\tilde{r}}{dr}(r_+)}{-\frac{(n-1)a}{r_+^{n-2}} + \frac{r_0^n}{r_+^{n-1}}}.$$

Recalling that r_+ is a root of $1 + \frac{a}{r^{n-1}} - \frac{r_0^n}{r^n} = 0$, this implies

$$\frac{d\tilde{r}}{dr}(r_+) = e^{2\hat{u}(r_+)} \frac{n\tilde{r}_0}{r_+(n-1 + \frac{1}{r_+^n})}.$$

By (1.12), we obtain

$$\frac{d\tilde{r}}{dr}(r_+) = e^{2\hat{u}(r_+)} \frac{\tilde{r}_0}{\tilde{r}_0}.$$

So the lemma follows. Q.E.D.

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Denote \tilde{W}^r , $\hat{\tilde{W}}^r$, $\tilde{\tilde{W}}^r$ for (3.6) the analogous quantities W^r , \hat{W}^r , \tilde{W}^r for (1.9). In the new coordinate function \tilde{r} , we have

$$\begin{aligned} R(g) = & -n(n-1) + R(\gamma) \\ & - 2e^{-\hat{v}-(n-1)\ln\tilde{r}}\partial_{\tilde{r}}\left[e^{-2\tilde{u}+\hat{v}}\tilde{r}^{n-1}(\partial_{\tilde{r}}\hat{v} + \hat{\tilde{W}}^r)\right] \\ & - \frac{n\tilde{r}_0^n}{\tilde{r}^{n-1}}\partial_{\tilde{r}}\hat{v} - 2e^{2\tilde{u}-\hat{v}}\partial_{\xi}(e^{-\hat{v}}\tilde{W}^{\xi}) \\ & - 2e^{-2\tilde{u}}\left[\frac{n-1}{2(n-2)}(\hat{\tilde{W}}^r)^2 + \frac{1}{8}|\tilde{\tilde{W}}^r|_{\gamma}^2\right] \\ & - 2e^{-2\tilde{v}}\left[\frac{n-1}{2(n-2)}(\tilde{W}^{\xi})^2 + \frac{1}{8}|\tilde{W}^{\xi}|_{\gamma}^2\right]. \end{aligned}$$

This equality can be rewritten as

$$\begin{aligned} -2\partial_{\tilde{r}}\left[e^{-2\tilde{u}+\hat{v}}\tilde{r}^{n-1}(\partial_{\tilde{r}}\hat{v} + \hat{\tilde{W}}^r)\right] = & n\tilde{r}_0^n\partial_{\tilde{r}}(e^{\hat{v}}) + 2e^{2\tilde{u}}\tilde{r}^{n-1}\partial_{\xi}(e^{-\hat{v}}\tilde{W}^{\xi}) \\ & - e^{\hat{v}}\tilde{r}^{n-1}R(\gamma) + A \end{aligned} \quad (3.14)$$

where

$$\begin{aligned} A = & e^{\hat{v}}\tilde{r}^{n-1}[R(g) + n(n-1)] \\ & + 2e^{-2\tilde{u}+\hat{v}}\tilde{r}^{n-1}\left[\frac{n-1}{2(n-2)}(\hat{\tilde{W}}^r)^2 + \frac{1}{8}|\tilde{\tilde{W}}^r|_{\gamma}^2\right] \\ & + 2e^{2\tilde{u}-\hat{v}}\tilde{r}^{n-1}\left[\frac{n-1}{2(n-2)}(\tilde{W}^{\xi})^2 + \frac{1}{8}|\tilde{W}^{\xi}|_{\gamma}^2\right] \geq 0. \end{aligned}$$

Integrating (3.14) on $[\tilde{r}_0, \infty)$, noting that $e^{-2\tilde{u}}$ vanishes at $\tilde{r} = \tilde{r}_0$, and using (3.13), we obtain

$$\begin{aligned} \lim_{\tilde{r} \rightarrow \infty} -2e^{-2\tilde{u}+\hat{v}}\tilde{r}^{n-1}(\partial_{\tilde{r}}\hat{v} + \hat{\tilde{W}}^r) = & n\tilde{r}_0^n\left(1 - \frac{\tilde{r}_0}{\tilde{r}_0}\right) \\ & + \int_{\tilde{r}_0}^{\infty} \left[2e^{2\tilde{u}}\tilde{r}^{n-1}\partial_{\xi}(e^{-\hat{v}}\tilde{W}^{\xi}) - e^{\hat{v}}\tilde{r}^{n-1}R(\gamma) + A\right] d\tilde{r}. \end{aligned} \quad (3.15)$$

Since

$$\begin{aligned}\hat{W}^r &= \frac{1}{2}\text{tr}_\gamma(\partial_{\tilde{r}}\gamma) - \frac{n-2}{\tilde{r}} \\ &= -\frac{n-1}{\tilde{r}^n}\text{tr}_{h_0}\tilde{w}_{n-1} - \frac{n}{\tilde{r}^{n+1}}\text{tr}_{h_0}\tilde{w}_n + O(\tilde{r}^{-2-n}).\end{aligned}$$

Combining this with (3.7) and (3.11), we obtain

$$\begin{aligned}\partial_{\tilde{r}}\hat{v} + \hat{W}^r &= -\frac{n-1}{\tilde{r}^n}(\tilde{v}_{n-1} + \text{tr}_{h_0}\tilde{w}_{n-1}) - \frac{n}{\tilde{r}^{n+1}}(\tilde{v}_n + \text{tr}_{h_0}\tilde{w}_n) + O(\tilde{r}^{-2-n}) \\ &= -\frac{n}{\tilde{r}^{n+1}}(\tilde{v}_n + \text{tr}_{h_0}\tilde{w}_n) + O(\tilde{r}^{-2-n}),\end{aligned}$$

which implies

$$\lim_{\tilde{r} \rightarrow \infty} -2e^{-2\tilde{u} + \hat{v}}\tilde{r}^{n-1}(\partial_{\tilde{r}}\hat{v} + \hat{W}^r) = 2n(\tilde{v}_n + \text{tr}_{h_0}\tilde{w}_n).$$

Put this into (3.15), and then integrate over $(\partial_\infty M, h_0)$. Noting that the term containing $\partial_\xi(e^{-\hat{v}}\tilde{W}^\xi)$ cancels away after integrating over ξ , and recalling the assumption (1.13), we obtain

$$\begin{aligned}2n \int_{\partial_\infty M} (\tilde{v}_n + \text{tr}_{h_0}\tilde{w}_n) dV_{h_0} &\geq n\tilde{r}_0^n \left(1 - \frac{\check{r}_0}{\tilde{r}_0}\right) \text{Vol}(\partial_\infty M, h_0) \\ &\quad + \int_{\partial_\infty M} \int_{\tilde{r}_0}^\infty A d\tilde{r} dV_{h_0}.\end{aligned}$$

Inserting this into (3.12), we obtain

$$\begin{aligned}E(g) - E(g_{\text{HM}}) &\geq \tilde{r}_0^n \left[n-1 + \left(\frac{\check{r}_0}{\tilde{r}_0}\right)^n - n\frac{\check{r}_0}{\tilde{r}_0} \right] \text{Vol}(\partial_\infty M, h_0) \\ &\quad + \int_{\partial_\infty M} \int_{\tilde{r}_0}^\infty A d\tilde{r} dV_{h_0}.\end{aligned}$$

It is easy to see that

$$n-1 + s^n - ns = (1-s)(n-1-s-\dots-s^{n-1}) \geq 0$$

for $s > 0$, and the equality occurs if and only if $s = 1$. Taking $s = \frac{\check{r}_0}{\tilde{r}_0}$, we obtain

$$E(g) - E(g_{\text{HM}}) \geq 0.$$

For the case of equality, we have

$$\tilde{r}_0 = \check{r}_0, \quad A \equiv 0,$$

which implies

$$R + n(n-1) = 0, \quad \hat{W}^r = \tilde{W}^\xi = 0, \quad \tilde{W}^r = 0, \quad \tilde{W}^\xi = 0.$$

It follows that

$$\tilde{W}^r = \frac{n-2}{\tilde{r}} + \hat{W}^r = \frac{n-2}{\tilde{r}}.$$

Hence

$$\partial_r(\tilde{\gamma}_{ij}) = \frac{2\tilde{W}^r}{n-2}\tilde{\gamma}_{ij} + \tilde{W}^r = \frac{2}{\tilde{r}}\tilde{\gamma}_{ij}.$$

Therefore

$$\partial_r\left(\frac{1}{\tilde{r}^2}\tilde{\gamma}_{ij}\right) = 0.$$

It implies that

$$\frac{1}{\tilde{r}^2}\tilde{\gamma}_{ij} = \frac{1}{\tilde{r}^2}\tilde{\gamma}_{ij}\Big|_{\tilde{r}=\infty} = \delta_{ij}, \quad \gamma = \tilde{r}^2 \sum_{i=3}^n (d\phi^i)^2.$$

Now we can simplify (3.14) and find that

$$\partial_{\tilde{r}}^2\left(e^{\hat{v}}\right) + \frac{(n+1)\tilde{r}^n + (\frac{n}{2}-1)\tilde{r}_0^n}{\tilde{r}(\tilde{r}^n - \tilde{r}_0^n)} \cdot \left(\partial_{\tilde{r}}e^{\hat{v}}\right)^2 = 0.$$

Since

$$e^{\hat{v}(\tilde{r}_0, \xi)} = \frac{\tilde{r}_0}{\tilde{r}_0} = 1 = \lim_{\tilde{r} \rightarrow \infty} e^{\hat{v}(\tilde{r})},$$

we can show $\hat{v} \equiv 0$ by the maximum principle. Finally, we obtain

$$g = \frac{1}{\tilde{r}^2\left(1 - \frac{\tilde{r}_0^n}{\tilde{r}^n}\right)} d\tilde{r}^2 + \tilde{r}^2\left(1 - \frac{1}{\tilde{r}^n}\right) d\xi^2 + \tilde{r}^2 \sum_{i=3}^n (d\phi^i)^2.$$

Thus g is the Horowitz–Myers metric g_{HM} .

Q.E.D.

Acknowledgement. The work is supported by Chinese National Natural Science of Foundation (Grant No. 11731001), the special foundation for Junwu and Guangxi Ba Gui Scholars, the Guangdong Basic and Applied Basic Research Foundation 2021A1515010379. The authors thank the referees for carefully reading the manuscript and for their helpful suggestions.

REFERENCES

- [1] H. Barzegar, P. Chruściel, M. Hörzinger, M. Maliborski, L. Nguyen. Remarks on the energy of asymptotically Horowitz-Myers metrics. Phys. Rev. D **101** (2020) 024007.
- [2] P. Chruściel, M. Herzlich. The mass of asymptotically hyperbolic Riemannian manifolds. Pac. J. Math. **212** (2003) 231-264.
- [3] G. Galloway, S. Surya, and E. Woolgar. On the Geometry and Mass of Static, Asymptotically AdS Spacetimes, and the Uniqueness of the AdS Soliton. Commun. Math. Phys. **241** (2003) 1-25.
- [4] G. Horowitz, R. Myers. AdS-CFT correspondence and a new positive energy conjecture for general relativity. Phys. Rev. D **59** (1998) 026005.
- [5] Z. Liang, X. Zhang. Metrics of Horowitz-Myers type with the negative constant scalar curvature. J. Math. Phys. **61** (2020) 112503.
- [6] X. Wang. The mass of asymptotically hyperbolic manifolds. J. Diff. Geom. **57** (2001) 273-299.

- [7] E. Woolgar. The rigid Horowitz-Myers conjecture. J. High Energ. Phys. **03** (2017) 104.

[†]DEPARTMENT OF MATHEMATICS, JINAN UNIVERSITY, GUANGZHOU, 510632, P. R. CHINA

Email address: `tzbliang@jnu.edu.cn`

^bGUANGXI CENTER FOR MATHEMATICAL RESEARCH, GUANGXI UNIVERSITY, NANNING, GUANGXI 530004, PR CHINA

^bINSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, PR CHINA AND SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, PR CHINA

Email address: `xzhang@gxu.edu.cn`, `xzhang@amss.ac.cn`