

Asymptotic behavior of solutions toward the rarefaction waves to the Cauchy problem for the generalized Benjamin-Bona-Mahony-Burgers equation with dissipative term

Natsumi Yoshida¹

^a*Graduate Faculty of Interdisciplinary Research Faculty of Education, University of Yamanashi, Kofu, Yamanashi 400-8510, Japan.*

Abstract

In this paper, we investigate the asymptotic behavior of solutions to the Cauchy problem with the far field condition for the generalized Benjamin-Bona-Mahony-Burgers equation with a fourth-order dissipative term. When the corresponding Riemann problem for the hyperbolic part admits a Riemann solution which consists of single rarefaction wave, it is proved that the solution of the Cauchy problem tends toward the rarefaction wave as time goes to infinity. We can further obtain the same global asymptotic stability of the rarefaction wave to the generalized Korteweg-de Vries-Benjamin-Bona-Mahony-Burgers equation with a fourth-order dissipative term as the former one.

Keywords: Benjamin-Bona-Mahony-Burgers equation, Korteweg-de Vries-Benjamin-Bona-Mahony-Burgers equation, convex flux, asymptotic behavior, rarefaction wave

AMS subject classifications: 35K55, 35B40, 35L65

1. Introduction and main theorems

In this paper, we consider the asymptotic behavior of solutions to the Cauchy problem for the generalized Benjamin-Bona-Mahony-Burgers equation with a fourth-order dissipative term

$$\begin{cases} \partial_t u + \partial_x(f(u)) - \alpha \partial_t \partial_x^2 u - \beta \partial_x^2 u + \gamma \partial_x^4 u = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty). \end{cases} \quad (1.1)$$

Email address: 14v00067@gmail.com (Natsumi Yoshida)

Here, $u = u(t, x)$ is the unknown function of $t > 0$ and $x \in \mathbb{R}$, and α, β, γ are positive constants u_0 is the initial data, and $u_{\pm} \in \mathbb{R}$ are the prescribed far field states. We suppose that f is a smooth function.

There are many results concerning with the mathematical structure, such as the global existence and time-decay properties of solutions, of the generalized Benjamin-Bona-Mahony-Burgers equation with the dissipative terms (see Kondo-Webster [21], [22], [23], [24], Wang [49], Xu-Li [51] Zhao-Xuan [64] and so on). The model (1.1) is closely related to the following Benjamin-Bona-Mahony-Burgers equation

$$\partial_t u + \delta \partial_x u + u \partial_x u - \alpha \partial_t \partial_x^2 u - \beta \partial_x^2 u = 0, \quad (1.2)$$

where $\delta \in \mathbb{R}$. The mathematical structure of (1.2) have also been investigated by Amick-Bona-Schonbek [1], Mei [37], [38], Mei-Schmeiser [39], Naumkin [40] and so on. When $\beta = 0$, then (1.2) becomes the following Benjamin-Bona-Mahony equation

$$\partial_t u + \delta \partial_x u + u \partial_x u - \alpha \partial_t \partial_x^2 u = 0, \quad (1.3)$$

which was advocated by [3] as a refinement of the following Korteweg de-Vries equation

$$\partial_t u + \delta \partial_x u + u \partial_x u - \alpha \partial_x^3 u = 0. \quad (1.4)$$

For the case $\alpha = 1$ and $\beta = 0$ of (1.2), (1.2) is the so-called regularized long wave equation, which was proposed by Peregrine [44] and [3], as follows.

$$\partial_t u + \delta \partial_x u + u \partial_x u - \partial_t \partial_x^2 u = 0. \quad (1.5)$$

We note that (1.3) and (1.5) are known as the approximated models for the long waves of small amplitude.

We are going to obtain the rarefaction stability of the solution to (1.1). Therefore we deal with the case where the flux function f is fully convex, that is,

$$f''(u) > 0 \quad (u \in \mathbb{R}), \quad (1.6)$$

and $u_- < u_+$. Then, since the corresponding Riemann problem (cf. [27])

$$\begin{cases} \partial_t u + \partial_x (f(u)) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0^R(x) := \begin{cases} u_- & (x < 0), \\ u_+ & (x > 0) \end{cases} \end{cases} \quad (1.7)$$

turns out to admit a single rarefaction wave solution, we expect that the solution of the Cauchy problem (1.1) tends toward the rarefaction wave as time goes to infinity. Here, the rarefaction wave connecting u_- to u_+ is given by

$$u^r \left(\frac{x}{t}; u_-, u_+ \right) = \begin{cases} u_- & (x \leq f'(u_-) t), \\ (f')^{-1} \left(\frac{x}{t} \right) & (f'(u_-) t \leq x \leq f'(u_+) t), \\ u_+ & (x \geq f'(u_+) t). \end{cases} \quad (1.8)$$

In particular, we also expect that if $u_- = u_+ =: \tilde{u}$, then the solution of the Cauchy problem (1.1) tends toward the constant state \tilde{u} as time goes to infinity.

There are many results concerning with the rarefaction stabilities. For viscous conservation law,

$$\begin{cases} \partial_t u + \partial_x (f(u) - \mu \partial_x u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases}$$

with the condition (1.6), Il'in-Oleĭnik [18] showed that the solution tends toward the single rarefaction wave under the condition $u_- < u_+$ (and the one does the single viscous shock wave under the condition $u_- > u_+$, for further studies, see [34], [35], [53], [61] and so on). Hattori-Nishihara [16] also obtained the pointwise and time-decay estimates of the difference $|u - u^*|$. Harabetian [14] further considered the following rarefaction problem for a quasilinear parabolic equation

$$\begin{cases} \partial_t u + \partial_x (f(u) - A'(u) \partial_x u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases}$$

where $A'(u) \geq 0$ ($u \in \mathbb{R}$), and obtained the precise time-decay estimates of global stability of the rarefaction wave with the aid of the arguments on monotone semigroups by Osher-Ralston [42]. For the following Cauchy problem of the Matsumura-Nishihara model

$$\begin{cases} \partial_t u + \partial_x (f(u) - \mu |\partial_x u|^{p-1} \partial_x u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases}$$

where $p > 1$ and the viscosity $\mu |\partial_x u|^{p-1} \partial_x u$ is the so-called Ostwald-de Waele-type viscosity advocated by de Waele [9] and Ostwald [43] (which is a typical example for the non-Newtonian viscosity, see also [6], [7], [8], [19], [25], [30], [31], [48] and so on), Matsumura-Nishihara [33] first investigated and proved the global stability of the rarefaction wave by using the technical energy method. Yoshida [54] further obtained its precise time-decay estimates by using the time-weighted energy method (for the stabilities of the multiwave pattern, see [55], [56], [57]). Furthermore, Matsumura-Yoshida [36] considered the following Cauchy problem of the non-Newtonian viscous conservation law

$$\begin{cases} \partial_t u + \partial_x (f(u) - \sigma(\partial_x u)) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases}$$

where viscosity function σ satisfies the conditions

$$\begin{cases} \sigma(0) = 0, \quad \sigma'(v) > 0 \quad (v \in \mathbb{R}), \\ |\sigma(v)| \sim |v|^p, \quad |\sigma'(v)| \sim |v|^{p-1} \quad (|v| \rightarrow \infty), \end{cases}$$

and obtained the global stability of the rarefaction wave for the case $0 < p < 3/7$. Recently, Yoshida [60] obtained this rarefaction stability for more general case

$0 < p < 1/3$ and its precise time-decay estimates. For the rarefaction problem of the Korteweg-de Vries equation

$$\begin{cases} \partial_t u + \partial_x(-3u^2 + \partial_x^2 u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \begin{cases} \rightarrow 0 & (x \rightarrow \infty), \\ \rightarrow u_- > 0 & (x \rightarrow -\infty), \end{cases} \end{cases}$$

Egorova-Grunert-Teschl [12] and Egorova-Teschl [13] obtained the existence and uniqueness in some class of the classical solution, and Andreiev-Egorova-Lange-Teschl [2] also obtained the valid asymptotic formula of the solution. For diffusive dispersive conservation laws, Wang-Zhu [50] obtained the local stability of the rarefaction wave for the following Cauchy problem of the generalized Korteweg-de Vries-Burgers equation

$$\begin{cases} \partial_t u + \partial_x(f(u) - \mu \partial_x u + \delta \partial_x^2 u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty). \end{cases}$$

Duan-Zhao [11] and Yoshida [59] further obtained the global stabilities of the rarefaction wave under some growth conditions for f (for stability and time-decay properties of a travelling wave, see [4], [5], [41]). For the Cauchy problem of the generalized Korteweg-de Vries-Burgers-Kuramoto equation

$$\begin{cases} \partial_t u + \partial_x(f(u) - \mu \partial_x u + \delta \partial_x^2 u + \nu \partial_x^3 u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases}$$

Ruan-Gao-Chen [47] first obtained the local stability of the rarefaction wave. Duan-Fan-Kim-Xie [10] and Yoshida [59] also obtained the global stability of the rarefaction wave under some growth conditions for f . Recently, Yoshida [63] (see also [62]) investigated the following Cauchy problem for a diffusive dispersive conservation law, that is, the generalized Korteweg-de Vries-Burgers-Kuramoto equation without the viscosity term as

$$\begin{cases} \partial_t u + \partial_x(f(u) + \delta \partial_x^2 u + \nu \partial_x^3 u) = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases}$$

and obtained the global stability of the rarefaction wave under some growth conditions for f .

Our main results of the present paper are as follows.

Theorem 1.1 (Main Theorem I). *Assume the far field states u_{\pm} satisfy $u_- = u_+ = \tilde{u}$, and the convective flux $f \in C^2(\mathbb{R})$. Further assume the initial data satisfy $u_0 - \tilde{u} \in L^2$ and $\partial_x u_0 \in H^2$. Then the Cauchy problem (1.1) has a unique global in time solution u satisfying*

$$\begin{cases} u - \tilde{u} \in C^0([0, \infty); H^3), \\ \partial_x u \in L^2(0, \infty; H^3), \\ \partial_t u \in L^2(0, \infty; H^2), \end{cases}$$

and the asymptotic behavior

$$\lim_{t \rightarrow \infty} \left(\sup_{x \in \mathbb{R}} |u(t, x) - \tilde{u}| + \sup_{x \in \mathbb{R}} |\partial_x u(t, x)| + \sup_{x \in \mathbb{R}} |\partial_x^2 u(t, x)| \right) = 0.$$

Theorem 1.2 (Main Theorem II). *Assume the far field states u_{\pm} satisfy $u_- < u_+$, and the convective flux $f \in C^5(\mathbb{R})$ satisfy (1.6). Further assume the initial data satisfy $u_0 - u_0^R \in L^2$ and $\partial_x u_0 \in H^2$. Then the Cauchy problem (1.1) has a unique global in time solution u satisfying*

$$\begin{cases} u - u_0^R \in C^0([0, \infty); H^3), \\ \partial_x u \in L_{\text{loc}}^2(0, \infty; H^3), \\ \partial_t u \in L_{\text{loc}}^2(0, \infty; H^2), \end{cases}$$

and the asymptotic behavior

$$\begin{cases} \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| u(t, x) - u^r \left(\frac{x}{t}; u_-, u_+ \right) \right| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} |\partial_x u(t, x) - \partial_x u^r(t, x; u_-, u_+)| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} |\partial_x^2 u(t, x) - \partial_x^2 u^r(t, x; u_-, u_+)| = 0, \end{cases}$$

where $\partial_x u^r$ and $\partial_x^2 u^r$ are given by

$$\partial_x u^r(t, x; u_-, u_+) = \begin{cases} 0 & (x \leq f'(u_-)t), \\ \frac{1}{f''\left((f')^{-1}\left(\frac{x}{t}\right)\right)} \frac{1}{t} & (f'(u_-)t \leq x \leq f'(u_+)t), \\ 0 & (x \geq f'(u_+)t), \end{cases}$$

$$\begin{aligned} & \partial_x^2 u^r(t, x; u_-, u_+) \\ &= \begin{cases} 0 & (x \leq f'(u_-)t), \\ \frac{1}{\left(f''\left((f')^{-1}\left(\frac{x}{t}\right)\right)\right)^3} \frac{x}{t^3} - \frac{1}{f''\left((f')^{-1}\left(\frac{x}{t}\right)\right)} \frac{1}{t^2} & (f'(u_-)t \leq x \leq f'(u_+)t), \\ 0 & (x \geq f'(u_+)t). \end{cases} \end{aligned}$$

Furthermore, for the Cauchy problem for the generalized Benjamin-Bona-Mahony-Burgers equation with third-order dispersive and fourth-order dissipative terms, the so-called generalized Korteweg-de Vries-Benjamin-Bona-Mahony-Burgers equation with a fourth-order dissipative term (see [46])

$$\begin{cases} \partial_t u + \partial_x(f(u)) - \alpha \partial_t \partial_x^2 u - \beta \partial_x^2 u + \delta \partial_x^3 u + \gamma \partial_x^4 u = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases} \quad (1.9)$$

where $\delta \in \mathbb{R}$, we can obtain the same stabilities as Theorems 1.1 and 1.2 in the next theorems.

Theorem 1.3 (Main Theorem III). *Assume the far field states u_{\pm} satisfy $u_- = u_+ = \tilde{u}$, and the convective flux $f \in C^2(\mathbb{R})$. Further assume the initial data satisfy $u_0 - \tilde{u} \in L^2$ and $\partial_x u_0 \in H^2$. Then for the Cauchy problem (1.9), the same result as in Theorem 1.1 holds true.*

Theorem 1.4 (Main Theorem IV). *Assume the far field states u_{\pm} satisfy $u_- < u_+$, and the convective flux $f \in C^5(\mathbb{R})$ satisfy (1.6). Further assume the initial data satisfy $u_0 - u_0^R \in L^2$ and $\partial_x u_0 \in H^2$. Then for the Cauchy problem (1.9), the same result as in Theorem 1.2 holds true.*

We finally remark that when $\gamma = \delta = 0$, the problem (1.9) becomes the following Cauchy problem for usual generalized Benjamin-Bona-Mahony-Burgers equation.

$$\begin{cases} \partial_t u + \partial_x(f(u)) - \alpha \partial_t \partial_x^2 u - \beta \partial_x^2 u = 0 & (t > 0, x \in \mathbb{R}), \\ u(0, x) = u_0(x) \rightarrow u_{\pm} & (x \rightarrow \pm\infty), \end{cases} \quad (1.10)$$

We can also obtain the similar stabilities as Theorems 1.1 and 1.2 in the next theorems.

Theorem 1.5. *Assume the far field states u_{\pm} satisfy $u_- = u_+ = \tilde{u}$, and the convective flux $f \in C^2(\mathbb{R})$. Further assume the initial data satisfy $u_0 - \tilde{u} \in L^2$ and $\partial_x u_0 \in H^2$. Then the Cauchy problem (1.10) has a unique global in time solution u satisfying*

$$\begin{cases} u - \tilde{u} \in C^0([0, \infty); H^3), \\ \partial_x u \in L^2(0, \infty; H^2), \\ \partial_t u \in L^2(0, \infty; H^2), \end{cases}$$

and the asymptotic behavior

$$\lim_{t \rightarrow \infty} \left(\sup_{x \in \mathbb{R}} |u(t, x) - \tilde{u}| + \sup_{x \in \mathbb{R}} |\partial_x u(t, x)| + \sup_{x \in \mathbb{R}} |\partial_x^2 u(t, x)| \right) = 0.$$

Theorem 1.6. *Assume the far field states u_{\pm} satisfy $u_- < u_+$, and the convective flux $f \in C^5(\mathbb{R})$ satisfy (1.6). Further assume the initial data satisfy $u_0 - u_0^R \in L^2$ and $\partial_x u_0 \in H^2$. Then the Cauchy problem (1.10) has a unique global in time solution u satisfying*

$$\begin{cases} u - u_0^R \in C^0([0, \infty); H^3), \\ \partial_x u \in L_{\text{loc}}^2(0, \infty; H^2), \\ \partial_t u \in L_{\text{loc}}^2(0, \infty; H^2), \end{cases}$$

and the asymptotic behavior

$$\begin{cases} \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| u(t, x) - u^r \left(\frac{x}{t}; u_-, u_+ \right) \right| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \partial_x u(t, x) - \partial_x u^r(t, x; u_-, u_+) \right| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \partial_x^2 u(t, x) - \partial_x^2 u^r(t, x; u_-, u_+) \right| = 0, \end{cases}$$

where $\partial_x u^r$ and $\partial_x^2 u^r$ are given in Theorem 1.2.

Because the proofs of Theorems 1.1 and 1.3-1.6 are similarly given as or easier than that for Theorem 1.2, we only show Theorem 1.2 in the following sections.

This paper is organized as follows. In Section 2, we construct the approximation of the rarefaction wave and prepare the basic properties of the rarefaction wave and the approximated one. We reformulate the problem in terms of the deviation from the asymptotic state in Section 3. In order to show the asymptotics, we establish the *a priori* estimates by using the technical energy method in Section 4. Finally in Section 5, we give several uniform estimates by using the *a priori* estimates in Sections 3 and 4.

Some Notation. We denote by C generic positive constants unless they need to be distinguished. In particular, use $C_{\alpha, \beta, \dots}$ when we emphasize the dependency on α, β, \dots .

For function spaces, $L^p = L^p(\mathbb{R})$ and $H^k = H^k(\mathbb{R})$ denote the usual Lebesgue space and k -th order Sobolev space on the whole space \mathbb{R} with norms $\|\cdot\|_{L^p}$ and $\|\cdot\|_{H^k}$, respectively.

2. Preliminaries

In this section, we prepare the several lemmas concerning with the basic properties of the rarefaction wave for the proof of the main Theorem 1.2. Since the rarefaction wave u^r is not smooth enough, we construct a smooth approximated one. To do that, we first consider the rarefaction wave solution w^r to the Riemann problem for the non-viscous Burgers equation

$$\begin{cases} \partial_t w + \partial_x \left(\frac{1}{2} w^2 \right) = 0 & (t > 0, x \in \mathbb{R}), \\ w(0, x) = w_0^R(x; w_-, w_+) := \begin{cases} w_+ & (x > 0), \\ w_- & (x < 0), \end{cases} \end{cases} \quad (2.1)$$

where $w_{\pm} \in \mathbb{R}$ ($w_- < w_+$) are the prescribed far field states. The unique global weak solution $w = w^r(x/t; w_-, w_+)$ of (2.1) is explicitly given by

$$w^r \left(\frac{x}{t}; w_-, w_+ \right) = \begin{cases} w_- & (x \leq w_- t), \\ \frac{x}{t} & (w_- t \leq x \leq w_+ t), \\ w_+ & (x \geq w_+ t). \end{cases} \quad (2.2)$$

Next, under the condition $f''(u) > 0$ ($u \in \mathbb{R}$) and $u_- < u_+$, the rarefaction wave solution $u = u^r\left(\frac{x}{t}; u_-, u_+\right)$ of the Riemann problem (1.2) for hyperbolic conservation law is exactly given by

$$u^r\left(\frac{x}{t}; u_-, u_+\right) = (\lambda)^{-1}\left(w^r\left(\frac{x}{t}; \lambda_-, \lambda_+\right)\right) \quad (2.3)$$

which is nothing but (1.6), where $\lambda_{\pm} := \lambda(u_{\pm}) = f'(u_{\pm})$. We define a smooth approximation of $w^r\left(\frac{x}{t}; w_-, w_+\right)$ by the unique classical solution

$$w = w(t, x; w_-, w_+)$$

to the Cauchy problem for the following non-viscous Burgers equation as

$$\begin{cases} \partial_t w + \partial_x \left(\frac{1}{2} w^2 \right) = 0 & (t > 0, x \in \mathbb{R}), \\ w(0, x) = w_0(x) := \frac{w_- + w_+}{2} + \frac{w_+ - w_-}{2} K_q \int_0^{\epsilon x} \frac{dy}{(1+y^2)^q} & (x \in \mathbb{R}), \end{cases} \quad (2.4)$$

where K_q is a positive constant such that

$$K_q \int_0^{\infty} \frac{dy}{(1+y^2)^q} = 1 \quad \left(q > \frac{1}{2} \right).$$

By applying the method of characteristics, we get the following formula

$$\begin{cases} w(t, x) = w_0(x_0(t, x)) = \frac{\lambda_- + \lambda_+}{2} + \frac{\lambda_+ - \lambda_-}{2} K_q \int_0^{\epsilon x_0(t, x)} \frac{dy}{(1+y^2)^q}, \\ x = x_0(t, x) + w_0(x_0(t, x)) t. \end{cases} \quad (2.5)$$

By making use of (2.5) similarly as in [32], we can obtain the properties of the smooth approximation $w(t, x; w_-, w_+)$ in the next lemma.

Lemma 2.1. *Assume that the far field states satisfy $w_- < w_+$. Then the classical solution $w(t, x) = w(t, x; w_-, w_+)$ given by (2.4) satisfies the following properties:*

- (1) $w_- < w(t, x) < w_+$ and $\partial_x w(t, x) > 0$ ($t > 0, x \in \mathbb{R}$).
- (2) For any $r \in [1, \infty]$, there exists a positive constant $C_{q,r}$ such that

$$\begin{aligned} & \|\partial_x w(t)\|_{L^r}^r \leq C_{q,r} \min \left\{ \epsilon^{r-1} \tilde{w}^r, \tilde{w} (1+t)^{-r+1} \right\} \quad (t \geq 0), \\ & \|\partial_t w(t)\|_{L^r}^r \leq C_{q,r} \tilde{w}^r \min \left\{ \epsilon^{r-1} \tilde{w}^r, \tilde{w} (1+t)^{-r+1} \right\} \quad (t \geq 0), \\ & \|\partial_x^2 w(t)\|_{L^r}^r \\ & \leq C_{q,r} \min \left\{ \epsilon^{2r-1} \tilde{w}^r, \epsilon^{(r-1)(1-\frac{1}{2q})} \tilde{w}^{-\frac{r-1}{2q}} (1+t)^{-r-\frac{r-1}{2q}} \right\} \quad (t \geq 0), \\ & \|\partial_x^3 w(t)\|_{L^r}^r \leq C_{q,r} \min \left\{ \epsilon^{3r-1} \tilde{w}^r, a(1+t, \epsilon, \tilde{w}) \right\} \quad (t \geq 0), \\ & \|\partial_x^4 w(t)\|_{L^r}^r \leq C_{q,r} \min \left\{ \epsilon^{4r-1} \tilde{w}^r, b(1+t, \epsilon, \tilde{w}) \right\} \quad (t \geq 0), \end{aligned}$$

where

$$\tilde{w} := \frac{w_+ - w_-}{2} > 0, \quad \tilde{w} := \max\{|w_-|, |w_+|\},$$

$$\begin{aligned} a(t, \epsilon, \tilde{w}) &:= \epsilon^{3r} \tilde{w}^r (1 + \epsilon \tilde{w} t)^{1-4r} + \epsilon^{2(r-1)(1-\frac{1}{2q})} \tilde{w}^{-\frac{r-1}{q}} t^{-1-(r-1)(1+\frac{1}{q})} \\ &\quad + \epsilon^{(2r-1)(1-\frac{1}{2q})} \tilde{w}^{-\frac{2r-1}{2q}} t^{-r-\frac{2r-1}{2q}}, \end{aligned}$$

and

$$\begin{aligned} b(t, \epsilon, \tilde{w}) &:= \epsilon^{3r} \tilde{w}^r (1 + \epsilon \tilde{w} t)^{1-5r} + \epsilon^{(3r-2)(1-\frac{1}{2q})} \tilde{w}^{-\frac{3r-2}{2q}} t^{-r(1+\frac{3}{2q})+\frac{1}{q}} \\ &\quad + \epsilon^{(3r-1)(1-\frac{1}{2q})} \tilde{w}^{-\frac{3r-1}{2q}} t^{-r-\frac{3r-1}{2q}}. \end{aligned}$$

(3) It follows that

$$\begin{cases} \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| w(t, x) - w^r \left(\frac{x}{t} \right) \right| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} |\partial_x w(t, x) - \partial_x w^r(t, x)| = 0, \end{cases}$$

where $\partial_x w^r$ is given by

$$\partial_x w^r(t, x; w_-, w_+) = \begin{cases} 0 & (x \leq w_- t), \\ \frac{1}{t} & (w_- t \leq x \leq w_+ t), \\ 0 & (x \geq w_+ t). \end{cases}$$

We now define the approximation for the rarefaction wave $u^r(x/t; u_-, u_+)$ by

$$U^r(t, x; u_-, u_+) = (\lambda)^{-1} (w(t, x; \lambda_-, \lambda_+)). \quad (2.6)$$

Using Lemma 2.1, we also have the next lemma.

Lemma 2.2. *Let $q = 1$. Assume that the far field states satisfy $u_- < u_+$, and the flux function $f \in C^5(\mathbb{R})$, $f''(u) > 0$ ($u \in [u_-, u_+]$). Then we have the following properties.*

(1) $U^r(t, x)$ defined by (2.6) is the unique C^4 -global solution in space-time of the Cauchy problem

$$\begin{cases} \partial_t U^r + \partial_x (f(U^r)) = 0 & (t > 0, x \in \mathbb{R}), \\ U^r(0, x) = (\lambda)^{-1} \left(\frac{\lambda_- + \lambda_+}{2} + \frac{\lambda_+ - \lambda_-}{2} K_q \int_0^{\epsilon x} \frac{dy}{(1+y^2)^q} \right) & (x \in \mathbb{R}), \\ \lim_{x \rightarrow \pm \infty} U^r(t, x) = u_{\pm} & (t \geq 0). \end{cases}$$

(2) $u_- < U^r(t, x) < u_+$ and $\partial_x U^r(t, x) > 0$ ($t > 0, x \in \mathbb{R}$).

(3) For any $r \in [1, \infty]$, there exists a positive constant $C_{\lambda_{\pm}, r}$ such that

$$\begin{aligned} \|\partial_x U^r(t)\|_{L^r}^r &\leq C_{\lambda_{\pm}, r} \min\{\epsilon^{r-1}, (1+t)^{-r+1}\} \quad (t \geq 0), \\ \|\partial_t U^r(t)\|_{L^r}^r &\leq C_{\lambda_{\pm}, r} \min\{\epsilon^{r-1}, (1+t)^{-r+1}\} \quad (t \geq 0), \\ \|\partial_x^2 U^r(t)\|_{L^r}^r &\leq C_{\lambda_{\pm}, r} \min\left\{\epsilon^{2r-1}, \epsilon^{\frac{r-1}{2}} (1+t)^{-\frac{3r-1}{2}}\right\} \quad (t \geq 0), \\ \|\partial_x^3 U^r(t)\|_{L^r}^r &\leq C_{\lambda_{\pm}, r} \min\{\epsilon^{3r-1}, \epsilon^{r-1} (1+t)^{-2r+1}\} \quad (t \geq 0), \\ \|\partial_x^4 U^r(t)\|_{L^r}^r &\leq C_{\lambda_{\pm}, r} \min\left\{\epsilon^{4r-1}, \epsilon^{\frac{3r-2}{2}} (1+t)^{-\min\{\frac{3}{2}r+1, \frac{5}{2}r-1\}}\right\} \quad (t \geq 0). \end{aligned}$$

(4) It follows that

$$\begin{cases} \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| U^r(t, x) - u^r\left(\frac{x}{t}\right) \right| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} |\partial_x U^r(t, x) - \partial_x u^r(t, x)| = 0, \\ \lim_{t \rightarrow \infty} \sup_{x \in \mathbb{R}} |\partial_x^2 U^r(t, x) - \partial_x^2 u^r(t, x)| = 0. \end{cases}$$

Because the proofs of Lemmas 2.1 and 2.2 are given in [15], [16], [29], [32], [35], [50], [53], [60], [63] and so on, we omit the proofs here.

3. Reformulation of the problem

In this section, we reformulate our problem (1.1) in terms of the deviation from the asymptotic state. Putting ϕ as

$$u(t, x) = U^r(t, x) + \phi(t, x), \quad (3.1)$$

we reformulate the problem (1.1) in terms of the deviation ϕ from U^r as

$$\begin{cases} \partial_t \phi + \partial_x (f(\phi + U^r) - f(U^r)) \\ \quad - \alpha \partial_t \partial_x^2 \phi - \beta \partial_x^2 \phi + \gamma \partial_x^4 \phi = F(U^r) \quad (t > 0, x \in \mathbb{R}), \\ \phi(0, x) = \phi_0(x) := u_0(x) - U^r(0, x) \rightarrow 0 \quad (x \rightarrow \pm\infty), \end{cases} \quad (3.2)$$

where

$$F(U^r) := \alpha \partial_t \partial_x^2 U^r + \beta \partial_x^2 U^r - \delta \partial_x^3 U^r - \gamma \partial_x^4 U^r.$$

Then we look for the unique global in time solution ϕ which has the asymptotic behavior

$$\sum_{k=1}^3 \sup_{x \in \mathbb{R}} |\partial_x^k \phi(t, x)| \rightarrow 0 \quad (t \rightarrow \infty). \quad (3.3)$$

Here we note that $\phi_0 \in H^3$ by the assumptions on u_0 and Lemma 2.2. Then the corresponding theorems for ϕ to Theorem 1.2 we should prove is as follows.

Theorem 3.1. (Global Existence). *Assume the far field states u_{\pm} satisfy $u_- < u_+$, and the convective flux $f \in C^5(\mathbb{R})$ satisfy (1.6). Further assume the initial data satisfy $\phi_0 \in H^3$. Then the Cauchy problem (3.2) has a unique global in time solution ϕ satisfying*

$$\begin{cases} \phi \in C^0([0, \infty); H^3), \\ \partial_x \phi \in L^2(0, \infty; H^3), \\ \partial_t \phi \in L^2(0, \infty; H^2), \end{cases}$$

and the asymptotic behavior

$$\lim_{t \rightarrow \infty} \sum_{k=0}^2 \sup_{x \in \mathbb{R}} |\partial_x^k \phi(t, x)| = 0.$$

In order to obtain Theorem 3.1, we prepare the local existence precisely, we formulate the problem (3.2) at general initial time $\tau \geq 0$:

$$\begin{cases} \partial_t \phi + \partial_x (f(\phi + U^r) - f(U^r)) \\ \quad - \alpha \partial_t \partial_x^2 \phi - \beta \partial_x^2 \phi + \gamma \partial_x^4 \phi = F(U^r) \quad (t > \tau, x \in \mathbb{R}), \\ \phi(\tau, x) = \phi_{\tau}(x) := u_{\tau}(x) - U^r(\tau, x) \rightarrow 0 \quad (x \rightarrow \pm\infty). \end{cases} \quad (3.4)$$

Theorem 3.2 (Local Existence). *For any $M > 0$, there exists a positive constant $t_0 = t_0(M)$ not depending on τ such that if $\phi_{\tau} \in H^3$ and*

$$\|\phi_{\tau}\|_{H^3} \leq M,$$

then the Cauchy problem (3.4) has a unique solution ϕ on the time interval $[\tau, \tau + t_0(M)]$ satisfying

$$\begin{cases} \phi \in C^0([\tau, \tau + t_0]; H^3), \\ \partial_x \phi \in L^2(\tau, \tau + t_0; H^3), \\ \partial_t \phi \in L^2(\tau, \tau + t_0; H^2), \\ \sup_{t \in [\tau, \tau + t_0]} \|\phi(t)\|_{H^3} \leq 2M. \end{cases}$$

Because the proof of Theorem 3.2 is standard, we omit the details here (cf. [52], [64]). The *a priori* estimates we establish in Section 4 are the following.

Theorem 3.3 (A Priori Estimates). *Under the same assumptions as in Theorem 3.1, for any initial data $\phi_0 \in H^3$, there exists a positive constant C_{ϕ_0} such that if the Cauchy problem (3.1) has a solution ϕ on the time interval $[0, T]$ satisfying*

$$\begin{cases} \phi \in C^0([0, T]; H^3), \\ \partial_x \phi \in L^2(0, T; H^3), \\ \partial_t \phi \in L^2(0, T; H^2), \end{cases}$$

for some positive constant T , then it holds that

$$\begin{aligned} & \|\phi(t)\|_{H^3}^2 + \int_0^t \|(\sqrt{\partial_x U^r} \phi)(\tau)\|_{L^2}^2 d\tau \\ & + \int_0^t \|\partial_x \phi(\tau)\|_{H^3}^2 d\tau + \int_0^t \|\partial_t \phi(\tau)\|_{H^2}^2 d\tau \leq C_{\phi_0} \quad (t \in [0, T]). \end{aligned} \quad (3.5)$$

Combining the local existence Theorem 3.2 together with the *a priori* estimates, Theorem 3.3, we can obtain global existence Theorem 3.1. In fact, we can obtain the unique global in time solutions ϕ to (3.2) in Theorem 3.1 satisfying

$$\begin{cases} \phi \in C^0([0, \infty); H^3), \\ \partial_x \phi \in L^2(0, \infty; H^3), \\ \partial_t \phi \in L^2(0, \infty; H^2), \end{cases}$$

and

$$\begin{aligned} & \sup_{t \geq 0} \|\phi(t)\|_{H^3}^2 + \int_0^\infty \|(\sqrt{\partial_x U^r} \phi)(t)\|_{L^2}^2 dt \\ & + \int_0^\infty \|\partial_x \phi(t)\|_{H^3}^2 dt + \int_0^\infty \|\partial_t \phi(t)\|_{H^2}^2 dt < \infty \end{aligned} \quad (3.6)$$

which yields

$$\int_0^\infty \left| \frac{d}{dt} \|\partial_x \phi(t)\|_{L^2}^2 \right| dt < \infty, \quad \int_0^\infty \left| \frac{d}{dt} \|\partial_x^2 \phi(t)\|_{L^2}^2 \right| dt < \infty. \quad (3.7)$$

In fact, by using (3.6), direct computation shows that

$$\begin{aligned} \int_0^\infty \left| \frac{d}{dt} \|\partial_x \phi(t)\|_{L^2}^2 \right| dt &= 2 \int_0^\infty \left| \int_{-\infty}^\infty \partial_x \phi \partial_t \partial_x \phi dx \right| dt \\ &\leq \int_0^\infty (\|\partial_x \phi(t)\|_{L^2}^2 + \|\partial_t \partial_x \phi(t)\|_{L^2}^2) dt < \infty, \end{aligned} \quad (3.8)$$

$$\begin{aligned} \int_0^\infty \left| \frac{d}{dt} \|\partial_x^2 \phi(t)\|_{L^2}^2 \right| dt &= 2 \int_0^\infty \left| \int_{-\infty}^\infty \partial_x^2 \phi \partial_t \partial_x^2 \phi dx \right| dt \\ &\leq \int_0^\infty (\|\partial_x^2 \phi(t)\|_{L^2}^2 + \|\partial_t \partial_x^2 \phi(t)\|_{L^2}^2) dt < \infty. \end{aligned} \quad (3.9)$$

From (3.8) and (3.9), we get (3.7). We immediately have from (3.7) that

$$\lim_{t \rightarrow \infty} \|\partial_x \phi(t)\|_{L^2} = 0, \quad \lim_{t \rightarrow \infty} \|\partial_x^2 \phi(t)\|_{L^2} = 0. \quad (3.10)$$

Further from (3.10), by using the Sobolev inequality, we obtain the desired asymptotic behavior (3.3) as follows.

$$\begin{aligned}\sup_{x \in \mathbb{R}} |\phi(t, x)| &\leq \sqrt{2} \|\phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{2}} \rightarrow 0 \quad (t \rightarrow \infty), \\ \sup_{x \in \mathbb{R}} |\partial_x \phi(t, x)| &\leq \sqrt{2} \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{2}} \rightarrow 0 \quad (t \rightarrow \infty), \\ \sup_{x \in \mathbb{R}} |\partial_x^2 \phi(t, x)| &\leq \sqrt{2} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{2}} \rightarrow 0 \quad (t \rightarrow \infty).\end{aligned}$$

Thus Theorems 3.1 is proved.

4. *A priori estimates*

In this section, we show the following *a priori* estimate for ϕ in Theorem 3.3. To do that, we prepare the following basic estimate.

Proposition 4.1. *There exists a positive constant C_{ϕ_0} such that*

$$\begin{aligned}\|\phi(t)\|_{H^1}^2 + \int_0^t \int_{-\infty}^{\infty} \int_0^{\phi} (f'(\eta + U^r) - f'(U^r)) \, d\eta \, \partial_x U^r \, dx \, d\tau \\ + \int_0^t \|\partial_x \phi(\tau)\|_{H^1}^2 \, d\tau \leq C_{\phi_0} \quad (t \in [0, T]).\end{aligned}$$

Proof of Proposition 4.1. Multiplying the equation in (3.2) by ϕ and integrating it with respect to x , we have, after integration by parts, that

$$\begin{aligned}\frac{1}{2} \frac{d}{dt} \|\phi(t)\|_{L^2}^2 + \frac{\alpha}{2} \frac{d}{dt} \|\partial_x \phi(t)\|_{L^2}^2 \\ + \int_{-\infty}^{\infty} \int_0^{\phi} (f'(\eta + U^r) - f'(U^r)) \, d\eta \, \partial_x U^r \, dx \\ + \beta \|\partial_x \phi(t)\|_{L^2}^2 + \gamma \|\partial_x^2 \phi(t)\|_{L^2}^2 = \int_{-\infty}^{\infty} \phi F(U^r) \, dx.\end{aligned}\tag{4.1}$$

By making use of the Sobolev inequality and the Young inequality, we estimate the right-hand-side of (4.1) as follows.

$$\begin{aligned}\left| \int_{-\infty}^{\infty} \phi F(U^r) \, dx \right| &\leq \sqrt{2} \|\phi\|_{L^2}^{\frac{1}{2}} \|\partial_x \phi\|_{L^2}^{\frac{1}{2}} \|F(U^r)\|_{L^1} \\ &\leq \frac{\beta}{2} \|\partial_x \phi\|_{L^2}^2 + C_{\beta} \|\phi\|_{L^2}^{\frac{2}{3}} \|F(U^r)\|_{L^1}^{\frac{4}{3}} \\ &\leq \frac{\beta}{2} \|\partial_x \phi\|_{L^2}^2 + C_{\beta} (1 + \|\phi\|_{L^2}^2) \|F(U^r)\|_{L^1}^{\frac{4}{3}}.\end{aligned}\tag{4.2}$$

Substituting (4.2) into (4.1), integrating the resultant inequality with respect to t , noting

$$\partial_t \partial_x U^r = -f''(U^r) |\partial_x U^r|^2 - f'(U^r) \partial_x^2 U^r,$$

$\|F(U^r)\|_{L^1}^{\frac{4}{3}} \leq C_{\alpha, u_{\pm}} \|\partial_x U^r\|_{L^2}^{\frac{8}{3}} + C_{\alpha, \beta, u_{\pm}} \|\partial_x^2 U^r\|_{L^1}^{\frac{4}{3}} + \gamma^{\frac{4}{3}} \|\partial_x^4 U^r\|_{L^1}^{\frac{4}{3}} \in L_t^1(0, \infty)$ from Lemma 2.2, and using the Gronwall inequality, we obtain the desired estimate.

Thus, we complete the proof of Proposition 4.1.

Only from Proposition 4.1, using the Sobolev inequality, we can easily get the uniform boundedness of ϕ in the next lemma (cf. [17], [20], [26]).

Lemma 4.2. *There exists a positive constant C_{ϕ_0} such that*

$$\sup_{t \in [0, T], x \in \mathbb{R}} |\phi(t, x)| \leq C_{\phi_0}.$$

By the uniform boundedness of ϕ , Lemma 4.2, we note that the second term on the left-hand side of the *a priori* estimate in Proposition 4.1 can be replaced by the left-hand side of the following inequality as

$$\begin{aligned} & \int_0^t \int_{-\infty}^{\infty} \int_0^{\phi} (f'(\eta + U^r) - f'(U^r)) \, d\eta \, \partial_x U^r \, dx \, d\tau \\ & \geq C_{\phi_0}^{-1} \int_0^t \|\sqrt{\partial_x U^r} \phi(\tau)\|_{L^2}^2 \, d\tau. \end{aligned} \quad (4.3)$$

We now prepare the following propieties for the *a priori* estimates for the derivatives of ϕ in the next lemma.

Lemma 4.3. *There exists a positive constant C_{ϕ_0} such that*

$$\begin{aligned} & \int_0^t \|(f(\phi + U^r) - f(U^r))(\tau)\|_{L^2}^2 \, d\tau \leq C_{\phi_0} \quad (t \in [0, T]), \\ & \int_0^t \|F(U^r)(\tau)\|_{L^2}^2 \, d\tau \leq C_{\phi_0} \quad (t \in [0, T]). \end{aligned}$$

Proof of Lemma 4.3. The second estimate, that is, the time-integrability of $\|F(U^r)\|_{L^2}^2$, is obtained by Lemma 2.2. Noting Proposition 4.1 and using Lagrange's mean-value theorem, we can also obtain the first estimate, that is, the time-integrability of $\|f(\phi + U^r) - f(U^r)\|_{L^2}^2$.

Thus, the proof of Lemma 4.3 is completed.

Next, we show the *a priori* estimate for $\partial_x \phi$ and $\partial_x^2 \phi$ as follows.

Proposition 4.4. *There exists a positive constant C_{ϕ_0} such that*

$$\|\partial_x \phi(t)\|_{H^1}^2 + \int_0^t (\|\partial_t \phi(\tau)\|_{H^1}^2 + \|\partial_x^2 \phi(\tau)\|_{H^1}^2) \, d\tau \leq C_{\phi_0} \quad (t \in [0, T]).$$

Proof of Proposition 4.4. Multiplying the equation in (3.2) by $\partial_t \phi - \partial_x^2 \phi$ and integrating it with respect to x , we have, after integration by parts, that

$$\begin{aligned} & \frac{\beta+1}{2} \frac{d}{dt} \|\partial_x \phi(t)\|_{L^2}^2 + \frac{\gamma+1}{2} \frac{d}{dt} \|\partial_x^2 \phi(t)\|_{L^2}^2 \\ & + \|\partial_t \phi(t)\|_{L^2}^2 + \alpha \|\partial_t \partial_x \phi(t)\|_{L^2}^2 + \beta \|\partial_x^2 \phi(t)\|_{L^2}^2 + \gamma \|\partial_x^3 \phi(t)\|_{L^2}^2 \\ & = - \int_{-\infty}^{\infty} (\partial_t \phi - \partial_x^2 \phi) (f(\phi + U^r) - f(U^r)) dx \\ & + \int_{-\infty}^{\infty} (\partial_t \phi - \partial_x^2 \phi) F(U^r) dx. \end{aligned} \quad (4.4)$$

By using the Young inequality, we estimate the each terms on the right-hand side of (4.4) as follows.

$$\begin{aligned} & \left| \int_{-\infty}^{\infty} (\partial_t \phi - \partial_x^2 \phi) (f(\phi + U^r) - f(U^r)) dx \right| \\ & \leq \epsilon (\|\partial_t \phi\|_{L^2}^2 + \|\partial_x^2 \phi\|_{L^2}^2) + C_{\epsilon, \beta} \|f(\phi + U^r) - f(U^r)\|_{L^2}^2, \end{aligned} \quad (4.5)$$

$$\begin{aligned} & \left| \int_{-\infty}^{\infty} (\partial_t \phi - \partial_x^2 \phi) F(U^r) dx \right| \\ & \leq \epsilon (\|\partial_t \phi\|_{L^2}^2 + \|\partial_x^2 \phi\|_{L^2}^2) + C_{\epsilon, \beta} \|F(U^r)\|_{L^2}^2, \end{aligned} \quad (4.6)$$

for $\epsilon > 0$. Choosing ϵ suitably small, substituting (4.5) and (4.6) into (4.4), and using Lemma 4.3, we obtain the desired estimate.

Thus, we complete the proof of Proposition 4.4.

Remark 4.5. Similarly to Lemma 4.2, we have the uniform boundedness of $\partial_x \phi$ by using Proposition 4.4 that

$$\sup_{t \in [0, T], x \in \mathbb{R}} |\partial_x \phi(t, x)| \leq C_{\phi_0}.$$

We further show the *a priori* estimate for $\partial_x^2 \phi$ and $\partial_x^3 \phi$ as follows.

Proposition 4.6. *There exists a positive constant C_{ϕ_0} such that*

$$\|\partial_x^2 \phi(t)\|_{H^1}^2 + \int_0^t (\|\partial_t \partial_x \phi(\tau)\|_{H^1}^2 + \|\partial_x^3 \phi(\tau)\|_{H^1}^2) d\tau \leq C_{\phi_0} \quad (t \in [0, T]).$$

Proof of Proposition 4.6. Multiplying the equation in (3.2) by $\partial_x^4 \phi - \partial_t \partial_x^2 \phi$

and integrating it with respect to x , we have, after integration by parts, that

$$\begin{aligned}
& \frac{\beta+1}{2} \frac{d}{dt} \|\partial_x^2 \phi(t)\|_{L^2}^2 + \frac{\alpha+\gamma}{2} \frac{d}{dt} \|\partial_x^3 \phi(t)\|_{L^2}^2 \\
& + \|\partial_t \partial_x \phi(t)\|_{L^2}^2 + \alpha \|\partial_t \partial_x^2 \phi(t)\|_{L^2}^2 + \beta \|\partial_x^3 \phi(t)\|_{L^2}^2 + \gamma \|\partial_x^4 \phi(t)\|_{L^2}^2 \\
& = - \int_{-\infty}^{\infty} (\partial_x^4 \phi - \partial_t \partial_x^2 \phi) (f(\phi + U^r) - f(U^r)) dx \\
& + \int_{-\infty}^{\infty} (\partial_x^4 \phi - \partial_t \partial_x^2 \phi) F(U^r) dx.
\end{aligned} \tag{4.7}$$

The each terms on the right-hand side of (4.7) can be estimated quite similarly to (4.5) and (4.6). Therefore, we obtain the desired estimate by using Lemma 4.3.

Thus, we complete the proof of Proposition 4.6.

Remark 4.7. Similarly to Lemma 4.2 and remark 4.5, we have the uniform boundedness of $\partial_x^2 \phi$ by using Proposition 4.6 that

$$\sup_{t \in [0, T], x \in \mathbb{R}} |\partial_x^2 \phi(t, x)| \leq C_{\phi_0}.$$

5. Remarks on the uniform estimates

It is worthwhile to mention the uniform estimates of the solution ϕ to (3.2). By using the Sobolev inequality to ϕ , $\partial_x \phi$, $\partial_x^2 \phi$, $\partial_x^3 \phi$, $\partial_x^4 \phi$, $\partial_t \phi$ and $\partial_t \partial_x \phi$, and integration by parts, we obtain

$$\begin{aligned}
\sup_{x \in \mathbb{R}} |\phi(t, x)| & \leq \sqrt{2} \|\phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{2}} \\
& \leq \min \left\{ \sqrt{2} \|\phi(t)\|_{L^2}^{\frac{3}{4}} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{4}}, \|\phi(t)\|_{H^1} \right\} \\
& \leq \sqrt{2} \|\phi(t)\|_{L^2}^{\frac{3}{4}} \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{8}} \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{8}} \\
& \leq \sqrt{2} \|\phi(t)\|_{L^2}^{\frac{3}{4}} \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{8}} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{16}} \|\partial_x^4 \phi(t)\|_{L^2}^{\frac{1}{16}},
\end{aligned} \tag{5.1}$$

$$\begin{aligned}
\sup_{x \in \mathbb{R}} |\partial_x \phi(t, x)| & \leq \sqrt{2} \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{2}} \\
& \leq \min \left\{ \sqrt{2} \|\partial_x \phi(t)\|_{L^2}^{\frac{3}{4}} \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{4}}, \|\partial_x \phi(t)\|_{H^1} \right\} \\
& \leq \sqrt{2} \|\partial_x \phi(t)\|_{L^2}^{\frac{3}{4}} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{8}} \|\partial_x^4 \phi(t)\|_{L^2}^{\frac{1}{8}},
\end{aligned} \tag{5.2}$$

$$\begin{aligned}
\sup_{x \in \mathbb{R}} |\partial_x^2 \phi(t, x)| & \leq \sqrt{2} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{2}} \\
& \leq \min \left\{ \sqrt{2} \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{3}{4}} \|\partial_x^4 \phi(t)\|_{L^2}^{\frac{1}{4}}, \|\partial_x^2 \phi(t)\|_{H^1} \right\},
\end{aligned} \tag{5.3}$$

$$\sup_{x \in \mathbb{R}} |\partial_x^3 \phi(t, x)| \leq \sqrt{2} \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_x^4 \phi(t)\|_{L^2}^{\frac{1}{2}} \leq \|\partial_x^3 \phi(t)\|_{H^1}, \quad (5.4)$$

$$\sup_{x \in \mathbb{R}} |\partial_t \phi(t, x)| \leq \sqrt{2} \|\partial_t \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_t \partial_x \phi(t)\|_{L^2}^{\frac{1}{2}} \leq \|\partial_t \phi(t)\|_{H^1}, \quad (5.5)$$

$$\sup_{x \in \mathbb{R}} |\partial_t \partial_x \phi(t, x)| \leq \sqrt{2} \|\partial_t \partial_x \phi(t)\|_{L^2}^{\frac{1}{2}} \|\partial_t \partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{2}} \leq \|\partial_t \partial_x \phi(t)\|_{H^1}, \quad (5.6)$$

Noting the first term on the *a priori* estimates in Theorem 3.3, that is,

$$\begin{aligned} & \|\phi(t)\|_{H^3}^2 + \int_0^t \|(\sqrt{\partial_x U^r} \phi)(\tau)\|_{L^2}^2 d\tau \\ & + \int_0^t \|\partial_x \phi(\tau)\|_{H^3}^2 d\tau + \int_0^t \|\partial_t \phi(\tau)\|_{H^2}^2 d\tau \leq C_{\phi_0}, \end{aligned} \quad (5.7)$$

and using (5.1)-(5.6), we arrive at

$$\sup_{x \in \mathbb{R}} |\phi(t, x)| \leq C_{\phi_0} \min \left\{ \min_{k=1,2,3,4} \|\partial_x^k \phi(t)\|_{L^2}^{\frac{1}{2k}}, \|\phi(t)\|_{H^1} \right\}, \quad (5.8)$$

$$\sup_{x \in \mathbb{R}} |\partial_x \phi(t, x)| \leq C_{\phi_0} \min \left\{ \min_{k=1,2,3} \|\partial_x^{k+1} \phi(t)\|_{L^2}^{\frac{1}{2k}}, \|\partial_x \phi(t)\|_{L^2}^{\frac{1}{2}}, \|\partial_x \phi(t)\|_{H^1} \right\}, \quad (5.9)$$

$$\sup_{x \in \mathbb{R}} |\partial_x^2 \phi(t, x)| \leq C_{\phi_0} \min \left\{ \|\partial_x^2 \phi(t)\|_{L^2}^{\frac{1}{2}}, \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{2}}, \|\partial_x^4 \phi(t)\|_{L^2}^{\frac{1}{4}}, \|\partial_x^2 \phi(t)\|_{H^1} \right\}, \quad (5.10)$$

$$\sup_{x \in \mathbb{R}} |\partial_x^3 \phi(t, x)| \leq C_{\phi_0} \min \left\{ \|\partial_x^3 \phi(t)\|_{L^2}^{\frac{1}{2}}, \|\partial_x^4 \phi(t)\|_{L^2}^{\frac{1}{2}} \right\}, \quad (5.11)$$

for some $C_{\phi_0} > 0$. Then, by (5.4), (5.5) and (5.8)-(5.11), noting the interpolation, we obtain the uniform estimates as follows.

Proposition 5.1. *There exists a positive constant C_{ϕ_0} such that*

$$\begin{aligned} & \int_0^t \left(\sup_{x \in \mathbb{R}} |\phi(\tau, x)| \right)^k d\tau \leq C_{\phi_0} \quad (2 \leq k \leq 32), \\ & \int_0^t \left(\sup_{x \in \mathbb{R}} |\partial_x \phi(\tau, x)| \right)^k d\tau \leq C_{\phi_0} \quad (2 \leq k \leq 16), \\ & \int_0^t \left(\sup_{x \in \mathbb{R}} |\partial_x^2 \phi(\tau, x)| \right)^k d\tau \leq C_{\phi_0} \quad (2 \leq k \leq 8), \\ & \int_0^t \left(\sup_{x \in \mathbb{R}} |\partial_x^3 \phi(\tau, x)| \right)^k d\tau \leq C_{\phi_0} \quad (2 \leq k \leq 4), \\ & \int_0^t \left(\sup_{x \in \mathbb{R}} |\partial_t \phi(\tau, x)| \right)^2 d\tau \leq C_{\phi_0}, \\ & \int_0^t \left(\sup_{x \in \mathbb{R}} |\partial_t \partial_x \phi(\tau, x)| \right)^2 d\tau \leq C_{\phi_0}, \end{aligned}$$

for $t \in [0, T]$.

References

- [1] C.J. Amick, J.L. Bona and M.E. Schonbek, *Decay of solutions of some nonlinear wave equations*, J. Differential Equations **81** (1989), pp. 1-49.
- [2] K. Andreiev, I. Egorova, T.L. Lange and G. Teschl, *Rarefaction waves of the Korteweg-de Vries equation via nonlinear steepest descent*, J. Differential Equations **10** (2016), pp. 5371-5410.
- [3] T.B. Benjamin, J.L. Bona and J.J. Mahony, *Model equations for long waves in nonlinear dispersive system*, Phil. Trans. R. Soc. Lond. Ser. A **272** (1972), pp. 47-78.
- [4] J.L. Bona and M.E. Schonbek, *Travelling-wave solutions to the Korteweg-de Vries-Burgers equation*, Proc. Roy. Soc. Edinburgh **101A** (1985), pp. 207-226.
- [5] J.L. Bona, S.V. Rajopadhye and M.E. Schonbek, *Models for the propagation of bores I. Two dimensional theory*, Differential Integral Equations **7** (1994), pp. 699-734.
- [6] R.P. Chhabra, *Bubbles, drops and particles in non-Newtonian Fluids*, CRC, Boca Raton, FL (2006).
- [7] R.P. Chhabra, *Non-Newtonian Fluids: An Introduction*, URL <http://www.physics.iitm.ac.in/~compflu/Lect-notes/chhabra.pdf>.
- [8] R.P. Chhabra and J.F. Richardson, *Non-Newtonian flow and applied rheology*, 2nd edn. Butterworth-Heinemann, Oxford (2008).
- [9] A. de Waele, *Viscometry and plastometry*, J. Oil Colour Chem. Assoc. **6** (1923), pp. 33-69.
- [10] R. Duan, L.-L. Fan, J.-S. Kim and L.-Q. Xie, *Nonlinear stability of strong rarefaction waves for the generalized KdV-Burgers-Kuramoto equation with large initial perturbation*, Nonlinear Anal. **73** (2010), pp. 3254-3267.
- [11] R. Duan and H.-J. Zhao, *Global stability of strong rarefaction waves for the generalized KdV-Burgers equation*, Nonlinear Anal. **66** (2007), pp. 1100-1117.
- [12] I. Egorova, K. Grunert and G. Teschl, *On the Cauchy problem for the Korteweg-de Vries equation with steplike finite-gap initial data I. Schwartz-type perturbations*, Nonlinearity **22** (2009), pp. 1431-1457.
- [13] I. Egorova and G. Teschl, *On the Cauchy problem for the Korteweg-de Vries equation with steplike finite-gap initial data II. Perturbations with finite moments*, J. Anal. Math. **115** (2011), pp. 71-101.
- [14] E. Harabetian, *Rarefaction and large time behavior for parabolic equations and monotone schemes*, Comm. Math. Phys. **114** (1988), pp. 527-536.

- [15] I. Hashimoto and A. Matsumura, *Large time behavior of solutions to an initial boundary value problem on the half space for scalar viscous conservation law*, Methods Appl. Anal. **14** (2007), pp. 45-59.
- [16] Y. Hattori and K. Nishihara, *A note on the stability of rarefaction wave of the Burgers equation*, Japan J. Indust. Appl. Math. **8** (1991), pp. 85-96.
- [17] A.M. Il'in, A.S. Kalašnikov and O.A. Oleĭnik, *Second-order linear equations of parabolic type*, Uspekhi Math. Nauk SSSR **17** (1962), pp. 3-146 (in Russian); English translation in Russian Math. Surveys **17** (1962), pp. 1-143.
- [18] A.M. Il'in and O.A. Oleĭnik, *Asymptotic behavior of the solutions of the Cauchy problem for some quasi-linear equations for large values of the time*, Mat. Sb. **51** (1960), pp. 191-216 (in Russian).
- [19] P. Jahangiri, R. Streblow and D. Müller, *Simulation of Non-Newtonian Fluids using Modelica*, Proceedings of the 9th International Modelica Conference September 3-5, Munich, Germany, (2012), pp. 57-62.
- [20] Y. Kanel', *On a model system of one-dimensional gas motion*, Differentsial'nye Uravneniya **4** (1968), pp. 374-380.
- [21] C.I. Kondo and C.M. Webler *The generalized BBM-Burger equations with non-linear dissipative term: existence and convergence results*, Appl. Anal. **87** (2008), pp. 977-995.
- [22] C.I. Kondo and C.M. Webler *Higher order for the generalized BBM-Burgers equation: existence and convergence results*, Appl. Anal. **88** (2009), pp. 1085-1101.
- [23] C.I. Kondo and C.M. Webler *Higher order for the generalized BBM-Burgers equation: existence and convergence results*, Acta Appl. Math. **111** (2010), pp. 45-64.
- [24] C.I. Kondo and C.M. Webler *The generalized BBM-Burgers equations: convergence results for conservation law with discontinuous flux function*, Appl. Anal. **95** (2016), pp. 503-523.
- [25] O.A. Ladyženskaja, *New Equations for the Description of the Viscous Incompressible Fluids and Solvability in the Large of the Boundary Value Problems for Them*, in "Boundary Value Problems of Mathematical Physics V", Amer. Math. Soc., Providence, Rhode. Island, 1970.
- [26] O. A. Ladyženskaja, V. A. Solonnikov and N. N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, Transl. Math. Monographs, vol. 23, Amer. Math. Soc., Providence, Rhode. Island, 1968.
- [27] P.D. Lax, *Hyperbolic systems of conservation laws II*, Comm. Pure Appl. Math. **10** (1957), pp. 537-566.

- [28] H.W. Liepmann and A. Roshko, *Elements of Gas Dynamics*, John Wiley & Sons, Inc., New York, 1957.
- [29] T.-P. Liu, A. Matsumura and K. Nishihara, *Behaviors of solutions for the Burgers equation with boundary corresponding to rarefaction waves*, SIAM J. Math. Anal. **29** (1998), pp. 293-308.
- [30] J. Málek, *Some frequently used models for non-Newtonian fluids*, URL <http://www.karlin.mff.cuni.cz/~malek/new/images/Lecture4.pdf>.
- [31] J. Málek, D. Pražák and M. Steinhauer, *On the existence and regularity of solutions for degenerate power-law fluids*, Differential Integral Equations **19** (2006), pp. 449-462.
- [32] A. Matsumura and K. Nishihara, *Asymptotic toward the rarefaction wave of solutions of a one-dimensional model system for compressible viscous gas*, Japan J. Appl. Math. **3** (1986), pp. 1-13.
- [33] A. Matsumura and K. Nishihara, *Asymptotics toward the rarefaction wave of the solutions of Burgers' equation with nonlinear degenerate viscosity*, Nonlinear Anal. TMA **23** (1994), pp. 605-614.
- [34] A. Matsumura and K. Nishihara, *Asymptotic stability of traveling waves for scalar viscous conservation laws with non-convex nonlinearity*, Comm. Math. Phys. **165** (1994), pp. 83-96.
- [35] A. Matsumura and N. Yoshida, *Asymptotic behavior of solutions to the Cauchy problem for the scalar viscous conservation law with partially linearly degenerate flux*, SIAM J. Math. Anal. **44** (2012), pp. 2526-2544.
- [36] A. Matsumura and N. Yoshida, *Global asymptotics toward the rarefaction waves for solutions to the Cauchy problem of the scalar conservation law with nonlinear viscosity*, Osaka J. Math. **57** (2020), pp. 187-205.
- [37] M. Mei, *Large-time behavior of solution for generalized Benjamin-Bona-Mahony-Burgers equations*, Nonlinear Anal. **33** (1998), pp. 699-714.
- [38] M. Mei, *L^q -decay rates of solutions for generalized Benjamin-Bona-Mahony-Burgers equations*, J. Differential Equations **158** (1999), pp. 314-340.
- [39] M. Mei and C. Schmeiser, *Asymptotic profiles of solutions for the BBM-Burgers equation*, Funkcialaj Ekvacioj **44** (2001), pp. 151-170.
- [40] P.I. Naumkin, *Large-time asymptotic of a step for the Benjamin-Bona-Mahony-Burgers equation*, Proc. Roy. Soc. Edinburgh **126A** (1996), pp. 1-18.
- [41] K. Nishihara and S.V. Rajopadhye, *Asymptotic behavior of solutions to the Korteweg-de Vries-Burgers equation*, Differential Integral Equations **11** (1998), pp. 85-93.

- [42] S. Osher and J. Ralston *L^1 stability of traveling waves with applications to convective porous media flow*, Comm. Pure Appl. Math. **35** (1982), pp. 737-751.
- [43] W. Ostwald, *Über die Geschwindigkeitsfunktion der Viskosität disperser Systeme*, I. Colloid Polym. Sci. **36** (1925), pp. 99-117 (in German).
- [44] D.H. Peregrine, *Calculations of the development of an undular bore*, J. Fluid Mech. **25** (1966), pp. 321-330.
- [45] S.V. Rajopadhye, *Decay rates for the solutions of model equations for bore propagation*, Proc. Roy. Soc. Edinburgh **125A** (1995), pp. 371-398.
- [46] J. Rashindinia, O. Nikan and L. Khoddam, *Numerically stable scheme to approximate the nonlinear KdV-Benjamin-Bona-Mahony-Burger's equation*, 4th International Conference on Combinatorics, Cryptography, Computer Science and Computing (2019), pp. 1-10.
- [47] L.-Z. Ruan, W.-L. Gao and J. Chen, *Asymptotic stability of the rarefaction wave for the generalized KdV-Burgers-Kuramoto equation*, Nonlinear Anal. **68** (2008), pp. 402-411.
- [48] T. Sochi, *Pore-Scale Modeling of Non-Newtonian Flow in Porous Media*, PhD thesis, Imperial College London, 2007.
- [49] Y. Wang, *On time periodic solutions to the generalized BBM-Burgers equation with time-dependent periodic external force*, Math. Model. Anal. **25** (2020), pp. 184-197.
- [50] Z.-A. Wang and C.-J. Zhu, *Stability of the rarefaction wave for the generalized KdV-Burgers equation*, Acta Math. Sci. **22B**(3) (2002), pp. 319-328.
- [51] H. Xu and B. Li, *Global existence and bounded estimate of solutions of the BBM-Burgers equation*, Wuhan Univ. J. Nat. Sci. **21** (2016), pp. 428-432.
- [52] H. Yin, H. Zhao and J. Kim, *Convergence rates of solutions toward boundary layer solutions for generalized Benjamin-Bona-Mahony-Burgers equations in the half-space*, J. Differential Equations **245** (2008), pp. 3144-3216.
- [53] N. Yoshida, *Decay properties of solutions toward a multiwave pattern for the scalar viscous conservation law with partially linearly degenerate flux*, Nonlinear Anal. **96** (2014), pp. 189-210.
- [54] N. Yoshida, *Decay properties of solutions to the Cauchy problem for the scalar conservation law with nonlinearly degenerate viscosity*, Nonlinear Anal. **128** (2015), pp. 48-76.
- [55] N. Yoshida, *Large time behavior of solutions toward a multiwave pattern for the Cauchy problem of the scalar conservation law with degenerate flux and viscosity*, in: Mathematical Analysis in Fluid and Gas Dynamics, Sūrikaiseikikenkyūsho Kōkyūroku **1947** (2015), pp. 205-222.

- [56] N. Yoshida, *Asymptotic behavior of solutions toward a multiwave pattern for the scalar conservation law with the Ostwald-de Waele-type viscosity*, SIAM J. Math. Anal. **49** (2017), pp. 2009-2036.
- [57] N. Yoshida, *Decay properties of solutions toward a multiwave pattern to the Cauchy problem for the scalar conservation law with degenerate flux and viscosity*, J. Differential Equations **263** (2017), pp. 7513-7558.
- [58] N. Yoshida, *Asymptotic behavior of solutions toward the viscous shock waves to the Cauchy problem for the scalar conservation law with nonlinear flux and viscosity*, SIAM J. Math. Anal. **50** (2018), pp. 891-932.
- [59] N. Yoshida, *Asymptotic behavior of solutions toward the rarefaction waves to the Cauchy problem for the scalar diffusive dispersive conservation laws*, Nonlinear Anal. **189** (2019), pp. 1-19.
- [60] N. Yoshida, *Global structure of solutions toward the rarefaction waves for the Cauchy problem of the scalar conservation law with nonlinear viscosity*, J. Differential Equations **269** (2020), pp. 10350-10394.
- [61] N. Yoshida, *Asymptotic behavior of solutions toward a multiwave pattern to the Cauchy problem for the dissipative wave equation with partially linearly degenerate flux*, Funkcialaj Ekvacioj **64** (2021), pp. 49-73.
- [62] N. Yoshida, *Asymptotic behavior of solutions toward the constant state to the Cauchy problem for the non-viscous diffusive dispersive conservation law*, Preprint, arXiv:2107.07874.
- [63] N. Yoshida, *Global asymptotic stability of the rarefaction waves to the Cauchy problem for the scalar non-viscous diffusive dispersive conservation laws*, Preprint.
- [64] H. Zhao and B. Xuan, *Existence and convergence of solutions for the generalized BBM-Burgers equations with dissipative term*, Nonlinear Anal. **28** (1997), pp. 1835-1849.