# A CHARACTERIZATION OF CONTINUOUS q-JACOBI, CHEBYSHEV OF THE FIRST KIND AND AL-SALAM CHIHARA POLYNOMIALS

K. CASTILLO, D. MBOUNA, AND J. PETRONILHO

ABSTRACT. The purpose of this note is to characterize those orthogonal polynomials sequences  $(P_n)_{n>0}$  for which

$$\pi(x)\mathcal{D}_q P_n(x) = (a_n x + b_n)P_n(x) + c_n P_{n-1}(x), \quad n = 0, 1, 2, \dots,$$

where  $\mathcal{D}_q$  is the Askey-Wilson operator,  $\pi$  is a polynomial of degree at most 2, and  $(a_n)_{n\geq 0}$ ,  $(b_n)_{n\geq 0}$  and  $(c_n)_{n\geq 0}$  are sequences of complex numbers such that  $c_n \neq 0$  for  $n = 1, 2, \ldots$ 

### 1. INTRODUCTION AND MAIN RESULTS

Let  $\pi$  be a nonzero polynomial of degree at most 2 and consider three sequences of complex numbers  $(a_n)_{n\geq 0}$ ,  $(b_n)_{n\geq 0}$  and  $(c_n)_{n\geq 0}$ . Al-Salam and Chihara [2] proved that the only orthogonal polynomial sequences (OPS), say  $(P_n)_{n\geq 0}$ , that satisfy

(1.1) 
$$\pi(x) DP_n(x) = (a_n x + b_n) P_n(x) + c_n P_{n-1}(x),$$

are those of Hermite, Laguerre, Jacobi, and Bessel. Here D denotes the standard derivative with respect to x. Consider now (1.1) with D replaced by the Askey-Wilson operator,

(1.2) 
$$(\mathcal{D}_q f)(x) = \frac{\check{f}(q^{1/2}z) - \check{f}(q^{-1/2}z)}{\check{e}(q^{1/2}z) - \check{e}(q^{-1/2}z)}, \quad z = e^{i\theta},$$

where  $\check{f}(z) = f((z+1/z)/2) = f(\cos\theta)$  for each polynomial f and e(x) = x. Here 0 < q < 1 and  $\theta$  is not necessarily a real number (see [6, p. 300]). The following conjecture is the first part of [6, Conjecture 24.7.8], rewritten using the (2.1) below.

**Conjecture 1.1.** Let  $(P_n)_{n\geq 0}$  be a monic OPS and  $\pi$  be a polynomial of degree at most 2 which does not depend on n. If  $(P_n)_{n\geq 0}$  satisfies

(1.3) 
$$\pi(x)\mathcal{D}_{q}P_{n}(x) = (a_{n}x + b_{n})P_{n}(x) + c_{n}P_{n-1}(x),$$

then  $(P_n)_{n\geq 0}$  are continuous q-Jacobi polynomials, Al-Salam-Chihara polynomials, or special or limiting cases of them.

Date: January 3, 2022.

<sup>2000</sup> Mathematics Subject Classification. Primary 42C05; Secondary 33C45.

Key words and phrases. Askey-Wilson operator, Al-Salam-Chihara polynomials, Chebyshev polynomials of the first kind, continuous q-Jacobi polynomials.

Our beloved friend and mentor J. Petronilho passed away unexpectedly on August 27, 2021 while we were preparing the final version of this manuscript. To him, our eternal gratitude.

Ismail himself proved that the continuous q-Jacobi polynomials indeed satisfy (1.3) for suitable polynomial  $\pi$  and parameters  $a_n$ ,  $b_n$ , and  $c_n$  (cf. [6, Theorem 15.5.2]). Al-Salam [1] proved Conjecture 1.1 for  $\pi(x) = 1$  (see [5, Proposition 5.3.1] for an alternative proof of this result based on the approach developed in this note), by characterizing the Rogers q-Hermite polynomials,  $P_n(x) = H_n(x|q)$ , as the only OPS that fulfill  $\mathcal{D}_q P_n = c_n P_{n-1}$  for  $n = 1, 2, \ldots$  Recall that the monic continuous q-Jacobi polynomials,  $\hat{P}_n^{(a,b)}(x|q)$ , depend on two real parameters a and b, and they are characterized by the three-term recurrence relation

$$x\widehat{P}_{n}^{(a,b)}(x|q) = \widehat{P}_{n+1}^{(a,b)}(x|q) + \frac{1}{2} \left( q^{(2a+1)/4} + q^{-(2a-1)/4} - y_n - z_n \right) \widehat{P}_{n}^{(a,b)}(x|q) + \frac{1}{4} y_{n-1} z_n \, \widehat{P}_{n-1}^{(a,b)}(x|q),$$

n = 0, 1, ..., being

$$y_n = \frac{(1-q^{n+a+1})(1-q^{n+a+b+1})(1+q^{n+(a+b+1)/2})(1+q^{n+(a+b+2)/2})}{q^{(2a+1)/4}(1-q^{2n+a+b+1})(1-q^{2n+a+b+2})}$$
$$z_n = \frac{q^{(2a+1)/4}(1-q^n)(1-q^{n+b})(1+q^{n+(a+b)/2})(1+q^{n+(a+b+1)/2})}{(1-q^{2n+a+b})(1-q^{2n+a+b+1})},$$

while the monic Al-Salam-Chihara polynomials,  $Q_n(x; c, d|q)$ , which also depend on two parameters c and d, are characterized by

$$xQ_n(x;c,d|q) = Q_{n+1}(x;c,d|q) + \frac{1}{2}(c+d)q^n Q_n(x;c,d|q) + \frac{1}{4}(1-cdq^{n-1})(1-q^n) Q_{n-1}(x;c,d;q),$$

 $n = 0, 1, \ldots$ , provided we define  $\widehat{P}_{-1}^{(a,b)}(x|q) = Q_{-1}(x;c,d|q) = 0$  (see e.g. [6]). Further, up to normalization, the Rogers q-Hermite polynomials are the special case c = d = 0 of the Al-Salam-Chihara polynomials.

The following two theorems summarize the main results of this note and, together with Al-Salam's theorem, give positive answer to Conjecture 1.1.

**Theorem 1.2.** The Al-Salam Chihara polynomials with nonzero parameters c and d such that  $c/d = q^{\pm 1/2}$  are the only OPS satisfying (1.3) for deg  $\pi = 1$ .

**Theorem 1.3.** The Chebyschev polynomials of the first kind and the continuous q-Jacobi polynomials are the only OPS satisfying (1.3) for deg  $\pi = 2$ .

The two previous theorems give more than a simple answer to Conjecture 1.1. In fact, we now know, for instance, that Al-Salam Chihara polynomials appear only when deg  $\pi = 1$  and for  $c/d = q^{\pm 1/2}$ , or that continuous q-Jacobi polynomials are exclusively related to the case deg  $\pi = 2$ , or that Chebyschev polynomials of the first type are the only "limiting case" that satisfy Conjecture 1.1.

Remark 1.4. Koornwinder [4, Section 4] gave a structure formula (and resulting lowering formula) for Askey-Wilson polynomials. Necessarily, because of Conjecture 1.1 settled in the present paper, the left-hand side in Koornwinder's formula has a more general form than in (1.3). With the usual parameters a, b, c, d for Askey-Wilson polynomials, substitution of  $c = q^{1/2}$ ,  $d = -q^{1/2}$  in these formulas gives similar formulas for continuous q-Jacobi polynomials with q replaced by  $q^2$  (see [4, Section 5]). In a personal communication Koornwinder pointed out to us that Ismail's and his formulas in the continuous q-Jacobi case are the same modulo polynomial multiples of the second order q-difference formula and the three-term recurrence relation for continuous q-Jacobi polynomials. Since this work is an application of the theory developed in our previous paper [3], in the next sections, we suppose that the reader has [3] at hand and we shall use its notation, definitions, and results.

# 2. Preliminary results

Taking  $e^{i\theta} = q^s$  in (1.2),  $\mathcal{D}_q$  reads

$$\mathcal{D}_q f(x(s)) = \frac{f\left(x(s+\frac{1}{2})\right) - f\left(x(s-\frac{1}{2})\right)}{x(s+\frac{1}{2}) - x(s-\frac{1}{2})}, \quad x(s) = \frac{1}{2}(q^s + q^{-s}).$$

We define an operator  $\mathcal{S}_q : \mathbb{R}[x] \to \mathbb{R}[x]$  by

$$S_q f(x(s)) = \frac{f(x(s+\frac{1}{2})) + f(x(s-\frac{1}{2}))}{2}.$$

Hereafter, we denote  $X = x(s) = (q^s + q^{-s})/2$  with 0 < q < 1. Recall that a monic OPS  $(P_n)_{n\geq 0}$  satisfies the following three term recurrence relation (TTRR):

(2.1) 
$$XP_n(X) = P_{n+1}(X) + B_n P_n(X) + C_n P_{n-1}(X), \quad n = 0, 1, 2, \dots,$$

with  $P_{-1}(X) = 0$  and  $B_n \in \mathbb{C}$  and  $C_{n+1} \in \mathbb{C} \setminus \{0\}$  for each  $n = 0, 1, 2, \ldots$  We start by showing that all monic OPS,  $(P_n)_{n\geq 0}$ , satisfying (1.3) belongs to a well known class of OPS and then we prove that the coefficients of the associated TTRR satisfy a system of non linear equations that will be solved in the next section.

**Lemma 2.1.** Let  $\mathbf{u} \in \mathcal{P}^*$  be a regular functional such that its corresponding monic OPS  $(P_n)_{n\geq 0}$  satisfies (1.3) subject to the condition  $c_n \neq 0$  for n = 1, 2, ... Then  $\mathbf{u}$  is x-classical; that is  $\mathcal{D}_q(\phi \mathbf{u}) = \mathcal{S}_q(\psi \mathbf{u})$  with  $\deg \phi \leq 2$  and  $\deg \psi \leq 1$ . Moreover,  $\psi$  and  $\phi$  are polynomials given by

(2.2) 
$$\psi(X) = X - B_0, \quad \phi(X) = (\mathfrak{a}X - \mathfrak{b})(X - B_0) - (\mathfrak{a} + \alpha)C_1 ,$$

where

(2.3) 
$$\mathfrak{a} = \frac{(a_2C_2 + c_2)C_1}{(a_1C_1 + c_1)C_2} - \alpha, \quad \mathfrak{b} = -B_0 + (\mathfrak{a} + \alpha)B_1 - \frac{b_1 + a_1B_1}{c_1 + a_1C_1}C_1.$$

(Here  $B_0$ ,  $B_1$ ,  $C_1$ , and  $C_2$  are coefficients of the TTRR (2.1) satisfied by  $(P_n)_{n\geq 0}$ .)

*Proof.* Let  $(\mathbf{a}_n)_{n\geq 0}$  be the dual basis associated to the monic OPS  $(P_n)_{n\geq 0}$ . We claim that

(2.4) 
$$\mathcal{D}_q(\pi \mathbf{u}) = R_1 \mathbf{u}, \quad R_1(X) = -\frac{a_1 C_1 + c_1}{C_1} (X - B_0),$$

with  $a_1C_1 + c_1 \neq 0$ . Indeed, using (1.3) and (2.1), we have

$$\langle \mathcal{D}_q(\pi \mathbf{a}_0), P_j \rangle = -\langle \mathbf{a}_0, \pi \mathcal{D}_q P_j \rangle = -a_j \delta_{0,j+1} - (a_j B_j + b_j) \delta_{0,j} - (c_j + a_j C_j) \delta_{1,j},$$

for fixed  $j \in \mathbb{N}_0$ . Taking n = 0 in (1.3), we find  $a_0 = b_0 = 0$ , and since  $\langle \mathbf{u}, P_n^2 \rangle \mathbf{a}_n = P_n \mathbf{u}$  and  $C_{n+1} = \langle \mathbf{u}, P_{n+1}^2 \rangle / \langle \mathbf{u}, P_n^2 \rangle$ , we obtain

$$\mathcal{D}_q(\pi \mathbf{a}_0) = \sum_{j=0}^{\infty} \left\langle \mathcal{D}_q(\pi \mathbf{a}_0), P_j \right\rangle \mathbf{a}_j = -(c_1 + a_1 C_1) \mathbf{a}_1.$$

If  $c_1 + a_1C_1 = 0$ , then  $\mathbf{D}_x(\pi \mathbf{u}) = 0$ , hence  $0 = \langle \mathcal{D}_q(\pi \mathbf{u}), f \rangle = -\langle \pi \mathbf{u}, \mathcal{D}_q f \rangle, \forall f \in \mathcal{P}$ . This implies  $\pi \mathbf{u} = \mathbf{0}$ . But this is impossible, since  $\pi \neq 0$  and  $\mathbf{u}$  is regular. So  $c_1 + a_1 C_1 \neq 0$ . Hence (2.4) holds. Applying  $\mathcal{D}_q$  to both sides of (2.1), and using [3, (2.28)], yields

$$\mathcal{S}_q P_n(X) = -\alpha X \mathcal{D}_q P_n(X) + \mathcal{D}_q P_{n+1}(X) + B_n \mathcal{D}_q P_n(X) + C_n \mathcal{D}_q P_{n-1}(X).$$

Multiplying both sides of this equality by  $\pi(X)$  and using (1.3) and (2.1), we obtain (2.5)

$$\pi(X)\mathcal{S}_q P_n(X) = r_n^{[1]} P_{n+2}(X) + r_n^{[2]} P_{n+1}(X) + r_n^{[3]} P_n(X) + r_n^{[4]} P_{n-1}(X) + r_n^{[5]} P_{n-2}(X)$$

for each n = 0, 1, 2, ..., where

$$r_n^{[1]} = a_{n+1} - \alpha a_n,$$
  

$$r_n^{[2]} = g_{n+1} - \alpha g_n + a_n (B_n - \alpha B_{n+1}),$$
  

$$r_n^{[3]} = s_{n+1} - \alpha s_n + g_n (1 - \alpha) B_n + a_{n-1} C_n - \alpha a_n C_{n+1},$$
  

$$r_n^{[4]} = (g_{n-1} - \alpha g_n) C_n + s_n (B_n - \alpha B_{n-1}),$$
  

$$r_n^{[5]} = C_n s_{n-1} - \alpha C_{n-1} s_n,$$

and  $g_n = b_n + a_n B_n$ ,  $s_n = c_n + a_n C_n$ . For a fixed  $j \in \mathbb{N}_0$ , using (2.5) we obtain

$$\langle \mathcal{S}_q(\pi \mathbf{a}_0), P_j \rangle = \langle \mathbf{a}_0, \pi \mathcal{S}_q P_j \rangle = r_j^{[1]} \delta_{0,j+2} + r_j^{[2]} \delta_{0,j+1} + r_j^{[3]} \delta_{0,j} + r_j^{[4]} \delta_{0,j-1} + r_j^{[5]} \delta_{0,j-2}.$$

Therefore,

$$S_q(\pi \mathbf{a}_0) = \sum_{j=0}^{\infty} \left\langle S_q(\pi \mathbf{a}_0), P_j \right\rangle \mathbf{a}_j = r_0^{[3]} \mathbf{a}_0 + r_1^{[4]} \mathbf{a}_1 + r_2^{[5]} \mathbf{a}_2,$$

and so

(2.6) 
$$S_q(\pi \mathbf{u}) = R_2 \mathbf{u}, \quad R_2(X) = r_0^{[3]} + \frac{r_1^{[4]}}{C_1} P_1(X) + \frac{r_2^{[5]}}{C_2 C_1} P_2(X).$$

Next, on the first hand, applying successively (2.6), [3, (2.47)] and (2.4), we obtain

(2.7) 
$$\mathcal{D}_q(R_2\mathbf{u}) = \frac{2\alpha^2 - 1}{\alpha} \mathcal{D}_q(R_1\mathbf{u}) + \frac{\mathbf{U}_1}{\alpha} \mathcal{D}_q(R_1\mathbf{u}).$$

On the other hand, using [3, (2.34)] with  $f = U_1$ , we obtain

 $\alpha \mathcal{D}_q(\mathbf{U}_1 R_1 \mathbf{u}) = \mathbf{U}_1 \mathcal{D}_q(R_1 \mathbf{u}) + (\alpha^2 - 1) \mathcal{S}_q(R_1 \mathbf{u}).$ 

Thus, from (2.7) we obtain  $\mathcal{D}_q((R_2 - U_1R_1)\mathbf{u}) = \mathcal{S}_q(\alpha R_1\mathbf{u})$ . This leads us to define

$$\psi(X) = X - B_0, \quad \phi(X) = -\frac{C_1}{\alpha(c_1 + a_1C_1)} \Big( R_2(X) - \mathbf{U}_1(X)R_1(X) \Big).$$

Clearly, deg  $\psi = 1$ , deg  $\phi \leq 2$  and  $\mathcal{D}_q(\phi \mathbf{u}) = \mathcal{S}_q(\psi \mathbf{u})$ . Finally, since  $a_0 = 0 = b_0$ , and setting (without lost of generality)  $c_0 = 0$  and  $C_0 = 0$ , we have

$$\phi''(0) = \frac{2(c_2 + a_2C_2)C_1}{(c_1 + a_1C_1)C_2} - 2\alpha, \ \phi'(0) = (1 - \mathfrak{a})B_0 - (\mathfrak{a} + \alpha)B_1 + \frac{b_1 + a_1B_1}{c_1 + a_1C_1}C_1,$$
  
$$\phi(0) = -(\mathfrak{a} + \alpha)C_1 - B_0\left(B_0 - (\mathfrak{a} + \alpha)B_1 + \frac{b_1 + a_1B_1}{c_1 + a_1C_1}C_1\right),$$

and the proof is complete.

**Lemma 2.2.** Let  $(P_n)_{n\geq 0}$  be a monic OPS satisfying (1.3). Then the coefficients  $B_n$  and  $C_n$  of the TTRR (2.1) satisfied by  $(P_n)_{n\geq 0}$  fulfill the following system of

difference equations:

$$(2.8) \quad a_{n+2} - 2\alpha a_{n+1} + a_n = 0, \ t_{n+2} - 2\alpha t_{n+1} + t_n = 0, \\ t_n = \frac{c_n}{C_n} = k_1 q^{n/2} + k_2 q^{-n/2}, \\ (2.9) \quad r_{n+3} B_{n+2} - (r_{n+2} + r_{n+1}) B_{n+1} + r_n B_n = 0, \quad r_n = t_n + a_n - a_{n-1}, \\ (2.10) \quad r_n \left( B_n^2 - 2\alpha B_n B_{n-1} + B_{n-1}^2 \right) \\ = (r_{n+1} + r_{n+2})(C_{n+1} - 1/4) - 2(1+\alpha)r_n(C_n - 1/4) \\ + (r_{n-1} + r_{n-2})(C_{n-1} - 1/4) \\ (2.11) \quad (1 - \alpha^2)b_n = 2(1 - \alpha)(a_n B_n + b_n)B_n^2 + (t_{n+1} + a_{n+1} - a_{n+2})B_{n+1}C_{n+1} + (t_n + a_{n-1} - a_{n-2})B_{n-1}C_n \\ + \left[ (2a_n - a_{n+2} - a_{n-1})C_{n+1} + (2a_n - a_{n+1} - a_{n-2})C_n \\ + (1 - 2\alpha)(c_n + c_{n+1}) + (\alpha^2 - 1)a_n \right] B_n + 2(b_n - \alpha b_{n+1})C_{n+1} \\ + 2(b_n - \alpha b_{n-1})C_n. \end{aligned}$$

In addition, the following relations hold:

(2.12) 
$$b_n = \gamma_n, \quad c_n = (b_n - b_{n-1}) \sum_{j=0}^{n-1} B_j + \pi(0) b_n, \quad \text{if } \deg \pi = 1,$$

(2.13) 
$$a_n = \gamma_n, \quad b_n = (a_n - a_{n-1}) \sum_{j=0}^{n-1} B_j + \pi'(0) a_n, \text{ if } \deg \pi = 2.$$

*Proof.* Applying the operator  $S_q$  to both sides of (2.1) and using [3, (2.29)], we deduce  $U_2(X)\mathcal{D}_qP_n(X) + \alpha XS_qP_n(X) = S_qP_{n+1}(X) + B_nS_qP_n(X) + C_nS_qP_{n-1}(X)$ . Multiplying both sides of this equality by  $\pi(X)$  and then using successively [3, (2.25)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), (1.3), (2.5), and (2.1), we obtain a vanishing linear combination of the polynomials  $P_{n+3}, P_{n+2}, \ldots, P_{n-3}$ . Thus, setting  $t_n = c_n/C_n$  for  $n = 1, 2, 3, \ldots$ , after straightforward computations we obtain (2.8)

together with the following equations:

(2.14) $(a_{n+1} - a_{n+2})B_{n+1} + (a_n - a_{n-1})B_n + b_{n+2} - 2\alpha b_{n+1} + b_n = 0.$ (2.15) $(a_{n+1} - a_{n+2} - t_{n+2})B_{n+1} + (a_n - a_{n-1} + t_{n+1} + t_n)B_n - t_{n-1}B_{n-1}$  $+ b_{n+1} - 2\alpha b_n + b_{n-1} = 0,$ (2.16) $(a_{n+1} - a_{n+2})B_{n+1}^2 + 2(1 - \alpha)a_n B_n^2 + (a_n - a_{n-1})B_n B_{n+1} + (a_n - a_{n+2})C_{n+1}$  $+ (b_{n+1} + b_n - 2\alpha b_{n+1})B_{n+1} + (b_{n+1} + b_n - 2\alpha b_n)B_n + (a_n - a_{n-2})C_n$  $+c_{n+2} - 2\alpha c_{n+1} + c_n = (1 - \alpha^2)a_n,$ (2.17) $(2(1-\alpha)a_n + t_n)B_n^2 + (t_n + a_{n-1} - a_{n-2})B_{n-1}^2 + (b_n + b_{n-1} - 2\alpha b_n)B_n$  $+(a_n - t_{n-1} - t_{n+1} - a_{n+1})B_nB_{n-1} + (b_{n-1} + b_n - 2\alpha b_{n-1})B_{n-1}$ +  $(a_n - a_{n+2} - t_{n+2} - t_{n+1})C_{n+1} + (2(1+\alpha)t_n + a_n - a_{n-2})C_n$  $-(t_{n-2}+t_{n-1})C_{n-1}+c_{n+1}-2\alpha c_n+c_{n-1}=(1-\alpha^2)(t_n+a_n),$ (2.18) $2(1-\alpha)a_nB_n^3 + 2(1-\alpha)b_nB_n^2 + [(2a_n - a_{n+2} - a_{n-1})C_{n+1}]$  $+(2a_n - a_{n+1} - a_{n-2})C_n + c_{n+1} - 2\alpha c_n + c_n - 2\alpha c_{n+1} - (1 - \alpha^2)a_n]B_n$  $+(c_{n+1}+a_{n+1}C_{n+1}-a_{n+2}C_{n+1})B_{n+1}+(c_n+a_{n-1}C_n-a_{n-2}C_n)B_{n-1}$  $+2(b_n - \alpha b_{n+1})C_{n+1} + 2(b_n - \alpha b_{n-1})C_n = (1 - \alpha^2)b_n.$ 

(2.9) (respectively, (2.10)) is obtained by shifting n to n+1 in (2.15) (respectively, (2.17)) and combining it with (2.14) (respectively, (2.16)) and by using (2.8). (2.11) follows from (2.8) and (2.18). Now suppose that  $deg\pi = 2$ . Using (2.1), we may write

$$P_n(X) = X^n - X^{n-1} \sum_{j=0}^{n-1} B_j + w_n X^{n-2} + \cdots,$$

for some complex sequence  $(w_n)_{n\geq 0}$ . Using [3, (2.40)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), we compare the two first coefficients of higher power of n in both side of (1.3) to deduce (2.13). (2.12) is obtained in a similar way and this completes the proof.

For the next results of this section, we distinguish two cases according as deg  $\pi = 1$  or deg  $\pi = 2$ .

2.1. Case deg  $\pi = 1$ . In this case, (1.3) can be rewritten as

(2.19) 
$$(X-r)\mathcal{D}_q P_n(X) = b_n P_n(X) + c_n P_{n-1}(X), \quad n = 0, 1, 2, \dots,$$

where  $r \in \mathbb{C}$ .

**Lemma 2.3.** Let  $(P_n)_{n>0}$  be a monic OPS satisfying (2.19). Then

(2.20) 
$$\left(c_2C_1 - q^{-1/2}c_1C_2\right)\left(c_2C_1 - q^{1/2}c_1C_2\right) = 0.$$

*Proof.* Since  $(P_n)_{n\geq 0}$  satisfies (2.19), then  $a_n = 0$  for each n = 0, 1, 2, ... and by (2.8) and (2.12), we have

(2.21) 
$$t_n = k_1 q^{n/2} + k_2 q^{-n/2}, \ k_1 = \frac{c_2 C_1 - q^{-1/2} c_1 C_2}{(q-1)C_1 C_2}, \ k_2 = \frac{c_2 C_1 - q^{1/2} c_1 C_2}{(q^{-1} - 1)C_1 C_2}.$$

Suppose, contrary to our claim, that (2.20) does not hold. This means that  $k_1k_2 \neq 0$ . Taking successively n = 1 and n = 2 in (2.19), and using (2.1), [3, (2.40)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), we have  $b_1 = 1$ ,  $b_2 = 2\alpha$ ,  $r = B_0 - c_1$ ,  $c_2 = (2\alpha - 1)(B_1 + B_0) - 2\alpha r$  and  $r(B_1 + B_0) = -c_2B_0 + 2\alpha(B_0B_1 - C_1)$ . Hence

(2.22) 
$$c_1 = B_0 - r, \quad B_1 = (2\alpha - 1)B_0 + 2\alpha \frac{C_1}{c_1}.$$

We claim that

(2.23) 
$$B_n = \frac{t_0 t_1 B_0}{t_n t_{n+1}}, \quad n = 0, 1, 2, \dots,$$

with  $B_0 \neq 0$ . Indeed, writing (2.9) as  $t_{n+3}B_{n+2} - t_{n+1}B_{n+1} = t_{n+2}B_{n+1} - t_nB_n$ and proceeding in a recurrent way, we have

(2.24) 
$$B_n = \frac{t_0 t_1 B_0 q^{n/2} + K_b \left(k_1 q^n + (k_2 - k_1 q^{-1/2}) q^{n/2} - k_2 q^{-1/2}\right)}{(k_1 q^n + k_2)(k_1 q^{n+1} + k_2)} q^{(n+1)/2},$$

where  $K_b = (t_2B_1 - t_0B_0)/(1 - q^{-1/2})$ . Since  $k_2 \neq 0$  and 0 < q < 1, we obtain  $\lim_{n\to\infty} q^{-n/2}B_n = -K_b/k_2$  and consequently we have  $K_b = 0$  by applying limit of the same expression using [3, (4.4)] ( for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ). This implies that  $B_1 = t_0B_0/t_2$ . If  $B_0 = 0$ , then we find  $B_1 = 0$  which is in contradiction with the second equation in (2.22). Then (2.23) is proved. Note that, from (2.3) and using (2.21), we obtain

(2.25) 
$$\mathfrak{a} = \frac{c_2 C_1}{c_1 C_2} - \alpha = \frac{t_2}{t_1} - \alpha = \frac{k_1 q^{1/2} - k_2 q^{-1/2}}{2u t_1}, \ u^{-1} = q^{1/2} - q^{-1/2},$$

since  $a_n = 0$ . Using (2.23), we obtain  $S_n = \sum_{j=0}^{n-1} B_j = t_1 B_0 \gamma_n / t_n$  for  $n = 0, 1, 2, \ldots$ . Thus using [3, (4.4)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) to evaluate the same sum, we have  $t_1 B_0(\mathfrak{a}\gamma_{2n-2} + \alpha_{2n-2}) = -t_n (\phi'(0)\gamma_{n-1} - B_0\alpha_{n-1})$ . This gives the following equations:

$$(2\mathfrak{a}ut_1 + k_2q^{-1/2})B_0 = -2uk_1q^{1/2}\phi'(0) \text{ and } (2\mathfrak{a}ut_1 - k_1q^{1/2})B_0 = -2uk_2q^{-1/2}\phi'(0).$$

Taking into account that  $k_1k_2 \neq 0$  and using (2.25), we get

$$|B_0 + 2u\phi'(0)| + |B_0 - 2u\phi'(0)| = 0,$$

which is impossible because  $B_0 \neq 0$ , and the lemma follows.

2.2. Case  $deg \pi = 2$ . In this case, we rewrite (1.3) as

(2.26)

$$(X - r)(X - s)\mathcal{D}_q P_n(X) = (a_n X + b_n)P_n(X) + c_n P_{n-1}(X), \ n = 0, 1, 2, \dots,$$

where  $r, s \in \mathbb{C}$  and  $c_n \neq 0$  for  $n = 1, 2, 3, \ldots$  From (2.13), (2.8) and (2.9), we obtain (2.27)

$$a_n = \gamma_n, \quad b_n = (\gamma_n - \gamma_{n-1}) \sum_{k=0}^{n-1} B_k - (r+s)\gamma_n$$

(2.28)

$$t_n = \frac{c_n}{C_n} = k_1 q^{n/2} + k_2 q^{-n/2}, \quad k_1 = \frac{c_2 C_1 - q^{-1/2} c_1 C_2}{(q-1)C_1 C_2}, \quad k_2 = \frac{c_2 C_1 - q^{1/2} c_1 C_2}{(q^{-1} - 1)C_1 C_2}$$
(2.29)  

$$r_n = \hat{a} q^{n/2} + \hat{b} q^{-n/2}, \quad \hat{a} = k_1 + u(1 - q^{-1/2}), \quad \hat{b} = k_2 - u(1 - q^{1/2}),$$

for  $n = 0, 1, 2, ..., u^{-1} = q^{1/2} - q^{-1/2}$ . Recall that  $t_0 = k_1 + k_2$  and so, we also define by compatibility  $r_0 = \hat{a} + \hat{b}$ .

**Lemma 2.4.** Let  $(P_n)_{n>0}$  be a monic OPS satisfying (2.26). Then

(2.30) 
$$\widehat{a}\ \widehat{b}(1-2\mathfrak{a} u)(1+2\mathfrak{a} u) \neq 0,$$

where  $\hat{a}$ ,  $\hat{b}$  are defined in (2.29) and  $\mathfrak{a}$  is given in (2.3).

*Proof.* Assume that (2.30) does not hold. Suppose, for instance, that  $\hat{a} = 0$ . Then (2.29) reduces to  $r_n = \hat{b}q^{-n/2}$  for each n = 0, 1, 2, ... Then (2.9) becomes

$$q^{-3/2}B_{n+2} - (q^{-1} + q^{-1/2})B_{n+1} + B_n = 0, \quad n = 0, 1, 2, \dots,$$

and so we may write

(2.31) 
$$B_n = r_0(1-q^{1/2})q^{n/2} + s_0(1-q)q^n, \quad n = 0, 1, 2, \dots,$$

for some complex numbers  $r_0$  and  $s_0$ . From (2.3), we also have

$$\mathfrak{a} = \frac{(c_2 + 2\alpha C_2)C_1}{(c_1 + C_1)C_2} - \alpha = \frac{1 + r_2}{r_1} - \alpha = -\frac{1}{2u} + \frac{1}{\widehat{b}q^{-1/2}}.$$

From (2.31), we get  $S_n = \sum_{j=0}^{n-1} B_j = r_0(1-q^{n/2}) + s_0(1-q^n)$  for  $n = 0, 1, 2, \dots$ Now we compute this sum using [3, (4.4)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) to obtain

(2.32) 
$$(r_0q^{n/2} + s_0q^n - r_0 - s_0)d_{2n} = \gamma_{n+1}e_n, \quad n = 1, 2, 3, \dots$$

where  $2d_{2n} = (1 + 2\mathfrak{a}u)q^n + (1 - 2\mathfrak{a}u)q^{-n}$  and  $2e_n = (-B_0 + 2u\phi'(0))q^{n/2} - (B_0 + 2u\phi'(0))q^{-n/2}$  for n = 0, 1, 2, ... It is easily seen that (2.32) implies  $r_0 = 0 = s_0$  as well as  $B_0 = 0 = \phi'(0)$ . Hence  $B_n = 0$  for n = 0, 1, 2, ... In addition, using (2.2), we obtain  $\mathfrak{b} = 0$ . Next we apply [3, (4.5)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) to obtain

(2.33) 
$$C_{n+1} = \frac{(1-q^{n+1})(B-q^n)\left(q^{2n+1} + (4(q+\hat{b})C_1 - q - B)q^n + B\right)}{4(B-q^{2n})(B-q^{2n+2})},$$

for n = 0, 1, 2, ..., with  $B = q + \hat{b}(1 - q)$ , while (2.10) reduces to

$$(q^{-1/2} + q^{-1})(C_{n+1} - 1/4) - 2(1 + \alpha)(C_n - 1/4) + (q^{1/2} + q)(C_{n-1} - 1/4) = 0.$$

Therefore, we may write  $C_{n+1} = \overline{r}_0 q^{n/2} + \overline{s}_0 q^n + 1/4$  for  $n = 2, 3, \ldots$ , with  $\overline{r}_0, \overline{s}_0 \in \mathbb{C}$ . This is compatible with (2.33) if and only if  $C_1 = 1/2$ ,  $\overline{r}_0 = \overline{s}_0 = 0$  and  $\hat{b} = 1$ . This implies that

$$B_{n-1} = 0, \quad C_1 = 1/2, \quad C_{n+1} = 1/4, \quad n = 1, 2, \dots$$

From (3.11)-(3.14) below we find r = -s and so

(2.34) 
$$c_1 = -r^2, \quad c_2 = \alpha(1 - 2r^2)$$

Since  $\hat{a} = 0$  and  $\hat{b} = 1$ , from (2.29) we find  $k_1 = -k_2 = -(1 + q^{1/2})^{-1}$  and so  $t_n = -(1 - q^{-1/2})\gamma_n$ . On the other hand we have

$$\mathfrak{a} = \frac{(c_2 + 2\alpha C_2)C_1}{(c_1 + C_1)C_2} - \alpha = \frac{2\alpha + t_2}{1 + t_1} - \alpha = \alpha.$$

Using (2.34), we also write  $\alpha = \mathfrak{a} = \alpha(5 - 6r^2)/(1 - 2r^2)$ , hence  $r^2 = 1$  so that  $c_1 = -1$  and  $c_2 = -\alpha$ . But using what is preceding, from (2.28) we have  $k_1 = -k_2 = -2u$  and therefore  $t_n = -2\gamma_n$ , which is in contradiction with the previous expression of  $t_n$ , which gives  $\hat{a} \neq 0$ . The case  $\hat{b} = 0$  can be treated similarly.

Assume now that  $1 + 2\mathfrak{a}u = 0$ . Since  $\mathfrak{a} = -\alpha + (1 + r_2)/r_1$ , we obtain  $\hat{a} = -uq^{-1/2} \neq 0$ . On the other hand, we use [3, (4.4)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) to obtain

$$B_n = q^n (1+q^{-1}) \Big( \mathfrak{b}u(q^n-1) + \frac{B_0}{1+q^{-1}} \Big),$$

for n = 0, 1, 2, ... This satisfies (2.9) if and only if  $\mathfrak{b} = 0$  and  $B_0 = 0$ , and so  $B_n = 0$ . Taking this into account, [3, (4.5)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) gives

$$C_{n+1} = \frac{1}{4}(1 - q^{n+1})\left(1 - q^n + \frac{4C_1}{1 - q}q^n\right), \quad n = 0, 1, 2, \dots$$

This does not satisfy (2.10) because  $\hat{a} \neq 0$  and  $B_n = 0$ . Hence  $1 + 2\mathfrak{a}u \neq 0$ . The case  $1 - 2\mathfrak{a}u = 0$  can be treated similarly, which proves the lemma.

# 3. Proof of theorems 1.2 and 1.3

Proof of Theorem 1.2. Note that (2.20) is equivalent to  $k_1k_2 = 0$ . Suppose that  $k_1 = 0$ . By (2.21), we have  $t_n = k_2q^{-n/2}$ , where  $k_2 = q^{1/2}c_1/C_1$ . We claim that

(3.1) 
$$B_n = B_0 q^n = (r + c_1) q^n, \quad n = 0, 1, 2, \dots$$

Indeed, (2.9) reduces to  $q^{-1/2}B_{n+2} + (1+q^{1/2})B_{n+1} + qB_n = 0$ , n = 0, 1, 2, ...and so we find  $B_n = vq^n + sq^{n/2}$  for some  $v, s \in \mathbb{C}$ . Moreover, since  $k_1 = 0$ , from (2.25) we get  $\mathfrak{a} = -1/(2u)$ . Hence, by (2.2),  $\phi(X) = -((X + 2\mathfrak{b}u)(X - B_0) + 2uq^{-1/2}C_1)/(2u)$  and  $\psi(X) = X - B_0$ . Therefore, using [3, (4.4)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), we obtain  $B_n = q^{(2n-1)/2}(2\alpha u\mathfrak{b}(q^n - 1) + q^{1/2}B_0)$ . Comparing the two previous expressions for  $B_n$ , we find  $s = 0 = \mathfrak{b}$  and  $v = B_0$ . Hence using the first equation in (2.22), (3.1) follows. As consequence, taking n = 1 in (3.1) and comparing the result with the expression for  $B_1$  given by (2.22), we obtain

(3.2) 
$$C_1 = (q^{1/2} - 1)(r + c_1)c_1.$$

Since  $C_n = c_n/t_n$ , from (2.12) and (3.1), we find

(3.3) 
$$C_{n+1} = \frac{C_1}{(q-1)c_1} \left(1 - q^{n+1}\right) \left(r - \frac{r+c_1}{1+q^{1/2}} (1+q^{(2n+1)/2})\right).$$

Taking into account that  $\mathfrak{a} = -1/(2u)$  and  $\mathfrak{b} = 0$ , using [3, (4.5)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), we also have

(3.4) 
$$C_{n+1} = (1 - q^{n+1}) \left( \frac{1}{4} (1 - q^n) + \frac{C_1}{1 - q} q^n \right), \quad n = 0, 1, 2, \dots$$

If  $c_1 = rq^{1/2}$  then (3.3) becomes  $C_{n+1} = C_1(1-q^{n+1})q^n/(1-q)$  which is incompatible with (3.4). Thus  $c_1 \neq rq^{1/2}$ . Comparing the expressions for  $C_{n+1}$  given in (3.3) and (3.4) yields

(3.5) 
$$C_1 = (1-q)\frac{(1+q^{1/2})c_1}{4(c_1-q^{1/2}r)}.$$

Therefore, combining (3.5) with (3.2), we see that  $r+c_1$  is a solution of the following quadratic equation

(3.6) 
$$2Y^2 - 2(1+q^{-1/2})c_1Y - 1 - \alpha = 0.$$

Let c and d be two complex numbers defined by

$$(c,d) \text{ or } (d,c) \in \left\{ \left( c_1 - \sqrt{\Delta}, q^{-1/2}(c_1 - \sqrt{\Delta}) \right), \left( c_1 + \sqrt{\Delta}, q^{-1/2}(c_1 + \sqrt{\Delta}) \right) \right\},$$

where  $\Delta = c_1^2 + q^{1/2}$ . Note that  $cd \neq 0$ . Set  $Y_1 = (c+d)/2$  and  $Y_2 = -(c^{-1}+d^{-1})/2$ . Hence  $Y_1$  and  $Y_2$  are solutions of (3.6). Without loss of generality we may set  $r + c_1 = Y_1$  and so  $Y_1 + Y_2 = (1 + q^{-1/2})c_1$ , which yields

$$r = (c+d)\frac{1+cdq^{-1/2}}{2cd(1+q^{-1/2})}, \quad c_1 = (c+d)\frac{cd-1}{2cd(1+q^{-1/2})}$$

Hence (3.5) (or (3.2)) becomes  $C_1 = (1 - q)(1 - cd)/4$ . Consequently, from (3.3) (or (3.4)) and (3.1), we obtain

(3.7) 
$$B_n = (c+d)q^n/2, \quad C_{n+1} = (1-q^{n+1})(1-cdq^n)/4,$$

together with  $k_2 = q^{1/2}c_1/C_1 = 2u(c+d)/(cd(1+q^{-1/2}))$ . Using (3.7), equation (2.10) now reads as

$$(q^{-1} + q^{-1/2})(C_{n+1} - 1/4) - 2(1 + \alpha)(C_n - 1/4) + (q + q^{1/2})(C_{n-1} - 1/4)$$
  
(3.8)  
$$= (\alpha - 1)(\alpha + 1/2)(c + d)^2 q^{2n-1}.$$

From  $c^2 + d^2 = 2\alpha cd$ , it is easy to see that  $B_n$  and  $C_{n+1}$ , in (3.7), satisfy (3.8). (2.11) in this case  $(a_n = 0 \text{ for } n = 0, 1, 2, ...)$  reads as

$$(3.9) \qquad 2(1-\alpha)b_nB_n^2 + (1-2\alpha)(c_n+c_{n+1})B_n + c_{n+1}B_{n+1} + c_nB_{n-1} + (b_n-b_{n+2})(C_{n+1}-1/4) + (b_n-b_{n-2})(C_n-1/4) = 0,$$

where  $c_n = t_n C_n = k_2 q^{-n/2} C_n$  for  $n = 1, 2, \ldots$  Similarly, one may check that (3.9) is also satisfied and, therefore, the system of equations (2.8)-(2.11) is fulfilled. By a similar argument, if  $k_2 = 0$ , we obtain (3.7) with q replaced by 1/q and  $c^2 + d^2 - 2\alpha cd = 0$  as solution of the system of difference equations (2.8)-(2.11). Thus

(3.10) 
$$P_n(X) = Q_n(X; c, d|q)$$
 or  $P_n(X) = Q_n(X; c, d|q^{-1}), \quad n = 0, 1, 2, \dots,$ 

with  $c^2 + d^2 - 2\alpha cd = 0$ , i.e.  $c/d = q^{\pm 1/2}$ , and this is precisely the assertion of Theorem 1.2.

Proof of Theorem 1.3. Taking successively n = 1 and n = 2 in (2.26) using (2.1) and [3, (2.40)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) we obtain the following:

$$(3.11) B_0 = b_1 + r + s, c_1 = (B_0 - r)(B_0 - s),$$

(3.12) 
$$b_2 = (2\alpha - 1)(B_0 + B_1) - 2\alpha(r+s),$$

$$(3.13) rs(B_0 + B_1) = c_2 B_0 - b_2 (B_0 B_1 - C_1),$$

$$(3.14) c_2 = b_2(B_0 + B_1) - 2\alpha(B_0B_1 - C_1) + (r+s)(B_0 + B_1) + 2\alpha rs.$$

Solving (2.9) we find

(3.15) 
$$B_n = \frac{r_0 r_1 B_0 q^{n/2} + \widehat{K}_b \left( \widehat{a} q^n + (\widehat{b} - \widehat{a} q^{-1/2}) q^{n/2} - \widehat{b} q^{-1/2} \right)}{(\widehat{a} q^n + \widehat{b}) (\widehat{a} q^{n+1} + \widehat{b})} q^{(n+1)/2} ,$$

for n = 0, 1, 2, ..., where  $\widehat{K}_b = (r_2 B_1 - r_0 B_0)/(1 - q^{-1/2})$ . Since  $\widehat{a} \ \widehat{b} \neq 0$  and 0 < q < 1, then  $\lim_{n\to\infty} q^{-n/2} B_n = -\widehat{K}_b/\widehat{b}$ . Evaluating this limit using [3, (4.4)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), we find  $\widehat{K}_b = 0$ , because  $1 - 2\mathfrak{a}u \neq 0$ . Hence (3.15) reduces to

(3.16) 
$$B_n = \frac{r_0 r_1 B_0}{r_n r_{n+1}}, \quad n = 0, 1, 2, \dots$$

It is immediate that  $S_n = \sum_{j=0}^{n-1} B_j = r_1 B_0 a_n / r_n$  for n = 0, 1, 2, ... Comparing this with the result obtained using [3, (4.4)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) we have (3.17)  $(2\mathfrak{a} u r_1 q^{-1/2} + \widehat{b} q^{-1}) B_0 = -2\widehat{a} u \phi'(0)$ ,  $(2\mathfrak{a} u r_1 q^{1/2} - \widehat{a} q) B_0 = -2\widehat{b} u \phi'(0)$ .

CASE 1. Suppose 
$$\phi'(0) = 0$$
. From (3.17) we obtain  $r_1B_0 = 0$ . But from (2.4) we obtain  $0 \neq c_1 + a_1C_1 = r_1C_1$ , i.e.  $r_1 \neq 0$ , and so  $B_0 = 0$ . (Conversely, if we assume  $R_1 = 0$  by (2.20) we obtain  $\phi'(0) = 0$ .) Hence  $R_2 = 0$  and  $h = 0$ . From

assume  $B_0 = 0$ , by (2.30), we obtain  $\phi'(0) = 0$ .) Hence  $B_0 = 0$  and  $\mathfrak{b} = 0$ . From this, we use [3, (4.4)-(4.5)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ) to obtain  $B_n = 0$  and

(3.18) 
$$C_{n+1} = \frac{(1-q^{n+1})(1-hq^{n-1})(1+wq^n+hq^{2n})}{4(1-hq^{2n-1})(1-hq^{2n+1})}$$

with  $h = -(1+2\mathfrak{a}u)/(1-2\mathfrak{a}u)$  and  $w = 4u(2(\mathfrak{a}+\alpha)C_1-\mathfrak{a})/(2\mathfrak{a}u-1)$ . Now defining  $\widehat{C}_n = C_n - 1/4$ , (2.10) reads as

$$(3.19) (r_{n+1}+r_{n+2})\widehat{C}_{n+1}-2(1+\alpha)r_n\widehat{C}_n+(r_{n-1}+r_{n-2})\widehat{C}_{n-1}=0.$$

Therefore, we may write

$$(3.20) C_{n+1} = \frac{1}{4} + \frac{\widehat{\theta}_0 \widehat{\theta}_1 \widehat{C}_1 q^{n/2} + \widehat{K}_c \left(\widehat{r}_0 q^n + (\widehat{r}_1 - \widehat{r}_0 q^{-1/2}) q^{n/2} - \widehat{r}_1 q^{-1/2}\right)}{(\widehat{r}_0 q^n + \widehat{r}_1)(\widehat{r}_0 q^{n+1} + \widehat{r}_1) q^{-(n+1)/2}}$$

for n = 2, 3, ..., for some complex numbers  $\hat{\theta}_0$ ,  $\hat{\theta}_1$  and  $\hat{K}_c$ , where  $\hat{\theta}_n = r_n + r_{n+1} = \hat{r}_0 q^{n/2} + \hat{r}_1 q^{-n/2}$ . Taking into account (2.30) one may see that (3.18) and (3.20) are compatible if and only if either

(3.21) 
$$C_1 = 1/4, \quad \widehat{K}_c = 0, \quad \mathfrak{a} = \alpha (4\alpha^2 - 3)/(4\alpha^2 - 1),$$

 $\mathbf{or}$ 

(3.22) 
$$C_1 = 1/2, \quad \mathfrak{a} = \alpha, \quad \widehat{K}_c = 0, \quad \widehat{\theta}_0 \widehat{\theta}_1 = 0.$$

In case of (3.21) holds, we get

$$B_n = 0, \quad C_{n+1} = 1/4, \quad n = 0, 1, \dots$$

This satisfies (2.8)-(2.11), taking into account that from (2.13) we have  $a_n = \gamma_n$ and  $b_n = -(r+s)\gamma_n$  for  $n = 0, 1, \ldots$  From (3.11)-(3.14), we obtain r+s = 0,  $c_1 = -r^2$  and  $c_2 = -\alpha(2r^2 - 1/2)$ . In addition,  $r^2 = \alpha^2$ . In fact, this follows by comparing the expression of  $\mathfrak{a}$  given in (3.21) and the one obtained from (2.3). So we now have  $c_1 = -\alpha^2$  and  $c_2 = -\alpha(2\alpha^2 - 1/2)$ . Next, with these expressions,  $k_1$ and  $k_2$  given in (2.28) become  $k_1 = -u(1+q)$  and  $k_2 = u(1+q^{-1})$  and consequently  $c_n = -\alpha\gamma_{n+1}/2$  for  $n = 0, 1, \ldots$  However taking n = 3 in (2.26) and using the fact that  $B_n = 0$  and r = -s, we obtain

(3.23)  

$$2(2\alpha^2 - 1)(C_1 + C_2) = \alpha^2 - 1 + c_3 + (4\alpha^2 - 1)r^2, \ c_3C_1 = (1 - \alpha^2 - C_1 - C_2)r^2.$$

It is then clear that the obtained values of  $C_1$ ,  $C_2$ ,  $c_3$  and  $r^2$  do not satisfy (3.23). For the case where conditions (3.22) meet, we obtain

$$B_{n-1} = 0, \ C_1 = 1/2, \ C_{n+1} = 1/4, \quad n = 1, 2, \dots$$

Again this satisfies (2.8)-(2.11) taking into account that from (2.13) we have  $a_n = \gamma_n$ and  $b_n = -(r+s)\gamma_n$  for  $n = 0, 1, \ldots$  From (3.11)-(3.14), we obtain r+s = 0,  $c_1 = -r^2$  and  $c_2 = \alpha(1-2r^2)$ . In addition, from the expression of  $\mathfrak{a}$  giving in (3.22), we write

$$\alpha = \mathfrak{a} = \frac{(c_2 + 2\alpha C_2)C_1}{(c_1 + C_1)C_2} - \alpha = \alpha \frac{5 - 6r^2}{1 - 2r^2},$$

Therefore  $r^2 = 1$ , and we have  $c_1 = -1$  and  $c_2 = -\alpha$ . Hence  $k_1$  and  $k_2$  given in (2.28) become  $k_1 = -k_2 = -2u$  and, consequently,  $t_n = -2\gamma_n$  so that  $c_1 = -1$  and  $c_n = -\gamma_n/2$  for  $n = 1, 2, \ldots$  We check at once that  $C_1$ ,  $C_2$ ,  $c_3$  and  $r^2$  satisfy (3.23). Hence  $P_n(X) = \hat{T}_n(X)$  for  $n = 0, 1, 2, \ldots$ , where  $(\hat{T}_n)_{n\geq 0}$  is the monic Chebyschev polynomials of the first kind, and this is precisely the first assertion of Theorem 1.3.

CASE 2. Suppose  $\phi'(0) \neq 0$ . (3.17), taking into account (2.30), implies  $B_0 \neq 0$ . The converse is also true. Hence

$$r_1 B_0(\widehat{a}q^{1/2} - \widehat{b}q^{-1/2}) \neq 0.$$

Solving (3.17), we get

(3.24) 
$$\mathfrak{a} = -\frac{1 + (q\widehat{a}/\widehat{b})^2}{2u\left(1 - (q\widehat{a}/\widehat{b})^2\right)}, \quad B_0 = 2u\phi'(0)\frac{1 - (q\widehat{a}/\widehat{b})}{1 + (q\widehat{a}/\widehat{b})}$$

Considering  $\hat{a}$ ,  $\hat{b}$ ,  $B_0$  and  $C_1$  as free parameters, let us define, without loss of generality, two complex numbers a and b such that  $-q^{a/2}$  and  $q^{b/2}$  are solutions of the following quadratic equation

(3.25) 
$$Y^2 + \frac{2r_1 B_0 q^{1/4}}{\widehat{b}(1+q^{1/2})} Y + \frac{\widehat{a}}{\widehat{b}} = 0.$$

Thus

(3.26) 
$$q^{a/2} - q^{b/2} = \frac{2r_1 B_0 q^{1/4}}{\widehat{b}(1+q^{1/2})}, \quad q^{(a+b)/2} = -\widehat{a}/\widehat{b}$$

On the other hand, we have

$$(3.27) \qquad (q^{a/2}, q^{b/2}) \in \left\{ \left( \frac{\widehat{a}(1+q^{1/2})}{r_1 B_0 q^{1/4} + \sqrt{\Delta}}, -\frac{r_1 B_0 q^{1/4} + \sqrt{\Delta}}{\widehat{b}(1+q^{1/2})} \right), \\ \left( \frac{\widehat{a}(1+q^{1/2})}{r_1 B_0 q^{1/4} - \sqrt{\Delta}}, -\frac{r_1 B_0 q^{1/4} - \sqrt{\Delta}}{\widehat{b}(1+q^{1/2})} \right) \right\},$$

where  $\Delta = q^{1/2}B_0^2 r_1^2 - \hat{a}\hat{b}(1+q^{1/2})^2$ . From (3.26), (3.24) and (2.2), we obtain

$$\begin{split} \mathfrak{a} &= -\frac{1+q^{a+b+2}}{2u(1-q^{a+b+2})}, \ B_0 = \frac{(1+q^{1/2})q^{1/4}(q^{a/2}-q^{b/2})}{2(1-q^{(a+b+2)/2})}, \\ \mathfrak{b} &= \frac{q^{3/4}(q^{a/2}-q^{b/2})q^{(a+b+2)/2}}{2u^2(q^{1/2}-1)(1-q^{a+b+2})(1-q^{(a+b+2)/2})}. \end{split}$$

Note that (2.29) can be written using (3.26) as

(3.28) 
$$r_n = \hat{b}(1 - q^{n+(a+b)/2})q^{-n/2}, \quad n = 0, 1, 2, \dots$$

Therefore (3.16) becomes

(3.29) 
$$B_n = q^{1/4} (1+q^{1/2}) \frac{(1-q^{(a+b)/2})(q^{a/2}-q^{b/2})q^n}{2(1-q^{(2n+a+b)/2})(1-q^{(2n+a+b+2)/2})} \\ = \left(q^{(2a+1)/4} + q^{-(2a+1)/4} - y_n - z_n\right)/2.$$

Taking into account the above, (2.2) becomes

$$\begin{split} \phi(X) &= -\frac{1+q^{a+b+2}}{2u(1-q^{a+b+2})} X^2 + q^{1/4}(1+q^{1/2}) \frac{(q^{a/2}-q^{b/2})}{4u(1+q^{(a+b+2)/2})} X \\ &+ \frac{(1+\alpha)q^{(a+b+4)/2}(q^{a/2}-q^{b/2})^2}{2u(1-q^{a+b+2})(1-q^{(a+b+2)/2})^2} - q^{-1/2} \frac{1-q^{a+b+3}}{1-q^{a+b+2}} C_1, \\ \psi(X) &= X - q^{1/4}(1+q^{1/2}) \frac{(q^{a/2}-q^{b/2})}{2(1-q^{(a+b+2)/2})}. \end{split}$$

Let **u** be the regular linear functional with respect to the monic OPS  $(P_n)_{n\geq 0}$ . By [3, (4.2)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), the regularity conditions for **u** are

$$(1 - q^{n+a+1})(1 - q^{n+b+1})(1 - q^{n+a+b+1})C_1 \neq 0, \quad n = 0, 1, 2, \dots$$

Moreover, by [3, (4.4)-(4.5)] (for  $\mathfrak{c}_3 = 0$  and  $\mathfrak{c}_1 = \mathfrak{c}_2 = 1/2$ ), we obtain the same expression for  $B_n$  and

$$C_{n+1} = \frac{K(1-q^{n+1})(1-q^{n+a+b+1})(1-q^{n+a+1})(1-q^{n+b+1})}{(1-q^{(2n+a+b+1)/2})(1-q^{(2n+a+b+2)/2})^2(1-q^{(2n+a+b+3)/2})},$$

where

$$K = -\frac{q^{-1/2}uC_1(1-q^{(a+b+3)/2})(1-q^{(a+b+2)/2})^2}{(1-q^{a+1})(1-q^{b+1})(1+q^{(a+b+1)/2})}.$$

After tedious calculations, we can see that this expression for  $C_{n+1}$  and (3.29) satisfy (2.10) if and only if

$$C_1 = \frac{(1-q)(1-q^{a+1})(1-q^{b+1})(1+q^{(a+b+1)/2})}{4(1-q^{(a+b+3)/2})(1-q^{(a+b+2)/2})^2}.$$

Alternatively,  $C_1$  follows by taking n = 2 in (2.10), using  $C_2$  and  $C_3$ . Note that (2.10) holds for each  $n = 2, 3, \ldots$ . Consequently, we obtain (3.30)

$$C_{n+1} = \frac{(1-q^{n+1})(1-q^{n+a+b+1})(1-q^{n+a+1})(1-q^{n+b+1})}{4(1-q^{(2n+a+b+1)/2})(1-q^{(2n+a+b+2)/2})^2(1-q^{(2n+a+b+3)/2})} = y_n z_{n+1}/4.$$

Taking into account that  $c_n = t_n C_n$ , (3.11)-(3.14) yields

$$\begin{aligned} \widehat{b}q^{-1/2}(1-q^{(a+b+2)/2})C_1 + (r+s)B_0 - rs &= B_0^2 + C_1, \\ \widehat{b}q^{-1}(1-q^{(a+b+4)/2})B_0C_2 - rs(B_0 + B_1) + 2\alpha(B_0B_1 - C_1)(r+s) \\ &= (2\alpha - 1)\Big((B_0B_1 - C_1)(B_0 + B_1) + B_0C_2\Big), \\ \widehat{b}q^{-1}(1-q^{(a+b+4)/2})C_2 - 2\alpha rs + (2\alpha - 1)(r+s - B_0 - B_1)(B_0 + B_1) \\ &= 2\alpha(C_1 - B_0B_1) + (2\alpha - 1)C_2. \end{aligned}$$

Solving firstly the above system for  $\hat{b}$ , r+s and rs, and secondly the obtained result for r and s, we get  $\hat{b} = uq^{1/2}(1 + q^{-(a+b+2)/2})$  and

$$(r,s)$$
 or  $(s,r) \in \left\{ \left( (q^{(2a+1)/4} + q^{-(2a+1)/4})/2, -(q^{(2b+1)/4} + q^{-(2b+1)/4}) \right)/2 \right\}$ .

So (3.28) becomes  $r_n = u(1 + q^{-(a+b+2)/2})(1 - q^{n+(a+b)/2})q^{(1-n)/2}$  and  $t_n = u(1 + q^{-(a+b+1)/2})(1 - q^{n+(a+b+1)/2})q^{-n/2}$ . Using (3.29), (2.27) becomes

(3.31) 
$$b_n = (q^{a/2} - q^{b/2})\gamma_n \frac{(1 + q^{n+a+b+1/2})}{2(1 - q^{n+(a+b)/2})} q^{-(2a+2b+1)/4},$$

for each  $n = 0, 1, 2, \dots$  Also, since  $c_n = t_n C_n$ , we obtain

(3.32) 
$$c_n = -\gamma_n \frac{(1-q^{n+a})(1-q^{n+b})(1-q^{n+a+b})(1+q^{-(a+b+1)/2})}{4(1-q^{n+(a+b-1)/2})(1-q^{n+(a+b)/2})^2}$$

for  $n = 0, 1, 2, \ldots$  Finally, again after tedious computations, from (3.29), (3.30), (3.31) and (3.32), we see that (2.11) holds. Hence (2.8)-(2.11) also hold. Note that (3.24) may also write as

$$\mathfrak{a} = \frac{1 + (q^{-1}\widehat{b}/\widehat{a})^2}{2u\left(1 - (q^{-1}\widehat{b}/\widehat{a})^2\right)}, \quad B_0 = -2u\phi'(0)\frac{1 - (q^{-1}\widehat{b}/\widehat{a})}{1 + (q^{-1}\widehat{b}/\widehat{a})}.$$

Proceeding similarly with the same parameters a and b as defined in (3.27), we obtain the same results with q replaced by  $q^{-1}$ . Thus

$$P_n(X) = \hat{P}_n^{(a,b)}(X|q) \text{ or } P_n(X) = \hat{P}_n^{(a,b)}(X|q^{-1}),$$

where  $\widehat{P}_n^{(a,b)}(\cdot|q)$  is the monic continuous q-Jacobi polynomial, and this is precisely the second assertion of Theorem 1.3.

#### Acknowledgments

The authors thank to Professor T. H. Koornwinder for helpful discussions and comments. This work is partially supported by the Centre for Mathematics of the University of Coimbra - UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES.

## References

- W. Al-Salam, A characterization of the Rogers q-Hermite polynomials, Internat. J. Math. & Math. Sci. 18 (1995), no. 4, 641-648
- [2] W. Al-Salam and T. S. Chihara, Another characterization of the classical orthogonal polynomials, SIAM J. Math. Anal. 3 (1972) 65-70.
- [3] K. Castillo, D. Mbouna, and J. Petronilho, On the functional equation for classical orthogonal polynomials on lattices, arXiv:2102.00033 [math.CA] (2021).
- [4] T. H. Koornwinder, The structure relation for Askey-Wilson polynomials, J. Comput. Appl. Math. 207 (2007), 214-226.
- [5] D. Mbouna, On some problems in the theory of orthogonal polynomials, Math. Dept. of the University of Coimbra, PhD Thesis (2021).
- [6] M. E. H. Ismail, Classical and quantum orthogonal polynomials in one variable. With two chapters by W. Van Assche. With a foreword by R. Askey., Encyclopedia of Mathematics and its Applications 98. Cambridge University Press, Cambridge, 2005.

CMUC, department of Mathematics, University of Coimbra, 3001-501 Coimbra, Portugal

 $Email \ address, \ K. \ Castillo: \ \texttt{kenier@mat.uc.pt}$ 

Email address, D. Mbouna: dmbouna@mat.uc.pt