

Central charge criticality of charged AdS black hole surrounded by different fluids

R. B. Alfaia,^{1,*} I. P. Lobo,^{2,1,†} and L. C. T. Brito^{1,‡}

¹*Physics Department, Federal University of Lavras, Caixa Postal 3037, 37200-000 Lavras-MG, Brazil*

²*Department of Chemistry and Physics, Federal University of Paraíba, Rodovia BR 079 - Km 12, 58397-000 Areia-PB, Brazil*

(Dated: September 22, 2021)

We analyze the extended phase space thermodynamics of Kiselev black hole introducing a central charge and allowing Newton's constant to vary. We also discuss the relation between the chemical potential and the size of the black hole, besides the new description of phase transitions. We obtain as a conclusion that the universality of the central charge does not remain valid in general.

I. INTRODUCTION

Since the fundamental works of Bekenstein [1, 2], Bardeen, Carter and Hawking [3, 4], black hole thermodynamics have played a prominent role in theoretical physics due to its property of bringing together gravity, quantum physics and thermodynamics into a unified framework. Since then, many developments have been achieved as the existence of phase transitions when different kinds of black hole elements are considered, like the presence of a cosmological constant [5], electric charge, and many other types of matter Lagrangians [6–12].

In particular, one of the most interesting aspects that emerged recently consists in the possibility of assuming a negative cosmological constant Λ as a thermodynamic variable, which plays the role of a pressure term. In this context, we now identify the mass parameter as an enthalpy instead of the internal energy, as was assumed previously [13, 14] (at constant pressure, these quantities coincide). In this case, intriguing features have been analyzed like the resemblance of the Reisser-Nordström-anti de Sitter spacetime with a van der Waals fluid as a thermodynamic system, a property that also allowed this approach to be named *black hole chemistry* [15]. We can see the productiveness of this research area by the features that have been extensively analyzed in the literature in the past years [16–20] as, for example, the existence and properties of phase transitions, classification of phases, conjectured geometric inequalities and the effect of different kinds of matter fields and quantum effects as conductors of transitions.

Due to the dependence of a negative cosmological constant, the relation between this approach and the holographic principle naturally emerges. Based on this, the system's behavior as a holographic heat engine [21] has also been studied in the literature considering the effect of different contributions on the efficiency of some cycles [22–26]. However, a satisfactory realization of this principle in black hole thermodynamics has been a matter of scrutiny [27]. Recently, advances in this direction have considered the assumption of Newton's constant G as a thermodynamic variable as a way how to present a Smarr formula that does not depend on the spacetime dimension nor on the bulk pressure [28, 29]. This approach relies on the introduction of a new thermodynamic quantity: the central charge C of the dual Conformal Field Theory.

The analysis presented in [29] contains interesting results that may open a new branch in black hole thermodynamics. From a fundamental point of view, some aspects which deserve attention are the conjectured role played by the central charge in phase transitions and the existence of a “universal” critical charge from which phase transitions can occur. Besides that, there are other interesting points to be highlighted: the realization of a holographic Smarr relation which leads to the first law of thermodynamics in the usual D -dimensional spacetime when G is constant; the verification of the validity of the reverse isoperimetric inequality for a class of black holes; the redefinition of some quantities like a new thermodynamic volume and the introduction of a “chemical potential”. In a few words, Ref.[29] is based on the thermodynamics of Kerr-Newman-anti de Sitter black hole with a variable G and the introduction of new thermodynamic variables that mix the cosmological constant, the spacetime dimension and Newton's constant in a new extended phase space. In this work, we investigate if those results remain valid when a different matter content is present and what is the impact of this matter content on the findings of the original paper. To achieve this objective, we will use the so-called Kiselev black hole in D -dimensions [30, 31] as our ansatz, and for simplicity reasons, we will assume the static configuration.

*Electronic address: alfaiaramon@gmail.com

†Electronic address: iarley_lobo@fisica.ufpb.br, lobofisica@gmail.com

‡Electronic address: lcbrito@ufla.br

Our paper is organized as follows. In section II, we verify the validity of holographic formulation of the thermodynamics of Kiselev (Reissner-Nordström-anti de Sitter) black hole. We consider a variable Newton's constant and the introduction of the central charge as a thermodynamic variable. In section III, we study the 4-dimensional case in this new extended phase space structure and verify the relation between the size of the black hole and its chemical potential. In section IV, we study its critical behavior, the conditions for the existence of the universal nature of the critical central charge. Then, in section V, we draw our final remarks. We assume natural units: $\hbar = c = k_B = 1$.

II. KISELEV BLACK HOLE IN ARBITRARY DIMENSIONS

In this section, we consider the so-called Kiselev black hole [30], that has been extensively analyzed in the literature, that aims to incorporate the effects of exotic matter that, in a cosmological setting, would be responsible for describing an accelerated expansion of the universe. We assume the D -dimensional spherically symmetric spacetime ansatz

$$ds^2 = f(r)dt^2 - g(r)dr^2 - r^2 d\theta_1^2 - r^2 \sin^2 \theta_1 d\theta_2^2 - \dots - r^2 \sin^2 \theta_1 \dots \sin^2 \theta_{D-3} d\theta_{D-2}^2, \quad (1)$$

and that the black hole is surrounded by matter with an average stress energy tensor of the form [31]

$$T_t^t = T_r^r = \rho(r), \quad (2)$$

$$T_{\theta_1}^{\theta_1} = T_{\theta_2}^{\theta_2} = \dots = T_{\theta_{D-2}}^{\theta_{D-2}} = -\frac{\rho(r)}{D-2}[(D-1)\omega + 1], \quad (3)$$

where ρ is interpreted as an average energy density of this fluid and ω is an average equation of state parameter, that labels different fluids.¹ For instance, in four dimensions, the so called average quintessence matter ($-1 < \omega < -1/3$) [35] is described by $\omega = -2/3$, $\omega = -1$ describes an average cosmological constant and $\omega = -4/3$ describes an average phantom field ($\omega < -1$) [36].

The solution of these field equations obeys additivity and linearity properties. It means that we can add fluid contributions to the field equations by adding related extra terms to the metric. In fact, this way, the solution of the Einstein field equations that endows the black hole mass, electric charge, cosmological constant and Kiselev's fluid has been found in [31], and its form that reduces to the Reissner-Nordström-Tangherlini metric [37–41] reads

$$f(r) = g(r)^{-1} = 1 - \frac{16\pi GM}{(D-2)\Omega_{D-2}r^{D-3}} + \frac{32\pi^2(\sqrt{G}Q)^2}{(D-2)(D-3)\Omega_{D-2}^2r^{2(D-3)}} - \frac{2\Lambda r^2}{(D-1)(D-2)} - \frac{Gb}{r^{(D-1)\omega+D-3}}, \quad (4)$$

$$\rho(r) = b \frac{\omega(D-1)(D-2)}{32\pi r^{(D-1)(\omega+1)}}, \quad (5)$$

where G is Newton's constant in arbitrary dimensions, M is the black hole's mass, Q is its electric charge, Λ is the cosmological constant (that we shall assume as negative in this paper), b is a parameter related to the energy density of Kiselev's fluid and $\Omega_{D-2} = 2\pi^{(D-1)/2}\Gamma((D-1)/2)$ is the area of the unit sphere \mathbb{S}^{D-2} (which in 4-dimensions gives 4π). We also have the form of the energy density, in which we assumed the standard matter coupling in Einstein equations $G_{\mu\nu} = 8\pi GT_{\mu\nu}$. If $\omega = 0$ and $b \propto M$, one reproduces the Schwarzschild contribution, when $\omega = (D-3)/(D-1)$ and $b \propto -Q^2$, one can also reproduce the Reissner-Nordström case, and so on.

A. Thermodynamics when G is a constant

In this first subsection, we are going to express the usual thermodynamic quantities when G is constant. It will be important to deduce the first law and Smarr formula for the D -dimensional Kiselev black hole as done in [31]. Then, from the form of the first law with G fixed, we shall verify what turns out when we let G varies as a thermodynamic quantity.

As usual, the temperature reads

$$T = \frac{\kappa}{2\pi} = \frac{1}{4\pi} \left[\frac{df}{dr} \right]_{r=r_+} = \frac{1}{4\pi} \left[\frac{D-3}{r_+} - \frac{32\pi^2 G Q^2}{(D-2)\epsilon_0 \Omega_{D-2}^2 r_+^{2D-5}} - 2\frac{\Lambda}{D-2}r_+ + (D-1)\frac{Gb\omega}{r_+^{D-2+\omega(D-1)}} \right], \quad (6)$$

¹ Let us be more explicit about the notation used: in fact, Kiselev's stress-energy tensor does not describe a perfect fluid, as highlighted in [32, 33], however an average equation of state parameter can be defined, which is given by ω . This issue is also discussed in the second section of [34].

where κ is the surface gravity.

The entropy is found by integrating the expression $dS = dM/T|_{Q,\Lambda,b}$ as

$$S = \frac{\Omega_{D-2}}{4G} r_+^{D-2} = \frac{A}{4G}, \quad (7)$$

where A is the area of the sphere of radius r_+ in D dimensions.

The electric potential reads

$$\phi = \left. \frac{\partial M}{\partial Q} \right|_{S,\Lambda,b} = \frac{4\pi Q}{(D-3)\Omega_{D-2}} r_+^{3-D}, \quad (8)$$

which gives exactly the electrostatic potential at $D = 4$ in Gaussian units.

The pressure is simply proportional to Λ and volume is its conjugate quantity:

$$P = -\frac{\Lambda}{8\pi G} \doteq \frac{(D-1)(D-2)}{16\pi G l^2}, \quad (9)$$

$$V = \left. \frac{\partial M}{\partial P} \right|_{S,Q,b} = \frac{\Omega_{D-2}}{D-1} r_+^{D-1}, \quad (10)$$

where, in fact, V is the Euclidean volume in D -dimensions, and we introduced the quantity l , that shall be convenient in the following sections.

The Kiselev (or quintessence) potential is

$$B = \left. \frac{\partial M}{\partial b} \right|_{S,Q,P} = -\frac{(D-2)\Omega_{D-2}}{16\pi} r_+^{\omega(1-D)}. \quad (11)$$

When $D = 4$, these quantities reduce to the ones found in [42]. From [31], we can see the Smarr formula and first law of thermodynamics takes the form

$$(D-3)M = (D-2)TS + (D-3)\phi Q - 2PV + [\omega(D-1) + D-3]Bb, \quad (12)$$

$$\delta M = T\delta S + V\delta P + \phi\delta Q + B\delta b. \quad (13)$$

In principle, these equations do not depend on G , but if we intend to investigate the effects of allowing it to vary, we need to start from the first law with, indeed, the black hole quantities that appear in the metric function (4). In fact, if we recover G , the first law assumes the form

$$\delta(GM) = \frac{\kappa}{8\pi}\delta A - \frac{V}{8\pi}\delta\Lambda + \sqrt{G}\phi\delta(\sqrt{G}Q) + B\delta(Gb). \quad (14)$$

This is the expression that is going to be analyzed in the following section.

B. Assuming G as a thermodynamic variable

It has been argued [28] that the Smarr relation that is supposed to be compatible with the holographic principle should be of the form $M = TS + \nu^i B_i + \mu C$, where C is the central charge that relates to the cosmological constant and Newton's constant as

$$C = k \frac{l^{D-2}}{16\pi G}, \quad (15)$$

where k depends on the specifics of the holographic model [27, 29] and μ is its corresponding potential, that we shall interpret as a chemical potential using the language of the black hole chemistry. The terms ν^i correspond to additional potentials of the problem [28], in this case the electric and Kiselev ones. Notice that the term of the bulk pressure (cosmological constant) does not appear on this relation, not even the spacetime dimension is made explicit. In this paper, we shall not discuss the details or soundness of this proposal, but instead we will investigate the consequences of assuming it in the Kiselev background. Also notice that C depends on the spacetime dimension D , Newton's constant G and also on the cosmological constant by means of the l term (see Eq.(9)).

From Eq.(14), if we let G to vary, we find

$$\delta M = \frac{\kappa}{8\pi G} \delta A - \frac{V}{8\pi G} \delta \Lambda + \phi \delta Q + B \delta b - \frac{\beta}{G} \delta G, \quad (16)$$

where

$$\beta = M - \frac{1}{2} \phi Q - bB = TS + \frac{1}{2} \phi Q + \tilde{\mu} C. \quad (17)$$

To obtain the last equation, we have used the proposed holographic Smarr relation

$$M = TS + \phi Q + bB + \mu C. \quad (18)$$

as a way to incorporate the central charge contribution. This is just a simple generalization of what was found in [29].

In order to go deeper in the exploration of this formalism, let us find the length dimensions L of the quantities involved in this framework (this can be seen from the thermodynamic quantities found in the previous section, from the metric function (4) and from the assumption of natural units):

$$\begin{aligned} [V] &= L^{D-1}, & [S] &= L^0, & [A] &= L^{D-2} = [G], & [\Lambda] &= [\mu]^2 = [M]^2 = L^{-2}, \\ [Q] &= L^{(D-4)/2}, & [\phi] &= L^{(2-D)/2}, & [b] &= L^{\omega(D-1)-1}, & [B] &= L^{\omega(1-D)}. \end{aligned} \quad (19)$$

From (16), (17) and (19), we derive the Smarr relation (now considering G and Λ) for variable G as

$$-M = \frac{V}{4\pi G} \Lambda - \phi Q - (D-2)\mu C + [\omega(D-1) - 1]bB, \quad (20)$$

from which we find

$$V = \frac{8\pi G l^2}{(D-1)(D-2)} \{M - \phi Q - (D-2)\mu C + [\omega(D-1) - 1]bB\}. \quad (21)$$

Again, using the holographic Smarr relation (18) (in which the information about D , G and Λ is inserted in the thermodynamic quantity C), this can be simplified as

$$\begin{aligned} V &= \frac{1}{2P} \{(D-2)TS - (D-3)(TS + \mu C + bB) + [\omega(D-1) + D-3]bB\} = \\ &= \frac{1}{2P} \{(D-2)TS - (D-3)\phi Q + [\omega(D-1) + D-3]bB - (D-3)M\}, \end{aligned} \quad (22)$$

where we used again (18). This is exactly the Smarr relation (12) when G is a constant. Therefore, we demonstrate the preservation, for different fluids, of the result found in [29] about the compatibility between the *holographic Smarr relation*, the *first law of black hole thermodynamics in D -dimensions with variable G* and the *Smarr relation in D -dimensions for constant G* .

In order to properly write down this approach using the pressure and the central charge, we see that from Eq.(15) we have

$$\frac{\delta G}{G} = -\frac{2}{D} \frac{\delta C}{C} - \frac{(D-2)}{D} \frac{\delta P}{P}, \quad (23)$$

which allows us to write the bulk first law (16) as

$$\delta M = T\delta S + \phi\delta Q + B\delta b + V_C\delta P + \mu\delta C, \quad (24)$$

where

$$V_C = \frac{2M + (D-4)\phi Q + 2[\omega(D-1) - 1]bB}{2DP}, \quad \mu = \frac{2P(V_C - V)}{C(D-2)}. \quad (25)$$

The term V_C is the new thermodynamic volume, that generalizes the one found in [29], and μ is the conjugate quantity that furnishes the response of the system to variations in the central charge. As can be verified, if $\delta G = 0$, one recovers the usual first law of thermodynamics (13), since δC and δP must be related in this case.

III. EXTENDED PHASE SPACE THERMODYNAMICS WITH VARIABLE G

In order to apply the techniques explored in the previous section, we resort to the 4-dimensional case. So, consider the metric function (4) when $D = 4$ and the relation between Λ and l from (9):

$$f(r) = 1 - \frac{2GM}{r} + \frac{GQ^2}{r^2} + \frac{r^2}{l^2} - \frac{Gb}{r^{3\omega+1}}. \quad (26)$$

The standard thermodynamic quantities can be straightforwardly calculated:

$$M = \frac{r_+}{2G} + \frac{Q^2}{2r_+} + \frac{r_+^3}{2Gl^2} - \frac{b}{2r_+^{3\omega}}, \quad T = \frac{3r_+^4 + l^2r_+^2 - GQ^2l^2 + 3Gbl^2\omega r_+^{1-3\omega}}{4\pi l^2 r_+^3}, \quad S = \frac{\pi r_+^2}{G} \quad (27)$$

$$V = \frac{4\pi}{3}r_+^3, \quad \phi = \frac{Q}{r_+}, \quad B = -\frac{1}{2r_+^{3\omega}}. \quad (28)$$

And the new variables defined in (25):

$$V_C = \frac{\pi}{3r_+}(GQ^2l^2 + l^2r_+^2 + r_+^4 - 3Gbl^2\omega r_+^{-3\omega+1}), \quad \mu = \frac{2\pi}{kl^4r_+}(GQ^2l^2 + l^2r_+^2 - 3r_+^4 - 3Gbl^2\omega r_+^{-3\omega+1}). \quad (29)$$

From the above expressions, we see that if $b\omega < 0$, the thermodynamics volume V_C is a positive definite quantity, and this condition coincides with the one that requires the positiveness of the energy density of Kiselev fluid (5). Since P and C are positive quantities, we see that in the phase space region in which $V_C > V$, the following reverse isoperimetric inequality [15, 29, 43]

$$\left[\frac{(D-1)V}{\Omega_{D-2}} \right]^{\frac{1}{D-1}} \geq \left(\frac{A}{\Omega_{D-2}} \right)^{\frac{1}{D-2}} \quad (30)$$

will remain valid for the new thermodynamic volume V_C . As shown in [44], for the usual Kiselev black hole volume, this inequality is valid for $1 \geq (-\omega)^{1/3}$.

Now, let us investigate the sign of the chemical potential μ and its relation to the black hole size and the reverse isoperimetric inequality for some specific cases of Kiselev fluids.

- $\omega = -1/3$. In this case, the standard isoperimetric inequality is satisfied and it is possible to directly relate the size of the black hole and the sign of the chemical potential. In fact, one has $\mu > 0$ for small black holes

$$r_+^2 < \frac{l^2}{6} \left[(1 + Gb) + \sqrt{1 + Gb + 12 \frac{GQ^2}{l^2}} \right], \quad (31)$$

which also satisfies the inequality (30) for the new thermodynamic volume V_C , since in this case $V_C > V$. On the other hand, large black holes are characterized by $\mu < 0$. This classification and its relation with the isoperimetric inequality is compatible with the results found in [29]. The behavior of $\mu(r_+)$ can be seen in Fig.(1) in the blue/dotted line, where we described the dimensionful functions in terms of a fiducial length ℓ_0 .


- **Average quintessence** $\omega = -2/3$. In this case, the inequality involving r_+ that separates positive and negative values of μ is quite complicated. Nevertheless, we can see that the term $-3r_+^4$ in (29) will dominate for very large black holes. Therefore, we verify the same classification above, i.e., that small black holes present $\mu > 0$, while large ones present $\mu < 0$. Also, the reverse isoperimetric inequality is satisfied for V and will be valid for V_C for small black holes. The behavior of $\mu(r_+)$ can be seen in Fig.(1) in the black/dashed line.

As a matter of fact, this behavior will occur whenever we have $-1 < \omega \leq 0$. If $\omega = -1$, this behavior will also occur as long as $Gb < l^{-2}$, in order to preserve the anti-de Sitter-like metric contribution (26).


- **Average phantom field** $\omega = -4/3$. In this case, one has $\mu > 0$ for

$$m(r_+) = GQ^2l^2 + l^2r_+^2 - 3r_+^4 + 4Gbl^2r_+^5 > 0, \quad (32)$$

which is achievable in a more diverse regime than the previous case. One can have $\mu > 0$ for small and very large black holes, depending on the parameters. Therefore, in this case, the sign of μ does not work for this classification of black hole sizes, as can be seen in Fig.(1) in the red/solid curve. Curiously in this case, the reverse isoperimetric inequality (30) is no longer valid for the standard volume V .



(a) We set $Q = G\ell_0^{-2} = 1$, $l/\ell_0 = 7.8$, $b\ell_0^{1-3\omega} = 10^{-3}$ and the central charge parameter $k = 16\pi$.



(b) We set $Q = 0.05$, $G\ell_0^{-2} = 1$, $l/\ell_0 = 2$, $b\ell_0^{1-3\omega} = 10^{-1}$ and the central charge parameter $k = 16\pi$. The red curve grows indefinitely.

FIG. 1: Depiction of $\mu(r_+) \times r_+$ in units of a fiducial length ℓ_0 .

IV. CRITICAL BEHAVIOR

As also happened in the original formulation of black hole chemistry, we can also study phase transitions induced by the presence of the black hole electric charge. The critical quantities are defined by the saddle points of our thermodynamic functions. In this case, it will be convenient to study the equation of state $T = T(r_+)$ found in (27) and to solve the following system of equations

$$\frac{\partial T(r_c, l_c, C_c)}{\partial r_+} = 0 = \frac{\partial^2 T(r_c, l_c, C_c)}{\partial r_+^2}, \quad (33)$$

where we use the definition of the central charge (15) and the relation (9) between l and P to find

$$G = \frac{1}{8\pi} \sqrt{\frac{3k}{2PC}}. \quad (34)$$

These critical outer horizon radius r_c , cosmological constant l_c , central charge C_c and temperature $T_c = T(r_c, l_c, C_c)$ can only be found for specific values of the Kiselev parameter ω . So, let us analyze the same specific cases of the previous section.

- $\omega = -1/3$: The critical radius, cosmological constant, temperature and central charge are found to be

$$r_c = \frac{\sqrt{6GQ}}{\sqrt{1-Gb}}, \quad l_c = \frac{6\sqrt{GQ}}{|1-Gb|}, \quad T_c = \frac{\sqrt{6}(1-Gb)^{3/2}}{18\pi\sqrt{GQ}}, \quad C_c = \frac{9kQ^2}{4\pi(1-Gb)^2}. \quad (35)$$

Notice that due to the coupling with the Kiselev parameter b , the critical central charge C_c now depends on G , which may indicate the violation of the universality that was presented in [29], since this may be translated into a pressure dependence. In fact, using Eq.(9) and l_c , we find

$$G = \frac{1}{b + \sqrt{96\pi PQ}}, \quad (36)$$

which can be independently varied for fixed Q and b . Besides that, the other critical quantities are the same that are found considering the usual extended phase space thermodynamics [42].

- **Average quintessence $\omega = -2/3$** : In this case, we have

$$r_c = \sqrt{6GQ}, \quad l_c = \sqrt{6}r_c, \quad T_c = \frac{\sqrt{6}}{18\pi\sqrt{GQ}} - \frac{Gb}{2\pi}, \quad C_c = \frac{9kQ^2}{4\pi}. \quad (37)$$

Notice that r_c and l_c do not couple to b , and that T_c only suffers a translation, which also is consistent with previous criticality studies in the standard extended phase space [42]. Now, C_c does not depend G and one recovers the universal behavior and also the same relation between G and P of [29], given by (36) if $b = 0$.




FIG. 2: Depiction of the Free Energy versus Temperature $F \times T$ in units of a fiducial length ℓ_0 for the average quintessence fluid $\omega = -2/3$. We set $Q = 1$, $P\ell_0^4 = 15$, $b\ell_0^3 = 0.3$ and the central charge parameter $k = 16\pi$. The values of the central charge are described in the figure.

- **Average phantom fluid $\omega = -4/3$:**

$$l_c = \frac{\sqrt{2}\sqrt{3Q^2r_c^2 + 2br_c^7}}{\sqrt{Q^2 + 4br_c^5}}, \quad T_c = \frac{2Q^2 + 3br_c^5}{6\pi Q^2r_c + 4\pi br_c^6}, \quad C_c = \frac{k(3Q^2 + 2br_c^5)^2}{4\pi(Q^2 + 4br_c^5)}, \quad (38)$$

where r_c satisfies the equation $4Gbr_c^5 - r_c^2 + 6GQ^2 = 0$ that cannot be solved analytically. In any case, one also loses universality for the critical radius, since variations in G imply in variations in r_c . Consequently, this implies in absence of universality of l_c , T_c and of C_c .

We can see these transitions from the behavior of the Helmholtz Free Energy and the Temperature

$$F = M - TS = \frac{3GQ^2l^2 + l^2r_+^2 - r_+^4}{4Gl^2r_+} - \frac{(3\omega + 2)b}{4r_+^3} = \frac{2\pi r_+}{3} \sqrt{\frac{6PC}{k}} - \frac{2\pi Pr_+^3}{3} + \frac{3Q^2}{4r_+} - \frac{(3\omega + 2)b}{4r_+^3}, \quad (39)$$

$$T = \frac{1}{4\pi r_+} + \sqrt{\frac{6kP}{C}} \frac{r_+}{8\pi} - \sqrt{\frac{6k}{CP}} \frac{Q^2}{64\pi^2 r_+^3} + \sqrt{\frac{6k}{CP}} \frac{3\omega}{64\pi^2} \frac{b}{r_+^{2+3\omega}}, \quad (40)$$

where we used equation (34) and (9) to express the terms with G and l in terms of P and C . In Fig.(2), we depict the free energy as a function of the temperature for the case of an average quintessence fluid $\omega = -2/3$. As can be seen, the blue/dotted curve describes the critical curve, from which phase transitions can occur in the region above in the phase space $F \times T$, as can be seen in the orange/solid line. Below the critical curve, for smaller values of the central charge, no transitions are allowed (green/dashed curve). From this figure, one realizes the fundamental role played by the central charge in the determination of criticality, which is a novel featured highlighted in Ref.[29]. For other values of parameter ω the free energy follows a similar behavior, therefore, we do not depict them here.

V. CONCLUDING REMARKS

Based on the work of Cong, Kubizňák and Mann [29], we analyzed a new proposal of extended phase space for the thermodynamics of black holes, which is compatible with the holographic program. It is possible, by the introduction of a novel thermodynamic variable that depends on Newton's constant (which is allowed to vary), the bulk cosmological constant and the spacetime dimension. In particular, we extended that work for the case of different matter contents that surround a static black hole, modeled by the Kiselev fluid, which is labeled by a free parameter.

Our work demonstrated the validity of the holographic technique for the class of Kiselev black holes. We generalized the bulk first law of thermodynamics by the introduction of an alternative thermodynamic volume and a chemical potential-like term μ , which is conjugate to the central charge. As an application, we analyzed the 4-dimensional

case assuming three different average equations of state parameters ($\omega = -1/3, -2/3, -4/3$). For which we studied their relationships between the sign of the chemical potential and the size of the black hole. Thus, when the reverse isoperimetric inequality is valid for the standard volume V , then the black holes follow a simple size classification based on the sign of the chemical potential: positive (negative) μ for small (large) black holes. On the other hand, if this inequality is no longer valid for V , then such classification is broken since the polynomial that regulates the sign of μ has a more diverse behavior.

We also analyzed phase transitions, in which the central charge plays a fundamental role in the determination of the criticality, thus in agreement with results of [29]. However, we verified an important divergence between ours and those previous results: a possible surrounding matter for which the universality of the critical central charge is no longer valid. This property is violated as a result of the dependence of the critical central charge on Newton's constant due to the coupling with Kiselev's fluid. We also illustrated the behavior of the phase transitions from the phase space diagram of the free energy versus the temperature. We observe that the critical central charge plays the role of separating the space between regions in which phase transition may or may not occur.

Acknowledgements

I. P. L. would like to acknowledge the contribution of the COST Action CA18108. I. P. L. was partially supported by the National Council for Scientific and Technological Development - CNPq grant 306414/2020-1. R. B. A. was partially supported by the Tutoring Program of the Federal University of Lavras.

-
- [1] J. D. Bekenstein, "Black holes and the second law," *Lett. Nuovo Cim.* **4** (1972) 737–740.
 - [2] J. D. Bekenstein, "Black holes and entropy," *Phys. Rev. D* **7** (1973) 2333–2346.
 - [3] J. M. Bardeen, B. Carter, and S. W. Hawking, "The Four laws of black hole mechanics," *Commun. Math. Phys.* **31** (1973) 161–170.
 - [4] S. W. Hawking, "Particle Creation by Black Holes," *Commun. Math. Phys.* **43** (1975) 199–220. [Erratum: *Commun. Math. Phys.* **46**, 206 (1976)].
 - [5] S. W. Hawking and D. N. Page, "Thermodynamics of Black Holes in anti-De Sitter Space," *Commun. Math. Phys.* **87** (1983) 577.
 - [6] R.-G. Cai, D.-W. Pang, and A. Wang, "Born-Infeld black holes in (A)dS spaces," *Phys. Rev. D* **70** (2004) 124034, [arXiv:hep-th/0410158](#).
 - [7] A. de la Cruz-Dombriz, A. Dobado, and A. L. Maroto, "Black Holes in f(R) theories," *Phys. Rev. D* **80** (2009) 124011, [arXiv:0907.3872 \[gr-qc\]](#). [Erratum: *Phys. Rev. D* **83**, 029903 (2011)].
 - [8] V. Faraoni, "Black hole entropy in scalar-tensor and f(R) gravity: An Overview," *Entropy* **12** (2010) 1246, [arXiv:1005.2327 \[gr-qc\]](#).
 - [9] J. P. Morais Graça, I. P. Lobo, and I. G. Salako, "Cloud of strings in f(R) gravity," *Chin. Phys. C* **42** (2018) no. 6, 063105, [arXiv:1708.08398 \[gr-qc\]](#).
 - [10] R. C. Myers and J. Z. Simon, "Black Hole Thermodynamics in Lovelock Gravity," *Phys. Rev. D* **38** (1988) 2434–2444.
 - [11] D. Blas and S. Sibiryakov, "Horava gravity versus thermodynamics: The Black hole case," *Phys. Rev. D* **84** (2011) 124043, [arXiv:1110.2195 \[hep-th\]](#).
 - [12] R. M. Wald, "The thermodynamics of black holes," *Living Rev. Rel.* **4** (2001) 6, [arXiv:gr-qc/9912119](#).
 - [13] D. Kastor, S. Ray, and J. Traschen, "Enthalpy and the Mechanics of AdS Black Holes," *Class. Quant. Grav.* **26** (2009) 195011, [arXiv:0904.2765 \[hep-th\]](#).
 - [14] D. Kubiznak and R. B. Mann, "P-V criticality of charged AdS black holes," *JHEP* **07** (2012) 033, [arXiv:1205.0559 \[hep-th\]](#).
 - [15] D. Kubiznak, R. B. Mann, and M. Teo, "Black hole chemistry: thermodynamics with Lambda," *Class. Quant. Grav.* **34** (2017) no. 6, 063001, [arXiv:1608.06147 \[hep-th\]](#).
 - [16] S. Fernando, "P-V criticality in AdS black holes of massive gravity," *Phys. Rev. D* **94** (2016) no. 12, 124049, [arXiv:1611.05329 \[gr-qc\]](#).
 - [17] M.-S. Ma and R.-H. Wang, "Peculiar P – V criticality of topological Hořava-Lifshitz black holes," *Phys. Rev. D* **96** (2017) no. 2, 024052, [arXiv:1707.09156 \[gr-qc\]](#).
 - [18] S. H. Hendi and Z. Armanfard, "Extended phase space thermodynamics and P – V criticality of charged black holes in Brans–Dicke theory," *Gen. Rel. Grav.* **47** (2015) no. 10, 125, [arXiv:1503.07070 \[gr-qc\]](#).
 - [19] B. R. Majhi and S. Samanta, "P-V criticality of AdS black holes in a general framework," *Phys. Lett. B* **773** (2017) 203–207, [arXiv:1609.06224 \[gr-qc\]](#).
 - [20] I. P. Lobo, H. Moradpour, J. P. Morais Graça, and I. G. Salako, "Thermodynamics of Black Holes in Rastall Gravity," *Int. J. Mod. Phys. D* **27** (2018) no. 07, 1850069, [arXiv:1710.04612 \[gr-qc\]](#).
 - [21] C. V. Johnson, "Holographic Heat Engines," *Class. Quant. Grav.* **31** (2014) 205002, [arXiv:1404.5982 \[hep-th\]](#).

- [22] B. Eslam Panah, K. Jafarzade, and S. H. Hendi, “Charged 4D Einstein-Gauss-Bonnet-AdS black holes: Shadow, energy emission, deflection angle and heat engine,” *Nucl. Phys. B* **961** (2020) 115269, [arXiv:2004.04058 \[hep-th\]](#).
- [23] D. V. Singh and S. Siwach, “Thermodynamics and P-v criticality of Bardeen-AdS Black Hole in 4D Einstein-Gauss-Bonnet Gravity,” *Phys. Lett. B* **808** (2020) 135658, [arXiv:2003.11754 \[gr-qc\]](#).
- [24] J. P. Morais Graça, I. P. Lobo, V. B. Bezerra, and H. Moradpour, “Effects of a string cloud on the criticality and efficiency of AdS black holes as heat engines,” *Eur. Phys. J. C* **78** (2018) no. 10, 823, [arXiv:1806.02913 \[gr-qc\]](#).
- [25] V. B. Bezerra, I. P. Lobo, J. P. Morais Graça, and L. C. N. Santos, “Effects of quantum corrections on the criticality and efficiency of black holes surrounded by a perfect fluid,” *Eur. Phys. J. C* **79** (2019) no. 11, 949, [arXiv:1908.08140 \[gr-qc\]](#).
- [26] C. V. Johnson, “Born–Infeld AdS black holes as heat engines,” *Class. Quant. Grav.* **33** (2016) no. 13, 135001, [arXiv:1512.01746 \[hep-th\]](#).
- [27] A. Karch and B. Robinson, “Holographic Black Hole Chemistry,” *JHEP* **12** (2015) 073, [arXiv:1510.02472 \[hep-th\]](#).
- [28] M. R. Visser, “Holographic Thermodynamics Requires a Chemical Potential for Color,” [arXiv:2101.04145 \[hep-th\]](#).
- [29] W. Cong, D. Kubiznak, and R. B. Mann, “Thermodynamics of AdS Black Holes: Critical Behavior of the Central Charge,” *Phys. Rev. Lett.* **127** (2021) no. 9, 091301, [arXiv:2105.02223 \[hep-th\]](#).
- [30] V. V. Kiselev, “Quintessence and black holes,” *Class. Quant. Grav.* **20** (2003) 1187–1198, [arXiv:gr-qc/0210040](#).
- [31] S. Chen, B. Wang, and R. Su, “Hawking radiation in a d -dimensional static spherically-symmetric black Hole surrounded by quintessence,” *Phys. Rev. D* **77** (2008) 124011, [arXiv:0801.2053 \[gr-qc\]](#).
- [32] M. Visser, “The Kiselev black hole is neither perfect fluid, nor is it quintessence,” *Class. Quant. Grav.* **37** (2020) no. 4, 045001, [arXiv:1908.11058 \[gr-qc\]](#).
- [33] P. Boonserm, T. Ngampitipan, A. Simpson, and M. Visser, “Decomposition of the total stress energy for the generalized Kiselev black hole,” *Phys. Rev. D* **101** (2020) no. 2, 024022, [arXiv:1910.08008 \[gr-qc\]](#).
- [34] I. P. Lobo, M. G. Richarte, J. P. Morais Graça, and H. Moradpour, “Thin-shell wormholes in Rastall gravity,” *Eur. Phys. J. Plus* **135** (2020) no. 7, 550, [arXiv:2007.05641 \[gr-qc\]](#).
- [35] V. Sahni, “Dark matter and dark energy,” *Lect. Notes Phys.* **653** (2004) 141–180, [arXiv:astro-ph/0403324](#).
- [36] R. R. Caldwell, “A Phantom menace?,” *Phys. Lett. B* **545** (2002) 23–29, [arXiv:astro-ph/9908168](#).
- [37] F. R. Tangherlini, “Schwarzschild field in n dimensions and the dimensionality of space problem,” *Nuovo Cim.* **27** (1963) 636–651.
- [38] G. Kunstatter, “ d -dimensional black hole entropy spectrum from quasinormal modes,” *Phys. Rev. Lett.* **90** (2003) 161301, [arXiv:gr-qc/0212014](#).
- [39] R.-J. Yang and H. Gao, “Constraints from accretion onto a Tangherlini-Reissner-Nordstrom black hole,” [arXiv:1801.07973 \[gr-qc\]](#).
- [40] B. Wu, W. Liu, H. Tang, and R.-H. Yue, “Destroying charged black holes in higher dimensions with test particles,” *Int. J. Mod. Phys. A* **32** (2017) no. 21, 1750125.
- [41] R. Emparan and H. S. Reall, “Black Holes in Higher Dimensions,” *Living Rev. Rel.* **11** (2008) 6, [arXiv:0801.3471 \[hep-th\]](#).
- [42] A. Jawad, I. Siddique, I. P. Lobo, and W. u. Salam, “Effects of Gauss–Bonnet entropy on thermodynamics of Kiselev black hole,” *Int. J. Mod. Phys. D* **29** (2020) no. 15, 2050101.
- [43] M. Cvetič, G. W. Gibbons, D. Kubiznak, and C. N. Pope, “Black Hole Enthalpy and an Entropy Inequality for the Thermodynamic Volume,” *Phys. Rev. D* **84** (2011) 024037, [arXiv:1012.2888 \[hep-th\]](#).
- [44] M. Azreg-Aïnou, “Charged de Sitter-like black holes: quintessence-dependent enthalpy and new extreme solutions,” *Eur. Phys. J. C* **75** (2015) no. 1, 34, [arXiv:1410.1737 \[gr-qc\]](#).