

## On symmetries and charges at spatial infinity

---

**Lennart Brocki<sup>a</sup> Jerzy Kowalski-Glikman<sup>a,b</sup>**

<sup>a</sup>*Institute for Theoretical Physics, University of Wrocław, pl. M. Borna 9, 50-204 Wrocław, Poland*

<sup>b</sup>*National Centre for Nuclear Research, Pasteura 7, 02-093 Warsaw, Poland*

*E-mail:* [lennart.brocki@uwr.edu.pl](mailto:lennart.brocki@uwr.edu.pl), [jerzy.kowalski-glikman@uwr.edu.pl](mailto:jerzy.kowalski-glikman@uwr.edu.pl)

**ABSTRACT:** Following the recent work of Henneaux and Troessaert [1], which revisits the problem of spacetime symmetries at spatial infinity, we analyze this problem using the Bondi metric without determinant condition as our starting point. It turns out that in this case the symmetries at spatial infinity form the BMS symmetry appended with an additional infinite set of abelian symmetries. We furthermore find that imposing the determinant condition to the Bondi metric would result in a drastic reduction of symmetries, with no spatial (super) translations present.

**KEYWORDS:** asymptotically flat spacetimes, charges at infinity

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Review of previous results</b>	<b>3</b>
2.1	Role of surface integrals and extension of the original treatment to include BMS	3
<b>3</b>	<b>Conjugate momenta in terms of spacetime metric components</b>	<b>8</b>
<b>4</b>	<b>Finiteness of symplectic structure</b>	<b>10</b>
<b>5</b>	<b>Leading order of constraints</b>	<b>12</b>
<b>6</b>	<b>Asymptotic symmetries</b>	<b>14</b>
6.1	Preservation of falloff and gauge conditions	14
6.2	Determinant condition and symmetry reduction	17
6.3	The charges	18
6.4	Discussion of the asymptotic symmetry	19
<b>7</b>	<b>Conclusions</b>	<b>20</b>
<b>A</b>	<b>3+1 decomposition</b>	<b>21</b>
<b>B</b>	<b>The algebra of asymptotic symmetries</b>	<b>22</b>

---

## 1 Introduction

Take a black box. There is some stuff in it and you want to know how to characterize its motion not looking inside the box. This can be done by measuring conserved charges given by integrals of specific functions of gravitational field. These charges are associated with physical symmetries, which in turn result from gauge symmetries of gravity being broken by the presence of boundaries (see [2] and references therein for in-depth discussion.) A particularly interesting physical situation arises when the boundary is situated far away from localized material sources, where the geometry is almost flat. In the seminal paper [3] Regge and Teitelboim formulated the well-defined variational principle for gravity on asymptotically flat space by adding an appropriate boundary term to the Einstein-Hilbert action and then show that the resulting boundary charges satisfy the Poincaré algebra and therefore measure the energy, momenta, angular momenta and boost charges of spacetime. This result was well expected and not surprising because the Poincaré symmetry is the physical symmetry of flat Minkowski space. A really surprising result was derived more

than a decade before the work of Regge and Teitelboim, in [4], [5], [6] where Bondi, van der Burg, Metzner and Sachs established that the physical symmetries at null infinity of asymptotically flat spacetime are much larger and form an infinite dimensional group, the BMS group, containing Poincaré group as its (maximal) finite-dimensional subgroup (see [7] for a review of exciting recent developments.) Then an immediate question arises, why the symmetries at null infinity are so much different from the ones at the spatial infinity? Or are they?

In the recent publication [1] Henneaux and Troessaert revisit the analysis [3] of Regge and Teitelboim of asymptotically flat spaces in the Hamiltonian (ADM) formulation and derived an infinite set of charges generating BMS supertranslations at spatial infinity. Therefore the asymptotic symmetry of asymptotically flat spacetimes, according to [1], is the (unextended) BMS group. This is in contrast to the original result of [3] where these charges were found to vanish and the asymptotic symmetry was determined to form the Poincaré group and not the BMS group. The key difference of the two approaches is that Henneaux and Troessaert make the asymptotic expansion in spherical coordinates that makes it possible for them to use parity conditions on the phase space functions different from the ones used by Regge and Teitelboim in their case of Cartesian coordinates expansion. These conditions are the crucial point of the analysis because they guarantee cancellation of divergences, generally plaguing expressions for the asymptotic symplectic form and the charges.

The treatments in [1] and [3] have in common that they take as a starting point a generic asymptotic expansion of a spatial metric and its conjugate momenta. In this paper we instead perform a 3+1 decomposition of a spacetime metric  $g$  in Bondi coordinates which is asymptotically flat at null infinity, off-shell and fulfills the Bondi gauge except for the determinant condition. We drop the determinant condition since, as it will be explained in detail below, its presence drastically reduces the asymptotic symmetries. Using the ADM formalism we express the expansion of the spatial metric and momenta in terms of the metric functions and their derivatives. These expressions are then substituted into the symplectic form, Hamiltonian and diffeomorphism constraint and charges given in [1]. This procedure leads to two interesting insights.

First, it shows that there exists a term in the symplectic form which is divergent and which can not be canceled by imposing parity conditions. We propose new boundary conditions which cancel this divergence and argue that they are very mild conditions because they amount to demanding that the time derivatives of some components of the metric vanish at the past boundary of future null infinity  $\mathcal{I}_-^+$  so that the total energy radiated by the system is finite. Additionally we find that the foliation of spacetime is constrained by the demand that the symplectic form be finite.

Second, we find an asymptotic symmetry at spatial infinity that is larger than the BMS symmetry. The crucial difference to the treatment of [1] lies in the different choice of boundary conditions which lead to finite charges corresponding to supertranslations which lie outside of the BMS symmetry. This larger-than-BMS algebra is isomorphic to the one found by Troessaert [11].

The plan of the paper is as follows. In the next section we recall the results presented in

[1] and [3]. In the following Sect. 3 we compute the conjugate momenta and express them in terms of the spacetime metric components. We then closely follow the treatment of [1] and analyze which differences arise when substituting these expressions for the momenta. Sect. 4 is devoted to the discussion of the asymptotic symplectic structure, while Sect. 5 concerns the Hamiltonian and diffeomorphism constraint. In Sect. 6 we discuss the consequences for the asymptotic symmetry. We end this paper with some concluding remarks in Sect. 7.

## 2 Review of previous results

In this section we recall some results derived in [1] and [3] that we are going to make use of in the paper.

### 2.1 Role of surface integrals and extension of the original treatment to include BMS

Since it is crucial for our discussion, we briefly review the Hamiltonian analysis made in [3]. The main result is that in order to have a well-defined Hamiltonian formalism one has to add surface integrals to the Hamiltonian which, for asymptotically flat spacetimes, turn out to be the Poincaré charges. Asymptotically flat spacetimes and the associated conjugate momenta in asymptotically Cartesian coordinates are assumed to obey the asymptotic expansion ( $r^2 = \sum_i x_i^2$ )

$$g_{ij} = \delta_{ij} + \frac{1}{r} \bar{h}_{ij} + O(r^2) \quad (2.1)$$

$$\pi^{ij} = \frac{1}{r^2} \bar{\pi}^{ij} + O(r^3). \quad (2.2)$$

Hamilton's equations are defined as the functional derivatives

$$\dot{g}_{ij} = \frac{\delta H}{\delta \pi^{ij}} \quad (2.3)$$

$$\dot{\pi}^{ij} = -\frac{\delta H}{\delta g_{ij}} \quad (2.4)$$

which are, by definition, the coefficients of  $\delta g_{ij}$  and  $\delta \pi^{ij}$  in the variation of the Hamiltonian

$$\delta H = \int d^3x [A^{ij} \delta g_{ij} + B_{ij} \delta \pi^{ij}]. \quad (2.5)$$

Therefore, for Hamilton's equations to be defined properly, it is necessary that the variation of the Hamiltonian can be put into the form of (2.5). The important observation made in [3] is that this is not the case for the Hamiltonian constraint of general relativity

$$H_0 = \int d^3x N \mathcal{H} + N^i \mathcal{H}_i \quad (2.6)$$

but that instead extra surface terms appear

$$\delta H_0 = \int d^3x [A^{ij} \delta g_{ij} + B_{ij} \delta \pi^{ij}] + \int d^2x K_{N, N^i}, \quad (2.7)$$

which arise as a result of partial integration moving a derivative away from the variations of the canonical variables. If the boundary term in the last expression can be rewritten as the variation of some quantity  $C$ , i.e. if it is exact

$$\delta H_0 = \int d^3x [A^{ij} \delta g_{ij} + B_{ij} \delta \pi^{ij}] - \delta C_{N,N^i} \quad (2.8)$$

this quantity can now be added to the Hamiltonian, such that the variation of the redefined Hamiltonian is of the form

$$\delta(H_0 + C_{N,N^i}) = \int d^3x [A^{ij} \delta g_{ij} + B_{ij} \delta \pi^{ij}] \quad (2.9)$$

and thus the functional derivatives are well defined. Then if the surface integral  $C$  is finite, it turns out to be a combination of Poincaré charges that satisfy the Poincaré algebra. Furthermore the transformations generated by these charges leave the asymptotic conditions (2.1), (2.2) invariant. Physically, this means that the asymptotic observer can measure the total momentum and angular momentum, as well as the boost, of asymptotically flat spacetime. In order to make the charges and asymptotic symplectic structure finite Regge and Teitelboim introduce certain parity conditions on the leading order terms  $\bar{h}_{ij}$  and  $\bar{\pi}^{ij}$  in the metric expansion [3], where parity here refers to antipodal points on the sphere, i.e. the map  $\mathbf{n} \rightarrow -\mathbf{n}$ , where  $\mathbf{n}$  is a unit vector. Explicitly, it is assumed that

$$\bar{h}_{ij}(-\mathbf{n}) = \bar{h}_{ij}(\mathbf{n}), \quad \bar{\pi}_{ij}(-\mathbf{n}) = -\bar{\pi}_{ij}(\mathbf{n}). \quad (2.10)$$

so that  $\bar{h}_{ij}$  is even under parity, while  $\bar{\pi}^{ij}$  is odd.

It is crucial to notice that approach of Regge and Teitelboim in addition to the standard Poincaré translations and Lorentz transformations the asymptotic conditions are also invariant under angle-dependent translations, however, the associated charges vanish due to the introduced parity conditions. This means that the actual asymptotic symmetry algebra is the Poincaré algebra since the symmetry under angle-dependent translations is pure gauge, i.e., the charges associated with these symmetries vanish as a consequence of the chosen parity conditions<sup>1</sup>.

In the recent paper [1] Henneaux and Troessaert propose different parity conditions which keep Hamilton's equations well-defined but are less restrictive in the sense that they render the charges associated with angle-dependent translations finite. They further show that the algebra of these charges is isomorphic to the BMS algebra.

Instead of Cartesian coordinates used by Regge and Teitelboim, in [1] Henneaux and

---

<sup>1</sup>We will discuss the relation between gauge symmetries, boundary charges and asymptotic conditions in more details below.

Troessaert employ spherical coordinates and the asymptotic conditions take the form

$$h_{rr} = 1 + \frac{1}{r}\bar{h}_{rr} + \frac{1}{r^2}h_{rr}^{(2)} + o(r^{-2}) \quad (2.11)$$

$$h_{rA} = \bar{h}_{rA} + \frac{1}{r}h_{rA}^{(2)} + o(r^{-1}) \quad (2.12)$$

$$h_{AB} = r^2\bar{\gamma}_{AB} + r\bar{h}_{AB} + h_{AB}^{(2)} + o(1) \quad (2.13)$$

$$\pi^{rr} = \bar{\pi}^{rr} + \frac{1}{r}\pi^{(2)rr} + o(r^{-1}) \quad (2.14)$$

$$\pi^{rA} = \frac{1}{r}\bar{\pi}^{rA} + \frac{1}{r^2}\pi^{(2)rA} + o(r^{-2}) \quad (2.15)$$

$$\pi^{AB} = \frac{1}{r^2}\bar{\pi}^{AB} + \frac{1}{r^3}\pi^{(2)AB} + o(r^{-3}) \quad (2.16)$$

while the parity conditions are

$$\bar{\lambda} \sim \bar{\pi}^{AB} = \text{even}, \quad \bar{p} \sim \bar{k}_{AB} \sim \bar{\pi}^{rA} = \text{odd}, \quad (2.17)$$

where

$$\bar{\lambda} = \frac{1}{2}\bar{h}_{rr}, \quad \bar{k}_{AB} = \frac{1}{2}\bar{h}_{AB} + \bar{\lambda}\gamma_{AB} \quad (2.18)$$

$$\bar{p} = 2(\bar{\pi}^{rr} - \bar{\pi}^A_A), \quad \pi_{(k)}^{AB} = 2\bar{\pi}^{AB}. \quad (2.19)$$

and  $\gamma_{AB}$  is the metric on the unit sphere. Although a generic expansion of an asymptotically flat metric allows the term  $\bar{h}_{rA}$  to be non-vanishing Henneaux and Troessaert assume [1] that

$$\bar{h}_{rA} = 0 \quad (2.20)$$

which is necessary in order for the boost charges to be integrable.

The parity conditions are introduced to cancel the following logarithmic divergences in the Hamiltonian kinetic term, i.e., the symplectic structure

$$\int dr \frac{1}{r} \int d\theta d\varphi \left( \bar{\pi}^{rr} \dot{\bar{h}}_{rr} + \bar{\pi}^{AB} \dot{\bar{h}}_{AB} \right) = \int dr \frac{1}{r} \int d\theta d\varphi \left( \bar{p} \dot{\bar{\lambda}} + \pi_{(k)}^{AB} \dot{\bar{k}}_{AB} \right), \quad (2.21)$$

which is zero because the integral over the sphere of a function with odd parity vanishes. It is furthermore demonstrated that all divergences occurring in the expressions for the charges can be canceled by imposing the Hamiltonian and diffeomorphism constraints to the leading order and that no parity conditions have to be imposed. The vanishing of the leading order of constraints imposes the following relations

$$\bar{\pi}^{rA} = -D_B \bar{\pi}^{BA} \quad (2.22)$$

$$D_A \bar{\pi}^{Ar} = \bar{\pi}^A_A \quad (2.23)$$

$$D_A D_B k^{AB} = D_A D^A k, \quad (2.24)$$

which arise from  $\mathcal{H}^A$ ,  $\mathcal{H}^r$  and  $\mathcal{H}$  respectively.

The transformations preserving the above boundary conditions are

$$\xi^\perp = rb + F + O(r^{-1}), \quad \xi^r = W + O(r^{-1}), \quad \xi^A = Y^A + \frac{1}{r}I^A + O(r^{-2}) \quad (2.25)$$

with

$$D_A D_B b + \gamma_{AB} b = 0, \quad \mathcal{L}_Y \gamma_{AB} = 0, \quad (2.26)$$

where  $b, F, W, Y^A$  are functions on the sphere and

$$I^A = \frac{2b}{\sqrt{\gamma}} \bar{\pi}^{rA} + D^A W \quad (2.27)$$

The vectors  $Y^A$  describe spatial rotations,  $b$  Lorentz boosts,  $f$  contains time translations through its zero mode and  $W$  contains spatial translations through the  $l = 1$  terms in an expansion in spherical harmonics. In order for the parity conditions (2.17) to be preserved as well Henneaux and Troessaert further assume [1]

$$F = -3b\bar{\lambda} - \frac{1}{2}b\bar{h} + T, \quad (2.28)$$

where  $T$  is an even function on the sphere and  $W$  is an odd function. The above defined transformations  $\xi$  form under the bracket

$$[\xi_1, \xi_2]_M = [\xi_1, \xi_2]_{SD} + \delta_2^{h,\pi} \xi_1 - \delta_1^{h,\pi} \xi_2 \quad (2.29)$$

the following algebra

$$\widehat{\xi}(\widehat{Y}, \widehat{b}, \widehat{T}, \widehat{W}) = [\xi_1(Y_1, b_1, T_1, W_1), \xi_2(Y_2, b_2, T_2, W_2)]_M, \quad (2.30)$$

with

$$\widehat{Y}^A = Y_1^B \partial_B Y_2^A + \bar{\gamma}^{AB} b_1 \partial_B b_2 - (1 \leftrightarrow 2) \quad (2.31)$$

$$\widehat{b} = Y_1^B \partial_B b_2 - (1 \leftrightarrow 2) \quad (2.32)$$

$$\widehat{T} = Y_1^A \partial_A T_2 - 3b_1 W_2 - \partial_A b_1 D^A W_2 - b_1 D_A D^A W_2 - (1 \leftrightarrow 2) \quad (2.33)$$

$$\widehat{W} = Y_1^A \partial_A W_2 - b_1 T_2 - (1 \leftrightarrow 2), \quad (2.34)$$

which is shown to be isomorphic to the BMS algebra. In (2.29)  $[\xi_1, \xi_2]_{SD}$  is the surface deformation bracket defined as

$$[\xi_1, \xi_2]_{SD}^\perp = \xi_1^i \partial_i \xi_2^\perp - \xi_2^i \partial_i \xi_1^\perp \quad (2.35)$$

$$[\xi_1, \xi_2]_{SD}^i = \xi_1^k \partial_k \xi_2^i - \xi_2^k \partial_k \xi_1^i + h^{ik} (\xi_1^\perp \partial_k \xi_2^\perp - \xi_2^\perp \partial_k \xi_1^\perp) \quad (2.36)$$

and  $\delta_2^{h,\pi} \xi_1$  is given by

$$\delta_2^{h,\pi} \xi_1 = \frac{\delta \xi_1}{\delta g_{ij}} \delta_{\xi_2} g_{ij} + \frac{\delta \xi_1}{\delta \pi^{ij}} \delta_{\xi_2} \pi^{ij}. \quad (2.37)$$

The terms of the form  $\delta_1^{g,\pi} \xi_2$  appear because  $\xi$  depends on phase-space functions and one therefore has to take into account the change of  $\xi$  induced by the variation of these functions.

The boundary terms given in [1], which correspond to the term  $\int d^2x K$  in (2.7), are

$$\begin{aligned} \mathcal{K}_\xi [\delta g_{ij}, \delta \pi^{ij}] = & \delta \oint d^2x \left\{ -2Y^A \left( \bar{h}_{AB} \bar{\pi}^{rB} + \bar{\gamma}_{AB} \pi^{(2)rB} + \bar{h}_{rA} \bar{\pi}^{rr} \right) \right. \\ & \left. - 2\sqrt{\gamma} b k^{(2)} - \sqrt{\gamma} \frac{1}{4} b (\bar{h}^2 + \bar{h}^{AB} \bar{h}_{AB}) \right\} \\ & + \oint d^2x \left\{ -2I^A \bar{\gamma}_{AB} \delta \bar{\pi}^{rB} - 2W \delta \bar{\pi}^{rr} - \sqrt{\gamma} (2F + \bar{h}b) \delta (2\bar{\lambda} + \bar{D}_A \bar{h}_{rA}) \right. \\ & \left. + \sqrt{\gamma} (\bar{h}^{rC} \partial_C b \bar{\gamma}^{AB} - b \bar{D}^A \bar{h}^{rB}) \delta \bar{h}_{AB} \right\} + o(r^0). \end{aligned} \quad (2.38)$$

It can be seen that the form of  $F$  in (2.28) guarantees integrability of the boundary terms. The expression for the charge, which corresponds to the term  $C$  in (2.8), is finally given by

$$\begin{aligned} \mathcal{B}_\xi [g_{ij}, \pi^{ij}] = & \oint d^2x \left\{ Y^A \left( 4\bar{k}_{AB} \bar{\pi}^{rB} - 4\bar{\lambda} \bar{\gamma}_{AB} \bar{\pi}^{rB} + 2\bar{\gamma}_{AB} \pi^{(2)rB} \right) + W (2\bar{\pi}^{rr} - 2\bar{D}_A \bar{\pi}^{rA}) \right. \\ & \left. + T 4\sqrt{\gamma} \bar{\lambda} + b\sqrt{\gamma} \left( 2k^{(2)} + \bar{k}^2 + \bar{k}_B^A \bar{k}_A^B - 6\bar{\lambda} \bar{k} \right) + b \frac{2}{\sqrt{\gamma}} \bar{\gamma}_{AB} \bar{\pi}^{rA} \bar{\pi}^{rB} \right\} \end{aligned} \quad (2.39)$$

and notice in particular that the charges proportional to higher modes of  $W$  and  $T$  are in general non-vanishing. Supertranslations are therefore part of the asymptotic symmetry for the new set of parity conditions.

To see how the cancellation of the divergences in the boundary terms works consider for instance the divergence proportional to  $Y^A$ , which appears in the term  $\int d^2x K$  in (2.7) (see [1] for details of derivation of the charges)

$$\begin{aligned} \int d^2x K_Y &= -2r \oint d^2x Y^A \gamma_{AB} \delta \bar{\pi}^{rB} + O(1) \\ &= 2r \oint d^2x Y_B D_C \delta \bar{\pi}^{BC} + O(1) \\ &= -2r \oint d^2x D_{(C} Y_{B)} \delta \bar{\pi}^{BC} + O(1), \end{aligned}$$

which vanishes since  $Y^A$  are the Killing vectors on the sphere and thus obey

$$D_{(C} Y_{B)} = 0. \quad (2.40)$$

This means that in order to remove potential  $Y$ -charges divergences we must assume that we cannot extend the rotation sector beyond the standard three rotational Killing vectors.

The aim of this paper is to express, in a first step, the expansions of the spatial metric and conjugate momenta (2.11)-(2.16) in terms of an asymptotically flat spacetime metric. This will be done by means of a 3+1 decomposition and the ADM formalism [15], which we briefly review in the Appendix A. In a second step we substitute these expressions into the symplectic structure (2.21), constraints (2.22)-(2.24) and charges (2.39) and analyze what are the consequences of this procedure for the asymptotic symmetries.



### 3 Conjugate momenta in terms of spacetime metric components

In this section we are going to perform a 3+1 decomposition of a spacetime metric and use the ADM formalism to express the momenta in terms of components of this metric and their derivatives. We consider a metric that is asymptotically flat at null infinity and is defined by the following expansion

$$\begin{aligned}
g_{\mu\nu}dx^\mu dx^\nu = & - \left( 1 - \frac{2M}{r} - \frac{\bar{g}_{uu}}{r^2} + O(r^{-3}) \right) du^2 - 2 \left( 1 - \frac{\bar{g}_{ur}}{r} - \frac{g_{ur}^{(2)}}{r^2} + O(r^{-3}) \right) dudr \\
& + \left( \psi_A + \frac{1}{r}F_A + O(r^{-2}) \right) dudx^A \\
& + \left( r^2\gamma_{AB} + rC_{AB} + D_{AB} + O(r^{-1}) \right) dx^A dx^B,
\end{aligned} \tag{3.1}$$

where  $\gamma_{AB}$  is the unit metric on the sphere and all other metric components are functions of  $(u, x^A)$ . This metric is subject only to the partial Bondi gauge condition

$$g_{rr} = 0, \quad g_{rA} = 0 \tag{3.2}$$

and no further assumptions are made at this stage.

The metric (3.1) is more general than the Bondi metric which is usually considered for the investigation of asymptotic symmetries at null infinity [4–6], [10], [14]. Usually the Bondi metric is assumed to additionally satisfy the Einstein field equations and to be subject to the determinant condition

$$\det g_{AB} = r^4 \det \gamma_{AB}, \tag{3.3}$$

which implies

$$\gamma^{AB}C_{AB} = 0, \quad \gamma^{AB}D_{AB} = \frac{1}{2}C^{AB}C_{AB}. \tag{3.4}$$

Here we instead start with an off-shell metric and the field equations will be partially imposed by demanding that the leading order of the Hamiltonian and diffeomorphism constraint has to vanish. We also are not imposing the determinant condition since it leads to a metric that is too rigid: in subsection 6.2 we will demonstrate that imposing this condition would drastically reduce the asymptotic symmetry by excluding spatial (super) translations.

We choose spacelike hypersurfaces  $\Sigma_t$  of constant ‘time’  $t$  defined to be  $t = u + r +$

$f(x^A) + \frac{g(x^A)}{r}$  and in coordinates  $(t, r, x^A)$  the metric (3.1) takes the form

$$\begin{aligned}
g_{\mu\nu}dx^\mu dx^\nu = & - \left( 1 - \frac{2M}{r} - \frac{\bar{g}_{uu}}{r^2} + O(r^{-3}) \right) dt^2 + 2 \left( \frac{\bar{g}_{ur} - 2M}{r} + O(r^{-2}) \right) dt dr \\
& + 2 \left( \partial_A f + \frac{1}{2} \psi_A + O(r^{-1}) \right) dt dx^A \\
& + \left( 1 + \frac{2M - 2\bar{g}_{ur}}{r} + \frac{\bar{g}_{uu} - 2g_{ur}^{(2)}}{r^2} + O(r^{-3}) \right) dr^2 \\
& + \left( -\psi_A + \frac{(4M - 2\bar{g}_{ur})\partial_A f - F_A}{2r} + O(r^{-2}) \right) dr dx^A \\
& + (r^2 \gamma_{AB} + r C_{AB} + D_{AB} - \partial_A f \partial_B f - \psi_A \partial_B f) dx^A dx^B. \tag{3.5}
\end{aligned}$$

By comparing the form of (3.5) with the decomposition (A.4) we find the following expressions for lapse and shift

$$N = 1 - \frac{M}{r} + O(r^{-2}), \quad N_r = \frac{\bar{g}_{ur} - 2M}{r} + O(r^{-2}), \quad N_A = \partial_A f + \frac{1}{2} \psi_A + O(r^{-1}) \tag{3.6}$$

and we identify the leading order terms in the metric expansion (2.11)-(2.13) as

$$\begin{aligned}
\bar{h}_{rr} = 2M - 2\bar{g}_{ur}, \quad \bar{h}_{rA} = -\frac{\psi_A}{2}, \quad \bar{h}_{AB} = C_{AB} \\
h_{rr}^{(2)} = \bar{g}_{uu} - 2g_{ur}^{(2)}, \quad h_{rA}^{(2)} = \frac{4\partial_A f - F_A}{2}, \quad h_{AB}^{(2)} = D_{AB} - \partial_A f \partial_B f - \psi_A \partial_B f. \tag{3.7}
\end{aligned}$$

The condition (2.20) now takes the form

$$\psi_A = 0, \tag{3.8}$$

and following the arguments of [1] we will eventually also make this assumption. For the sake of obtaining a general form for the expressions of the momenta, however, we will keep  $\psi_A$  non-zero for now and assume that it is vanishing from Sect. 5 on.

Notice that the choice  $t = u + r + f(x^A) + \frac{g(x^A)}{r}$  is the most general one that is compatible with the fall-off conditions (2.11)-(2.13). Choosing  $t = u + rk(x^A) + O(1)$  would lead to

$$h_{ab}dy^a dy^b = dr^2(2k - k^2 + O(r^{-1})) + .. \tag{3.9}$$

which does not agree with the falloff condition (2.11).

The unit normal on the spacelike hypersurfaces is given by

$$n_\alpha = -N \partial_\alpha t = - \left( 1 - \frac{M}{r} + O(r^{-2}) \right) \partial_\alpha \left( u + r + f(x^A) + \frac{g(x^A)}{r} \right) \tag{3.10}$$

and using this expression to evaluate extrinsic curvature (A.8) we find the asymptotic expressions

$$K_{rr} = -\frac{\partial_u M}{r} + O(r^{-2}), \quad (3.11)$$

$$K_{rA} = -\frac{1}{4r} \left( -2\partial_A \bar{g}_{ur} + 4\partial_A M - 2(\psi_A + 2\partial_A f) \right. \\ \left. - (\psi_B + 2\partial_B f)\gamma^{BC}\partial_u C_{CA} - 4\partial_A f\partial_u M \right) + O(r^{-2}) \quad (3.12)$$

$$K_{AB} = \frac{1}{2}\partial_u C_{AB} + O(r^{-2}). \quad (3.13)$$

We have now all the expressions at hand we need to write the momenta in terms of the components of (3.1) and find from evaluating (A.1)

$$\pi^{rr} = -\frac{r}{2}\sqrt{\gamma}\gamma^{AB}\partial_u C_{AB} + \frac{1}{2}\sqrt{\gamma}(4\mathcal{M} - \gamma^{AB}\partial_u D_{AB} \\ + \gamma^{AB}D_A(2D_B f + \psi_B) + G[\partial_u C_{AB}]) + O(r^{-1}) \quad (3.14)$$

$$\pi^{rA} = \frac{1}{4r}\sqrt{\gamma}(-2\gamma^{AB}D_B \mathcal{M} + 2\gamma^{AB}(2D_B f + \psi_B) + G[\partial_u M, \partial_u C_{AB}]) + O(r^{-2}) \quad (3.15)$$

$$\pi^{AB} = \frac{1}{2r}\sqrt{\gamma}(2\gamma^{AB}\partial_u M - (\gamma^{AB}\gamma^{CD}\partial_u C_{CD} - \partial_u C^{AB})) \\ + \frac{1}{2r^2}\sqrt{\gamma}\left[\gamma^{AB}(\partial_u \bar{g}_{uu} - 2\mathcal{M}\partial_u \bar{g}_{ur}) - (\gamma^{AB}\gamma^{CD}\partial_u D_{CD} - \partial_u D^{AB}) \right. \\ \left. - \frac{\gamma^{AB}}{2}(\gamma^{CD}(2D_C f + \psi_C)\partial_u \psi_D) - (2D^A D^B f + D^{(A}\psi^{B)}) \right. \\ \left. + \gamma^{CD}(2D_C D_D f + D_C \psi_D)\gamma^{AB} + G[\partial_u M, \partial_u C_{AB}]\right] + O(r^{-3}), \quad (3.16)$$

where

$$\mathcal{M} = \bar{g}_{ur} - 2M \quad (3.17)$$

and  $G$  stands for lengthy terms proportional to either  $\partial_u C_{AB}$  or  $\partial_u M$ . Notice that the leading order of  $\pi^{rr}$  and  $\pi^{AB}$  is of lower order than what is demanded by the falloff conditions (2.14) and (2.16). In order to get rid off these unwanted contributions we require

$$\partial_u C_{AB} \sim u^{-(1+\varepsilon)}, \quad \partial_u M \sim u^{-(1+\varepsilon)}, \quad \varepsilon > 0, \quad (3.18)$$

which provides an extra damping factor since the above expansion of the momenta is defined on a hypersurface  $t = \text{const.}$  in the limit  $r \rightarrow \infty$  which implies  $u \rightarrow -\infty$ . The terms denoted with  $G$  are therefore also suppressed. The conditions (3.18) imply that there is no gravitational radiation at the very distant past of null infinity  $\mathcal{I}_-^+$ , which is a natural physical requirement.

#### 4 Finiteness of symplectic structure

The falloff conditions defined in (2.11)-(2.16) are not sufficient to remove divergences in the symplectic structure

$$\int d^3x \pi^{ab} \dot{h}_{ab}, \quad (4.1)$$

since terms of order  $O(r^{-1})$  appear which are logarithmically divergent. As was explained above, after eq. (2.21), the authors of [1] remove these divergences by introducing parity conditions on the leading order terms in the expansion of the metric and momenta. However, as we are going to show now, when expressing (4.1) in terms of the spacetime metric (3.1) and its derivatives a divergent term appears which cannot be cancelled by any choice of parity conditions. Instead, here we assume the damping condition (3.18) and we show that an appropriate choice of function  $f(x^A)$  defining the foliation suffices to remove all the divergences.

Indeed, the terms which are potentially divergent are the following ones

$$\int dr \frac{1}{r} \int d\phi d\theta \left( \bar{\pi}^{rr} \dot{h}_{rr} + \bar{\pi}^{AB} \dot{h}_{AB} + \bar{\pi}^{rA} \dot{h}_{rA} \right) \quad (4.2)$$

and using the definition of  $\dot{h}_{ab}$  (A.9) we find the expressions

$$\dot{h}_{rr} = -2\partial_u M, \quad \dot{h}_{rA} = 0, \quad (4.3)$$

$$\dot{h}_{AB} = \partial_u C_{AB} + 2(D_A D_B f + \gamma_{AB} \mathcal{M}) + D_{(A} \psi_{B)}. \quad (4.4)$$

The first thing to notice is that the term  $\bar{\pi}^{rr} \dot{h}_{rr}$  contains a logarithmic divergence proportional to  $M \partial_u M$ . Since the product of two odd functions is an even function and since such a function is in general non-vanishing when integrated over the sphere this term can not be canceled by introducing parity conditions. We therefore do not rely on parity conditions to make the symplectic structure finite but observe that the damping condition (3.18) together with the equation

$$D_A D_B f + \gamma_{AB} \mathcal{M} + \frac{1}{2} D_{(A} \psi_{B)} = 0 \quad (4.5)$$

is sufficient to guarantee its finiteness. Since  $f(x^A)$  defines the foliation of spacetime this can be considered a mild condition since it does not constrain the class of spacetimes which we allow, as long as there exists a solution for  $f$ . Since  $f$  is a function of  $x^A$  only, equation (4.5) can only be fulfilled if

$$\partial_u (\gamma_{AB} \mathcal{M} + \frac{1}{2} D_{(A} \psi_{B)}) = 0 \quad (4.6)$$

and we therefore assume this as an additional boundary condition.

There is an important case in which (4.5) does not have a solution for  $f$ , namely  $\mathcal{M} = \mathcal{M}_c = \text{const}$ ,  $\psi_A = 0$  which includes the Schwarzschild metric. All functions referring to this special case will be denoted with an index  $c$ . A solution would require

$$D_A D_B f_c = \mathcal{M}_c \gamma_{AB} \rightarrow D^2 f_c = 2\mathcal{M}_c, \quad (4.7)$$

but such a function does not exist since  $D^2 Y_{l,m} = -l(l+1)Y_{l,m}$  and  $D^2$  therefore annihilates the  $l=0$  mode. In this case we thus have to demand instead  $\bar{\pi}_c^{AB} = 0$  so that the divergence in (4.2) still gets canceled, where we will and from this it follows that  $f_c$  has to fulfill

$$-2D^A D^B f_c + 2D^2 f_c \gamma^{AB} = \partial_u P_c^{AB} \quad (4.8)$$

which can only have a solution if

$$\partial_u^2 P_c^{AB} = 0 \quad (4.9)$$

and where

$$P^{AB} = \sqrt{\gamma}\gamma^{AB}(\bar{g}_{uu} - 2\mathcal{M}\bar{g}_{ur} - \gamma^{CD}D_{CD}) + \sqrt{\gamma}D^{AB}. \quad (4.10)$$

For the Schwarzschild case  $\partial_u P_c^{AB} = 0$  and (4.8) is solved simply by  $f_c = \text{const.}$

## 5 Leading order of constraints

As was explained in subsection 2.1 the vanishing of the leading order of the Hamiltonian (2.24) and diffeomorphism constraint (2.22), (2.23) is crucial for canceling divergences which arise in the expression of the charges. In this section we show which restrictions on the form of the momenta it implies. We are also going to assume  $\psi_A = 0$  from now on.

We need to distinguish here two different cases, which were discussed at the end of section 4. First, consider spacetimes with  $\mathcal{M} = \mathcal{M}_c = \text{const.}$  in which case  $f_c$  has to fulfill (4.8) and  $\bar{\pi}_c^{AB} = 0$ . We find from (2.22) and (3.15)

$$\bar{\pi}_c^{rA} = 0 \rightarrow f_c = \text{const.} \quad (5.1)$$

and (4.8) then implies

$$\partial_u P_c^{AB} = 0. \quad (5.2)$$

With this condition the next constraint (2.23) is fulfilled as well and there is another condition coming from (2.24) which introduces a relation between  $C_{AB}$  and  $\bar{h}_{rr}$

$$D_A D_B C^{AB} - D^2 \gamma^{AB} C_{AB} = D^2 \bar{h}_{rr}, \quad (5.3)$$

where this relation also holds for the case of general  $\mathcal{M}$ .

Second, spacetimes where  $\mathcal{M}$  is allowed to be a general function on the sphere. First notice that from (4.6) it follows that

$$\partial_u \mathcal{M} = 0. \quad (5.4)$$

From (2.22) we find the condition

$$\frac{\sqrt{\gamma}}{4} \gamma^{AB} D_B (4f - 2\mathcal{M}) = -D_B \bar{\pi}^{AB} \quad (5.5)$$

which implies

$$\frac{\sqrt{\gamma}}{4} \gamma^{AB} (4f - 2\mathcal{M}) = -\bar{\pi}^{AB} + A \gamma^{AB} \sqrt{\gamma}, \quad (5.6)$$

with  $A$  being an arbitrary constant. Substituting this expression for  $\bar{\pi}^{AB}$  into (2.23) yields

$$(D^2 + 2)(4f - 2\mathcal{M}) = 8A \quad (5.7)$$

and when we now use that from (4.5) it follows that  $D^2 f = -2\mathcal{M}$  we have

$$(D^2 + 2)(D^2 + 4)f(x^A) = 8A. \quad (5.8)$$

Upon expanding  $f$  in spherical harmonics  $Y_{lm}$  satisfying  $D^2 Y_{lm} = -l(l+1)Y_{lm}$  we find that this equation has a general solution of the form

$$f = A + \sum_{m=-1}^1 a_m Y_{1m} \quad (5.9)$$

with  $a_m$  being arbitrary constants. Then

$$\mathcal{M} = -\frac{1}{2}D^2 f = \sum_{m=-1}^1 a_m Y_{1m}, \quad (5.10)$$

which means that the vanishing of the leading order of constraints demands  $\mathcal{M}$  to consist solely of  $l = 1$  modes. Substituting this to (5.6) we find that  $\bar{\pi}^{AB}$  is required to take the form

$$\bar{\pi}^{AB} = -\frac{\sqrt{\gamma}}{2}\gamma^{AB}\mathcal{M} \quad (5.11)$$

and comparing this with the expression for  $\bar{\pi}^{AB}$  in (3.16)

$$\bar{\pi}^{AB} = -\sqrt{\gamma}\gamma^{AB}\mathcal{M} + \partial_u P^{AB}, \quad (5.12)$$

with  $P^{AB}$  given by

$$P^{AB} = \sqrt{\gamma}\gamma^{AB}(\bar{g}_{uu} - 2\mathcal{M}\bar{g}_{ur} - \gamma^{CD}D_{CD}) + \sqrt{\gamma}D^{AB} \quad (5.13)$$

we find that for consistency we need to demand that

$$\partial_u P^{AB} = \frac{\sqrt{\gamma}}{2}\gamma^{AB}\mathcal{M}. \quad (5.14)$$

To summarize, in the case  $\mathcal{M} = const.$  the expressions for the momenta now read

$$\bar{\pi}_c^{rr} = \frac{\sqrt{\gamma}}{2}(4\mathcal{M}_c - \gamma^{AB}\partial_u D_{AB}) \quad (5.15)$$

$$\bar{\pi}_c^{rA} = \bar{\pi}_c^{AB} = 0 \quad (5.16)$$

$$\begin{aligned} \pi_c^{(2)rA} = \frac{\sqrt{\gamma}}{4} & \left( 3F^A + 8D^A g - 2D^A g_{ur}^{(2)} + 2D^A \bar{g}_{uu} - 4MD^A \bar{g}_{ur} \right. \\ & \left. + 16MD^A M - 2\bar{g}_{ur}D^A \bar{g}_{ur} \right) \end{aligned} \quad (5.17)$$

$$(5.18)$$

and otherwise  $\mathcal{M} = \sum_{m=-1}^1 a_m Y_{1m}$  and momenta are given by the following expressions

$$\bar{\pi}^{rr} = \frac{\sqrt{\gamma}}{2} (2\mathcal{M} - \gamma^{AB} \partial_u D_{AB}) \quad (5.19)$$

$$\bar{\pi}^{rA} = \frac{\sqrt{\gamma}}{2} \gamma^{AB} D_B \mathcal{M} \quad (5.20)$$

$$\begin{aligned} \pi^{(2)rA} = \frac{\sqrt{\gamma}}{4} & \left( D^A \mathcal{M} \gamma^{AB} C_{AB} - 4C^{AB} D_B \mathcal{M} + 3F^A + 8D^A g - 2D^A g_{ur}^{(2)} \right. \\ & \left. + 2D^A \bar{g}_{uu} - 2D^A \mathcal{M} \partial_u \bar{g}_{uu} - 4MD^A \bar{g}_{ur} + 16MD^A M - 2\bar{g}_{ur} D^A \bar{g}_{ur} \right) \end{aligned} \quad (5.21)$$

$$\bar{\pi}^{AB} = -\frac{\sqrt{\gamma}}{2} \gamma^{AB} \mathcal{M}, \quad (5.22)$$

where conditions (3.8) and (4.5) have been used.

## 6 Asymptotic symmetries

In this section we discuss the asymptotic symmetries of  $h_{ab}$  and  $\pi^{ab}$ , in particular we will analyze which transformations preserve the falloff conditions (2.11)-(2.16) and the gauge condition (2.20). This will reproduce the expressions (2.25) and (2.26) which were already given in [1]. We are giving here the details of this derivation to stress the fact that, as we are going to show, the preservation of the falloff conditions allows for a large group of supertranslations parametrized by two arbitrary functions on the sphere. These details are furthermore needed for our discussion of the determinant condition (3.3) in subsection 6.2.

To this end we will evaluate the change in the canonical variables generated by the Hamiltonian  $H = \int d^3x (\xi^\perp \mathcal{H} + \xi^a \mathcal{H}_a)$  which is given by, see [15]

$$\delta_\xi h_{ab} = 2\xi^\perp h^{-1/2} \left( \pi_{ab} - \frac{1}{2} h_{ab} \pi \right) + \mathcal{L}_\xi h_{ab}. \quad (6.1)$$

$$\begin{aligned} \delta \pi^{ab} = & -\xi^\perp h^{\frac{1}{2}} \left( R^{ab} - \frac{1}{2} h^{ab} R \right) + \frac{1}{2} \xi^\perp h^{ab} h^{-\frac{1}{2}} \left( \pi_{mn} \pi^{mn} - \frac{1}{2} \pi^2 \right) \\ & - 2\xi^\perp h^{-\frac{1}{2}} \left( \pi^{am} \pi_m^b - \frac{1}{2} \pi^{ab} \pi \right) + h^{\frac{1}{2}} \left( \xi^\perp |^{ab} - h^{ab} \xi^\perp |^m \right) \\ & + \left( \pi^{ab} \xi^m \right) |^m - \xi^a |^m \pi^{mb} - \xi^b |^m \pi^{am}, \end{aligned} \quad (6.2)$$

where  $\mathcal{L}_\xi h_{ab}$  is the Lie derivative

$$\mathcal{L}_\xi h_{ab} = \xi_{a|b} + \xi_{b|a}. \quad (6.3)$$

### 6.1 Preservation of falloff and gauge conditions

From the preservation of the falloff conditions (2.11)-(2.13) we obtain the demands

$$\delta h_{rr} = O(r^{-1}), \quad \delta h_{rA} = O(1), \quad \delta h_{AB} = O(r) \quad (6.4)$$

and we now want to find such  $\xi^\perp, \xi^a$  that the change of  $h_{ab}$  defined in (6.1) obeys these demands.

Using the expansion of Christoffel symbols associated with  $h_{ab}$

$$\Gamma_{AB}^r = -r\bar{\gamma}_{AB} + O(1) \quad (6.5)$$

$$\Gamma_{BC}^A = \bar{\Gamma}_{BC}^A + O(r^{-1}) \quad (6.6)$$

$$\Gamma_{rA}^r = \frac{1}{2r} (\partial_A \bar{h}_{rr} + \psi_A) + O(r^{-2}) \quad (6.7)$$

$$\Gamma_{rr}^r = -\frac{\bar{h}_{rr}}{2r^2} + O(r^{-3}) \quad (6.8)$$

$$\Gamma_{rB}^A = \frac{1}{r} \delta_B^A + O(r^{-2}) \quad (6.9)$$

$$\Gamma_{rr}^A = -\frac{\bar{\gamma}^{AB} \partial_B \bar{h}_{rr}}{2r^3} + O(r^{-3}), \quad (6.10)$$

we find the following transformation of  $h_{rr}$

$$\delta h_{rr} = \frac{2\xi^\perp}{\sqrt{\gamma}r^2} \left( \bar{\pi}_{rr} - \frac{1}{2} \bar{h}_{rr} \bar{\pi} \right) + \xi^A \partial_A \frac{\bar{h}_{rr}}{r} + 2\partial_r \xi^r - 2\psi_A \partial_r \xi^A + (\text{subleading}), \quad (6.11)$$

where

$$\bar{\pi} = \bar{\pi}^{rr} + \gamma_{AB} \bar{\pi}^{AB}. \quad (6.12)$$

Comparing this with (6.4) we find that a large  $r$  expansion of  $\xi^\perp, \xi^a$  has to be of the form

$$\xi^\perp = rb + F + O(r^{-1}), \quad \xi^A = Y^A + \frac{1}{r} I^A + O(r^{-2}), \quad \xi^r = W + \frac{1}{r} \xi_1^r + O(r^{-2}), \quad (6.13)$$

where all terms in this expansion are functions on the sphere.

The transformation of  $h_{rA}$

$$\delta h_{rA} = \frac{2b}{\sqrt{\gamma}} \bar{\pi}_{rA} - \frac{1}{2} Y^B D_B \psi_A + \partial_A W - \frac{1}{2} \psi_B D_A Y_0^B - \gamma_{AB} I^B + (\text{subleading}) \quad (6.14)$$

does not give any further restrictions on  $\xi^\perp, \xi^a$ .

To fulfill the demand that  $\delta h_{AB}$

$$\begin{aligned} \delta h_{AB} = r^2 D_{(A} Y_{B)} + r \left( 2W \gamma_{AB} + Y^C D_C C_{AB} + \psi_{(B} \gamma_{A)C} Y^C + 2C_{C(B} D_{A)} Y^C \right. \\ \left. + 2\gamma_{C(B} D_{A)} I^C + \frac{2b}{\sqrt{\gamma}} \left( \bar{\pi}_{AB} - \frac{1}{2} \gamma_{AB} \bar{\pi} \right) \right) \end{aligned} \quad (6.15)$$

is of order  $O(r)$  we have to assume that  $Y^B$  are the Killing vectors on the 2-sphere

$$D_{(A} Y_{B)} = 0. \quad (6.16)$$

So far there are no restrictions on  $b$  but it is fixed from the preservation of the asymptotic form of momenta which demands

$$\delta \pi^{rr} = O(1), \quad \delta \pi^{rA} = O(r^{-1}), \quad \delta \pi^{AB} = O(r^{-2}). \quad (6.17)$$



One can check that there are no new restrictions coming from the first two conditions in (6.17). The third one, however, does lead to a new restriction and reads

$$\begin{aligned}\delta\pi^{AB} &= r^2\sqrt{\gamma}\left(\nabla^A\nabla^B\xi^\perp - h^{AB}\nabla^i\nabla_i\xi^\perp\right) + (\text{subleading}) \\ &= \frac{\sqrt{\gamma}}{r^2}\left(\gamma^{AC}\gamma^{AD}\nabla_C\nabla_D\xi^\perp - \gamma^{AB}\gamma^{CD}\nabla_C\nabla_D\xi^\perp\right) + (\text{subleading}).\end{aligned}\quad (6.18)$$

Now consider (we use the notation  $\nabla_A = D_A + (\text{subleading})$ )

$$\begin{aligned}\nabla_C\nabla_D\xi^\perp &= r\partial_C\partial_D b - \Gamma_{CD}^r\partial_r(rb) - r\Gamma_{CD}^A\partial_A b + O(1) \\ &= rD_C D_B b + r\gamma_{CD}b,\end{aligned}\quad (6.19)$$

where we used (6.5) and plugging this back into the previous equation we obtain

$$\delta\pi^{AB} = \frac{1}{r}\left(D^A D^B b - \gamma^{AB}D^2 b - \gamma^{AB}b\right) + O(r^{-2}).\quad (6.20)$$

Vanishing of the leading order therefore imposes the condition

$$D_A D_B b - \gamma_{AB}D^2 b - \gamma_{AB}b = 0 \Rightarrow D^2 b = -2b\quad (6.21)$$

and therefore we find that  $b$  has to fulfill the condition

$$D_A D_B b + \gamma_{AB}b = 0,\quad (6.22)$$

whose only solution is  $b$  being a linear combination of three  $l = 1$  harmonics with constant coefficient.

Now we consider the preservation of the gauge condition  $\bar{h}_{rA} = 0$ . From (6.14) one can directly see that in order to have  $\delta\bar{h}_{rA} = 0$  we need to assume

$$I^A = D^A W + \frac{2b}{\sqrt{\gamma}}\bar{\pi}^{rA},\quad (6.23)$$

which means that the preservation of the gauge choice

$$\bar{h}_{rA} = -\frac{1}{2}\psi_A = 0,\quad (6.24)$$

which we are going to adapt from now on, determines the subleading term of  $\xi^A$ .

To summarize,  $F$  and  $W$  are not constrained by the boundary conditions and are associated with angle-dependent translations, temporal and spatial ones, respectively.  $Y^A$  are the three Killing vectors on the sphere parametrizing rotations and  $b$  contains only  $l = 1$  harmonics and parametrizes three boosts. Except for the assumptions of parity on  $W$  and  $F$  we have therefore reproduced the asymptotic symmetry transformations given in [1] and the corresponding surface terms are therefore identical with (2.38), which has been derived for  $F, W$  having no definite parity. Integrability of these surface terms demands that  $F$  is of the form  $F = -\frac{1}{2}\bar{h}b + T(x^A)$ , where  $T(x^A)$  is a general function on the sphere. Any function of  $\bar{\lambda}$  could be added to  $F$  without spoiling integrability, which introduces an ambiguity in the expression of the charges. We choose for  $F$  the form (2.28).

In [1] it is shown that transformations with  $W$ =odd,  $T$ =even form an algebra isomorphic to the BMS algebra found at null infinity. The odd and even functions combine to give the single function parametrizing supertranslations at null infinity. Since we do not involve any parity conditions it appears that the resulting asymptotic symmetry is larger than the BMS symmetry, as long as the associated charges are finite. Before proceeding with the discussion of the charges, however, we turn to the aforementioned determinant condition.

## 6.2 Determinant condition and symmetry reduction

In the previous subsection we have found that a large group of supertranslations and Lorentz transformations preserve the falloff conditions (2.11)-(2.16). We are now going to demonstrate that additionally imposing the determinant condition (3.3) breaks the invariance under all spatial translations, including the Poincaré ones.

The determinant condition implies

$$\gamma^{AB}C_{AB} = 0 \tag{6.25}$$

and transformations preserving this condition must fulfill

$$\delta(\gamma^{AB}C_{AB}) = \gamma^{AB}\delta\bar{h}_{AB} = 0, \tag{6.26}$$

where we have used that  $\delta\gamma_{AB} = 0$  and  $C_{AB} = \bar{h}_{AB}$ . Substituting (6.15) we find

$$\gamma^{AB}\delta\bar{h}_{AB} = 2(D^2 + 2)W + \frac{4}{\sqrt{\gamma}}D_A(b\bar{\pi}^{rA}) - \frac{2b}{\sqrt{\gamma}}\bar{\pi}^{rr} = 0 \tag{6.27}$$

and for the case  $\mathcal{M} = \text{const.}$  we plug in the expressions for the momenta (5.15) and (5.16) and obtain

$$2(D^2 + 2)W = b(4\mathcal{M}_c - \gamma^{AB}\partial_u D_{AB}). \tag{6.28}$$

This equation has no solution for  $W$  since the RHS is in general non-vanishing and contains  $l = 1$  harmonics<sup>2</sup> which can not be produced by the LHS.

The other case we need to consider is  $\mathcal{M} = \sum_{m=-1}^1 a_m Y_{1m}$  and substituting the momenta (5.19) and (5.20) into (6.27) yields

$$\begin{aligned} 2(D^2 + 2)W &= -2D_A(bD^A\mathcal{M}) + b(2\mathcal{M} - \gamma^{AB}\partial_u D_{AB}) \\ &= -2D_A bD^A\mathcal{M} + 6b\mathcal{M} - b\gamma^{AB}\partial_u D_{AB} \end{aligned} \tag{6.29}$$

and by the same argument as in the first case we see that this equation has no solution for  $W$ .

We can therefore conclude that spatial translations, by which we mean standard Poincaré ones and supertranslations, do not preserve the condition (6.25), which shows that imposing the determinant condition excludes spatial translations from the group of asymptotic symmetries.

---

<sup>2</sup>In fact  $b$  purely consists of  $l = 1$  harmonics.

This result is in contrast with the situation at null infinity. The transformation of  $C_{AB}$  there is given by (see for instance section 2.2 in [9] for details concerning the asymptotic symmetries at null infinity)

$$-\delta C_{AB} = \left[ f\partial_u + \mathcal{L}_R - \frac{1}{2}D_A R^A \right] C_{AB} - 2D_A D_B f + D_A D^A f \gamma_{AB}, \quad (6.30)$$

where  $R^A = Y^A - D^A b$  and  $f = T + \frac{1}{2}u D_A R^A$  parametrize Lorentz transformations and supertranslations which leave the asymptotic form of the Bondi metric at null infinity invariant. One can recognize from this expression that the condition  $\gamma^{AB} C_{AB} = 0$  is automatically preserved, since  $\gamma^{AB} \delta C_{AB} = 0$  under this condition. We therefore find that the symmetry algebra at spatial infinity is not obtained as the continuous limit  $u \rightarrow -\infty$  of the one at null infinity. This is also reflected by the fact that the asymptotic symmetry transformations at null infinity themselves are in fact divergent in that limit, as is exemplified by the  $u$ -component of these transformations,  $\xi^u = f$ .

### 6.3 The charges

In the previous subsections we found that the asymptotic symmetry appears to be larger than the BMS symmetry at null infinity because there are two arbitrary functions on the sphere parametrizing translations. In the treatment of Henneaux and Troessaert [1] the charges associated with transformations outside of BMS are vanishing due to the imposed parity conditions and these transformations are therefore pure gauge. Here we show that for our boundary conditions all the charges associated with  $T$  and  $W$  are in fact non-vanishing. To this end we can use the expression (2.39) derived for general  $T, W$ , which corresponds to the case that we are considering. We then only need to substitute the expressions for  $h_{ab}$  and  $\pi^{ab}$  in terms of (3.1), which we have already obtained from the 3+1 decomposition. From (2.39) the charges associated with translations are given in [1] as

$$\mathcal{B}_{T,W} = \oint d^2x \{ 2T \sqrt{\gamma} \bar{h}_{rr} + 2W (\bar{\pi}^{rr} - \bar{\pi}_A^A) \}. \quad (6.31)$$

For completeness we will also give the expressions for boost charges

$$\mathcal{B}_b = \oint d^2x \left[ b \sqrt{\gamma} \left( 2k^{(2)} + \bar{k}^2 + \bar{k}_B^A \bar{k}_A^B - 6\bar{\lambda} \bar{k} \right) + b \frac{2}{\sqrt{\gamma}} \bar{\gamma}_{AB} \bar{\pi}^{rA} \bar{\pi}^{rB} \right] \quad (6.32)$$

and rotational charges

$$\mathcal{B}_Y = \oint d^2x Y^A \left( 4\bar{k}_{AB} \bar{\pi}^{rB} - 4\bar{\lambda} \bar{\gamma}_{AB} \bar{\pi}^{rB} + 2\bar{\gamma}_{AB} \pi^{(2)rB} \right), \quad (6.33)$$

where  $k^{(2)}$  is defined via the expansion

$$K_B^A = h^{AC} K_{BC} = -\frac{1}{r} \delta_B^A + \frac{1}{r^2} \bar{k}_B^A + \frac{1}{r^3} k_B^{(2)A} + O(r^{-3}) \quad (6.34)$$

$$K_{AB} = \frac{1}{2\lambda} (-\partial_r h_{AB} + \nabla_A h_{rB} + \nabla_B h_{rA}) \quad (6.35)$$

$$\lambda = \frac{1}{\sqrt{h^{rr}}}. \quad (6.36)$$

Substituting the expressions for the momenta in the case  $\mathcal{M} = \text{const.}$  (5.15),(5.16) or for  $\mathcal{M} = \sum_{m=-1}^1 a_m Y_{1m}$  (5.19), (5.22) into (6.31) yields for the translational charge the same expression

$$\mathcal{B}_{W,T} = \oint d^2x \sqrt{\gamma} \{ T 2 \bar{h}_{rr} + W (4\mathcal{M} - \gamma^{AB} \partial_u D_{AB}) \}. \quad (6.37)$$

Upon expanding  $W$  in spherical harmonics we see that the term proportional to  $W\mathcal{M}$  is finite only for the  $l = 0$  or  $l = 1$  modes of  $W$  due to the orthonormality of the harmonics and all other modes are vanishing. The two other terms have in general finite contributions for all modes of  $T$  and  $W$  because we did not have to constrain the functions  $\bar{h}_{rr}$  and  $D_{AB}$  in any way. We can therefore see that indeed all the charges associated with supertranslations are non-vanishing.

For the rotational charge (6.33) we find upon substituting the expressions for the momenta

$$\begin{aligned} \mathcal{B}_Y = \oint d^2x \sqrt{\gamma} Y^A & \left( 2D_A \gamma^{AB} C_{AB} \mathcal{M} - C_{AB} D^B \mathcal{M} + \frac{3}{2} F_A + 4D_A g - D_A g_{ur}^{(2)} \right. \\ & \left. + D_A \bar{g}_{uu} - D_A \mathcal{M} \partial_u \bar{g}_{uu} - 2M D_A \bar{g}_{ur} + 4D_A M^2 - \frac{1}{2} D_A \bar{g}_{ur}^2 \right), \end{aligned} \quad (6.38)$$

where in the case  $\mathcal{M} = \text{const.}$  one just has to replace  $\mathcal{M}$  with  $\mathcal{M}_c$ . To obtain the expression for the boost charge (6.32) we first evaluate (6.35) and (6.36) and find

$$K_{AB} = -2r \gamma_{AB} - C_{AB} + \frac{2}{r} D_{(A} [2\mathcal{M} D_{B)} \mathcal{M} - \frac{1}{2} F_{B)}] + O(r^{-2}) \quad (6.39)$$

and

$$\frac{1}{\lambda} = 1 - \frac{\bar{h}_{rr}}{r} + \frac{L}{r^2} + O(r^{-3}), \quad (6.40)$$

where

$$L = -4\bar{g}_{uu} + 12(\mathcal{M} \bar{g}_{ur} + M^2) + 8g_{ur}^{(2)}. \quad (6.41)$$

To calculate  $k^{(2)}$  from (6.34) we also need

$$h^{AB} = \frac{1}{r^2} \gamma^{AB} - \frac{1}{r^3} C^{AB} + \frac{1}{r^4} (C_D^A C^{DB} - D^{AB} + D^A \mathcal{M} D^B \mathcal{M}) + O(r^{-5}) \quad (6.42)$$

and finally find for the boost charge

$$\begin{aligned} \mathcal{B}_b = \oint d^2x \sqrt{\gamma} b & \left( -D_A \mathcal{M} D^A \mathcal{M} - \frac{1}{2} D^2 \mathcal{M} - \frac{1}{2} D_A F^A - 2L - \frac{3}{4} C^{AB} C_{AB} \right. \\ & \left. - \bar{h}_{rr} \gamma^{AB} C_{AB} + \frac{3}{2} \bar{h}_{rr}^2 + \frac{1}{4} (\gamma^{AB} C_{AB})^2 + 2\gamma^{AB} D_{AB} \right). \end{aligned} \quad (6.43)$$

#### 6.4 Discussion of the asymptotic symmetry

From our discussion of the charges it has become clear that the asymptotic symmetry we find at spatial infinity is larger than the BMS symmetry. The crucial difference to the results of [1] is that the charges associated with even  $W$  and odd  $T$  are non-vanishing.

A larger-than-BMS asymptotic symmetry at spatial infinity has been found previously by Ashtekar and Hansen [17]. The Spi algebra they find has the same structure as BMS, namely a semi-direct product of the abelian ideal of supertranslations and the Lorentz algebra. The difference lies in the size of the supertranslation ideal, which for the BMS algebra corresponds to functions on the 2-sphere whereas for Spi it corresponds to functions on the three-dimensional hyperboloid.

Also Troessaert [11] finds an asymptotic symmetry at spatial infinity which is larger than BMS, but smaller than the Spi algebra. By additionally assuming that the spacetime metric considered therein has to be asymptotically flat not only at spatial infinity but also at null infinity conditions on the metric functions are found which reduce the algebra to one that is isomorphic to the BMS algebra. This algebra is then in turn shown by [1] to be isomorphic to the algebra (2.30) with odd  $W$  and even  $T$ . For arbitrary  $W$  and  $T$  the algebra (2.30) is in fact also isomorphic to the one found by [11], before cutting it down to BMS. We recall some details of this construction in Appendix B.

To summarize, the boundary conditions we propose allow for an asymptotic symmetry at spatial infinity that is larger than the BMS algebra but smaller than the Spi algebra. Our result would therefore suggest that the tension arising from the presence of different asymptotic symmetries at spatial infinity and null infinity still remains.

## 7 Conclusions

We have analyzed the asymptotic symmetries of asymptotically flat spacetimes in the Hamiltonian formulation of GR. In contrast to previous treatments we have expressed the asymptotic expansion of the spatial metric and conjugate momenta in terms of a Bondi spacetime metric 3.1 using a 3+1 decomposition. The central insight of this procedure was that there exist divergent terms in the symplectic structure that cannot be canceled by parity conditions. To cancel these terms we proposed new boundary conditions, which require certain retarded time derivatives of the spacetime metric functions to vanish close to spatial infinity. Additionally, the foliation of spacetime has to be restricted in a particular way.

The charges which are generating supertranslations are parametrized by two arbitrary functions on the sphere and are finite for every mode of the two functions. The supertranslation sector is therefore larger than the one of the BMS algebra, which is parametrized by a single arbitrary function on the sphere.

A result that remains to be understood better is that spatial translations do not preserve the Bondi determinant condition in our treatment. Why is it that at null infinity this condition is fulfilled automatically whereas at spatial infinity it turns out to be too rigid? Another intriguing question in this context is whether the supertranslation sector at null infinity can be enlarged by dropping the determinant condition. Possibly this enlarged algebra at null infinity is isomorphic to the one we found at spatial infinity? We hope to address these questions in the future.

## Acknowledgment

This work was supported by funds provided by the National Science Center, projects number 2017/27/B/ST2/01902 and 2019/33/B/ST2/00050.

## A 3+1 decomposition

Following [8] we briefly recall the main features of the 3+1 decomposition. Such a decomposition is obtained by introducing a foliation of spacetime into spacelike hypersurfaces  $\Sigma_t$ , defined by  $t = \text{const.}$ , where the only condition on  $t$  is that the unit normal  $n_\alpha \propto \partial_\alpha t$  has to be a future directed timelike vector field. We will show in the next section how different choices of  $t$  affect the conjugate momenta and the parity conditions. Further, one introduces a time-evolution vector field  $t^\alpha$  to define the direction of time derivatives. The defining condition is  $t^\alpha \partial_\alpha t = 1$ , which allows to interpret the directional derivative  $t^\alpha \partial_\alpha$  as  $\partial_t$  and thus ensures that the direction of time derivatives is compatible with the meaning of time provided by  $t$ . Using  $t^\alpha$  one can define  $\dot{h}_{ab} = \mathcal{L}_t h_{ab}$  and conjugate momenta

$$\pi^{ab} = \frac{\partial}{\partial \dot{h}_{ab}} \mathcal{L}_G, \quad (\text{A.1})$$

where  $h_{ab}$  is the induced metric on  $\Sigma_t$  and  $\mathcal{L}_G$  is the gravitational Lagrangian. The vector field  $t^\alpha$  is usually decomposed into parts tangential and orthogonal to the spatial hypersurfaces

$$t^\alpha = N n^\alpha + N^a e_a^\alpha, \quad (\text{A.2})$$

where  $e_a^\alpha = \frac{\partial x^\alpha}{\partial y^a}$  are the tangent vectors on  $\Sigma_t$ ,  $x^\alpha$  are coordinates in the full spacetime and  $y^a$  are coordinates intrinsic to  $\Sigma_t$ .  $N$  is referred to as lapse and  $N^i$  as shift. Using this decomposition of  $t^\alpha$  one can write

$$dx^a = \frac{\partial x^a}{\partial t} dt + \frac{\partial x^a}{\partial y^\alpha} dy^\alpha = t^\alpha dt + e_a^\alpha dy^a \quad (\text{A.3})$$

and for the line element of a generic metric it follows

$$ds^2 = g_{\alpha\beta} dx^\alpha dx^\beta = -N^2 dt^2 + h_{ab} (dy^a + N^a dt)(dy^b + N^b dt), \quad (\text{A.4})$$

where the definition of the induced metric

$$h_{ab} = g_{\alpha\beta} e_a^\alpha e_b^\beta \quad (\text{A.5})$$

has been used. Note, that asymptotic flatness demands for lapse and shift to behave asymptotically as [3],[12]

$$N = 1 + O(1/r), \quad N^r = O(1/r), \quad N^A = O(1/r^2). \quad (\text{A.6})$$

The conjugate momenta can be expressed as

$$\pi^{ab} = \sqrt{h}(K^{ab} - Kh^{ab}) \quad (\text{A.7})$$

and we are going to use this expression to write the conjugate momenta in terms of  $g_{\alpha\beta}$  and its derivatives. The extrinsic curvature  $K_{ab}$  is defined as

$$K_{ab} = n_{\alpha;\beta} e_a^\alpha e_b^\beta, \quad (\text{A.8})$$

where semicolon denotes the covariant derivative associated with  $g_{\alpha\beta}$  and  $K = h^{ab}K_{ab}$  is the trace of extrinsic curvature. Finally,  $\dot{h}_{ab}$  can be written as

$$\dot{h}_{ab} = \mathcal{L}_t h_{ab} = 2NK_{ab} + N_{a|b} + N_{b|a}, \quad (\text{A.9})$$

where  $N_{a|b}$  denotes the covariant derivative associated with  $h_{ab}$ .

## B The algebra of asymptotic symmetries

This can be seen as follows [1]. In [11] the following asymptotic algebra is found

$$[(\mathcal{Y}_1, \omega'_1), (\mathcal{Y}_2, \omega'_2)] = \left( [\mathcal{Y}_1, \mathcal{Y}_2], \mathcal{Y}_1^a \partial_a \omega'_2 - \frac{s}{2} \psi_1 \omega'_2 - (1 \leftrightarrow 2) \right) = (\widehat{\mathcal{Y}}, \widehat{\omega}'), \quad (\text{B.1})$$

where  $x^a = (s, x^A)$  are coordinates on the unit hyperboloid,  $\mathcal{Y}^a$  represents the Lorentz algebra,  $\omega' = \sqrt{1-s^2}\omega$  and  $\omega(x^a)$  parametrize a sub-set of Spi supertranslations [17]. Full Spi supertranslations would be given by general functions  $\omega(x^a)$  but, as [11] explains, to remove divergences in the symplectic structure one has to demand

$$(D^a D_a + 3)\omega = 0. \quad (\text{B.2})$$

The general solution of this equation is shown to be of the form

$$\omega = \frac{1}{\sqrt{1-s^2}} (\widehat{\omega}^{\text{even}} + \widehat{\omega}^{\text{odd}}) \quad (\text{B.3})$$

$$\widehat{\omega}^{\text{even}} = \sum_{l,m} \widehat{\omega}_{lm}^V V_l(s) Y_{lm}^0(x^A), \quad \widehat{\omega}^{\text{odd}} = \sum_{l,m} \widehat{\omega}_{lm}^W W_l(s) Y_{lm}^0(x^A),$$

where odd and even refers to the combination of time reversal  $s \rightarrow -s$  and antipodal mapping  $x^A \rightarrow -x^A$  and  $V_l(s), W_l(s)$  are defined in terms of Legendre polynomials and Legendre functions of the second kind.

Rotations are parametrized by Killing vectors on the 2-sphere  $\mathcal{Y}_R^A(x^A)$

$$\mathcal{Y}^s = 0, \quad \mathcal{Y}^A = \mathcal{Y}_R^A \quad (\text{B.4})$$

and boosts by  $\psi(x^A)$  such that  $D^2\psi + 2\psi = 0$

$$\mathcal{Y}^s = -\frac{1}{2}(1-s^2)\psi, \quad \mathcal{Y}^A = -\frac{1}{2}s\gamma^{AB}\partial_B\psi. \quad (\text{B.5})$$

One can check that the Lorentz algebras in (B.1) and (2.30) are isomorphic under the identification  $\mathcal{Y}_R^A = Y^A$  and  $\psi = 2b$ . The action of the Lorentz algebra on  $\omega'$  in (B.1) can then be written as

$$\widehat{\omega}' = Y_1^A \partial_A \omega'_2 - s b_1 \omega'_2 - s \partial^A b_1 \partial_A \omega'_2 - (1-s^2) b_1 \partial_s \omega'_2 - (1 \leftrightarrow 2). \quad (\text{B.6})$$

The connection with the ADM description in (2.30) can be made by defining  $W, T$  as initial conditions at  $s = 0$

$$\omega|_{s=0} = \omega'|_{s=0} = W(x^A), \quad \partial_s \omega|_{s=0} = \partial_s \omega'|_{s=0} = T(x^A). \quad (\text{B.7})$$

One can check from the definitions of  $V_l(s)$  and  $W_l(s)$  that  $\omega|_{s=0}$  and  $\partial_s \omega|_{s=0}$  contain contributions from all modes of spherical harmonics and that therefore  $W$  and  $T$  such defined are arbitrary functions on the sphere. Substituting these definitions in (B.6) one obtains

$$\widehat{W} = Y_1^A \partial_A W_2 - b_1 T_2 - (1 \leftrightarrow 2). \quad (\text{B.8})$$

Acting with the  $s$  derivative on (B.6) yields

$$\begin{aligned} \partial_s \widehat{\omega}' = & Y_1^A \partial_A \partial_s \omega'_2 - b_1 \omega'_2 - s b_1 \partial_s \omega'_2 - \partial^A b_1 \partial_A \omega'_2 - s \partial^A b_1 \partial_A \omega'_2 \\ & + 2s b_1 \partial_s \omega'_2 - (1 - s^2) b_1 \partial_s^2 \omega'_2 - (1 \leftrightarrow 2). \end{aligned} \quad (\text{B.9})$$

The expression for  $\partial_s^2 \omega'|_{s=0}$  can be obtained from

$$(\mathcal{D}_a \mathcal{D}^a + 3)\omega = -(1 - s^2)^2 \partial_s^2 \omega + (1 - s^2) D^2 \omega + 3\omega = 0 \quad (\text{B.10})$$

and first one has

$$\partial_s^2 \omega|_{s=0} = D^2 \omega|_{s=0} + 3\omega|_{s=0} \quad (\text{B.11})$$

and together with

$$\partial_s^2 \omega'|_{s=0} = -\omega|_{s=0} + \partial_s^2 \omega|_{s=0} \quad (\text{B.12})$$

one finally has

$$\partial_s^2 \omega'|_{s=0} = 2\omega|_{s=0} + D^2 \omega|_{s=0}. \quad (\text{B.13})$$

Substituting this and the above definitions in (B.9) and evaluating at  $s = 0$  one finally obtains

$$\widehat{T} = Y_1^A \partial_A T_2 - 3b_1 W_2 - \partial_A b_1 D^A W_2 - b_1 D^2 W_2 - (1 \leftrightarrow 2). \quad (\text{B.14})$$

## References

- [1] M. Henneaux and C. Troessaert, “BMS Group at Spatial Infinity: the Hamiltonian (ADM) approach,” *JHEP* **03** (2018), 147 [arXiv:1801.03718 [gr-qc]].
- [2] L. Freidel, M. Geiller and D. Pranzetti, “Edge modes of gravity. Part I. Corner potentials and charges,” *JHEP* **11** (2020), 026 [arXiv:2006.12527 [hep-th]].
- [3] T. Regge and C. Teitelboim, “Role of Surface Integrals in the Hamiltonian Formulation of General Relativity,” *Annals Phys.* **88** (1974), 286
- [4] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, “Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems,” *Proc. Roy. Soc. Lond. A* **269** (1962) 21.



- [5] R. K. Sachs, “Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times,” *Proc. Roy. Soc. Lond. A* **270** (1962), 103-126
- [6] R. Sachs, “Asymptotic symmetries in gravitational theory,” *Phys. Rev.* **128**, 2851 (1962).
- [7] A. Strominger, “Lectures on the Infrared Structure of Gravity and Gauge Theory,” [arXiv:1703.05448 [hep-th]].
- [8] E. Poisson, “A Relativist’s Toolkit: The Mathematics of Black-Hole Mechanics,” Cambridge University Press, Cambridge, 2004. doi:10.1017/CBO9780511606601
- [9] G. Barnich and C. Troessaert, “BMS charge algebra,” *JHEP* **12** (2011), 105 doi:10.1007/JHEP12(2011)105 [arXiv:1106.0213 [hep-th]].
- [10] É. É. Flanagan and D. A. Nichols, “Conserved charges of the extended Bondi-Metzner-Sachs algebra,” *Phys. Rev. D* **95** (2017) no.4, 044002 doi:10.1103/PhysRevD.95.044002 [arXiv:1510.03386 [hep-th]].
- [11] C. Troessaert, *Class. Quant. Grav.* **35** (2018) no.7, 074003 doi:10.1088/1361-6382/aaae22 [arXiv:1704.06223 [hep-th]].
- [12] M. Henneaux and C. Troessaert, “Hamiltonian structure and asymptotic symmetries of the Einstein-Maxwell system at spatial infinity,” *JHEP* **07** (2018), 171 doi:10.1007/JHEP07(2018)171
- [13] G. Barnich and C. Troessaert, “Aspects of the BMS/CFT correspondence,” *JHEP* **05** (2010), 062 doi:10.1007/JHEP05(2010)062 [arXiv:1001.1541 [hep-th]].
- [14] G. Barnich and C. Troessaert, “Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited,” *Phys. Rev. Lett.* **105** (2010), 111103 doi:10.1103/PhysRevLett.105.111103 [arXiv:0909.2617 [gr-qc]].
- [15] R. L. Arnowitt, S. Deser and C. W. Misner, “The Dynamics of general relativity,” *Gen. Rel. Grav.* **40** (2008), 1997-2027 doi:10.1007/s10714-008-0661-1 [arXiv:gr-qc/0405109 [gr-qc]].
- [16] G. Compère and A. Fiorucci, “Advanced Lectures on General Relativity,” [arXiv:1801.07064 [hep-th]].
- [17] A. Ashtekar and R. O. Hansen, *J. Math. Phys.* **19** (1978), 1542-1566 doi:10.1063/1.523863