

# On the energy of gravitational waves

Rong-Gen Cai,<sup>1,2,3</sup> Xing-Yu Yang,<sup>1,2,\*</sup> and Long Zhao<sup>1,2</sup>

<sup>1</sup>*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,  
Chinese Academy of Sciences, Beijing 100190, China*

<sup>2</sup>*School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China*

<sup>3</sup>*School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study,  
University of Chinese Academy of Sciences, Hangzhou 310024, China*

The energy of gravitational waves (GWs) is a fundamental problem in gravity theory. The existing descriptions for the energy of GWs, such as the well-known Isaacson energy-momentum tensor, suffer from several defects. Due to the equivalence principle, the gravitational energy-momentum can only be defined quasilocally, being associated with a closed spacelike 2-surface bounding a region. We propose a new approach to derive the energy of GWs directly from the quasilocal gravitational energy. Such an approach is natural and consistent with the quasilocal energy-momentum, and it is valid for GWs with any wavelengths in any order of perturbations.

*Introduction.*—One of great predictions of general relativity is the existence of gravitational wave (GW). The observation of GWs from a binary black hole merger [1] has launched a new era of astronomy and cosmology. When discussing GWs, a fundamental problem is their energy. Back to the 1950s, there was a controversy on whether or not GWs can carry energy. The controversy was ultimately resolved by Bondi using a simple thought experiment [2].

A mathematical description for the energy of GWs was not devised until the works of Isaacson, in which an effective energy-momentum tensor of GWs was obtained by averaging the square of gradient of the wave field over several wavelengths with the shortwave approximation [3, 4]. In applications to physics of the very early Universe, the fluctuations of interest have wavelengths larger than the Hubble radius, an effective energy-momentum tensor was derived by Mukhanov, Abramo and Brandenberger [5, 6]. In these approaches named as geometric approach, the gravitational field is divided into the background part and wave part, the effective energy-momentum tensor comes from the back reaction of wave to the background. Another kind of approaches is named as field-theoretical approach, in which the effective energy-momentum tensor is derived by the Lagrange-Belinfante-Rosenfeld procedure [7–9]. The results are various expressions of pseudotensors [10–15].

Although different approaches for obtaining the energy of GWs were proposed in literature, they have several defects. In the geometric approach, an artificial partition of the gravitational field is needed, while in the field-theoretical approach, the pseudotensor depends on coordinates. Besides, an additional elaborate averaging scheme is necessary for both approaches in order to obtain a meaningful effective energy-momentum tensor of GWs. The dependence on these artificial objects leads to some ambiguities. As a result, different approaches are matched only in the linear order of metric perturba-

tions. These difficulties root in the equivalence principle of general relativity, which leads to the fact that the gravitational energy-momentum can not be defined locally.

The modern concept, introduced by Penrose [16], is that a proper energy-momentum of gravity is quasilocal, being associated with a closed spacelike 2-surface bounding a region. Chang, Nester and Chen found a natural quasilocal Hamiltonian interpretation of the pseudotensor [17]. Many proposals of quasilocal energy were made [18–28], of which an important one is the work of Brown and York in which they proposed their definition by using the Hamiltonian formulation of general relativity [22]. The Brown-York energy has the right asymptotic behavior but is not nonnegative in general, this defect motivated the seminal works [28, 29], in which Wang and Yau proposed a well-defined quasilocal energy (for a comprehensive review see [30]).

GWs are just parts of the gravitational field, their energy should also be quasilocal. A natural thought is that one should derive the energy of GWs directly from the quasilocal gravitational energy instead of the ill-defined gravitational energy-momentum tensor. In this paper, we give a new approach to derive the energy of GWs directly from the quasilocal gravitational energy. Such an approach is more natural and more consistent with the quasilocal energy-momentum, and it is valid for GWs with any wavelengths in any order of perturbations.

*Quasilocal energy.*—Due to the equivalence principle, the gravitational energy can not be defined locally. However when there is an asymptotic symmetry, the total energy can be defined, which is called ADM energy and Bondi energy when viewed from spatial infinity and null infinity, respectively [31, 32]. It was proposed to measure the energy of a system by enclosing it with a closed spacelike 2-surface, which is the idea behind the definition of quasilocal energy of the surface [16]. There are several conditions the quasilocal energy should satisfy: 1) the ADM or Bondi energy should be recovered as spatial or null infinity is approached (‘correct large-sphere behavior’), 2) the expected limits should be obtained when the surface converges to a point (‘correct small-sphere behav-

\* Corresponding author. [yangxingyu@itp.ac.cn](mailto:yangxingyu@itp.ac.cn)

ior'), 3) the quasilocal energy should be nonnegative in general and vanish for Minkowski spacetime. Brown and York obtained the quasilocal energy of a closed spacelike 2-surface including contributions from both gravitational and matter fields by employing a Hamilton-Jacobi analysis of the action functional [22]. Their definition has the right asymptotic behavior and was proved to be positive when the spacelike 3-region which the surface bounds is time symmetric [33], but it depends on the choice of the 3-region and is not nonnegative in general. Motivated by geometric consideration, Liu and Yau introduced a definition which is independent of the 3-region, and proved that it is always positive [27], however it can be strictly positive even when the surface is in a flat spacetime as pointed out by Murchadha, Szabados and Tod [34]. After that, Wang and Yau rectified that defect and obtained a well-defined quasilocal energy [28, 29], which is the most satisfactory one so far, and we will briefly revisit it.

Given a closed spacelike 2-surface  $\mathcal{S}$  which bounds a spacelike 3-region  $\mathcal{V}$  in the physical spacetime  $(\mathcal{M}, g_{\mu\nu})$ . Along  $\mathcal{S}$ , let  $u^\mu$  be the future-directed timelike unit normal to  $\mathcal{V}$  and  $v^\mu$  be the outward spacelike unit normal to  $\mathcal{S}$  such that  $u^\mu v_\mu = 0$ . Given a future-directed timelike unit vector field  $t^\mu$  along  $\mathcal{S}$ , it can be decomposed as  $t^\mu = Nu^\mu + N^\mu$ . Through the Hamilton-Jacobi analysis of the gravitational action, one can obtain the surface Hamiltonian

$$\mathfrak{H}(t^\mu, u^\mu) = -\frac{1}{\kappa} \int_{\mathcal{S}} [Nk - N^\mu v^\nu (K_{\mu\nu} - K_\lambda^\lambda \gamma_{\mu\nu})], \quad (1)$$

where  $\kappa = 8\pi G/c^4$ ,  $k$  is the trace of the two-dimensional extrinsic curvature of  $\mathcal{S}$  in  $\mathcal{V}$  with respect to  $v^\mu$ ,  $K_{\mu\nu}$  is the extrinsic curvature of  $\mathcal{V}$  in  $\mathcal{M}$  with respect to  $u^\mu$ , and  $\gamma_{\mu\nu}$  is the induced Riemannian metric on  $\mathcal{V}$ .

To define the quasilocal energy, one needs to find a reference action that corresponds to fixing the metric on the timelike boundary of the time history of the bounded region, and compute the corresponding reference surface Hamiltonian  $\bar{\mathfrak{H}}$ . The quasilocal energy is then the difference of the two surface Hamiltonian  $E = \mathfrak{H} - \bar{\mathfrak{H}}$ . The reference spacetime is arbitrary, and the freedom to choose different reference spacetime is just the freedom to choose the zero point of energy for the system. Therefore considering the conditions that quasilocal energy should satisfy, a natural choice for the reference spacetime  $(\bar{\mathcal{M}}, \bar{g}_{\mu\nu})$  is Minkowski spacetime, which is used through the paper.

Given an isometric embedding  $\varphi : \mathcal{S} \rightarrow \bar{\mathcal{M}}$ , i.e.  $\sigma_{\mu\nu} = (\varphi^* \bar{\sigma})_{\mu\nu}$ , where  $\sigma_{\mu\nu}$  and  $\bar{\sigma}_{\mu\nu}$  are the induced Riemannian metric on  $\mathcal{S}$  and  $\bar{\mathcal{S}} \equiv \varphi(\mathcal{S})$  respectively, and a future-directed timelike unit translational Killing vector field  $\bar{t}^\mu$  in  $\bar{\mathcal{M}}$ . Viewing  $\bar{t}^\mu$  as a vector field along  $\bar{\mathcal{S}}$ , one can decompose  $\bar{t}^\mu$  as  $\bar{t}^\mu = {}^\perp \bar{t}^\mu + {}^\parallel \bar{t}^\mu$ , where  ${}^\parallel \bar{t}^\mu$  is tangent to  $\bar{\mathcal{S}}$  and  ${}^\perp \bar{t}^\mu$  is normal to  $\bar{\mathcal{S}}$ . Choose  $\bar{u}^\mu = {}^\perp \bar{t}^\mu / |{}^\perp \bar{t}^\mu|$ ,  $\bar{N} = |{}^\perp \bar{t}^\mu|$  and  $\bar{v}^\mu$  be the outward spacelike unit normal vector of  $\bar{\mathcal{S}}$  that is orthogonal to  $\bar{u}^\mu$ , then one can obtain the corresponding surface Hamiltonian  $\bar{\mathfrak{H}}(\bar{t}^\mu, \bar{u}^\mu)$ , which obviously depends on the choice of the pair  $(\varphi, \bar{t}^\mu)$ .

Having picked  $\bar{t}^\mu$  and  $\bar{u}^\mu$  along  $\bar{\mathcal{S}}$  in  $\bar{\mathcal{M}}$ , one chooses a

corresponding pair of  $\check{t}^\mu$  and  $\check{u}^\mu$  along  $\mathcal{S}$  in  $\mathcal{M}$  as follows. First, via the embedding  $\varphi$ ,  ${}^\parallel \bar{t}^\mu$  can be pulled back to  $\mathcal{S}$  and be viewed as a vector field tangent to  $\mathcal{S}$  in  $\mathcal{M}$ . Next, define  $\check{t}^\mu = \check{N}\check{u}^\mu + {}^\parallel \check{t}^\mu$ , where  $\check{N} \equiv \varphi^* \bar{N}$ ,  ${}^\parallel \check{t}^\mu \equiv (\varphi^* {}^\parallel \bar{t})^\mu$ , and  $\check{u}^\mu$  is the unique future-directed timelike unit normal vector along  $\mathcal{S}$  such that

$$\mathfrak{h}_\mu \check{u}^\mu = \bar{\mathfrak{h}}_\mu \bar{u}^\mu. \quad (2)$$

Here  $\mathfrak{h}_\mu$  is the mean curvature vector of  $\mathcal{S}$  in  $\mathcal{M}$  which is assumed to be spacelike, and  $\bar{\mathfrak{h}}^\mu$  is the mean curvature vector of  $\bar{\mathcal{S}}$  in  $\bar{\mathcal{M}}$ . The mean curvature vector can be calculated as  $\mathfrak{h}^\mu = (K_\lambda^\lambda - K_{\rho\sigma} v^\rho v^\sigma) u^\mu - k v^\mu$ , which is indeed independent of the choice of  $u^\mu$  and  $v^\mu$ . Physically, Eq. (2) means the expansions of  $\mathcal{S} \subset \mathcal{M}$  and  $\bar{\mathcal{S}} \subset \bar{\mathcal{M}}$  along the respective directions  $\check{u}^\mu$  and  $\bar{u}^\mu$  are the same. Therefore, we obtain a quasilocal observer  $\check{t}^\mu$  along  $\mathcal{S}$  with the same lapse and shift as  $\bar{t}^\mu$  along  $\bar{\mathcal{S}}$ . Take  $\check{v}^\mu$  to be the spacelike unit normal vector of  $\mathcal{S}$  that is orthogonal to  $\check{u}^\mu$  and satisfies  $\check{v}^\mu \mathfrak{h}_\mu < 0$ . The quasilocal energy of  $\mathcal{S}$  with respect to the quasilocal observer  $(\varphi, \check{t}^\mu)$  is  $E(\mathcal{S}; \varphi, \check{t}^\mu) = \mathfrak{H}(\check{t}^\mu, \check{u}^\mu) - \bar{\mathfrak{H}}(\bar{t}^\mu, \bar{u}^\mu)$ .

In [29], Wang and Yau proved the following existence theorem. Given a Riemannian metric  $\sigma_{\mu\nu}$  and a function  $\tau$  on  $\mathcal{S}$  (which is topologically  $S^2$ ) such that the metric  $\sigma + d\tau \otimes d\tau$  is a Riemannian metric with positive Gaussian curvature. There exists a unique spacelike isometric embedding  $\varphi : \mathcal{S} \rightarrow \bar{\mathcal{M}}$  such that on  $\bar{\mathcal{S}}$  the time function (with respect to  $\bar{t}^\mu$ ) restricts to  $\tau$ , which shall be called the canonical embedding of  $(\mathcal{S}; \tau)$ . One can find that  $\mathfrak{h}_\mu \check{u}^\mu = -\mathcal{D}^\rho \mathcal{D}_\rho \tau / \sqrt{1 + \mathcal{D}^\sigma \tau \mathcal{D}_\sigma \tau}$  and  ${}^\parallel \check{t}^\mu = -\mathcal{D}^\mu \tau$  where  $\mathcal{D}$  is covariant derivative of  $\sigma_{\mu\nu}$ , which indicates that  $E(\mathcal{S}; \varphi, \check{t}^\mu)$  depends on  $(\varphi, \check{t}^\mu)$  only through  $\tau$ , therefore  $E(\mathcal{S}; \varphi, \check{t}^\mu)$  can also be written as  $E(\mathcal{S}; \tau)$ .

We summarize here the physical essence briefly without any cumbersome mathematical description. Due to the equivalence principle, the energy of a physical system should be measured by enclosing it with a closed spacelike 2-surface  $\mathcal{S}$ . Note that energy is a conserved quantity, but not an invariant quantity, it depends on the observer. Given  $\mathcal{S}$  in physical spacetime  $(\mathcal{M}, g_{\mu\nu})$ , one can only obtain its energy with respect to certain quasilocal observer which is depicted equivalently by  $(\varphi, \bar{t}^\mu)$  or  $\check{t}^\mu$  or  $\tau$ . Having the closed spacelike 2-surface and the quasilocal observer, the energy of the physical system with respect to the observer is determined, which can be calculated by the aforementioned mathematical procedure.

*Energy of GWs.*—Once the total energy of a physical system is obtained, the energy of GWs can be derived directly, since GWs are just parts of the system. Although only the closed spacelike 2-surface  $\mathcal{S}$  is involved when concerning the total energy, the spacelike 3-region  $\mathcal{V}$  must be specified when GWs are concerned, which can be easily seen as follows. Suppose a source emits a pulse of GWs, then for two different  $\mathcal{V}_1$  and  $\mathcal{V}_2$  with same boundary  $\mathcal{S}$  as shown in Fig. 1, they have same total energy but different energy of GWs.

For a given  $\mathcal{V}$ , it can be foliated into a family of closed spacelike 2-surfaces  $\{\mathcal{S}_\tau\}$ . Let the normal vector  $u^\mu$  of  $\mathcal{V}$




FIG. 1. Schematic diagram for a pulse of GWs. The shaded part denotes the worldtube of the source and the wavy lines denote the GWs.  $\mathcal{V}_1$  and  $\mathcal{V}_2$  are two different spacelike 3-regions with same boundary and same normal vector on the boundary.

be the  $\tilde{u}^\mu$  for each  $\mathcal{S}_r$ , then a family of isometric embeddings  $\phi \equiv \{\varphi_r : \mathcal{S}_r \rightarrow \bar{\mathcal{M}}\}$  is uniquely determined, which is an embedding  $\phi : \mathcal{V} \rightarrow \bar{\mathcal{M}}$  and gives a unique fleet of observers  $\tilde{t}^\mu$  on  $\mathcal{V}$ . For different foliation of  $\mathcal{V}$ ,  $\tilde{t}^\mu$  could be different, which indicates  $\mathcal{V}$  is observed by different observer. One can extend  $\phi$  into a mapping  $\Phi : \mathcal{M} \rightarrow \bar{\mathcal{M}}$  such that  $g_{\mu\nu} = (\Phi^* \bar{g})_{\mu\nu}$  on  $\mathcal{S}$  and  $\tilde{t}^\mu = (\Phi^* \bar{t})^\mu$  on  $\mathcal{V}$  by exponential map [35], which shall be called the canonical reference with respect to  $(\mathcal{V}; \phi, \tilde{t}^\mu)$ . In order to obtain the energy of GWs, one needs a method to depict the contributions of GWs. There are several commonly used methods [36], here we introduce two of them.

One method is through the scalar-vector-tensor (SVT) decomposition. In order to compare the tensors in  $\mathcal{M}$  and  $\bar{\mathcal{M}}$ , a prescription for identifying points of these spacetimes must be given, since it is a basic fact of differential geometry that only the comparison of tensors at the same point is meaningful. The tensors on these spacetimes can be compared after pullback  $\Phi^*$  or pushforward  $\Phi_*$ , we will omit these operators for brevity when no confusion arises. Then the physical metric can be decomposed as

$$g_{\mu\nu} = \bar{u}_\mu \bar{u}_\nu (\bar{u}^\rho \bar{u}^\sigma g_{\rho\sigma}) - 2\bar{u}_{(\mu} (\bar{u}^\rho \bar{\gamma}_{\nu)}^\sigma g_{\rho\sigma}) + \bar{\gamma}_\mu^\rho \bar{\gamma}_\nu^\sigma g_{\rho\sigma}, \quad (3)$$

with

$$\bar{u}^\rho \bar{u}^\sigma g_{\rho\sigma} = A, \quad (4a)$$

$$\bar{u}^\rho \bar{\gamma}_\nu^\sigma g_{\rho\sigma} = \bar{D}_\nu B + S_\nu, \quad (4b)$$

$$\bar{\gamma}_\mu^\rho \bar{\gamma}_\nu^\sigma g_{\rho\sigma} = C \bar{\gamma}_{\mu\nu} + (\bar{D}_\mu \bar{D}_\nu - \frac{1}{3} \bar{\gamma}_{\mu\nu} \bar{D}^\rho \bar{D}_\rho) E + \bar{D}_{(\mu} F_{\nu)} + H_{\mu\nu}, \quad (4c)$$

where  $\bar{u}^\mu$  is the normal vector on  $\bar{\mathcal{V}} \equiv \phi(\mathcal{V})$  satisfying  $\tilde{u}^\mu = (\Phi^* \bar{u})^\mu$ ,  $\bar{\gamma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{u}_\mu \bar{u}_\nu$  and  $\bar{D}$  is its corresponding covariant derivative.  $\{A, B, C, E\}$ ,  $\{S_\mu, F_\mu\}$  which satisfy  $\bar{D}^\mu S_\mu = \bar{D}^\mu F_\mu = 0$ , and  $H_{\mu\nu}$  which

satisfies  $\bar{D}^\mu H_{\mu\nu} = 0$  and  $H_\mu^\mu = 0$ , are called scalar-, vector-, and tensor-type components of  $g_{\mu\nu}$ , respectively. Here and hereafter the indices of these component quantities are raised and lowered with  $\bar{g}_{\mu\nu}$ . In order to ensure that the decomposition can be inverted (and therefore that there is a one-to-one correspondence between  $g_{\mu\nu}$  and the variables  $\{A, B, C, E, S_\mu, F_\mu, H_{\mu\nu}\}$ ), one must also impose the boundary conditions that  $\{B|_{\mathcal{S}}, E|_{\mathcal{S}}, (\bar{D}^\mu \bar{D}_\mu E)|_{\mathcal{S}}, (F_\mu)|_{\mathcal{S}}\}$  are equal to zero for finite  $\mathcal{V}$ , or approach to zero sufficiently fast at spatial infinity. These boundary conditions ensure the uniqueness of the decomposition.

It is found that the quasilocal energy derived by Wang and Yau is closely related to that derived by Chen, Nester and Tung from a covariant Hamiltonian formalism [24, 35, 37], which is closely related to the conserved currents obtained by Katz, Bičák and Lynden-Bell by applying the Noether theorem to the Lagrangian [26, 38]. Inspired by such relations, one can obtain

$$E(\mathcal{S}; \varphi, \tilde{t}^\mu) = \int_{\mathcal{S}} J^{\mu\nu} \tilde{v}_\nu \tilde{u}_\mu = \int_{\mathcal{V}} I^\mu \tilde{u}_\mu \quad (5)$$

where  $I^\mu = \nabla_\nu J^{\mu\nu}$  is the conserved Noether current which satisfies  $\nabla_\mu I^\mu \equiv 0$ . Furthermore,  $I^\mu$  can be written as  $I^\mu = (T_\nu^\mu + \mathcal{T}_\nu^\mu) \tilde{t}^\nu$ , where  $T_\nu^\mu$  is the energy-momentum tensor of matter fields which satisfies the Einstein equation  $G_\nu^\mu = \kappa T_\nu^\mu$ , and

$$2\kappa \mathcal{T}_\nu^\mu = g^{\rho\sigma} [(\Delta_{\rho\lambda}^\lambda \Delta_{\sigma\nu}^\mu + \Delta_{\rho\sigma}^\mu \Delta_{\lambda\nu}^\lambda - 2\Delta_{\rho\lambda}^\mu \Delta_{\sigma\nu}^\lambda) - \delta_\nu^\mu (\Delta_{\rho\sigma}^\eta \Delta_{\eta\lambda}^\lambda - \Delta_{\rho\lambda}^\eta \Delta_{\eta\sigma}^\lambda)] + g^{\mu\lambda} (\Delta_{\rho\sigma}^\sigma \Delta_{\lambda\nu}^\rho - \Delta_{\lambda\sigma}^\sigma \Delta_{\rho\nu}^\rho), \quad (6)$$

with  $\Delta_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\rho} (\bar{\nabla}_\mu g_{\rho\nu} + \bar{\nabla}_\nu g_{\rho\mu} - \bar{\nabla}_\rho g_{\mu\nu})$  is the difference between the Christoffel symbols in  $\mathcal{M}$  and  $\bar{\mathcal{M}}$ . One can find that the energy of a physical system contains contributions from matter fields and gravitational fields, and the energy of GWs shall contribute the part of the gravitational energy.

Under the SVT decomposition, the quasilocal energy  $E(\mathcal{S}; \varphi, \tilde{t}^\mu)$  as a functional of  $g_{\mu\nu}$  can be written as functional of  $\{A, B, C, E, S_\mu, F_\mu, H_{\mu\nu}\}$ . The energy of GWs in  $\mathcal{V}$  with respect to observer  $(\phi, \tilde{t}^\mu)$  is the part, which is contributed only by the tensor component  $H_{\mu\nu}$ , of the gravitational energy,

$$E_{\text{GW}}(\mathcal{V}; \phi, \tilde{t}^\mu) = \int_{\mathcal{V}} \mathfrak{P}_H \mathcal{T}_\nu^\mu \tilde{t}^\nu \tilde{u}_\mu, \quad (7)$$

where the operator  $\mathfrak{P}_H$  picks out the parts of  $\mathcal{T}_\nu^\mu$  that only depend on  $H_{\mu\nu}$ . Note that  $\mathfrak{P}_H \mathcal{T}_\nu^\mu$  can be taken as an effective energy-momentum tensor of GWs, but it depends on the observer unlike the energy-momentum tensor of matter fields. In our approach, the energy of GWs is derived *directly* from the quasilocal gravitational energy, then one can call a tensor as effective energy-momentum tensor when the volume integration of its time-time component equals to the energy. This is a significant difference between our approach and the existing

approaches in literature, in which one derives an effective energy-momentum tensor (which suffers from several defects) firstly, then calls the volume integration of its time-time component as energy.

Another method is through the Newman-Penrose (NP) formalism [39]. Choosing an orthonormal tetrad  $\{(e_\alpha)^\mu\}$  such that on  $\mathcal{V}$  one has  $(e_0)^\mu = \tilde{u}^\mu$  and  $(e_3)^\mu = \tilde{v}^\mu$ , one can construct a tetrad of four null basis vectors  $\{m^\mu, \bar{m}^\mu, l^\mu, k^\mu\}$  by defining  $m^\mu \equiv [(e_1)^\mu - i(e_2)^\mu]/\sqrt{2}$ ,  $\bar{m}^\mu \equiv [(e_1)^\mu + i(e_2)^\mu]/\sqrt{2}$ ,  $l^\mu \equiv [(e_0)^\mu - (e_3)^\mu]/\sqrt{2}$ ,  $k^\mu \equiv [(e_0)^\mu + (e_3)^\mu]/\sqrt{2}$  (here and hereafter we follow the notations in [40]). Of this null vectors,  $l^\mu$  and  $k^\mu$  are real and point along “ingoing” and “outgoing” directions respectively,  $m^\mu$  and  $\bar{m}^\mu$  are complex conjugates of each other and span “angular” directions. With this tetrad, the Weyl tensor  $C_{\mu\nu\rho\sigma}$  can be recast to five complex scalars, of which  $\Psi_4 \equiv C_{\mu\nu\rho\sigma}\bar{m}^\mu l^\nu \bar{m}^\rho l^\sigma$  represents outgoing GWs [41].

One can rewrite the surface Hamiltonian as functional of the spin coefficients, for instance, when  $\tilde{t}^\mu = \tilde{u}^\mu$ ,

$$\mathfrak{H}(\tilde{t}^\mu, \tilde{u}^\mu) = \frac{\sqrt{2}}{\kappa} \int_{\mathcal{S}} \Re(\rho + \mu), \quad (8)$$

where  $\rho \equiv -m^\mu \bar{m}^\nu \nabla_\nu k_\mu$  and  $\mu \equiv \bar{m}^\mu m^\nu \nabla_\nu l_\mu$  are spin coefficients,  $\Re$  denotes the real part. By solving the NP equations, the spin coefficients can be written as functional which contains  $\Psi_4$  terms. The energy of GWs is the part, which only contains contributions from  $\Psi_4$ , of the gravitational energy. One can find the reference surface Hamiltonian  $\bar{\mathfrak{H}}$  will contribute nothing, since the Minkowski reference spacetime has vanishing Weyl tensor. A straight conclusion is that the conformally flat spacetime has vanishing energy of GWs due to its vanishing Weyl tensor. This is quite reasonable physically, since the conformally flat spacetime has no GWs. Note that as the zero point of energy, a natural choice for the reference spacetime is Minkowski spacetime. However, once only the energy of GWs is concerned, any spacetime with vanishing Weyl tensor can be used as reference spacetime. Such a choice could give different value of energy of the gravitational field compared with the Minkowski choice, but it should not change the value of energy of GWs. The relation of the NP formalism and SVT decomposition can be seen from that in the linear regime  $\Psi_4$  is directly related to the metric perturbations as  $\Psi_4 = \partial_t^2(H_+ - iH_\times)$  [36].

*Calculations in perturbation theory.*—Usually in practice, it is hard to get the physical metric  $g_{\mu\nu}$  including the GW part by solving the Einstein equation directly. Regarding that the physical metric deviates smally from a known exact solution which is the so-called background spacetime  $(\mathcal{M}_0, {}^{(0)}g_{\mu\nu})$ , one can apply the perturbation theory to find the approximate solution of the Einstein equation. In order to make the comparison of tensors in the physical and background spacetime meaningful, a prescription for identifying points of these spacetimes must be given. A gauge choice is precisely this, i.e., a mapping  $\Upsilon$  between the physical and

background spacetime. Using a gauge  $\Upsilon$ , the physical metric  $g_{\mu\nu}$  can be perturbed on the background metric  ${}^{(0)}g_{\mu\nu}$  as  $g_{\mu\nu} = \sum_{n=0}^{\infty} \frac{1}{n!} {}^{(n)}g_{\mu\nu}$ , then the energy of GWs can be calculated order by order,  $E_{\text{GW}} = \sum_{n=0}^{\infty} {}^{(n)}E_{\text{GW}} = \sum_{n=0}^{\infty} \int_{\mathcal{V}} \mathfrak{P}_H {}^{(n)}\mathcal{T}_\nu^\mu \tilde{t}^\nu \tilde{u}_\mu$ .

Here we show an explicit calculation for GWs in asymptotically flat spacetime with a vacuum Einstein equation. For such a spacetime, a convenient choice for the background spacetime is Minkowski spacetime. Consider a 3-region  $\mathcal{V}$  with Eulerian observer  $(\phi, \bar{t}^\mu)$ , i.e., the worldline of observer is orthogonal to  $\mathcal{V}$ , then  $\tilde{t}^\mu = \tilde{u}^\mu$ . Calculating  $\mathfrak{P}_H \mathcal{T}_\nu^\mu$  order by order gives  $\mathfrak{P}_H {}^{(0)}\mathcal{T}_\nu^\mu = \mathfrak{P}_H {}^{(1)}\mathcal{T}_\nu^\mu = 0$ , and

$$2\kappa \mathfrak{P}_H {}^{(2)}\mathcal{T}_\nu^\mu \tilde{t}^\nu \tilde{t}_\mu = \frac{1}{4} {}^{(0)}\dot{H}_{\mu\nu} {}^{(0)}\dot{H}^{\mu\nu} + \frac{1}{4} \bar{D}_\rho {}^{(0)}H_{\mu\nu} \bar{D}^\rho {}^{(0)}H^{\mu\nu} - \frac{1}{2} \bar{D}_\mu {}^{(0)}H_{\rho\nu} \bar{D}^\nu {}^{(0)}H^{\rho\mu}, \quad (9)$$

where the over dot denotes the Lie derivative  $\mathcal{L}_{\tilde{t}}$ . The first and second terms are equal due to the first order Einstein equation. The volume integration of last term can be converted to a surface integral after integration by parts, which vanishes due to the isometric condition  $g_{\mu\nu} = \bar{g}_{\mu\nu}$  on the surface. Then one has

$${}^{(2)}E_{\text{GW}}(\mathcal{V}; \phi, \bar{t}^\mu) = \int_{\mathcal{V}} \frac{1}{4\kappa} {}^{(0)}\dot{H}_{\mu\nu} {}^{(0)}\dot{H}^{\mu\nu}, \quad (10)$$

which is the commonly used form in literature. In a general case, the choice of background spacetime is quite arbitrary as long as the perturbation theory is valid. For instance, the FLRW metric is commonly used as the background metric in the cosmological perturbation theory.

One now has three spacetimes, they are the physical, reference and background spacetimes, respectively. Reference  $\Phi$  (gauge  $\Upsilon$ ) is a one-to-one correspondence between points in the physical spacetime and points in the reference (background) spacetime. The schematic relations of these three spacetimes are shown in Fig. 2.




FIG. 2. Schematic relations of physical spacetime  $\mathcal{M}$ , reference spacetime  $\bar{\mathcal{M}}$  and background spacetime  $\mathcal{M}_0$ . The canonical reference  $\Phi$  is related to the observer, while the gauge  $\Upsilon$  is arbitrary. The black parts are physically relevant and the grey parts are just mathematically relevant.

Note that only the physical and reference spacetimes are physically relevant, as they depict the dynamics and ground state respectively, while the background spacetime is just a mathematical auxiliary tool which is useful in the perturbation theory. The canonical reference is related to the observer, thus it is also physically relevant,



while the gauge is just auxiliary mapping in perturbation theory. Therefore it is obvious that the physical quantities, of course including the energy of GWs, do not depend on the choice of the background spacetime and the gauge.

*Conclusions.*—The energy of GWs is a fundamental problem in gravity theory. The existing descriptions for the energy of GWs, such as the well-known Isaacson energy-momentum tensor, suffer from several defects. Due to the equivalence principle, the gravitational energy-momentum can not be defined locally in general relativity. The proper gravitational energy-momentum is quasilocal, being associated with a closed spacelike 2-surface bounding a region, as introduced by Penrose. In this paper, we proposed a new approach to derive the energy of GWs directly from the quasilocal gravitational

energy through the SVT decomposition and NP formalism. Such a quasilocal approach is more natural and more consistent with the quasilocal nature of gravitational energy-momentum. Although we only showed an explicit calculation for vacuum asymptotic flat spacetime in linear order, our approach is valid for GWs with any wavelengths in any order of perturbations. As a byproduct, with our approach the gauge dependence issue on the energy spectrum of GWs disappears naturally [42].

We thank Misao Sasaki and Shing-Tung Yau for helpful communications. This work is supported in part by the National Natural Science Foundation of China Grants No.11690022, No.11821505, No.11991052, No.11947302 and by the Strategic Priority Research Program of the CAS Grant No.XDPB15, and by the Key Research Program of Frontier Sciences of CAS.

- 
- [1] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **116**, 061102 (2016), [arXiv:1602.03837 \[gr-qc\]](#).
  - [2] H. Bondi, *Nature* **179**, 1072 (1957).
  - [3] R. A. Isaacson, *Phys. Rev.* **166**, 1263 (1968).
  - [4] R. A. Isaacson, *Phys. Rev.* **166**, 1272 (1968).
  - [5] V. F. Mukhanov, L. R. W. Abramo, and R. H. Brandenberger, *Phys. Rev. Lett.* **78**, 1624 (1997), [arXiv:gr-qc/9609026](#).
  - [6] L. R. W. Abramo, R. H. Brandenberger, and V. F. Mukhanov, *Phys. Rev. D* **56**, 3248 (1997), [arXiv:gr-qc/9704037](#).
  - [7] F. J. Belinfante, *Physica* **6**, 887 (1939).
  - [8] F. J. Belinfante, *Physica* **7**, 449 (1940).
  - [9] L. Rosenfeld, *Mém. Acad. Roy. Belg.* **18** (1940).
  - [10] A. Einstein, *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften (Berlin)*, 844 (1915).
  - [11] A. Papapetrou, *Proc. Roy. Irish Acad.* **A 52**, 11 (1948).
  - [12] P. G. Bergmann and R. Thomson, *Phys. Rev.* **89**, 400 (1953).
  - [13] J. N. Goldberg, *Phys. Rev.* **111**, 315 (1958).
  - [14] C. Möller, *Annals of Physics* **4**, 347 (1958).
  - [15] L. D. Landau and E. M. Lifschits, *The Classical Theory of Fields*, Course of Theoretical Physics, Vol. Volume 2 (Pergamon Press, Oxford, 1975).
  - [16] R. Penrose, *Proc. Roy. Soc. Lond. A* **381**, 53 (1982).
  - [17] C.-C. Chang, J. M. Nester, and C.-M. Chen, *Phys. Rev. Lett.* **83**, 1897 (1999), [arXiv:gr-qc/9809040](#).
  - [18] J. Katz and A. Ori, *Classical and Quantum Gravity* **7**, 787 (1990).
  - [19] J. Jezierski and J. Kijowski, *Gen. Rel. Grav.* **22**, 1283 (1990).
  - [20] A. J. Dougan and L. J. Mason, *Phys. Rev. Lett.* **67**, 2119 (1991).
  - [21] G. Bergqvist, *Classical and Quantum Gravity* **9**, 1917 (1992).
  - [22] J. D. Brown and J. W. York, Jr., *Phys. Rev. D* **47**, 1407 (1993), [arXiv:gr-qc/9209012](#).
  - [23] S. A. Hayward, *Phys. Rev. D* **49**, 831 (1994), [arXiv:gr-qc/9303030](#).
  - [24] C.-M. Chen, J. M. Nester, and R.-S. Tung, *Phys. Lett. A* **203**, 5 (1995), [arXiv:gr-qc/9411048](#).
  - [25] C.-M. Chen and J. M. Nester, *Class. Quant. Grav.* **16**, 1279 (1999), [arXiv:gr-qc/9809020](#).
  - [26] C.-M. Chen and J. M. Nester, *Grav. Cosmol.* **6**, 257 (2000), [arXiv:gr-qc/0001088](#).
  - [27] C.-C. M. Liu and S.-T. Yau, *Phys. Rev. Lett.* **90**, 231102 (2003), [arXiv:gr-qc/0303019](#).
  - [28] M.-T. Wang and S.-T. Yau, *Phys. Rev. Lett.* **102**, 021101 (2009), [arXiv:0804.1174 \[gr-qc\]](#).
  - [29] M.-T. Wang and S.-T. Yau, *Communications in Mathematical Physics* **288**, 919 (2009).
  - [30] L. B. Szabados, *Living Rev. Rel.* **12**, 4 (2009).
  - [31] R. L. Arnowitt, S. Deser, and C. W. Misner, *Phys. Rev.* **122**, 997 (1961).
  - [32] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, *Proc. Roy. Soc. Lond. A* **269**, 21 (1962).
  - [33] Y. Shi and L.-F. Tam, *Journal of Differential Geometry* **62**, 79 (2002).
  - [34] N. Ó. Murchadha, L. B. Szabados, and K. P. Tod, *Phys. Rev. Lett.* **92**, 259001 (2004), [arXiv:gr-qc/0311006 \[gr-qc\]](#).
  - [35] J.-L. Liu and C. Yu, *J. Geom. Phys.* **120**, 330 (2017), [arXiv:1604.05302 \[math.DG\]](#).
  - [36] N. T. Bishop and L. Rezzolla, *Living Rev. Rel.* **19**, 2 (2016), [arXiv:1606.02532 \[gr-qc\]](#).
  - [37] C.-M. Chen, J. M. Nester, and R.-S. Tung, *Phys. Rev. D* **72**, 104020 (2005), [arXiv:gr-qc/0508026](#).
  - [38] J. Katz, J. Bičák, and D. Lynden-Bell, *Phys. Rev. D* **55**, 5957 (1997), [arXiv:gr-qc/0504041 \[gr-qc\]](#).
  - [39] E. Newman and R. Penrose, *J. Math. Phys.* **3**, 566 (1962).
  - [40] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers, and E. Herlt, *Exact solutions of Einstein's field equations*, Cambridge Monographs on Mathematical Physics (Cambridge Univ. Press, Cambridge, 2003).
  - [41] P. Szekeres, *J. Math. Phys.* **6**, 1387 (1965).
  - [42] R.-G. Cai, X.-Y. Yang, and L. Zhao, (2021), [arXiv:2109.06865 \[astro-ph.CO\]](#).