Optical Observations and Modeling of the Superluminous Supernova 2018lfe

YAO YIN,¹ SEBASTIAN GOMEZ,^{2,1} EDO BERGER,¹ GRIFFIN HOSSEINZADEH,^{3,1} MATT NICHOLL,⁴ AND PETER K. BLANCHARD⁵

¹Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

³Steward Observatory, University of Arizona, 933 N. Cherry Ave., Tucson, AZ 85721, USA

⁴Birmingham Institute for Gravitational Wave Astronomy and School of Physics and Astronomy, University of Birmingham, Birmingham B15 2TT, UK

⁵Center for Interdisciplinary Exploration and Research in Astrophysics and Department of Physics and Astronomy,

Northwestern University, 1800 Sherman Ave, Evanston, IL 60201, USA

ABSTRACT

We present optical imaging and spectroscopy of SN 2018lfe, which we classify as a Type I superluminous supernova (SLSN-I) at a redshift of z = 0.3501 with a peak absolute magnitude of $M_r \approx -22.1$ mag, one of the brightest SLSNe discovered. SN 2018lfe was identified for follow-up using our FLEET machine learning pipeline. Both the light curve and the spectra of SN 2018lfe are consistent with the broad population of SLSNe. We fit the light curve with a magnetar central engine model and find an ejecta mass of $M_{\rm ej} \approx 3.8 \, {\rm M}_{\odot}$, a magnetar spin period of $P \approx 2.9 \, {\rm ms}$ and a magnetic field strength of $B_{\perp} \approx 2.8 \times 10^{14} \, {\rm G}$. The magnetic field strength is near the top of the distribution for SLSNe, while the spin period and ejecta mass are near the median values of the distribution for SLSNe. From late-time imaging and spectroscopy we find that the host galaxy of SN 2018lfe has an absolute magnitude of $M_r \approx -17.85$, ($L_B \approx 0.029 \, L^*$), and an inferred metallicity of $Z \approx 0.3 \, {\rm Z}_{\odot}$, and star formation rate of $\approx 0.8 \, {\rm M}_{\odot} \, {\rm yr}^{-1}$.

Keywords: supernova: general – supernova: individual (SN 2018lfe)

1. INTRODUCTION

Superluminous supernovae (SLSNe) are a rare class of core-collapse supernovae (SNe) that can exceed the luminosities of normal SNe by two orders of magnitude (Chomiuk et al. 2011; Quimby et al. 2011). Type I SLSNe (hereafter, SLSNe-I) are classified based on their hydrogen-free spectra, strong O II absorption lines at early times, and blue continuum (Chomiuk et al. 2011; Quimby et al. 2011; Nicholl et al. 2014a; Gal-Yam 2019; Inserra et al. 2013). Despite their increasing discovery rate over the past decade thanks to systematic wide-field optical surveys, SLSNe-I still remain a small population, with only about 150 spectroscopically classified events to date (Gomez et al. 2020); moreover, not all of these events have well observed light curves and spectroscopic time series.

Corresponding author: Yao Yin yyin@college.harvard.edu

Several possible mechanisms that can power SLSNe-I have been explored in the literature: large radioactive ⁵⁶Ni mass (Pastorello et al. 2010), circumstellar interaction (Chevalier & Irwin 2011), and a magnetar central engine (Kasen & Bildsten 2010; Woosley 2010). The bulk of the observational evidence points to a magnetar engine as the dominant mechanism. This includes the diverse light curve behavior (Nicholl et al. 2017), the early-time UV spectra (Mazzali et al. 2016; Nicholl et al. 2017; Howell et al. 2013; Dessart 2019; Yan et al. 2017), the late-time light curve flattening (Nicholl et al. 2018; Blanchard et al. 2021), and their nebular spectra (Dessart et al. 2012; Nicholl et al. 2019; Jerkstrand 2017). The preference of SLSNe-I to low metallicity host galaxies (similar to the hosts of the engine-powered long gamma-ray bursts) also supports a magnetar engine energy source (Lunnan et al. 2014; Leloudas et al. 2015; Ørum et al. 2020; Angus et al. 2019; Perley et al. 2016; Chen et al. 2017; Schulze et al. 2018).

Here, we present detailed optical observations of SN 2018lfe, a transient first detected by the Zwicky Transient Facility (ZTF; Bellm et al. 2019) on 2018




Figure 1. Optical light curve of SN 2018lfe in the gwri bands using data from FLWO, PSST, and ZTF. The photometry is corrected for Galactic extinction, and time is shown in rest-frame days relative to the time of peak brightness in r-band. The blue vertical lines mark the times when the first three spectra were taken (see Section 3), the date of the last spectrum is outside the range of this plot.

December 31 and first reported by the Pan-STARRS Survey for Transients (PSST; Chambers et al. 2019a) on 2019 January 1. Our follow-up observations of the SN and its host galaxy continue up to a year after discovery and classify the event as a SLSN-I at a redshift of $z = 0.3501 \pm 0.0004$, with a peak absolute magnitude of $M_r \approx -22.1$. We use this dataset to investigate the properties of SN2018lfe and its host, in the context of magnetar-powered SLSNe, to gain a better understanding of the larger SLSNe population.

The paper is structured as follows. In §2 we present SN 2018lfe and our multi-band follow-up observations. In §3 we describe the observed features of the spectra and compare to other SLSNe-I. In §4 we present our modeling of the light curve using a magnetar engine model, the bolometric light curve, and compare the parameters of SN 2018lfe to the existing SLSNe-I population. In §5 we explore the host galaxy properties. Throughout the paper we assume a flat Λ CDM cosmology with $H_0 = 69.3$ km s⁻¹ Mpc⁻¹, $\Omega_m = 0.286$, and $\Omega_{\Lambda} = 0.712$ (Hinshaw et al. 2013).

2. OBSERVATIONS

2.1. Discovery and Classification

SN 2018lfe was first reported by PSST on 2019 January 1 (MJD = 58484.0) with a magnitude of g = 19.66 at coordinates R.A.= $09^{h}33^{m}29^{s}.556$, decl.= + $00^{\circ}03'08''.39$ (J2000) (Chambers et al. 2019b). We selected SN 2018lfe for follow-up observations as a probable SLSN-I candidate using our *Finding Luminous* and Exotic Extragalactic Transients (FLEET) machine learning pipeline (Gomez et al. 2020), which determined a probability of SN 2018lfe being a SLSN-I of $P_{\rm SLSN-I} \approx 0.71$. We obtained a spectrum on 2019 January 31 with the Inamori-Magellan Areal Camera and Spectrograph (IMACS; Dressler et al. 2011) on the Magellan Baade 6.5-m Telescope, and classified it as a SLSN-I (Gomez 2019) with aid from template matching to known SN spectra with the Supernovae Identification (SNID, Blondin & Tonry 2007) program. We calculated the redshift $z = 0.3501 \pm 0.0004$ by measuring the center of the H β and [O III] $\lambda\lambda$ 4959,5007 host galaxy emission lines visible in this spectrum in addition to three additional follow-up spectra.

2.2. Optical Photometry

We obtained images of SN 2018lfe with the KeplerCam imager on the 1.2-m telescope at Fred Lawrence Whipple Observatory (FLWO) in the g, r, and i bands, starting on 2019 January 11 (MJD = 58494). SN 2018lfe was also observed at several earlier epochs by ZTF in the g and r bands, for which we obtained the raw images from the NASA/IPAC Infrared Science Archive¹. The SN was also observed by PSST in the w and i bands, and we obtained the photometry from the publicly available data on the Open Supernova Catalog² (Guillochon et al. 2017).

We processed the FLWO and ZTF images using standard IRAF³ routines. We modeled the point-spread function from each image using reference stars and subtracted the model from the target to measure the instrumental magnitudes. We combined ZTF images obtained within ≈ 2 days of each other to improve the signal-tonoise ratio of the measured fluxes. Similarly, for data obtained from FLWO after a phase of 110 days (MJD = 58575), we combined images in the r and i bands that were obtained within ≈ 2 days of each other. For the PSST photometry, we averaged the reported magnitudes obtained on the same night.

We calibrated all the photometry relative to $PS1/3\pi$ magnitudes and applied a correction for Galactic extinction with $E(B - V) = 0.033 \pm 0.001$ mag and $R_V = 3.1$ (Schlaffy & Finkbeiner 2011), using the Cardelli et al. (1989) Milky Way extinction law. We define phase 0 to be the date of the brightest magnitude in the *r*band, MJD = 58465, with peak apparent magnitude of $r = 19.2 \pm 0.1$ and $g = 19.1 \pm 0.1$ after correcting for Galactic extinction. We use this reference point throughout the paper, with all days quoted in the rest frame unless otherwise specified. The full light curve of

¹ irsa.ipac.caltech.edu

² https://sne.space/

³ IRAF is written and supported by the National Optical Astronomy Observatories, operated by the Association of Universities for Research in Astronomy, Inc. under cooperative agreement with the National Science Foundation.

SN 2018lfe is shown in Figure 1. The photometry table is provided in the supplementary materials in machine readable format, as well as on the Open Supernova Catalog (Guillochon et al. 2017).

2.3. Optical Spectroscopy

We obtained four epochs of low-resolution optical spectroscopy at phases of 35, 41, 84, and 256 days. We used the IMACS and the Low Dispersion Survey Spectrograph (LDSS3c, Stevenson et al. (2016)) spectrographs on the Magellan 6.5-m telescopes; and the Blue Channel (Schmidt et al. 1989) and Binospec (Fabricant et al. 2019) spectrographs on the MMT 6.5-m telescope. All the spectra were obtained with the slit oriented along the parallactic angle. Details of the spectroscopic observations are provided in Table 1. We processed all spectra using standard IRAF routines. A wavelength calibration was applied using a HeNeAr lamp spectrum taken near the time of each science image. Relative flux calibration was applied to each spectrum using a standard star taken on the same night. The spectra were then calibrated to absolute flux scaling using the available gri photometry. Lastly, the spectra were corrected for Galactic extinction and transformed to the rest frame of SN 2018lfe.

The resulting spectra are shown in Figure 2, with prominent emission features from the host galaxy, absorbtion features of the SN, and telluric absorption features marked. The spectra are also available on the Weizmann Interactive Supernova Data Repository (WISeREP⁴; Yaron & Gal-Yam 2012).

Table 1. Optical Spectroscopy of SN 2018lfe

UT Date	MJD	$Phase^{a}$	Exp. Time	Airmass	Wavelength	Telescope + Instrument	Grating	Slit Width
		(d)	(s)		(Å)		(lines/mm)	('')
2019 Jan 30	58513	35	1200	1.17	4500-9400	Baade + IMACS	300	0.9
$2019~{\rm Feb}~7$	58521	41	1200	1.50	4500 - 9500	Clay + LDSS	300	1
$2019~{\rm Apr}~6$	58579	84	1800	1.16	4500 - 9400	Baade + IMACS	1600	1
$2019~{\rm Nov}~24$	58811	256	900	1.22	3800-9200	MMT + Binospec	300	0.9

^aRest-frame days since peak r-band luminosity.

3. SPECTRAL FEATURES AND COMPARISONS

The earliest spectrum of SN 2018lfe (35 days) exhibits a blue continuum and broad absorption features. We detect a broad feature at the location of Mg I] λ 4500 and measure its width with a Gaussian profile, leading to velocity widths of 11190±660, 11770±600, and 10180±470 km s⁻¹ for phases of 35, 41, and 84 days, respectively. There is thus no evidence for a significant velocity evolution. The three early spectra also show Si II λ 6355 regions with an average velocity center of \approx 11700 km s⁻¹. This is consistent with the ejecta velocity obtained from fitting the light curve with a magnetar model (see §4). As is common for SLSNe-I, the continuum becomes redder at later phases. No obvious SN features are detected in the final spectrum at 256 days, which shows only narrow emission lines from the host galaxy.

We compare the spectra of SN 2018lfe to those of three other SLSNe-I in Figure 2. The objects SN 2010md

(=PTF10hgi) (Inserra et al. 2013), SN 2016eay (=Gaia16apd) (Nicholl et al. 2017), and PTF12dam (Nicholl et al. 2014b) were chosen for having similar values of magnetar engine and ejecta parameters⁵ as SN 2018lfe, and for their abundance of spectral data, specifically at phases that match those of SN 2018lfe.

The emission regions of Mg I] λ 4571, Ca II H λ 3933, Ca II K λ 3968 in the early spectrum of SN 2018lfe at phase of 35 – 41 days also match with the early phase data of SN 2010md and PTF12dam. As demonstrated in Figure 2, the spectroscopic evolution of SN 2018lfe follows that of SN 2010md. In particular, both objects display prominent Mg I] and Ca II emissions features that are similar in shape at similar phases, namely 41 – 47, 75 – 95, and 241 – 256 days from explosion. However, SN2010md shows an unusually prominent absorption line near 5700 Å, unlike SN 2018lfe and other typical SLSNe-I.

⁴ https://wiserep.weizmann.ac.il

 $^{^{5}}$ The values of these parameters are from our previous work in Nicholl et al. (2017).




Figure 2. Optical spectra of SN 2018lfe spanning a rest frame phase of 35 to 256 days, with the last spectrum being dominated by host galaxy emission. The spectra are corrected for Galactic extinction and shifted to the rest-frame of SN 2018lfe using z = 0.3501. We bin the spectra for clarity and show the unbinned spectra as shaded regions. The labeled vertical lines mark spectral features from the SN, narrow lines from the host galaxy, and telluric absorption. The different colors in the vertical lines represent different elements. We compare the evolution of SN 2018lfe to three other SLSNe-I: SN 2010md (PTF10hgi; Inserra et al. 2013), SN 2016eay (Gaia16apd; Nicholl et al. 2017), and PTF12dam (Nicholl et al. 2013), with their spectra colored gray.

4. LIGHT CURVE MODELING

We model the multi-band light curve with a magnetar spin-down model implemented in the Modular Open-Source Fitter for Transients (MOSFiT) Python package, a Markov chain Monte Carlo (MCMC) code that can fit the light curves of transients using a variety of power sources (Guillochon et al. 2018). We use the emcee (Foreman-Mackey et al. 2013) implementation of MCMC to run each sampler. Since MOSFiT implements a simplified analytical one-zone model, the uncertainties presented here represent only the statistical errors on the fit. The priors used in the model are listed in Table 2, following our previous work in Nicholl et al. (2017).

The magnetar model fits are shown in Figure 3, and the parameter posterior distributions are shown in Figure 4 and listed in Table 2. The model provides an excellent fit to the data, with the following primary physical parameters: an initial spin period of $P \approx 2.9$ ms, a magnetic field of $B_{\perp} \approx 2.8 \times 10^{14}$ G, an ejecta mass of $M_{\rm ej} \approx 3.8 \, {\rm M}_{\odot}$, and an ejecta velocity of $v_{\rm ej} \approx 10^4$ km s⁻¹ (in good agreement with the velocity inferred from the spectra in §3). We calculate a total kinetic energy of $E_K \approx 4 \times 10^{51}$ erg. The mass of the progenitor star is estimated to be $M_{\rm NS} + M_{\rm ej} \approx 5.7 \, {\rm M}_{\odot}$, which is consistent with the range of $3.6 - 40 \, {\rm M}_{\odot}$ inferred from the SLSNe-I sample of Blanchard et al. (2020).

We compare the resulting parameters for SN 2018lfe to the distribution of 62 SLSNe from Nicholl et al. (2015); Blanchard et al. (2020); Villar et al. (2018) in Figure 6. As seen in the Figure, all of the parameters for SN 2018lfe fall within the range covered by previous SLSNe. However, we do find that the magnetic field value inferred for SN 2018lfe is in the top 16th percentile of the SLSN distribution. The combination of the strong

Parameter	Definition	Prior	Full Light Curve	Units
$\log(M_{\rm ej})$	Ejecta Mass	[-1, 2]	$0.58^{+0.11}_{-0.11}$	${\rm M}_{\odot}$
B_{\perp}	Magnetar Magnetic Field Strength	[0.1, 10]	$2.8^{+2.1}_{-1.0}$	$10^{14}\mathrm{G}$
$P_{\rm spin}$	Magnetar Spin	[1, 10]	$2.89^{+0.47}_{-0.58}$	
$M_{\rm NS}$	Mass of Neutron Star	[0.1, 2.2]	$1.87^{+0.24}_{-0.41}$	${ m M}_{\odot}$
$v_{ m ej}$	Ejecta Velocity	[5.0e3, 2.0e4]	10062_{-949}^{+867}	${\rm km~s^{-1}}$
$E_{m{k}}^{\dagger}$	Ejecta Kinetic Energy		$3.9^{+1.3}_{-1.2}$	$10^{51} {\rm ~erg~s^{-1}}$
$t_{\rm exp}$	Explosion Time Relative To First Data Point	[-500, 0]	$-5.4^{+0.86}_{-1.0}$	days
$\log(n_{H,\text{host}})$	Column density in the host galaxy	[16, 23]	$18.4^{+1.7}_{-1.7}$	$\rm cm^{-2}$
T_{\min}	Photospheric Temperature	[3.0e3, 1.0e4]	7683^{+1532}_{-1731}	Κ
κ	Optical opacity	[0.05, 0.2]	$0.16\substack{+0.02\\-0.04}$	$\rm cm^2g^{-1}$
$\log(\kappa_{\gamma})$	Gamma-ray opacity	[-1, 4]	1.6 ± 1.6	$\rm cm^2g^{-1}$

 Table 2.
 Best-fit Parameters for the Magnetar Models

NOTE—Best model parameters, definitions, prior ranges, and 1σ error bars for the realizations shown in Figure 3.

[†] These parameters were not fit for, but were calculated using all the posterior distribution samples of the fitted parameters.




Figure 3. *Top Panel:* Light curve of SN 2018lfe with a magnetar engine model. Each line is a sample realization of the most likely models generated from MOSFiT.

magnetic field, fast spin $(P_{\rm spin} < 3 \text{ ms})$, and high ejecta velocity is most likely responsible for significant heating of the ejecta, which explains the high optical luminosity of SN 2018lfe.

We further use the MOSFiT model to construct a bolometric light curve, photospheric radius evolution, and temperature evolution of SN 2018lfe; see Figure 5. The shaded regions in the Figure represent the 1σ values across all 500 walkers. We find a peak bolometric luminosity of $(4.0 \pm 1.1) \times 10^{44}$ erg s⁻¹. The integrated radiated energy over the observed portion of the light curve (phase 0 to 103 days) is $E_{\rm rad} \approx 1.2 \times 10^{51}$ erg, consistent with the known population of SLSNe, despite its bright peak luminosity (Lunnan et al. 2018; Nicholl et al. 2020). The photospheric radius exhibits an increase up to a maximum of $4.6^{+1.1}_{-0.5} \times 10^{15}$ cm at a phase of $\approx 30 - 50$ d, with a gradual decline thereafter. Finally, the photospheric temperature does shows a steady decline from an inferred peak of $\approx 3.5 \times 10^4$ K down to $\approx 10^4$ K at a phase of $\gtrsim 30$ days and suggest a temperature $< 10^4$ K at 40 days. Although due to the lack




Figure 4. Posterior distributions and correlations of the magnetar model parameters. The median and 1σ values are marked and labeled. A summary of all the parameter values is presented in Table 2.




Figure 5. Bolometric light curve, blackbody temperature evolution, and photospheric radius evolution obtained from the MOSFiT models. The plots show the mean model (blue line) and the 1σ regions of 500 walkers used in the MCMC process.

of data in the UV range, there is large uncertainty at early times when the spectral energy distribution peaks in this wavelength range.

5. HOST GALAXY

No galaxy is detected at the position of SN 2018lfe in the PS1/3 π catalog down to the nominal 5 σ limiting magnitudes of $q \approx 23.3$, $r \approx 23.2$, $i \approx 23.1$, and $z \approx 22.3$ (Chambers et al. 2016). We obtained deeper images on 2020 November 15 using Binospec and detected an extended source with a half-light radius of $\approx 0.5''$, measured using sep (Barbary et al. 2015). From extrapolating the observed light curve (see Section 2.2), we expect the apparent magnitude of the SLSN to be dimmer than ≈ 25 mag in the r-band, therefore the host should be the only source present. We extracted the host magnitudes by performing aperture photometry using the photutils package (Bradley et al. 2020) with a circular aperture of radius 2'' across all filters. At the measured redshift of z = 0.3501, the observed r-band is closest to rest-frame B-band. The apparent host magnitude of $r \approx 23.52$ mag thus corresponds to a $L_B \approx 0.029 \text{ L}^*$, determined using the DEEP2 luminosity function at z = 0.3 (Willmer et al. 2006). The host galaxy properties are summarized in Table 3.

We measured the strength of the galaxy emission lines by fitting narrow Gaussian profiles (FWHM ≈ 40 Å) to




Figure 6. Median values and 1σ errors of key parameters (P, B_{\perp} , $M_{\rm ej}$, E_K) for all SLSNe in Blanchard et al. 2020; Villar et al. 2018 and Nicholl et al. 2015 (labeled blue) compared with SN 2018lfe (labeled orange). SN 2018lfe lies consistent with the magnetar model.

the three spectra at phases of 35, 41, and 84 days. The flux ratio of $L_{\text{H}\alpha}/L_{\text{H}\beta} = 2.75 \pm 0.37$ is consistent with the expectation for case B recombination ($L_{\text{H}\alpha}/L_{\text{H}\beta} =$ 2.86), suggesting no significant host galaxy extinction (in agreement with the lack of extinction inferred from modeling of the SN light curves). The H α line luminosity is $L_{\text{H}\alpha} = (9.8 \pm 0.3) \times 10^{43} \text{ erg s}^{-1}$. This corresponds to a star formation rate (SFR) of $0.77 \pm 0.02 \text{ M}_{\odot}$ yr⁻¹ (Kennicutt 1998)

To calculate the metallicity we use the R_{23} method (Kobulnicky et al. 1999) and adopt the lower branch solution due to the lack of an N II detection. We determine a value of $12 + \log(O/H) = 8.18 \pm 0.12$, or $Z = 0.31 \pm 0.05 Z_{\odot}$ (Asplund et al. 2009). Coincidentally, this is also where the two values from the R_{23} method converge, and therefore our metallicity is not affected by the choice of the lower or upper branch solution. The low metallicity is consistent with the consensus that SLSNe preferentially occur in galaxies with sub-solar metallicity (Lunnan et al. 2014; Chen et al. 2017; Perley et al. 2016; Schulze et al. 2017).

 Table 3. Host Galaxy Properties

	Value (mag)	Units
g	24.07 ± 0.20	mag
r	23.52 ± 0.24	mag
i	24.01 ± 0.19	mag
z	24.36 ± 0.19	mag
M_g	-17.30 ± 0.2	mag
M_r	-17.85 ± 0.3	mag
M_i	-17.36 ± 0.4	mag
M_z	-17.01 ± 0.3	mag
L_B	0.029 ± 0.01	L^*
$12 + \log(O/H)$	8.18 ± 0.12	
Z	0.31 ± 0.05	Z_{\odot}
SFR	0.77 ± 0.02	${\rm M}_{\odot}~{\rm yr}^{-1}$

NOTE—Apparent magnitudes, not corrected for extinction, obtained from observations using the Binospec (Fabricant et al. 2003) imaging camera on the MMT on November 15, 2020. Absolute magnitudes are corrected for extinction. The star formation rate and the metallicities (in terms of $12 + \log(O/H)$ and Z_{\odot}) are estimated using the flux immensities of the host galaxy emission lines. L_B is the host luminosities relative to L^* in the *B* band (Willmer et al. 2006).

, due to the fact that at z=0.35, the observed r-band is closest to rest-frame B-band, see Section 5.

6. CONCLUSIONS

We presented the classification and detailed optical follow-up observations of the Type-I SLSN 2018lfe and its host galaxy at z = 0.3501. We model the light curve using a spin down magnetar model and find that SN 2018lfe has a best fit ejecta mass of $\approx 3.8 \, M_{\odot}$, a magnetar spin period of $\approx 2.9 \, \text{ms}$ and a magnetic field strength of $\approx 2.8 \times 10^{14} \, \text{G}$, consistent with the population of known SLSNe. We determine the host galaxy properties through spectral analysis and follow

- Angus, C. R., Smith, M., Sullivan, M., et al. 2019, MNRAS, 487, 2215
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARA&A, 47, 481
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123
- Barbary, K. 2016, extinction, v0.3.0, Zenodo, doi:10.5281/zenodo.804967
- Barbary, K., Boone, K., & Deil, C. 2015, sep: v0.3.0, v0.3.0, Zenodo, doi:10.5281/zenodo.15669

up photometric observations and find a low metallicity of $12 + \log(O/H) = 8.18 \pm 0.12$, typical for SLSNe-I hosts. Therefore, SN 2018lfe is a typical SLSN-I in terms of its explosion and host galaxy properties and contributes to a small population of SLSNe-I with a well-defined explosion time and peak absolute magnitude.

We thank Y. Beletsky for carrying out some of the Magellan observations. The Berger Time-Domain Group at Harvard is supported in part by NSF under grant AST-1714498 and by NASA under grant NNX15AE50G. M. Nicholl is supported by a Royal Astronomical Society Research Fellowship and by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 948381). Operation of the Pan-STARRS1 telescope is supported by the National Aeronautics and Space Administration under grant No. NNX12AR65G and grant No. NNX14AM74G issued through the NEO Observation Program. This paper includes data gathered with the 6.5 meter Magellan Telescopes located at Las Campanas Observatory, Chile. Observations reported here were obtained at the MMT Observatory, a joint facility of the University of Arizona and the Smithsonian Institution. This research has made use of NASA's Astrophysics Data System. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France.

Software: Astropy(AstropyCollaborationetal.2018), MOSFiT(Guillochon et al. 2018), PyRAF(Science Software Branch at STScI 2012), SAOImage DS9 (Smithsonian Astrophysical Observatory 2000), emcee(Foreman-Mackey et al. 2013), corner (Foreman-Mackey 2016), Matplotlib(Hunter 2007), SciPy(van der Walt et al. 2011), NumPy(Oliphant 2007), SEP(Barbary et al. 2015) extinction(Barbary 2016), FLEET(Gomez et al. 2020)).

REFERENCES

- Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, PASP, 131, 018002
- Blanchard, P. K., Berger, E., Nicholl, M., et al. 2021, arXiv:2105.03475
- Blanchard, P. K., Berger, E., Nicholl, M., & Villar, V. A. 2020, ApJ, 897, 114
- Blondin, S., & Tonry, J. L. 2007, ApJ, 666, 1024
- Bradley, L., Sipőcz, B., Robitaille, T., et al. 2020, astropy/photutils: 1.0.0, 1.0.0, Zenodo, doi:10.5281/zenodo.4044744

- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245
- Chambers, K. C., Magnier, E. A., Metcalfe, N., et al. 2016, arXiv e-prints, arXiv:1612.05560
- Chambers, K. C., Huber, M. E., Flewelling, H., et al. 2019 a, Transient Name Server Discovery Report, 2019-5, 1
- Chambers, K. C., Huber, M. E., Flewelling, H., et al. 2019 b, Transient Name Server Discovery Report, 2019-5, 1
- Chen, T.-W., Schady, P., Xiao, L., et al. 2017, The Astrophysical Journal, 849, L4
- Chevalier, R. A., & Irwin, C. M. 2011, ApJ, 729, L6
- Chomiuk, L., Chornock, R., Soderberg, A. M., et al. 2011, ApJ, 743, 114
- Dessart, L. 2019, A&A, 621, A141
- Dessart, L., Hillier, D. J., Waldman, R., Livne, E., & Blondin, S. 2012, MNRAS, 426, L76
- Dressler, A., Bigelow, B., Hare, T., et al. 2011, PASP, 123, 288
- Fabricant, D., Fata, R., Epps, H., et al. 2019, Publications of the Astronomical Society of the Pacific, 131, 075004
- Fabricant, D. G., Epps, H. W., Brown, W. L., Fata, R. G., & Mueller, M. 2003, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes, ed. M. Iye & A. F. M. Moorwood, 1134–1144
- Foreman-Mackey, D. 2016, The Journal of Open Source Software, 1
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, PASP, 125, 306
- Gal-Yam, A. 2019, ARA&A, 57, 305
- Gomez, S. 2019, Transient Name Server Classification Report, 2019-1246, 1
- Gomez, S., Berger, E., Blanchard, P. K., et al. 2020, The Astrophysical Journal, 904, 74
- Gomez, S., Berger, E., Blanchard, P. K., et al. 2020, FLEET Finding Luminous and Exotic Extragalactic Transients, 1.0.0, Zenodo, doi:10.5281/zenodo.4013965
- Guillochon, J., Nicholl, M., Villar, V. A., et al. 2018, ApJS, 236, 6
- Guillochon, J., Parrent, J., Kelley, L. Z., & Margutti, R. 2017, ApJ, 835, 64
- Hinshaw, G., Larson, D., Komatsu, E., et al. 2013, ApJS, 208, 19
- Howell, D. A., Kasen, D., Lidman, C., et al. 2013, ApJ, 779, 98
- Hunter, J. D. 2007, Computing in Science and Engineering, 9, 90
- Inserra, C., Smartt, S. J., Jerkstrand, A., et al. 2013, ApJ, 770, 128

- Jerkstrand, A. 2017, Spectra of Supernovae in the Nebular Phase, 795
- Kasen, D., & Bildsten, L. 2010, The Astrophysical Journal, 717, 245–249
- Kennicutt, Robert C., J. 1998, ARA&A, 36, 189
- Kobulnicky, H. A., Kennicutt, Robert C., J., & Pizagno, J. L. 1999, ApJ, 514, 544
- Leloudas, G., Schulze, S., Krühler, T., et al. 2015, Monthly Notices of the Royal Astronomical Society, 449, 917–932
- Lunnan, R., Chornock, R., Berger, E., et al. 2014, ApJ, 787, 138
- Lunnan, R., Fransson, C., Vreeswijk, P. M., et al. 2018, Nature Astronomy, 2, 887
- Mazzali, P. A., Sullivan, M., Pian, E., Greiner, J., & Kann, D. A. 2016, MNRAS, 458, 3455
- Nicholl, M., Berger, E., Blanchard, P. K., Gomez, S., & Chornock, R. 2019, ApJ, 871, 102
- Nicholl, M., Berger, E., Margutti, R., et al. 2017, ApJL, 835, L8
- Nicholl, M., Berger, E., Margutti, R., et al. 2017, The Astrophysical Journal, 835, L8
- Nicholl, M., Guillochon, J., & Berger, E. 2017, ApJ, 850, 55
- Nicholl, M., Smartt, S. J., Jerkstrand, A., et al. 2013, Nature, 502, 346
- Nicholl, M., Smartt, S. J., Jerkstrand, A., et al. 2014a, MNRAS, 444, 2096
- Nicholl, M., Smartt, S. J., Jerkstrand, A., et al. 2014b, MNRAS, 444, 2096
- Nicholl, M., Smartt, S. J., Jerkstrand, A., et al. 2015, ApJ, 807, L18
- Nicholl, M., Blanchard, P. K., Berger, E., et al. 2018, ApJL, 866, L24
- Nicholl, M., Blanchard, P. K., Berger, E., et al. 2020, Nature Astronomy, arXiv:2004.05840
- Oliphant, T. E. 2007, CSE, 9, 10
- Ørum, S. V., Ivens, D. L., Strandberg, P., et al. 2020, A&A, 643, A47
- Pastorello, A., Smartt, S. J., Botticella, M. T., et al. 2010, ApJL, 724, L16
- Perley, D. A., Quimby, R. M., Yan, L., et al. 2016, ApJ, 830, 13
- Quimby, R. M., Kulkarni, S. R., Kasliwal, M. M., et al. 2011, Nature, 474, 487
- Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103
- Schmidt, G. D., Weymann, R. J., & Foltz, C. B. 1989, PASP, 101, 713
- Schulze, S., Krühler, T., Leloudas, G., et al. 2017, Monthly Notices of the Royal Astronomical Society, 473, 1258–1285

- Schulze, S., Krühler, T., Leloudas, G., et al. 2018, MNRAS, 473, 1258
- Science Software Branch at STScI. 2012, PyRAF: Python alternative for IRAF, Astrophysics Source Code Library, ascl:1207.011
- Smithsonian Astrophysical Observatory. 2000, SAOImage DS9: A utility for displaying astronomical images in the X11 window environment, ascl:0003.002
- Stevenson, K. B., Bean, J. L., Seifahrt, A., et al. 2016, ApJ, 817, 141
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, CSE, 13, 22
- Villar, V. A., Nicholl, M., & Berger, E. 2018, ApJ, 869, 166
- Willmer, C. N. A., Faber, S. M., Koo, D. C., et al. 2006, The Astrophysical Journal, 647, 853–873
- Woosley, S. E. 2010, ApJL, 719, L204
- Yan, L., Lunnan, R., Perley, D. A., et al. 2017, ApJ, 848, 6
- Yaron, O., & Gal-Yam, A. 2012, PASP, 124, 668