

BV right-continuous solutions of second-order differential inclusions governed by maximal monotone operators

Dalila Azzam-Laouir *

November 30, 2021

Abstract

This paper concerns existence of right-continuous with bounded variation solutions of a perturbed second-order differential inclusion governed by time and state-dependent maximal monotone operators.

Keywords: Bounded variation, differential measure, Lipschitz mapping, maximal monotone operator, pseudo-distance, right-continuous, second-order.

AMS Subject Classifications: 2010: 34A60, 28A25, 28C20

1 Introduction

Let $I = [0, T]$ ($T > 0$) and \mathcal{H} a separable Hilbert space. Consider, for each $(t, x) \in I \times \mathcal{H}$, a maximal monotone operator of \mathcal{H} ; $A(t, x)$, for which the set-valued map $t \mapsto A(t, x)$ is right-continuous with bounded variation (BVRC) w.r.t the time t , and Lipschitz continuous w.r.t the state variable x , in the sense that there exists a function $\rho : I \rightarrow [0, \infty[$, which is right-continuous on $[0, T[$, and a nonnegative real number γ such that for $0 \leq s \leq t \leq T$, $x, y \in \mathcal{H}$,

$$\text{dis}(A(t, x), A(s, y)) \leq d\rho([s, t]) + \gamma\|x - y\| = \rho(t) - \rho(s) + \gamma\|x - y\|,$$

where $\text{dis}(\cdot, \cdot)$ is the pseudo-distance between maximal monotone operators introduced by Vladimirov [49]; see relation (2.8), and $f : I \times \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ a Carathéodory mapping satisfying a linear growth condition.

Let λ be the Lebesgue measure on I and $d\rho$ the Stieljes-measure associated with ρ . We set $\nu := \lambda + d\rho$ and $\frac{d\lambda}{d\nu}$ the density of λ w.r.t ν .

*LAOTI, FSEI, Université Mohamed Seddik Benyahia de Jijel, Algérie. E-mail: laouir.dalila@gmail.com, dalilalaouir@univ-jijel.dz

In this paper, we are mainly interested by the existence of BVRC mappings $u, v : I \longrightarrow \mathcal{H}$ satisfying

$$(P_f) \begin{cases} u(0) = u_0, v(0) = v_0 \in D(A(0, u_0)); \\ v(t) \in D(A(t, u(t))) \quad \forall t \in I; \\ \frac{du}{d\nu}(t) = v(t) \quad d\nu - a.e. t \in I; \\ -\frac{dv}{d\nu}(t) \in A(t, u(t))v(t) + f(t, u(t), v(t))\frac{d\lambda}{d\nu}(t) \quad d\nu - a.e. t \in I. \end{cases}$$

This work constitutes an extension to the second-order of few results dealing with first-order BVRC time-dependent maximal monotone operators differential inclusions, established in [9, 16, 31].

For some investigations on first-order differential inclusions governed by time-dependent and time and state dependent maximal monotone operators, where the variation of these operators is absolutely continuous or continuous with bounded variation; we refer for instance to [12, 13, 14, 15, 20, 24, 30, 31, 32, 41, 45, 48, 49]. Concerning the second order, we refer to the recent works [28, 43].

It is well known that the normal cone to closed and convex sets is a maximal monotone operator, so differential inclusions governed by this normal cone, called "sweeping processes" ([34, 35, 33]) are particular cases of the evolutions with general maximal monotone operators. Existence of BVRC solutions to the first-order sweeping process was discussed in many interesting papers; we cite for instance [1, 5, 18, 19, 27, 29, 36, 39, 46, 47]. However, to the best of our knowledge, there are no results for the BVRC second-order sweeping process in comparison with the vast literature concerning absolutely continuous and continuous with bounded variation solutions; see [2, 3, 4, 6, 7, 8, 10, 11, 21, 22, 25, 26, 39, 40, 44].

The paper is organized as follows. In section 2, we give notations and recall the needed preliminary results. In section 3, we prove the existence of right-continuous with bounded variation solutions to the evolution problem (P_f) when $t \mapsto A(t, \cdot)$ is BVRC and the perturbation $f : I \times \mathcal{H} \times \mathcal{H} \longrightarrow \mathcal{H}$ is measurable on I and Lipschitz-continuous on $\mathcal{H} \times \mathcal{H}$.

2 Notations and Preliminaries

Throughout the paper, we will denote by $\langle \cdot, \cdot \rangle$ the inner product of \mathcal{H} , $\| \cdot \|$ the associated norm and $r\overline{B}_{\mathcal{H}}$ the closed ball of center 0 and radius $r > 0$. We will denote by $\mathcal{B}(I)$ the Borel σ -algebra on I . The identity mapping of \mathcal{H} will be denoted by $Id_{\mathcal{H}}$. For a subset K of \mathcal{H} , $\overline{\text{co}}(K)$ will be the closed convex hull of K , which is characterized by:

$$\overline{\text{co}}(K) = \{y \in \mathcal{H} : \langle y, x \rangle \leq \sup_{z \in K} \langle z, x \rangle \quad \forall x \in K\}. \quad (2.1)$$

If μ is a positive measure on I , we will denote by $L^p(I, \mathcal{H}; \mu)$ $p \in [1, +\infty[$, the Banach space of classes of p - μ -integrable mappings, equipped with its classical norm $\| \cdot \|_p$.

The following results concern definitions and some properties of functions with bounded variation and general vector measures, taken from references [33, 36, 37, 38].

Let $u : I \longrightarrow \mathcal{H}$. The variation of u in I is the nonnegative extended real number

$$\text{var}(u; I) := \sup \sum_{k=1}^n \|u(t_k) - u(t_{k-1})\|, \quad (2.2)$$

where the supremum is taken in $[0, +\infty]$ w.r.t all the finite sequences $t_0 < t_1 < \dots < t_n$ of points of I (n is arbitrary). The function u is said with bounded variation (BV) if and only if $\text{var}(u; I) < +\infty$. In this case we have

$$\lim_{s \uparrow t} \|u(t) - u(s)\| = \lim_{s \uparrow t} \text{var}(u; [s, t]).$$

Now, let μ be a positive Radon measure on I and $\hat{\mu}$ be an \mathcal{H} -valued measure on I admitting a density $\frac{d\hat{\mu}}{d\mu} \in L^1_{\text{loc}}(I, \mathcal{H}; \mu)$. Then for $d\mu$ -almost every $t \in I$, we have

$$\frac{d\hat{\mu}}{d\mu}(t) = \lim_{\varepsilon \downarrow 0} \frac{d\hat{\mu}([t, t + \varepsilon])}{d\mu([t, t + \varepsilon])} = \lim_{\varepsilon \downarrow 0} \frac{d\hat{\mu}([t - \varepsilon, t])}{d\mu([t - \varepsilon, t])}. \quad (2.3)$$

The measure $\hat{\mu}$ is absolutely continuous w.r.t μ if and only if $\hat{\mu} = \frac{d\hat{\mu}}{d\mu}\mu$, i.e., $\frac{d\hat{\mu}}{d\mu}$ is a density of $\hat{\mu}$ w.r.t μ . In this case a mapping $u : I \longrightarrow \mathcal{H}$ is $\hat{\mu}$ -integrable if and only if the mapping $t \mapsto u(t) \frac{d\hat{\mu}}{d\mu}(t)$ is μ -integrable, and we have

$$\int_I u(t) d\hat{\mu}(t) = \int_I u(t) \frac{d\hat{\mu}}{d\mu}(t) d\mu(t). \quad (2.4)$$

If $u : I \longrightarrow \mathcal{H}$ is BVRC and du is its differential measure, then we have

$$u(t) = u(s) + \int_{]s, t]} du \quad \forall s, t \in I \ (s \leq t). \quad (2.5)$$

Conversely, if there exists $v \in L^1(I, \mathcal{H}; \mu)$ such that $u(t) = u(0) + \int_{]0, t]} v d\mu$ for all $t \in I$, then u is BVRC and $du = v d\mu$, that is v is a density of the vector measure du w.r.t the measure μ . So that by (2.3), we get

$$v(t) = \frac{du}{d\mu}(t) = \lim_{\varepsilon \downarrow 0} \frac{du([t, t + \varepsilon])}{d\mu([t, t + \varepsilon])} = \lim_{\varepsilon \downarrow 0} \frac{du([t - \varepsilon, t])}{d\mu([t - \varepsilon, t])}. \quad (2.6)$$

If $\mu(\{t\}) > 0$, this last relation shows that

$$v(t) = \frac{du}{d\mu}(t) = \frac{du(\{t\})}{d\mu(\{t\})} \quad \text{and} \quad \frac{d\lambda}{d\mu}(t) = 0. \quad (2.7)$$

Next, we give definition and some properties of maximal monotone operators. We refer the reader to [17, 23, 50] for more details.

Let $A : \mathcal{H} \rightrightarrows \mathcal{H}$ be a set-valued map. We denote by $D(A)$ and $Gr(A)$ its domain and graph. We say that the operator A is monotone, if $\langle y_1 - y_2, x_1 - x_2 \rangle \geq 0$ for all $(x_i, y_i) \in Gr(A)$ ($i = 1, 2$), and we say that A is a maximal monotone operator of \mathcal{H} , if it is monotone and its graph could not be contained strictly in the graph of any other monotone operator.

If A is a maximal monotone operator of \mathcal{H} , then for every $x \in D(A)$, $A(x)$ is nonempty closed and convex. We denote the projection of the origin on the set $A(x)$ by $A^0(x)$.

For $\eta > 0$, we denote by $J_\eta^A = (Id_{\mathcal{H}} + \eta A)^{-1}$ the resolvent of A . It is well-known that this operator is single-valued and defined on all of \mathcal{H} , furthermore $J_\eta^A(x) \in D(A)$, for all $x \in \mathcal{H}$.

Let $A : D(A) \subset \mathcal{H} \rightrightarrows \mathcal{H}$ and $B : D(B) \subset \mathcal{H} \rightrightarrows \mathcal{H}$ be two maximal monotone operators. Then, $dis(A, B)$ is the Vladimirov's pseudo-distance between A and B , defined as follows

$$dis(A, B) = \sup \left\{ \frac{\langle y - y', x' - x \rangle}{1 + \|y\| + \|y'\|} : (x, y) \in Gr(A), (x', y') \in Gr(B) \right\}. \quad (2.8)$$

The following lemmas are needed for our proof. We refer to [31] for their proofs.

Lemma 2.1 *Let A be a maximal monotone operator of \mathcal{H} . If $x \in \overline{D(A)}$ and $y \in \mathcal{H}$ are such that*

$$\langle A^0(z) - y, z - x \rangle \geq 0 \quad \forall z \in D(A),$$

then $x \in D(A)$ and $y \in A(x)$.

Lemma 2.2 *Let A_n ($n \in \mathbb{N}$) and A be maximal monotone operators of \mathcal{H} such that $dis(A_n, A) \rightarrow 0$. Suppose also that $x_n \in D(A_n)$ with $x_n \rightarrow x$ and $y_n \in A_n(x_n)$ with $y_n \rightarrow y$ weakly for some $x, y \in E$. Then $x \in D(A)$ and $y \in A(x)$.*

Lemma 2.3 *Let A and B be maximal monotone operators of \mathcal{H} . Then*

1) for $\eta > 0$ and $x \in D(A)$

$$\|x - J_\eta^B(x)\| \leq \eta \|A^0(x)\| + dis(A, B) + \sqrt{\eta(1 + \|A^0(x)\|)dis(A, B)}.$$

2) For $\eta > 0$ and $x, x' \in E$

$$\|J_\eta^A(x) - J_\eta^A(x')\| \leq \|x - x'\|.$$

Lemma 2.4 *Let A_n ($n \in \mathbb{N}$) and A be maximal monotone operators of \mathcal{H} such that $dis(A_n, A) \rightarrow 0$ and $\|A_n^0(x)\| \leq c(1 + \|x\|)$ for some $c > 0$, all $n \in \mathbb{N}$ and $x \in D(A_n)$. Then for every $z \in D(A)$ there exists a sequence (ζ_n) such that*

$$\zeta_n \in D(A_n), \quad \zeta_n \rightarrow z \quad \text{and} \quad A_n^0(\zeta_n) \rightarrow A^0(z).$$

We end this section by the following discrete Gronwall's lemma.

Lemma 2.5 *Let (α_i) , (β_i) , (γ_i) and (a_i) be sequences of nonnegative real numbers such that $a_{i+1} \leq \alpha_i + \beta_i(a_0 + a_1 + \dots + a_{i-1}) + (1 + \gamma_i)a_i$ for $i \in \mathbb{N}$. Then*

$$a_j \leq \left(a_0 + \sum_{k=0}^{j-1} \alpha_k \right) \exp \left(\sum_{k=0}^{j-1} (k\beta_k + \gamma_k) \right) \quad \text{for } j \in \mathbb{N}^*.$$

We will establish our main result under the following hypotheses.

(H₁) There exist a function $\rho : I \rightarrow [0, +\infty[$, which is right-continuous on $[0, T[$ and nondecreasing with $\rho(0) = 0$ and $\rho(T) < +\infty$, and nonnegative constant γ such that

$$\text{dis}(A(t, x), A(s, y)) \leq d\rho([s, t]) + \gamma \|x - y\| \frac{d\lambda}{d\nu}(s) \quad \text{for } 0 \leq s \leq t \leq T, \quad x, y \in \mathcal{H},$$

and since $\frac{d\lambda}{d\nu}(s) \leq 1$, (H₁) implies that

$$\text{dis}(A(t, x), A(s, y)) \leq d\rho([s, t]) + \gamma \|x - y\| \quad \text{for } 0 \leq s \leq t \leq T, \quad x, y \in \mathcal{H}.$$

(H₂) There exists a nonnegative real constant c such that

$$\|A^0(t, x)y\| \leq c(1 + \|x\| + \|y\|) \quad \text{for } t \in I, \quad x \in \mathcal{H}, \quad y \in D(A(t, x)).$$

(H₃) For any bounded subset E of \mathcal{H} , $D(A(I \times E))$ is relatively ball-compact, i.e., its intersection with any closed ball of \mathcal{H} is relatively compact.

(H₄) $t \mapsto f(t, \cdot, \cdot)$ is $\mathcal{B}(I)$ -measurable.

(H₅) There is a nonnegative function $k \in L^1(I, \mathbb{R}; \lambda)$ such that

$$\|f(t, x, y) - f(t, x', y')\| \leq k(t)(\|x - x'\| + \|y - y'\|) \quad \forall t \in I, \quad \forall (x, y), (x', y') \in \mathcal{H} \times \mathcal{H}.$$

(H₆) There exists a nonnegative real constant m such that

$$\|f(t, x, y)\| \leq m(1 + \|x\| + \|y\|) \quad \forall (t, x, y) \in I \times \mathcal{H} \times \mathcal{H}.$$

3 Main result

Now we are able to state our main theorem.

Theorem 3.1 *Let for every $(t, x) \in I \times \mathcal{H}$, $A(t, x) : D(A(t, x)) \subset \mathcal{H} \rightrightarrows \mathcal{H}$ be a maximal monotone operator satisfying (H₁), (H₂) and (H₃). Let $f : I \times \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ such that (H₄), (H₅) and (H₆) are satisfied. Then for any $(u_0, v_0) \in \mathcal{H} \times D(A(0, u_0))$, there exists a BVRC solution $(u, v) : I \rightarrow \mathcal{H} \times \mathcal{H}$ to problem (P_f) .*

Proof. Following Castaing et al [27], we can choose a sequence $(\varepsilon_n)_{n \geq 1} \subset]0, 1]$ such that $\varepsilon_n \downarrow 0$ and a partition $0 = t_0^n < t_1^n < \dots < t_{q_n}^n = T$ of I (with $\lim_{n \rightarrow \infty} q_n = +\infty$), for which we have

$$\nu([t_i^n, t_{i+1}^n]) = |t_{i+1}^n - t_i^n| + d\rho([t_i^n, t_{i+1}^n]) \leq \varepsilon_n \quad \text{for } i = 0, \dots, q_n - 1, \quad (3.1)$$

we can take, for example, $\varepsilon_n = \frac{T}{q_n}$. For each $i \in \{0, \dots, q_n - 1\}$, put $I_i^n =]t_i^n, t_{i+1}^n]$,

$$\delta_{i+1}^n = d\rho([t_i^n, t_{i+1}^n]), \quad \eta_{i+1}^n = t_{i+1}^n - t_i^n, \quad \beta_{i+1}^n = \nu([t_i^n, t_{i+1}^n]). \quad (3.2)$$

For every $n \in \mathbb{N}$, put $u_0^n = u_0$, $v_0^n = v_0 \in D(A(0, u_0))$, and let us define by induction, sequences $(v_i^n)_{0 \leq i \leq q_n - 1}$, $(u_i^n)_{0 \leq i \leq q_n - 1}$ such that

$$v_{i+1}^n = J_{i+1}^n \left(v_i^n - \int_{t_i^n}^{t_{i+1}^n} f(s, u_i^n, v_i^n) d\lambda(s) \right) \quad (3.3)$$

where $J_{i+1}^n := J_{\beta_{i+1}^n}^{A(t_{i+1}^n, u_i^n)} = (Id_{\mathcal{H}} + \beta_{i+1}^n A(t_{i+1}^n, u_i^n))^{-1}$, and

$$u_{i+1}^n = u_i^n + \beta_{i+1}^n v_{i+1}^n. \quad (3.4)$$

From the properties of the resolvent, we have that $v_{i+1}^n \in D(A(t_{i+1}^n, u_i^n))$ and

$$-\frac{1}{\beta_{i+1}^n} \left(v_{i+1}^n - v_i^n + \int_{t_i^n}^{t_{i+1}^n} f(s, u_i^n, v_i^n) d\lambda(s) \right) \in A(t_{i+1}^n, u_i^n) v_{i+1}^n. \quad (3.5)$$

For $t \in [t_i^n, t_{i+1}^n[$, $i = 0, \dots, q_n - 1$, set

$$v_n(t) = v_i^n + \frac{\nu([t_i^n, t])}{\nu([t_i^n, t_{i+1}^n])} \left(v_{i+1}^n - v_i^n + \int_{t_i^n}^{t_{i+1}^n} f(s, u_i^n, v_i^n) d\lambda(s) \right) - \int_{t_i^n}^t f(s, u_i^n, v_i^n) d\lambda(s), \quad (3.6)$$

$$u_n(t) = u_i^n + \frac{\nu([t_i^n, t])}{\nu([t_i^n, t_{i+1}^n])} (u_{i+1}^n - u_i^n), \quad (3.7)$$

and $v_n(T) = v_{q_n}^n$, $u_n(T) = u_{q_n}^n$. So that $v_n(t_i^n) = v_i^n$, $u_n(t_i^n) = u_i^n$ and v_n, u_n are BVRC mappings on I .

Step 1. We prove in this step that the sequences (v_n) , (u_n) are bounded in norm and in variation.

Using Lemma 2.3, we have from (3.3), (H_1) , (H_2) and (H_6) , for $i = 0, 1, \dots, q_n - 1$ (keeping in mind that $\sqrt{ab} \leq \frac{a}{2} + b$ for nonnegative real numbers a, b)

$$\begin{aligned} \|v_{i+1}^n - v_i^n\| &\leq \left\| J_{i+1}^n \left(v_i^n - \int_{t_i^n}^{t_{i+1}^n} f(s, u_i^n, v_i^n) d\lambda(s) \right) - J_{i+1}^n(v_i^n) \right\| + \|J_{i+1}^n(v_i^n) - v_i^n\| \\ &\leq \int_{t_i^n}^{t_{i+1}^n} \|f(s, u_i^n, v_i^n)\| d\lambda(s) + \beta_{i+1}^n \|A^0(t_i^n, u_{i-1}^n) v_i^n\| + \text{dis}(A(t_{i+1}^n, u_i^n), A(t_i^n, u_{i-1}^n)) \\ &\quad + \sqrt{\beta_{i+1}^n (1 + \|A^0(t_i^n, u_{i-1}^n) v_i^n\|) \text{dis}(A(t_{i+1}^n, u_i^n), A(t_i^n, u_{i-1}^n))} \\ &\leq m(1 + \|u_i^n\| + \|v_i^n\|) \beta_{i+1}^n + (1 + c(1 + \|u_{i-1}^n\| + \|v_i^n\|)) \beta_{i+1}^n + \gamma \|u_i^n - u_{i-1}^n\| \\ &\quad + \sqrt{\beta_{i+1}^n (1 + c(1 + \|u_{i-1}^n\| + \|v_i^n\|)) (\beta_{i+1}^n + \gamma \|u_i^n - u_{i-1}^n\|)} \\ &\leq m(1 + \|u_i^n\| + \|v_i^n\|) \beta_{i+1}^n + (1 + c(1 + \|u_{i-1}^n\| + \|v_i^n\|)) \beta_{i+1}^n + \gamma \|u_i^n - u_{i-1}^n\| \\ &\quad + \beta_{i+1}^n \frac{1}{2} (1 + c(1 + \|u_{i-1}^n\| + \|v_i^n\|)) + \beta_{i+1}^n + \gamma \|u_i^n - u_{i-1}^n\|, \end{aligned}$$

that is,

$$\begin{aligned} \|v_{i+1}^n - v_i^n\| &\leq (m+1)(1 + \|u_i^n\| + \|v_i^n\|) \beta_{i+1}^n + \frac{3}{2}(1+c)(1 + \|u_{i-1}^n\| + \|v_i^n\|) \beta_{i+1}^n + 2\gamma \|u_i^n - u_{i-1}^n\| \\ &=: m_1(1 + \|u_i^n\| + \|v_i^n\|) \beta_{i+1}^n + c_1(1 + \|u_{i-1}^n\| + \|v_i^n\|) \beta_{i+1}^n + \gamma_1 \|u_i^n - u_{i-1}^n\|, \end{aligned} \quad (3.8)$$

which entails

$$\|v_{i+1}^n\| \leq \|v_i^n\| + m_1(1 + \|u_i^n\| + \|v_i^n\|) \beta_{i+1}^n + c_1(1 + \|u_{i-1}^n\| + \|v_i^n\|) \beta_{i+1}^n + \gamma_1 \|u_i^n - u_{i-1}^n\|. \quad (3.9)$$

On the other hand, we have from (3.4),

$$\|u_i^n - u_{i-1}^n\| = \beta_i^n \|v_i^n\| \quad (3.10)$$

and (using (3.1))

$$\begin{aligned} \|u_i^n\| &= \|u_{i-1}^n + \beta_i^n v_i^n\| = \|u_0 + \beta_1^n v_1^n + \beta_2^n v_2^n + \cdots + \beta_i^n v_i^n\| \\ &\leq \|u_0\| + \varepsilon_n (\|v_1^n\| + \|v_2^n\| + \cdots + \|v_i^n\|). \end{aligned} \quad (3.11)$$

Replacing these two last relations in (3.9), we get by the use of (3.1),

$$\begin{aligned} \|v_{i+1}^n\| &\leq \|v_i^n\| + m_1 \left(1 + \|u_0\| + \varepsilon_n (\|v_1^n\| + \|v_2^n\| + \cdots + \|v_i^n\|) + \|v_i^n\| \right) \varepsilon_n \\ &\quad + c_1 (1 + \|u_0\| + \varepsilon_n (\|v_1^n\| + \|v_2^n\| + \cdots + \|v_{i-1}^n\|) + \|v_i^n\|) \varepsilon_n + \gamma_1 \|v_i^n\| \varepsilon_n \\ &\leq (m_1 + c_1) (1 + \|u_0\|) \varepsilon_n + (m_1 + c_1) (\|v_1^n\| + \|v_2^n\| + \cdots + \|v_{i-1}^n\|) \varepsilon_n^2 \\ &\quad + \left(1 + (m_1 \varepsilon_n + m_1 + c_1 + \gamma_1) \varepsilon_n \right) \|v_i^n\|. \end{aligned}$$

Applying Lemma 2.5, we get for all $i \leq q_n$,

$$\begin{aligned} \|v_i^n\| &\leq \left(\|v_0\| + \sum_{j=0}^{i-1} (m_1 + c_1) (1 + \|u_0\|) \varepsilon_n \right) \exp \left(\sum_{j=0}^{i-1} (j(m_1 + c_1) \varepsilon_n^2 + (m_1 \varepsilon_n + m_1 + c_1 + \gamma_1) \varepsilon_n) \right) \\ &\leq \left(\|v_0\| + (m_1 + c_1) (1 + \|u_0\|) T \right) \exp \left((2 + T) m_1 + 2c_1 + \gamma_1 \right) =: m_2, \end{aligned}$$

using this last estimate in (3.11), we obtain

$$\|u_i^n\| \leq \|u_0\| + m_2 T,$$

and replacing in (3.10) and in (3.8), it comes that

$$\|u_{i+1}^n - u_i^n\| \leq m_2 \beta_{i+1}^n,$$

and

$$\|v_{i+1}^n - v_i^n\| \leq (m_1 + c_1) (1 + \|u_0\| + m_2 T + m_2) \beta_{i+1}^n + \gamma_1 m_2 \beta_i^n =: m_3 \beta_{i+1}^n + m_4 \beta_i^n.$$

Consequently,

$$\|u_i^n\| \leq M, \quad \|v_i^n\| \leq M \quad \forall 0 \leq i \leq q_n \quad (3.12)$$

and for all $0 \leq i < q_n - 1$,

$$\|u_{i+1}^n - u_i^n\| \leq M \nu([t_i^n, t_{i+1}^n]), \quad \|v_{i+1}^n - v_i^n\| \leq M (\nu([t_i^n, t_{i+1}^n]) + \nu([t_{i-1}^n, t_i^n])), \quad (3.13)$$

where $M = \max(m_2, m_3, m_4)$. Now, for $t \in [t_i^n, t_{i+1}^n[$, we have from (3.1), (3.6), (3.12), (3.13) and (H_6) ,

$$\|v_n(t) - v_i^n\| \leq 2M \varepsilon_n + 2m(1 + 2M) \varepsilon_n \leq 2M + 2m(1 + 2M) =: M_1,$$

and from (3.1), (3.7), (3.12) and (3.13), we get

$$\|u_n(t) - u_i^n\| \leq M.$$

These last relations give us, for all $n \in \mathbb{N}$,

$$\|v_n(t)\| \leq M + M_1 =: M_2 \quad \text{and} \quad \|u_n(t)\| \leq 2M \quad \forall t \in I,$$

that is

$$\sup_n \|v_n\| := \sup_n \left(\sup_{t \in I} \|v_n(t)\| \right) \leq M_2 \quad \text{and} \quad \sup_n \|u_n\| \leq 2M.$$

That is (u_n) , (v_n) are bounded in norm and in variation.

Step 2. Convergence of the sequences (u_n) , (v_n) and $(\frac{dv_n}{d\nu})$.
Define the functions $\phi_n, \theta_n : I \rightarrow I$ by

$$\phi_n(t) = t_i^n, \quad \theta_n(t) = t_{i+1}^n \quad \text{for } t \in]t_i^n, t_{i+1}^n], \quad i = 0, 1, \dots, q_n - 1, \quad \text{and} \quad \phi_n(0) = \theta_n(0) = 0,$$

and set for all $t \in I$,

$$B_n(t) = \sum_{i=0}^{q_n-1} \frac{1}{\nu([t_i^n, t_{i+1}^n])} \left(v_{i+1}^n - v_i^n + \int_{t_i^n}^{t_{i+1}^n} f(s, u_i^n, v_i^n) d\lambda(s) \right) \chi_{[t_i^n, t_{i+1}^n]}(t),$$

where $\chi_J(\cdot)$ is the characteristic function of the set $J \subset I$. Whence, by (3.6), it is clear that

$$v_n(t) = v_0 + \int_{[0,t]} B_n(\tau) d\nu(\tau) - \int_{[0,t]} f(\tau, u_n(\phi_n(\tau)), v_n(\phi_n(\tau))) d\lambda(\tau).$$

Since $\frac{d\lambda}{d\nu}$ is a density of λ w.r.t ν , then by (2.4), we get for every $t \in I$,

$$v_n(t) = v_0 + \int_{[0,t]} \left(B_n(\tau) - f(\tau, u_n(\phi_n(\tau)), v_n(\phi_n(\tau))) \frac{d\lambda}{d\nu}(\tau) \right) d\nu(\tau),$$

so that, $\frac{dv_n}{d\nu}$ is a density of the vector measure dv_n w.r.t ν and for ν -almost every $t \in I$,

$$\frac{dv_n}{d\nu}(t) = B_n(t) - f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t),$$

that is, by the definition of B_n , for ν -almost every $t \in I$,

$$\begin{aligned} & \frac{dv_n}{d\nu}(t) + f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) \\ &= \sum_{i=0}^{q_n-1} \frac{1}{\nu([t_i^n, t_{i+1}^n])} \left(v_{i+1}^n - v_i^n + \int_{t_i^n}^{t_{i+1}^n} f(s, u_i^n, v_i^n) d\lambda(s) \right) \chi_{[t_i^n, t_{i+1}^n]}(t). \end{aligned} \quad (3.14)$$

From (H_6) , (3.12) and (3.14), it results that

$$\left\| \frac{dv_n}{d\nu}(t) + f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) \right\| \leq M + m(1 + 2M) =: M_3 \quad \nu - a.e. t \in I. \quad (3.15)$$

On the other hand, we know that $\nu = d\rho + \lambda$, whence $\frac{d\lambda}{d\nu}(t) \leq 1$. So that, by (H_6) and (3.12), it follows that

$$\left\| f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) \right\| \leq m(1 + 2M) =: M_4 \quad \nu - a.e. t \in I, \quad (3.16)$$

consequently, there is a Borel subset $J \subset I$ with $\nu(J) = 0$, such that

$$\left\| \frac{dv_n}{d\nu}(t) \right\| \leq M_3 + M_4 =: M_5 \quad \forall t \in I \setminus J. \quad (3.17)$$

Now, observe that by (3.5) and (3.14), for each $n \in \mathbb{N}$, there is a Borel subset $J_n \subset I$ with $\nu(J_n) = 0$ such that

$$\frac{dv_n}{d\nu}(t) + f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) \in -A(\theta_n(t), u_n(\phi_n(t)))v_n(\theta_n(t)) \quad \forall t \in I \setminus J_n, \quad (3.18)$$

moreover,

$$v_n(\theta_n(t)) \in D(A(\theta_n(t), u_n(\phi_n(t)))) \quad \forall t \in I. \quad (3.19)$$

We will show in the following, that (v_n) converges uniformly on I . On the one hand, since $\frac{dv_n}{d\nu}$ is a density of the vector measure dv_n w.r.t the measure ν , we have from (2.5), for all $s, t \in I$ ($s \leq t$),

$$v_n(t) = v_n(s) + \int_{]s, t]} \frac{dv_n}{d\nu}(s) d\nu(s),$$

that is, using relation (3.17), we get

$$\|v_n(t) - v_n(s)\| \leq M_5 \nu([s, t]), \quad (3.20)$$

which means that (v_n) is equi-right-continuous with bounded variation. On the other hand, from (3.12) and (3.19), it is clear that for any $t \in I$, we have

$$(v_n(\theta_n(t))) \subset D(I \times M\overline{B}_{\mathcal{H}}) \cap M\overline{B}_{\mathcal{H}},$$

which shows that the sequence $(v_n(\theta_n(t)))$ is relatively compact (see (H_3)). From (3.20), we have that for all $t \in I$,

$$\|v_n(\theta_n(t)) - v_n(t)\| \leq M_5 d\nu([t, \theta_n(t)]) \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (3.21)$$

whence, we conclude that $(v_n(t))$ is also relatively compact. Using Helly-Banach's Theorem (see [42]), we may assume the existence of a BVRC mapping v such that $(v_n(t))$ converges strongly to $v(t)$ for every $t \in I$. This together with (3.20), show that

$$\|v(t) - v(s)\| \leq M_5 \nu([s, t]), \quad (3.22)$$

and from (3.21)

$$\|v_n(\theta_n(t)) - v(t)\| \leq \|v_n(\theta_n(t)) - v_n(t)\| + \|v_n(t) - v(t)\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (3.23)$$

As regards the sequence (u_n) , let us set for all $t \in I$,

$$\alpha_n(t) = \sum_{i=0}^{q_n-1} \frac{1}{\nu([t_i^n, t_{i+1}^n])} (u_{i+1}^n - u_i^n) \chi_{[t_i^n, t_{i+1}^n]}(t),$$

and remark that by relation (3.4), $\alpha_n(t) = v_n(\theta_n(t))$ for ν -a.e. $t \in I$. Consequently, from (3.7), it comes that for all $t \in I$,

$$u_n(t) = u_0 + \int_{]0,t]} \alpha_n(\tau) d\nu(\tau) = u_0 + \int_{]0,t]} v_n(\theta_n(\tau)) d\nu(\tau). \quad (3.24)$$

Using relations (3.12) and (3.21) and the strong convergence in \mathcal{H} of $(v_n(t))$ to $v(t)$, we obtain by Lebesgue dominated convergence, that

$$\lim_{n \rightarrow \infty} u_n(t) = u_0 + \int_{]0,t]} v(\tau) d\nu(\tau) =: u(t)$$

so that, u is a BVRC mapping, $(u_n(t))$ converges strongly to $u(t)$ for all $t \in I$, and $\frac{du}{d\nu}$ is a density of the vector measure du w.r.t ν and for ν -almost every $t \in I$,

$$\frac{du}{d\nu}(t) = v(t). \quad (3.25)$$

Now, observe that the sequence $(\frac{dv_n}{d\nu})$ is bounded in $L^2(I, \mathcal{H}; \nu)$ due to relation (3.17), so that it converges weakly in $L^2(I, \mathcal{H}; \nu)$ to some mapping $w \in L^2(I, \mathcal{H}; \nu)$. In particular, for all $t \in I$

$$\int_{]0,t]} \frac{dv_n}{d\nu}(s) d\nu(s) \longrightarrow \int_{]0,t]} w(s) d\nu(s) \quad \text{weakly in } \mathcal{H}.$$

Since $\frac{dv_n}{d\nu}$ is a density of the vector measure dv_n w.r.t the measure ν , we have for all $t \in I$,

$$v_n(t) = v_0 + \int_{]0,t]} \frac{dv_n}{d\nu}(s) d\nu(s)$$

and since for any $t \in I$, $(v_n(t))$ converges strongly in \mathcal{H} and then weakly to $v(t)$, we deduce by what precedes that

$$v(t) = v_0 + \int_{]0,t]} w(s) d\nu(s) \quad \forall t \in I.$$

This shows that $\frac{dv}{d\nu} = w$ for ν -a.e. $t \in I$. Consequently, $(\frac{dv_n}{d\nu})$ converges weakly in $L^2(I, \mathcal{H}; \nu)$ to $\frac{dv}{d\nu}$.

Step 3. Existence of solution.

First, observe that from (3.19), $v_n(\theta_n(t)) \in D(A(\theta_n(t), u_n(\phi_n(t))))$ for all $t \in I$. Also, from (H_1) , we have that for all $t \in I$,

$$\text{dis}(A(\theta_n(t), u_n(\phi_n(t))), A(t, u(t))) \leq |\rho(\theta_n(t)) - \rho(t)| + \gamma \|u_n(\phi_n(t)) - u(t)\| \frac{d\lambda}{d\nu}(t),$$

but from (3.24), it is clear that

$$\|u_n(\phi_n(t)) - u(t)\| \leq \|u_n(\phi_n(t)) - u_n(t)\| + \|u_n(t) - u(t)\| \leq M\nu([\phi_n(t), t]) + \|u_n(t) - u(t)\|,$$

so, from (2.7), it comes that

$$\|u_n(\phi_n(t)) - u(t)\| \frac{d\lambda}{d\nu}(t) \leq (M\nu([\phi_n(t), t]) + \|u_n(t) - u(t)\|) \frac{d\lambda}{d\nu}(t) \longrightarrow 0 \text{ as } n \rightarrow \infty. \quad (3.26)$$

This shows that

$$\text{dis}(A(\theta_n(t), u_n(\phi_n(t))), A(t, u(t))) \longrightarrow 0 \text{ as } n \rightarrow \infty. \quad (3.27)$$

On the other hand, by (3.12) and (H_2) , we have that the sequence $(A^0(\theta_n(t), u_n(\phi_n(t)))v_n(\theta_n(t)))$ is bounded in \mathcal{H} , and hence it is weakly relatively compact. Using this fact, (3.23) and (3.26), we conclude by Lemma 2.2, that $v(t) \in D(A(t, u(t)))$ for all $t \in I$.

Next, remember that the sequence $(\frac{dv_n}{d\nu})$ converges weakly in $L^2(I, \mathcal{H}; \nu)$ to $\frac{dv}{d\nu}$, so that by Mazur's theorem, there is a sequence (ξ_n) such that for each $n \in \mathbb{N}$, $\xi_n \in \text{co}\{\frac{dv_k}{d\nu}; k \geq n\}$, and (ξ_n) converges strongly in $L^2(I, \mathcal{H}; \nu)$ to $\frac{dv}{d\nu}$. Whence, there is a subsequence (ξ_{n_j}) , which converges ν -almost every where to $\frac{dv}{d\nu}$. This means the existence of a Borel subset $J' \subset I$, with $\nu(J') = 0$ and for $t \in I \setminus J'$,

$$\xi_{n_j}(t) \longrightarrow \frac{dv}{d\nu}(t) \in \bigcap_n \overline{\text{co}}\{\frac{dv_k}{d\nu}; k \geq n\}.$$

This implies, by (2.1), that for any fixed $\eta \in \mathcal{H}$,

$$\langle \frac{dv}{d\nu}(t), \eta \rangle \leq \limsup_{n \rightarrow \infty} \langle \frac{dv_n}{d\nu}(t), \eta \rangle. \quad (3.28)$$

On the other hand, we have from (H_5) and (3.20)

$$\begin{aligned} & \|f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) - f(t, u(t), v(t)) \frac{d\lambda}{d\nu}(t)\| \\ & \leq k(t)(\|u_n(\phi_n(t)) - u(t)\| + \|v_n(\phi_n(t)) - v(t)\|) \frac{d\lambda}{d\nu}(t) \\ & \leq k(t)(\|u_n(\phi_n(t)) - u(t)\| + \|v_n(\phi_n(t)) - v_n(t)\| + \|v_n(t) - v(t)\|) \frac{d\lambda}{d\nu}(t) \\ & \leq k(t)(\|u_n(\phi_n(t)) - u(t)\| + M_5\nu([\phi_n(t), t]) + \|v_n(t) - v(t)\|) \frac{d\lambda}{d\nu}(t), \end{aligned}$$

whence by the convergence of $(v_n(t))$ to $v(t)$ and relations (2.7), (3.26), it results that for all $t \in I$,

$$f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) \longrightarrow f(t, u(t), v(t)) \frac{d\lambda}{d\nu}(t) \text{ as } n \rightarrow \infty. \quad (3.29)$$

Now, to prove that (u, v) is a solution to our considered problem, we will use Lemma 2.1. Since $v(t) \in D(A(t, u(t)))$ for all $t \in I$, we have to show that for ν -almost every fixed $t \in I$ and for any $z \in D(A(t, u(t)))$

$$\langle A^0(t, u(t))z + \frac{dv}{d\nu}(t) + f(t, u(t), v(t)) \frac{d\lambda}{d\nu}(t), z - v(t) \rangle \geq 0.$$

Indeed, let $t \in I$. By (H_2) , using Lemma 2.4, we can ensure the existence of a sequence $(\zeta_n)_n$, such that

$$\zeta_n \in D(A(\theta_n(t), u_n(\phi_n(t)))), \quad \zeta_n \longrightarrow z \quad \text{and} \quad A^0(\theta_n(t), u_n(\phi_n(t)))\zeta_n \longrightarrow A^0(t, u(t))z. \quad (3.30)$$

Since $A(\theta_n(t), u_n(\phi_n(t)))$ is monotone, using (3.18), we have for each $n \in \mathbb{N}$ and $t \in I \setminus J_n$,

$$\langle v_n(\theta_n(t)) - \zeta_n, \frac{dv_n}{d\nu}(t) + f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) + A^0(\theta_n(t), u_n(\phi_n(t)))\zeta_n \rangle \leq 0. \quad (3.31)$$

Next, let $t \in I \setminus (J \cup J' \cup (\bigcup_n J_n))$. From (3.28), we have

$$\langle \frac{dv}{d\nu}(t), v(t) - z \rangle \leq \limsup_{n \rightarrow \infty} \langle \frac{dv_n}{d\nu}(t), v(t) - z \rangle,$$

and since from (3.17) and (3.31), we have for each $n \in \mathbb{N}$,

$$\begin{aligned} & \langle \frac{dv_n}{d\nu}(t), v(t) - z \rangle \leq \langle \frac{dv_n}{d\nu}(t), v(t) - v_n(\theta_n(t)) \rangle + \langle \frac{dv_n}{d\nu}(t), v_n(\theta_n(t)) - \zeta_n \rangle + \langle \frac{dv_n}{d\nu}(t), \zeta_n - z \rangle \\ & \leq M_5(\|v(t) - v_n(\theta_n(t))\| + \|\zeta_n - z\|) + \langle \frac{dv_n}{d\nu}(t), v_n(\theta_n(t)) - \zeta_n \rangle \\ & \leq M_5(\|v(t) - v_n(\theta_n(t))\| + \|\zeta_n - z\|) \\ & + \langle f(t, u_n(\phi_n(t)), v_n(\phi_n(t))) \frac{d\lambda}{d\nu}(t) + A^0(\theta_n(t), u_n(\phi_n(t)))\zeta_n, \zeta_n - v_n(\theta_n(t)) \rangle \end{aligned}$$

we conclude, using (3.23), (3.29) and (3.30) that

$$\langle \frac{dv}{d\nu}(t), v(t) - z \rangle \leq \langle f(t, u(t), v(t)) \frac{d\lambda}{d\nu}(t) + A^0(t, u(t))z, z - v(t) \rangle, \quad (3.32)$$

which, with the fact that $v(t) \in D(A(t, u(t)))$, $u(0) = u_0$, $v(0) = v_0$ and (3.25), means that (u, v) is a BVRC solution to our problem (P_f) . \square

Remark 3.1 *If the operators depend only on the time and satisfy the following hypotheses:*
 (H'_1) *There exists a function $\rho : I \longrightarrow [0, +\infty[$, which is right-continuous on $[0, T[$ and nondecreasing with $\rho(0) = 0$ and $\rho(T) < +\infty$ such that*

$$\text{dis}(A(t), A(s)) \leq d\rho([s, t]) \quad \text{for } 0 \leq s \leq t \leq T.$$

(H'_2) *There exists a nonnegative real constant c such that*

$$\|A^0(t)x\| \leq c(1 + \|x\|) \quad \text{for } t \in I, x \in D(A(t)).$$

(H'_3) *For any $t \in I$, $D(A(t))$ is relatively ball-compact.*

Our theorem reads as

Corollary 3.1 *Let for every $t \in I$, $A(t) : D(A(t)) \subset \mathcal{H} \rightrightarrows \mathcal{H}$ be a maximal monotone operator satisfying (H'_1) , (H'_2) and (H'_3) . Let $f : I \times \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ such that (H_4) , (H_5) and (H_6) are satisfied. Then for any $(u_0, v_0) \in \mathcal{H} \times D(A(0))$, there exists a BVRC solution $(u, v) : I \rightarrow \mathcal{H} \times \mathcal{H}$ to the problem*

$$\begin{cases} u(0) = u_0, v(0) = v_0 \in D(A(0)); \\ v(t) \in D(A(t)) \quad \forall t \in I; \\ \frac{du}{d\nu}(t) = v(t) \quad d\nu - \text{a.e. } t \in I; \\ -\frac{dv}{d\nu}(t) \in A(t)v(t) + f(t, u(t), v(t))\frac{d\lambda}{d\nu}(t) \quad d\nu - \text{a.e. } t \in I. \end{cases}$$

Remark 3.2 *In particular, if we consider $A(t, x) = N_{C(t, x)}$, the normal cone, in the sense of convex analysis, to the closed convex set $C(t, x)$, then under the following hypotheses: (H''_1) There exists a function $\rho : I \rightarrow [0, +\infty[$, which is right-continuous on $[0, T[$ and nondecreasing with $\rho(0) = 0$ and $\rho(T) < +\infty$ and a nonnegative real constant γ such that*

$$|d_{C(t, x)}(z) - d_{C(s, y)}(z)| \leq d\rho([s, t]) + \gamma\|x - y\|\frac{d\lambda}{d\nu}(s) \quad \text{for } 0 \leq s \leq t \leq T, \quad x, y, z \in \mathcal{H}.$$

(H''_2) *For any bounded subset E of \mathcal{H} , $C(I \times E)$ is relatively ball-compact. We can derive the following corollary.*

Corollary 3.2 *Let $C : I \times \mathcal{H} \rightrightarrows \mathcal{H}$ be a set-valued map with nonempty, closed and convex values satisfying (H''_1) and (H''_2) . Let $f : I \times \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{H}$ such that (H_4) , (H_5) and (H_6) are satisfied. Then for any $(u_0, v_0) \in \mathcal{H} \times C(0, u_0)$, there exists a BVRC solution $(u, v) : I \rightarrow \mathcal{H} \times \mathcal{H}$ to the problem*

$$\begin{cases} u(0) = u_0, v(0) = v_0; \\ v(t) \in C(t, u(t)) \quad \forall t \in I; \\ \frac{du}{d\nu}(t) = v(t) \quad d\nu - \text{a.e. } t \in I; \\ -\frac{dv}{d\nu}(t) \in N_{C(t, u(t))}(v(t)) + f(t, u(t), v(t))\frac{d\lambda}{d\nu}(t) \quad d\nu - \text{a.e. } t \in I. \end{cases}$$

References

- [1] S. Adly, T. Haddad and L. Thibault, Convex sweeping process in the framework of measure differential inclusions and evolution variational inequalities. Math. Program, 148 (1-2, Ser. B) (2014) 5-47.
- [2] S. Adly and B. K. Le, Unbounded second-order state-dependent Moreau's sweeping processes in Hilbert spaces. J. Optim. Theory Appl. 169, (2016) 407-423.
- [3] S. Adly and B. K. Le, Unbounded state-dependent sweeping process with perturbations in uniformly convex and q-uniformly smooth Banach spaces. Numer. Algebra Control Optim. 8, 8195 (2018).

- [4] S. Adly and F. Nacry, An existence result for discontinuous second-order nonconvex state-dependent sweeping process. *Appl. Math. Optim.* DOI 10.1007/s00245-017-9446-9 (2017).
- [5] S. Adly, F. Nacry and L. Thibault, Discontinuous sweeping process with prox-regular sets. *ESAIM: Control, Optimizatio and Calculus of Variation.* 23 (4) (2017) 1293-1329.
- [6] F. Aliouane, D. Azzam-Laouir, A second order differential inclusion with proximal normal cone in Banach spaces, *Topol Methods Nonlinear Anal.* 44(1) 143-160 (2014).
- [7] F. Aliouane, D. Azzam-Laouir, Second-order sweeping process with a lipschitz perturbation. *J. Math. Anal. Appl.* 452, 729-746 (2017).
- [8] F. Aliouane, D. Azzam-Laouir, C. Castaing and M.D.P. Monteiro Marques, Second-order time and state sweeping process in Hilbert space, *J Optimiz Theory App.* 182, 153-188 (2019).
- [9] D. Azzam-Laouir, A measure differential inclusion with time-dependent maximal monotone operators. *ArXiv preprint, arXiv:2103.07376* (2021) 1-15.
- [10] D. Azzam-Laouir, Mixed semicontinuous perturbation of a second order nonconvex sweeping process. *Electronic Journal of Qualitative Theory of Differential Equations*, 37, 1-9 (2008).
- [11] D. Azzam-Laouir and S. Izza, Existence of solutions for second-order perturbed nonconvex sweeping process. *Computers & Mathematics with Applications.* 62, 1736-1744 (2011).
- [12] D. Azzam-Laouir, W. Belhoula, C. Castaing and M.D.P. Monteiro Marques, Perturbed evolution problems with absolutely continuous variation in time and applications. *J. Fixed Point Theory Appl.* (2019) 21: 40.
- [13] D. Azzam-Laouir, W. Belhoula, C. Castaing and M.D.P. Monteiro Marques, Multi-valued perturbation to evolution problems involving time dependent maximal monotone operators. *Evolution Equations & Control Theory*, 9(1) (2020) 219-254.
- [14] D. Azzam-Laouir, I. Boutana-Harid, Mixed semicontinuous perturbation to an evolution problem with time-dependent maximal monotone operator. *J. Nonlinear and Convex Analysis.* Vol 20, Number 1 (2018) 39-52.
- [15] D. Azzam-Laouir, C. Castaing and M.D.P. Monteiro Marques, Perturbed evolution problems with continuous bounded variation in time and applications. *Set-Valued Var. Anal.* Volume 26, No. 3 (2018) 693-728.
- [16] D. Azzam-Laouir, C. Castaing and M.D.P. Monteiro Marques, BV right-continuous solutions of differential inclusions involving time dependent maximal monotone operators. *ArXiv: 2103.01113v1* (2021) 1-43.
- [17] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach Spaces.* Noordhoff Int. Publ. Leyden, (1976).

- [18] H. Benabdellah, C. Castaing and M.A. Gamal Ibrahim, BV Solutions of multivalued differential equations on closed moving sets in Banach spaces. Banach center publications, Vol 32, Institute of Mathematics, Polish academy of Sciences, Warszawa (1995).
- [19] H. Benabdellah, C. Castaing, A. Salvadori and A. Syam, Nonconvex sweeping process. Journal of Applied Analysis, Vol 2, No2 (1996) 17-40.
- [20] M. Benguessoum, D. Azzam-Laouir and C. Castaing, On a time and state dependent maximal monotone operator coupled with a sweeping process with perturbations. Set-Valued Var. Anal, 29 (2021) 191-219.
- [21] M. Bounkhel, Existence results for second order convex sweeping processes in p-uniformly smooth and q-uniformly convex Banach spaces, Electron J Qual Theo. 27, 1-10 (2012).
- [22] M. Bounkhel, D. Laouir-Azzam, Existence results for second-order nonconvex sweeping processes. Set-Valued Var Anal. 12(3), 291-318 (2004).
- [23] H. Brezis, Opérateurs maximaux monotones, North Holland Publ. Compagny, Amsterdam-London, (1973).
- [24] Camlibel K, Ianelli L, Tanwani A. Convergence of proximal solutions for evolution inclusions with time-dependent maximal monotone operators. Math. Programming, Series A. <https://doi.org/10.1007/s10107-021-01666-7> (2021)
- [25] C. Castaing, Quelques problèmes d'évolution du second ordre. Sem. Anal. Convexe. Montpellier. Expose No. 5. (1988).
- [26] C. Castaing, A.G. Ibrahim, M.F. Yarou, Some contributions to nonconvex sweeping process. J. Nonlinear Convex Anal. Vol. 10, No. 1, 1-20 (2009).
- [27] C. Castaing, M.D.P. Monteiro Marques, Evolution problems associated with nonconvex closed moving sets with bounded variation. Portugal. Math. 53 (1996) 73-87.
- [28] C. Castaing, M. D. P. Monteiro Marques and P. Raynaud de Fitte, Second order evolution problems with time dependent maximal monotone operator. Adv. Math. Econ. Vol 23 (2018).
- [29] J.F. Edmond and L. Thibault, BV solutions of non convex sweeping process differential inclusion with perturbation. J. Differential Equations. 22 (2006) 135-179.
- [30] N. Kenmochi, Solvability of nonlinear evolution equations with time-dependent constraints and applications. Bull. Fac. Educ. Chiba Univ. 30 (1981).
- [31] M. Kunze and M.D.P. Monteiro Marques, BV solutions to evolution problems with time-dependent domains. Set-Valued Anal. 5 (1997) 57-72.
- [32] B.K. Le, Well-posedness and nonsmooth Lyapunov pairs for state-dependent maximal monotone differential inclusions. Optimization, 69 (6) (2020) 1187-1217.

- [33] M.D.P. Monteiro Marques, Differential inclusions nonsmooth mechanical problems, shocks and dry friction. Progress in Nonlinear Differential Equations and Their Applications, Birkhauser, Vol 9 (1993).
- [34] J.J. Moreau, Raffle par un convexe variable I, Sem. Anal. Convexe Montpellier (1971) Exposé 15.
- [35] J.J. Moreau, Evolution problem associated with a moving convex set in a Hilbert space. J. Differential. Equations. Vol 26 (1977) 347-374.
- [36] J.J. Moreau, Bounded variation in time, in Topics in Non-smooth Mechanics (J.J. Moreau, P.D. Panagiotopoulos, G. Strang, Eds.), Birkhäuser Verlag, Basel-Boston-Berlin (1988).
- [37] J.J. Moreau, Sur les mesures différentielles des fonctions vectorielles à variation bornée. Sem. Anal. Convexe. Montpellier (1975) Exposé 17.
- [38] J.J. Moreau and M. Valadier, A chain rule involving vector functions of bounded variations. J. Functional Analysis, Vol 74, 2 (1987) 333-345.
- [39] F. Nacry, J. Noel and L. Thibault, On first and second order state-dependent prox-regular sweeping process. Preprint (2020).
- [40] J. Noel, Second-order general perturbed sweeping process differential inclusion. J. Fixed Point Theory Appl. 20, 133 (2018).
- [41] N.H. Pavel, Nonlinear evolution operators and semigroups, Lecture Notes in Mathematics, 1260. Springer, Nex York (1987).
- [42] J.E. Porter, Helly selection principle for functions of bounded P -variation. Rocky Mountain Journal of Mathematics, Vol 25, Number 2 (2005).
- [43] S. Saidi, On a second-order functional evolution problem with time and state dependent maximal monotone operators. Evolution Equations & Control Theory, (2021).
- [44] M. Schatzman, A class of nonlinear differential equations of second order in time, Non-linear Anal-Theor. 2 (1978).
- [45] F. Selamnia, D. Azzam-Laouir, M.D.P. Monteiro Marques, Evolution problems involving state-dependent maximal monotone operators. Appl. Anal. (2020) <https://doi.org/10.1080/00036811.2020.1738401>
- [46] L. Thibault, Moreau sweeping process with bounded retraction. J. Convex Anal. 23, 4 (2016) 1051-1098.
- [47] A.A. Tolstonogov, BV solutions of a convex sweeping process with a composed perturbation. Evolution Equations & Control theory. (2021).

- [48] A.A. Tolstonogov, BV continuous solutions of an evolution inclusion with maximal monotone operator and non-convex valued perturbation. Existence theorem. Set-Valued Var. Anal. 29 (2021) 29-60.
- [49] A.A. Vladimirov, Nonstationnary dissipative evolution equation in Hilbert space. Non-linear Anal. 17 (1991) 499-518
- [50] I.I. Vrabie, Compactness methods for nonlinear evolution equations. Pitman Monographs and Surveys in Pure and Applied mathematics, Longman Scientific and Technical, John Wiley and Sons, Inc. New York, Vol 32 (1987).