

On noncommutative weak Orlicz-Hardy spaces

Turdebek N. Bekjan^{a,*}, Madi Raikhan^b

^a*Faculty of Mechanics and Mathematics, L.N. Gumilyov Eurasian National University,
Nur-Sultan 010008, Kazakhstan.*

^b*Astana IT University, Nur-Sultan 010000, Kazakhstan*

Abstract

We introduce noncommutative weak Orlicz spaces associated with a weight and study their properties. We also define noncommutative weak Orlicz-Hardy spaces and characterize their dual spaces.

Keywords: noncommutative Lorentz space, noncommutative Marcinkiewicz space, weak noncommutative Orlicz space, noncommutative weak Orlicz-Hardy space.

2010 MSC: 46L52, 47L05.

1. Introduction

Al-Rashed and Zegarliński[1] introduced the noncommutative Orlicz spaces associated to a normal faithful state on a semifinite von Neumann algebra. In [2], the authors considered a certain class of noncommutative Orlicz spaces, associated with arbitrary faithful normal locally-finite weights on a semi-finite von Neumann algebra \mathcal{M} . In [20], the authors have investigated weak version of Orlicz spaces and proved the Burkholder-Gundy inequalities of martingales for this weak Orlicz spaces. The weak noncommutative Orlicz spaces were investigated in [3] and it was used for the theory of noncommutative martingales. In this paper, we extend the results of [2] to the weak noncommutative Orlicz space case.

The dual spaces of commutative weak L_p -spaces were characterized in [10, 11], its noncommutative versions proved in [9, 16]. In [8, 9], Ciach introduced noncommutative Lorentz space and noncommutative Marcinkiewicz space, and discussed their dual spaces. The aim of this paper is to define noncommutative weak Orlicz-Hardy spaces and characterize their dual spaces.

The paper is organized as follows. In Section 2 some necessary definitions and notations are collected including the weak Orlicz spaces and the noncommutative weak Orlicz spaces. Using relationship between noncommutative weak Orlicz spaces and noncommutative Marcinkiewicz space, and Ciach's results to give dual spaces of weak noncommutative Orlicz spaces. The noncommutative weak Orlicz spaces associated with a weight are studied in Section 3. In Section 4, we characterized the dual spaces of noncommutative weak Orlicz-Hardy spaces.

*Corresponding author

Email addresses: bekjant@yahoo.com (Turdebek N. Bekjan), raikhan\$_-\$m@enu.kz (Madi Raikhan)

2. Preliminaries

Let $\Omega = [0, \gamma)$ ($0 < \gamma \leq \infty$) be equipped with the usual Lebesgue measure μ . We denote by $L_0(\Omega)$ the space of μ -measurable real-valued functions f on (Ω) such that $\mu(\{\omega \in \Omega : |f(\omega)| > s\}) < \infty$ for some s . The decreasing rearrangement function $f^* : [0, \infty) \mapsto [0, \infty]$ for $f \in L_0(\Omega)$ is defined by

$$f^*(t) = \inf\{s > 0 : \mu(\{\omega \in \Omega : |f(\omega)| > s\}) \leq t\}$$

for $t \geq 0$.

The classical weak L_p -space $L_{p,\infty}(\Omega)$ ($0 < p < \infty$) is defined as the set of all measurable functions f on Ω such that

$$\|f\|_{L_{p,\infty}} = \sup_{t>0} t^{\frac{1}{p}} f^*(t) < \infty.$$

However, for $p > 1$ $L_{p,\infty}(\Omega)$ can be renormed as a Banach space by

$$f \mapsto \sup_{t>0} t^{-1+\frac{1}{p}} \int_0^t f^*(s) ds.$$

We refer to [15] for more information about weak L_p -spaces.

A function $\Phi : (-\infty, \infty) \rightarrow [0, \infty)$ is called an N-function if it satisfies the following conditions: (i) Φ is even and convex, (ii) $\Phi(t) = 0$ iff $t = 0$, (iii) $\lim_{t \rightarrow 0} \frac{\Phi(t)}{t} = 0$, $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} = +\infty$.

Let $\phi(t)$ be the left derivative of Φ . Then $\phi(t)$ is left continuous, nondecreasing on $(0, \infty)$ and satisfies: $0 < \phi(t) < \infty$ for $0 < t < \infty$, $\phi(0) = 0$ and $\lim_{t \rightarrow \infty} \phi(t) = \infty$. The left inverse of ϕ ($\psi(s) = \inf\{t > 0 : \phi(t) > s\}$ for $s > 0$) will be denoted by ψ . We define a complementary N-function Ψ of Φ by

$$\Psi(|s|) = \int_0^{|s|} \psi(v) dv.$$

It is clear that Φ is the complementary N-function of Ψ . We call (Φ, Ψ) a pair of complementary N-functions.

Let (Φ, Ψ) be a pair of complementary N-functions, with inverses Φ^{-1} , Ψ^{-1} (which are uniquely defined on $[0, \infty)$). Then

$$t < \Phi^{-1}(t)\Psi^{-1}(t) < 2t, \quad t > 0. \quad (2.1)$$

An N-function Φ is said to satisfy the Δ_2 -condition for all t , written as $\Phi \in \Delta_2$, if there is $K > 2$ such that $\Phi(2t) \leq K\Phi(t)$ for all $t \geq 0$. Φ is called to satisfy the ∇_2 -condition for all t , written as $\Phi \in \nabla_2$, if there is a constant $c > 1$ such that $\Phi(t) \leq \frac{1}{2c}\Phi(ct)$ for all $t \geq 0$. For a pair of complementary N-functions (Φ, Ψ) , we have that $\Phi \in \Delta_2$ if and only if $\Psi \in \nabla_2$ (see [26, Theorem 2]).

Let (Φ, Ψ) is a pair of complementary N-functions. Then the Orlicz space on Ω associated with Φ defined by

$$L_\Phi(\Omega) = \left\{ f \in L_0(\Omega) : \int_0^\infty \Phi(|af(t)|) dt < \infty \text{ for some } a > 0 \right\}.$$

We define

$$\|f\|_\Phi = \inf \left\{ c > 0 : \int_0^\infty \Phi\left(\frac{f(t)}{c}\right) dt \leq 1 \right\}.$$

Then for any $f \in L_\Phi(\Omega)$,

$$\|f\|_\Phi \leq \sup \left\{ \left| \int_0^\infty f(t)g(t)dt \right| : \int_0^\infty \Psi(|g(t)|)dt \leq 1, g \in L^\Psi(\Omega) \right\} \leq 2\|f\|_\Phi.$$

For an N-function Φ , we define

$$a_\Phi = \inf_{t>0} \frac{t\Phi'(t)}{\Phi(t)} \quad \text{and} \quad b_\Phi = \sup_{t>0} \frac{t\Phi'(t)}{\Phi(t)}.$$

Then $1 \leq a_\Phi \leq b_\Phi \leq \infty$ and $\Phi \in \Delta_2$ if and only if $b_\Phi < \infty$. It is well-known that

$$L_\Phi(\Omega)^* = L_\Psi(\Omega), \quad L_\Psi(\Omega)^* = L_\Phi(\Omega), \quad (2.2)$$

with equivalent norms. We refer to [26] for the details on Orlicz spaces.

Now we consider the set of all measurable functions

$$L_{\Phi,\infty}(\Omega) = \{f \in L_0(\Omega) : \exists c > 0, \Phi\left(\frac{t}{c}\right)\mu(|f| > t) \leq 1, \forall t > 0\}$$

and denote

$$\|f\|_{\Phi,\infty} = \inf \{c > 0 : \Phi\left(\frac{t}{c}\right)\mu(|f| > t) \leq 1, \forall t > 0\}.$$

We call $L_{\Phi,\infty}(\Omega)$ is a weak Orlicz space. If $\Phi(t) = t^p$, then $L_{\Phi,\infty}(\Omega) = L_{p,\infty}(\Omega)$ (see [20] for more details).

Recall that

$$L_{\Phi,\infty}(\Omega) = \left\{ x \in L_0(\Omega) : \exists c > 0 \text{ such that } \sup_{t>0} t\Phi(f^*(t)/c) < \infty \right\},$$

and

$$\begin{aligned} \|x\|_{\Phi,\infty} &= \inf \{c > 0 : t\Phi(f^*(t)/c) \leq 1, \forall t > 0\} \\ &= \inf \left\{ c > 0 : \frac{1}{\Phi^{-1}(\frac{1}{t})} \mu_t(x)/c \leq 1, \forall t > 0 \right\} \end{aligned}$$

(see [3, Proposition 3.1]).

2.1. Noncommutative weak L_p spaces

We keep all notations introduced in the above. In rest of this paper, Φ will always denote an N-function and Ψ denote a complementary N-function of Φ , \mathcal{M} always denote a semifinite von Neumann algebra acting on a Hilbert space \mathbb{H} with a normal semifinite faithful trace τ ($\tau(1) = \gamma$).

For $0 < p < \infty$ let $L_p(\mathcal{M})$ denote the noncommutative L_p space with respect to (\mathcal{M}, τ) . As usual, we set $L_\infty(\mathcal{M}, \tau) = \mathcal{M}$ equipped with the operator norm. Also,

let $L_0(\mathcal{M})$ denote the topological $*$ -algebra of measurable operators with respect to (\mathcal{M}, τ) .

For $x \in L_0(\mathcal{M})$, we define

$$\lambda_s(x) = \tau(e_s^\perp(|x|)) \quad (s > 0) \quad \text{and} \quad \mu_t(x) = \inf\{s > 0 : \lambda_s(x) \leq t\} \quad (t > 0),$$

where $e_s^\perp(|x|) = e_{(s, \infty)}(|x|)$ is the spectral projection of $|x|$ associated with the interval (s, ∞) . We call the function $s \mapsto \lambda_s(x)$ is the distribution function of x and $\mu_t(x)$ is the generalized singular number of x . For simplicity, we denote by $\lambda(x)$ and $\mu(x)$ the two functions $s \mapsto \lambda_s(x)$ and $t \mapsto \mu_t(x)$, respectively. It is clear that both functions $\lambda(x)$ and $\mu(x)$ are decreasing and continuous from the right on $(0, \infty)$ (for further information, see [14]).

For $0 < p < \infty$, the noncommutative weak L_p space $L_{p, \infty}(\mathcal{M})$ is defined as the space of all measurable operators x such that

$$\|x\|_{p, \infty} = \sup_{t > 0} t^{\frac{1}{p}} \mu_t(x) < \infty.$$

Equipped with $\|\cdot\|_{p, \infty}$, $L_{p, \infty}(\mathcal{M})$ is a quasi-Banach space. However, for $p > 1$ $L_{p, \infty}(\mathcal{M})$ can be renormed as a Banach space by

$$x \mapsto \sup_{t > 0} t^{-1 + \frac{1}{p}} \int_0^t \mu_s(x) ds.$$

On the other hand, the quasi-norm admits the following useful description

$$\|x\|_{p, \infty} = \inf \{c > 0 : t(\mu_t(x)/c)^p \leq 1, \forall t > 0\}. \quad (2.3)$$

Also, we have a description in terms of distribution function as follows

$$\|x\|_{p, \infty} = \sup_{s > 0} s \lambda_s(x)^{\frac{1}{p}}. \quad (2.4)$$

Recall that noncommutative weak L_p spaces can be presented through noncommutative Lorenz spaces, for details see [12] and Xu [28].

2.2. Noncommutative weak Orlicz spaces

Let

$$L_\Phi(\mathcal{M}) = \{x \in L_0(\mathcal{M}) : \tau(\Phi(|x|)) = \int_0^{\tau(1)} \Phi(\mu_t(x)) dt < \infty\}$$

and

$$\|x\|_\Phi = \inf\{\lambda > 0 : \tau(\Phi(\frac{|x|}{\lambda})) \leq 1\}, \quad \forall x \in L_\Phi(\mathcal{M}).$$

Then $L_\Phi(\mathcal{M})$ is a Banach space. We call it is the noncommutative Orlicz space on (\mathcal{M}, τ) .

Definition 2.1. The noncommutative weak Orlicz space $L_{\Phi,\infty}(\mathcal{M})$ is defined as following:

$$L_{\Phi,\infty}(\mathcal{M}) = \left\{ x \in L_0(\mathcal{M}) : \sup_{t>0} t\Phi(\mu_t(x)) < \infty \right\},$$

equipped with

$$\|x\|_{\Phi,\infty} = \inf \{ c > 0 : t\Phi(\mu_t(x)/c) \leq 1, \forall t > 0 \}.$$

If $\Phi(t) = t^p$ with $1 \leq p < \infty$, then $L_{\Phi,\infty}(\mathcal{M})$ is the noncommutative weak L_p -space.

Recall that $L_{\Phi,\infty}(\mathcal{M})$ is a quasi-Banach space, and for any $x \in L_{\Phi,\infty}(\mathcal{M})$

$$\|x\|_{\Phi,\infty} = \inf \left\{ c > 0 : \frac{1}{\Phi^{-1}(\frac{1}{t})} \mu_t(x)/c \leq 1, \forall t > 0 \right\} = \sup_{t>0} \frac{1}{\Phi^{-1}(\frac{1}{t})} \mu_t(x).$$

For any $c > 0$ we have that

$$\sup_{t>0} t\Phi(\mu_t(x)/c) = \sup_{s>0} \lambda_s(x)\Phi(s/c), \quad \forall x \in L_0(\mathcal{M}). \quad (2.5)$$

For more information on noncommutative weak Orlicz spaces, see [3].

For any $x \in L_0(\mathcal{M})$, set $\tilde{\mu}_t(x) = \int_0^t \mu_s(x) ds$. Then $\mu_t(x) \leq \tilde{\mu}_t(x)$ for all $t > 0$ and the map $x \mapsto \tilde{\mu}(x)$ is a sublinear operator from $L_0(\mathcal{M})$ to $L_0(\Omega)$ ($a = \tau(1)$).

Proposition 2.1. If Φ is an Orlicz function with $1 < a_\Phi \leq b_\Phi < \infty$, then there exists a constant $C > 0$ such that

$$\sup_{t>0} t\Phi[\tilde{\mu}_t(x)] \leq C \sup_{t>0} t\Phi[\mu_t(x)] \quad (2.6)$$

for all $x \in L_{\Phi,\infty}(\mathcal{M})$. Consequently,

$$\sup_{t>0} \frac{1}{\Phi^{-1}(\frac{1}{t})} \tilde{\mu}_t(x) \leq C' \sup_{t>0} \frac{1}{\Phi^{-1}(\frac{1}{t})} \mu_t(x), \quad \forall x \in L^{\Phi,\infty}(\mathcal{M}). \quad (2.7)$$

Proof. Let $1 < p_0 < a_\Phi \leq b_\Phi < p_1 < \infty$. By [7, Theorem III.3.8 and III.3.10], the map $x \mapsto \tilde{\mu}(x)$ is bounded from $L_{p_i}(\mathcal{M})$ to $L_{p_i}(\Omega)$, $i = 0, 1$. Using [3, Corollary 4.4], we obtain the desired result. \square

We use (2.1) and the above proposition to obtain the following result.

Corollary 2.1. Let Φ be an Orlicz function with $1 < a_\Phi \leq b_\Phi < \infty$. Set

$$\|x\|_{(\Phi)',\infty} = \sup_{t>0} \Psi^{-1}\left(\frac{1}{t}\right) \int_0^t \mu_s(x) ds, \quad \forall x \in L^{\Phi,\infty}(\mathcal{M}).$$

Then $\|x\|_{(\Phi)',\infty}$ is an equivalent norm on $L_{\Phi,\infty}(\mathcal{M})$.

Set $\varphi(t) = 1/\Psi^{-1}(\frac{1}{t})$. Then φ is an increasing concave function on $(0, \infty)$ with $\lim_{t \rightarrow 0} \varphi(t) = 0$ and $\lim_{t \rightarrow \infty} \varphi(t) = \infty$. Let $\Lambda_\varphi(\Omega)$, $M_\varphi(\Omega)$ be the usual Lorentz and Marcinkiewicz spaces with norms defined by

$$\Lambda_\varphi(\Omega) = \{f \in L_0(\Omega) : \|f\|_{\Lambda_\varphi} = \int_0^\infty f^*(t)\varphi'(t)dt < \infty\}$$

and

$$M_\varphi(\Omega) = \{f \in L_0(\Omega) : \|f\|_{M_\varphi} = \sup_{t>0} \frac{1}{\varphi(t)} \int_0^t f^*(s)ds < \infty\}.$$

The Lorentz space $\Lambda_\varphi(\Omega)$ has order continuous norm and $\Lambda_\varphi(\Omega)^* = M_\varphi(\Omega)$. If $M_\varphi^0(\Omega)$ denotes the linear subspace of $M_\varphi(\Omega)$ consisting of all $f \in M_\varphi(\Omega)$ for which

$$\limsup_{t \rightarrow 0} \frac{1}{\varphi(t)} \int_0^t \mu_s(x)ds = 0 \quad \text{and} \quad \limsup_{t \rightarrow \infty} \frac{1}{\varphi(t)} \int_0^t \mu_s(x)ds = 0.$$

Then $M_\varphi^0(\Omega)^* = \Lambda_\varphi(\Omega)$ (for more details see [18, Chapter II.5]). Since $M_\varphi^0(\Omega)$ is separable,

$$M_\varphi^0(\mathcal{M}) = \text{closure of } S(\mathcal{M}) \text{ in } M_\varphi(\mathcal{M}).$$

Hence,

$$\Lambda_\varphi(\mathcal{M})^* = M_\varphi(\mathcal{M}), \quad M_\varphi^0(\mathcal{M})^* = \Lambda_\varphi(\mathcal{M})$$

(see [12, Proposition 5.3], also see [8, Theorem 2.1] and [9, Proposition 2.1]).

Let $1 < a_\Phi \leq b_\Phi < \infty$. By Proposition 2.1,

$$M_\varphi(\mathcal{M}) = L_{\Phi, \infty}(\mathcal{M}).$$

We denote the closure of $S(\mathcal{M})$ in $L_{\Phi, \infty}(\mathcal{M})$ by $L_{\Phi, \infty}^0(\mathcal{M})$ and $\Lambda_\varphi(\mathcal{M})$ by $L_{1, \Psi}(\mathcal{M})$, respectively. Then

$$L_{1, \Psi}(\mathcal{M})^* = L_{\Phi, \infty}(\mathcal{M}) \quad \text{and} \quad L_{\Phi, \infty}^0(\mathcal{M})^* = L_{1, \Psi}(\mathcal{M}) \quad (2.8)$$

It is clear that

$$\lim_{t \rightarrow 0} \frac{t}{\varphi(t)} = \lim_{t \rightarrow 0} t\Psi^{-1}(\frac{1}{t}) = 0, \quad \lim_{t \rightarrow \infty} \frac{t}{\varphi(t)} = \lim_{t \rightarrow \infty} t\Psi^{-1}(\frac{1}{t}) = \infty.$$

We define continuous seminorms N_0 and N_∞ on $L_{\Phi, \infty}(\mathcal{M})$ by

$$N_0(x) = \limsup_{t \rightarrow 0} \frac{1}{\varphi(t)} \int_0^t \mu_s(x)ds$$

and

$$N_\infty(x) = \limsup_{t \rightarrow \infty} \frac{1}{\varphi(t)} \int_0^t \mu_s(x)ds,$$

for all $x \in L_{\Phi, \infty}(\mathcal{M})$. Using main result in [9], we obtain that

$$L_{\Phi, \infty}(\mathcal{M})^* = L_{1, \Psi}(\mathcal{M}) \oplus S_0 \oplus S_\infty, \quad (2.9)$$

where

$$\begin{aligned} S_0 &= \{\ell \in L_{\Phi, \infty}(\mathcal{M})^* : \ell \text{ annihilates all } x \in \mathcal{M}\} \\ &= \{\ell \in L_{\Phi, \infty}(\mathcal{M})^* : \exists C > 0, |\ell(x)| \leq CN_0(x), \quad \forall x \in L_{\Phi, \infty}(\mathcal{M})\}, \end{aligned}$$

$$\begin{aligned} S_\infty &= \{\ell \in L_{\Phi, \infty}(\mathcal{M})^* : \ell \text{ annihilates all } x \in L_{\Phi, \infty}(\mathcal{M}) \text{ with } r(x) \in S(\mathcal{M})\} \\ &= \{\ell \in L_{\Phi, \infty}(\mathcal{M})^* : \exists C > 0, |\ell(x)| \leq CN_\infty(x), \quad \forall x \in L_{\Phi, \infty}(\mathcal{M})\}. \end{aligned}$$

Let

$$M(t, \varphi) = \sup_{s>0} \frac{\varphi(ts)}{\varphi(s)}, \quad t > 0.$$

Define

$$p_\varphi = \lim_{t \searrow 0} \frac{\log M(t, \varphi)}{\log t}, \quad q_\varphi = \lim_{t \nearrow \infty} \frac{\log M(t, \varphi)}{\log t}.$$

Then

$$[p_\varphi, q_\varphi] \subset \left[\frac{1}{b_\Phi}, \frac{1}{a_\Phi}\right] \quad (2.10)$$

(For more details, see [22, Remarks 3 (p.84) and Theorem 11.11] and [21, Theorem 4.2]).

3. Noncommutative weak Orlicz spaces associated with a weight

We denote by $L_{loc}(\mathcal{M})$ by the set of all measurable locally-measurable operators affiliated with \mathcal{M} . It is well-known that $L_{loc}(\mathcal{M})$ is a $*$ -algebra with respect to the strong sum and strong product and $L_0(\mathcal{M})$ is a $*$ -subalgebra in $L_{loc}(\mathcal{M})$ (see [23, 24]). Set $M^+ = \{x \in \mathcal{M} : x \geq 0\}$ and $L_{loc}(\mathcal{M})^+ = \{x \in L_{loc}(\mathcal{M}) : x \geq 0\}$. Let

$$\tilde{\tau}(x) = \sup\{\tau(y) : y \in \mathcal{M}^+, y \leq x\}, \quad x \in L_{loc}(\mathcal{M})^+.$$

Then $\tilde{\tau}$ is an extension of τ to $L_{loc}(\mathcal{M})^+$ (see [24, §4.1]). The extension will be denoted still by τ .

Definition 3.1. (i) A weight on \mathcal{M} is a map $\omega : \mathcal{M}^+ \rightarrow [0, \infty]$ satisfying

$$\omega(x + \lambda y) = \omega(x) + \lambda \omega(y), \quad \forall x, y \in \mathcal{M}^+, \quad \forall \lambda \in \mathbb{R}$$

(where $0 \cdot \infty = 0$).

(ii) A weight ω is said to be normal if $\sup_i \omega(x_i) = \omega(\sup_i x_i)$ for any bounded increasing net (x_i) in \mathcal{M}^+ , faithful if $\omega(x) = 0$ implies $x = 0$, semifinite if the linear span \mathcal{M}_ω of the cone $\mathcal{M}_\omega^+ = \{x \in \mathcal{M}^+ : \omega(x) < \infty\}$ is dense in \mathcal{M} with respect to the ultra-weak topology, and locally finite if for any non-zero $x \in \mathcal{M}^+$ there is a non-zero $y \in \mathcal{M}^+$ such that $y \leq x$ and $0 < \omega(y) < \infty$.

Let ω be a faithful normal semifinite weight on \mathcal{M} . Then ω has a Radon-Nikodym derivative D_ω with respect to τ such that $\omega(\cdot) = \tau(D_\omega \cdot)$ (see [25]). The weight ω is locally finite if and only if the operator D_ω is locally measurable (see [27]). In the sequel, unless otherwise specified, we always denote by ω a faithful normal locally finite weight on \mathcal{M} . Let $\Phi^{-1} : [0, \infty) \rightarrow [0, \infty)$ be the inverse of Φ (which is uniquely defined on \mathbb{R}^+).

Let

$$\mathcal{M}_{\Phi, \infty}^{\alpha, \omega} = \left\{ x \in \mathcal{M} : \sup_{t>0} t\mu_t(\Phi(|\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha}|)) < \infty \right\}$$

and

$$\|x\|_{\Phi, \infty, \alpha, \omega} = \inf \left\{ c > 0 : \sup_{t>0} t\Phi(\mu_t(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha})/c) \leq 1 \right\}.$$

Lemma 3.1. $\mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$ is linear subspace in \mathcal{M} .

Proof. Let $x \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$ and $\eta \in \mathbb{C}$. If $|\eta| \leq 1$, by Lemma 2.5 in [14] and convexity of Φ ,

$$\begin{aligned} & \sup_{t>0} t\mu_t(\Phi(|\Phi^{-1}(D_\omega)^\alpha \eta x \Phi^{-1}(D_\omega)^{1-\alpha}|)) \\ &= \sup_{t>0} t\Phi(\mu_t(\Phi^{-1}(D_\omega)^\alpha \eta x \Phi^{-1}(D_\omega)^{1-\alpha})) \\ &= \sup_{t>0} t\Phi(|\eta|\mu_t(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha})) \\ &\leq |\eta| \sup_{t>0} t\Phi(\mu_t(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha})) \\ &= |\eta| \sup_{t>0} t\mu_t(\Phi(|\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha}|)) < \infty. \end{aligned}$$

Hence, $\eta x \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$. If $|\eta| > 1$, since $\Phi \in \Delta_2$, there exists a constant $k = k(|\eta|) > 0$ such that $\Phi(|\eta|t) \leq k\Phi(t)$ for all $t > 0$. Similar to the above, we obtain that $\eta x \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$.

Now let $x, y \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$. Using Lemma 2.5 in [14], convexity of Φ and $\Phi \in \Delta_2$, we get

$$\begin{aligned} & \sup_{t>0} t\mu_t(\Phi(|\Phi^{-1}(D_\omega)^\alpha x + y \Phi^{-1}(D_\omega)^{1-\alpha}|)) \\ &= \sup_{t>0} t\Phi(\mu_t(\Phi^{-1}(D_\omega)^\alpha x + y \Phi^{-1}(D_\omega)^{1-\alpha})) \\ &\leq \sup_{t>0} t\Phi(\mu_{t/2}(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha}) \\ &\quad + \mu_{t/2}(\Phi^{-1}(D_\omega)^\alpha y \Phi^{-1}(D_\omega)^{1-\alpha})) \\ &\leq c \sup_{t>0} t/2\Phi(\mu_{t/2}(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha})) \\ &\quad + c \sup_{t>0} t/2\Phi(\mu_{t/2}(\Phi^{-1}(D_\omega)^\alpha y \Phi^{-1}(D_\omega)^{1-\alpha})) < \infty, \end{aligned}$$

and so $x + y \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$. □

Similar to [3, Proposition 3.2], we have the following result.

Proposition 3.1. (i) If $\|x\|_{\Phi, \infty, \alpha, \omega} > 0$ then

$$\sup_{t>0} t\Phi(\mu_t(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha})/\|x\|_{\Phi, \infty, \alpha, \omega}) \leq 1.$$

(ii) $\|x\|_{\Phi, \infty, \alpha, \omega}$ is a quasi-norm on the linear space $\mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$ and

$$\|x + y\|_{\Phi, \infty, \alpha, \omega} \leq 2(\|x\|_{\Phi, \infty, \alpha, \omega} + \|y\|_{\Phi, \infty, \alpha, \omega}), \quad \forall x, y \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}. \quad (3.1)$$

(iii) If $\|x\|_{\Phi, \infty, \alpha, \omega} \leq 1$, then

$$\sup_{t>0} t\Phi(\mu_t(\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha})) \leq \|x\|_{\Phi, \infty, \alpha, \omega}.$$

(iv) $\|x\|_{\Phi, \infty, \alpha, \omega} \leq \|x\|_{\Phi, \alpha, \omega}$ for any $x \in \mathcal{M}_\Phi^{\alpha, \omega}$, where

$$\mathcal{M}_\Phi^{\alpha, \omega} = \{x \in \mathcal{M} : \Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha} \in L_\Phi(\mathcal{M})\}$$

and $\|x\|_{\Phi, \alpha, \omega} = \|\Phi^{-1}(D_\omega)^\alpha x \Phi^{-1}(D_\omega)^{1-\alpha}\|_\Phi$. Consequently, $\mathcal{M}_\Phi^{\alpha, \omega} \subset \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$.

Definition 3.2. Let ω be a faithful normal semifinite weight on \mathcal{M} and $\alpha \in [0, 1]$. We call the completion of $(\mathcal{M}_{\Phi, \infty}^{\alpha, \omega}, \|\cdot\|_{\Phi, \infty, \alpha, \omega})$ is the weak noncommutative Orlicz space associated with Φ, \mathcal{M} and ω , denote by $L_{\Phi, \infty}^{\alpha, \omega}(\mathcal{M}, \tau)$.

Lemma 3.2. Let D be a positive nonsingular operator in $L^1(\mathcal{M})$. If $\alpha \in [0, 1]$, then $\Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}$ is dense in $L_{\Phi, \infty}^0(\mathcal{M})$.

Proof. Set $e_n = e_{(\frac{1}{n}, n]}(D)$, for any $n \in \mathbb{N}$. Then e_n increases strongly to 1 and $\tau(e_n) < \infty$, for any $n \in \mathbb{N}$. Let $x \in S(\mathcal{M})$. Then there is a projection e in \mathcal{M} such that $\tau(e) < \infty$ and $ex = xe = x$. Hence, $x \in L^1(\mathcal{M})$. By [17, Lemma 2.1], we get $\lim_{n \rightarrow \infty} \|xe_n - x\|_1 = 0$. It follows that $xe_n - x \rightarrow 0$ in measure as $n \rightarrow \infty$. Using [14, Lemma 3.1], we get for any $t > 0$, $\mu_t(xe_n - x) \rightarrow 0$ as $n \rightarrow \infty$. On the other hand, by Lemma 2.5 in [14], $\mu_t(xe_n - x) \leq \mu_t(x)$ for all $t > 0$. Applying Lebesgue dominated convergence theorem, we get

$$\lim_{n \rightarrow \infty} \tau(\Phi(|xe_n - x|)) = \lim_{n \rightarrow \infty} \int_0^{\tau(1)} \Phi(\mu_t(xe_n - x)) dt = 0.$$

Therefore, $\lim_{n \rightarrow \infty} \|xe_n - x\|_\Phi = 0$. Similarly, $\lim_{n \rightarrow \infty} \|e_n x - x\|_\Phi = 0$. Using (iv) of Proposition 3.1, we obtain that $\lim_{n \rightarrow \infty} \|xe_n - x\|_{\Phi, \infty} = 0$ and $\lim_{n \rightarrow \infty} \|e_n x - x\|_{\Phi, \infty} = 0$, and so

$$\lim_{n \rightarrow \infty} \|e_n x e_n - x\|_{\Phi, \infty} \leq 2[\lim_{n \rightarrow \infty} \|xe_n - x\|_{\Phi, \infty} + \lim_{n \rightarrow \infty} \|e_n x - x\|_{\Phi, \infty}] = 0,$$

i.e., the closure of $\cup_{n=1}^\infty e_n \mathcal{M} e_n$ in $L_{\Phi, \infty}(\mathcal{M})$ contains $S(\mathcal{M})$. Thus, $\cup_{n=1}^\infty e_n \mathcal{M} e_n$ is dense in $L_{\Phi, \infty}^0(\mathcal{M})$.

Next, we prove that $\Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha} \subset L_{\Phi, \infty}^0(\mathcal{M})$. Set $\Phi^{(p)}(t) = \Phi(t^p)$, for $1 < p < \infty$. Let $y \in \mathcal{M}$. If $\alpha \in (0, 1)$, then

$$\lim_{n \rightarrow \infty} \tau(\Phi^{(\frac{1}{\alpha})}(|\Phi^{-1}(D)^\alpha - e_n \Phi^{-1}(D)^\alpha|)) = \lim_{n \rightarrow \infty} \tau(D - D e_n) = 0.$$

It follows that $\lim_{n \rightarrow \infty} \|\Phi^{-1}(D)^\alpha - e_n \Phi^{-1}(D)^\alpha\|_{\Phi^{(\frac{1}{\alpha})}, \infty} = 0$. Since $\Phi^{-1}(\frac{s}{2}) \geq \frac{1}{2} \Phi^{-1}(s)$ for all $s > 0$, by Lemma 2.5 in [14], we get

$$\begin{aligned} & \|\Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} - e_n \Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha}\|_{\Phi, \infty} \\ &= \sup_{t>0} \frac{1}{\Phi^{-1}(\frac{1}{2t})} \mu_{2t}(\Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} - e_n \Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha}) \\ &\leq 2\|y\| \sup_{t>0} \frac{1}{\Phi^{-1}(\frac{1}{t})} \mu_t((1 - e_n) \Phi^{-1}(D)^\alpha) \mu_t(\Phi^{-1}(D)^{1-\alpha}) \\ &\leq 2\|y\| \|(1 - e_n) \Phi^{-1}(D)^\alpha\|_{\Phi^{(\frac{1}{\alpha})}, \infty} \|\Phi^{-1}(D)^{1-\alpha}\|_{\Phi^{(\frac{1}{1-\alpha})}, \infty}. \end{aligned}$$

Hence, $\lim_{n \rightarrow \infty} \|\Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} - e_n \Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha}\|_{\Phi, \infty} = 0$. Similar to the above, $\lim_{n \rightarrow \infty} \|\Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} - e_n \Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} e_n\|_{\Phi, \infty} = 0$. On the other hand, $e_n \Phi^{-1}(D)^\alpha$, $\Phi^{-1}(D)^{1-\alpha} e_n \in \mathcal{M}$, and so for all $n \in \mathbb{N}$,

$$e_n \Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} e_n \in e_n \mathcal{M} e_n.$$

Therefore, $\Phi^{-1}(D)^\alpha y \Phi^{-1}(D)^{1-\alpha} \in L_{\Phi, \infty}^0(\mathcal{M})$. In the case $\alpha = 0, 1$, this result also holds. Thus $\Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha} \subset L_{\Phi, \infty}^0(\mathcal{M})$.

Finally, we prove that $\Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}$ is dense in $L_{\Phi, \infty}^0(\mathcal{M})$. For this it is sufficient to prove that $\cup_{n=1}^\infty e_n \mathcal{M} e_n \subset \Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}$. Since for any $n \in \mathbb{N}$, $e_n \Phi^{-1}(D)^{-\alpha}$, $\Phi^{-1}(D)^{\alpha-1} e_n \in \mathcal{M}$, we have that

$$\begin{aligned} e_n \mathcal{M} e_n &= \Phi^{-1}(D)^\alpha \Phi^{-1}(D)^{-\alpha} e_n \mathcal{M} e_n \Phi^{-1}(D)^{\alpha-1} \Phi^{-1}(D)^{1-\alpha} \\ &= \Phi^{-1}(D)^\alpha e_n \Phi^{-1}(D)^{-\alpha} \mathcal{M} \Phi^{-1}(D)^{\alpha-1} e_n \Phi^{-1}(D)^{1-\alpha} \\ &\subset \Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}. \end{aligned}$$

It follows that $\cup_{n=1}^\infty e_n \mathcal{M} e_n \subset \Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}$. \square

Theorem 3.1. *Let ω be a faithful normal semifinite weight on \mathcal{M} such that its the Radon-Nikodym derivative D_ω with respect to satisfy $D_\omega \in L^1(\mathcal{M})$. If $\alpha \in [0, 1]$, then $L_{\Phi, \infty}^{\alpha, \omega}(\mathcal{M}, \tau)$ and $L_{\Phi, \infty}^0(\mathcal{M})$ are isometrically isomorphic.*

Proof. We define $T : \mathcal{M}_{\Phi, \infty}^{\alpha, \omega} \rightarrow L_{\Phi, \infty}^{\alpha, \omega}(\mathcal{M}, \tau)$ by

$$T(x) = \Phi^{-1}(D)^\alpha x \Phi^{-1}(D)^{1-\alpha}, \quad x \in \mathcal{M}_{\Phi, \infty}^{\alpha, \omega}.$$

Then T is a linear isometry from $\mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$ to $\Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}$. By the definition of $L_{\Phi, \infty}^{\alpha, \omega}(\mathcal{M}, \tau)$ and Lemma 3.2, we know that $\mathcal{M}_{\Phi, \infty}^{\alpha, \omega}$ is dense in $L_{\Phi, \infty}^{\alpha, \omega}(\mathcal{M}, \tau)$ and $\Phi^{-1}(D)^\alpha \mathcal{M} \Phi^{-1}(D)^{1-\alpha}$ is dense in $L_{\Phi, \infty}^0(\mathcal{M})$. Hence, we can extend to an isometric isomorphism between $L_{\Phi, \infty}^{\alpha, \omega}(\mathcal{M}, \tau)$ and $L_{\Phi, \infty}^0(\mathcal{M})$. \square

4. Noncommutative weak Orlicz-Hardy spaces

We will assume that \mathcal{D} is a von Neumann subalgebra of \mathcal{M} such that the restriction of τ to \mathcal{D} is still semifinite. Let \mathcal{E} be the (unique) normal faithful conditional expectation of \mathcal{M} with respect to \mathcal{D} which leaves τ invariant.

Definition 4.1. *A w^* -closed subalgebra \mathcal{A} of \mathcal{M} is called a subdiagonal subalgebra of \mathcal{M} with respect to \mathcal{E} (or \mathcal{D}) if*

- (i) $\mathcal{A} + J(\mathcal{A})$ is w^* -dense in \mathcal{M} , where $J(\mathcal{A}) = \{x^* : x \in \mathcal{A}\}$,
- (ii) $\mathcal{E}(xy) = \mathcal{E}(x)\mathcal{E}(y)$, $\forall x, y \in \mathcal{A}$,
- (iii) $\mathcal{A} \cap J(\mathcal{A}) = \mathcal{D}$.

\mathcal{D} is then called the diagonal of \mathcal{A} .

In this section \mathcal{M} always denotes a semifinite von Neumann algebra with a normal semi-finite faithful trace τ satisfying $\tau(1) = \gamma$ and \mathcal{A} denotes a subdiagonal subalgebra of \mathcal{M} with respect to \mathcal{E} (or \mathcal{D}). We keep all notations introduced in the previous section.

Let

$$H_{\Phi, \infty}(\mathcal{A}) = \{x \in L_{\Phi, \infty}(\mathcal{M}) : \tau(xa) = 0, \quad \forall a \in \mathcal{A}_0\}.$$

Then $H_{\Phi, \infty}(\mathcal{A})$ is called noncommutative weak Orlicz-Hardy space associated with \mathcal{A} . Similarly, we define $H_{1, \Psi}(\mathcal{A})$ by

$$H_{1, \Psi}(\mathcal{A}) = \{x \in L_{1, \Psi}(\mathcal{M}) : \tau(xa) = 0, \quad \forall a \in \mathcal{A}_0\}.$$

Let \mathcal{M} be finite ($\tau(1) = \alpha < \infty$). By Propositions 4.3 in [4], we know that $H_{1, \Psi}(\mathcal{A})$ is the closure of \mathcal{A} in $L_{1, \Psi}(\mathcal{M})$.

Proposition 4.1. *Let $1 < a_{\Phi} \leq b_{\Phi} < \infty$. Then*

$$L_{1, \Psi}(\mathcal{M}) = H_{1, \Psi}(\mathcal{A}) \oplus J(H_{1, \Psi}(\mathcal{A})_0),$$

where $H_{1, \Psi}(\mathcal{A})_0 = \{x \in H_{1, \Psi}(\mathcal{A}) : \mathcal{E}(x) = 0\}$.

Proof. Since the lower Boyd index $p_{L_{1, \Psi}}$ and upper Boyd index $q_{L_{1, \Psi}}$ of $L_{1, \Psi}(\Omega)$ are $\frac{1}{q_{\varphi}}$ and $\frac{1}{p_{\varphi}}$, respectively (see [21, Theorem 4.2]). Using (2.10), we get that $1 < p_{L_{1, \Psi}} \leq q_{L_{1, \Psi}} < \infty$. If $\tau(1) < \infty$, then by [5, Theorem 5], we obtain the desired result. If $\tau(1) = \infty$. Choose that $1 < p < p_{L_{1, \Psi}} \leq p_{L_{1, \Psi}} < q < \infty$. Since there is a bounded projection operator P from $L_p(\mathcal{M})$ onto $H_p(\mathcal{A})$ and from $L_q(\mathcal{M})$ onto $H_q(\mathcal{A})$ (see [6, Theorem 4.2]), by Theorem 3.4 in [13], we know that P is a bounded projection from $L_{1, \Psi}(\mathcal{M})$ onto $H_{1, \Psi}(\mathcal{A})$. \square

Theorem 4.1. *Let $1 < a_{\Phi} \leq b_{\Phi} < \infty$. Then*

$$H_{\Phi, \infty}(\mathcal{A})^* = H_{1, \Psi}(\mathcal{A}) \oplus S_0|_{H_{\Phi, \infty}(\mathcal{A})} \oplus S_{\infty}|_{H_{\Phi, \infty}(\mathcal{A})}. \quad (4.1)$$

Proof. It is clear that

$$H_{1, \Psi}(\mathcal{A}) \oplus S_0|_{H_{\Phi, \infty}(\mathcal{A})} \oplus S_{\infty}|_{H_{\Phi, \infty}(\mathcal{A})} \subset H_{\Phi, \infty}(\mathcal{A})^*.$$

Let $\ell \in H_{\Phi, \infty}(\mathcal{A})^*$. By Hahn-Banach theorem, there is a functional $\tilde{\ell} \in L_{\Phi, \infty}(\mathcal{M})^*$ such that $\ell = \tilde{\ell}|_{H_{\Phi, \infty}(\mathcal{A})}$. Using (2.9), we get that

$$\tilde{\ell}(x) = \tau(xy^*) + \tilde{\ell}_1(x) + \tilde{\ell}_2(x), \quad \forall x \in L_{\Phi, \infty}(\mathcal{M}),$$

where $y \in L_{1, \Psi}(\mathcal{M})$, $\tilde{\ell}_1 \in S_0$ and $\tilde{\ell}_2 \in S_{\infty}$. Using Proposition 4.1, we obtain that there exist $h \in H_{1, \Psi}(\mathcal{A})$ and $z \in H_{1, \Psi}(\mathcal{A})_0$ such that $y = h + z^*$. Hence,

$$\tau(ay^*) = \tau(ah^*) + \tau(az) = \tau(ah^*), \quad a \in H_{\Phi, \infty}(\mathcal{A}).$$

Therefore, $\ell = h + \tilde{\ell}_1|_{H_{\Phi, \infty}(\mathcal{A})} + \tilde{\ell}_2|_{H_{\Phi, \infty}(\mathcal{A})}$. From this follows (4.2). \square

If $\Phi(t) = \frac{t^p}{p}$ ($1 < p < \infty$), then $\Psi(t) = \frac{t^q}{q}$ and $\varphi(t) = \frac{1}{q^{1/q}}t^{1/q}$, where $\frac{1}{p} + \frac{1}{q} = 1$. Hence,

$$N_0(x) = \limsup_{t \rightarrow 0} q^{1/q} \frac{1}{t^{1/q}} \int_0^t \mu_s(x) ds = q^{1/q} \limsup_{t \rightarrow 0} t^{1/p-1} \int_0^t \mu_s(x) ds,$$

and

$$N_\infty(x) = \limsup_{t \rightarrow \infty} q^{1/q} \frac{1}{t^{1/q}} \int_0^t \mu_s(x) ds = q^{1/q} \limsup_{t \rightarrow \infty} t^{1/p-1} \int_0^t \mu_s(x) ds.$$

Corollary 4.1. *Let $1 < p < \infty$. Then*

$$H_{p,\infty}(\mathcal{A})^* = H_{1,q}(\mathcal{A}) \oplus S_0|_{H_{p,\infty}(\mathcal{A})} \oplus S_\infty|_{H_{p,\infty}(\mathcal{A})}. \quad (4.2)$$

Acknowledgement

T. B. Bekjan and M. Raikhan are partially supported by project AP09259802 of the Science Committee of Ministry of Education and Science of the Republic of Kazakhstan.

References

- [1] M. H. A. Al-Rashed and B. Zegarliński Noncommutative Orlicz spaces associated to a state, *Studia Math.* **180** (2007), no. 3, 199-207.
- [2] Sh. A. Ayupov, V. I. Chilin and R. Z. Abdullaev, Orlicz spaces associated with a semi-finite von Neumann algebra, *Comment. Math. Univ. Carolin.* **53** (2012), no. 4, 519-533.
- [3] T. N. Bekjan, Z. Chen, P. Liu and Y. Jiao, Noncommutative weak Orlicz spaces and martingale inequalities, *Studia Math.* **204** (2011), 195-212.
- [4] T. N. Bekjan, Noncommutative symmetric Hardy spaces, *Inter. Equat. Oper. Th.* **81** (2015), 191-212.
- [5] T. N. Bekjan and M. Mustafa, On interpolation of noncommutative symmetric Hardy spaces, *Positivity* **21** (2017), 1307-1317.
- [6] T. N. Bekjan, Noncommutative Hardy space associated with semi-finite subdiagonal algebras, *J. Math. Anal. Appl.* **429** (2015), 1347-1369.
- [7] C. Bennett and R. Sharpley, *Interpolation of operators*, Academic Press Inc., Boston, MA, 1988.
- [8] L. J. Ciach, On the conjugates of some operator spaces I, *Demonstratio Math.* **18** (1985), 537-553.
- [9] L. J. Ciach, On the conjugates of some operator spaces, II, *Demonstratio Math.* **21** (1988), 357-367.
- [10] M. Cwikel, The dual of weak L^p , *Ann. Inst. Fourier* **25** (1975) 81-126.

- [11] M. Cwikel and Y. Sagher, $L(p, \infty)^*$, *Indiana Math. J.* **21** (1972) 781-786.
- [12] P. G. Dodds, T. K. Dodds and B. de Pagter, Noncommutative Köthe duality, *Trans. Amer. Math. Soc.* **339** (1993), 717-750.
- [13] P. G. Dodds, T. K. Dodds, and B. de Pagter, Fully symmetric operator spaces, *Integ. Equ. Oper. Theory* **15** (1992), 942-972.
- [14] T. Fack and H. Kosaki, Generalized s -numbers of τ -measurable operators, *Pac. J. Math.* **123** (1986), 269-300.
- [15] L. Grafakos, *Classical and Modern Fourier Analysis*, Pearson Education, London, 2004
- [16] Y. Han and J. Shao, The dual of $L_{p,\infty}(\mathcal{M})$, *J. Math. Anal. Appl.* **398** (2013) 814-821.
- [17] M. Junge, Doob's inequality for noncommutative martingales, *J. Reine. Angew. Math.* **549** (2002) 149-190.
- [18] S. G. Krein, J. I. Petunin and E. M. Semenov, *Interpolation of linear operators*, Translations of Mathematical Monographs, vol.54, Amer. Math. Soc., 1982.
- [19] J. Lindenstrauss and L. Tzafriri, *Classical Banach space II*, Springer-Verlag, Berlin, 1979.
- [20] P. Liu, Y. Hou and M. Wang, Weak Orlicz space and its applications to martingale theory, *Sci. China Math.* **53** (4) (2010), 905-916.
- [21] L. Maligranda, Indices and interpolation, *Dissert. Math.* **234**, Polska Akademia Nauk, Inst. Mat., 1985.
- [22] L. Maligranda, Orlicz spaces and interpolation, Seminars in Mathematics, Departamento de Matemática, Universidade Estadual de Campinas, Brasil, 1989.
- [23] M. A. Muratov and V. I. Chilin, $*$ -algebras of unbounded operators affiliated with a von Neumann algebra, *J. Math. Sci.* **140**(2007), 445-451.
- [24] M. A. Muratov and V. I. Chilin, *Algebras of measurable operators and locally measurable operators*, Kyev. Institute of Math., Ukrainian Academy of Sciences, 2007.
- [25] G. K. Pedersen and M. Takesaki, The Radon-Nikodym theorem for von Neuman algebras, *Acta Math.* **130** (1973) 53-87.
- [26] M. Rao and Z. Ren, *Application of Orlicz Spaces*, New York: Marcel Dekker, 2002.
- [27] N. V. Trunov, To the theory normal weights on von Neumann algebras, *Izv. Vuzov. Math.*, no. 8, 1982, 61-70.
- [28] Q. Xu, Analytic functions with values in lattices and symmetric spaces of measurable operators, *Math. Proc. Cambridge Phil. Soc.* **109** (1991), 541-563.