Rota-Baxter C^* -algebras

Zhonghua Li^{*} and Shukun Wang[†]

School of Mathematical Sciences, Tongji University No. 1239 Siping Road, Shanghai 200092, China

Abstract

This paper introduces the notion of Rota-Baxter C^* -algebras. Here a Rota-Baxter C^* -algebra is a C^* -algebra with a Rota-Baxter operator. Symmetric Rota-Baxter operators, as special cases of Rota-Baxter operators on C^* -algebra, are defined and studied. A theorem of Rota-Baxter operators on concrete C^* -algebras is given, deriving the relationship between two kinds of Rota-Baxter algebras. As a corollary, some connection between *-representations and Rota-Baxter operators is given. The notion of representations of Rota-Baxter C^* -algebras are constructed, and a theorem of representations of direct sums of Rota-Baxter representations is derived. Finally using Rota-Baxter operators, the notion of quasidiagonal operators on C^* -algebra is reconstructed.

Keywords Rota-Baxter operators, C^{*}-algebras, Rota-Baxter C^{*}-algebras. 2010 Mathematics Subject Classification 17B38, 46L05, 16G99.

1 Introduction

Let \mathcal{A} be an associative algebra over a given field F. A linear operator P on \mathcal{A} is called a Rota-Baxter operator of weight $\lambda \in F$ if it satisfies

$$P(a)P(b) = P(aP(b)) + P(P(a)b) + \lambda P(ab), \qquad \forall a, b \in \mathcal{A}.$$
(1.1)

An associative algebra with a Rota-Baxter operator is called a Rota-Baxter algebra, which can be regarded as an analogue of a differential algebra. In fact, when taking $\lambda = 0$, Eq. (1.1) is an algebraic abstraction of the formula of integration by parts.

The study of Rota-Baxter algebras originated from probability theory [2] and has found applications in many areas of mathematics and physics. In the late 1990s, Rota-Baxter algebras were found their applications in the work of Connes and Kremier [4] regarding the renormalization of perturbative quantum field theory. Since 2000, the connection between the classical Yang-Baxter equation in mathematical physics and Rota-Baxter operators has been found in [1].

^{*} E-mail address: zhonghua_li@tongji.edu.cn

[†]E-mail address: 2010165@tongji.edu.cn

The representations of Rota-Baxter algebras were studied in [12], where some basic concepts and properties were established. However, it is still in the early stages of development. In [12], Lin and Qiao studied the representations and regular-singular decomposition of Laurent series Rota-Baxter algebras. In [9], regular-singular decomposition of Rota-Baxter modules were obtained under the condition of quasi-idempotency. And in [15], representations of the polynomial Rota-Baxter algebras were studied.

The theory of operator algebras in Hilbert spaces was initiated by von Neumann [20] in 1929. In [13], Murray and von Neumann laid the foundation of the theory of W^* -algebras. The notion of C^* -algebras was introduced by Gelfand and Naimark [6] in 1943. Basic theory of representations of C^* -algebras was established in [7, 10, 14, 19]. Derivations on C^* -algebras and W^* -algebras were studied deeply in the 1960s and 1970s in [10, 11, 16, 17]. Especially, in [17], it was shown that all derivations on W^* -algebras are inner derivations. While Rota-Baxter operators on C^* -algebras have not been studied.

In this note, we establish basic concepts of Rota-Baxter C^* -algebras. Some special Rota-Baxter operators in C^* -algebras, which are called symmetric Rota-Baxter operators and Rota-Baxter operators matching projections are studied. And representations of Rota-Baxter C^* -algebras are also established, which are special cases of traditional Rota-Baxter representations.

A C^* -algebra \mathcal{A} is a special algebra in the complex field. Combined with a Rota-Baxter operator on \mathcal{A} , the basic concepts of Rota-Baxter C^* -algebras were established in Subsection 2.2. A Rota-Baxter operator P on the C^* -algebra \mathcal{A} is symmetric if $P(a^*) = P(a)^*$ for any $a \in \mathcal{A}$. In Subsection 2.3, We show that a C^* -algebra \mathcal{A} can be decomposed into a direct sum of two C^* -subalgebras if and only if there is a bounded idempotent symmetric Rota-Baxter operator of weight -1 on \mathcal{A} .

In Section 3, we study Rota-Baxter operators on the C^* -algebra $B(\mathcal{H})$, where \mathcal{H} is a Hilbert space. In Subsection 3.2, the notion of Rota Baxter operators that match projections are introduced. We find that the Rota-Baxter operators matching projections on Hilbert spaces have many good properties. In fact, we can construct Rota-Baxter operators of this kind from Rota-Baxter operators on C^* -subalgebras of $B(\mathcal{H})$. As a corollary, we study the relationship between the Rota-Baxter operators matching projections and *-representations of C^* -algebras on a Hilbert space. In Subsection 3.3, we construct representations of Rota-Baxter C^* -algebras.

At last in Section 4, we reconstruct the notion of quasidiagonal operators of C^* algebras with the help of Rota-Baxter operators.

2 Rota-Baxter C*-algebras

2.1 C^* -algebras

We first recall some basic concepts of C^* -algebras from [18]. Let F be the complex field \mathbb{C} or the real number field \mathbb{R} .

Let \mathcal{A} be an associative algebra over F. The algebra \mathcal{A} is called a normed algebra if associated to each element a in \mathcal{A} there is a real number ||a||, called the norm of a, with the properties:

(i) $||a|| \ge 0$ for any $a \in \mathcal{A}$, and ||a|| = 0 if and only if a = 0;

(ii) $||a + b|| \le ||a|| + ||b||$ for any $a, b \in \mathcal{A}$;

(iii) $\|\lambda a\| = |\lambda| \|a\|$ for any $a \in \mathcal{A}$ and $\lambda \in F$;

(iv) $||ab|| \leq ||a|| ||b||$ for any $a, b \in \mathcal{A}$.

The topology defined by the norm $\|\cdot\|$ on \mathcal{A} is called the uniform topology. If \mathcal{A} is complete with respect to the norm, then \mathcal{A} is called a Banach algebra. A map $\mathcal{A} \to \mathcal{A}; a \mapsto a^*$ is called an involution if it satisfies the following conditions:

(i) $(a^*)^* = a$ for any $a \in \mathcal{A}$;

- (ii) $(a+b)^* = a^* + b^*$ for any $a, b \in \mathcal{A}$;
- (iii) $(\lambda a)^* = \overline{\lambda} a^*$ for any $a \in \mathcal{A}$ and $\lambda \in F$;

(iv) $(ab)^* = b^*a^*$ for any $a, b \in \mathcal{A}$.

An algebra with an involution * is called a *-algebra. Finally, a Banach *-algebra \mathcal{A} is called a (*F*-linear) C^* -algebra if it satisfies

$$||a^*a|| = ||a||^2$$

for any $a \in \mathcal{A}$. Note that a \mathbb{C} -linear C^* -algebra is natural a \mathbb{R} -linear C^* -algebra.

A subset S of a C^* -algebra \mathcal{A} is called self-adjoint if $a^* \in S$ for any $a \in S$. In particular, an element $a \in \mathcal{A}$ is called self-adjoint if $a^* = a$. A self-adjoint, uniformly closed subalgebra of \mathcal{A} is called a C^* -subalgebra of \mathcal{A} , which is also a C^* -algebra.

As there is a new structure * on a C^* -algebra, it is natural to consider the *-homomorphisms between two C^* -algebras. Let \mathcal{A} and \mathcal{B} be two C^* -algebras. A map $\phi : \mathcal{A} \to \mathcal{B}$ is called a *-homomorphism if it satisfies

(i)
$$\phi(a+b) = \phi(a) + \phi(b)$$
 for any $a, b \in \mathcal{A}$

- (ii) $\phi(\lambda a) = \lambda \phi(a)$ for any $a \in \mathcal{A}$ and $\lambda \in F$;
- (iii) $\phi(ab) = \phi(a)\phi(b)$ for any $a, b \in \mathcal{A}$;
- (iv) $\phi(a^*) = \phi(a)^*$ for any $a \in \mathcal{A}$.

We recall the definition of the direct sum of C^* -algebras. Let $\{\mathcal{A}_k\}_{k\in\Lambda}$ be a family of C^* -algebras. We define the direct sum

$$\bigoplus_{k\in\Lambda}\mathcal{A}_k = \left\{ (a_k)_{k\in\Lambda} \mid a_k \in \mathcal{A}_k \text{ for any } k \in \Lambda, \sup_{k\in\Lambda} ||a_k|| < \infty \right\}.$$

Then $\bigoplus_{k \in \Lambda} \mathcal{A}_k$ is a C^* -algebra under the following operators:

(i) $(a_k) + (b_k) = (a_k + b_k);$ (ii) $\lambda(a_k) = (\lambda a_k), \quad (\lambda \in F);$ (iii) $(a_k)(b_k) = (a_k b_k);$ (iv) $||(a_k)|| = \sup_k ||a_k||;$ (v) $(a_k)^* = (a_k^*).$

For any $k \in \Lambda$, it is natural to regard \mathcal{A}_k as a C^* -subalgebra of $\bigoplus \mathcal{A}_k$.

2.2 Rota-Baxter C*-algebras

Now we introduce the concept of Rota-Baxter C^* -algebras.

Definition 2.1. Let \mathcal{A} be a C^* -algebra. A linear operator $P : \mathcal{A} \to \mathcal{A}$ is called a Rota-Baxter operator of weight $\lambda \in F$ on \mathcal{A} if it satisfies:

$$P(a)P(b) = P(aP(b)) + P(P(a)b) + \lambda P(ab), \qquad \forall a, \ b \in \mathcal{A}.$$

If P is a Rota-Baxter operator of weight λ , then it is easy to verify that

$$P = -\lambda \operatorname{id}_{\mathcal{A}} - P$$

is also a Rota-Baxter operator of weight λ .

Definition 2.2. A Rota-Baxter C^* -algebra of weight λ is a pair (\mathcal{A}, P) consisting of a C^* -algebra \mathcal{A} , and a Rota-Baxter operator $P : \mathcal{A} \to \mathcal{A}$ of weight λ on \mathcal{A} .

We give an example of Rota-Baxter C^* -algebras.

Example 2.3. Let C([0,1]) be the set of all complex valued, continuous functions on the closed interval [0,1]. Then C([0,1]) becomes a commutative algebra over \mathbb{C} under pointwise addition and multiplication. It was showed in [18] that C([0,1]) is a commutative C^* -algebra, where for any $f \in C([0,1])$, the norm of f is defined by

$$||f|| = \sup_{x \in [0,1]} |f(x)|,$$

and the involution f^* of f is given by

$$f^*(x) = \overline{f(x)}, \quad \forall x \in [0, 1].$$

The linear operator T on C([0,1]) defined by

$$T(f)(x) = \int_0^x f(s)ds, \quad \forall f \in C([0,1]), \; \forall x \in [0,1]$$

is called the Volterra operator. It is easy to verify that T is a Rota-Baxter operator of weight 0 on C([0,1]). Then (C([0,1]),T) is a Rota-Baxter C^{*}-algebra.

As in [18], basic concepts on C^* -algebras can be similarly defined for Rota-Baxter C^* -algebras. Particularly, a Rota-Baxter C^* -subalgebra of a Rota-Baxter C^* -algebra \mathcal{A} is a C^* -subalgebra I of \mathcal{A} such that $P(I) \subseteq I$. A Rota-Baxter C^* -algebra homomorphism $\phi : (\mathcal{A}_1, P_1) \to (\mathcal{A}_2, P_2)$ between two Rota-Baxter C^* -algebras of the same weight λ is a *-homomorphism such that $\phi \circ P_1 = P_2 \circ \phi$.

In below, we always take $F = \mathbb{C}$. Hence all the Hilbert spaces and algebras are assumed to be over the complex field \mathbb{C} .

We recall the definition of derivations of C^* -algebras from [18]. Let \mathcal{A} be a C^* algebra. A linear map $\delta : \mathcal{A} \to \mathcal{A}$ is called a derivation if

$$\delta(ab) = \delta(a)b + a\delta(b), \qquad \forall a, \ b \in \mathcal{A}.$$

It is easy to see that if P is an invertible Rota-Baxter operator of weight 0 on A, then the inverse operator of P is a derivation of A. As an application, we have the following proposition. **Proposition 2.4.** If \mathcal{A} is a commutative C^* -algebra, then there is no invertible Rota-Baxter operator of weight 0 on \mathcal{A} .

Proof. From [18], we know that if δ is a derivation on \mathcal{A} , then $\delta = 0$. While if there is an invertible Rota-Baxter operator P of weight 0 on \mathcal{A} , then the inverse of P is an invertible derivation on \mathcal{A} , which is a contradiction.

2.3 Symmetric Rota-Baxter operators

In this subsection, we introduce the notion of symmetric Rota-Baxter operators.

Definition 2.5. Let \mathcal{A} be a C^* -algebra and let \mathcal{A}_1 be a C^* -subalgebra of \mathcal{A} . A Rota-Baxter operator P on \mathcal{A} is called symmetric on \mathcal{A}_1 if

$$P(a^*) = P(a)^*$$

for any $a \in A_1$. If P is symmetric on A, we just call that P is symmetric.

Example 2.6. The Volterra operator T on C([0,1]) in Example 2.3 is symmetric.

Let \mathcal{A} be a commutative C^* -algebra. Let \mathcal{A}^s be the set of all self-adjoint elements in \mathcal{A} . It is easy to verify that \mathcal{A}^s is a \mathbb{R} -linear subalgebra of \mathcal{A} . We have the following result.

Proposition 2.7. Let \mathcal{A} be a commutative C^* -algebra and $\lambda \in \mathbb{R}$. Then there is a 1-1 correspondence between the following two sets:

(1) the set of symmetric Rota-Baxter operators of weight λ on \mathcal{A} ;

(2) the set of \mathbb{R} -linear Rota-Baxter operators of weight λ on \mathcal{A}^s .

Proof. Denote by Γ the set of symmetric Rota-Baxter operators of weight λ on \mathcal{A} , and by Γ_1 the set of \mathbb{R} -linear Rota-Baxter operators of weight λ on \mathcal{A}^s .

Let $P \in \Gamma$. Then for any $a \in \mathcal{A}^s$, we have

$$P(a) = P(a^*) = P(a)^*.$$

Hence we get a restriction $P|_{\mathcal{A}^s} : \mathcal{A}^s \to \mathcal{A}^s$, which is an element of Γ_1 . Therefore we have a map $\Phi : \Gamma \to \Gamma_1$, such that $\Phi(P) = P|_{\mathcal{A}^s}$.

Conversely, let $P_1 \in \Gamma_1$. For any $a \in \mathcal{A}$, we have $a = a_1 + ia_2$, where $i = \sqrt{-1}$ and

$$a_1 = \frac{1}{2}(a + a^*) \in \mathcal{A}^s, \qquad a_2 = \frac{1}{2i}(a - a^*) \in \mathcal{A}^s.$$

Then we get a map $P : \mathcal{A} \to \mathcal{A}$ by setting $P(a) = P_1(a_1) + iP_1(a_2)$. It is obvious that P is linear.

We show that P satisfies the Rota-Baxter relation. Let $b = b_1 + ib_2$ be another element of \mathcal{A} with $b_1, b_2 \in \mathcal{A}^s$ defined similarly as above. We have

$$P(a)P(b) = (P_1(a_1) + iP_1(a_2))(P_1(b_1) + iP_1(b_2))$$

= $P_1(a_1)P_1(b_1) - P_1(a_2)P_1(b_2) + i(P_1(a_2)P_1(b_1) + P_1(a_1)P_1(b_2)).$

Since P_1 is a Rota-Baxter operator on \mathcal{A}^s , we find

$$\begin{split} P(a)P(b) = & P_1(a_1P_1(b_1) + P_1(a_1)b_1 + \lambda a_1b_1) - P_1(a_2P_1(b_2) + P_1(a_2)b_2 + \lambda a_2b_2) \\ &+ iP_1(a_2P_1(b_1) + P_1(a_2)b_1 + \lambda a_2b_1) + iP_1(a_1P_1(b_2) + P_1(a_1)b_2 + \lambda a_1b_2) \\ = & P_1(a_1P_1(b_1)) - P_1(a_2P_1(b_2)) + iP_1(a_2P_1(b_1)) + iP_1(a_1P_1(b_2)) \\ &+ P_1(P_1(a_1)b_1) - P_1(P_1(a_2)b_2) + iP_1(P_1(a_2)b_1) + iP_1(P_1(a_1)b_2) \\ &+ \lambda P_1(a_1b_1 - a_2b_2) + i\lambda P_1(a_2b_1 + a_1b_2). \end{split}$$

According to the following decompositions

$$ab = (a_1b_1 - a_2b_2) + i(a_2b_1 + a_1b_2),$$

$$aP(b) = (a_1P_1(b_1) - a_2P_1(b_2)) + i(a_2P_1(b_1) + a_1P_1(b_2)),$$

$$P(a)b = (P_1(a_1)b_1 - P_1(a_2)b_2) + i(P_1(a_2)b_1 + P_1(a_1)b_2),$$

we finally get

$$P(a)P(b) = P(aP(b)) + P(P(a)b) + \lambda P(ab),$$

which means that P is a Rota-Baxter operator of weight λ on \mathcal{A} .

Now we show that P is symmetric. Since we have the decomposition $a^* = a_1 - ia_2$, we get

$$P(a^*) = P_1(a_1) - iP_1(a_2).$$

On the other hand, we have

$$P(a)^* = (P_1(a_1) + iP_1(a_2))^* = P_1(a_1)^* - iP_1(a_2)^* = P_1(a_1) - iP_1(a_2).$$

Hence $P(a^*) = P(a)^*$ and then P is symmetric.

Therefore we get a map $\Psi : \Gamma_1 \to \Gamma$, such that $\Psi(P_1) = P$.

Finally, it is easy to verify that $\Phi \circ \Psi$ and $\Psi \circ \Phi$ are identity maps. And the proposition is proved.

The following result gives an equivalent condition for the idempotent and symmetric Rota-Baxter operators of weight -1 on C^* -algebras.

Theorem 2.8. Let \mathcal{A} be a C^* -algebra and let P be a bounded linear operator on \mathcal{A} . Then the following two statements are equivalent:

- (1) P is an idempotent and symmetric Rota-Baxter operator of weight -1 on \mathcal{A} ;
- (2) There is a direct sum decomposition $\mathcal{A} = \mathcal{A}_1 \oplus \mathcal{A}_2$, such that \mathcal{A}_1 and \mathcal{A}_2 are C^* -subalgebras of \mathcal{A} and P is the projection of \mathcal{A} onto \mathcal{A}_1 .

Proof. Assume that P is an idempotent and symmetric Rota-Baxter operator of weight -1 on \mathcal{A} . Let $\mathcal{A}_1 = P(\mathcal{A})$ and $\mathcal{A}_2 = \tilde{P}(\mathcal{A})$, where $\tilde{P} = \mathrm{id}_{\mathcal{A}} - P$. Then from [8, Theorem 1.1.13], we know that \mathcal{A}_1 and \mathcal{A}_2 are subalgebras of \mathcal{A} , $\mathcal{A} = \mathcal{A}_1 \oplus \mathcal{A}_2$ is a direct sum decomposition, and P is the projection of \mathcal{A} onto \mathcal{A}_1 . For any $a = P(b) \in \mathcal{A}_1$ with $b \in \mathcal{A}$, we have

$$a^* = P(b)^* = P(b^*) \in \mathcal{A}_1.$$

Hence \mathcal{A}_1 is closed under the involution * and then self-adjoint. Now let $\{P(b_n)\}$ be a sequence in \mathcal{A}_1 that uniformly converges to $a \in \mathcal{A}$. Then since P is bounded and idempotent, we get $\{P(b_n)\}$ uniformly converges to $P(a) \in \mathcal{A}_1$. Therefore \mathcal{A}_1 is a C^* -subalgebra of \mathcal{A} . It is similar to show that \mathcal{A}_2 is a C^* -subalgebra of \mathcal{A} .

Conversely, assume that we have a direct sum decomposition $\mathcal{A} = \mathcal{A}_1 \oplus \mathcal{A}_2$, such that \mathcal{A}_1 and \mathcal{A}_2 are C^* -subalgebras of \mathcal{A} , and P is the projection of \mathcal{A} onto \mathcal{A}_1 . From [8, Theorem 1.1.13], we have P is an idempotent Rota-Baxter operator of weight -1 on \mathcal{A} . For any $a \in \mathcal{A}$, assume that $a = a_1 + a_2$ with $a_1 \in \mathcal{A}_1$ and $a_2 \in \mathcal{A}_2$, then we have $a^* = a_1^* + a_2^*$. Since \mathcal{A}_i is a C^* -subalgebra of \mathcal{A} , we get $a_i^* \in \mathcal{A}_i$. Hence

$$P(a^*) = a_1^* = P(a)^*,$$

which means that P is symmetric. Finally, from [18, Corollary 1.2.6] and the definition of the direct sum of C^* -algebras in Subsection 2.1, we know that

$$||a|| = \max(||a_1||, ||a_2||) \ge ||a_1|| = ||P(a)||.$$

Hence we have

$$||P|| = \sup_{a \in \mathcal{A}, a \neq 0} \frac{||P(a)||}{||a||} \le 1,$$

which means that P is bounded.

3 Rota-Baxter operators matching projections on Hilbert spaces

3.1 The C^* -algebra $B(\mathcal{H})$

We recall the notion of Hilbert spaces. A complex linear space \mathcal{H} is an inner product space if associated to any $x, y \in \mathcal{H}$ there is a product $\langle x, y \rangle$ which satisfies the following properties:

(i)
$$\langle x, y \rangle = \overline{\langle y, x \rangle}$$
 for any $x, y \in \mathcal{H}$;

(ii)
$$\langle \lambda_1 x_1 + \lambda_2 x_2, y \rangle = \lambda_1 \langle x_1, y \rangle + \lambda_2 \langle x_2, y \rangle$$
 for any $\lambda_1, \lambda_2 \in \mathbb{C}$ and $x_1, x_2, y \in \mathcal{H}$;

(iii) $\langle x, x \rangle = 0$ for x = 0 and $\langle x, x \rangle > 0$ for any nonzero x in \mathcal{H} .

Then the norm of \mathcal{H} induced from the inner product is defined by $||x|| = \sqrt{\langle x, x \rangle}$ for any $x \in \mathcal{H}$. We call \mathcal{H} a Hilbert space if it is complete with respect to the norm.

Let $B(\mathcal{H})$ be the set of all bounded linear operators on the Hilbert space \mathcal{H} . For any $f \in B(\mathcal{H})$ and any $x \in \mathcal{H}$, we sometimes denote f(x) by f.x. From [18], we know that $B(\mathcal{H})$ is a C^* -algebra. In fact, the algebra $B(\mathcal{H})$ is a Banach algebra with the norm given by

$$||f|| = \sup_{\|x\| \le 1, x \in \mathcal{H}} ||f(x)||, \quad \forall f \in B(\mathcal{H}).$$

And then $B(\mathcal{H})$ is a C^{*}-algebra with the involution * determined by

$$\langle f(x), y \rangle = \langle x, f^*(y) \rangle, \quad \forall f \in B(\mathcal{H}), \quad \forall x, y \in \mathcal{H}.$$

For a projection p on the Hilbert space \mathcal{H} , we mean that p is a linear operator on \mathcal{H} satisfying $p^2 = p$ and $p^* = p$. We set $p^{\perp} = \mathrm{id}_H - p$. Then we have $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$,

where $\mathcal{H}_1 = p\mathcal{H}$ and $\mathcal{H}_2 = p^{\perp}\mathcal{H}$ are orthogonal closed subspaces of \mathcal{H} . We call the projection p on \mathcal{H} nontrivial if p is not equal to 0 and $\mathrm{id}_{\mathcal{H}}$.

Using the projections on \mathcal{H} , we can construct Rota-Baxter operators on the C^* -algebra $B(\mathcal{H})$.

Example 3.1. Let p be a projection on \mathcal{H} . We define a linear operator L_p on $B(\mathcal{H})$ by

$$L_p(a) = pa$$

for any $a \in B(\mathcal{H})$. It is easy to verify that L_p is a Rota-Baxter operator of weight -1on $B(\mathcal{H})$. Then $(B(\mathcal{H}), L_p)$ is a Rota-Baxter C^{*}-algebra.

3.2 Rota-Baxter operators on $B(\mathcal{H})$

From here to the end of this subsection, without further mention, we will always assume that \mathcal{H} is a Hilbert space, p is a projection on \mathcal{H} with $\mathcal{H}_1 = p\mathcal{H}$ and $\mathcal{H}_2 = p^{\perp}\mathcal{H}$, \mathcal{A} is a C^* -subalgebra of $B(\mathcal{H})$ and \mathcal{A}_1 is a C^* -subalgebra of \mathcal{A} . Note that we have $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$.

Definition 3.2. Let f be a linear operator on \mathcal{H} and let P be a Rota-Baxter operator of weight $\lambda \in \mathbb{C}$ on \mathcal{A} . The operator P is called a Rota-Baxter operator matching f on \mathcal{A}_1 if

$$P(a).f(x) = f(P(a).x) + f(a.f(x)) + \lambda f(a.x), \qquad \forall a \in \mathcal{A}_1, \ \forall x \in \mathcal{H}.$$

As a special case of the above definition, Rota-Baxter operators matching projections on the Hilbert space \mathcal{H} are the central notion of the article. We first give an equivalent condition for a Rota-Baxter operator of weight -1 which matching projections. Recall that for a Rota-Baxter operator P of weight λ on \mathcal{A} , we set $\tilde{P} = -\lambda \operatorname{id}_{\mathcal{A}} - P$.

Lemma 3.3. Let P be a Rota-Baxter operator of weight -1 on \mathcal{A} . Then P is a Rota-Baxter operator matching p on \mathcal{A}_1 if and only if for any $a \in \mathcal{A}_1$ and any $x \in \mathcal{H}$, we have $P(a).p(x) \in \mathcal{H}_1$ and $\tilde{P}(a).p^{\perp}(x) \in \mathcal{H}_2$.

Proof. Assume that P matches p on \mathcal{A}_1 . For any $x \in \mathcal{H}$ and any $a \in \mathcal{A}_1$, we have

$$P(a).p^{2}(x) = p(P(a).p(x)) + p(a.p^{2}(x)) - p(a.p(x)).$$

As $p^2 = p$, we have

$$P(a).p(x) = p(P(a).p(x)) \in \mathcal{H}_1$$

Similarly, one can show that $\widetilde{P}(a).p^{\perp}(x) \in \mathcal{H}_2$.

Conversely, assume that for any $a \in \mathcal{A}_1$ and any $x \in \mathcal{H}$, we have $P(a).p(x) \in \mathcal{H}_1$ and $\tilde{P}(a).p^{\perp}(x) \in \mathcal{H}_2$. We write $x = x_1 + x_2$ with $x_1 = p(x) \in \mathcal{H}_1$ and $x_2 = p^{\perp}(x) \in \mathcal{H}_2$. Then we have

$$p(P(a).x) + p(a.p(x)) - p(a.x)$$

= $p(P(a).(x_1 + x_2)) + p(a.x_1) - p(a.(x_1 + x_2))$
= $p(P(a).x_1) + p(P(a).x_2) - p(a.x_2)$
= $p(P(a).x_1) - p(\widetilde{P}(a).x_2).$

Since $P(a).x_1 \in \mathcal{H}_1$ and $\widetilde{P}(a).x_2 \in \mathcal{H}_2$, we have

$$p(P(a).x) + p(a.p(x)) - p(a.x) = P(a).x_1 = P(a).p(x),$$

which shows that P matches p on \mathcal{A}_1 .

As p^{\perp} is also a projection, the above lemma has an immediate corollary.

Corollary 3.4. Let P be a Rota-Baxter operator of weight -1 on \mathcal{A} . Then P matches p on \mathcal{A}_1 if and only if \widetilde{P} matches p^{\perp} on \mathcal{A}_1 .

Now we write an element $x = x_1 + x_2 \in \mathcal{H}$ with $x_1 = p(x) \in \mathcal{H}_1$ and $x_2 = p^{\perp}(x) \in \mathcal{H}_2$ as a vector $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$. As shown in [3,Theorem 2.10], for any $a \in B(\mathcal{H})$, since

$$a.x = a(p(x)) + a(p^{\perp}(x))$$

= $p(a(p(x))) + p^{\perp}(a(p(x))) + p(a(p^{\perp}(x))) + p^{\perp}(a(p^{\perp}(x)))$
= $(pap)(x_1) + (pap^{\perp})(x_2) + (p^{\perp}ap)(x_1) + (p^{\perp}ap^{\perp})(x_2),$

we may write a as a matrix

$$a = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix},$$

with

$$a_{11} = pap : \mathcal{H}_1 \to \mathcal{H}_1, \qquad a_{12} = pap^{\perp} : \mathcal{H}_2 \to \mathcal{H}_1,$$

$$a_{21} = p^{\perp}ap : \mathcal{H}_1 \to \mathcal{H}_2, \qquad a_{22} = p^{\perp}ap^{\perp} : \mathcal{H}_2 \to \mathcal{H}_2.$$

It is easy to show that the operations in $B(\mathcal{H})$ coincides with the corresponding matrix operations. For example, we have

$$a^* = \begin{bmatrix} a_{11}^* & a_{21}^* \\ a_{12}^* & a_{22}^* \end{bmatrix},$$

which comes from

$$(a^*)_{11} = pa^*p = (pap)^* = a_{11}^*,$$

$$(a^*)_{12} = pa^*p^{\perp} = (p^{\perp}ap)^* = a_{21}^*,$$

$$(a^*)_{21} = p^{\perp}a^*p = (pap^{\perp})^* = a_{12}^*,$$

$$(a^*)_{22} = p^{\perp}a^*p^{\perp} = (p^{\perp}ap^{\perp})^* = a_{22}^*$$

And if $b = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \in B(\mathcal{H})$, then $ab = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}.$

In fact, we have

$$ab = (pap + pap^{\perp} + p^{\perp}ap + p^{\perp}ap^{\perp})(pbp + pbp^{\perp} + p^{\perp}bp + p^{\perp}bp^{\perp})$$

$$= pappbp + pap^{\perp}p^{\perp}bp + pappbp^{\perp} + pap^{\perp}p^{\perp}bp^{\perp} + p^{\perp}appbp + p^{\perp}ap^{\perp}p^{\perp}bp + p^{\perp}appbp^{\perp} + p^{\perp}ap^{\perp}p^{\perp}bp^{\perp} = p(a_{11}b_{11} + a_{12}b_{21})p + p(a_{11}b_{12} + a_{12}b_{22})p^{\perp} + p^{\perp}(a_{21}b_{11} + a_{22}b_{21})p + p^{\perp}(a_{21}b_{12} + a_{22}b_{22})p^{\perp}.$$

Using the above matrix notation, we may restate Lemma 3.3 as in the following proposition.

Proposition 3.5. Let P be a Rota-Baxter operator of weight -1 on \mathcal{A} . Then P matches p on \mathcal{A}_1 if and only if for any $a \in \mathcal{A}_1$, we have $P(a)_{21} = 0$ and $P(a)_{12} = a_{12}$.

Proof. Assume that P matches p on \mathcal{A}_1 . For any $x_1 \in \mathcal{H}_1$, we have

$$P(a)_{21}(x_1) = (p^{\perp}P(a)p)(p(x)) = p^{\perp}(P(a).p(x)).$$

By Lemma 3.3, $P(a).p(x) \in \mathcal{H}_1$. Hence $P(a)_{21}(x_1) = 0$ and then $P(a)_{21} = 0$. Similarly, for any $x_2 \in \mathcal{H}_2$, we have

$$P(a)_{12}(x_2) = (pP(a)p^{\perp})(p^{\perp}(x)) = p(P(a).p^{\perp}(x)).$$

Since $\widetilde{P} = \operatorname{id} - P$, we have

$$P(a)_{12}(x_2) = p(a.p^{\perp}(x)) - p(\widetilde{P}(a).p^{\perp}(x)) = a_{12}(x_2) - p(\widetilde{P}(a).p^{\perp}(x)).$$

By Lemma 3.3, $\tilde{P}(a).p^{\perp}(x) \in \mathcal{H}_2$. Therefore $P(a)_{12}(x_2) = a_{12}(x_2)$ and then $P(a)_{12} = a_{12}$.

Conversely, assume that for any $a \in \mathcal{A}_1$, we have $P(a)_{21} = 0$ and $P(a)_{12} = a_{12}$. For any $x \in \mathcal{H}$, we have

$$P(a).p(x) = pP(a)p(x_1) + p^{\perp}P(a)p(x_1)$$

= pP(a)p(x_1) + P(a)_{21}(x_1) = pP(a)p(x_1) \in \mathcal{H}_1,

and

$$\widetilde{P}(a).p^{\perp}(x) = p\widetilde{P}(a)p^{\perp}(x_2) + p^{\perp}\widetilde{P}(a)p^{\perp}(x_2) = pap^{\perp}(x_2) - pP(a)p^{\perp}(x_2) + p^{\perp}\widetilde{P}(a)p^{\perp}(x_2) = (a_{12} - P(a)_{12})(x_2) + p^{\perp}\widetilde{P}(a)p^{\perp}(x_2) = p^{\perp}\widetilde{P}(a)p^{\perp}(x_2) \in \mathcal{H}_2.$$

Hence by Lemma 3.3, the Rota-Baxter operator P matches p on \mathcal{A}_1 .

Note that the direct sum $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$ can be regarded as a C^* -subalgebra of $B(\mathcal{H})$ by the following embedding

$$\iota: B(\mathcal{H}_1) \oplus B(\mathcal{H}_2) \to B(\mathcal{H}); (a_1, a_2) \mapsto \begin{bmatrix} a_1 & 0\\ 0 & a_2 \end{bmatrix}$$

The following is the main result of this note.

Theorem 3.6. Let \mathcal{H} be a Hilbert space and let p be a projection on \mathcal{H} . Set $\mathcal{H}_1 = p\mathcal{H}$ and $\mathcal{H}_2 = p^{\perp}\mathcal{H}$. Then there is a 1-1 correspondence between the following two sets: (1) the set of Rota-Baxter operators of weight -1 on $B(\mathcal{H})$ matching p on $B(\mathcal{H})$;

(2) the set of Rota-Baxter operators of weight -1 on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$.

Proof. Denote by Γ the set of Rota-Baxter operators of weight -1 on $B(\mathcal{H})$ matching p on $B(\mathcal{H})$, and by Λ the set of Rota-Baxter operators of weight -1 on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$.

Assume that $P \in \Gamma$. For any $a_1 \in B(\mathcal{H}_1)$ and $a_2 \in B(\mathcal{H}_2)$, we have $a = \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix} \in B(\mathcal{H})$ and

$$P(a) = \begin{bmatrix} P(a)_{11} & 0\\ 0 & P(a)_{22} \end{bmatrix} \in B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$$

by Proposition 3.5. Hence we get a linear operator P' on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$ defined by $P' = P|_{B(\mathcal{H}_1)\oplus B(\mathcal{H}_2)}$. It is obvious that P' is a Rota-Baxter operator of weight -1 on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$. Hence we have a map $\Phi : \Gamma \to \Lambda$ defined by $\Phi(P) = P'$.

Conversely, assume that $P' \in \Lambda$. For any $a = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in B(\mathcal{H})$, we have $\begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix} \in B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$. We then define an operator P on $B(\mathcal{H})$ by

$$P(a) = P'\left(\begin{bmatrix} a_{11} & 0\\ 0 & a_{22} \end{bmatrix} \right) + \begin{bmatrix} 0 & a_{12}\\ 0 & 0 \end{bmatrix}.$$

It is obvious that P is linear.

We show that P is a Rota-Baxter operator of weight -1 on $B(\mathcal{H})$. Assume that $P'\left(\begin{bmatrix}a_{11} & 0\\0 & a_{22}\end{bmatrix}\right) = \begin{bmatrix}a'_{11} & 0\\0 & a'_{22}\end{bmatrix}$, then we have $P(a) = \begin{bmatrix}a'_{11} & 0\\0 & a'_{22}\end{bmatrix} + \begin{bmatrix}0 & a_{12}\\0 & 0\end{bmatrix}$.

For another element $b = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \in B(\mathcal{H})$, we set $P'\left(\begin{bmatrix} b_{11} & 0 \\ 0 & b_{22} \end{bmatrix}\right) = \begin{bmatrix} b'_{11} & 0 \\ 0 & b'_{22} \end{bmatrix}$. Hence we have

$$P(a)P(b) = \begin{bmatrix} a'_{11} & a_{12} \\ 0 & a'_{22} \end{bmatrix} \begin{bmatrix} b'_{11} & b_{12} \\ 0 & b'_{22} \end{bmatrix} = \begin{bmatrix} a'_{11}b'_{11} & a'_{11}b_{12} + a_{12}b'_{22} \\ 0 & a'_{22}b'_{22} \end{bmatrix}.$$

Since

$$aP(b) = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b'_{11} & b_{12} \\ 0 & b'_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b'_{11} & a_{11}b_{12} + a_{12}b'_{22} \\ a_{21}b'_{11} & a_{21}b_{12} + a_{22}b'_{22} \end{bmatrix},$$

we get

$$P(aP(b)) = P'\left(\begin{bmatrix} a_{11}b'_{11} & 0\\ 0 & a_{21}b_{12} + a_{22}b'_{22} \end{bmatrix} \right) + \begin{bmatrix} 0 & a_{11}b_{12} + a_{12}b'_{22}\\ 0 & 0 \end{bmatrix}.$$

Similarly, we have

$$P(P(a)b) = P'\left(\begin{bmatrix} a'_{11}b_{11} + a_{12}b_{21} & 0\\ 0 & a'_{22}b_{22} \end{bmatrix} \right) + \begin{bmatrix} 0 & a'_{11}b_{12} + a_{12}b_{22}\\ 0 & 0 \end{bmatrix}$$

and

$$P(ab) = P'\left(\begin{bmatrix}a_{11}b_{11} + a_{12}b_{21} & 0\\ 0 & a_{21}b_{12} + a_{22}b_{22}\end{bmatrix}\right) + \begin{bmatrix}0 & a_{11}b_{12} + a_{12}b_{22}\\ 0 & 0\end{bmatrix}.$$

Therefore we find

$$P(aP(b)) + P(P(a)b) - P(ab)$$

= $P'\left(\begin{bmatrix}a_{11}b'_{11} + a'_{11}b_{11} - a_{11}b_{11} & 0\\ 0 & a_{22}b'_{22} + a'_{22}b_{22} - a_{22}b_{22}\end{bmatrix}\right) + \begin{bmatrix}0 & a_{12}b'_{22} + a'_{11}b_{12}\\ 0 & 0\end{bmatrix}.$

On the other hand, since $P' \in \Lambda$, we have

$$\begin{bmatrix} a'_{11} & 0\\ 0 & a'_{22} \end{bmatrix} \begin{bmatrix} b'_{11} & 0\\ 0 & b'_{22} \end{bmatrix}$$

= $P' \left(\begin{bmatrix} a_{11} & 0\\ 0 & a_{22} \end{bmatrix} \begin{bmatrix} b'_{11} & 0\\ 0 & b'_{22} \end{bmatrix} + \begin{bmatrix} a'_{11} & 0\\ 0 & a'_{22} \end{bmatrix} \begin{bmatrix} b_{11} & 0\\ 0 & b_{22} \end{bmatrix} - \begin{bmatrix} a_{11}b_{11} & 0\\ 0 & a_{22}b_{22} \end{bmatrix} \right).$

Then we get

$$P(aP(b)) + P(P(a)b) - P(ab) = \begin{bmatrix} a'_{11}b'_{11} & 0\\ 0 & a'_{22}b'_{22} \end{bmatrix} + \begin{bmatrix} 0 & a_{12}b'_{22} + a'_{11}b_{12}\\ 0 & 0 \end{bmatrix},$$

which implies that

$$P(a)P(b) = P(aP(b)) + P(P(a)b) - P(ab).$$

So P is a Rota-Baxter operator of weight -1 on $B(\mathcal{H})$. And from Proposition 3.5, we see that P matches p on $B(\mathcal{H})$. Then we have a map $\Psi : \Lambda \to \Gamma$ defined by $\Psi(P') = P$.

It is easy to verify that $\Psi \circ \Phi$ and $\Phi \circ \Psi$ are identity maps. And so there is a 1-1 correspondence between Λ and Γ .

Remark 3.7. We can't extend the above theorem to the C^* -subalgebra \mathcal{A} of $B(\mathcal{H})$. For a Rota-Baxter operator P' of weight -1 on $p\mathcal{A}p \oplus p^{\perp}\mathcal{A}p^{\perp}$, as p and p^{\perp} may not be in \mathcal{A} , we can't construct a Rota-Baxter operator P of weight -1 matching p on \mathcal{A} as above.

From Theorem 3.6, we can construct Rota-Baxter operators of weight -1 on $B(\mathcal{H})$ matching p on $B(\mathcal{H})$ from the Rota-Baxter operators of weight -1 on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$. The following lemma gives a method to construct Rota-Baxter operators on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$.

Lemma 3.8. Let \mathcal{A}_1 and \mathcal{A}_2 be two C^* -algebras. If P_1 and P_2 are Rota-Baxter operators of weight $\lambda \in \mathbb{C}$ on \mathcal{A}_1 and \mathcal{A}_2 , respectively, then the operator

$$P = P_1 \oplus P_2 : \mathcal{A}_1 \oplus \mathcal{A}_2 \to \mathcal{A}_1 \oplus \mathcal{A}_2; (a_1, a_2) \mapsto (P_1(a_1), P_2(a_2))$$

is a Rota-Baxter operator of weight λ on $\mathcal{A}_1 \oplus \mathcal{A}_2$.

Proof. Is is trivial to check.

Then using Lemma 3.8 and Theorem 3.6, we get

Corollary 3.9. Let P_1 be a Rota-Baxter operator of weight -1 on $B(\mathcal{H}_1)$ and let P_2 be a Rota-Baxter operator of weight -1 on $B(\mathcal{H}_2)$. Then the operator

$$P = \begin{bmatrix} P_1 & \text{id} \\ 0 & P_2 \end{bmatrix} : B(\mathcal{H}) \to B(\mathcal{H}); \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \mapsto \begin{bmatrix} P_1(a_{11}) & a_{12} \\ 0 & P_2(a_{22}) \end{bmatrix}$$

is a Rota-Baxter operator of weight -1 matching p on $B(\mathcal{H})$.

Recall that a Rota-Baxter operator P of weight λ on \mathcal{A} is called symmetric on \mathcal{A}_1 provided that

$$P(a^*) = P(a)^*, \quad \forall a \in \mathcal{A}_1.$$

Proposition 3.10. If there is a Rota-Baxter operator P of weight -1 on \mathcal{A} which is symmetric and matches p on \mathcal{A}_1 , then we have $\mathcal{A}_1 \subseteq B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$.

Proof. For any $a = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \in \mathcal{A}_1$, from Proposition 3.5, we have

$$P(a) = \begin{bmatrix} P(a)_{11} & a_{12} \\ 0 & P(a)_{22} \end{bmatrix}.$$

On the other hand, since P is symmetric on \mathcal{A}_1 , we have

$$P(a) = P(a^*)^* = \begin{bmatrix} (P(a^*)_{11})^* & 0\\ a_{21} & (P(a^*)_{22})^* \end{bmatrix}.$$

Then we get $a_{12} = 0$ and $a_{21} = 0$. So $\mathcal{A}_1 \subseteq B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$.

To state the last result of this subsection, we recall the definition of *-representation from [18]. A *-representation π of the C*-algebra \mathcal{A} is a *-homomorphism $\pi : \mathcal{A} \to \mathcal{B}(\mathcal{H})$ with \mathcal{H} a Hilbert space. We denote this *-representation of \mathcal{A} by $\{\pi, \mathcal{H}\}$. We say $\{\pi, \mathcal{H}\}$ is topologically irreducible if $\pi(\mathcal{A})$ has no proper invariant subspaces. It is called algebraically irreducible if it has no proper invariant manifolds (subspaces of \mathcal{H} that are not necessarily closed). These two notions coincide for C*-algebras from [5]. Hence we call $\{\pi, \mathcal{H}\}$ irreducible when either of the two conditions holds.

Corollary 3.11. Let $\{\pi, \mathcal{H}\}$ be a *-representation of a C*-algebra \mathcal{A} . Then the representation $\{\pi, \mathcal{H}\}$ is irreducible if and only if there is no Rota-Baxter operator of weight -1 on $B(\mathcal{H})$ which is symmetric and matches p on $\pi(\mathcal{A})$ for some nontrivial projection p on \mathcal{H} .

Proof. If there is such a Rota-Baxter operator on $\pi(\mathcal{A})$, then from Proposition 3.10, we have $\pi(\mathcal{A}) \subseteq B(p\mathcal{H}) \oplus B(p^{\perp}\mathcal{H})$. Hence $\pi(\mathcal{A})$ has a proper invariant closed subspace $p\mathcal{H}$ of \mathcal{H} . Therefore $\{\pi, \mathcal{H}\}$ is not irreducible.

Conversely, if the representation $\{\pi, \mathcal{H}\}$ is not irreducible, then $\pi(\mathcal{A})$ has a proper invariant closed subspace \mathcal{H}_1 . There exists a projection p on \mathcal{H} such that $\mathcal{H}_1 = p\mathcal{H}$. Therefore $\pi(\mathcal{A}).p\mathcal{H} \subseteq p\mathcal{H}$. Define a linear operator P on $B(\mathcal{H})$ by

$$P(a) = pap + p^{\perp}ap^{\perp} + pap^{\perp}$$

for any $a \in B(\mathcal{H})$. Then by the proof of Theorem 3.6, P is a Rota-Baxter operator of weight -1 matching p on $B(\mathcal{H})$. For any $b \in \pi(\mathcal{A})$, we have b = P(b) and $b^* = P(b^*)$ from Proposition 3.5. Hence we find

$$P(b) = b = (b^*)^* = P(b^*)^*,$$

which means that P is symmetric on $\pi(\mathcal{A})$.

At the end of this subsection, we see an example.

Example 3.12. Let $f : \mathcal{A} \to \mathbb{C}$ be a linear function, where \mathcal{A} is a C^* -algebra. The function f is called a positive linear functional if $f(a) \ge 0$ for any $a \ge 0$ in \mathcal{A} (the notion of $a \ge 0$ can be found in Section 4). The norm of f is defined by

$$||f|| = \sup_{a \in \mathcal{A}, a \neq 0} \frac{|f(a)|}{||a||}.$$

As in [5], the function f is called a state on \mathcal{A} if f is a positive linear functional and ||f|| = 1. The state f is called a pure state if there is not a $\lambda \in (0, 1)$ and two states f_1 and f_2 on \mathcal{A} , such that $f = \lambda f_1 + (1 - \lambda) f_2$.

From [5, Theorem I.9.6], for any state f on \mathcal{A} , we can construct a Hilbert space \mathcal{H}_f and a *-representation $\{\pi_f, \mathcal{H}_f\}$ of \mathcal{A} . This method of building representations from states is called GNS-constructions. From [5, Theorem I.9.8], we know that the representation $\{\pi_f, \mathcal{H}_f\}$ is an irreducible representation of \mathcal{A} if and only if f is a pure state. Hence if f is not a pure state, then from Corollary 3.11, there is at least one Rota-Baxter operator of weight -1 which is symmetric and matches p on $\pi_f(\mathcal{A})$ for some nontrivial projection p on \mathcal{H}_f .

3.3 Representations of Rota-Baxter C^{*}-algebras

We introduce the notion of representations of Rota-Baxter C^* -algebras.

Definition 3.13. Let \mathcal{A} be a C^* -algebra and let P be a Rota-Baxter operator of weight λ on \mathcal{A} . Let \mathcal{H} be a Hilbert space and let P' be a Rota-Baxter operator of weight λ on $B(\mathcal{H})$. Let $\pi : (\mathcal{A}, P) \to (B(\mathcal{H}), P')$ be a Rota-Baxter C^* -algebra homomorphism and let f be a linear operator on \mathcal{H} . Then $\{\pi, \mathcal{H}, f\}$ is a Rota-Baxter *-representation of (\mathcal{A}, P) into $(B(\mathcal{H}), P')$ if P' matches f on $\pi(\mathcal{A})$.

Hence $\{\pi, \mathcal{H}, f\}$ is a Rota-Baxter *-representation of (\mathcal{A}, P) into $(B(\mathcal{H}), P')$ if for any $h \in \mathcal{H}$ and any $a \in \mathcal{A}$ we have

$$\pi(P(a))(f(h)) = f(\pi(P(a))(h)) + f(\pi(a)(f(h))) + \lambda f(\pi(a)(h)).$$

We first prepare two lemmas.

Lemma 3.14. Let \mathcal{H} be a Hilbert space and let p be a projection on \mathcal{H} . Let P' be a Rota-Baxter operator of weight -1 on $B(p\mathcal{H}) \oplus B(p^{\perp}\mathcal{H})$ and let P be the Rota-Baxter operator of weight -1 on $B(\mathcal{H})$ matching p that corresponding to P' (See Theorem 3.6.). Then the embedding $\iota : B(p\mathcal{H}) \oplus B(p^{\perp}\mathcal{H}) \to B(\mathcal{H})$ is a Rota-Baxter C^* -algebra homomorphism.

Proof. It needs to show that $\iota \circ P' = P \circ \iota$, which is obvious.

Lemma 3.15. Let $\pi_1 : (\mathcal{A}_1, P_1) \to (\mathcal{A}_2, P_2)$ and $\pi_2 : (\mathcal{A}'_1, P'_1) \to (\mathcal{A}'_2, P'_2)$ be two Rota-Baxter C^{*}-algebra homomorphisms. Then the map

 $\pi = \pi_1 \oplus \pi_2 : \mathcal{A}_1 \oplus \mathcal{A}'_1 \to \mathcal{A}_2 \oplus \mathcal{A}'_2; (a_1, b_1) \mapsto (\pi_1(a_1), \pi_2(b_1))$

is a Rota-Baxter C*-algebra homomorphism.

Proof. It is easy to verify that π is a Rota-Baxter C^* -algebra homomorphism. Then we state the main theorem of this subsection.

Theorem 3.16. Let \mathcal{H}_1 and \mathcal{H}_2 be two Hilbert spaces. For i = 1, 2, let P'_i be a Rota-Baxter operator of weight -1 on $B(\mathcal{H}_i)$, \mathcal{A}_i be a C^* -subalgebra of $B(\mathcal{H}_i)$, and let P_i be a Rota-Baxter operator of weight -1 on \mathcal{A}_i . Set $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$. Let P' be the unique Rota-Baxter operator of weight -1 on $B(\mathcal{H})$ corresponding to $P'_1 \oplus P'_2$ on $B(\mathcal{H}_1) \oplus B(\mathcal{H}_2)$ (See Theorem 3.6 and Lemma 3.8.). Let p be the projection from \mathcal{H} onto \mathcal{H}_1 . Then the following two statements are equivalent:

- (a) There are Rota-Baxter *-representations $\{\pi_i, \mathcal{H}_i, \mathrm{id}_{\mathcal{H}_i}\}\$ from (\mathcal{A}_i, P_i) into $(B(\mathcal{H}_i), P'_i)$ with i = 1, 2;
- (b) There is a Rota-Baxter *-representation $\{\pi, \mathcal{H}, p\}$ from $(\mathcal{A}_1 \oplus \mathcal{A}_2, P_1 \oplus P_2)$ into $(B(\mathcal{H}), P')$ such that $\pi(\mathcal{A}_i) \subseteq B(\mathcal{H}_i)$ for i = 1, 2.

Proof. Set $\mathcal{A} = \mathcal{A}_1 \oplus \mathcal{A}_2$ and $P = P_1 \oplus P_2$.

(a) \Rightarrow (b) Assume that (a) holds. Using Lemmas 3.14 and 3.15, we have a Rota-Baxter C^* -algebra homomorphism

$$\pi: (\mathcal{A}_1 \oplus \mathcal{A}_2, P_1 \oplus P_2) \xrightarrow{\pi_1 \oplus \pi_2} (B(\mathcal{H}_1) \oplus B(\mathcal{H}_2), P'_1 \oplus P'_2) \xrightarrow{\iota} (B(\mathcal{H}), P').$$

It is obvious that $\pi(\mathcal{A}_i) \subseteq B(\mathcal{H}_i)$ for i = 1, 2.

We check that P' matches p on $\pi(\mathcal{A})$. For any $a = (a_1, a_2) \in \mathcal{A}$ with $a_i \in \mathcal{A}_i$ and any $h = (h_1, h_2) \in \mathcal{H}$ with $h_i \in \mathcal{H}_i$, we have

$$p(\pi(P(a))(h)) + p(\pi(a)(p(h))) - p(\pi(a)(h))$$

= $p(\pi_1(P_1(a_1))(h_1), \pi_2(P_2(a_2))(h_2)) + p(\pi_1(a_1)(h_1), 0) - p(\pi_1(a_1)(h_1), \pi_2(a_2)(h_2))$
= $\pi_1(P_1(a_1))(h_1),$

and

$$\pi(P(a))(p(h)) = (\pi_1(P_1(a_1)), \pi_2(P_2(a_2)))(h_1, 0) = \pi_1(P_1(a_1))(h_1).$$

Hence P' matches p on $\pi(\mathcal{A})$.

(a) \Leftarrow (b) Assume that (b) holds. As π is a Rota-Baxter C*-algebra homomorphism, for any $a_1 \in \mathcal{A}_1$ and $a_2 \in \mathcal{A}_2$, we have

$$\pi \circ P(a_1, a_2) = P' \circ \pi(a_1, a_2).$$

Taking $a_2 = 0$, we get $\pi \circ P_1(a_1) = P' \circ \pi(a_1)$. Since $P_1(a_1) \in \mathcal{A}_1$ and $\pi(a_1) \in B(\mathcal{H}_1)$, we have

$$\pi \circ P_1(a_1) = P_1' \circ \pi(a_1).$$

Hence we get a Rota-Baxter C^* -algebra homomorphism

$$\pi_1 = \pi|_{\mathcal{A}_1} : (\mathcal{A}_1, P_1) \to (B(\mathcal{H}_1), P_1').$$

Similarly, we have a Rota-Baxter C^* -algebra homomorphism

$$\pi_2 = \pi|_{\mathcal{A}_2} : (\mathcal{A}_2, P_2) \to (B(\mathcal{H}_2), P_2').$$

Finally, since P' matches p on $\pi(\mathcal{A})$, one can show that P'_i matches $\mathrm{id}_{\mathcal{H}_i}$ on $\pi_i(\mathcal{A}_i)$ for i = 1, 2.

4 Quasidiagonal operators and Rota-Baxter operators

In this section, all Hilbert spaces and C^* -algebras are assumed to be separable. Here a topology space X is separable if it contains a countable dense subset.

We recall some basic definitions of quasidiagonal linear operators from [3]. Let \mathcal{A} be a C^* -algebra and let a be a self-adjoint element of \mathcal{A} . If \mathcal{A} is unital, the spectrum of a is defined by

 $\operatorname{Spec}(a) = \{\lambda \in \mathbb{C} | a - \lambda 1_{\mathcal{A}} \text{ is not invertible} \},\$

where $1_{\mathcal{A}}$ is the identity element of \mathcal{A} . If \mathcal{A} is nonunital, then from [18, Proposition 1.1.7], there is a unital C^* -algebra \mathcal{A}_1 such that $\mathcal{A}_1 \simeq \mathcal{A} \bigoplus \mathbb{C}$. Then for any $a \in \mathcal{A}$, the spectrum of a is the spectrum of a as an element of \mathcal{A}_1 . The element a is called positive if Spec(a) is contained in the non-negative reals. For any $a, b \in \mathcal{A}$, we say that a is not bigger than b if b - a is positive, and we denote this by $a \leq b$. We also denote ab - ba by the Lie bracket [a, b].

Let \mathcal{H} be a Hilbert space. A projection p on \mathcal{H} is of finite rank if $p\mathcal{H}$ is of finite dimension. For any sequence $a_n \in B(\mathcal{H})$ and $a \in B(\mathcal{H})$, we say that $a_n \to a$ as $n \to \infty$ in the strong operator topology if $||a_n.h - a.h|| \to 0$ as $n \to \infty$ for any $h \in \mathcal{H}$.

We first give the definition of block diagonal linear operators.

Definition 4.1. A bounded linear operator d on a Hilbert space \mathcal{H} is called block diagonal if there exists an increasing sequence of finite rank projections $p_1 \leq p_2 \leq p_3 \leq \cdots$ on \mathcal{H} , such that $\|[d, p_n]\| = 0$ and $p_n \to id_{\mathcal{H}}$ as $n \to \infty$ in the strong operator topology.

We can construct an equivalent condition of block diagonal operators with the help of Rota-Baxter operators. Let \mathcal{A} be a C^* -algebra. For any $a \in \mathcal{A}$, there is a C^* subalgebra of \mathcal{A} generated by a, which is denoted by $C^*(a)$. In fact, $C^*(a)$ is the intersection of all C^* -subalgebras of \mathcal{A} containing a, and $C^*(a)$ is the closure of the algebra generated by a and a^* .

Proposition 4.2. Let \mathcal{H} be a Hilbert space and $d \in B(\mathcal{H})$. Then the following statements are equivalent:

- (a) The operator d is block diagonal;
- (b) There exists an increasing sequence of finite rank projections p₁ ≤ p₂ ≤ p₃ ≤ ··· on H and a sequence of Rota-Baxter operators {P_n} of weight −1 on B(H), such that p_n → id_H as n → ∞ in the strong operator topology, and P_n are symmetric on C^{*}(d) and matches p_n on B(H) for any n ≥ 1.

Proof. For any $x \in \mathcal{H}, n \in \mathbb{N}$ and $b \in B(\mathcal{H})$, we have $p_n^{\perp} b p_n(x) \in p_n^{\perp} \mathcal{H}$ and $p_n b p_n^{\perp}(x) \in p_n \mathcal{H}$. Therefore we find

$$||[b, p_n]|| = ||p_n^{\perp}bp_n - p_nbp_n^{\perp}|| \ge ||p_n^{\perp}bp_n||.$$

And similarly we have

$$||[b, p_n]|| \ge ||p_n b p_n^{\perp}||.$$

On the other hand, it holds

$$||[b, p_n]|| = ||p_n^{\perp} b p_n - p_n b p_n^{\perp}|| \le ||p_n^{\perp} b p_n|| + ||p_n b p_n^{\perp}||.$$

Therefore $||[b, p_n]|| = 0$ if and only if $||p_n^{\perp}bp_n|| = 0$ and $||p_nbp_n^{\perp}|| = 0$.

(a) \Rightarrow (b) Assume that *d* is block diagonal. Then there exists an increasing sequence of finite rank projections $p_1 \leq p_2 \leq p_3 \leq \cdots$ on \mathcal{H} , such that $||[d, p_n]|| = 0$ for any $n \in \mathbb{N}$.

Let A be the set of elements which are finite products of d and d^* in $C^*(d)$. For any $b \in A$, if $b = b_1 b_2 \cdots b_n$ with $b_k = d$ or d^* for $1 \leq k \leq n$, we set |b| = n. We first prove that for any $b \in A$, $||[b, p_n]|| = 0$ by induction on |b|. The case of |b| = 1 is trivial. For $|b| \geq 2$, we can assume that there is a $b_1 \in A$ such that $|b_1| = |b| - 1$, and $b = b_1 d$ without loss of generality. We have

$$\begin{aligned} \|p_n^{\perp}bp_n\| &= \|p_n^{\perp}b_1p_nd + p_n^{\perp}b_1[d, p_n]\| \\ &\leq \|p_n^{\perp}b_1p_nd\| + \|p_n^{\perp}b_1[d, p_n]\| \\ &\leq \|p_n^{\perp}b_1p_n\|\|d\| + \|p_n^{\perp}\|\|b_1\|\|[d, p_n]\| \end{aligned}$$

Hence using the inductive assumption $||p_n^{\perp}b_1p_n|| = 0$ and the fact $||[d, p_n]|| = 0$, we find $||p_n^{\perp}bp_n|| = 0$. Similarly we get $||p_nbp_n^{\perp}|| = 0$. Therefore we have $||[b, p_n]|| = 0$.

Using the above result, it is easy to know that for any $b \in C^*(d)$, we have $||[b, p_n]|| = 0$. Define a linear operator P_n on $B(\mathcal{H})$ such that for any $a \in B(\mathcal{H})$ we have

$$P_n(a) = p_n a p_n + p_n^{\perp} a p_n^{\perp} + p_n a p_n^{\perp}.$$

Then P_n is a Rota-Baxter operator of weight -1 matching p_n on $B(\mathcal{H})$ by Corollary 3.9. Finally, for any $b \in C^*(d)$, as

$$\begin{aligned} \|P_{n}(b) - P_{n}(b^{*})^{*}\| \\ &= \|p_{n}bp_{n} + p_{n}^{\perp}bp_{n}^{\perp} + p_{n}bp_{n}^{\perp} - p_{n}bp_{n} - p_{n}^{\perp}bp_{n}^{\perp} - p_{n}^{\perp}bp_{n}\| \\ &= \|p_{n}bp_{n}^{\perp} - p_{n}^{\perp}bp_{n}\| \\ &\leq \|p_{n}bp_{n}^{\perp}\| + \|p_{n}^{\perp}bp_{n}\| = 0, \end{aligned}$$

we have $P_n(b) = P_n(b^*)^*$.

(b) \Rightarrow (a) Assume that (b) holds. Then using Proposition 3.10, we know that $C^*(d) \subseteq B(p_n \mathcal{H}) \bigoplus B(p_n^{\perp} \mathcal{H})$ for any $n \in \mathbb{N}$. Hence $||p_n dp_n^{\perp}|| = ||p_n^{\perp} dp_n|| = 0$. Therefore $||[d, p_n]|| = 0$, and (a) holds.

Now we give the definition of quasidiagonal linear operators.

Definition 4.3. A bounded linear operator d on a Hilbert space \mathcal{H} is called quasidiagonal if there exists an increasing sequence of finite rank projections $p_1 \leq p_2 \leq p_3 \leq \cdots$ on \mathcal{H} , such that $\|[d, p_n]\| \to 0$ as $n \to \infty$ and $p_n \to id_{\mathcal{H}}$ as $n \to \infty$ in the strong operator topology.

As above, we have an equivalent condition of quasidiagonal operators in terms of Rota-Baxter operators.

Proposition 4.4. Let \mathcal{H} be a Hilbert space and $d \in B(\mathcal{H})$. Then the following statements are equivalent:

- (a) The operator d is quasidiagonal;
- (b) There exists an increasing sequence of finite rank projections $p_1 \leq p_2 \leq p_3 \leq \cdots$ on \mathcal{H} and a sequence of Rota-Baxter operators $\{P_n\}$ of weight -1 on $B(\mathcal{H})$, such that P_n matches p_n on $B(\mathcal{H})$ for any $n \geq 1$, $p_n \to \mathrm{id}_{\mathcal{H}}$ and $||P_n(b) - P_n(b^*)^*|| \to 0$ for any $b \in C^*(d)$ as $n \to \infty$.

Proof. For any $x \in \mathcal{H}$ and $b \in B(\mathcal{H})$, similarly as in the proof of Proposition 4.2, we have

$$||[b, p_n]|| \ge ||p_n^{\perp}bp_n||, \quad ||[b, p_n]|| \ge ||p_nbp_n^{\perp}||,$$

and

$$||[b, p_n]|| \le ||p_n^{\perp} b p_n|| + ||p_n b p_n^{\perp}||.$$

Therefore $||[b, p_n]|| \to 0$ as $n \to \infty$ if and only if $||p_n^{\perp}bp_n|| \to 0$ and $||p_nbp_n^{\perp}|| \to 0$ as $n \to \infty$.

(a) \Rightarrow (b) Assume that d is quasidiagonal. Then there exists an increasing sequence of finite rank projections $p_1 \leq p_2 \leq p_3 \leq \cdots$ on \mathcal{H} , such that $||[d, p_n]|| \rightarrow 0$ and $p_n \rightarrow \mathrm{id}_{\mathcal{H}}$ as $n \rightarrow \infty$.

Let A be the set of elements which are finite products of d and d^* in $C^*(d)$. We first prove that for any $b \in A$, $||[b, p_n]|| \to 0$ as $n \to \infty$ by induction on |b|. The case of |b| = 1 is trivial. For $|b| \ge 2$, we can assume that there is a $b_1 \in A$ such that $|b_1| = |b| - 1$, and $b = b_1 d$ without loss of generality. We have

$$\|p_n^{\perp}bp_n\| \le \|p_n^{\perp}b_1p_n\| \|d\| + \|p_n^{\perp}\| \|b_1\| \|[d, p_n]\|.$$

Hence using the inductive assumption, we find $\|p_n^{\perp}bp_n\| \to 0$ as $n \to \infty$. Similarly we get $\|p_n bp_n^{\perp}\| \to 0$ as $n \to \infty$. Therefore we have $\|[b, p_n]\| \to 0$ as $n \to \infty$.

Using the above result, it is easy to know that for any $b \in C^*(d)$, we have $||[b, p_n]|| \to 0$ as $n \to \infty$.

Define a linear operator P_n on $B(\mathcal{H})$ such that for any $a \in B(\mathcal{H})$ we have

$$P_n(a) = p_n a p_n + p_n^{\perp} a p_n^{\perp} + p_n a p_n^{\perp}.$$

Then P_n is a Rota-Baxter operator of weight -1 matching p_n on $B(\mathcal{H})$ by Corollary 3.9. Finally, for any $b \in C^*(d)$, as

$$||P_n(b) - P_n(b^*)^*|| \le ||p_n b p_n^{\perp}|| + ||p_n^{\perp} b p_n||,$$

we have $||P_n(b) - P_n(b^*)^*|| \to 0$ as $n \to \infty$.

(a) \Leftarrow (b) Assume that (b) holds. For any $\epsilon > 0$, there is a N such that for any n > N we have

$$\|P_n(d) - P_n(d^*)^*\| < \epsilon.$$

Then we find

$$\|p_n^{\perp} dp_n\| = \left\| \begin{bmatrix} 0 & 0 \\ -d_{21} & 0 \end{bmatrix} \right\| \le \left\| \begin{bmatrix} 0 & d_{12} \\ -d_{21} & 0 \end{bmatrix} \right\|$$
$$= \|P_n(d) - P_n(d^*)^*\| \le \epsilon.$$

Hence $\|p_n^{\perp} dp_n\| \to 0$ as $n \to \infty$. We can prove that $\|p_n dp_n^{\perp}\| \to 0$ similarly. Finally we have $\|[d, p_n]\| \to 0$ as $n \to \infty$.

Such a sequence $\{P_n\}$ of Rota-Baxter operators of weight -1 on $B(\mathcal{H})$ in (b) of the above proposition is called a quasi symmetric Rota-Baxter operator sequence on $B(\mathcal{H})$.

References

- [1] M. Aguiar, Pre-Poisson algebras, Lett. Math. Phys. 54 (4) (2000), 263-277.
- [2] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.* 10 (1960), 731-742.
- [3] N. P. Brown, On quasidiagonal C*-algebras, Operator algebras and applications, 19-64, Adv. Stud. Pure Math. 38, Math. Soc. Japan, Tokyo, 2004.
- [4] A. Connes, D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Comm. Math. Phys.* **210** (1) (2000), 249-273.
- [5] K. R. Davidson, C^{*}-algebras by example, Fields Institute Monographs, 6, American Mathematical Society, Providence, RI, 1996.
- [6] I. Gelfand, M. Neumark, On the imbedding of normed rings into the ring of operators in Hilbert space, *Rec. Math. [Mat. Sbornik] N.S.* 12 (54) (1943), 197-213.
- [7] I. Gelfand, D. Raikov, Irreducible unitary representations of locally bicompact groups, *Rec. Math. [Mat. Sbornik] N. S.* 13 (55) (1943), 301-316.
- [8] L. Guo, An introduction to Rota-Baxter algebra, Surveys of Modern Mathematics,
 4, International Press, Somerville, MA; Higher Education Press, Beijing, 2012.
- [9] L. Guo, Z. Lin, Representations and modules of Rota-Baxter algebras, arXiv:1905.01531v2.
- [10] R. V. Kadison, Isometries of operator algebras, Ann. of Math. 54 (2) (1951), 325-338.
- [11] R. V. Kadison, Derivation of operator algebras, Ann. of Math. 83 (2) (1966), 280-293.
- [12] Z. Lin, L. Qiao, Representations of Rota-Baxter algebras and regular singular decompositions, arXiv:1603.05912.
- [13] F. J. Murray, J. von Neumann, On rings of operators, Ann. of Math. 37 (1) (1936), 116-229.
- [14] R. T. Prosser, On the ideal structure of operator algebras, Mem. Amer. Math. Soc, 45 (1963), ii+28 pp.
- [15] L. Qiao, J. Pei, Representations of polynomial Rota-Baxter algebras, J. Pure Appl. Algebra 222 (7) (2018), 1738-1757.
- [16] C. E. Rickart, The uniqueness of norm problem in Banach algebras, Ann. of Math. 51 (1950), 615-628.
- [17] S. Sakai, Derivations of W^{*}-algebras, Ann. of Math. 83 (1966), 273-279.

- [18] S. Sakai, C^* -algebras and W^* -algebras, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band **60**, Springer-Verlag, New York-Heidelberg, 1971.
- [19] I. E. Segal, Irreducible representations of operator algebras, Bull. Amer. Math. Soc. 53 (1947), 73-88.
- [20] J. von Neumann, Zur Algebra der Funktionaloperationen und Theorie der normalen Operatoren, Math. Ann. 102 (1) (1930), 370-427.