

GARSDIE THEORY: A COMPOSITION–DIAMOND LEMMA POINT OF VIEW

VIKTOR LOPATKIN

ABSTRACT. This paper shows how to obtain the key concepts and notations of Garside theory by using machinery of a Composition–Diamond lemma. We also show in some cases the greedy normal form is exactly a Gröbner–Shirshov normal form and a family of a left-cancellative category is to be a Garside family if and only if a suitable set of reductions is to be confluent up to some congruence on words.

INTRODUCTION

Throughout this note, all rings are nonzero, associative with identity, all categories are small (objects and morphisms are sets), unless otherwise stated.

The Braid Theory has been formalized in the classical E. Artin’s work [2]. Later, in Garside’s works it has been showed that the braid groups has a positive representation. This concept has been considered by S.I. Adjan [1] and W. Thurston [16] (independently), as a result we have the Adjan–Thurston representation of the the braid group.

W. Thurston [16] constructed a finite state automaton (see also [11, Chapter 9]), having as the set of states the positive non-repeating braids, *i.e.*, any two of its strands cross at most once. The concept of non-repeating braids is very useful, because the total (algebraic) number of crossings of two given strands in a braid is clearly an invariant of isotopy. Since a positive braid has only positive crossing, the absolute number of crossings of two strands in a positive braid is an invariant of isotopy. This idea is very useful, because we can forget about isotopic equivalence and, moreover, there is a bijection between the set of non-repeating braids and permutations.

An interesting characteristic of Thurston’s automaton is that, after a word is imputed, the state is the maximal head of the word that lies in the set of non-repeating braids. This automaton allowed proving that the braid group is automatic. Moreover, this automaton rewrites any word into a canonical form which is called the (left or right) greedy normal form.

Roughly speaking, the describing of rewriting procedure of braids can be described as follows. We have a binary operation. This operation is resulted from the construction of this automaton, that is, the Thurston automaton works in the following way. Suppose we have two non-repeating braids a and b , and we want to rewrite the word ab . The braid a looks for a new crossing of the braid b , and if the braid b allows to take this crossing (*i.e.*, if there is a presentation $b = b'b''$ such that b' is a braid which exactly contains the needed crossing for the braid a : ab' is still non-repeating braid), then the braid a takes this crossing. So, the operation “add the needed crossing” from the braid b to the braid a can be interpreted as a “head” function of the braid ab , denoted by $H(ab)$, *i.e.*, we can say that, the braid a is hungry and greedy for new crossing every time. Then the rewriting procedure of a word ab can be written as follows $ab = H(ab)T(ab)$, where a function T is also called the tail function [10, p. 294].

The natural generalization of these ideas to some other monoids and categories is called now Garside Theory which has been developed by P. Dehornoy, F. Digne, E. Godele, D. Krammer, and J. Michel [10]. This theory deals with left-cancellative categories (as a special case, one object, this includes the case of monoids). We refer the reader to book [10] for complete historical background and some details.

If we look at the standard presentation (=the Coxeter presentation) of the symmetric group \mathfrak{S}_n , which is generated by s_1, \dots, s_{n-1} with relations

- (1) $s_i^2 = 1$, for all $1 \leq i \leq n-1$,
- (2) $s_i s_j = s_j s_i$ for all $1 \leq i, j \leq n-1$ such that $|i-j| > 1$,
- (3) $s_{i+1} s_i s_{i+1} = s_i s_{i+1} s_i$ for all $1 \leq i \leq n-2$,

and at the Artin's presentation of the braid group B_n , which is generated by $\sigma_1, \dots, \sigma_{n-1}$ with relations

- (1) $\sigma_i \sigma_j = \sigma_j \sigma_i$ for all $1 \leq i, j \leq n-1$ such that $|i-j| > 1$,
- (2) $\sigma_{i+1} \sigma_i \sigma_{i+1} = \sigma_i \sigma_{i+1} \sigma_i$ for all $1 \leq i \leq n-2$,

we can see a similarity between these presentations. One obvious invariant of an isotopy of a braid is the permutation it induces on the order of the strands: given a braid b , the strands define a map $P(b)$ from the top set of endpoints to the bottom set of endpoints, which we interpret as a permutation of $\{1, \dots, n\}$. In this way we get a homomorphism (epimorphism) $P_n : B_n \rightarrow \mathfrak{S}_n$, $\sigma_i \mapsto s_i$ for $1 \leq i \leq n-1$. The inverse map (=a section) can be constructed by the Garside's result, from which follows that the braid group can be presented as a monoid which is generated by a set of divisors (left and right) of the braid Δ_n (=the Garside braid which can be described physically by rotating the n strands together 180° clockwise) and by the element Δ_n^{-1} . For any permutation $\pi \in \mathfrak{S}_n$, we correspond the set $R_\pi := \{(i, j) : i < j \text{ \& } \pi(i) > \pi(j)\}$, thus we obtain Adjan–Thurston's generators (or divisors of Δ_n).

However, this idea can be done via the following interesting way which has been suggested by L.A. Bokut' (see [5, 3.1.4]). Using the Gröbner–Shirshov normal form NF for the elements of \mathfrak{S}_n (see [7, 8]), we consider a monoid B'_n generated by the following set of generators $\cup_{\pi \in \mathfrak{S}_n} \{E_{\text{NF}(\pi)}\}$ with relations $E_{\text{NF}(\pi)} E_{\text{NF}(\tau)} = E_{\text{NF}(\pi\tau)}$ whenever $\ell(\text{NF}(\pi\tau)) = \ell(\text{NF}(\pi)) + \ell(\text{NF}(\tau))$, here $\ell(u)$ means the length of a word u . It is easy to show that the monoid B'_n is isomorphic to the positive braid monoid B_n^+ . Next, as it was shown in [5, 3.1.4] the elements $E_{\text{NF}(\pi)}$ are exactly the Adjan–Thurston generators. Further it was also shown that a set of polynomials $S = \{ab - H(ab)T(ab)\}$, where a, b run all over the simple braids, is a Gröbner–Shirshov basis relative to some monomial order, and then, by the Composition–Diamond lemma, the corresponding Gröbner–Shirshov normal form is exactly the greedy normal form.

In [10, VI] the concept of a Garside germ was introduced. Further in the book this concept was applied to Garside families etc. In particular, using this concept, in [10, A, VI, Example 2.72] it was shown how to obtain the braid monoid (and greedy normal form) via the symmetric group. This approach is very similar to Bokut's approach. In particular the elements of $\pi, \tau \in \mathfrak{S}_n$ are called *tight* if $\ell(\pi\tau) = \ell(\pi) + \ell(\tau)$ (in a sense described there).

In this paper, using the concept of germ in the sense of [10, A, VI], we generalize L.A. Bokut' ideas and W. Thurston approach for braid groups to consider Garsdie theory via a Composition–Diamond lemma. We consider two categories, say, \mathcal{C} and \mathcal{A} with a surjective functor $P : \mathcal{C} \rightarrow \mathcal{A}$. Assume further that \mathcal{A} can be presented as follows $\mathcal{A} = \text{Cat}\langle \Gamma \mid R \rangle$ (=a presentation of the \mathcal{A}). By the Composition–Diamond lemma for categories (see [3, 6]) we can calculate a normal form NF (=a Gröbner–Shirshov normal form) and hence a \mathbb{K} -basis \mathfrak{B} for the algebra $\mathbb{K}\mathcal{A}$ (= a category algebra, see Definition 2.1). We then consider a section $E : \mathcal{A} \rightarrow \mathcal{C}$ for the functor P and we wonder know whether $E(a)E(b) = E(\text{NF}(ab))$ for $a, b \in \mathfrak{B}$. This leads us to the concept of a germ (see Construction 3.1) in the sense of [10, A, VI] and we take an interest on whether this germ, denoted by $\Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$, is a Garside germ. The main result of this paper is Theorem 3.12 answers whether such germs are Garside germs in terms of the confluence (up to a congruence) of the corresponding reduction system $\overline{S}_\tau(\mathfrak{B})$ (see Construction 3.5).

For these reasons we need a new version (see Theorem 2.9) of the classical Composition–Diamond lemma for categories (or for monoids). We consider a rewriting system S on a set

of words (=morphisms of the corresponding free category) with a congruence which satisfies some suitable conditions (S -admissibility, see Definition 2.3) and then we require that this system of reductions is to be confluent up to this congruence. This allows us to deal with the fact that an element of a left-cancellative category with a Garside family S , in general, may have more than one S -normal decompositions. On the other hand, according to [10, A, III, Proposition 15] (see Proposition 1.10 in this paper), any two S -normal decomposition of an element are deformations by invertible elements of one another. This is why we need a “deformed” version of a Composition–Diamond lemma.

In particular (see 3.1), when $\mathcal{C} = \mathcal{A} = \text{Cat}\langle\Gamma|R\rangle$, and P is assumed to be an identity functor $\mathbf{id}_{\mathcal{C}}$, a criteria for a germ $\Upsilon(\mathcal{A}, \mathfrak{B}) := \Upsilon_E(\mathcal{C}, \mathcal{C}, \mathbf{id}_{\mathcal{C}}, \mathfrak{B})$ to be a Garside germ is equivalent to a choosing a suitable subset of a set \mathfrak{B} of basic elements of \mathcal{C} (= irreducible elements). In other words to calculate a Garside family of a left-cancellative category \mathcal{C} we first of all have to calculate a Gröbner–Shirshov basis for the ideal $I(R)$, this gives a basis $\text{Irr}(R)$, and then we chose a subset of of this basis to make a Garside family. This can be done by considering the corresponding system of reductions (see Construction 3.5) and then we have to check that all ambiguity are resolvable. In the case when a category \mathcal{C} has nontrivial invertible elements we have to require that all such ambiguities are to be resolvable up to congruence \approx , where \approx is a deformation of elements of \mathcal{C} by invertible elements (see Definition 1.9). We demonstrate how this works for the Klein bottle monoid (Example 3.15).

1. BASIC NOTATIONS OF THE GARSIDE THEORY

In this section we recall the basic concept of the Garside theory. We refer to [10] (see also the survey [9]) for details.

We deal with categories and correspondence terminology, let us remind some basic definitions and concepts.

Definition 1.1. A *precategory* \mathcal{P} is a pair $(\text{Ob}(\mathcal{P}), \text{Hom}(\mathcal{P}))$ with two maps $s, t : \text{Hom}(\mathcal{P}) \rightarrow \text{Ob}(\mathcal{P})$ which are called *source* and *target*, respectively. The elements of $\text{Ob}(\mathcal{P})$ are called the objects, those of $\text{Hom}(\mathcal{P})$ are called the elements or morphisms.

By definition, a category is a precategory, plus composition map that obeys certain rules. Since for every object x there is a morphism $1_x : x \rightarrow x$, we can consider only morphisms. Identity elements 1_x are also called *trivial*, and the collection of all trivial elements in \mathcal{C} is denoted by $\mathbf{1}_{\mathcal{C}}$.

Assume that \mathcal{P} is a precategory. For $p \geq 1$ and $x, y \in \mathcal{P}$, an \mathcal{P} -*path* of length p with source x and target y is a finite sequence g_1, \dots, g_p of elements of \mathcal{P} such that the source of g_1 is x , the target of g_p is y , and $g_1 \cdots g_p$ is defined. The family of all \mathcal{P} -paths of length p is denoted by $\mathcal{P}^{[p]}$.

Definition 1.2 ([13, II, 7,8]). Let $\Gamma = (V, E)$ be an oriented graph. We say that a *graph* $\Gamma = (V, E)$ *generates a category* \mathcal{C} if $\text{Ob}(\mathcal{C}) = V$, and any morphism of \mathcal{C} will be the strings of composable paths of Γ , so that a morphism of $\alpha : b \rightarrow a$ may be pictured as a path from b to a , consisting of successive edges, say e_1, \dots, e_n , of Γ , we then say that the α has length n , and we write $\ell(\alpha) = n$. This category \mathcal{C} will be written $\text{Cat}\langle\Gamma\rangle$ and called the *free category* generated by the graph Γ .

Example 1.3. Let Γ have a single vertex, say, v , and $n \geq 1$ edges e_1, \dots, e_n . Then $\text{Cat}\langle\Gamma\rangle$ has the following semigroup presentation (= as a presentation of a semigroup)

$$\text{Cat}\langle\Gamma\rangle = \text{Smg}\langle 1_v, e_1, \dots, e_n \mid 1_v^2 = 1_v, 1_v e_1 = e_1 1_v = e_1, \dots, 1_v e_n = e_n 1_v = e_n \rangle,$$

it follows that we get the free monoid generated by e_1, \dots, e_n and where 1_v is its identity element (=empty word).

Definition 1.4. Let \mathcal{C} be a category. A function \mathcal{R} which assigns to each pair of objects a, b of \mathcal{C} a binary relation $\mathcal{R}_{a,b}$ on the set $\mathcal{C}(a, b)$ (=all morphism from a to b) is called a *congruence* on $\text{Cat}\langle\Gamma\rangle$ i.e.,

- (1) for each pair x, y of objects, $\mathcal{R}_{a,b}$ is a reflexive, symmetric, and transitive relation on $\text{Cat}\langle\Gamma\rangle(x, y)$.
- (2) if $f, f' : x \rightarrow y$ have $f\mathcal{R}_{x,y}f'$, then for all $g : x' \rightarrow x$ and all $h : y \rightarrow y'$ one has $(hfg)\mathcal{R}_{x',y'}(hf'g)$.

In case $\text{Cat}\langle\Gamma\rangle$ is the free category generated by a graph Γ we shall call $\text{Cat}\langle\Gamma|\mathcal{R}\rangle := \text{Cat}\langle\Gamma\rangle/\mathcal{R}$ the category with generators Γ and relations \mathcal{R} .

□

As a special case (one object), this includes the case of a monoid given by generators and relations.

Throughout this note we frequently use the following notations; $\text{Cat}\langle\Gamma|\mathcal{R}\rangle$ (*resp.* $\text{Smg}\langle\Gamma|\mathcal{R}\rangle$, *resp.* $\text{Gr}\langle\Gamma|\mathcal{R}\rangle$) means a category (*resp.* semigroup, *resp.* group) generated by a graph (*resp.* a set) Γ and relations \mathcal{R} .

Let \mathcal{C} be a category. Every subset $\mathcal{S} \subseteq \mathcal{C}$ is called subfamily, i.e., \mathcal{S} is the precategory made of \mathcal{S} together with the restriction of the source and target maps to \mathcal{S} .

Definition 1.5. [10, A, III, Definitions 1.1, 1.17]

A category \mathcal{C} is called *left-cancellative* if every relation $fg = fh$ with $f, g, h \in \mathcal{C}$ implies $g = h$.

A subfamily \mathcal{S} of a left-cancellative category \mathcal{C} is to be *closed under right-divisor* if every right-divisor of an element \mathcal{S} is an element of \mathcal{S} .

A length-two \mathcal{C} -path $b \cdot c \in \mathcal{C}^{[2]}$ is called \mathcal{S} -greedy if each relation $s \leq abc$ with s in \mathcal{S} implies $s \leq ab$.

A path $a_1 \cdots a_n$ is called \mathcal{S} -greedy if $a_i \cdot a_{i+1}$ is \mathcal{S} -greedy for each $i < n$.

The family of all invertible elements in a category \mathcal{C} is denoted by \mathcal{C}^\times . Let $\mathcal{S} \subseteq \mathcal{C}$ be a family in \mathcal{C} , we set

$$\mathcal{S}^\# := \mathcal{S}\mathcal{C}^\times \cup \mathcal{C}^\times,$$

here $\mathcal{S}\mathcal{C}^\times$ consists of all elements $f \in \mathcal{C}$ of the form $f = sc$ where $s \in \mathcal{S}$ and $c \in \mathcal{C}^\times$.

A path is \mathcal{S} -normal if it is \mathcal{S} -greedy and its entries lie in $\mathcal{S}^\#$.

A procedure of finding \mathcal{S} -normal form is called *Garside normalization*.

Example 1.6. Let $W(X)$ be a free monoid generated by nonempty set X . Let S be the family of all squarefree words in $W(X)$, i.e., the words that admit no factors of the form x^2 . It is obviously that every word $w \in W(X)$ admits a longest prefix that is squarefree.

For instance, let $X = \{a, b, c\}$, and take $w = ab^2cab^2babca$, then its S -normalization is

$$w = ab \cdot bcac \cdot cbabca.$$

Example 1.7 (Free Abelian Monoids [10, A, I.1.1]). Take $n \geq 1$, and consider the free abelian monoid \mathbb{N}^n . It is clear that any its element g can be viewed as a map $g : \{1, \dots, n\} \rightarrow \mathbb{N}$. Denote by $g(k)$ the k th entry of g .

For $g, g' \in \mathbb{N}^n$, we define $g \cdot g'(k) := g(k) + g'(k)$ for each k .

Set Δ_n by $\Delta_n(k) = 1$ for every k , and put $N_n := \{\eta \in \mathbb{N}^n \mid \eta(k) \in \{0, 1\} \text{ for any } k\}$, and, finally, for $f, g \in \mathbb{N}^n$, say that $f \leq g$ is true if $f(i) \leq g(i)$ holds for every $i \in \{1, \dots, n\}$.

Proposition 1.8 ([10, A, I, Proposition 1.1]). *Every element of \mathbb{N}^n admits a unique decomposition of the form $\Delta_n^d \eta_1 \cdots \eta_p$ with $d \in \mathbb{N}$ and $\eta_1, \dots, \eta_p \in N_n$ satisfying $\eta_1 \neq \Delta_n$, $\eta_p \neq 1$, and, for every $i < p$, and for every $g \in \mathbb{N}^n \setminus \{1\}$ we have $g \leq \eta_{i+1}$ implies $\eta_i g \not\leq \Delta_n$.*

For instance, let $n = 3$, and take $f = (5, 4, 3)$, then its N_n -normalization is

$$\begin{aligned} (5, 4, 3) &= (3, 3, 3) + (2, 1, 0) \\ &= (3, 3, 3) + (1, 1, 0) + (1, 0, 0), \end{aligned}$$

thus we obtain $f = \Delta_3^3 \cdot \eta_1 \cdot \eta_2$

Definition 1.9 ([10, A, III, Definition 1.20]). (See Fig.1) Assume that \mathcal{C} is a left-cancellative category. A \mathcal{C} -path $a = a_1 \cdots a_n$ is said to be a *deformation by invertible elements* or \mathcal{C}^\times -*deformation*, of another \mathcal{C} -path $b = b_1 \cdots b_m$, we then write $a \approx b$, if there exist $\epsilon_0, \dots, \epsilon_\ell \in \mathcal{C}^\times$, $\ell = \max(n, m)$, such that ϵ_0 and ϵ_ℓ are identity elements and $\epsilon_{i-1}b_i = a_i\epsilon_i$ holds for $1 \leq i \leq \ell$, where, for $n \neq m$, the shorter path is expanded by identity elements.




FIGURE 1. Deformation by invertible elements: invertible elements connect the corresponding entries; if one path is shorter (here we are in the case $n < m$), it is extended by identity elements.

Proposition 1.10 ([10, A, III, Proposition 1.25]). *If S is a subfamily of a left-cancellative category \mathcal{C} , any two S -normal decomposition of an element of \mathcal{C} are \mathcal{C}^\times -deformations of one another.*


Example 1.11. Let us consider a symmetric group \mathfrak{S}_n , $n \geq 3$, in Coxeter presentation:

$$\mathfrak{S}_n = \text{Smg}\langle s_1, \dots, s_{n-1} \mid \mathcal{R} \rangle,$$


where \mathcal{R} is the following set of relations:

- (1) $s_i^2 = 1$, for all $1 \leq i \leq n - 1$,
- (2) $s_i s_j = s_j s_i$ for all $1 \leq i, j \leq n - 1$ such that $|i - j| > 1$,
- (3) $s_{i+1} s_i s_{i+1} = s_i s_{i+1} s_i$ for all $1 \leq i \leq n - 2$.

We have $s_i s_j \approx s_j s_i$ for any $1 \leq i, j \leq n - 1$, $|i - j| > 1$ because the diagram



is commutative, and $s_k s_{k+1} s_k \approx s_{k+1} s_k s_{k+1}$ for all $1 \leq k \leq n - 1$ because the diagram



is commutative. □

Definition 1.12. [10, A, III, Definition 1.31] A subfamily \mathcal{S} of a left-cancellative category \mathcal{C} is called a *Garside family* in \mathcal{C} if every element of \mathcal{C} admits at least one \mathcal{S} -normal decomposition.

A subfamily \mathcal{S} of a category \mathcal{C} is said to be *solid* in \mathcal{C} if \mathcal{S} includes $\mathbf{1}_{\mathcal{C}}$ and it is closed under right-divisor.

Definition 1.13. [10, A, VI, Definitions 1.3, 1.23]

A germ is a triple $(\Upsilon, \bullet, \mathbf{1}_{\Upsilon})$ where Υ is a precategory, $\mathbf{1}_{\Upsilon}$ is a subfamily of an elements x with source and target x for each object x , and the partial map \bullet that satisfies

- (1) if $\alpha \bullet \beta$ is defined, the source of $\alpha \bullet \beta$ is the source of α , and its target is the target of β ;
- (2) the relation $1_x \bullet \alpha = \alpha = \alpha \bullet 1_y$ hold for each α in $\Upsilon(x, y)$;
- (3) if $\alpha \bullet \beta$ and $\beta \bullet \gamma$ are defined, then $(\alpha \bullet \beta) \bullet \gamma$ is defined if and only if $\alpha \bullet (\beta \bullet \gamma)$ is, in which case they are equal.

The germ is said to be left-associative (*resp.* right-associative) if

- (4) $(\alpha \bullet \beta) \bullet \gamma$ is defined, then so is $\beta \bullet \gamma$;
- (5) (*resp.* $\alpha \bullet (\beta \bullet \gamma)$ is defined, then so is $\alpha \bullet \beta$).

If $(\Upsilon, \bullet, \mathbf{1}_{\Upsilon})$ is a germ, we denote by $\text{Cat}\langle \Upsilon, \bullet, \mathbf{1}_{\Upsilon} \rangle$ (or shortly $\text{Cat}\langle \Upsilon \rangle$) the category $\text{Cat}\langle \Upsilon \mid \mathcal{R}_{\bullet} \rangle$, where \mathcal{R}_{\bullet} is the family of all relations $\alpha \bullet \beta = \alpha\beta$ with $\alpha, \beta \in \Upsilon$ and $\alpha \bullet \beta$ is defined.

A germ Υ is called a *Garside germ* if it is a Garside family for the category $\text{Cat}\langle \Upsilon \rangle$.

Example 1.14. Let us consider the positive braid monoid in three strands

$$B_3^+ = \text{Smg}\langle \sigma_1, \sigma_2, \mid \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \rangle,$$

and let Υ be the following family in B_3^+ , $\Upsilon = \{1, \sigma_1, \sigma_2, \sigma_1\sigma_2, \sigma_2\sigma_1, \Delta_3\}$, where $\Delta_3 := \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2$.

Define the partial map $\bullet : \Upsilon^{[2]} \rightarrow \Upsilon$ as follows

\bullet	1	σ_1	σ_2	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	Δ_3
1	1	σ_1	σ_2	$\sigma_1\sigma_2$	$\sigma_2\sigma_1$	Δ_3
σ_1	σ_1		$\sigma_1\sigma_2$		Δ_3	
σ_2	σ_2	$\sigma_2\sigma_1$		Δ_3		
$\sigma_1\sigma_2$	$\sigma_1\sigma_2$	Δ_3				
$\sigma_2\sigma_1$	$\sigma_2\sigma_1$		Δ_3			
Δ_3	Δ_3					

By the straightforward verification, it is easy to see that (Υ, \bullet) is a left-cancellative and left-associative germ.

Definition 1.15 ([10, A, VI, Definitions 2.1, 2.27]). For Υ a germ and $\alpha, \beta \in \Upsilon$, we put

$$\mathcal{I}_{\Upsilon}(\alpha, \beta) := \{\delta \in \Upsilon \mid \exists \gamma \in \Upsilon (\delta = \alpha \bullet \gamma \text{ and } \gamma \preceq_{\Upsilon} \beta)\},$$

$$\mathcal{J}_{\Upsilon}(\alpha, \beta) := \{\gamma \in \Upsilon \mid \beta \bullet \gamma \in \Upsilon \text{ \& } \gamma \preceq_{\Upsilon} \beta\}.$$

A map $\mathfrak{I} : \Upsilon^{[2]} \rightarrow \Upsilon$ is called an \mathcal{I} -function (*resp.* a \mathcal{J} -function) if, for every $\alpha \cdot \beta \in \Upsilon^{[2]}$, the value at $\alpha \cdot \beta$ lies in $\mathcal{I}_{\Upsilon}(\alpha, \beta)$ (*resp.* in $\mathcal{J}_{\Upsilon}(\alpha, \beta)$).

An \mathcal{I} -function $\mathfrak{I} : \Upsilon^{[2]} \rightarrow \Upsilon$ is called greatest \mathcal{I} -function if and only if for every $\alpha \cdot \beta \in \Upsilon^{[2]}$, we have $\mathfrak{I}(\alpha, \beta) = \alpha \bullet \gamma$ for some $\gamma \in \Upsilon$ satisfying $\gamma \preceq_{\Upsilon} \beta$, and $\delta \preceq_{\Upsilon} \mathfrak{I}(\alpha, \beta)$ holds for every $\delta \in \mathcal{I}_{\Upsilon}(\alpha, \beta)$.

Proposition 1.16 ([10, A, VI, Proposition 2.28]). A germ Υ is a Garside germ if and only if Υ is left-associative, left-cancellative, and admits a greatest \mathcal{I} -function.

Definition 1.17 (\mathcal{I} -law). [10, A, VI, Definition 2.5] If Υ is a germ and \mathfrak{I} is a map from $\Upsilon^{[2]}$ to Υ , we say that \mathfrak{I} obeys \mathcal{I} -law if, for every $\alpha, \beta, \gamma \in \Upsilon$ with $\alpha \bullet \beta$ defined, we have

$$\mathfrak{I}(\alpha, \mathfrak{I}(\beta, \gamma)) =^\times \mathfrak{I}(\alpha \bullet \beta, \gamma). \quad (1.1)$$

If the counterpart of this equality with $=$ replacing $=^\times$, we say that I obeys the sharp \mathcal{I} -law.

Theorem 1.18 ([10, A, VI, Proposition 2.8]). *A germ Υ is a Garside germ if and only if it is left-associative, left-cancellative, and admits an \mathcal{I} -function obeying the sharp \mathcal{I} -law.*

Proposition 1.19 ([10, A, VI, Lemma 2.4]). *If Υ is a Garside germ, the following are equivalent for every $\alpha, \beta \in \Upsilon$:*

- (1) *The path $\alpha \cdot \beta$ is Υ -normal.*
- (2) *The element α is $<_\Upsilon$ -maximal in $\mathcal{I}_\Upsilon(\alpha, \beta)$.*
- (3) *Every element of $\mathcal{I}_\Upsilon(\alpha, \beta)$ is invertible in Υ .*

Proposition 1.20. *Let \mathcal{S} be a solid family generating a left-cancellative category \mathcal{C} , then \mathcal{S} equipped with the induced partial product is a germ $\Upsilon(\mathcal{S})$. If the $\Upsilon(\mathcal{S})$ is a Garside germ then \mathcal{C} is isomorphic to $\text{Cat}(\Upsilon(\mathcal{S}))$.*

Proof. To prove this we need the following results

Lemma 1.21 ([10, A, VI, Proposition 1.1]). *If \mathcal{S} is a solid Garside family in a left-cancellative category \mathcal{C} , then \mathcal{S} equipped with the induced partial product is a germ $\Upsilon(\mathcal{S})$ and \mathcal{C} is isomorphic to $\text{Cat}(\Upsilon(\mathcal{S}))$,*

Lemma 1.22 ([10, A, IV, Proposition 2.24]). *A solid generating subfamily \mathcal{S} of a left-cancellative category \mathcal{C} is a Garside family if and only if there exists $\mathfrak{I} : \mathcal{S}^{[2]} \rightarrow \mathcal{S}$ satisfying $\alpha \leq \mathfrak{I}(\alpha, \beta) \leq \alpha \cdot \beta$ for all α, β and $\mathfrak{I}(\alpha, \mathfrak{I}(\beta, \gamma)) = \mathfrak{I}(\alpha\beta, \gamma)$ for every $\alpha, \beta, \gamma \in \mathcal{S}$ satisfying $\alpha \cdot \beta \in \mathcal{S}$.*

Since $\Upsilon(\mathcal{S})$ is assumed to be a Garside germ then there is a \mathcal{I} -greatest function \mathfrak{I} . It is clear that it satisfies the statement above (see also proof of [10, A, IV, Lemma 2.23, Proposition 2.24] and [10, IV, Proposition 2.8]). Hence, by [10, A, IV, Proposition 2.24], \mathcal{S} is a Garside family in \mathcal{C} . Therefore, by Proposition [10, A, VI, Proposition 1.1], the statement follows. \square

Theorem 1.23 (germ from Garside, [10, A, VI, Proposition 1.1]). *If \mathcal{S} is a solid Garside family in a left-cancellative category \mathcal{C} , then \mathcal{S} equipped with the induced partial product is a germ $\Upsilon(\mathcal{S})$ and \mathcal{C} is isomorphic to $\text{Cat}(\Upsilon(\mathcal{S}))$.*

Remark 1.24. *If a solid family \mathcal{S} in a left-cancellative category \mathcal{C} generates \mathcal{C} and $\Upsilon(\mathcal{S})$ is a Garside germ then $\mathcal{C} \cong \text{Cat}(\Upsilon(\mathcal{S}))$*

2. A COMPOSITION–DIAMOND LEMMA FOR CATEGORIES

In this section we present a “deformed” version of a Composition–Diamond lemma for categories. We essentially follows [3]. A sketch of a Composition–Diamond lemma for categories has been appeared in [3, 9.3] and it has been developed in detail in [6] (see also [5, 2.4]).

Definition 2.1. Let \mathbb{K} be a commutative ring and \mathcal{C} a category. Define the *category algebra* $\mathbb{K}\mathcal{C}$ to be the the free \mathbb{K} -module with the morphisms of the category \mathcal{C} as a basis. The product of morphisms a and b as elements of $\mathbb{K}\mathcal{C}$ is defined to be

$$a \cdot b := \begin{cases} ab & \text{if } a \text{ and } b \text{ can be composed,} \\ 0 & \text{otherwise,} \end{cases}$$

and this product is extended to the whole of $\mathbb{K}\mathcal{C}$ using bilinearity of multiplication.

Given a graph Γ , consider the corresponding free category $\text{Cat}\langle\Gamma\rangle$. Let S be a set of pairs of the form $\sigma = (W_\sigma, \omega_\sigma)$, where $W_\sigma, \omega_\sigma \in \text{Cat}\langle\Gamma\rangle$. For any $\sigma \in S$ and $A, B \in \text{Cat}\langle\Gamma\rangle$, let $r_{A\sigma B} : \text{Cat}\langle\Gamma\rangle \rightarrow \text{Cat}\langle\Gamma\rangle$ be a functor defined as follows:

$$r_{A\sigma B}(W) := \begin{cases} A\omega_\sigma B & \text{if } W = W_\sigma \\ W & \text{if } W \neq W_\sigma. \end{cases}$$

We call the given set S a *reduction system*, and the maps $r_{A\sigma B}$ *reductions*, if $r_{A\sigma B}(W) = W$, we say a reduction $r_{A\sigma B}$ acts *trivially* on W , and we shall call W *irreducible under S* if every reduction is trivial on W , and we say that a word W is *reducible* in otherwise.

We shall frequently write $r_{A\sigma B} : W \rightarrow W'$ if $r_{A\sigma B}(W) = W'$.

Denote by $\text{Irr}(S)$ the subset of all irreducible elements of $\text{Cat}\langle\Gamma\rangle$ under all reductions from the set S . A finite sequence of reductions r_1, \dots, r_n will be said to be *final* on $W \in \text{Cat}\langle\Gamma\rangle$ if $r_n \cdots r_1(W) \in \text{Irr}(S)$.

An element W of $\text{Cat}\langle\Gamma\rangle$ will be called *reduction-finite* if for every infinite sequence r_1, r_2, \dots , of reductions, r_i acts trivially on $r_{i-1} \cdots r_1(W)$ for all sufficiently large i .

Definition 2.2. A 5-tuple (σ, τ, A, B, C) with $\sigma, \tau \in S$, and $A, B, C \in \text{Cat}\langle\Gamma\rangle \setminus \{\mathbf{1}\}$, is called *overlap (resp. inclusion) ambiguity of S* if $W_\sigma = AB$, $W_\tau = BC$ (resp. if $\sigma \neq \tau$ and $W_\sigma = B$, $W_\tau = ABC$).

Definition 2.3. Let \approx be a congruence on a free category $\text{Cat}\langle\Gamma\rangle$ and S a set of reductions. We call a congruence \approx *S -admissible* if the following hold:

- (1) if $W \notin \text{Irr}(S)$ then $W' \notin \text{Irr}(S)$ for any $W' \approx W$,
- (2) if $\omega \in \text{Irr}(S)$ then $\omega' \in \text{Irr}(S)$ for any $\omega' \approx \omega$.

An element $W \in \text{Cat}\langle\Gamma\rangle$ is called *reduction-unique up to \approx under S* if it is reduction-finite, and if its images under all final sequences reductions and under using an S -admissible congruence are the same. This common value will be denoted by $r_S(W)$, and we also write $r : W \rightarrow r_S(W)$ if r is a composition of reductions of S only and we write $r : W \rightsquigarrow r_S(W)$ in otherwise.

Remark 2.4. Note that if all elements of $\text{Cat}\langle\Gamma\rangle$ are reduction unique up to \approx then the following condition holds; if $W \rightsquigarrow U$, $W' \rightsquigarrow U'$ then from $W \approx W'$ it follows that $U \approx U'$.

Definition 2.5. By a *\approx -admissible categorical partial preorder* on a free category $\text{Cat}\langle\Gamma\rangle$ we shall mean a partial preorder \lesssim such that for $A, B, B', C \in \text{Cat}\langle\Gamma\rangle$ the following hold:

- (1) if $B < B'$ then $ABC < AB'C$,
- (2) whenever $B < B'$, $B' < B$ it must have that $B \approx B'$,
- (3) if $B < A$ and $B \approx C$ then $C < A$,
- (4) if $A < B$, $B \approx C$ then $A < C$.

Next, we say that a \approx -admissible categorical partial preorder is *compatible* with S if for all $(W_\sigma, \omega_\sigma) = \sigma \in S$, $\omega_\sigma < W_\sigma$, $W_\sigma \not< \omega_\sigma$.

In the case with a single object, which includes the case of a free monoid, we also call a \approx -admissible categorical partial preorder as a *\approx -admissible monomial partial preorder*.

For a given set of reductions $S = \cup_{\sigma \in S} \{(W_\sigma, \omega_\sigma)\}$ on a free category $\text{Cat}\langle\Gamma\rangle$ we consider the following set of polynomials $\cup_{\sigma \in S} \{W_\sigma - \omega_\sigma\}$, and we say that (S) is *the corresponding set of polynomials for a given set of reductions S* .

Next, let \approx be an S -admissible congruence on $\text{Cat}\langle\Gamma\rangle$. Set $\widetilde{(S)} := (S) \cup (S')$, where $(S') = \cup_{\sigma \in S} \{W_\sigma - W'_\sigma \mid W_\sigma \approx W'_\sigma, W_\sigma, W'_\sigma \notin \text{Irr}(S)\}$ and consider a two-sided ideal, denoted by $I_\approx(S)$, generated by $\widetilde{(S)}$.

Remark 2.6. Note that an ideal $I_{\approx}(S)$ cannot be described as a two-sided ideal generated by the following set of polynomials $\cup_{\sigma \in S} \{W_{\sigma} - \omega'_{\sigma} \mid W_{\sigma} \rightsquigarrow \omega'_{\sigma}\}$. Indeed, let $r_1 : W_{\sigma} \rightarrow \omega_{\sigma}$, and $r_s : W'_{\sigma} \rightarrow \omega'_{\sigma}$ be reductions of S . Then we have $r' : W_{\sigma} \rightsquigarrow \omega'_{\sigma}$ but $r' \notin S$, $W_{\sigma} \notin \text{Irr}(S)$, and $\omega_{\sigma} \in \text{Irr}(S)$, hence $W_{\sigma} - \omega'_{\sigma} \notin I_{\approx}(S)$.

Definition 2.7. An overlap ambiguity (σ, τ, A, B, C) (resp. inclusion ambiguity) is called *resolvable up to \approx* if there exist compositions of reductions $r : \omega_{\sigma}C \rightsquigarrow r(\omega_{\sigma}C)$ and $r' : A\omega_{\tau} \rightsquigarrow r'(A\omega_{\tau})$ (resp. $r : \omega_{\sigma} \rightsquigarrow r(\omega_{\sigma})$, $r' : A\omega_{\tau}C \rightsquigarrow r'(A\omega_{\tau}C)$) such that $r(\omega_{\sigma}C) \approx r'(A\omega_{\tau})$ (resp. $r(\omega_{\sigma}) \approx r'(A\omega_{\tau}C)$).

If all ambiguities of S are resolvable up to \approx under S we then say that S is *confluent up to \approx* .

Let \lesssim be a categorical partial preorder on a free category $\text{Cat}\langle\Gamma\rangle$ compatible with a reduction system S . Consider $f_{\sigma} = W_{\sigma} - \omega_{\sigma}$, $f_{\tau} = W_{\tau} - \omega_{\tau} \in (S)$. There are two kinds of compositions:

- (1) if $ABC = W_{\sigma}C = AW_{\tau}$ with $\ell(W_{\sigma}) + \ell(W_{\tau}) > \ell(ABC)$, then the polynomial $(f_{\sigma}, f_{\tau})_{ABC} := f_{\sigma}C - Af_{\tau}$ is called the *intersection composition* of f_{σ} and f_{τ} with respect to ABC ,
- (2) if $ABC = W_{\sigma} = AW_{\tau}C$, then the polynomial $(f_{\sigma}, f_{\tau})_{ABC} := f_{\sigma} - Af_{\tau}C$ is called the *inclusion composition* of f_{σ} and f_{τ} with respect to ABC .

Consider $g \in \mathbb{K}\text{Cat}\langle\Gamma\rangle$. Set $g \approx 0$ if g can be presented as follows $g = \sum_{i=1}^n (U_i - V_i)$, where for any $1 \leq i \leq n$, $U_i, V_i \in \text{Cat}\langle\Gamma\rangle$ and all $U_i \approx V_i$. We say $g \in \mathbb{K}\text{Cat}\langle\Gamma\rangle$ is *trivial modulo $((S), W)$ up to \approx* , denoted by $g \approx 0 \pmod{(S), W}$, here $W \in \text{Cat}\langle\Gamma\rangle$, if g can be presented as follows $g = \sum_{i=1}^n \sum_{\sigma_i \in S} U_i f_{\sigma_i} V_i + g'$, where all $U_i, V_i \in \text{Cat}\langle\Gamma\rangle$, $UW_{\sigma}V < W$, $g' \in \mathbb{K}\text{Cat}\langle\Gamma\rangle$, $g' \approx 0$, and for all its monomials, say G'_j , we have $G'_j < W$.

Definition 2.8. Let S be a set of reductions on a free category $\text{Cat}\langle\Gamma\rangle$, \approx an S -admissible congruence, and \lesssim a \approx -admissible categorical partial preorder compatible with S . The corresponding set of polynomials (S) is called a *\approx -closed Gröbner–Shirshov basis* in $\mathbb{K}\text{Cat}\langle\Gamma\rangle$ with respect to \lesssim if every composition $(f_{\sigma}, f_{\tau})_{ABC}$ of polynomials $f_{\sigma}, f_{\tau} \in (S)$ is trivial modulo $((S), ABC)$ up to \approx .

To stress that a set $X \subseteq \text{Cat}\langle\Gamma\rangle$ is considered as the set with a congruence \approx on $\text{Cat}\langle\Gamma\rangle$ we write (X, \approx) .

Theorem 2.9 (a \approx -closed Composition–Diamond Lemma). *Let \mathbb{K} be a commutative ring with unit. Let $\text{Cat}\langle\Gamma\rangle$ be a free category, S a reduction system, (S) the corresponding set of polynomials, \approx an S -admissible congruence on $\text{Cat}\langle\Gamma\rangle$, and \lesssim a \approx -admissible categorical partial preordering compatible with S having descending chain condition. Then the following conditions are equivalent:*

- (1) *All ambiguities of S are \approx -resolvable.*
- (2) *(S) is a \approx -closed Gröbner–Shirshov basis relative to \lesssim .*
- (3) *All elements of $\mathbb{K}\text{Cat}\langle\Gamma\rangle$ are reduction-unique up to \approx under S .*
- (4) *$(\text{Irr}(S), \approx)$ is a linear \mathbb{K} -basis of the algebra $\mathbb{K}\text{Cat}\langle\Gamma \mid \widetilde{(S)}\rangle = \mathbb{K}\text{Cat}\langle\Gamma\rangle / I_{\approx}(S)$, where $\text{Irr}(S) := \{W \in \text{Cat}\langle\Gamma\rangle \mid W \neq UW_{\sigma}V, \sigma \in S, U, V \in \text{Cat}\langle\Gamma\rangle\}$.*

Proof. We easily see from our general hypothesis, by induction with respect to the partial preordering with descending chain condition \lesssim , that every element of $\mathbb{K}\text{Cat}\langle\Gamma\rangle$, is reduction-finite up to \approx .

If $(S) = \cup_{i \in I} \{W_i - \omega_i\}$ is a \approx -closed Gröbner–Shirshov basis relative to \lesssim it then gives the following system of reduction $S = \cup_{i \in I} \{\sigma_i = (W_i, \omega_i)\}$ is compatible with \lesssim .

(1) \iff (2) Take $f_{\sigma}, f_{\tau} \in (S)$ and let $AW_{\sigma} = W_{\tau}C = ABC$ we then get

$$\begin{aligned} (f_{\sigma}, f_{\tau})_{ABC} &= A(W_{\sigma} - \omega_{\sigma}) - (W_{\tau} - \omega_{\tau})C \\ &= -A\omega_{\sigma} + \omega_{\tau}C \end{aligned}$$

Let $r_S^{(1)} : A\omega_\sigma \rightsquigarrow A_1W_{\sigma_1}C_1$ and $r_S^{(1)'} : \omega_\tau C \rightsquigarrow A'_1W_{\tau_1}C'_1$, we then get

$$\begin{aligned} (f_\sigma, f_\tau)_{ABC} &= -A\omega_\sigma + \omega_\tau C \\ &\approx -A_1(f_{\sigma_1} + \omega_{\sigma_1})C_1 + A'_1(f_{\tau_1} + \omega_{\tau_1})C'_1 \\ &= -A_1f_{\sigma_1}C_1 + A'_1f_{\tau_1}C'_1 - A_1\omega_{\sigma_1}C_1 + A'_1\omega_{\tau_1}C'_1. \end{aligned}$$

Next, setting $r_S^{(2)} : A_1\omega_{\sigma_1}C_1 \rightsquigarrow A_2W_{\sigma_2}C_2$, $r_S^{(2)'} : A'_1\omega_{\tau_1}C'_1 \rightsquigarrow A'_2W_{\tau_2}C'_2$ and so on, after some steps, we get

$$(f_\sigma, f_\tau)_{ABC} \approx -\sum_{i=1}^N A_i f_{\sigma_i} C_i + \sum_{j=1}^M A'_j f_{\tau_j} C'_j - A_N \omega_{\sigma_N} C_N + A'_M \omega_{\tau_M} C'_M.$$

It is easy to see that from Definition 2.5 it follows that $A_i W_{\sigma_i} C_i, A'_j W_{\tau_j} C'_j < ABC$, for any $1 \leq i \leq N, 1 \leq j \leq M$.

If (S) is a \approx -completed Gröbner–Shirshov basis relative to \lesssim then $N, M < \infty$, and $A_N \omega_{\sigma_N} C_N \approx A'_M \omega_{\tau_M} C'_M$. It follows that the ambiguity (σ, τ, AB, BC) is resolvable up to \approx . Similarly one can easily get the converse statement.

(2) \implies (3). It is clear that it is sufficient to prove all elements $W \in \text{Cat}\langle\Gamma\rangle$ reduction-unique up to \approx . We must show that given any two reductions $r_{L\sigma M'}$ and $r_{L'\tau M}$ each acting nontrivially on W we shall have $r_S(r_{L\sigma M'}(W)) \approx r_S(r_{L'\tau M}(W))$. There are three cases, according to the relative locations of the subwords W_σ and W_τ in the element W . We may assume without loss of generality that $\ell(L) \leq \ell(L')$.

Case 1. The subwords W_σ and W_τ , overlap in W , but neither contains the other. Then $W = UABC V$, where (σ, τ, A, B, C) is an overlap ambiguity of S . Let us consider the corresponding composition for the polynomials $f_\sigma, f_\tau \in (S)$, we have

$$(f_\sigma, f_\tau)_{ABC} \approx -\sum_{i=1}^N A_i f_{\sigma_i} C_i + \sum_{j=1}^M A'_j f_{\tau_j} C'_j - A_N \omega_{\sigma_N} C_N + A'_M \omega_{\tau_M} C'_M.$$

Since (S) has been assumed to be \approx -completed Gröbner–Shirshov basis relative to \lesssim we then obtain, $N, M < \infty$, $A_i W_{\sigma_i} C_i, A'_j W_{\tau_j} C'_j < ABC$ for all $1 \leq i \leq N, 1 \leq j \leq M$, and $A_{N+1} \omega_{\sigma_{N+1}} C_{N+1} \approx A'_{M+1} \omega_{\tau_{M+1}} C'_{M+1}$. Next, since \approx is assumed to be congruence on $\text{Cat}\langle\Gamma\rangle$ then $UA_{N+1} \omega_{\sigma_{N+1}} C_{N+1} V \approx UA'_{M+1} \omega_{\tau_{M+1}} C'_{M+1} V$, as required.

Case 2. One of the subwords W_σ, W_τ of W is contained in the other. Thus case is handled like the preceding, using the resolvability up to \approx of inclusion composition.

Case 3. Finally, W_σ, W_τ are disjoint subwords of W , but in this case the statement is obvious.

(3) \iff (4) Any reduction r_S induces the corresponding \mathbb{K} -module homomorphism on $\mathbb{K}\text{Cat}\langle\Gamma\rangle$, denote it also by r_S .

Assuming (3). Take $f \in \mathbb{K}\text{Cat}\langle\Gamma\rangle$, say, $f = \sum_{i=1}^M \kappa_i u_i$, where all $\kappa_i \in \mathbb{K}$, $u_i \in \text{Cat}\langle\Gamma\rangle$, and $u_1 > u_2 > \dots > u_M$. Since all reductions of S are assumed to be unique up to \approx then for any $1 \leq i \leq M$, $u_i \approx A_1^{(i)} W_{\sigma_1}^{(i)} B_1^{(i)}$ for some $A_1^{(i)}, B_1^{(i)} \in \text{Cat}\langle\Gamma\rangle$. Set $\varphi_\sigma := W_\sigma - \omega_\sigma$ for any $\sigma = (W_\sigma, \omega_\sigma) \in S$. We can write $W_\sigma = \varphi_\sigma + \omega_\sigma$, thus, by induction, we get

$$u_i \approx \sum_{k_i=1}^{m_i} A_{k_i}^{(i)} \varphi_{\sigma_{k_i}}^{(i)} B_{k_i}^{(i)} + C_{k_{m_i}}^{(i)},$$

where $C_{k_{m_i}}^{(i)} \in \text{Irr}(S)$, and all $A_{k_i}^{(i)} \text{LT}(\varphi_{\sigma_{k_i}}^{(i)}) B_{k_i}^{(i)}, C_{k_{m_i}}^{(i)} \lesssim u_p$, here $\text{LT}(\varphi)$ denotes a leading term of φ related to \lesssim . It follows that (cf. [5, 2.1, Lemma 2]) for any polynomial $f \in \mathbb{K}\text{Cat}\langle\Gamma\rangle$ we have

$$f \approx \sum_{p=1}^N \alpha_p U_p \varphi_{\sigma_p} U'_p + \sum_{q=1}^{N'} \beta_q V_q$$

where all $\alpha_p, \beta_q \in \mathbb{K}$, $U_p, U'_p \in \text{Cat}\langle\Gamma\rangle$, $V_q \in \text{Irr}(S)$, and $U_p \text{LT}(\varphi_p) U'_p, V_q \approx \text{LT}(f)$.

Next, since \approx is assumed to be S -admissible then $\mathbb{K}\text{Cat}\langle\Gamma\rangle \cong I_{\approx}(S) \oplus (\text{Irr}(S), \approx)$ and the statement (4) follows.

Conversely, assume (4) and suppose $W \in \text{Cat}\langle\Gamma\rangle$ can be reduced to either of $\omega, \omega' \in (\text{Irr}(S), \approx)$. Then $W - \omega, W - \omega' \in I_{\approx}(S)$ and hence $\omega - \omega' \in I_{\approx}(S)$. On the other hand, $\omega - \omega' \in (\text{Irr}(S), \approx)$, it follows that $\omega - \omega' \in (\text{Irr}(S), \approx) \cap I_{\approx}(S) = \{0\}$, because of \approx is assumed to be S -admissible, and proving (3).

Finally, the implication (3) \implies (1) is immediate. This completes the proof. \square

Remark 2.10. So we see that for a given free category $\text{Cat}\langle\Gamma\rangle$ we district two relations on it; reductions and congruence \approx . We do not consider an ambiguity up to \approx , *i.e.*, let S be a set of reductions, then we would define an ambiguity up to \approx as follows, say a 5-tuple (σ, τ, A, B, C) is called overlap (*resp.* inclusion) ambiguity if $W_{\sigma} \approx AB$ and $w_{\tau} \approx BC$ (*resp.* $W_{\sigma} \approx B, W_{\tau} \approx ABC$). But it is easy to see that in this case we just add to a reduction system new reductions of form $W_{\sigma} \rightarrow AB$ and $W_{\tau} \rightarrow BC$. This is why we district this two relations.

Remark 2.11. It is well known that similar to the Buchberger algorithm of completion for commutative polynomials, there is a Shirshov algorithm for noncommutative polynomials. However, in contrast to the case of commutative algebras, the Shirshov algorithm in general does not terminate in a finite number of steps (we cannot use Noetherianity in noncommutative case). In our case we of course can use the same way. If a subset $(S) \subseteq \mathbb{K}\text{Cat}\langle\Gamma\rangle$ is not a \approx -closed Gröbner–Shirshov basis then one can add to (S) all nontrivial up to \approx compositions of polynomials from (S) . Continuing this process repeatedly we finally obtain a \approx -closed Gröbner–Shirshov basis (\widehat{S}) that contains S . As in the case of classical Buchberger–Shirshov algorithm we have to assume that the initial set (S) is recursively enumerable (*e.g.*, finite) and that the underlying ring K is computable. The resulting set (\widehat{S}) is also recursively enumerable but not necessarily finite even if (S) was.

Example 2.12. Let us consider the symmetric group $\mathfrak{S}_n, n \geq 3$, in a Coxeter presentation

$$\mathfrak{S}_n = \text{Smg}\langle s_1, \dots, s_{n-1} \mid \mathcal{R} \rangle,$$

where \mathcal{R} is the following set of relations:

- (1) $s_i^2 = 1$, for all $1 \leq i \leq n - 1$,
- (2) $s_i s_j = s_j s_i$ for all $1 \leq i, j \leq n - 1$ such that $|i - j| > 1$,
- (3) $s_{i+1} s_i s_{i+1} = s_i s_{i+1} s_i$ for all $1 \leq i \leq n - 2$.

We have already remarked that $s_k s_{k+1} s_k \approx s_{k+1} s_k s_{k+1}$ and $s_i s_j \approx s_j s_i$ because the corresponding diagrams

$$\begin{array}{ccccc} \cdot & \xrightarrow{s_k} & \cdot & \xrightarrow{s_{k+1}} & \cdot & \xrightarrow{s_k} & \cdot \\ \parallel & & \downarrow s_k s_{k+1} & & \downarrow s_k s_{k+1} & & \parallel \\ \cdot & \xrightarrow{s_{k+1}} & \cdot & \xrightarrow{s_k} & \cdot & \xrightarrow{s_{k+1}} & \cdot \end{array}$$

and

$$\begin{array}{ccccc} \cdot & \xrightarrow{s_i} & \cdot & \xrightarrow{s_j} & \cdot \\ \parallel & & \downarrow s_i s_j & & \parallel \\ \cdot & \xrightarrow{s_j} & \cdot & \xrightarrow{s_i} & \cdot \end{array}$$

are commutative.

Consider the following set of reductions $S = \bigcup_{i=1}^{n-1} \{(s_i s_i, 1)\}$ on the free monoid $W(S)$ generated by s_1, \dots, s_{n-1} . Thus a word $U \in W(S)$ is reducible if and only if there is $U_1, U_2 \in W(S)$

such that $U = U_1 s_i s_i U_2$ for some $s_i \in \{s_1, \dots, s_{n-1}\}$. It is easy to see that \approx is S -admissible congruence.


Let us consider a partial preoder $<$ on $W(S)$ defined as follows: $U < V$ if and only if $\ell(U) < \ell(V)$, where $\ell(U)$ is a length of a word $U \in W(S)$. We see that if $U \approx U'$ then $\ell(U) = \ell(U')$, and if $r : U \rightarrow V$, for some reduction r , then $\ell(U) > \ell(V)$. Thus, it is clear that $<$ is a \approx -admissible monomial partial preordering on $W(S)$.

Next, by the straightforward computations, it is easy to see that all ambiguities of S are resolvable up to \approx and hence $(S) = \cup_{i=1}^{n-1} \{s_i^2 - 1\}$ is a \approx -closed Gröbner–Shirshov basis with respect to the $<$. Thus, by Theorem 2.9, the corresponding set $(\text{Irr}(S), \approx)$ is a set of squarefree words in s_i , $1 \leq i \leq n - 1$.

□


Remark 2.13. Since “the usual equality”, *i.e.*, $W \approx W'$ if and only if $W = W'$ for all elements of a free category, is, of course, a congruence then $=$ replacing \approx we then get the classical Composition–Diamond Lemma for categories (see [3, 9.3], [6], [5]). In such cases we shall say just a Gröbner–Shirshov basis instead of \approx -closed Gröbner–Shirshov basis.

Example 2.14. Let us consider the following graph Γ shown below




Set $\mathcal{C} = \text{Cat}\langle \Gamma \mid d^2 = ab, c^2 = ba \rangle$, and let R be the ideal $(d^2 - ab, c^2 - ba)$ in $\mathbb{K}\text{Cat}\langle \Gamma \rangle$. Let $a < b < c < d$ and consider the corresponding deg-lex ordering on the free category $\text{Cat}\langle \Gamma \rangle$.

Put $\sigma = (d^2, ab)$, $\rho = (c^2, ba)$. We have the following ambiguities $(\sigma, \sigma, d, d, d)$, (ρ, ρ, c, c, c) . We have




Thus, by the Buchberger–Shirshov algorithm, to calculate a Gröbner–Shirshov basis relative to the \leq we have to add to S these two reductions $\mu_1 = (dab, abd)$, $\mu_2 = (cba, bca)$. We have the following ambiguities $(\sigma, \rho_1, d, d, ab)$, (ρ, μ_2, c, c, ba) .

We get



and



Thus all ambiguities of the reduction system $S' = \{\sigma, \rho, \mu_1, \mu_2\}$ are resolvable, hence by the Composition–Diamond lemma the set $(d^2 - ab, c^2 - ba, dab - abd, cba - bca)$ is a Gröbner–Shirshov basis of the ideal (R) , and, therefore all basic elements of the category \mathcal{C} (a \mathbb{K} -basis of the algebra $\mathbb{K}\mathcal{C}$) are elements of form $w \in \text{Cat}\langle\Gamma\rangle$ such that $w \neq uw'v$ where $u, v, w' \in \text{Cat}\langle\Gamma\rangle$ and $w' \neq d^2, c^2, dab, cba$.

□

3. GARSIDE THEORY AND GRÖBNER–SHIRSHOV BASIS

This is a key section of this paper. We show how the main concepts of Garside theory can be obtained by using Theorem 3.12. We shall also see that in some cases the corresponding greedy normal form is exactly a Gröbner–Shirshov normal form. For a left-cancellative category and for its arbitrary subfamily we construct a set of reductions and we then show that this subfamily is Garside if and only if the set of reductions is confluent up to a congruence \approx (=deformations of paths by invertible elements).

We start with the following main construction which will be frequently used.

Construction 3.1. Let \mathcal{C}, \mathcal{A} be categories with surjective functor $P : \mathcal{C} \rightarrow \mathcal{A}$, i.e., both its restrictions to objects and elements (=morphisms) are surjective. Let us assume that \mathcal{A} has a presentation $\mathcal{A} = \text{Cat}\langle\Gamma | R\rangle$. Consider a categorical order \leq on the free category $\text{Cat}\langle\Gamma\rangle$. Let us assume that we know a Gröbner–Shirshov basis of the ideal (R) relative to \leq . Hence, by the Composition–Diamond lemma (= the classical verse, see Theorem 2.9 and Remark 2.13), we know a basis $\text{Irr}(R)$ (= a set of all irreducible elements) of \mathcal{A} .

Let us consider a map (=a section of P) $E : \mathcal{A} \rightarrow \mathcal{C}$ that is not a functor in general but $P \circ E = \text{id}_{\mathcal{A}}$. For a given subset $\mathfrak{B} \subseteq \text{Irr}(R)$ we construct the following germ

$$\Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B}) := \bigcup_{a \in \mathfrak{B}} \{E_a, | E_a \bullet E_b := E_{\text{NF}(ab)} \in \Upsilon \text{ whenever } E(a \bullet b) = E(a)E(b)\}.$$

In the case $\mathcal{A} = \mathcal{C}$ and $P = \text{id}_{\mathcal{C}}$ is an identity functor we then denote the germ $\Upsilon_E(\mathcal{C}, \mathcal{C}, \text{id}_{\mathcal{C}}, \mathfrak{B})$ by $\Upsilon_E(\mathcal{C}, \mathfrak{B})$. We consider this partial case in details later in this section.

Definition 3.2. For any $E_x, E_y \in \Upsilon$ we set $E_x \gtrsim E_y$ if $E_x \leq E_y$ and $E_y \not\leq E_x$.

Lemma 3.3. If a germ $\Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$ is left-associative then the binary operation \gtrsim is a partial preorder on the set $\{E_x, x \in \mathfrak{B}\}$.

Proof.

(1) It is clear that $E_a \gtrsim E_a$ because of $a = a\mathbf{1}_{s(a)}$.

(2) Let $E_a \gtrsim E_b$ and $E_b \gtrsim E_c$. Then we have $b = aa'$ and $a = bb'$ in \mathcal{A} with $E_a \bullet E_{a'}, E_b \bullet E_{b'} \in \Upsilon$ that contradicts to the definition of \gtrsim .

(3) Let $E_a \gtrsim E_b$ then $b = aa'$, $E_a \bullet E_{a'} \in \Upsilon$, there is no any relation $a = bb'$ in \mathcal{A} with $E_b \bullet E_{b'} \in \Upsilon$. Let $E_b \gtrsim E_c$ then $c = bb'$ with $E_b \bullet E_{b'} \in \Upsilon$ and there is no any relation $b = cc'$ in \mathcal{A} with $E_c \bullet E_{c'} \in \Upsilon$.

It follows that $c = bb' = aa'b'$. $E_c = E_b \bullet E_{b'} = (E_a \bullet E_{a'}) \bullet E_{b'}$ and by left-associativity of Υ , $E_{a'} \bullet E_{b'} \in \Upsilon$. Hence $E_c = E_a \bullet E_{a'b'}$.

Assume now that $a = cc''$ with $E_c \bullet E_{c''} \in \Upsilon$. Then $b = aa' = cc''a'$. $E_b = E_a \bullet E_{a'} = (E_c \bullet E_{c''}) \bullet E_{a'}$ and by left-associativity of Υ , $E_{c''} \bullet E_{a'} \in \Upsilon$ thus $E_b = E_c \bullet E_{\text{NF}(c''a')}$ that gives a contradiction to the assumption $E_b \gtrsim E_c$. It implies that $E_a \gtrsim E_c$. This completes the proof. □

Definition 3.4. Let $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$ be a left-associative and left-cancellative germ. An element $E_x E_y \in \Upsilon^{[2]}$ is called *reducible* if $y = y_1 y_2$ with $E_x \bullet E_{y_1}, E_{y_1} \bullet E_{y_2} \in \Upsilon$ and $E_x \gtrsim E_{\text{NF}(xy_1)}$. We write $E_x E_y \rightarrow E_{\text{NF}(xy_1)} E_{y_2}$ (of course, the case $E_{y_2} = \mathbf{1}_{t(y_1)}$ is also allowed). In otherwise the element $E_x E_y$ is called *irreducible*.

Construction 3.5. Keep the notations used in the previous Definition and Construction 3.1. Introduce the following set of reductions

$$S_{\Upsilon}(\mathfrak{B}) := \bigcup_{E_x E_y \in \Upsilon^{[2]}} \{r_{x,y} : E_x E_y \rightarrow E_{\text{NF}(xy_1)} E_{y_2}\},$$

and set

$$\overline{S}_{\Upsilon}(\mathfrak{B}) := \bigcup_{\substack{E_x E_y \in \Upsilon^{[2]} \\ E_{\text{NF}(xy_1)} E_{y_2} \in \text{Irr}(S_{\Upsilon}(\mathfrak{B}))}} \{r_{x,y} : E_x E_y \rightarrow E_{\text{NF}(xy_1)} E_{y_2}\}.$$

It is clear that $\overline{S}_{\Upsilon}(\mathfrak{B}) \subseteq S_{\Upsilon}(\mathfrak{B})$.

Proposition 3.6. *Let us consider a germ $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$. An element $E_a E_b \in \Upsilon^{[2]}$ is irreducible if and only if the set $\mathcal{J}_{\Upsilon}(E_a, E_b)$ consists of invertible elements of Υ .*

Proof. By Definition 1.15,

$$\mathcal{J}_{\Upsilon}(E_a, E_b) := \{E_c \in \Upsilon \mid E_a \bullet E_c \in \Upsilon, E_b = E_c \bullet E_d \text{ for some } E_d \in \Upsilon\}.$$

We assume that $E_c \in \mathcal{J}_{\Upsilon}(E_a, E_b)$, $E_b = E_c \bullet E_d$ with $E_a \bullet E_c, E_c \bullet E_d \in \Upsilon$.

(1) Let $E_a E_b$ be irreducible. By, $E_b = E_c \bullet E_d$, $E_a \bullet E_c \in \Upsilon$, and Definition 3.4, $E_a, E_{\text{NF}(ac)}$ must be not \cong -comparable. It follows that $E_{\text{NF}(ac)} \ll E_a$ because of $E_a \ll E_{\text{NF}(ac)}$. Hence $a = acc'$ with $E_{ac} \bullet E_{c'} \in \Upsilon$. We have $E_a = E_{\text{NF}(ac)} E_{c'} = (E_a E_c) E_{c'}$ and by left-associativity of Υ , $E_c \bullet E_{c'} \in \Upsilon$. Hence, $E_a = E_{\text{NF}(acc')} = E_a (E_c E_{c'})$ and by left-cancelativity of Υ , $E_c E_{c'} = \mathbf{1}_{s(c)}$.

Next, $E_a E_c = (E_{\text{NF}(ac)} \bullet E_{c'}) \bullet E_c$ and by left-associativity of Υ , $E_{c'} E_c \in \Upsilon$. Let us consider the element $E_c E_{c'} E_c$. We have $E_c E_{c'} E_c = (E_c E_{c'}) E_c = E_c$ on the other have $E_c E_{c'} E_c = E_c (E_{c'} E_c)$ and by the left-cancelativity of Υ , $E_{c'} E_c = \mathbf{1}_{s(c')}$. Thus $\mathcal{J}_{\Upsilon}(E_a, E_b) \subseteq \Upsilon^{\times}$ as claimed.

(2) Let $\mathcal{J}_{\Upsilon}(E_a, E_b) \subseteq \Upsilon^{\times}$, for some element $E_a E_b \in \Upsilon^{[2]}$. Suppose that $E_a E_b \rightarrow E_{\text{NF}(ac)} E_d$ where $E_{\text{NF}(ac)} E_d$ is irreducible. It is clear $E_c \in \mathcal{J}_{\Upsilon}(E_a, E_b)$ and thus by assumption E_c is invertible. Therefore $E_{\text{NF}(ac)} \ll E_a$ i.e., $E_a \not\ll E_{\text{NF}(ac)}$. This completes the proof. \square

Lemma 3.7. *Let \approx be a deformation of elements of $\text{Cat}(\Upsilon)$ by invertible elements. Then \approx is $\overline{S}_{\Upsilon}(\mathfrak{B})$ -admissible.*

Proof. We have to prove that if whenever $E_a E_b$ is reducible (resp. irreducible) then any $E_{a'} E_{b'} \approx E_a E_b$ is so.

By $E_a E_b \approx E_{a'} E_{b'}$, $E_a = E_{a'} E_e$, $E_{b'} = E_b E_e$, where $E_e \in \Upsilon^{\times}$,

$$\begin{array}{ccc} \cdot & \xrightarrow{E_{a'}} & \cdot \\ \parallel & \left(\begin{array}{c} \xrightarrow{E_b'} \\ \xrightarrow{E_e^{-1}} \end{array} \right) & \parallel \\ \cdot & \xrightarrow{E_a} & \cdot \\ & & \xrightarrow{E_b} \end{array}$$

(1) Let $E_a E_b$ be reducible, say $r_S : E_a E_b \rightarrow E_{\text{NF}(ab_1)} E_{b_2}$. $\Upsilon \ni E_a \bullet E_{b_1} = (E_{a'} E_e) E_{b_1}$, by left-associativity of Υ , $E_e E_{b_1} \in \Upsilon$. We thus have $E_{a'} \bullet E_e, E_e \bullet E_{b_1}, (E_{a'} \bullet E_e) \bullet E_{b_1} \in \Upsilon$. Hence, by Definition 1.13 (3), $E_{a'} \bullet (E_e \bullet E_{b_1}) = (E_{a'} \bullet E_e) \bullet E_{b_1} \in \Upsilon$, i.e., $E_{a'} \bullet E_{\text{NF}(eb_1)} \in \Upsilon$. Therefore, by $E_{b'} = E_e E_b = E_e E_{b_1} E_{b_2} = E_{\text{NF}(eb_1)} E_{b_2}$, there is a reduction $r : E_{a'} E_{b'} \rightarrow E_{\text{NF}(a'eb_1)} E_{b_2}$, i.e., $E_{a'} E_{b'}$ is reducible.

(2) Let $E_a E_b$ be irreducible. Then the statement immediately follows from the Proposition 3.6 and Definition 1.9, because of $E_a E_b = E_{a'} E_{b'}$ with $E_a = E_{a'} E_e$ and $E_b = E_e^{-1} E_{b'}$. \square

Remark 3.8. Let us consider a germ $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$. It seems that we also have to add a set of reduction of the form $E_x E_x^{-1} = \mathbf{1}_{s(x)}$ but it equivalences to $E_x E_x^{-1} \approx \mathbf{1}_{s(x)} \cdot \mathbf{1}_{s(x)}$, indeed,

$$\begin{array}{ccc} \cdot & \xrightarrow{E_x} & \cdot \\ \cdot & \xrightarrow{E_x^{-1}} & \cdot \\ \cdot & \xrightarrow{\mathbf{1}_{s(x)}} & \cdot \\ \cdot & \xrightarrow{\mathbf{1}_{s(x)}} & \cdot \end{array} \quad \begin{array}{c} \parallel \\ \parallel \\ \parallel \\ \parallel \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \xrightarrow{E_x^{-1}} \\ \xrightarrow{E_x} \\ \xrightarrow{\mathbf{1}_{s(x)}} \\ \xrightarrow{\mathbf{1}_{s(x)}} \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \\ \cdot \end{array}$$

thus according to Remark 2.10 we do not do that. However, it is useful to consider an ambiguity of form (E_a, E_b, E_c) with $E_b = E_c^{-1}$ and $\{E_a E_b \rightarrow E_{\text{NF}(ab_1)} E_{b_2}\} \in S_\Upsilon(\mathfrak{B})$. First of all let us note that $E_{b_1} = E_{\text{NF}(b_2c)}$. Indeed, we have $\Upsilon \ni \mathbf{1}_{s(b)} = E_b \bullet E_c = (E_{b_1} \bullet E_{b_2}) \bullet E_c$, hence, by left left-associativity of Υ , $E_{b_2} \bullet E_c \in \Upsilon$. Next, we have $E_b = E_{b_1} \bullet E_{b_2}, E_{b_2} \bullet E_c, E_b \bullet E_c = (E_{b_1} \bullet E_{b_2}) \bullet E_c = \mathbf{1}_{s(b)} \in \Upsilon$ then by Definition 1.13 (3), $\mathbf{1}_{s(b)} = (E_{b_1} \bullet E_{b_2}) \bullet E_c = E_{b_1} \bullet (E_{b_2} \bullet E_c)$, i.e., $E_{b_1} = E_{\text{NF}(b_2c)}^{-1}$, as claimed. Thus we have

$$\begin{array}{ccc} \cdot & \xrightarrow{E_a} & \cdot \\ \cdot & \xrightarrow{\mathbf{1}_{s(b_1)}} & \cdot \\ \cdot & \xrightarrow{E_{\text{NF}(ab_1)}} & \cdot \\ \cdot & \xrightarrow{E_{\text{NF}(b_2c)}} & \cdot \end{array} \quad \begin{array}{c} \parallel \\ \parallel \\ \parallel \\ \parallel \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \\ \cdot \end{array} \begin{array}{c} \xrightarrow{E_{b_1}} \\ \xrightarrow{E_{\text{NF}(b_2c)}} \\ \xrightarrow{E_{\text{NF}(ab_1)}} \\ \xrightarrow{E_{\text{NF}(b_2c)}} \end{array} \begin{array}{c} \cdot \\ \cdot \\ \cdot \\ \cdot \end{array}$$

i.e., $E_a \cdot \mathbf{1}_{s(b)} \approx E_{\text{NF}(ab_1)} \cdot E_{\text{NF}(b_2c)}$, because of $s(b) = s(b_1)$. On the other hand, the ambiguity (E_a, E_b, E_c) gives

$$\begin{array}{ccc} & E_a E_b E_c & \\ & \swarrow \quad \searrow & \\ E_{\text{NF}(ab_1)} E_{b_2} E_c & & E_a \mathbf{1}_{s(b)} \\ \downarrow & & \parallel \\ E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c)} & & E_a \mathbf{1}_{s(b)} \end{array}$$

i.e., according to the \approx -version of the Buchberger–Shirshov algorithm (see Remark 2.11) we get a new reduction $E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c)} \rightarrow E_a \mathbf{1}_{s(b)}$ which is equivalent to $E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c)} \approx E_a \mathbf{1}_{s(b)}$. \square

Lemma 3.9. Let $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$ be a left-associative and left-cancellative germ, then Υ is a Garside germ if and only if a set $\mathcal{I}(E_x, E_y)$ is totally preordered by \preceq .

Proof. (1) Let $\mathcal{I}(E_x, E_y)$ be a totally preordered set by \preceq . Define a map $\mathfrak{F} : \Upsilon^{[2]} \rightarrow \Upsilon$ as follows

$$\mathfrak{F}(E_a, E_b) := \begin{cases} E_{\text{NF}(ab_1)}, & \text{if } E_a E_b \rightarrow E_{\text{NF}(ab_1)} E_{b_2}, \text{ where } E_{\text{NF}(ab_1)} E_{b_2} \text{ is irreducible,} \\ E_a, & \text{in otherwise.} \end{cases} \quad (3.2)$$

Let $E_c \in \mathcal{I}(E_a, E_b)$, then $E_c = E_{\text{NF}(ab')}$ where $E_b = E_{b'} E_{b''}$. Then either $E_c \preceq E_{\text{NF}(ab_1)}$ or $E_{\text{NF}(ab_1)} \preceq E_c$.

Let $E_{\text{NF}(ab_1)} \preceq E_c$, say $E_c = E_{\text{NF}(ab_1)} E_d$. Since $E_c = E_{\text{NF}(ab')}$ then $E_a E_{b'} = E_a E_{b_1} E_d$, by left-cancellativity of Υ , $E_{b'} = E_{b_1} E_d$. Next, we have $E_b = E_{b_1} E_{b_2} = E_{b'} E_{b''}$ then $E_{b_2} = E_d E_{b''}$. Since $E_{\text{NF}(ab_1)} E_{b_2}$ is assumed to be irreducible then by Proposition 3.6, E_d is an invertible element, because of $E_{\text{NF}(ab_1)} E_{b_2} = E_{\text{NF}(ab_1)} E_d E_{b''}$ and $\Upsilon \ni E_{\text{NF}(ab')} = E_{\text{NF}(ab_1)} E_d$. It follows that $E_c = E_{\text{NF}(ab_1)} E_d$ and hence $E_c \preceq E_{\text{NF}(ab_1)}$ for any $E_c \in \mathcal{I}(E_a, E_b)$. Thus \mathfrak{F} is a greatest \mathcal{I} -function on Υ . Hence Υ is a Garside germ.

(2) Conversely, let Υ be a Garside germ. We then get a function \mathfrak{I} obeying \mathcal{I} -law, say, \mathfrak{I} , i.e., we have $\mathfrak{I}(E_a \bullet E_b, E_c) = \mathfrak{I}(E_a, \mathfrak{I}(E_b, E_c))$, where $E_a \bullet E_b \in \Upsilon$. We have

$$\begin{array}{ccc} & E_a E_b E_c & \\ & \swarrow \quad \searrow & \\ E_{\text{NF}(ab)} E_c & & E_a E_{\text{NF}(bc_3)} E_{c_4} \\ \downarrow & & \downarrow \\ E_{\text{NF}(abc_1)} E_{c_2} & & E_{\text{NF}(ad_1)} E_{d_2} E_{c_4} \end{array}$$

then $E_{\text{NF}(abc_1)} = E_{\text{NF}(ad_1)}$, where $E_{\text{NF}(bc_3)} = E_{d_1} E_{d_2}$. By the left-associativity of Υ , $E_{\text{NF}(bc_1)} = E_{d_1}$. Next, we have $E_b E_{c_3} = E_{d_1} E_{d_2} = E_b E_{c_1} E_{d_2}$ then $E_{c_3} = E_{\text{NF}(c_1 d_2)}$. Further, by $E_c = E_{c_1} E_{c_2} = E_{c_3} E_{c_4}$, $E_{c_2} = E_{d_2} E_{c_4}$. We thus get $c_3 = c_1 d_2$ and $c_2 = d_2 c_4$ and the statement follows. \square

Lemma 3.10. *If for any $E_a, E_b \in \Upsilon$ the set $(\mathcal{I}(E_a, E_b), \leq)$ is totally preordered by \leq then all elements of $\Upsilon^{[2]}$ are reduction unique up to \approx under $S_\Upsilon(\mathfrak{B})$.*

Proof. In other words we have to prove that the following diagrams

$$\begin{array}{ccc} E_x E_{\text{NF}(yz)} & \xrightarrow{\bar{r}_{x,yz}} & E_{\text{NF}(xy_1)} E_{\text{NF}(y_2 z)} \\ \bar{r}_{x,yz} \downarrow & \searrow^{\bar{r}_{xy_1 y_2 z} \circ \bar{r}_{x,yz}} & \downarrow \bar{r}_{xy_1 y_2 z} \\ E_{\text{NF}(xw_1)} E_{w_2} & \xrightarrow{\approx} & E_{\text{NF}(xy_1 d_1)} E_{d_2} \end{array} \quad \begin{array}{ccc} E_{y_2} E_z & \xrightarrow{\bar{r}_{y_2,z}} & E_{\text{NF}(y_2 z_1)} E_{z_2} \\ \bar{r}_{y_2,z} \downarrow & \searrow^{\bar{r}_{y_2 z_1 z_2} \circ \bar{r}_{y_2,z}} & \downarrow \bar{r}_{y_2 z_1 z_2} \\ E_{\text{NF}(y_2 z')} E_{z''} & \xrightarrow{\approx} & E_{\text{NF}(y_2 z_1 z_2)} E_{z_{22}} \end{array}$$

are commutative, where all \bar{r} are elements of $\bar{S}_\Upsilon(\mathfrak{B})$.

(1) By the assumption the set $(\mathcal{I}(E_x, E_{\text{NF}(yz)}), \leq)$ is totally preordered. We have $E_{w_1}, E_{\text{NF}(y_1 d)} \in \mathcal{I}(E_x, E_{\text{NF}(yz)})$.

Case 1. Let $E_{w_1} \leq E_{\text{NF}(y_1 d)}$ then $E_{\text{NF}(y_1 d)} = E_{w_1} E_e$. Since $E_{\text{NF}(yz)}, E_{\text{NF}(y_2 z)}$ have been assumed to be equal to $E_{w_1} E_{w_2}$ and $E_{d_1} E_{d_2}$ respectively then we get

$$\begin{aligned} E_{w_1} E_{w_2} &= E_{\text{NF}(y_1 d)} = E_{y_1} E_{y_2} E_z = E_{y_1} E_{\text{NF}(y_2 z)} \\ &= E_{y_1} E_{d_1} E_{d_2} = E_{\text{NF}(y_1 d_1)} E_{d_2} = E_{w_1} E_e E_{d_2}, \end{aligned}$$

hence, by left-cancellativity of Υ , $E_{w_2} = E_e E_{d_2}$, i.e., $E_e \leq E_{w_2}$. Further, $\Upsilon \ni E_{\text{NF}(xy_1 d_1)} = E_x E_{\text{NF}(y_1 d_1)} = E_x E_{w_1} E_e$. Since $E_{w_1} E_e = E_{\text{NF}(y_1 d_1)}$ then $E_{w_1} \bullet E_e \in \Upsilon$. It follows that we get a reduction $E_{\text{NF}(xw_1)} E_{w_2} \rightarrow E_{\text{NF}(xw_1 e)} E_{d_2}$. But $E_{\text{NF}(xw_1)} E_{w_2}$ is assumed to be irreducible then by Proposition 3.6, E_e is invertible. We thus obtain

$$E_{\text{NF}(xy_1 d_1)} = E_x E_{\text{NF}(y_1 d_1)} = E_x E_{w_1} E_e = E_{\text{NF}(xw_1)} E_e,$$

therefore $E_{\text{NF}(xy_1 d_1)} =^{\times} E_{\text{NF}(xw_1)}$, we thus get

$$\begin{array}{ccc} \cdot & \xrightarrow{E_{\text{NF}(xw_1)}} & \cdot \\ \parallel & \begin{array}{c} \xrightarrow{E_{w_2}} \\ \left(\begin{array}{c} E_e \\ \downarrow \\ E_e^{-1} \end{array} \right) \\ \xrightarrow{E_{d_2}} \end{array} & \parallel \\ \cdot & \xrightarrow{E_{\text{NF}(xy_1 d_1)}} & \cdot \end{array}$$

as claimed.

Case 2. Let $E_{\text{NF}(y_1 d_1)} \leq E_{w_1}$, say $E_{w_1} = E_{\text{NF}(y_1 d_1)} E_e$. By the left-associativity of Υ , $E_{d_1} \bullet E_e \in \Upsilon$. Since we have assumed that $E_{\text{NF}(yz)} = E_{w_1} E_{w_2}$, $E_{\text{NF}(y_2 z)} = E_{d_1} E_{d_2}$ then

$$E_{\text{NF}(yz)} = E_{w_1} E_{w_2} = E_{\text{NF}(y_1 d_1)} E_e E_{w_2} = E_{y_1} E_{d_1} E_e E_{w_2}$$

by $E_y = E_{y_1}E_{y_2}$ and left-associativity of Υ , $E_{\text{NF}(y_2z)} = E_{\text{NF}(d_1e)}E_{w_2}$. $E_{\text{NF}(y_2z)} = E_{d_1}E_{d_2}$ implies that $E_{d_2} = E_eE_{w_2}$. Next, by $E_{\text{NF}(xw_1)} = E_{\text{NF}(xy_1d_1)}E_e$ and $E_e \leq E_{d_2}$ then there is a reduction $E_{\text{NF}(xy_1d_1)}E_{d_2} \rightarrow E_{\text{NF}(xy_1d_1e)}E_{w_2}$. Hence, by Proposition 3.6, E_e is invertible because of $E_{\text{NF}(xy_1d_1)}E_{d_2}$ has been assumed to be irreducible. Thus $E_{\text{NF}(xw_1)} =^{\times} E_{\text{NF}(xy_1d_1)}$, we obtain

$$\begin{array}{ccc} \cdot & \xrightarrow{E_{\text{NF}(xw_1)}} & \cdot \\ \parallel & \left(\begin{array}{c} \uparrow E_e \\ \downarrow E_e^{-1} \end{array} \right) & \parallel \\ \cdot & \xrightarrow{E_{\text{NF}(xy_1d_1)}} & \cdot \end{array}$$

and then the statement follows.

(2) We have $E_{z'}, E_{\text{NF}(z_1z_2)} \in \mathcal{I}(E_{y_2}, E_z)$.

Case 1. Let $E_{\text{NF}(z_1z_2)} \leq E_{z'}$, i.e., $E_{z'} = E_{\text{NF}(z_1z_2)}E_e$, hence by left-cancellativity of Υ and $E_z = E_{z'}E_{z''} = E_{\text{NF}(z_1z_2)}E_{z_2z}$, $E_{z_2z} = E_eE_{z''}$. Since $E_{\text{NF}(y_2z')} = E_{\text{NF}(y_2z_1z_2)e}$ then, by Proposition 3.6, $E_e \in \Upsilon^{\times}$. It implies that

$$\begin{array}{ccc} \cdot & \xrightarrow{E_{\text{NF}(y_2z_1z_2)}} & \cdot \\ \parallel & \left(\begin{array}{c} \uparrow E_e^{-1} \\ \downarrow E_e \end{array} \right) & \parallel \\ \cdot & \xrightarrow{E_{y_2z'}} & \cdot \end{array}$$

and the statement follows. □

Case 2. $E_{z'} \leq E_{\text{NF}(z_1z_2)}$, i.e., $E_{\text{NF}(z_1z_2)} = E_{z'}E_e$, then by the left-associativity of Υ and $E_z = E_{z'}E_{z''} = E_{\text{NF}(z_1z_2)}E_{z_2z}$, $E_{z_2z} = E_eE_{z''}$. Since $E_{\text{NF}(y_2z_1z_2)} = E_{\text{NF}(y_2z'e)}$ then by Proposition 3.6, $E_e \in \Upsilon^{\times}$, it implies that

$$\begin{array}{ccc} \cdot & \xrightarrow{E_{\text{NF}(y_2z_1z_2)}} & \cdot \\ \parallel & \left(\begin{array}{c} \uparrow E_e \\ \downarrow E_e^{-1} \end{array} \right) & \parallel \\ \cdot & \xrightarrow{E_{y_2z'}} & \cdot \end{array}$$

and thus the statement follows.

Lemma 3.11. *If a set $(\mathcal{I}(E_a, E_b), \leq)$ is totally preordered for any $E_a, E_b \in \Upsilon$ then all ambiguities of $\overline{S}_{\Upsilon}(\mathfrak{B})$ are \approx -resolvable.*

Proof. Let us consider an ambiguity $(\sigma, \tau, E_a, E_b, E_c)$, we have

$$\begin{array}{ccc} & E_a E_b E_c & \\ \swarrow \iota_{a,b} & & \searrow \iota_{b,c} \\ E_{\text{NF}(ab_1)} E_{b_2} E_c & & E_a E_{\text{NF}(bc_1)} E_{c_2} \end{array}$$

Using Lemma 3.10, we get the following commutative diagrams

$$\begin{array}{ccc} E_{\text{NF}(ab_1)} E_{b_2} E_c & \xrightarrow{\iota_{b_2,c}} & E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1)} E_{c_2} \\ \downarrow \overline{\iota_{b_2,c}} & & \downarrow \overline{\iota_{b_2c_1,c_2}} \\ E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c')} E_{c''} & \xrightarrow{\approx} & E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1c_2)} E_{c_2} \end{array}$$

and

$$\begin{array}{ccc} E_a E_{\text{NF}(bc_1)} E_{c_2} & \xrightarrow{\overline{v_{abc_1}}} & E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1)} E_{c_2} \\ \overline{v_{abc_1}} \downarrow & & \downarrow \overline{v_{ab_1, b_2c_1}} \\ E_{\text{NF}(ad_1)} E_{d_2} E_{c_2} & \xrightarrow{\approx} & E_{\text{NF}(ab_1f_1)} E_{f_2} E_{c_2} \end{array}$$

Let us consider $E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1)} E_{c_2}$. Since $E_{\text{NF}(b_2c_1c_{21})} = E_{f_1} E_{f_2} E_{c_{21}}$, $E_{\text{NF}(ab_1)} \bullet E_{f_1} \in \Upsilon$ we then have reductions $r_1 : E_{f_2} E_{c_2} \rightarrow E_{\text{NF}(f_2c_{21})} E_{c_{22}}$, and $r_2 : E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1c_{21})} \rightarrow E_{\text{NF}(ab_1f_1)} E_{\text{NF}(f_2c_{21})}$. Next, by left-associativity of Υ , $E_{f_2} \bullet E_{c_{21}} \in \Upsilon$ and we get a reduction $r_3 : E_{f_2} E_{c_2} \rightarrow E_{\text{NF}(f_2c_{21})} E_{c_{22}}$. Thus we get

$$\begin{array}{ccc} E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1)} E_{c_2} & \xrightarrow{\overline{v_{b_2c_1, c_2}}} & E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c_1c_{21})} E_{c_{22}} \\ r_2 \downarrow & & \downarrow r_1 \\ E_{\text{NF}(ab_1f_1)} E_{f_2} E_{c_2} & \xrightarrow{r_3} & E_{\text{NF}(ab_1f_1)} E_{\text{NF}(f_2c_{21})} E_{c_{22}} \end{array}$$

Therefore we obtain the following commutative diagram up to \approx

$$\begin{array}{ccccc} & & E_a E_b E_c & & \\ & \swarrow & & \searrow & \\ E_{\text{NF}(ab_1)} E_{b_2} E_c & & & & E_a E_{\text{NF}(bc_1)} E_{c_2} \\ \downarrow & & & & \downarrow \\ E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c')} E_{c''} & & & & E_{\text{NF}(ad_1)} E_{d_2} E_{c_2} \\ & \swarrow \rightsquigarrow & & \nwarrow \rightsquigarrow & \\ & E_{\text{NF}(ab_1f_1)} E_{\text{NF}(f_2c_{21})} E_{c_{22}} & & & \end{array}$$

where the left and right dotted arrows denote the reductions

$$\begin{array}{ccc} E_{\text{NF}(ab_1)} E_{\text{NF}(b_2c')} E_{c''} & \rightsquigarrow & E_{\text{NF}(ab_1f_1)} E_{\text{NF}(f_2c_{21})} E_{c_{22}}, \\ E_{\text{NF}(ad_1)} E_{d_2} E_{c_2} & \rightsquigarrow & E_{\text{NF}(ab_1f_1)} E_{\text{NF}(f_2c_{21})} E_{c_{22}}, \end{array}$$

respectively. It implies that an ambiguity $(\sigma, \tau, E_a, E_b, E_c)$ is resolvable up to \approx , as claimed. \square

Theorem 3.12 (The Main Result). *Let \mathcal{C}, \mathcal{A} be categories, $P : \mathcal{C} \rightarrow \mathcal{A}$ a surjective functor, E its section, and the corresponding germ $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$ (see Construction 3.1) left-cancelative and left-associative.*

Let \approx be a deformation of paths by invertible elements, and consider the correspond set $\overline{S}_\Upsilon(\mathfrak{B})$ of reductions (see Construction 3.5).

The germ Υ is a Garside germ if and only if all ambiguities of the set $\overline{S}_\Upsilon(\mathfrak{B})$ are resolvable up to \approx .

Moreover, the set of all irreducible elements $\text{Irr}(\overline{S}_\Upsilon(\mathfrak{B}))$ coincides with the set of all Υ -normal elements of the germ Υ , and any two Υ -normal decompositions of an element of $\text{Cat}\langle \Upsilon \rangle$ are \approx -equivalent (i.e., are Υ^\times -deformations of one another).

Finally, if Υ is a solid generating family in \mathcal{C} then $\mathcal{C} \cong \text{Cat}\langle \Upsilon \rangle$.

Proof. Let us prove the first statement.

(1) Let Υ be a Garside germ. Then by Lemma 3.9, the set $\mathcal{I}(E_a, E_b)$ is totally preordered by \leq for any E_a, E_b , thus, by Lemma 3.10 and Lemma 3.11 the statement follows.

(2) Conversely, let all ambiguities of the set $\overline{S}_\Upsilon(\mathfrak{B})$ are \approx -resolvable. Define a map $\mathfrak{S} : \Upsilon^{[2]} \rightarrow \Upsilon$ as follows

$$\mathfrak{S}(E_a, E_b) := \begin{cases} E_{\text{NF}(ab_1)}, & \text{if } E_a E_b \rightarrow E_{\text{NF}(ab_1)} E_{b_2} \text{ and } E_{\text{NF}(ab_1)} E_{b_2} \text{ is irreducible,} \\ E_a, & \text{in otherwise.} \end{cases}$$

Let $E_c \in \mathcal{I}(E_a, E_b)$, then $E_c = E_{\text{NF}(ab')}$ where $E_b = E_{b'} E_{b''}$. Consider $E_a E_b$, by assumption, this element is reduction-unique up to \approx under $\overline{S}_\Upsilon(\mathfrak{B})$, i.e., we have the following commutative diagram

$$\begin{array}{ccc} E_a E_b & \xrightarrow{r_{a,b}} & E_{\text{NF}(ab')} E_{b''} \\ \overline{r_{a,b}} \downarrow & & \downarrow \overline{r_{ab',b''}} \\ E_{\text{NF}(ab_1)} E_{b_2} & \xrightarrow{\approx} & E_{\text{NF}(ab'd_1)} E_{d_2} \end{array}$$

where $E_{\text{NF}(ab'd_1)} E_{d_2}$ is assumed to be irreducible, $E_{b''} = E_{d_1} E_{d_2}$, $E_{\text{NF}(ab')} \bullet E_{d_2} \in \Upsilon$. Thus we have $E_{\text{NF}(ab_1)} = E_{\text{NF}(ab'd_1)} E_e$, for some $E_e \in \Upsilon^\times$. We have $E_{\text{NF}(ab'd_1)} E_e = E_{\text{NF}(ab')} E_{d_1} E_e$, i.e., $E_{\text{NF}(ab_1)} = E_{\text{NF}(ab')} E_{d_1} E_e$, hence $E_{\text{NF}(ab')} \preceq E_{\text{NF}(ab_1)}$. Thus \mathfrak{S} is a greatest \mathcal{I} -function on Υ . Hence Υ is a Garside germ.

The last statement immediately follows from Proposition 1.10, Proposition 1.19 (3), Proposition 3.6, and Theorem 1.23. \square

Corollary 3.13. *If a preorder \preceq has a descending chain condition then $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$ is a Garside germ if and only if the corresponding set of polynomials $(\overline{S}_\Upsilon(\mathfrak{B})) = \{E_x E_y - E_{\text{NF}(xy_1)} E_{y_2}\}$, where E_x, E_y run over all generators of the Υ , and whenever all $E_{\text{NF}(xy_1)} E_{y_2}$ is irreducible, is a \approx -closed Gröbner–Shirshov basis relative to \preceq (see Definition 3.2) and then the corresponding Gröbner–Shirshov normal form is exactly the Υ -normal form.*

Proof. It immediately follows from Theorems 2.9, 3.12. \square

Remark 3.14. Consider a germ $\Upsilon = \Upsilon_E(\mathcal{C}, \mathcal{A}, P, \mathfrak{B})$ is assumed to be a left-associative and left-cancellative. We aim to show that a deformations of paths by invertible elements can be also deduced as follows. Let us assume that we should use the classical version of Composition–Diamond lemma. Consider an ambiguity, say, $(\sigma, \tau, E_a, E_b, E_c)$, with $E_a \bullet E_b, E_b \bullet E_c \in \Upsilon$. We then get

$$\begin{array}{ccc} & E_a E_b E_c & \\ r_\tau \swarrow & & \searrow r_\sigma \\ E_{\text{NF}(ab)} E_c & & E_a E_{\text{NF}(bc)} \end{array}$$

hence, adding this relation $E_a E_{\text{NF}(bc)} = E_{\text{NF}(ab)} E_c$, and if $E_a \preceq E_{\text{NF}(ab)}$ we then get a reduction mentioned before. But let us assume now that $E_a \not\preceq E_{\text{NF}(ab)}$, thus we get $E_a =^\times E_{\text{NF}(ab)}$, say $E_a = E_{\text{NF}(ab)} E_e$. Hence, by left-cancellativity of Υ , $E_b E_e = \mathbf{1}_{s(b)}$, then $E_b = E_e^{-1}$, i.e., $E_e \in \Upsilon^\times$. Then adding new relation we get deformations of paths. It is clear that we can, without loss of generality, assume that $E_{\text{NF}(ab)} E_c, E_a E_{\text{NF}(bc)}$ are assumed to be irreducible.

3.1. The Case $\mathcal{C} = \mathcal{A}, P = \mathbf{id}_\mathcal{C}$. Let us consider a left-cancellative category \mathcal{C} with a presentation $\mathcal{C} = \text{Cat}(\Gamma | R)$ and let \approx be a deformation of paths by invertible elements. Let S_R be a set of reductions on $\text{Cat}(\Gamma)$ such that the corresponding set of polynomials (S_R) is to be a \approx -closed Gröbner–Shirshov basis of the ideal $I(R)$ of the category algebra $\mathbb{K}\mathcal{C}$ relative to a \approx -admissible categorical preorder \preceq compatible with S_R having descending chain condition. Thus, by Theorem 2.9, a basis $\mathfrak{B} = \mathfrak{B}(\mathcal{C})$ of \mathcal{C} is the set $\text{Irr}(S_R, \approx)$ and $\mathcal{C} \cong \text{Cat}(\Gamma | a \cdot b = r_{S_R}(ab))$.

Next, for an identity functor $\mathbf{id}_\mathcal{C} : \mathcal{C} \rightarrow \mathcal{C}$ we consider its arbitrary section E and by Construction 3.1 we construct the corresponding germ $\Upsilon_E(\mathcal{C}, \mathfrak{B}')$ and the corresponding set of

reductions $\overline{S}_\Upsilon(\mathfrak{B}')$ for some $\mathfrak{B}' \subseteq \mathfrak{B}$. It is easy to see that for an identity functor $\mathbf{id}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$ considering its section E is the same as choosing the corresponding subset of basis elements of \mathcal{C} , hence it is enough to consider a subset \mathfrak{B}' of basis elements of \mathcal{C} . We thus denote the corresponding germ by $\Upsilon(\mathcal{C}, \mathfrak{B}')$, and instead of E_a we just write a for any $a \in \text{Irr}(S_R)$.

Thus, by Theorem 3.12, we can say that there is the following one-to-one correspondence

$$\left(\begin{array}{l} \text{Garside families in} \\ \text{a left-cancellative category } \mathcal{C} \\ \text{with a presentation } \mathcal{C} = \text{Cat}(\Gamma | R). \end{array} \right) \Leftrightarrow \left(\begin{array}{l} \text{subsets } \mathfrak{B}' \subseteq \text{Irr}(S_R) \text{ such that} \\ \text{the corresponding set } \overline{S}_\Upsilon(\mathfrak{B}') \text{ of reductions} \\ \text{is to be confluent up to } \approx. \end{array} \right)$$

Example 3.15 (The Klein bottle monoid, (cf. [10, A, I. 3.2 and Example IV, 2.35])). Let us consider the Klein bottle monoid

$$K^+ = \text{Smg}\langle a, b \mid bab = a \rangle$$

Set $a < b$ and consider the corresponding deg-lex order \leq on the free monoid $W = W(a, b)$ generated by a, b . We have only one ambiguity (ρ, ρ, ba, b, ab) , where $\rho = (bab, a)$. We have

$$\begin{array}{ccc} & babab & \\ \rho \swarrow & & \searrow \rho \\ a^2b & & ba^2 \end{array}$$

hence we get a new relation $\sigma = (ba^2, a^2b)$ and then the ambiguity $(\rho, \sigma, ba, b, a^2)$. We obtain

$$\begin{array}{c} \begin{array}{c} baa^2b \equiv ba^2ab \xrightarrow{\sigma} a^2bab \xrightarrow{\rho} a^2a \equiv a^3 \\ \sigma \nearrow \\ baba^2 \\ \rho \searrow \\ a^3 \end{array} \end{array}$$

i.e., this ambiguity is resolvable. It is clear that we have no any other ambiguity and thus by the Composition–Diamond lemma the polynomials $bab - a$, $ba^2 - a^2b$ form a Gröbner–Shirshov basis of the ideal $R = (bab - a)$ relative to the order \leq . Hence, by the Composition–Diamond lemma, $\text{Irr}(R)$ is a \mathbb{K} -basis for the semigroup algebra $\mathbb{K}[K^+]$, therefore the basis elements of K^+ are all words w of the W which does not contain subwords of form bab and ba^2 . It is easy to verify that

$$\text{Irr}(R) = \bigcup_{n,m \geq 1} \{1, a^n, b^m, a^n b^m, b^m a, a^n b^m a\}.$$

For any $m \geq 0$ let us consider the following subset \mathfrak{B}_m of $\text{Irr}(R)$

$$\mathfrak{B}_m = \bigcup_{k \geq 0} \{1, a, b^k, b^k a, ab^{m+k}, ab^{m+k} a\}.$$

Let us consider the corresponding germ $\Upsilon_m := \Upsilon(K^+, \mathfrak{B}_m)$ where the partial map $\bullet : \Upsilon_m^{[2]} \rightarrow \Upsilon_m$ is defined as follows

\bullet	1	a	b^k	$b^n a$	ab^{m+p}	$ab^{m+q} a$
1	1	a	b^k	$b^n a$	ab^{m+p}	$ab^{m+q} a$
a	a		$a \bullet b^k$	$a \bullet b^n a$		
b^r	b^r	$b^r a$	b^{r+k}	$b^{r+n} a$	$b^r \bullet ab^{m+p}$	$b^r \bullet ab^{m+q} a$
$b^s a$	$b^s a$		$b^s a \bullet b^k$	$b^s a \bullet b^n a$		
ab^{m+t}	ab^{m+t}	$ab^{m+t} a$	ab^{m+t+k}	$ab^{m+t+n} a$	$ab^{m+t} \bullet ab^{m+p}$	
$ab^{m+l} a$	$ab^{m+l} a$		$ab^{m+l} a \bullet b^k$			

here

$$\begin{aligned}
a \bullet b^k &:= ab^k, & \text{if } k \geq m, \\
a \bullet b^n a &:= ab^n a, & \text{if } n \geq m, \\
b^r \bullet ab^{m+p} &:= \begin{cases} a, & \text{if } r = m + p, \\ b^{r-m-p} a, & \text{if } r > m + p, r - p \geq 2m, \\ ab^{m+p-r}, & \text{if } r < m + p, p \geq r, \end{cases} \\
b^r \bullet ab^{m+q} a &:= ab^{q+m-r} a, & \text{if } r < q + m, q \geq r, \\
b^s a \bullet b^k &:= \begin{cases} a, & \text{if } s = k, \\ b^{s-k} a, & \text{if } s > k, s - k \geq m, \\ ab^{k-s}, & \text{if } s < k, k - s \geq m, \end{cases} \\
ab^{m+t} \bullet ab^{m+p} &:= ab^{t-p} a, & \text{if } t - p \geq m, \\
ab^{m+p} \bullet ab^k &:= ab^{m+p-k} a, & \text{if } m + p > k, p \geq k,
\end{aligned}$$

Set $S = \cup_{u,v \in \mathfrak{B}_m} \{(uv, u \bullet v)\}$. By the straightforward verification it is easy to see that any ambiguity of the set S of reductions are resolvable with respect to the preorder \leq . Hence, by Theorem 3.12, the germ Υ_m is Garside germ for any $m \geq 0$, and hence, \mathfrak{B}_m is a Garside family in K^+ for any $m \geq 0$. Then, by (3.2), the corresponding \mathcal{I} -greatest function is defined as follows $\mathfrak{I}(u, v) := u \bullet v$ for all $u, v \in \mathfrak{B}_m$, it can be also identify with a function H defined as $H(g) := \min_{\leq}(g, ab^m a)$ (see [10, A, IV, Example 2.35]).

4. EXAMPLES

In this section we consider some interesting cases and examples.

4.1. A Free Abelian Monoid. We show how free abelian monoid can be obtained by using our method. We refer to [10, 3–5 pp.] for details.

Take $n \geq 1$, and consider the free abelian monoid \mathbb{N}^n . It is clear that any its element g can be viewed as a map $g : \{1, \dots, n\} \rightarrow \mathbb{N}$. Denote by $g(k)$ the k th entry of g .

For $g, g' \in \mathbb{N}^n$, we define $g \cdot g'(k) := g(k) + g'(k)$ for each k . We define Δ_n by $\Delta_n(k) = 1$ for every k , and put $N_n := \{\eta \in \mathbb{N}^n \mid \eta(k) \in \{0, 1\} \text{ for any } k\}$. For $i \leq n$, define α_i in \mathbb{N}^n by $\alpha_i(k) = \delta_{i,j}$.

Thus we get

$$\mathbb{N}^n = \text{Smg}\langle \alpha_i \alpha_j = \alpha_j \alpha_i, \text{ for all } 1 \leq i, j \leq n \rangle.$$

For $f, g \in \mathbb{N}^n$, say that $f \lesssim g$ is true if $f(i) \lesssim g(i)$ holds for every $1 \leq i \leq n$. We have (see [10, A, I.1, Proposition 1.1])

Every element of \mathbb{N}^n admits a unique decomposition of the form $\Delta_n^d \eta_1 \cdots \eta_p$ with d in \mathbb{N} and $\eta_1, \dots, \eta_p \in N_n$ satisfying $\eta_1 \neq \Delta_n, \eta_p \neq 1$, and, for every $i < p$, $g \lesssim \eta_{i+1}$ implies that $\eta_i g \not\lesssim \Delta_n$, for any $g \in \mathbb{N}^n$.

For instance, let $n = 3$, and take $f = (5, 4, 3)$, we have

$$\begin{aligned}
(5, 4, 3) &= (3, 3, 3) + (2, 1, 0) \\
&= (3, 3, 3) + (1, 1, 0) + (1, 0, 0),
\end{aligned}$$

thus we obtain $f = \Delta_3^3 \cdot (\alpha_1 \alpha_2) \cdot \alpha_1$

Let us consider the following monoid (group) $M = (\mathbb{Z}/2)^n$, it can be also presented as follows

$$(\mathbb{Z}/2)^n = \text{Sgr}\langle \theta, \xi_1, \dots, \xi_n \mid \xi_i^2 = \theta, 1 \leq i \leq n \rangle,$$

where $\theta(k) = 0 \pmod 2$ for all $1 \leq k \leq n$, $\xi_i(k) = \delta_{i,k}$, and for any $\varphi, \psi \in Z$, $\varphi \cdot \psi(k) := \varphi(k) + \psi(k) \pmod 2$.

We have an epimorphism $P : \mathbb{N}^n \rightarrow (\mathbb{Z}/2)^n$, $f \mapsto f \bmod 2$, where $(f \bmod 2)(k) := f(k) \bmod 2$, for all $1 \leq k \leq n$.

Let us consider, for instance, the following order; set $\xi_i > \xi_j$ if $i < j$, and consider then the corresponding deg-lex ordering on all elements of $(\mathbb{Z}/2)^n$. It is clear that the set of polynomials $\cup_{1 \leq i \leq n} \{\xi_i^2 - \theta\}$ is a Gröbner–Shirshov basis. Hence, by the Composition–Diamond lemma (see Theorem 2.9 and Remark 2.13) the set of squarefree words in ξ_i forms a basis \mathfrak{B} for $(\mathbb{Z}/2)^n$. On the other hand, it is clear that any basic element is also an arbitrary map $\varphi : \{1, \dots, n\} \rightarrow \mathbb{Z}/2$. Set $\varphi \cap \psi = \emptyset$ if $\varphi(k) \neq \psi(k)$ for all $1 \leq k \leq n$, and $\varphi \cap \psi \neq \emptyset$ in otherwise.

Take a section $E : (\mathbb{Z}/2)^n \rightarrow \mathbb{N}$ of P defined by $E(\varphi)(k) := E_\varphi(k) := \varphi(k \bmod 2)$, $1 \leq k \leq n$. By Constructions 3.1, 3.5, we thus have a germ

$$\begin{aligned} \Upsilon &= \Upsilon_E(\mathbb{N}, \mathbb{Z}/2, P, \mathfrak{B}) \\ &= \Upsilon = \{E_\varphi, \varphi \in (\mathbb{Z}/2)^n \mid E_\varphi \bullet E_\psi = E_{\varphi \cdot \psi} \text{ whenever } \varphi \cap \psi = \emptyset\} \end{aligned}$$

and a set of reductions

$$\overline{S}_\Upsilon(\mathfrak{B}) = \bigcup_{\substack{\varphi \cap \psi \cdot \chi = \emptyset \\ \varphi \cdot \psi \cap \chi \neq \emptyset}} \{E_\varphi E_{\psi \cdot \chi} \rightarrow E_{\varphi \cdot \psi} E_\chi\}.$$

Next, we have $E_\varphi \approx E_\psi$ if there exists $\varphi' \in (\mathbb{Z}/2)^n$ such that $\psi = \varphi \cdot \varphi'$ and $\varphi \cap \varphi' = \emptyset$. It is easy to see that $\Upsilon^\times = \{E_\theta\}$. Since the number of all generators of Υ is finite then the preorder \approx has descending chain condition.

By the straightforward computation one can get that all ambiguities of $\overline{S}_\Upsilon(\mathfrak{B})$ are resolvable, hence by Theorem 2.9, the corresponding set of polynomials

$$(\overline{S}_\Upsilon(\mathfrak{B})) = \cup_{\substack{\varphi \cap \psi \cdot \chi = \emptyset \\ \varphi \cdot \psi \cap \chi \neq \emptyset}} \{E_\varphi E_{\psi \cdot \chi} - E_{\varphi \cdot \psi} E_\chi\}$$

is a Gröbner–Shirshov basis relative to \approx . Therefore, by Theorem 3.12, Υ is a Garside germ and Υ -normal form is the Gröbner–Shirshov normal form.

Finally, it is clear that the family $\cup_{\varphi \in (\mathbb{Z}/2)^n} \{E_\varphi\}$ is a generating family for \mathbb{N}^n and hence by Theorem 3.12,

$$\mathbb{N}^n \cong \text{Smg}\langle E_\varphi, \varphi \in (\mathbb{Z}/2)^n \mid E_\varphi E_{\psi \cdot \chi} = E_{\varphi \cdot \psi} E_\chi \text{ whenever } \varphi \cap \psi \cdot \chi = \emptyset \text{ and } \varphi \cdot \psi \cap \chi \neq \emptyset \rangle,$$

and $\cup_{\varphi \in (\mathbb{Z}/2)^n} \{E_\varphi\}$ is a Garside family in it.

4.2. “A Natural Appearing of Invertible Elements”. Let us consider the following monoid

$$M = \text{Smg}\langle p, p', r, q \mid prp' = p \rangle.$$

Using an arbitrary order on generators and consider the corresponding deg-lex order on all elements of the free monoid $W = W(p, p', r, q)$ generated by p, p', r, q . We see that $\{prp' - p\}$ is a Gröbner–Shirshov basis for the ideal $(prp' - p)$ in $\mathbb{K}\langle p, p', r \rangle$. Hence by the classical version of the Composition–Diamond lemma (see Theorem 2.9, Remark 2.13),

$$\mathfrak{B} = \bigcup_{u, v, w \in F} \{w \neq uprp'v\}$$

is a basis for M .

Next, let \widetilde{M} be a left-cancellative monoid such that there is a surjective homomorphism $P : \widetilde{M} \rightarrow M$.

Let us consider a section $E : M \rightarrow \widetilde{M}$ of the P and set


$$\begin{aligned} \Upsilon &:= \Upsilon_E(\widetilde{M}, M, P, \mathfrak{B}) \\ &= \{E_p, E_r, E_{p'}, E_q E_{pr}, E_{rq} \mid E_p \bullet E_r = E_{pr}, E_r \bullet E_q = E_{rq}, E_{pr} \bullet E_{p'} = E_p\}. \end{aligned}$$

and


$$\overline{S}_\Upsilon(\mathfrak{B}) = \{E_p E_r \rightarrow E_{pr}, E_r E_q \rightarrow E_{rq}, E_{pr} E_{p'} \rightarrow E_p\}.$$

Suppose further that Υ is a left-associative and left-cancellative germ. By $E_{pr} \bullet E_{p'} \in \Upsilon$, $E_r \bullet E_{p'} \in \Upsilon$, and by $\Upsilon \ni E_p \bullet E_r = (E_{pr} \bullet E_{p'}) \bullet E_r$, $E_{p'} \bullet E_r \in \Upsilon$. Therefore, by Definition 1.13 (3), $E_p \bullet (E_r \bullet E_{p'}) = (E_p \bullet E_r) \bullet E_{p'}$. Hence, $E_r E_{p'} = \mathbf{1}$ because of $(E_p \bullet E_r) \bullet E_{p'} = E_p \bullet (E_r \bullet E_{p'}) = E_p$.

Next, let us consider an ambiguity $(E_r, E_{p'}, E_r)$, we obtain




then by left-cancellativity of Υ , $E_{p'} E_r = \mathbf{1}$. The ambiguity $(E_{p'}, E_r, E_q)$ gives



i.e., we have $E_{p'} E_{rq} = E_q$.

Finally let us consider an ambiguity (E_p, E_r, E_q) , we have



but $E_{pr} \leq E_p$ because of $E_p = E_{pr} E_{p'}$, i.e., $E_p \not\approx E_{pr}$. Thus if \approx is a deformation of elements by invertible elements (see Definition 1.9), we then get that all ambiguities of the system $\overline{S}_\Upsilon(\mathfrak{B})$ are resolvable up to \approx because of

$$\begin{array}{ccc} \cdot & \xrightarrow{E_p} & \cdot \xrightarrow{E_{rq}} \cdot \\ \parallel & \left(\begin{array}{c} \nearrow \\ E_r \\ \searrow \end{array} \right) & \parallel \\ \cdot & \xrightarrow{E_{pr}} & \cdot \xrightarrow{E_q} \cdot \end{array}$$

i.e., $E_{pr} E_q \approx E_p E_{rq}$. Hence, by Theorem 3.12, Υ is a Garside germ.

4.3. The Artin-Tits Monoid.

4.3.1. *Coxeter Systems and Groups.* Let \mathcal{S} be a set. A matrix $M_{\mathcal{S}} : \mathcal{S} \times \mathcal{S} \rightarrow \{1, 2, \dots, \infty\}$ is called a *Coxeter matrix* if it satisfies

$$\begin{aligned} m(a, b) &= m(b, a); \\ m(a, b) &= 1 \text{ if and only if } a = b, \end{aligned}$$

here $a, b \in \mathcal{S}$.

Equivalently, $M_{\mathcal{S}}$ can be represented by a *Coxeter graph* whose node set is \mathcal{S} and whose edges are the unordered pairs $\{a, b\}$ such that $m(a, b) \geq 3$. The edges with $m(a, b) \geq 4$ are labeled by that numbers.

Definition 4.1. Let \mathcal{S} be a set, \mathcal{W} a group. We say that $(\mathcal{W}, \mathcal{S})$ is a *Coxeter system*, and that \mathcal{W} is a *Coxeter group*, if \mathcal{W} admits the presentation

$$\mathcal{W} = \text{Smg}\langle \mathcal{S} \mid s^2 = 1, \text{ for all } s \in \mathcal{S}, \text{ and whenever } m(a, b) \neq \infty, (ab)^{m(a,b)} = (ba)^{m(b,a)} \rangle.$$




FIGURE 2. The Coxeter matrix and its corresponding graph.

Given a Coxeter system $(\mathcal{W}, \mathcal{S})$, each element $w \in \mathcal{W}$ can be written as a product of generators: $w = s_1 s_2 \cdots s_k$, $s_i \in \mathcal{S}$. If k is minimal among all such expressions for w , then k is called the *length* of w (written $\ell(w) = k$) and the word $s_1 s_2 \cdots s_k$ is called a *reduced word* (or *reduced decomposition* or *reduced expression*) for w .

The following properties are fundamental in the combinatorial theory of Coxeter groups: they characterize such groups.

(Deletion Property). *If $w = s_1 s_2 \cdots s_k$ and $\ell(w) < k$, then $w = s_1 \cdots \widehat{s_i} \cdots \widehat{s_j} \cdots s_k$ for some $1 \leq i < j \leq k$.*

(Exchange Property). *Let $w = s_1 s_2 \cdots s_k$ be reduced expression and $s \in \mathcal{S}$. If $\ell(sw) \leq \ell(w)$, then $sw = s_1 \cdots \widehat{s_i} \cdots s_k$ for some $1 \leq i < j \leq k$, for some $1 \leq i \leq k$.*

Theorem 4.2 ([4, Theorem 1.5.1]). *Let \mathcal{W} be a group and \mathcal{S} a set of generators of order 2. Then the following are equivalent.*

- (1) $(\mathcal{W}, \mathcal{S})$ is a Coxeter system.
- (2) $(\mathcal{W}, \mathcal{S})$ has the Exchange Property.
- (3) $(\mathcal{W}, \mathcal{S})$ has the Deletion Property.

For given $u, v \in \mathcal{W}$ we write $u \perp v$ if $\ell(uv) = \ell(u) + \ell(v)$, and $u \not\perp v$ if $\ell(uv) \neq \ell(u) + \ell(v)$.

Lemma 4.3. *Assume that $\ell(uv) = \ell(u) + \ell(v)$ and $\ell(vw) = \ell(v) + \ell(w)$ for some reduced $u, v, w \in \mathcal{W}$, i.e., $u \perp v$ and $v \perp w$. Then $uv \perp w$ and $u \perp vw$.*

Proof. Let $u = s_{i_1} \cdots s_{i_p}$, $v = s_{j_1} \cdots s_{j_q}$, and $w = s_{k_1} \cdots s_{k_r}$.

Let us assume that $\ell((uv)w) < \ell(uv) + \ell(w)$ or $\ell(u(vw)) < \ell(u) + \ell(vw)$. Then by the Deletion Property, $uvw = s_{i_1} \cdots \widehat{s_\alpha} \cdots \widehat{s_\beta} \cdots s_{k_r}$. Since $\ell(uv) = \ell(u) + \ell(v)$ and $\ell(vw) = \ell(v) + \ell(w)$ then 1) either $\alpha \in \{i_1, \dots, i_p\}$, $\beta \in \{j_1, \dots, j_q\}$, or 2) $\alpha \in \{i_1, \dots, i_p\}$, $\beta \in \{k_1, \dots, k_r\}$, or 3) $\alpha \in \{j_1, \dots, j_q\}$, $\beta \in \{k_1, \dots, k_r\}$. On the other hand, u, v, w are assumed to be reduced, hence we get a contradiction and the statement follows. \square

Corollary 4.4. *If $uv \perp w$ and $u \perp v$, then $u \perp vw$ and $v \perp w$.*

Proof. By $uv \perp w$, and $u \perp v$, $\ell(uvw) = \ell(u) + \ell(v) + \ell(w)$, and using Lemma 4.3, the statement follows. \square

Definition 4.5 ([4, 3.4]). Let $(\mathcal{W}, \mathcal{S})$ be a Coxeter system, $\mathfrak{R}(w)$ be the set of all reduced decompositions of an element w . The *normal form* of an element $w \in \mathcal{W}$ is $\min \mathfrak{R}(w)$, where the minimal is taken with respect to lexicographic order. Denote by $\mathfrak{B}(\mathcal{W})$ the set of all elements $w \in \mathcal{W}$ such that $\min \mathfrak{R}(w) = w$.

As before we denote by $\text{NF}(w)$ a normal form of an element w . We refer to [4] (especially section 3.4) for details and how to compute a normal form of an element of an arbitrary Coxeter group. However, just for reader convenience, we recall an elegant algorithm, see Appendix, to compute a normal form for any Coxeter group. This algorithm is based on so-called “the numbers game” (see [4, 4.3]) offers a general method for finding combinatorial representatives of the group elements.

4.3.2. Artin–Tits monoids.

Definition 4.6. Let $(\mathcal{W}, \mathcal{S})$ be a Coxeter system given by a Coxeter matrix M , an *Artin–Tits monoid* associated with $(\mathcal{W}, \mathcal{S})$, is a monoid, denoted by $B^+(\mathcal{W}, \mathcal{S})$ or shortly B^+ , admits the presentation

$$B^+(\mathcal{W}, \mathcal{S}) := \langle \mathcal{S} \mid \langle a, b \rangle^{m(a,b)} = \langle b, a \rangle^{m(b,a)} \text{ whenever } m(a, b) \neq \infty \rangle$$

here $m(a, b)$ are elements of the Coxeter matrix M . If $m(a, b) = \infty$, then there is no relation for a and b .

Let us turn to the preorder \succcurlyeq (Definition 3.2). It is clear the $E_{\text{NF}(u)} \succcurlyeq E_{\text{NF}(v)}$ if and only if $\ell(u) < \ell(v)$.

Proposition 4.7 (cf. [10, B, IX.1.3, Proposition 1.35]). *Let $(\mathcal{W}, \mathcal{S})$ be a Coxeter system. Consider the corresponding Artin–Tits monoid $B^+(\mathcal{W}, \mathcal{S})$. Let $P : B^+ \rightarrow \mathcal{W}$ be the corresponding surjective, and $E : \mathcal{W} \rightarrow B^+$ its sections which is defined as follows: any $s \in \mathcal{S}$ maps to the same s in B^+ . Define the corresponding germ $\Upsilon = \Upsilon_E(B^+, \mathcal{W}, P, \mathfrak{B}(\mathcal{W}))$ as follows*

$$\Upsilon := \text{Smg}\langle E_{\text{NF}(w)}, w \in \mathcal{W} \mid E_{\text{NF}(u)} \bullet E_{\text{NF}(v)} := E_{\text{NF}(uv)}, \text{ whenever } u \perp v \rangle.$$

Then the germ Υ is a Garside germ, and

$$B^+(\mathcal{W}, \mathcal{S}) \cong \text{Smg}\langle E_{\text{NF}(w)}, w \in \mathcal{W} \mid E_{\text{NF}(u)} E_{\text{NF}(vw)} = E_{\text{NF}(uv)} E_{\text{NF}(w)}, u \perp v \perp w \rangle,$$

and finally a basis of $B^+(\mathcal{W}, \mathcal{S})$ can be described as the following set

$$\bigcup_{\substack{u, v, \dots, w \in \mathcal{W} \\ u \perp v \perp \dots \perp w}} \{E_{\text{NF}(\pi_1)} E_{\text{NF}(\pi_2)} \cdots E_{\text{NF}(\pi_k)}\}.$$

Proof. Let us prove that the germ Υ is a Garside germ.

First of all we have to show that Υ is a left-cancellative and left-associative germ.

(1) Let $E_u \bullet E_v, E_v \bullet E_w \in \Upsilon$, then $u \perp v$ and $v \perp w$. By Lemma 4.3, $E_u \bullet (E_v \bullet E_w) \in \Upsilon$ if and only if $(E_u \bullet E_v) \bullet E_w \in \Upsilon$, and if so then they are equal.

(2) Since \mathcal{W} is assumed to be a group then Υ is left and right-cancellative. Next, by Corollary 4.4, Υ is left-associative.

(3) By Construction 3.5 we have the following set of reductions

$$\overline{\mathfrak{S}}_{\Upsilon}(\mathfrak{B}(\mathcal{W})) = \bigcup_{u \perp v \perp w} \{r_{u,vw} : E_{\text{NF}(u)} E_{\text{NF}(vw)} \rightarrow E_{\text{NF}(uv)} E_{\text{NF}(w)}\}.$$

It is clear that all possible ambiguities are $(r_{u,vw}, r_{vw,hg}, E_{\text{NF}(u)}, E_{\text{NF}(vw)}, E_{\text{NF}(hg)})$ where $u \perp v \perp w$ and $vw \perp h \perp g$.

By Corollary 4.4, $w \perp h$, $v \perp wh$ and we get

$$\begin{array}{ccc}
& & E_{\text{NF}(uv)}E_wE_{\text{NF}(hg)} \xrightarrow{r_{w,hg}} E_{\text{NF}(uv)}E_{\text{NF}(wh)}E_g \\
& \nearrow^{r_{u,vw}} & \\
E_uE_{\text{NF}(vw)}E_{\text{NF}(hg)} & & \\
& \searrow_{r_{vw,hg}} & \\
& & E_uE_{\text{NF}(vwh)}E_g \xrightarrow{r_{uv,wh}} E_{\text{NF}(uv)}E_{\text{NF}(wh)}E_g
\end{array}$$

Thus all ambiguities of $\overline{S}_\Upsilon(\mathfrak{B}(\mathcal{W}))$ are resolvable and hence by Theorem 3.12 the first statement follows.

Next, it is easy to see that the set $\cup_{w \in \mathcal{W}} \{E_{\text{NF}(w)}\}$ is a generating family for $B^+(\mathcal{W}, \mathcal{S})$. Hence, by Proposition 1.20, the second statement follows.

Finally, by Theorem 3.12, the last statement follows. \square

It is clear that if a Coxeter group \mathcal{W} is finite then the corresponding preorder \approx has the descending chain condition, thus, by Corollary 3.13, the corresponding Υ -normal form of an Artin monoid is exactly the corresponding Gröbner–Shirshov normal form.

4.4. A Greedy normal form on the braid monoids B_n^+ .

Let us consider the partial case $\mathcal{W} = \mathfrak{S}_n$ (= the symmetric group). It is well known that a symmetric group has the following Coxeter presentation.

Proposition 4.8 ([7, 8]). *Let $\mathcal{S} = \{s_1, \dots, s_{n-1}\}$ be the set of generators (transpositions) of the symmetric group \mathfrak{S}_n . Set $s_i > s_j$ whenever $i > j$ and consider the corresponding deg-lex ordering $>$ on the free monoid generated by s_1, \dots, s_{n-1} .*

A Gröbner–Shirshov basis for the symmetric group \mathfrak{S}_n , with respect to the order $>$, is the following set of relations:

- (1) $s_i^2 = 1$, for any $1 \leq i \leq n$
- (2) $s_i s_j = s_j s_i$, for $i - j \geq 2$ and $1 \leq i, j \leq n$,
- (3) $s_{i+1} s_i s_{i-1} \cdots s_j s_{i+1} = s_i s_{i+1} s_i s_{i-1} \cdots s_j$, if $i + 1 \geq j$ and $1 \leq i, j \leq n$.

As a consequence, using the Composition–Diamond lemma, we obtain the following

Corollary 4.9 ([7]). *The set $\mathfrak{B}(\mathfrak{S}_n) := \{s_{1i_1} s_{2i_2} \cdots s_{ni_n} \mid i_k \leq k + 1\}$ consists of Gröbner–Shirshov normal forms for \mathfrak{S}_n in the generators $s_i := (i, i + 1)$ relative to the deg-lex ordering, where $s_{\alpha\beta} := s_\beta s_{\beta-1} \cdots s_\alpha$ for $\beta \geq \alpha$ and $s_{\beta, \beta+1} := 1$.*

Let us consider a braid monoid B_n^+ , i.e., a monoid generated by $\sigma_1, \dots, \sigma_n$; its elements are called positive braids. We have a homomorphism $P : B_n^+ \rightarrow \mathfrak{S}_n$; given a positive braid B , the strands define a permutation $p(B)$ from the top set of endpoints to the bottom set of endpoints.

Take $\pi \in \mathfrak{S}_n$ with the normal form $\text{NF}(\pi) = s_{1i_1} s_{2i_2} \cdots s_{mi_m} \in \text{NF}(\mathfrak{S}_n)$, i.e.,

$$\text{NF}(\pi) = (s_{i_1} s_{i_1-1} \cdots s_1)(s_{i_2} s_{i_2-1} \cdots s_2) \cdots (s_{i_m} s_{i_m-1} \cdots s_m)$$

and, as above, set $\ell(\pi) := \ell(\text{NF}(\pi))$ (= the length). Define then a map $E : \mathfrak{S}_n \rightarrow B_n^+$ as follows

$$E(\text{NF}(\pi)) = (\sigma_{i_1} \sigma_{i_1-1} \cdots \sigma_1)(\sigma_{i_2} \sigma_{i_2-1} \cdots \sigma_2) \cdots (\sigma_{i_m} \sigma_{i_m-1} \cdots \sigma_m),$$

it is clear that E is a section for P .

Next, as above, we see that the corresponding germ $\Upsilon = \Upsilon_E(B_n^+, \mathfrak{S}_n, P, \mathfrak{B}(\mathfrak{S}_n))$ can be also described as follows

$$\Upsilon = \{E_{\text{NF}(\pi)}, \pi \in \mathfrak{S}_n \mid E_{\text{NF}(\pi)} \bullet E_{\text{NF}(\tau)} := E_{\text{NF}(\pi\tau)}, \text{ whenever } \pi \perp \tau\}.$$

Thus, by Proposition 4.7, *the corresponding set of reductions*

$$\overline{S}_\Upsilon(\mathfrak{B}(\mathfrak{S}_n)) = \bigcup_{\pi, \tau, \xi \in \mathfrak{S}_n} \{r_{\pi, \tau \xi} : E_{\text{NF}(\pi)} E_{\text{NF}(\tau \xi)} \rightarrow E_{\text{NF}(\pi \tau)} E_{\text{NF}(\xi)}, \pi \perp \tau \perp \xi\},$$

is resolvable and, by Theorem 2.9, *the corresponding set of polynomials*

$$(\overline{S}_\Upsilon(\mathfrak{B}(\mathfrak{S}_n))) = \bigcup_{\pi, \tau, \xi \in \mathfrak{S}_n} \{r_{\pi, \tau \xi} : E_{\text{NF}(\pi)} E_{\text{NF}(\tau \xi)} - E_{\text{NF}(\pi \tau)} E_{\text{NF}(\xi)}, \pi \perp \tau \perp \xi\},$$

is a Gröbner–Shirshov basis relative to the \succsim , and then the corresponding Gröbner–Shirshov normal form is normal form is exactly the greedy normal form, i.e., the set $\text{Irr}(\overline{S}_\Upsilon(\mathfrak{B}(\mathfrak{S}_n)))$ of irreducible elements under the $\overline{S}_\Upsilon(\mathfrak{B}(\mathfrak{S}_n))$

$$\text{Irr}(\overline{S}_\Upsilon(\mathfrak{B}(\mathfrak{S}_n))) := \bigcup_{\substack{\pi_1, \dots, \pi_k \in \mathfrak{S}_n \\ \pi_1 \perp \pi_2 \perp \dots \perp \pi_k}} \{E_{\text{NF}(\pi_1)} E_{\text{NF}(\pi_2)} \cdots E_{\text{NF}(\pi_k)}\},$$

coincides with the set of the greedy normal form of elements of B_n^+ . Finally for any $\pi \in \mathfrak{S}_n$, the element $E_{\text{NF}(\pi)}$ coincides with the Adjan–Thurston generator R_π .

Remark 4.10. In [10, A, VI, Example 2.72] it was shown how to obtain the braid monoid (and greedy normal form) via the symmetric group. This approach is very similar to this way. In particular the elements of $\pi, \tau \in \mathfrak{S}_n$ are called *tight* if $\ell(\pi \tau) = \ell(\pi) + \ell(\tau)$ (in a sense described there).

Remark 4.11. In [5, 3.1.4] it was said that the corresponding set of polynomials

$$(\overline{S}_\Upsilon(\mathfrak{B}(\mathfrak{S}_n))) = \bigcup_{\pi \perp \tau \perp \xi} \{E_{\text{NF}(\pi)} E_{\text{NF}(\tau \xi)} - E_{\text{NF}(\pi \tau)} E_{\text{NF}(\xi)}\}$$

is a Gröbner–Shirshov basis relative to the following ordering \leq . We assume that $s_1 < s_2 < \dots < s_{n-1}$ and define $E_{\text{NF}(\pi)} \leq E_{\text{NF}(\tau)}$ if and only if $\ell(\text{NF}(\pi)) > \ell(\text{NF}(\tau))$ or $\ell(\text{NF}(\pi)) = \ell(\text{NF}(\tau))$ and $\text{NF}(\pi) < \text{NF}(\tau)$ (lexicographical order). It is easy to see that this order is an extension of the preorder \succsim .

APPENDIX: THE NUMBER GAME

Let us recall the “number game” that first appeared in a somewhat restricted version related to Kac–Moody Lie algebras in [14] and the general version given in [4] that is due to Eriksson [12]. We refer to [4, Ch.I,4] for more details.

Let $(\mathcal{W}, \mathcal{S})$ be a Coxeter system given by a Coxeter matrix M . Define a function $\kappa : \mathcal{S} \times \mathcal{S} \rightarrow \mathbb{R}$ as follows;

$$\left\{ \begin{array}{l} \kappa_{s,s} = -2, \quad s \in \mathcal{S}; \\ \kappa_{a,b} = 0, \quad m(a,b) = 2, \\ \kappa_{a,b} > 0, \\ \kappa_{a,b} \kappa_{b,a} = 4 \cos^2 \frac{\pi}{m(a,b)}, \quad m(a,b) \neq \infty, \\ \kappa_{a,b} \kappa_{b,a} \geq 4, \quad m(a,b) = \infty \end{array} \right. , \quad m(-, =) < 3, \quad m(-, =) \geq 3 \quad (4.3)$$

Now we present (see [4, I.4.3]) the Number Game to calculate normal form. We label each node of the corresponding Coxeter graph with some real numbers, and each such assignment thought of as a position in a certain “game”. The “moves” in the game are local rearrangements of the assigned values at a chosen node s and its neighbors, governed by the labels of the edges surrounding in s in the Coxeter graph. The point of this game is that it gives a combinatorial model of the Coxeter group, where group elements correspond to positions and reduced decompositions correspond to play sequences.

The starting position for the game can be any distribution $\mathcal{S} \ni s \rightarrow p_s \in \mathbb{R}$ or real numbers p_s to the nodes $s \in \mathcal{S}$ of the Coxeter graph. A position is called positive if $p_s > 0$ for all $s \in \mathcal{S}$. The special position with $p_s = 1$ for all $s \in \mathcal{S}$ is called the unit position and denoted by $\mathbf{1}$.

Next, moves are defined as follows. A *firing of node s* changes a position $p \in \mathbb{R}^{\mathcal{S}}$ in the following way

- (1) Switch sign of the value at s .
- (2) Add $k_{s,a}p_a$ to the value at each neighbor a of s .
- (3) Leave all other values unchanged.

Such a move is called *positive* if $p_s > 0$, and *negative* if $p_s < 0$. A *positive game* is one that is played with positive moves from a given starting position, and similarly for a negative game. A *play sequence* is a word $s_{i_1}s_{i_2}\cdots s_{i_k}$, $s_{i_j} \in \mathcal{S}$, recording a game in which s_{i_1} was fired first, then s_{i_2} , then s_{i_3} , and so on. Similarly, a *positive play sequence* records a positive game and a *negative play sequence* records a negative game.


Theorem 4.12 ([4, Theorem 4.3.1]). *Let $p \in \mathbb{R}^{\mathcal{S}}$ be a starting position, s_{i_1}, \dots, s_{i_k} a play sequence, denote by $p^{s_{i_1}\cdots s_{i_k}}$ a position reached from p by this play sequence. Let $\mathfrak{P}_p \subseteq \mathbb{R}^{\mathcal{S}}$ denote the set of all positions that can be reached this way.*

- *Two play sequences $s_{i_1}s_{i_2}\cdots s_{i_p}$ and $s_{j_1}s_{j_2}\cdots s_{j_q}$ lead to the same position if and only if $s_{i_1}s_{i_2}\cdots s_{i_p} = s_{j_1}s_{j_2}\cdots s_{j_q}$ as elements of \mathcal{W} .*
- *The induced mapping $w \mapsto p^w$ is a bijection $\mathcal{W} \rightarrow \mathfrak{P}_p$.*
- *The play sequence $s_{i_1}s_{i_2}\cdots s_{i_k}$ is positive if and only if $s_{i_1}s_{i_2}\cdots s_{i_k}$ is a reduced decomposition.*

This Theorem implies the following algorithm for finding normal form $\text{NF}(w)$ of an element w of a Coxeter group \mathcal{W} .

- (1) Take an expression $w = s_{i_1}s_{i_2}\cdots s_{i_k}$.
- (2) Play from $\mathbf{1}$ according to the play sequence $s_{i_k}, \dots, s_{i_2}, s_{i_1}$.
- (3) Set $p := p^{s_{i_k}\cdots s_{i_2}s_{i_1}}$.
- (4) Play from p to $\mathbf{1}$ by firing at each step the minimal negative node.
- (5) Record the obtained play sequence $s_{j_1}, s_{j_2}, \dots, s_{j_n}$.
- (6) $\text{NF}(w) = s_{j_1}, s_{j_2}, \dots, s_{j_n}$.

Example 4.13. Let us consider the following Coxeter graph



Put $a < b < c < d$. Let us find the normal form of the word $w = babcdb$. Let us find the starting position (see fig.3). We have $p = (-3, -6, -4, 7)$. Now we have to play from this position to position $\mathbf{1}$ by firing at each step the minimal negative node (see fig.4). Thus we obtain $\text{NF}(w) = abacbd$.




FIGURE 3. The start position p^{babca} is found.




FIGURE 4. We are firing at each step the minimal negative node.

REFERENCES

- [1] S.I. Adjan, Fragments of the word Delta in a braid group, *Mat. Zam. Acad. Sci. SSSR* **36**(1) (1984), 25–34, (Russian); English translation in *Math. Notes of the Acad. Sci. USSR* **36**(1) (1984), 505–510.
- [2] E. Artin, Theory der Zöpfe, *Abh. Math. Seminar, Hamburg Univ.* **4** (1925), 47–72.
- [3] G. M. Bergman, The Diamond Lemma for ring theory, *Adv. in Math.* **29** (1978) 178–218.
- [4] A. Björner and F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics, vol. 231, Springer, New York, (2005)
- [5] L.A. Bokut and Y. Chen, Gröbner–Shirshov basis and their calculation, *Bull. Math. Sci.* **4**, (2014), 325–395.

- [6] L.A. Bokut, Y.Q. Chen and Y. Li, Gröbner–Shirshov Bases for Categories. Nankai Series in Pure, Applied Mathematics and Theoretical Physical, Operads and Universal Algebra, **9**, 1–23, (2012)
- [7] L.A. Bokut and L.S. Shiao, Gröbner–Shirshov bases for Coxeter groups, *Commun. Algebra*, **29**, 4305–4319, (2001).
- [8] M.A. Borges–Trenard, M. Borges–Quintana, J.A. Castellanos–Garzón and E. Martíns–Moro, The Symmetric Group given by a Gröbner basis, *J. Pure and App. Algebra*, **207** (2006), 149–154.
- [9] P. Dehornoy, Garside and Quadratic Normalization: a survey, <http://arxiv.org/abs/1504.07788>
- [10] P. Dehornoy with F. Digne, E. Godele, D. Krammer and J. Michel, *Foundations of Garside Theory*. EMS Tracts in Mathematics vol. 22 (2015).
- [11] D.B.A. Epstein, I.W. Cannon, D.E. Holt, S.V.F. Levy, M.S. Paterson and W.P. Thurston, Word Processing in Groups, Jones and Bartlett Publishers, INC., 1992.
- [12] K. Eriksson, *Strongly convergent games and Coxeter groups*, Ph.D. Thesis, KTH, Stockholm, Sweden, 1993.
- [13] S. Mac Lane, “*Categories for the Working Mathematician*,” Graduate Texts in Mathematics, **5**, Springer, Berlin/New Yourk, (1971).
- [14] S. Mozes, Reflection process on graphs and Weyl groups, *J. Combin. Theory, Ser. A* **53** (1990), 128–142.
- [15] E. Szpilrajn, “Sur léxtension de l’ordre partiel”, *Fundamenta mathematicae.*, **16**, 386–38 (1930).
- [16] W. Thurston, Finite state algorithm for the braid group, *Circulated notes*, 1988.

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, FACULTY OF COMPUTER SCIENCE, POKROVSKY BOULEVARD 11, MOSCOW, 109028 RUSSIA

Email address: wickktor@gmail.com