Shear free barotropic perfect fluids cannot rotate and expand simultaneously

Rituparno Goswami*

Astrophysics Research Centre and School of Mathematics, Statistics and Computer Science, University of KwaZulu-Natal, Private Bag X54001, Durban 4000, South Africa.

George F. R. Ellis^{\dagger}

Department of Mathematics and Applied Mathematics, University of Cape Town, Cape Town, Western Cape, South Africa. (Dated: January 27, 2022)

We prove here a long standing conjecture in general relativity, that if barotropic perfect fluid is moving in a shear free way, then it must be either expansion free or rotation free.

I. INTRODUCTION

Shear free perfect fluid solutions in General Relativity have been the subject of considerable study [1]. In general relativity fluid solutions, the shear σ_{ab} is the link between the free gravitational field represented by the Weyl Tensor C_{abcd} and the fluid represented by the matter stress tensor T_{ab} , because the electric part E_{ab} of the Weyl tensor is a tidal force [2] that generates shear. That is (see [3] [4]),

$$\dot{\sigma}_{ab} = \mathcal{F}_{ab} + E_{ab} \tag{1}$$

where \mathcal{F}_{ab} depends on the shear, expansion, acceleration, vorticity, and anisotropic pressure π_{ab} . This means that shear-free solutions (which include self-similar solutions) are very special: these terms must be just right to cancel out the contribution from E_{ab} so that (1) implies

$$\mathcal{C}_1: \{\sigma_{ab} = 0\} \Rightarrow \mathcal{F}_{ab} = -E_{ab}.$$
 (2)

That is, the propagation equation (1) becomes the constraint equation (2) (which is just (34) below).

To test for consistency, one takes the time derivative of (2) and substitutes from the propagation equations (1) and constraint equation (2) to either get (a) an identity, (b) an inconsistency, or (c) a new constraint equation C_2 . In the latter case one repeats, thus deriving a sequence of constrains C_n . One eventually either ends up with an identity or a contradiction, showing such solutions are either possible or not possible under certain circumstances.

In the case of pressure free (baryonic) matter, following a pioneering paper by Kurt Gödel for a spatially homogeneous case [5], it was shown in [6] by using this method that it is an exact property of the Einstein Field Equations that shear-free solutions cannot both expand and rotate. Generically when p = 0,

$$\{\sigma = 0\} \Rightarrow \omega \Theta = 0, \tag{3}$$

where σ is the fluid shear, ω its vorticity, and Θ its expansion [7] [3] [4]. This is true irrespective of the value of the cosmological constant Λ . In the case of perfect fluids many generalisations of this result have been obtained, as summarised in [8] [9] [11], but the generic result analogous to (3) has not yet been obtained.

In this paper we derive that generic result for a general barotropic perfect fluid, i.e. the case when there is no heat flux or anisotropic stress:

$$\{q_a = 0 = \pi_{ab}\} \Rightarrow T_{ab} = (\mu + p)u_a u_b + pg_{ab}, \quad (4)$$

where $u^a : u_a u^a = -1$ is the fluid 4-velocity, and the fluid isotropic pressure p and energy density μ obey a generic barotropic equation of state

$$p = p(\mu), \ \mu \neq 0, \ (\mu + p) \neq 0.$$
 (5)

Here the inequalities are to ensure that we are not considering a vacuum solution, and that u^a is a covariantly determined 4-vector: it is the unique timelike eigenvector of both T_{ab} and R_{ab} .

We prove the conjecture by considering a shear free perfect fluid that is rotating:

$$\omega^a \neq 0. \tag{6}$$

Since the vorticity is non-zero, we can decompose the instantaneous tangential rest space of an observer with respect to the vortex congruences and by doing so we get a series of geometrical results that force the expansion scalar to vanish.

II. SEMITETRAD 1+3 DESCRIPTION OF SPACETIME

We use the local 1+3 threading of spacetime manifold, using the unit timelike unit vector u^a along the fluid flow lines, as developed by Heckmann, Schücking, Ehlers [7], and Trumper, as summarised in [3] [4]. This formalism gives a clear covariant geometrical description of the matter evolution and a clear physical understanding of relativistic fluid flows. Then any tensor can be projected along the the fluid flow lines using the projection tensor

 $^{^{\}ast}$ goswami@ukzn.ac.za

[†] george.ellis@uct.ac.za

 $U_b^a = -u^a u_b$, while the tensor $h^a{}_b = g^a{}_b + u^a u_b$ projects onto the instantaneous 3-dimensional tangent rest spaces of the observers moving with 4-velocity u^a . This local 1+3 splitting then naturally defines two types of derivative operators. The vector u^a defines the *covariant time derivative* along the observers' worldlines (denoted by a dot) for any tensor $S^{a..b}{}_{c..d}$, given by

$$\dot{S}^{a..b}{}_{c..d} = u^e \nabla_e T^{a..b}{}_{c..d} \tag{7}$$

and the tensor h_{ab} defines the fully orthogonally projected covariant derivative D:

$$D_{e}S^{a..b}{}_{c..d} = h^{a}{}_{f}h^{p}{}_{c...}h^{b}{}_{g}h^{q}{}_{d}h^{r}{}_{e}\nabla_{r}S^{f..g}{}_{p..q} , \quad (8)$$

with total projection on all the free indices. Angle brackets denote orthogonal projections of vectors, and the orthogonally *projected symmetric trace-free* (PSTF) part of tensors: for example,

$$V^{\langle a \rangle} = h^{a}{}_{b}V^{b} , \ S^{\langle ab \rangle} = \left[h^{(a}{}_{c}h^{b)}{}_{d} - \frac{1}{3}h^{ab}h_{cd}\right]S^{cd} .$$
(9)

We can naturally define the 3 volume element of the instantaneous tangent rest spaces of the observers moving along the fluid flow lines as

$$\epsilon_{abc} = -\sqrt{|g|} \delta^0_{[a} \delta^1_b \delta^2_c \delta^3_{d]} u^d , \qquad (10)$$

that satisfies the following identities

$$\epsilon_{abc}\epsilon^{def} = 3!h^d_{[a}h^e_b h^f_{c]} , \quad \epsilon_{abc}\epsilon^{dec} = 2!h^d_{[a}h^e_{b]}. \tag{11}$$

The covariant derivative of the time-like vector u^a can now be decomposed into irreducible parts as

$$\nabla_a u_b = -\dot{u}_a u_b + \frac{1}{3} h_{ab} \Theta + \sigma_{ab} + \epsilon_{abc} \omega^c , \qquad (12)$$

where $\dot{u}_a \equiv \dot{u}_{\langle a \rangle} = u^b \nabla_b u^a$ is the acceleration, $\Theta = D_a u^a$ is the expansion, $\sigma_{ab} = D_{\langle a} u_{b \rangle}$ is the shear tensor and $\omega^a \equiv \omega^{\langle a \rangle} = \epsilon^{abc} D_b u_c$ is the vorticity vector. Similarly the Weyl curvature tensor can be decomposed irreducibly into Gravito-Electric and Gravito-Magnetic parts as

$$E_{ab} = C_{abcd} u^c u^d = E_{\langle ab \rangle}, \ E_{ab} u^b = 0, \ E^a{}_a = 0\,(13)$$

$$H_{ab} = \frac{1}{2} \epsilon_{acd} C_{be}^{cd} u^e = H_{\langle ab \rangle}, \ H_{ab} u^b = 0, \ H^a_{\ a} = 0 \, (14)$$

which allows a covariant description of tidal forces and gravitational radiation [2]. For perfect fluid matter obeying (4), (5), the vector field u^a is uniquely defined as the timelike eigenvector of T^{ab} . Thus the complete set of geometrical and thermodynamic variables that describe a perfect fluid spacetime to 1st order in derivatives of u^a and algebraically in R_{ab} and C_{abcd} is

$$\mathcal{D} = \{\Theta, \mu, p, \dot{u}^a, \omega^a, \sigma^{ab}, E^{ab}, H^{ab}\}.$$
 (15)

For brevity, we will use the following notation in this paper: for any 3-vector V^a and second rank 3-tensor A_{ab} ,

$$\operatorname{div} V = D_a V^a, \ (\operatorname{curl} V)_a = \epsilon_{abc} D^b V^c, \qquad (16)$$

$$(\operatorname{div} A)_a = D^b A_{ab}, (\operatorname{curl} A)_{ab} = \epsilon_{cd\langle a} D^c A^d_{b\rangle}.$$
 (17)

Also, the cross product of two 3-vectors X^a and Y^a is denoted by $[X, Y]^a$, where

$$[X,Y]^a = \epsilon^{abc} X_b Y_c , \qquad (18)$$

while that between a 3-vector V^a and a PSTF tensor A^{ab} is denoted as

$$[V,A]^{ab} = \epsilon^{cd\langle a} V_c A^{b\rangle}{}_d .$$
⁽¹⁹⁾

III. FIELD EQUATIONS FOR SHEAR FREE PERFECT FLUID

The Einstein Field Equations (EFE) are

$$G_{ab} \equiv R_{ab} - \frac{1}{2}Rg_{ab} = \kappa T_{ab} \tag{20}$$

where T_{ab} is the matter stress tensor and κ the gravitational constant. The Ricci identities for u^a are

$$\nabla_{[c}\nabla_{d]}u^{a} = 2R^{a}_{\ bcd}u^{b}.$$
 (21)

The once and doubly contracted Bianchi identities are

$$\nabla_c R_{ab} - \nabla_b R_{ac} + \nabla_d R^d_{\ abc} = 0, \qquad (22)$$

$$\nabla^a (R^b_{\ a} - \frac{1}{2} Rg^b_{\ a}) = 0 \Leftrightarrow T^{ab}_{\ ;a} = 0, \qquad (23)$$

where the equivalence follows from the EFE (20).

These equations can all be projected along the fluid flow lines and onto the instantaneous 3 dimensional tangential rest frames orthogonal to the flow lines, to give a set of propagation and constraint equations (see [6] [4]) for the variables in \mathcal{D} (see (15)).

For a shear-free barotropic perfect fluid $\sigma_{ab} = 0$, conditions (4), (5) hold, and the following set of equations that follow from (20)-(23) must be satisfied.

Propagation equations

The propagation equations are,

$$\dot{\Theta} - (\operatorname{div} \dot{u}) = -\frac{1}{3}\Theta^2 + \dot{u}_a \dot{u}^a + 2(\omega_a \omega^a) - \frac{1}{2}(\mu + 3p), \quad (24)$$

$$\dot{\omega}^{\langle a \rangle} = -\frac{2}{3}\Theta\omega^a + \frac{1}{2}(\operatorname{curl}\dot{u})^a , \qquad (25)$$

$$\dot{E}^{\langle ab\rangle} - (\operatorname{curl} H)^{ab} = -\Theta E^{ab} + 2[\dot{u}, H]^{ab} + [\omega, E]^{ab} , \quad (26)$$

$$\dot{H}^{\langle ab \rangle} + (\operatorname{curl} E)^{ab} = -\Theta H^{ab} - 2[\dot{u}, E]^{ab} + [\omega, H]^{ab} , \ (27)$$

$$\dot{\mu} = -\Theta(\mu + p) . \tag{28}$$

Constraint equations

The constraint equations are,

$$(C_1)^a \equiv \frac{2}{3} D^a \Theta - (\operatorname{curl} \omega)^a - 2[\dot{u}, \omega] = 0 , \qquad (29)$$

$$(C_2) \equiv (\operatorname{div} \omega) - \dot{u}^a \omega_a = 0 , \qquad (30)$$

$$(C_3)^{ab} \equiv H^{ab} + D^{\langle a}\omega^{b\rangle} + 2\dot{u}^{\langle a}\omega^{b\rangle} = 0 , \qquad (31)$$

$$(C_4)^a \equiv (\operatorname{div} E)^a - \frac{1}{3}D^a\mu - 3\omega_b H^{ab} = 0$$
, (32)

$$(C_5)^a \equiv (\operatorname{div} H)^a + (\mu + p)\omega^a + 3\omega_b E^{ab} = 0$$
, (33)

$$(\tilde{C}_6)^{ab} \equiv E^{ab} - D^{\langle a} \dot{u}^{b\rangle} - \dot{u}^{\langle a} \dot{u}^{b\rangle} + \omega^{\langle a} \omega^{b\rangle} = 0 , \quad (34)$$

$$(\tilde{C}_7)^a \equiv D^a p + (\mu + p) \dot{u}^a = 0.$$
 (35)

We note here three important points:

- 1. First of all, the above system of equations is not closed. To close the system we must supply a thermodynamic relation between fluid pressure and fluid energy density as an *equation of state*, which we specify to be (5) (see [7] and section V).
- 2. Secondly, the constraints C_i 's (29)-(33) are known to be consistently *time propagated* along u^a locally in the case of a general fluid flow, but are now specialised to the shear-free perfect fluid case. This gives two new constraints: \tilde{C}_6 , which is the shear constraint (2), and \tilde{C}_7 , which is the heat flux propagation equation restricted to the perfect fluid case.
- 3. Thirdly, although the spatial derivative operators D^a are orthogonal to u^a ($u^a D_a = 0$), because the vorticity is non-vanishing the local instantaneous rest frames of a family of observers do not mesh together to give 3-surfaces in spacetime orthogonal to u^a [7], but rather are just a collection of tangent planes. As the operators D^a do not span a three dimensional surface, their commutators acting on a scalar do not vanish (see (36)).

IV. IMPORTANT COMMUTATORS AND IDENTITIES FOR SHEAR FREE PERFECT FLUIDS

In this section, we list the important commutators and tensor identities that will be used in this paper. We note that these commutators and identities are specialised for shear free perfect fluid motion. For any scalar function 'f' the spatial derivative commutator can be written as

$$[D_a D_b - D_b D_a]f = 2\epsilon_{abc}\omega^c \dot{f} ,$$

$$\Leftrightarrow \epsilon^{abc} D_b D_c f = 2\omega^a \dot{f} .$$
(36)

and the *time-space commutator* is as follows

$$h^{a}_{b} \left[D^{b} f \right] = \left[D^{a} + \dot{u}^{a} \right] \dot{f} - \left[\frac{1}{3} \Theta h^{ab} + \epsilon^{abc} \omega_{c} \right] D_{b} f .$$
(37)

Similarly, the spatial derivative commutator for a projected 3-vector $V^{\langle a\rangle}$ is given as

$$D_{[a}D_{b]}V_{c} = \left[E_{c[a} + \frac{1}{3}\Theta\omega^{d}\epsilon_{cd[a} + \omega_{c}\omega_{[a} + \frac{1}{3}(\mu - \frac{1}{3}\Theta^{2} - 3\omega_{d}\omega^{d})h_{c[a}\right]V_{b]} + \left[h_{c[a}E_{b]d} - \frac{1}{3}\Theta h_{c[a}\epsilon_{b]de}\omega^{e} + h_{c[a}\omega_{b]}\omega_{d}\right]V^{d} + \epsilon_{abd}\omega^{d}\dot{V}_{\langle c\rangle} .$$
(38)

The curl of the cross product of two projected 3-vectors can be calculated using (11) to be

$$(\operatorname{curl}[X,Y])^a = D_b(X^a Y^b) - D_b(X^b Y^a)$$
. (39)

From the above relation we get the curl of the curl of a projected 3-vector as

$$(\operatorname{curl}\operatorname{curl} V)^a = D_b D^a V^b - D^2 (V^a) ,$$
 (40)

where $D^2 \equiv D^a D_a$ is the 3 dimensional Laplacian operator (which is meaningful even though no orthogonal 3-spaces exist in spacetime).

V. SHEAR FREE BAROTROPIC PERFECT FLUIDS AND VORTICITY

A barotropic perfect fluid (4) (5) is a useful model of fluid behaviour in a wide variety of scientific fields, from meteorology to astrophysics. In this case we can write

$$\dot{p} = c_s^2(\mu)\dot{\mu} , \ D^a p = c_s^2(\mu)D^a \mu ,$$
 (41)

where $c_s(\mu) \equiv \sqrt{(dp)/(d\mu)}$ is the local speed of sound (which is zero if p = 0). From (35), for a barotropic perfect fluid with $\mu + p \neq 0$ as assumed,

$$\dot{u}_c = -\left(\frac{D_c p}{\mu + p}\right) = -c_s^2(\mu) \left(\frac{D_c \mu}{\mu + p}\right).$$
(42)

Then the curl of the acceleration vector is given by

$$(\operatorname{curl} \dot{u})^a = -\epsilon^{abc} D_b \left(\frac{D_c p}{\mu + p}\right) ,$$
 (43)

which can be expanded as

$$(\operatorname{curl} \dot{u})^{a} = -\frac{\epsilon^{abc} D_{b} D_{c} p}{(\mu + p)} + \frac{\epsilon^{abc} D_{b} p D_{c} p}{(\mu + p)^{2}} + c_{s}^{2} \frac{\epsilon^{abc} D_{b} \mu D_{c} \mu}{(\mu + p)^{2}}$$
(44)

The last two terms vanish identically due to the antisymmetry of the 3-volume element. Using the commutator (36) and the propagation equation (28) this simplifies to

$$(\operatorname{curl} \dot{u})^a = -c_s^2 \frac{2\omega^a \dot{\mu}}{(\mu+p)} = 2c_s^2 \Theta \omega^a.$$
 (45)

The vorticity propagation equation (25) now simplifies to

$$\dot{\omega}^{\langle a \rangle} = \left(c_s^2 - \frac{2}{3}\right)\Theta\omega^a. \tag{46}$$

When a shear free barotropic perfect fluid is rotating, the projected time evolution of the vorticity vector is always parallel to the vorticity vector itself: it is Fermi propagated along the world line.

VI. FURTHER 1+2 DECOMPOSITION OF THE 3-SPACE USING THE VORTEX LINES

By (12) the vorticity vector is uniquely defined by the 4-velocity u^a , and so provides a preferred direction in the instantaneous 3-space of observers moving along the fluid flow lines. Thus this 3-space can be further decomposed with respect to the vortex vector direction e^a , defined by

$$\omega^a = \Omega e^a, \ e_a e^a = 1, \ \Omega \neq 0, \ e_a u^a = 0 \tag{47}$$

(the inequality, and the uniqueness of e_a , follow from (5), (6), and (12)). Then the projection tensor

$$N_a{}^b \equiv h_a{}^b - e_a e^b = g_a{}^b + u_a u^b - e_a e^b \qquad (48)$$

projects vectors onto tangential 2-spaces orthogonal to u^a and the vorticity vector which, following [10], we refer to as 2-sheets. It thus follows from this that

$$e^a N_{ab} = 0 = u^a N_{ab} , \quad N^a{}_a = 2 .$$
 (49)

Any spacetime 3-vector $\psi^a = h^a{}_b\psi^b$ can now be irreducibly split into a scalar, Ψ , which is the component along the vortex line and a 2-vector, Ψ^a , which is a 2-sheet component orthogonal to e^a , i.e.,

$$\psi^a = \Psi \, e^a + \Psi^{\bar{a}} \,\,, \tag{50}$$

where the bar on a particular index denotes projection with N_{ab} on that index. A similar decomposition can be done for a PSTF 3-tensor, ψ_{ab} , which can be split into scalar, 2-vector and 2-tensor parts as follows:

$$\psi_{ab} = \psi_{\langle ab \rangle} = \Psi\left(e_a \, e_b - \frac{1}{2}N_{ab}\right) + 2\Psi_{(a} \, e_{b)} + \Psi_{\{ab\}} ,$$
(51)

where the curly brackets denote the part of a tensor which is PSTF with respect to e^a . The tangential 2-sheet element determines a natural 2-volume element, the alternating Levi-Civita 2-tensor

$$\varepsilon_{ab} \equiv \epsilon_{abc} \, e^c = \eta_{dabc} \, e^c \, u^d \,, \tag{52}$$

induced by the volume element ϵ_{abc} of the tangential 3spaces. Note that this is a meaningful definition even though N_{ab} does not form 2-surfaces in spacetime. From the definition of ε_{ab} and N_{ab} , the following relations hold

$$\varepsilon_{ab} e^b = 0 = \varepsilon_{(ab)} , \qquad (53)$$

$$\epsilon_{abc} = e_a \ \varepsilon_{bc} - e_b \varepsilon_{ac} + e_c \varepsilon_{ab} \ , \tag{54}$$

$$\varepsilon_{ab}\,\varepsilon^{cd} = N_a{}^c N_b{}^d - N_a{}^d N_b{}^c \,, \tag{55}$$

$$\varepsilon_a{}^c \varepsilon_{bc} = N_{ab} , \qquad (56)$$

$$\varepsilon^{ab}\,\varepsilon_{ab} = 2\;.\tag{57}$$

Apart from the 'time' (dot) derivative of an object (scalar, vector or tensor), which is the derivative along the timelike congruence u^a , we introduce two new derivatives which the congruence e^a defines for any tensor $\psi_{a.b}^{c..d}$:

$$\hat{\psi}_{a..b}{}^{c..d} \equiv e^f D_f \psi_{a..b}{}^{c..d} , \qquad (58)$$

$$\delta_f \psi_{a..b}{}^{c..d} \equiv N_f{}^j N_a{}^l \cdots N_b{}^g N_h{}^c \cdots N_i{}^d D_j \psi_{l..g}{}^{h..i}(59)$$

The hat-derivative is the spatial derivative along vortex lines in the instantaneous rest frame of an observer, while the δ -derivative is the projected spatial derivative onto the tangent 2-sheet, with the projection on every free index. By these definitions, one obtains the following relations for the derivatives of the sheet-projection N_{ab} and the sheet volume element ε_{ab} :

$$\dot{N}_{ab} = 2u_{(a} \,\dot{u}_{b)} - 2e_{(a} \,\dot{e}_{b)} = 2u_{(a} \,\mathcal{A}_{b)} - 2e_{(a} \,\alpha_{b)}, (60)$$

$$N_{ab} = -2e_{(a} a_{b)} , (61)$$

$$\delta_c N_{ab} = 0 av{62}$$

$$\dot{\varepsilon}_{ab} = -2u_{[a}\,\varepsilon_{b]c}\,\mathcal{A}^c + 2e_{[a}\,\varepsilon_{b]c}\,\alpha^c \,\,, \tag{63}$$

$$\hat{\varepsilon}_{ab} = 2e_{[a}\,\varepsilon_{b]c}\,a^c \ , \tag{64}$$

$$\delta_c \varepsilon_{ab} = 0 , \qquad (65)$$

where we define the following 2-sheet quantities:

$$\mathcal{A}_a \equiv \dot{u}_{\bar{a}} , \alpha_a \equiv \dot{e}_{\bar{a}}, \ a_a = \hat{e}_{\bar{a}} . \tag{66}$$

With this decomposition relative to the vortex lines, the 4-acceleration splits as

$$\dot{u}^a = \mathcal{A} \, e^a + \mathcal{A}^a \,, \tag{67}$$

and the momentum equation (42) gives

$$\mathcal{A} = -\frac{\hat{p}}{(\mu+p)} = -c_s^2 \frac{\hat{\mu}}{(\mu+p)},$$
 (68)

and

$$\mathcal{A}^a = -\frac{\delta^a p}{(\mu+p)} = -c_s^2 \frac{\delta^a \mu}{(\mu+p)}.$$
 (69)

For the electric and magnetic part of the Weyl tensors we get

$$E_{ab} = \mathcal{E}\left(e_a \, e_b - \frac{1}{2}N_{ab}\right) + 2\,\mathcal{E}_{(a} \, e_{b)} + \mathcal{E}_{ab} \,\,, \qquad (70)$$

$$H_{ab} = \mathcal{H}\left(e_a e_b - \frac{1}{2}N_{ab}\right) + 2\mathcal{H}_{(a} e_{b)} + \mathcal{H}_{ab} . \quad (71)$$

Just like the covariant derivative of fluid 4-velocity can be irreducibly split into geometrical objects like expansion, shear, vorticity and acceleration, the projected 3derivative of the unit vector along the vortex lines can be irreducibly decomposed as

$$D_a e_b = e_a a_b + \frac{1}{2} \phi N_{ab} + \xi \varepsilon_{ab} + \zeta_{ab} , \qquad (72)$$

where analogous to the 1+3 splitting, we have defined

$$\phi = \delta_a e^a , \qquad (73)$$

$$\zeta_{ab} = \delta_{\{a} e_{b\}} , \qquad (74)$$

$$\xi = \frac{1}{2} \varepsilon^{ab} \delta_a e_b . \tag{75}$$

From the above equation we see that along the vortex lines, ϕ represents the spatial expansion of the sheet, ζ_{ab} is the shear (i.e., the spatial distortion of the sheet) and a_a its spatial acceleration, while ξ represents the spatial twist associated with the vortex lines; like Ω , it is a spacetime scalar that cannot be altered by any change of basis. Similar to the kinematical variables of u^a in the 1+3 approach (see (12)), the new kinematic variables a_a, ϕ, ξ , ζ_{ab} , \mathcal{A} and α_a are fundamental objects in spacetime, and their dynamics give us information about the spacetime geometry. Further to this, the complete set of the 1+3field equations (24-35) can be further decomposed into the equations along the vortex lines and those projected onto the 2-sheets. We refer to [10], corrected by [12], for the complete set of 1+1+2 evolution, propagation and constraint equations.

VII. GEOMETRIC RESULTS FOR ROTATING SHEAR FREE BAROTROPIC PERFECT FLUIDS

Having fixed the timelike vector of our 1+1+2 semitetrad frame along the fluid flow lines and the preferred spacelike vector along the vortex line, we are now in a position to prove some general geometric results for a rotating shear-free barotropic perfect fluid spacetime. We note that some of these results were already proved in [6], by constructing a complete orthonormal tetrad for dustlike matter.

Lemma 1. For a rotating shear-free barotropic perfect fluid, the projection of the time evolution of the unit vector along the vortex line onto the 2-sheet vanishes. In other words we must have $\alpha_a = 0$. *Proof.* The proof is straightforward from equation (46), which states that the projected time evolution of the vorticity vector on the instantaneous rest 3-space of an observer must be parallel to the vorticity. In other words we must have $\dot{\omega}^{\langle a \rangle} \equiv h^a_{\ b} \left[\Omega e^b \right]^{\cdot}$ parallel to e^a . But we know by (66) that

$$h^{a}_{\ b} \left[\Omega e^{b}\right] = \dot{\Omega} e^{a} + \Omega \alpha^{a}, \tag{76}$$

where the second term is the component in the 2-sheet and must vanish. Since Ω is strictly non zero by (47), this makes $\alpha_a = 0$.

The above Lemma has important consequences for a set of commutation relations, and a set of equations for derivatives.

Commutation relations When we decompose the scalar commutators (36,37) along the vortex lines and the 2-sheets, using the above Lemma, we get the following commutation relation for a scalar ψ ,

$$\hat{\psi} - \dot{\hat{\psi}} = -\mathcal{A}\,\dot{\psi} + \frac{1}{3}\Theta\hat{\psi},\tag{77}$$

$$\delta_a \dot{\psi} - (\delta_a \psi)_{\perp}^{\ } = -\mathcal{A}_a \, \dot{\psi} + \frac{1}{3} \Theta \delta_a \psi + \Omega \, \varepsilon_{ab} \delta^b \psi, \qquad (78)$$

$$\delta_a \hat{\psi} - (\delta_a \psi)_{\perp} = a_a \hat{\psi} + \frac{1}{2} \phi \, \delta_a \psi + (\zeta_{ab} + \xi \, \varepsilon_{ab}) \, \delta^b \psi(79)$$

$$\delta_{[a}\delta_{b]}\psi = \varepsilon_{ab}\left(\Omega\,\dot{\psi} - \xi\,\hat{\psi}\right) \ . \tag{80}$$

Here the symbol \perp denotes projection onto the 2-sheet.

A momentum relation Equation (69) shows

$$\frac{1}{2}\varepsilon^{ab}\delta_a\mathcal{A}_b = \frac{1}{2}\varepsilon^{ab}\frac{\delta_a\delta_bp}{(\mu+p)}.$$
(81)

The terms involving the δ -derivative of the denominator fall away because of the antisymmetry of ε^{ab} (just like (45)). Now using (80), (68) and (28) in (81), we have

$$\frac{1}{2}\varepsilon^{ab}\delta_a\mathcal{A}_b = \frac{\xi\hat{p} - \Omega\dot{p}}{(\mu + p)} = -\xi\mathcal{A} + \frac{2}{3}\Omega\Theta.$$
(82)

2-sheet relations From equation (80) it is clear that the 2-sheets are not genuine two surfaces but just a collection of tangent planes, as the sheet derivatives do not commute and have components along the fluid flow lines and possibly along the vortex lines (if $\xi \neq 0$). By contrast, the following theorem follows from these commutators (cf.[6] for the pressure-free case).

Theorem 1. For a rotating shear free barotropic perfect fluid the fluid flow lines and the vortex lines are always 2-surface forming.

Proof. The Lie bracket of the derivatives along the fluid flow lines and along the unit preferred spacelike vector along the vortex lines have no 2-sheet components. \Box

Derivative equations The vanishing of α_a results in the following set of derivative equations, which are free of any 2-sheet components:

$$\dot{\Omega} = \left(c_s^2 - \frac{2}{3}\right)\Omega\Theta , \qquad (83)$$

$$\hat{\Omega} = (\mathcal{A} - \phi)\Omega , \qquad (84)$$

$$\xi = -\frac{1}{3}\Theta\xi , \qquad (85)$$

$$\dot{\phi} = \frac{2}{3}\Theta\left(\mathcal{A} - \frac{1}{2}\phi\right) + 2\xi\Omega. \tag{86}$$

$$\hat{\Theta} = 3\xi\Omega . \tag{87}$$

Note that for the equation (86), we used the corrected equation (48) of [10] that was presented in [12].

The system of propagation equations (24)-(28) does not include an equation for the evolution of the acceleration \dot{u}^a , and so we do not have one for \mathcal{A} . To obtain one we take the time evolution of the momentum conservation equation (35), which depends on the equation of state. Imposing the barotropic equation of state (5) and using the commutator (77) for the pressure p, we get the evolution equation of \mathcal{A} as

$$\dot{\mathcal{A}} = -\mathcal{A}\Theta\left(\frac{1}{c_s^2}\frac{d^2p}{d\mu^2}(\mu+p) - c_s^2 + \frac{1}{3}\right) + 3c_s^2\xi\Omega. \quad (88)$$

We are now in a position to state and prove the main result of this paper:

Theorem 2. A shear free rotating barotropic perfect fluid must be expansion free: $\Theta = 0$.

Proof. The proof consists of two parts. In the first part we use standard separation of variables methods for the vorticity scalar, and derive resulting constraints. In the second part we use these constraints to prove the result.

Part1: It follows from (83)-(88) that the scalar Ω identically satisfies commutator relation (77) in the 2-surface spanned by the fluid flow lines and the vortex lines. With (83) and (84) this shows that the first order variation of this scalar in the [u, e] surface is independent of it's dependence on the tangent 2-sheet. On using standard separation of variables methods, Ω can be factorised:

$$\Omega = \Omega_1 \Omega_2 \neq 0, \, \delta^a \Omega_1 = 0, \, \Omega_2 = \Omega_2 = 0.$$
 (89)

The equations governing both the factors Ω_1 and Ω_2 can then be determined by the Ricci identities and commutation relations as follows.

The 2-sheet dependence factors out from the equations (83), (84) and (77) and we find

$$\hat{\dot{\Omega}}_1 - \dot{\hat{\Omega}}_1 = -\mathcal{A}\,\dot{\Omega}_1 + \frac{1}{3}\Theta\hat{\Omega}_1,\tag{90}$$

$$\dot{\Omega}_1 = \left(c_s^2 - \frac{2}{3}\right)\Omega_1\Theta, \qquad (91)$$

$$\hat{\Omega}_1 = (\mathcal{A} - \phi)\Omega_1 . \tag{92}$$

Furthermore putting $\psi = \Omega_1$ in (78) and (79), we get

$$\delta_a \dot{\Omega}_1 = -\mathcal{A}_a \dot{\Omega}_1 , \quad \delta_a \hat{\Omega}_1 = a_a \hat{\Omega}_1 , \qquad (93)$$

which gives the second order dependence of Ω_1 on the tangent 2-sheet variables. For the variable Ω_2 , the 2-vector part of shear divergence equation $(C_1)^a$ (eqn. (29)) becomes

$$\frac{2}{3}\frac{\delta_a\Theta}{\Omega_1} - \varepsilon_{ab}\delta^b\Omega_2 - 2\Omega_2\varepsilon_{ab}\mathcal{A}^b + \Omega_2\varepsilon_{ab}a^b = 0.$$
(94)

Also putting $\psi = \Omega_2$ in (78), (79) and (80) gives

$$\left(\delta_a \Omega_2\right)_{\perp}^{\cdot} = -\frac{1}{3} \Theta \delta_a \Omega_2 - \Omega \varepsilon_{ab} \delta^b \Omega_2 \,, \tag{95}$$

$$\left(\delta_a \Omega_2\right)_{\perp}^{\hat{}} = -\frac{1}{2} \phi \,\delta_a \Omega_2 - \left(\zeta_{ab} + \xi \,\varepsilon_{ab}\right) \delta^b \Omega_2, \quad (96)$$

$$\delta_{[a}\delta_{b]}\Omega_2 = 0. (97)$$

From equations (89-97), it is clear that the complete spacetime behaviours of both Ω_1 and Ω_2 are well determined by other geometrical variables and hence the factorisation (89) is well posed. Thus the dynamics now splits into those variables and derivatives \mathcal{D}_1 defined in or by the $\{u^a, \omega^b\}$ 2-surfaces, and those \mathcal{D}_2 defined in the orthogonal 2-sheets:

$$\mathcal{D}_1 = \{\mu, p, \Theta, \dot{u}^a, \Omega, \mathcal{A}\},$$
(98)

$$\mathcal{D}_2 = \{\mathcal{A}^a, a_a, \mathcal{H}^b, \xi, \phi, \zeta_{ab}\}.$$
(99)

Putting $\psi = \Omega_1$ in (80), the LHS vanishes identically due to (89), and RHS generates a new constraint:

$$C_8 \equiv \Omega \dot{\Omega}_1 - \xi \hat{\Omega}_1 = 0 . \tag{100}$$

Simplifying the above using (91) and (92) gives

$$C_9 \equiv \Omega\left(c_s^2 - \frac{2}{3}\right)\Theta - \xi(\mathcal{A} - \phi) = 0.$$
 (101)

Part2: We now look at the cases $c_s^2 \neq 2/3$ and $c_s^2 = 2/3$ separately.

Case 1: $c_s^2 \neq 2/3$ The constraint (101) must be consistently time propagated. Hence taking the derivative of the constraint along the fluid flow lines, using equations (83), (88), (86) (85) and simplifying using (101) we get a new constraint

$$C_{10} \equiv \xi \left[\dot{\Theta} - 3\xi^2 + F\Theta\xi \left(-\frac{3}{2}c_s^2\phi + \mathcal{A} \right) \right] = 0 , \quad (102)$$

where

$$F = \frac{2}{3} \frac{\left(-c_s^2 + \frac{2}{3} + \frac{1}{c_s^2} \frac{d^2 p}{d\mu^2} (\mu + p)\right)}{\left(c_s^2 - \frac{2}{3}\right)^2 \Omega} .$$
 (103)

This gives rise to two separate subcases:

- 1. $\xi = 0$. Using this in (101) immediately gives $\Theta = 0$ for the entire spacetime (as $\Omega \neq 0$ and $c_s^2 \neq 2/3$), and our result is proved.
- 2. $\xi \neq 0$. Then by (102), the expansion Θ obeys the evolution equation

$$\dot{\Theta} = 3\xi^2 + F\Theta\xi \left(-\frac{3}{2}c_s^2\phi + \mathcal{A}\right) . \tag{104}$$

Equation (104) and the Raychaudhuri equation (24) are independent equations for the evolution of the expansion of the fluid flow lines along themselves. The Raychaudhuri equation (24) is true for all shear-free fluids, whereas equation (104) is true for the special case of rotating shear-free barotropic perfect fluids (it holds when (5) and (6) are satisfied). Equations (24) and (104) must be compatible everywhere in that case.

We now check the dependence of the relevant set of linear equations on the quantity ξ . We do so by taking partial derivatives of these equations with respect to ξ , where the partial derivatives are not space time derivatives but derivatives in the space of variables $\{\mathcal{D}_1 \cup \mathcal{D}_2\}$ with coordinates given by (98), (99). Thus $\partial/\partial \xi$ means that in those equations, we keep constant all other variables in those sets; in particular, variables that are determined independent of the vector e^a . We make this explicit in the next few paragraphs.

Using the equations we have so far employed, the compatibility of (104) and (24) holds for arbitrary equations of state (5) such that $c_s^2 \neq 2/3$, including p = 0. For a barotropic fluid the sound speed is independent of any spatial frame choice, so

$$\partial c_s^2 / \partial \xi = 0 . \tag{105}$$

Since the kinematical variables related to the fluid flow lines u^a do not depend on the definition of e^a and $\omega^a \omega_a = \Omega^2$ is a scalar independent of e_a , we must also have

$$\frac{\partial \dot{u}^a}{\partial \xi} = \frac{\partial \Theta}{\partial \xi} = 0 = \frac{\partial \Omega}{\partial \xi} = \frac{\partial (\omega^a \omega_a)}{\partial \xi} .$$
(106)

The only way the Raychaudhuri equation (24) could depend on kinematical quantities related to the vortex line direction e^a is via the (div \dot{u}) and $\dot{u}_a \dot{u}^a$ terms. But these are scalars defined without any reference to that direction, so

$$\partial \dot{\Theta} / \partial \xi = 0. \tag{107}$$

Since the fluid flow lines are independent of the spatial vector e^a , $\partial u^a/\partial \xi = 0$ and the operator $\partial/\partial \xi$ commutes with the dot operator. Hence if $\partial \psi/\partial \xi = 0$ where ψ is any scalar, then $\partial \dot{\psi}/\partial \xi = 0$. Also, we know by definition that $\delta_{(a}e_{b)}$ and $\delta_{[a}e_{b]}$ are linearly independent tensors, and ϕ is the trace of the former while ξ is the only independent component of the later. Therefore we must have

$$\partial \phi / \partial \xi = \partial \dot{\phi} / \partial \xi = 0. \tag{108}$$

Now applying $\partial/\partial\xi$ to the constraint (101), we get

$$\frac{\partial C_9}{\partial \xi} \equiv (\mathcal{A} - \phi) + \xi \frac{\partial \mathcal{A}}{\partial \xi} = 0.$$
 (109)

Again applying (107) to (104), we have

$$6\xi - \frac{3}{2}c_s^2\phi F\Theta + F\Theta\left(\mathcal{A} + \xi\frac{\partial\mathcal{A}}{\partial\xi}\right) = 0 , \qquad (110)$$

which using (109) becomes,

$$C_{11} \equiv 6\xi - \frac{3}{2}F\Theta\left(c_s^2 - \frac{2}{3}\right)\phi = 0$$
 (111)

Applying $\partial/\partial\xi$ to this equation gives 6 = 0, which is a contradiction. Hence the subcase $\xi \neq 0$ is not possible by (101).

Therefore we have demonstrated that for a shear free barotropic perfect fluid with $c_s^2 \neq 2/3$, we must have $\xi = 0$ for the entire spacetime, which makes Θ vanish.

Case2: $c_s^2 = 2/3$. This is a curious case, where the sound speed is a constant ($c_s^2 = 2/3$) over the entire spacetime, and by equation (46) we have

$$\dot{\omega}^{\langle a \rangle} = 0 \Rightarrow \dot{\Omega} = \dot{\Omega}_1 = 0 \tag{112}$$

In other words, the vortex congruences are frozen in time for all tangent 3-spaces along a fluid flow line. The constraint (101), in this case, becomes

$$C_{12} \equiv \xi(\mathcal{A} - \phi) = 0. \tag{113}$$

This gives two subcases: either the first or second factor in that equation vanishes.

1. $\xi = 0$, $(\mathcal{A} - \phi) \neq 0$. If the spatial twist ξ vanishes in the entire spacetime we must have $\dot{\xi} = \hat{\xi} = 0$ and (85) is satisfied. Also in this case from equation (87), we have $\hat{\Theta} = 0$. Now the Ricci identity for the vector e^a can be written

$$R_{abc} \equiv 2\nabla_{[a}\nabla_{b]}e_c - R_{abcd}e^d = 0, \qquad (114)$$

Calculating the contraction $e^a \varepsilon^{bc} R_{abc}$ and putting $\hat{\xi} = 0 = \xi$, we get the constraint equation

$$\frac{1}{3}\Theta\Omega = -\frac{1}{2}\varepsilon_{ab}\delta^a a^b .$$
 (115)

Here the LHS only contains kinematical quantities related to the vector u^a , with no reference to e^a , while the RHS only contains kinematical quantities of the vector e^a (given u^a), This is only possible if both LHS and RHS are equal to the same spacetime constant k. This gives

$$\frac{1}{3}\Theta\Omega = k \tag{116}$$

Taking the hat derivative on both sides, we get

$$\Theta\Omega + \Theta\Omega = 0 \Rightarrow \Theta(\mathcal{A} - \phi)\Omega = 0 \tag{117}$$

by (84) and (87). As $\Omega \neq 0$ and $(\mathcal{A} - \phi) \neq 0$ in this case, we find $\Theta = 0$ for the entire spacetime.

2. $(\mathcal{A} - \phi) = 0$. If this case using equation (108), we have

$$\frac{\partial(\mathcal{A} - \phi)}{\partial \xi} = \frac{\partial \mathcal{A}}{\partial \xi} = 0.$$
 (118)

However by (67),

$$\mathcal{A}^2 = \dot{u}^a \dot{u}_a - \mathcal{A}^a \mathcal{A}_a. \tag{119}$$

From (118) and (106), this gives

$$\mathcal{A}^a \frac{\partial \mathcal{A}_a}{\partial \xi} = 0.$$
 (120)

We consider in turn the options $\{\mathcal{A}^a = 0\}$ and $\{\mathcal{A}^a \neq 0, \partial \mathcal{A}_a / \partial \xi = 0\}.$

A) Setting $\mathcal{A}^a = 0$ in (82) gives

$$-\xi \mathcal{A} + \frac{2}{3}\Omega \Theta = 0. \qquad (121)$$

Taking $\partial/\partial\xi$ of this equation gives $\mathcal{A} = 0$; substituting back gives $\Omega \Theta = 0 \Rightarrow \Theta = 0$, as desired.

B) Assuming $\{\mathcal{A}^a \neq 0, \partial \mathcal{A}_a / \partial \xi = 0\}$, take $\partial / \partial \xi$ of (82) and using (118) gives $\mathcal{A} = 0$. This is the case when the vorticity and acceleration vectors are orthogonal at an instant. Then (88) gives

$$3c_s^2 \xi \Omega = 0 \Rightarrow \xi = 0 \tag{122}$$

which is the condition that this remains true. Then in this special case, we have

$$\mathcal{A}^a \neq 0, \, \xi = \phi = \mathcal{A} = \hat{\Omega} = \dot{\Omega} = \hat{\Theta} = 0.$$
 (123)

As $p = (2/3)\mu$, from equation (68) we also have

$$\hat{p} = \hat{\mu} = 0 \tag{124}$$

and the scalar μ obeys the commutator relation (77) in the [u, e] plane identically. Furthermore, since $\dot{\mu}$ does not contain any 2-sheet term by (28), we can do a similar separation of variables for the scalar μ as we did above for Ω :

$$\mu = \mu_1 \mu_2 \neq 0, \, \delta^a \mu_1 = 0, \, \dot{\mu}_2 = \dot{\mu}_2 = 0.$$
 (125)

Then we immediately have from (28),

$$\dot{\mu}_1 = -\frac{5}{3}\Theta\mu_1 \ . \tag{126}$$

Putting $\psi = \mu_1$ in (80) we find $\dot{\mu}_1 = 0$ (as $\Omega \neq 0$) from (126), which again gives $\Theta = 0$ as desired.

We note that the theorem in the case $c_s^2 = 2/3$ is included in those proven in [13], and the case $\mathcal{A} = 0$ in [11].

Therefore for both cases $c_s^2(\mu) \neq 2/3$ and $c_s^2(\mu) = 2/3$ we have shown we must have $\Theta = 0$ in the entire spacetime, which completes the proof.

Note 1: The backbone of the proof is as follows.

- 1. By introducing the variables \mathcal{D} listed in (15), the equations (24)-(35) we use as the basis of our proof only involve first order linear differential operators.
- 2. The barotropic condition (5) gives the acceleration equation (42), from which the vorticity propagation equation (46) follows. That relation underlies Lemma 1 which then give the simplified commutation relations (77)-(80), the first order differential equations (83)-(88), and Theorem 1.
- 3. In Theorem 2, we identify an important geometric property of the vorticity scalar Ω , that is it's first order variation of this scalar on the [u, e] surface is independent of it's dependence on the tangent 2sheet. This prompted us to perform the usual separation of variables for the vorticity scalar as given in (89) and using the commutators and the Ricci identities of the vector u^a , we demostrated that such separation of variables is indeed well posed.
- 4. The third step gave rise to an independent constraint (101) that must be consistently time propagated. This consistency condition constrained the expansion to vanish in the entire spacetime.

Note 2: The key point is that the shear evolution equation is given by the combination of the Ricci identities for e^a and Raychaudhuri. In our analysis, we have used $\dot{\xi}$ and $\hat{\xi}$ equations, which are the different contractions of the Ricci identities of e^a . So effectively we are using the shear evolution constraint (34)

Three direct corollaries to Theorem 2 are as follows.

Corollary 1. For a shear free motion of a barotropic perfect fluid, the vorticity vector ω^a is Fermi propagated along the fluid flow lines, and its magnitude is constant.

Proof. From (46) and $\Theta = 0$,

$$\dot{\omega}^{\langle a \rangle} = (\operatorname{curl} \dot{u})^a = 0. \tag{127}$$

Then
$$(\Omega^2) = 2\omega_a \dot{\omega}^{\langle a \rangle} = 0.$$

Corollary 2. If a barotropic perfect fluid is rotating in a shear free way, then the vortex lines must be twist free.

Proof. From equation(101), if $\Theta = 0$, then

$$\xi(\mathcal{A} - \phi) = 0. \tag{128}$$

There are two cases.

- 1. $\xi = 0$, and the result is proved,
- 2. $A = \phi$, which must be true for all times. Then using (86) $\dot{\mathcal{A}} = 2\xi\Omega$ so (88) shows $\xi = 0$ or $2\Omega = 3c_s^2\Omega$, which is $c_s^2 = 2/3$ as $\Omega \neq 0$. If $c_s^2 = 2/3$, then from the equation (123) and the discussion after equation (121), we have $\mathcal{A} = 0$ for the entire spacetime. This makes $\dot{\mathcal{A}} = 0$. Using equation (88) then gives $\xi = 0$ (as $\Omega \neq 0$).

This proves the stated result.

Corollary 3. A shear free and rotating barotropic perfect fluid spacetime must be stationary.

Proof. To show this, following [7], we look for a vector

$$K_a = e^{U(x^i)} u_a, \tag{129}$$

in the direction of u^a , such that Killing's equations hold:

$$K_{a;b} + K_{b;a} = 0. (130)$$

Equation (129) gives

$$K_{a;b} = e^{U(x^{i})}(u_{a}U_{;b} + u_{a;b}).$$
(131)

Contracting (130) with $u^a u^b$, $u^a h^b_c$ and $h^a_c h^b_d$ and using (131), the Killing equations (130) are

$$u^{a}u^{b}(u_{a}U_{;b}+u_{a;b})=0, \qquad (132)$$

$$u^{a}h^{b}_{c}(u_{a}U_{;b} + u_{a;b} + u_{b}U_{;a} + u_{b;a}) = 0, \qquad (133)$$

$$h_c^a h_d^b (u_a U_{;b} + u_b U_{;a} + u_{a;b} + u_{b;a}) = 0.$$
(134)

First $u^a u_a = -1 \Rightarrow u^a u_{a;b} = 0$ so (132) is just

$$\dot{U} = 0. \tag{135}$$

Second, on using (135) and $u_a \dot{u}^a = 0$, (133) is

$$\dot{u}_c = h_c^b U_{;b} \,. \tag{136}$$

Now \dot{u}_a is determined by (35) where in the case of a barotropic perfect fluid with $\Theta = 0$, (28) and (41) show

$$\{\dot{\mu} = 0\} \Rightarrow \{\dot{p} = c_s^2 \dot{\mu} = 0\}.$$
 (137)

Using comoving coordinates $\{x^i\} = \{t, x^{\nu}\}, (x^{\nu}) = 0$ (see [4]) and (137), determine the function $W(x^{\nu})$ by integrating the time independent equation

$$\frac{\partial W(x^{\nu})}{\partial x^{\nu}} = w(p(x^{\nu}))\frac{\partial \mu(x^{\nu})}{\partial x^{\nu}}, \qquad (138)$$

$$w(\mu(x^{\nu})) := -\frac{(dp(x^{\nu})/d\mu(x^{\nu}))}{\mu(x^{\nu}) + p(\mu(x^{\nu}))}$$
(139)

in a surface $\{t = const\}$. This is always integrable because (138) implies $\partial^2 W / \partial x^{\mu} \partial x^{\nu} = \partial^2 W / \partial x^{\nu} \partial x^{\mu}$. Then (42) becomes

$$\dot{u}_c = h_c^b W_{;b}, \ \dot{W} = 0 \tag{140}$$

(which includes $c_s^2(\mu) = 0$ and so p = 0 as special cases). Thus because we are considering a barotropic fluid there is an acceleration potential (see [7]), and (135), (136) are satisfied with U = W.

Finally (134) is

$$\sigma_{ab} + \frac{1}{3}\Theta h_{ab} = 0 \tag{141}$$

which is true by Theorem 2.

Equations (132)-(134) are all satisfied. Consequently $K_a = e^{W(x^i)}u^a$ satisfies Killing's equations, the fluid flow lines are a Killing direction, and the spacetime must be stationary.

VIII. DISCUSSION

We have extended the result (3) to include the case of a generic barotropic perfect fluid (4), (5). A series of points arise.

1. The proof required only the inequalities $\mu \neq 0$, $\mu + p \neq 0$, not an energy condition like $\mu + p > 0$, and holds for all values of the speed of sound c_s^2 (thus including pressure-free matter, when $c_s^2 = 0$).

2. An interesting question is whether the result is true in the case of a scalar field $\Phi(x^j)$ satisfying the Klein Gordon equation, where $\dot{\Phi} \neq 0$ so that $\mu \neq 0, (\mu + p) \neq 0$. We conjecture it will be true in that case also.

3. This theorem is an exact result following from the Einstein Field Equations (20) and the shear-free condition. It is an open question if it holds approximately when the shear is and remains almost zero.

4. This result has interesting consequences for the Newtonian limit of General Relativity, see [9], because there is no corresponding result in Newtonian gravitational theory but Theorem 2 is an exact result applying even to quasi-Newtonian geometries.

Note that self-similar solutions are shear-free. Thus the result applies to all GR self similar solutions for barotropic perfect fluids (including pressure free matter), but not to Newtonian self-similar solutions for such matter.

5. There is a hidden symmetry in these equations, with the following result:

Lemma 2. The result remains true if the cosmological constant $\Lambda \neq 0$.

Proof. If we add a cosmological constant Λ by making the change

$$p \to p' = p + \Lambda, \ \mu \to \mu' = \mu - \Lambda, \ \nabla_a \Lambda = 0$$
 (142)

then

$$\mu + p \to (\mu + p)' = \mu + p \tag{143}$$

and (4) holds but with p changed by an additive constant, while (5) remains true. Then by (35) this does not alter the fluid acceleration, while (28) leaves $\dot{\mu}$ unchanged, and (32) and (33) remain true. The only one of the propagation and constraint equations (24)-(35) that is affected is the Raychaudhuri equation (24), but we only used a derivative of that equation in the above proof and Λ will not enter that derivative.

ACKNOWLEDGMENTS

RG thanks the South African National Research Foundation (NRF) for support.

GE thanks the UCT Research Committee (URC) for support.

We thank Norbert Van den Bergh for pointing out an important loophole in a previous draft, and Muzikayise Sikhonde for correcting a sign error in an earlier version.

Date: 2021/09/17

- N Van den Bergh, N., and R Slobodeanu (2016) "Shearfree perfect fluids with a barotropic equation of state in general relativity: the present status" *Classical and Quantum Gravity* 33:085008.
- [2] R Goswami and G F R Ellis (2021) "Tidal forces are gravitational waves". *Classical and Quantum Gravity* 38:085023.
- [3] G F R Ellis (2009) "Republication of Relativistic Cosmology" Gen Rel Grav 41: 581-660.
- [4] G F R Ellis and H van Elst (1999) "Cosmological Models", in *Theoretical and Observational Cosmology*, Ed. M Lachze-Rey, (Dordrecht: Kluwer), 1. [arXiv:gr-qc/9812046].
- [5] K Gödel (1952) "Rotating Universes in General Relativity Theory" Proceedings of the International Congress of Mathematicians, Cambridge, Mass., 1950, vol. 1, 175-181 (Amer. Math. Soc., R.I.).
- [6] G F R Ellis (1967) "The dynamics of pressure-free matter in general relativity", J. Math. Phys. 8, 1171-1194.

- [7] J Ehlers (1993) "Republication of Contributions to the relativistic mechanics of continuous media." Gen Rel Grav 25: 1225-1266.
- [8] J M M Senovilla, C F Sopuerta, and P Szekeres (1998) "Theorems on shear-free perfect fluids with their Newtonian analogues" *Gen. Rel. Grav.* **30**, 389-411 [arXiv:gr-qc/9702035v1].
- [9] G F R Ellis (2011) "Shear free solutions in general relativity theory" Gen Rel Grav 43:3253-3268.
- [10] C Clarkson (2007) "Covariant approach for perturbations of rotationally symmetric spacetimes", *Phys. Rev. D* 76, 104034.
- [11] M E Sikhonde and P K S Dunsby (2017) "Reviving the shear free perfect fluid conjecture in general relativity", Class. Quant. Grav. 34: 245007.
- [12] C Hansraj, R Goswami and S D Maharaj (2021) "A semi-tetrad decomposition of the Kerr spacetime", arXiv:2109.04162 (gr-qc).
- [13] R Slobodeanu (2014) "Shear-free perfect fluids with linear equation of state" Class. Quantum Grav 31 125012.