

PROFILE DECOMPOSITION IN SOBOLEV SPACES AND DECOMPOSITION OF INTEGRAL FUNCTIONALS II: HOMOGENEOUS CASE

MIZUHO OKUMURA

ABSTRACT. The present paper is devoted to a theory of profile decomposition for bounded sequences in *homogeneous* Sobolev spaces, and it enables us to analyze the lack of compactness of bounded sequences. For every bounded sequence in homogeneous Sobolev spaces, the sequence is asymptotically decomposed into the sum of profiles with dilations and translations and a double suffix residual term. One gets an energy decomposition in the homogeneous Sobolev norm. The residual term becomes arbitrarily small in the critical Lebesgue or Sobolev spaces of lower order, and then, the results of decomposition of integral functionals are obtained, which are important strict decompositions in the critical Lebesgue or Sobolev spaces where the residual term is vanishing.

CONTENTS

1. Introduction	2
1.1. Prologue	2
1.2. Notation and settings	3
1.3. Strategy of the paper	5
1.4. Organization of the paper	5
2. Profile decomposition in homogeneous Sobolev spaces	5
2.1. Settings	5
2.2. Main theorem	8
2.3. Proof of Theorem 2.7	11
2.4. Relations among profile decompositions in inhomogeneous and homogeneous Sobolev spaces	20
3. Decomposition of integral functionals in critical cases	23
3.1. Brief Summary	23
3.2. Main theorems	24
3.3. Proof of Theorem 3.1	26
3.4. Proof of Theorem 3.2	29
Acknowledgment	31
Appendix A. Abstract theory of profile decomposition	31
Appendix B. Abbreviated form of profile decomposition	32
Appendix C. Preliminaries	34

2020 *Mathematics Subject Classification.* 46B50, 46B06, 46E35.

Key words and phrases. Profile decomposition, decomposition of integral functionals, lack of compactness, concentration-compactness, Brezis-Lieb lemma, G -weak convergence, dislocation space, G -complete continuity.

1. Introduction

1.1. Prologue

The lack of compactness of bounded (mainly function) sequences in infinite dimensional normed vector spaces has been a significant and critical difficulty in mathematical analysis. To overcome the difficulty, a lot of researchers have been establishing a variety of ways to discuss the lack of compactness and the recovery of compactness. Classical and well-known results are, for instance, the Brezis-Lieb lemma [4], the concentration-compactness principles by Lions [9, 10, 11, 12], the global compactness results by Struwe [16], Brezis-Coron [3] and Bahri-Coron [2], and so on.

Profile decomposition is a way of asymptotic analysis of general bounded sequences that may have the defect of compactness, and originated in attempts to give an asymptotic decomposition of bounded sequences in some function spaces such as Lebesgue spaces, Sobolev spaces and so on. It states that every bounded sequence in a certain Banach space is asymptotically decomposed into a sum of *moving* profiles and a residual term, where that *movements* of profiles are considered to be descriptions of the defect of compactness.

Roughly speaking, there are two types of principles of profile decomposition in terms of the residual term: one with a single-suffix residual term as is treated in, e.g., Solimini [14], Tintarev-Fieseler [19] and Tintarev [18]; another with a double-suffix residual term as is treated in, e.g., Gérard [6], Jaffard [8] and Bahouri-Cohen-Koch [1]. The differences and relations between the above two types of principles are described in, e.g., [5, 13]. Most profile decomposition theorems offered in the above literature focus on isometric group actions on function spaces, named “dislocations”, which fairly prevents bounded sequences from converging strongly. By making use of such group actions appropriately, generic principles of profile decomposition have been successfully established.

Here we briefly review the results of Solimini and Jaffard. Solimini [14] developed the profile decomposition with a single-suffix residual term in the following sense. For any bounded sequence (u_n) in $\dot{W}^{1,p}(\mathbf{R}^N)$ ($1 < p < N$), there exist $w^l \in \dot{W}^{1,p}(\mathbf{R}^N)$ ($l \in \mathbf{Z}_{\geq 0}$) and $(y_{l,n}, j_{l,n}) \in \mathbf{R}^N \times \mathbf{R}$ ($l, n \in \mathbf{Z}_{\geq 0}$) such that, up to a subsequence,

$$(1.1) \quad u_n = \sum_{l=0}^{\infty} 2^{j_{l,n}N/p^*} w^l(2^{j_{l,n}}(\cdot - y_{l,n})) + r_n, \quad n \in \mathbf{Z}_{\geq 0},$$

with

$$\begin{aligned} 2^{-j_{l,n}N/p^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) &\rightarrow w^l \text{ weakly in } \dot{W}^{1,p}(\mathbf{R}^N), \\ |j_{l,n} - j_{k,n}| + 2^{j_{k,n}} |y_{l,n} - y_{k,n}| &\rightarrow \infty \text{ if } l \neq k \text{ (} n \rightarrow \infty \text{),} \\ \lim_{n \rightarrow \infty} \|r_n\|_{L^{p^*,q}(\mathbf{R}^N)} &= 0, \quad q > p^*, \\ \limsup_{n \rightarrow \infty} \|u_n\|_{\dot{W}^{1,p}(\mathbf{R}^N)}^p &\geq \sum_{l=0}^{\infty} \|\phi^l\|_{\dot{W}^{1,p}(\mathbf{R}^N)}^p, \end{aligned}$$

where $p^* := pN/(N - p)$ denotes the Sobolev critical exponent and $L^{p^*,q}(\mathbf{R}^N)$ denotes the Lorentz space.

On the other hand, Jaffard [8] developed the profile decomposition with a double-suffix residual term in the following sense. For any bounded sequence (u_n) in $\dot{H}^{s,p}$, there exist $w^l \in \dot{H}^{s,p}(\mathbf{R}^N)$ ($l \in \mathbf{Z}_{\geq 0}$) and $(y_{l,n}, j_{l,n}) \in \mathbf{R}^N \times \mathbf{R}$ ($l, n \in \mathbf{Z}_{\geq 0}$) such that, up to a subsequence,

$$(1.2) \quad u_n = \sum_{l=0}^L 2^{j_{l,n}N/p^*} w^l(2^{j_{l,n}}(\cdot - y_{l,n})) + r_n^L, \quad L, n \in \mathbf{Z}_{\geq 0},$$

with

$$\begin{aligned} |j_{l,n} - j_{k,n}| + 2^{j_{k,n}} |y_{l,n} - y_{k,n}| &\rightarrow \infty \text{ if } l \neq k \text{ (} n \rightarrow \infty \text{),} \\ \lim_{L \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|r_n^L\|_{L^{p_s^*}(\mathbf{R}^N)} &= 0, \\ \limsup_{n \rightarrow \infty} \|u_n\|_{\dot{H}^{s,p}(\mathbf{R}^N)}^p &\geq \sum_{l=0}^{\infty} \|\phi^l\|_{\dot{H}^{s,p}(\mathbf{R}^N)}^p. \end{aligned}$$

For more details of profile decomposition and applications of it, we refer the reader to [1, 6, 8, 13, 14, 17, 18, 19] and references therein. Also the relations between profile decomposition and classical ways of analysis of the lack of compactness are shown in [13].

The theory of Solimini and its extensions to Banach and Hilbert spaces by Tintarev et al. are successful in characterizing the profiles w^l with u_n by the use of the weak convergence. However, the well-definedness or the convergence of the infinite sum on (1.1) matter delicately and need some complicated arguments for verification, named “*routing procedure*” in [5]. On the other hand, in the theory of Gérard, Jaffard and Bahouri et al. appears the finite sum of scaled profiles (also called *dislocated profiles*) and the well-definedness and convergence do not matter. Also, Jaffard [8, p.386] remarks that the finite sum on (1.2) cannot be replaced with the infinite sum as in (1.1) because it does not usually converge.

Aim of the paper. Under these circumstances, in [13] the author developed a profile decomposition theorem in *inhomogeneous* Sobolev spaces, which was a *hybrid type* profile decomposition of [14, 18, 19] and [1, 6, 8]. In his theory, each profile of a bounded sequence is well characterized by the weak convergence and an isometric group action on inhomogeneous Sobolev spaces, that is, *translations* as in [14, 18, 19], but only a finite sum appears so that the well-definedness and convergence are always valid as in [1, 6, 8]. He also investigated significant results of decomposition of integral functionals (also regarded as the iterated Brezis-Lieb lemma and investigated by, e.g., [19, 18]), revealing that the profile decomposition leads to an asymptotic strict decomposition of integral functionals in suitable Lebesgue or Sobolev spaces where the residual term of profile decomposition is vanishing.

In the present paper, we shall develop a “*homogeneous*” version of a theory of profile decomposition and decomposition of integral functionals in contrast to [13], proving them in the same spirit. As for the energy decomposition in the Sobolev norm (see (2.2) below), it will be a sharper version of the ordinary energy decompositions provided by precursors which do not include the “residual term”. Results provided below will be well applied to studies of PDEs and Calculus of Variations.

1.2. Notation and settings

As is mentioned above, the way of profile decomposition needs a suitable setup of group actions responsible for the lack of compactness. Those group actions are called “*dislocations*” and they are built up from functional analytic viewpoints inspired by [18, 19].

Throughout the paper, we often use the following notation.

- (i) We write $(n_j) \preceq (n)$ when the left-hand side is a subsequence of the right-hand side.
- (ii) The set of integers greater than or equal to $l \in \mathbf{Z}$ is denoted by $\mathbf{Z}_{\geq l}$.
- (iii) For $\Lambda \in \mathbf{N} \cup \{0, +\infty\}$, the set of integers at least zero and at most Λ is denoted by $\mathbf{Z}_{\geq 0}^{\leq \Lambda+1}$, where we assume that $\Lambda + 1 = +\infty$ when $\Lambda = +\infty$, i.e., $\mathbf{Z}_{\geq 0}^{\leq \Lambda+1} = \{0, \dots, \Lambda\}$ if $\Lambda < +\infty$ and $\mathbf{Z}_{\geq 0}^{\leq \Lambda+1} = \mathbf{Z}_{\geq 0}$ if $\Lambda = +\infty$. Moreover, for real numbers $(a_L)_L$, we often write $\lim_{L \rightarrow \Lambda} a_L = a_\Lambda$ whenever $\Lambda < \infty$.
- (iv) We denote by C a non-negative constant, which does not depend on the elements of the corresponding space or set and may vary from line to line.
- (v) For an exponent $p \in [1, \infty]$, we denote by p' the Hölder conjugate exponent of p : $p' = p/(p-1)$ if $p \in]1, \infty[$; $p' = \infty$ if $p = 1$; $p' = 1$ if $p = \infty$.
- (vi) For a normed space X , we denote its norm by $\|\cdot\|$ or $\|\cdot\|_X$, denote its dual space by X^* and denote the duality pairing between X, X^* by $\langle \cdot, \cdot \rangle$ or $\langle \cdot, \cdot \rangle_X$.
- (vii) For a normed space X , we denote by $B_X(r)$ the closed ball of radius $r > 0$ centered at the origin, that is, $B_X(r) = \{u \in X; \|u\|_X \leq r\}$. Meanwhile, the closed ball in \mathbf{R}^N of radius $r > 0$ centered at $p \in \mathbf{R}^N$ is denoted by $B(p, r)$, that is, $B(p, r) := \{x \in \mathbf{R}^N; |x - p| \leq r\}$.
- (viii) For a normed space X , we denote by $\mathcal{B}(X)$ the normed space of all bounded linear operators on X equipped with the operator norm.
- (ix) For a Banach space X and for a bounded operator $T \in \mathcal{B}(X)$, the adjoint operator of T is denoted by T^* .

We shall follow and employ the setup for the framework of profile decomposition as in [13], and we here recall them briefly. Let $(X, \|\cdot\|_X)$ be a Banach space and let $G \subset \mathcal{B}(X)$ be a group (under operator composition) of bijective isometries.

- Definition 1.1.**
- (i) (*Operator convergence*) For a sequence (g_n) in $\mathcal{B}(X)$, the operator-strong convergence of (g_n) is defined as the pointwise strong convergence in X , while the operator-weak convergence of (g_n) is defined as the pointwise weak convergence in X .
 - (ii) (*G-weak convergence*) A sequence (u_n) in X is said to be G -weakly convergent to $u \in X$ provided that

$$\limsup_{n \rightarrow \infty} \sup_{g \in G} |\langle \phi, g^{-1}(u_n - u) \rangle| = 0 \quad \text{for all } \phi \in X^*.$$

- (iii) (*Dislocation group*) A group G of linear bijective isometries is called a *dislocation group* if the following two conditions are satisfied.
 - (a) For any sequence (g_n) in G with $g_n \not\rightarrow 0$, there exists a subsequence $(n_j) \preceq (n)$ such that (g_{n_j}) is convergent operator-strongly.
 - (b) For any sequence (g_n) in G with $g_n \not\rightarrow 0$ and for any sequence (u_n) in X with $u_n \rightarrow 0$ weakly in X , there exists a subsequence $(n_j) \preceq (n)$ such that $g_{n_j} u_{n_j} \rightarrow 0$ weakly in X .
- (iv) (*Dislocation space*) The pair (X, G) is called a *dislocation space* when G is a dislocation group.
- (v) (*G-complete continuity*) For a normed space Y , a linear operator $T : X \rightarrow Y$ is said to be G -completely continuous if every G -weakly convergent sequence in X is mapped to a strongly convergent sequence in Y .

Regarding the above settings, we shall make some remarks as in [13]:

- Remark 1.2.**
- (i) If $G = \{\text{Id}_X\}$, then the G -weak convergence is nothing but the ordinary weak convergence, i.e., the G -weak convergence is an extension of the weak convergence.
 - (ii) Suppose, in addition, that G satisfies that for every sequence (g_n) in G with $g_n \not\rightarrow 0$, the adjoint (g_n^*) has an operator-strongly convergent subsequence. Then the condition (b) in (iii) of Definition 1.1 above is always satisfied.
 - (iii) If $(X, \|\cdot\|_1)$ is equipped with another equivalent and complete norm $\|\cdot\|_2$, if the action of G on X is isometric in both $\|\cdot\|_1$ and $\|\cdot\|_2$, and if $((X, \|\cdot\|_1), G)$ is a dislocation space, then the pair $((X, \|\cdot\|_2), G)$ is also a dislocation space. In other words, the definition of dislocation spaces is irrelevant to the choice of equivalent norms.
 - (iv) The G -complete continuity is also called G -cocompactness in, e.g., [18] and some Tintarev's papers.

For more details of properties or examples of dislocation spaces or G -complete continuity, we refer the reader to [13, 18, 19] and references therein.

1.3. Strategy of the paper

As is developed in [13], we here briefly review a *recipe* of a theory of profile decomposition and decomposition of integral functionals. To construct the theory, one should prepare a pair (X, G, Y) where (X, G) is a dislocation Sobolev space which is embedded into some Lebesgue or Sobolev space Y G -completely continuously. The profile decomposition in (X, G) is obtained in three steps: (i) finding profiles by the use of [13, Theorem 2.1] (which will be exhibited in Appendix A below); (ii) a decomposition in the Sobolev norm and the exactness condition of profile decomposition; (iii) vanishing of the residual term in Y . The spirit of decomposition of integral functionals is: they are developed in Lebesgue or Sobolev spaces Y into which (X, G) is embedded G -completely continuously. Here the essential difficulty and difference in homogeneous case compared to the inhomogeneous case [13] are calculations in the energy decomposition in the homogeneous Sobolev norm. To this end, we shall employ an appropriate separation of the domain (see Lemmas 2.11, 2.14 and 2.15).

1.4. Organization of the paper

In Section 2, we shall establish a theorem of profile decomposition in homogeneous Sobolev spaces, which is our main interest of the paper. In Section 3, the results of decomposition of integral functionals subordinated to the profile decomposition in homogeneous Sobolev spaces will be discussed. Results and their proofs will be given in the same sections.

2. Profile decomposition in homogeneous Sobolev spaces

2.1. Settings

We begin with the definition of *homogeneous* Sobolev spaces. Let Ω be an open set in \mathbf{R}^N . For $1 \leq p < N/m$, $m, N \in \mathbf{N}$, the *homogeneous* Sobolev space $\dot{W}^{m,p}(\Omega)$ is defined by

$$\dot{W}^{m,p}(\Omega) = \overline{C_c^\infty(\Omega)}^{\|\cdot\|_{\dot{W}^{m,p}(\Omega)}},$$

where

$$\|u\|_{\dot{W}^{m,p}(\Omega)} = \left(\sum_{|\alpha|=m} \|\partial^\alpha u\|_{L^p(\Omega)}^p \right)^{1/p}.$$

Here for a multi-index $\alpha = (\alpha_1, \dots, \alpha_N) \in (\mathbf{Z}_{\geq 0})^N$,

$$|\alpha| = \sum_{n=1}^N \alpha_n \quad \text{and} \quad \partial^\alpha = \frac{\partial^{|\alpha|}}{\partial x_1^{\alpha_1} \dots \partial x_N^{\alpha_N}},$$

and derivatives are in the sense of distributions. We denote by p_m^* the Sobolev critical exponent: $p_m^* = pN/(N - mp)$ if $N > mp$; $p_m^* = \infty$ if $N \leq mp$.

The reader ought to be careful in reading [18] because Tintarev uses other notation for Sobolev spaces. He denotes the Sobolev spaces defined as above by $\dot{H}^{m,p}(\Omega)$ instead. The assumption $p < N/m$ in considering homogeneous Sobolev spaces is attributed to the fact that otherwise homogeneous Sobolev spaces are no longer spaces of measurable functions, i.e., there are no longer injections from $\dot{W}^{m,p}(\mathbf{R}^N)$ into $L_{\text{loc}}^1(\mathbf{R}^N)$ (see also [19, Remark 2.2]).

We next provide *actions of dislocations* defined on homogeneous Sobolev spaces.

Definition 2.1. Let $q \in \mathbf{R}$. Define a group action on $L_{\text{loc}}^1(\mathbf{R}^N)$ as follows:

$$G[\mathbf{R}^N, \mathbf{Z}; q] := \left\{ g[y, j; q] : L_{\text{loc}}^1(\mathbf{R}^N) \rightarrow L_{\text{loc}}^1(\mathbf{R}^N) \left| \begin{array}{l} g[y, j; q]u(\cdot) := 2^{jN/q}u(2^j(\cdot - y)), \\ u \in L_{\text{loc}}^1(\mathbf{R}^N), y \in \mathbf{R}^N, j \in \mathbf{Z} \end{array} \right. \right\}$$

When $q = p_m^*$, this is a group of bijective isometries on the Sobolev space $\dot{W}^{m,p}(\mathbf{R}^N)$ defined as above:

$$\|g[y, j; p_m^*]u\|_{\dot{W}^{m,p}(\mathbf{R}^N)} = \|u\|_{\dot{W}^{m,p}(\mathbf{R}^N)}, \quad u \in \dot{W}^{m,p}(\mathbf{R}^N), y \in \mathbf{R}^N, j \in \mathbf{Z}.$$

Moreover, inverse mappings are given as follows. For $g[y, j; p_m^*], g[z, k; p_m^*] \in G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$, $u \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($1 \leq p < N/m$),

$$\begin{aligned} g[y, j; p_m^*]^{-1}u(\cdot) &= 2^{-jN/p_m^*}u(2^{-j} \cdot + y) = g[-2^j y, -j; p_m^*]u(\cdot), \\ g[z, k; p_m^*]^{-1}g[y, j; p_m^*]u(\cdot) &= 2^{(j-k)N/p_m^*}u(2^{j-k}(\cdot - 2^k(y - z))) \\ &= g[2^k(y - z), j - k; p_m^*]u(\cdot). \end{aligned}$$

Remark 2.2. One can always replace $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ with $G[\mathbf{R}^N, \mathbf{R}; p_m^*]$ in all of results in this paper.

The following lemma characterizes the operator-weak convergence of dislocations on Sobolev spaces.

Lemma 2.3. *Let $1 < p < N/m$, $m, N \in \mathbf{N}$, and let $(g[y_n, j_n; p_m^*])$ and $(g[z_n, k_n; p_m^*])$ be sequences in $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$. As bounded operators on $\dot{W}^{m,p}(\mathbf{R}^N)$, the operator-weak convergence $g[y_n, j_n; p_m^*] \rightarrow 0$ is equivalent to $|y_n| + |j_n| \rightarrow \infty$, and $g[z_n, k_n; p_m^*]^{-1}g[y_n, j_n; p_m^*] \rightarrow 0$ is equivalent to $|j_n - k_n| + 2^{k_n}|y_n - z_n| \rightarrow \infty$.*

Proof. See [19, Lemmas 3.1 and 5.1]. □

Under these settings, homogeneous Sobolev spaces form dislocation spaces.

Proposition 2.4. *Let $p \in]1, N/m[$ and let $m, N \in \mathbf{N}$ with $m < N$. Then $(\dot{W}^{m,p}(\mathbf{R}^N), G[\mathbf{R}^N, \mathbf{Z}; p_m^*])$ is a dislocation space.*

Proof. We shall show that $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ is a dislocation group. For a sequence $(g[y_n, j_n; p_m^*])$ and $g[y, j; p_m^*]$ in $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$, it is easily checked that $g[y_n, j_n; p_m^*] \rightarrow g[y, j; p_m^*]$ operator-strongly if and only if $|j_n - j| + 2^j |y_n - y| \rightarrow 0$ as $n \rightarrow \infty$, and thus, one sees that $g[y_n, j_n; p_m^*] \rightarrow g[y, j; p_m^*]$ operator-strongly if and only if $g[y_n, j_n; p_m^*]^{-1} \rightarrow g[y, j; p_m^*]^{-1}$ operator-strongly. Let $(g[y_n, j_n; p_m^*])$ be a bounded sequence in $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ such that $g[y_n, j_n; p_m^*] \not\rightarrow 0$, which equivalently means that (y_n) and (j_n) are both bounded due to [Lemma 2.3](#). Then there exist a subsequence, still denoted by n , $y \in \mathbf{R}^N$ and $j \in \mathbf{Z}$ such that $y_n \rightarrow y$, $j_n = j$ ($n \rightarrow \infty$). Hence it follows that $g[y_n, j_n; p_m^*] \rightarrow g[y, j; p_m^*]$ operator-strongly. Finally, we shall show that $(g[y_n, j_n; p_m^*])^* \rightarrow (g[y, j; p_m^*])^*$ operator-strongly (recall (ii) of [Remark 1.2](#)). Indeed, let $u \in \dot{W}^{m,p}(\mathbf{R}^N)$ and let $\phi \in [\dot{W}^{m,p}(\mathbf{R}^N)]^*$. Take functions $\phi_\alpha \in L^{p'}(\mathbf{R}^N)$, $|\alpha| = m$, given by [Lemma C.1](#). It follows from the change of variables and the Hölder inequality that

$$\begin{aligned}
 & \| (g[y_n, j_n; p_m^*])^* \phi - (g[y, j; p_m^*])^* \phi \|_{[\dot{W}^{m,p}(\mathbf{R}^N)]^*} \\
 &= \sup_{\substack{u \in \dot{W}^{m,p}(\mathbf{R}^N) \\ \|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} = 1}} | \langle (g[y_n, j_n; p_m^*])^* \phi - (g[y, j; p_m^*])^* \phi, u \rangle | \\
 &= \sup_{\substack{u \in \dot{W}^{m,p}(\mathbf{R}^N) \\ \|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} = 1}} | \langle \phi, g[y_n, j_n; p_m^*] u - g[y, j; p_m^*] u \rangle | \\
 &= \sup_{\substack{u \in \dot{W}^{m,p}(\mathbf{R}^N) \\ \|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} = 1}} \left| \sum_{|\alpha|=m} \int_{\mathbf{R}^N} \phi_\alpha \partial^\alpha (g[y_n, j_n; p_m^*] u - g[y, j; p_m^*] u) \, dx \right| \\
 &= \sup_{\substack{u \in \dot{W}^{m,p}(\mathbf{R}^N) \\ \|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} = 1}} \left| \sum_{|\alpha|=m} \int_{\mathbf{R}^N} (2^{-j_n \frac{N}{p'}} \phi_\alpha (2^{-j_n} x + y_n) - 2^{-j \frac{N}{p'}} \phi_\alpha (2^{-j} x + y)) \partial^\alpha u \, dx \right| \\
 &= \sup_{\substack{u \in \dot{W}^{m,p}(\mathbf{R}^N) \\ \|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} = 1}} \left| \sum_{|\alpha|=m} \int_{\mathbf{R}^N} (g[y_n, j_n; p']^{-1} \phi_\alpha - g[y, j; p']^{-1} \phi_\alpha) \partial^\alpha u \, dx \right| \\
 &\leq \sup_{\substack{u \in \dot{W}^{m,p}(\mathbf{R}^N) \\ \|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} = 1}} \left(\sum_{|\alpha|=m} \|g[y_n, j_n; p']^{-1} \phi_\alpha - g[y, j; p']^{-1} \phi_\alpha\|_{L^{p'}(\mathbf{R}^N)} \right) \|u\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \\
 &= \sum_{|\alpha|=m} \|g[y_n, j_n; p']^{-1} \phi_\alpha - g[y, j; p']^{-1} \phi_\alpha\|_{L^{p'}(\mathbf{R}^N)}.
 \end{aligned}$$

The last term is convergent to zero as $n \rightarrow \infty$ since $y_n \rightarrow y$, $j_n \rightarrow j$ implies that $\|g[y_n, j_n; p']^{-1} u - g[y, j; p']^{-1} u\|_{L^{p'}(\mathbf{R}^N)} \rightarrow 0$ for all $u \in L^{p'}(\mathbf{R}^N)$. Hence one gets $(g[y_n, j_n; p_m^*])^* \phi - (g[y, j; p_m^*])^* \phi \rightarrow 0$ strongly in $[\dot{W}^{m,p}(\mathbf{R}^N)]^*$, whence follows $(g[y_n, j_n; p_m^*])^* \rightarrow (g[y, j; p_m^*])^*$ operator-strongly. This completes the proof. \square

Notice that the homogeneous Sobolev space $\dot{W}^{m,p}(\mathbf{R}^N)$ with the above dislocation group $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ forms a dislocation space irrelevant to the equivalent Sobolev norms. We finally exhibit an example of G -completely continuous embedding of the homogeneous Sobolev space

into the critical Lebesgue space. The definition of homogeneous Besov spaces will be described in [Appendix C](#) below for the sake of the reader's convenience.

Lemma 2.5. *Let $(\dot{W}^{m,p}(\mathbf{R}^N), G[\mathbf{R}^N, \mathbf{Z}; p_m^*])$ be as in [Proposition 2.4](#). Then the continuous embedding $\dot{W}^{m,p}(\mathbf{R}^N) \hookrightarrow L^{p_m^*}(\mathbf{R}^N)$ is $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ -completely continuous. Indeed, it holds that*

$$\|u\|_{L^{p_m^*}(\mathbf{R}^N)} \leq C \|u\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^{p/p_m^*} \|u\|_{\dot{B}_{\infty,\infty}^{-N/p_m^*}(\mathbf{R}^N)}^{1-p/p_m^*}$$

for all $u \in \dot{W}^{m,p}(\mathbf{R}^N)$, and that for a sequence (u_n) ,

$$\|u_n\|_{\dot{B}_{\infty,\infty}^{-N/p_m^*}(\mathbf{R}^N)} \rightarrow 0$$

whenever $u_n \rightarrow 0$ $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ -weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$.

Proof. See [\[18, Theorems 1.1.9, and 3.2.1\]](#). □

This lemma is generalized as follows:

Lemma 2.6. *Let $(\dot{W}^{m,p}(\mathbf{R}^N), G[\mathbf{R}^N, \mathbf{Z}; p_m^*])$ be as in [Proposition 2.4](#), and let $k \in \mathbf{Z}_{\geq 0}$ with $k < m$ (yielding $p_{m-k}^* < p_m^*$). Then the continuous embedding $\dot{W}^{m,p}(\mathbf{R}^N) \hookrightarrow \dot{W}^{k,p_{m-k}^*}(\mathbf{R}^N)$ is $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ -completely continuous. Moreover, it holds that*

$$\|u\|_{\dot{W}^{k,p_{m-k}^*}(\mathbf{R}^N)} \leq C \|u\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^{p/p_{m-k}^*} \|u\|_{\dot{B}_{\infty,\infty}^{-N/p_{m-k}^*}(\mathbf{R}^N)}^{1-p/p_{m-k}^*}$$

for all $u \in \dot{W}^{m,p}(\mathbf{R}^N)$, and that for a sequence (u_n) ,

$$\|u_n\|_{\dot{B}_{\infty,\infty}^{-N/p_{m-k}^*}(\mathbf{R}^N)} \rightarrow 0$$

whenever $u_n \rightarrow 0$ $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ -weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$.

Proof. See [\[18, Corollary 3.2.2\]](#). □

2.2. Main theorem

Now we move on to a fundamental theorem of profile decomposition in the *homogeneous* Sobolev space $\dot{W}^{m,p}(\mathbf{R}^N)$. Namely, every bounded sequence in $\dot{W}^{m,p}(\mathbf{R}^N)$ has a fine profile decomposition in the sense that the residual term is vanishing in $\dot{W}^{k,p_{m-k}^*}(\mathbf{R}^N)$.

Theorem 2.7. *Let $m, N \in \mathbf{N}$, let $1 < p < N/m$ and let (u_n) be a bounded sequence in $\dot{W}^{m,p}(\mathbf{R}^N)$. Then there exist $\Lambda \in \mathbf{N} \cup \{0, +\infty\}$, a subsequence $(N(n)) \preceq (n)$, $w^l \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$), $(y_{l,N(n)}, j_{l,N(n)}) \in \mathbf{R}^N \times \mathbf{Z}$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$), and residual terms $r_{N(n)}^L \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq L}$) with the relation of a double-suffix profile decomposition*

$$u_{N(n)} = \sum_{l=0}^L 2^{j_{l,N(n)} N/p_m^*} w^l (2^{j_{l,N(n)}} (\cdot - y_{l,N(n)})) + r_{N(n)}^L, \quad L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}, \quad n \in \mathbf{Z}_{\geq L},$$

such that the following hold:

- (i) $y_{0,N(n)} = 0$, $j_{0,N(n)} = 0$ ($n \geq 0$), $w^l \neq 0$ ($1 \leq l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$).

(ii) For each $l \neq k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, it holds that

$$(2.1) \quad |j_{l,N(n)} - j_{k,N(n)}| + 2^{j_{k,N(n)}} |y_{l,N(n)} - y_{k,N(n)}| \rightarrow \infty$$

as $n \rightarrow \infty$. Furthermore, one can assume without loss of generality that, for $l \neq k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, one and only one of the following three cases occurs:

- (a) $j_{l,N(n)} - j_{k,N(n)} \rightarrow +\infty$ ($n \rightarrow \infty$);
- (b) $j_{l,N(n)} - j_{k,N(n)} \rightarrow -\infty$ ($n \rightarrow \infty$);
- (c) $(j_{l,N(n)} - j_{k,N(n)})$ is convergent.

One can also assume without loss of generality that, for $l \neq k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, one and only one of the following two cases occurs:

- (d) $2^{j_{k,N(n)}} |y_{l,N(n)} - y_{k,N(n)}| \rightarrow \infty$ ($n \rightarrow \infty$);
- (e) $(2^{j_{k,N(n)}} (y_{l,N(n)} - y_{k,N(n)}))$ is convergent.

But the cases (c) and (e) do not occur simultaneously due to (2.1).

- (iii) $2^{-j_{l,N(n)}N/p_m^*} u_{N(n)} (2^{-j_{l,N(n)}} \cdot + y_{l,N(n)}) \rightarrow w^l$ as $n \rightarrow \infty$ weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$ and a.e. on \mathbf{R}^N ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$).
- (iv) For $k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$,

$$2^{-j_{k,N(n)}N/p_m^*} r_{N(n)}^L (2^{-j_{k,N(n)}} \cdot + y_{k,N(n)}) \rightarrow \begin{cases} 0 & \text{if } k = 0, \dots, L, \\ w^k & \text{if } k \geq L+1, \end{cases}$$

weakly in $\dot{W}^{1,p}(\mathbf{R}^N)$ and a.e. on \mathbf{R}^N as $n \rightarrow \infty$.

Moreover, the following energy relations hold:

$$(2.2) \quad \begin{aligned} & \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p \\ & \geq \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p + \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{R \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N \setminus \mathcal{B}_{n,R,L})}^p, \end{aligned}$$

$$(2.3) \quad \lim_{L \rightarrow \Lambda} \sup_{\phi \in U} \overline{\lim}_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^N, j \in \mathbf{Z}} \left| \left\langle \phi, 2^{-jN/p_m^*} r_{N(n)}^L (2^{-j} \cdot + y) \right\rangle \right| = 0,$$

$$(2.4) \quad \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \leq 2 \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)} < +\infty,$$

$$(2.5) \quad \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{L^{p_m^*}(\mathbf{R}^N)} = 0,$$

$$(2.6) \quad \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{k,p_m^*-k}(\mathbf{R}^N)} = 0, \quad k \in \mathbf{Z}_{\geq 0}, k < m,$$

where $U := B_{[\dot{W}^{m,p}(\mathbf{R}^N)]^*}(1)$ and $\mathcal{B}_{n,R,L} := \bigcup_{l=0}^L B(y_{l,N(n)}, 2^{-j_{l,N(n)}} R)$.

Remark 2.8. (i) The meaning of each assertion above is as follows: (2.2) indicates that the sum of (p -powered) Sobolev norms of all profiles is bounded by $\overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}$, which is, so to speak, an *energy estimate* or an *energy decomposition*; (2.3) implies the completion of performing the profile decomposition and called the exactness condition; (2.4) ensures that the residual term does not diverge as the number of subtracted dislocated profiles increases; (2.5) and (2.6) show that lower order derivatives of the residual term are vanishing strongly in appropriate Lebesgue or Sobolev spaces.

- (ii) When one considers profile decomposition in Sobolev spaces $\dot{W}^{m,2}(\mathbf{R}^N)$, which is a Hilbert space, one should employ the profile decomposition theorem in Hilbert spaces described in [13, Theorem 2.9] rather than the above theorem, since it provides a more

precise energy decomposition for the Sobolev norm, i.e., one obtains

$$\overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,2}(\mathbf{R}^N)}^2 = \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{m,2}(\mathbf{R}^N)}^2 + \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{m,2}(\mathbf{R}^N)}^2.$$

- (iii) Regarding the energy estimate (2.2), it is noteworthy that this is a sharper version than other authors' energy decompositions because other types of one do not include the "residual part" as in (2.2).
- (iv) Relations (2.3)–(2.6) are also obtained in the previous researches, e.g., [6, 8, 14, 18, 19].

Once the number of nontrivial profiles turns out to be finite, then the assertions above become simpler.

Corollary 2.9. *Suppose that the same conditions as in Theorem 2.7 are satisfied. In addition, assume that Λ is finite, i.e., the number of nontrivial profiles is finite. Then regarding the final residual term given by*

$$r_{N(n)}^\Lambda = u_{N(n)} - \sum_{l=0}^{\Lambda} 2^{j_{l,N(n)}N/p_m^*} w^l(2^{j_{l,N(n)}}(\cdot - y_{l,N(n)})), \quad n \geq \Lambda,$$

the relations (2.3)–(2.5) turn to

$$(2.7) \quad \begin{aligned} & \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p \\ & \geq \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p + \overline{\lim}_{R \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^\Lambda\|_{\dot{W}^{m,p}(\mathbf{R}^N \setminus \mathcal{B}_{n,R,\Lambda})}^p \\ & \lim_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^N, j \in \mathbf{Z}} \left| \left\langle \phi, 2^{-jN/p_m^*} r_{N(n)}^\Lambda(2^{-j} \cdot + y) \right\rangle \right| = 0, \quad \phi \in U, \\ & \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^\Lambda\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \leq 2 \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}, \\ & \lim_{n \rightarrow \infty} \|r_{N(n)}^\Lambda\|_{L^{p_m^*}(\mathbf{R}^N)} = 0, \\ & \lim_{n \rightarrow \infty} \|r_{N(n)}^\Lambda\|_{\dot{W}^{k,p_m^*}(\mathbf{R}^N)} = 0, \quad k \in \mathbf{Z}_{\geq 0}, \quad k < m, \end{aligned}$$

where $U := B_{[\dot{W}^{m,p}(\mathbf{R}^N)]^*}(1)$ and $\mathcal{B}_{n,R,\Lambda} := \bigcup_{l=0}^{\Lambda} B(y_{l,N(n)}, 2^{-j_{l,N(n)}}R)$. In particular, the relation (2.7) implies that the final residual term is $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ -weakly convergent to zero in $\dot{W}^{m,p}(\mathbf{R}^N)$.

The profile decomposition remains true in $\dot{W}^{m,p}(\mathbf{R}^N)$ equipped with an equivalent norm that is often used.

Corollary 2.10. *Under the same conditions as in Theorem 2.7, all of assertions in the theorem hold true even when the norm of $\dot{W}^{1,p}(\mathbf{R}^N)$ is replaced by*

$$\|u\|_{\dot{W}^{1,p}(\mathbf{R}^N)} := \left(\int_{\mathbf{R}^N} |\nabla u|^p dx \right)^{1/p}, \quad u \in \dot{W}^{1,p}(\mathbf{R}^N),$$

where $|\nabla u| = \sqrt{|\frac{\partial u}{\partial x_1}|^2 + \cdots + |\frac{\partial u}{\partial x_N}|^2}$.

Proof. The corollary can be proved in the same way as the proof of Theorem 2.7. \square

2.3. Proof of [Theorem 2.7](#)

We shall employ [Theorem A.1](#) to obtain profiles, and it remains to show the estimates for the Sobolev and Lebesgue norms. In what follows, for the sake of simplicity, we often denote $g[y, j; p_m^*], g[y_{l,n}, j_{l,n}; p_m^*] \in G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ by $g, g_{l,n}$, respectively, for short. Throughout this proof, when a sequence (u_n) converges to u weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$ and almost everywhere on \mathbf{R}^N , we denote it by

$$u_n \rightarrow u \quad \text{weakly and a.e.}$$

for short.

Step 1 (Finding profiles). [Theorem A.1](#) leads us to the existence of profile elements $(w^l, y_{l,N(n)}, j_{l,N(n)}, \Lambda) \in \dot{W}^{m,p}(\mathbf{R}^N) \times \mathbf{R}^N \times \mathbf{Z} \times (\mathbf{N} \cup \{0, \infty\})$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$) satisfying all assertions in [Theorem A.1](#). As in the proof of [Theorem A.1](#) in [\[13\]](#), each profile w^l is obtained as the weak limit of $g_{l,i(l,n)}^{-1} u_{i(l,n)}$, where $(i(l,n))$ denotes the l -th subsequence for which the l -th profile and the l -th dislocations are considered in an iterative process. However, with the help of the Sobolev compact embeddings, the weak convergence $u_n \rightarrow u$ in $\dot{W}^{m,p}(\mathbf{R}^N)$ also leads us to the pointwise convergence $u_n \rightarrow u$ a.e. on \mathbf{R}^N up to a subsequence. Moreover, for the L -th subsequence $(i(L,n))$ ($L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$), one has

$$(2.8) \quad |j_{l,i(L,n)} - j_{k,i(L,n)}| + 2^{j_{k,i(L,n)}} |y_{l,i(L,n)} - y_{k,i(L,n)}| \rightarrow \infty \\ (n \rightarrow \infty, 0 \leq l \neq k \leq L).$$

Thus, on a subsequence, still denoted by $i(L,n)$, one can always assume that, for each $0 \leq l \neq k \leq L$, one and only one of the following three conditions occurs:

- (i) $j_{l,i(L,n)} - j_{k,i(L,n)} \rightarrow +\infty$ ($n \rightarrow \infty$);
- (ii) $j_{l,i(L,n)} - j_{k,i(L,n)} \rightarrow -\infty$ ($n \rightarrow \infty$);
- (iii) $(j_{l,i(L,n)} - j_{k,i(L,n)})$ is convergent in \mathbf{Z} as $n \rightarrow \infty$,

and one can also assume that, for $0 \leq l \neq k \leq L$, one and only one of the following two cases occurs:

- (iv) $2^{j_{k,i(L,n)}} |y_{l,i(L,n)} - y_{k,i(L,n)}| \rightarrow \infty$ ($n \rightarrow \infty$);
- (v) $(2^{j_{k,i(L,n)}} (y_{l,i(L,n)} - y_{k,i(L,n)}))$ is convergent in \mathbf{R}^N as $n \rightarrow \infty$.

But (iii) and (v) do not occur simultaneously due to (2.8). Hence by extracting additional subsequences in each step of the proof, one can obtain both the weak convergence and the pointwise convergence for profiles and residual terms, and also get the above trichotomy and dichotomy regarding dislocations.

Therefore, according to [Theorem A.1](#) we have:

$$(2.9) \quad y_{0,N(n)} = 0, \quad j_{l,N(n)} = 0 \quad (n \geq 0), \\ w^l \neq 0 \quad (1 \leq l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}), \\ |j_{l,N(n)} - j_{k,N(n)}| + 2^{j_{k,N(n)}} |y_{l,N(n)} - y_{k,N(n)}| \rightarrow \infty$$

$$(2.10) \quad (n \rightarrow \infty, k \neq l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}), \\ 2^{-j_{l,N(n)}} u_{N(n)} (2^{-j_{l,N(n)}} \cdot + y_{l,N(n)}) \rightarrow w^l \\ \text{weakly and a.e. } (l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}).$$

Set the residual term by

$$r_{N(n)}^L := u_{N(n)} - \sum_{l=0}^L 2^{j_{l,N(n)}N/p_m^*} w^l (2^{j_{l,N(n)}}(\cdot - y_{l,N(n)})), \quad L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}, \quad n \in \mathbf{Z}_{\geq L}.$$

Then the triangle inequality yields

$$(2.11) \quad \sup_{n \geq 0} \|r_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \leq \sup_{n \geq 0} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)} + \sum_{l=0}^L \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}.$$

Also, [Theorem A.1](#) implies

$$\begin{aligned} 2^{-j_{l,N(n)}N/p_m^*} r_{N(n)}^L (2^{-j_{l,N(n)}} \cdot + y_{l,N(n)}) &\rightarrow 0 \quad \text{weakly and a.e.} \\ &(n \rightarrow \infty, 0 \leq l \leq L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}), \\ 2^{-j_{L,N(n)}N/p_m^*} r_{N(n)}^{L-1} (2^{-j_{L,N(n)}} \cdot + y_{L,N(n)}) &\rightarrow w^L \quad \text{weakly and a.e.} \\ &(n \rightarrow \infty, 1 \leq L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}). \end{aligned}$$

Moreover, one can assume that, for $l \neq k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, one and only one of the following three cases holds:

- (i) $j_{l,N(n)} - j_{k,N(n)} \rightarrow +\infty$ ($n \rightarrow \infty$);
- (ii) $j_{l,N(n)} - j_{k,N(n)} \rightarrow -\infty$ ($n \rightarrow \infty$);
- (iii) $j_{l,N(n)} - j_{k,N(n)}$ is convergent in \mathbf{Z} as $n \rightarrow \infty$,

and one can also assume that, for $l \neq k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, one and only one of the following two cases holds:

- (iv) $2^{j_{k,N(n)}} |y_{l,N(n)} - y_{k,N(n)}| \rightarrow \infty$ ($n \rightarrow \infty$);
- (v) $(2^{j_{k,N(n)}}(y_{l,N(n)} - y_{k,N(n)}))$ is convergent in \mathbf{R}^N as $n \rightarrow \infty$.

But the cases (iii) and (v) do not occur simultaneously due to (2.9). The remaining assertions (2.2)–(2.5) will be proved in the following five lemmas.

Step 2 (Energy decomposition and exactness condition).

Lemma 2.11 (Estimate (2.2) and the exactness condition (2.3)). *It holds that*

$$(2.12) \quad \begin{aligned} &\overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p \\ &\geq \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p + \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{R \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}(\mathbf{R}^N \setminus \mathcal{B}_{n,R,L})}^p, \end{aligned}$$

where $\mathcal{B}_{n,R,L} := \bigcup_{l=0}^L B(y_{l,N(n)}, 2^{-j_{l,N(n)}}R)$. Furthermore, it follows that

$$(2.13) \quad \limsup_{L \rightarrow \Lambda} \overline{\lim}_{\phi \in U} \overline{\lim}_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^N, j \in \mathbf{Z}} \left| \left\langle \phi, 2^{-jN/p_m^*} r_{N(n)}^L (2^{-j} \cdot + y) \right\rangle \right| = 0,$$

where $U := B_{[\dot{W}^{m,p}(\mathbf{R}^N)]^*}(1)$.

Proof. Once one has proved the energy estimate (2.12), then the latter condition (2.13) will be readily obtained due to [Theorem A.1](#). Hence we shall only prove (2.12). Since we have the trichotomy on dislocations $j_{l,N(n)}$, we can define the following disjoint sets: for $0 \leq l \leq L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$,

$$\mathcal{I}_{l,L}^+ := \{l' \in \{0, \dots, L\}; j_{l',N(n)} - j_{l,N(n)} \rightarrow +\infty \text{ as } n \rightarrow \infty\},$$

$$\begin{aligned}\mathcal{J}_{l,L}^- &:= \{l' \in \{0, \dots, L\}; j_{l',N(n)} - j_{l,N(n)} \rightarrow -\infty \text{ as } n \rightarrow \infty\}, \\ \mathcal{J}_{l,L}^0 &:= \{l' \in \{0, \dots, L\}; (j_{l',N(n)} - j_{l,N(n)}) \text{ is convergent}\}, \\ \mathcal{J}_{l,L}^+ &\sqcup \mathcal{J}_{l,L}^- \sqcup \mathcal{J}_{l,L}^0 = \{0, \dots, L\},\end{aligned}$$

where \sqcup denotes a disjoint union of sets. Roughly speaking, $\mathcal{J}_{l,L}^+$ stands for the set of profile numbers up to L whose profiles concentrate faster than the l -th profile w^l ; $\mathcal{J}_{l,L}^-$ for the set of profile numbers up to L whose profiles concentrate slower than the l -th profile w^l ; $\mathcal{J}_{l,L}^0$ for the set of profile numbers up to L whose profiles concentrate at the same speed as the l -th profile w^l .

Let χ be a characteristic function supported on the unit ball $B(0,1)$ and set $\chi_R(\cdot) := \chi(R^{-1}\cdot) = \chi_{B(0,R)}$. Also we define the scaling action $\sigma[y, j]$ for $y \in \mathbf{R}^N$ and $j \in \mathbf{Z}$ by

$$(\sigma[y, j]u)(\cdot) := u(2^j(\cdot - y)), \quad u \in L_{\text{loc}}^1(\mathbf{R}^N).$$

It is easily checked that

$$\begin{aligned}g[y, j; p_m^*]u &= 2^{jN/p_m^*}u(2^j(\cdot - y)) = 2^{jN/p_m^*}\sigma[y, j]u, \\ \partial^\alpha(g[y, j; p_m^*]u) &= 2^{jN/p_m^*}2^{mj}\sigma[y, j](\partial^\alpha u) \\ &= 2^{jN/p}\sigma[y, j](\partial^\alpha u) = g[y, j; p](\partial^\alpha u),\end{aligned}$$

where $\alpha \in (\mathbf{Z}_{\geq 0})^N$ with $|\alpha| = m$. For dislocations $(y_{l,n}, j_{l,n}) \in \mathbf{R}^N \times \mathbf{Z}$, we write

$$\sigma_{l,n}u := \sigma[y_{l,n}, j_{l,n}]u$$

for short. Then the characteristic function supported on $B(y_{l,N(n)}, 2^{-j_{l,N(n)}}R)$ can be written by $\sigma_{l,N(n)}\chi_R$.

Now we provide the following identity: for sufficiently large $n \gg 1$,

$$(2.14) \quad 1 \equiv \sum_{l=0}^L \sigma_{l,N(n)}\chi_R \prod_{l' \in \mathcal{J}_{l,L}^+} (1 - \sigma_{l',N(n)}\chi_R) + \prod_{l=0}^L (1 - \sigma_{l,N(n)}\chi_R).$$

When $\mathcal{J}_{l,L}^+ = \emptyset$, we always assume $\prod_{l' \in \mathcal{J}_{l,L}^+} (1 - \sigma_{l',N(n)}\chi_R) \equiv 1$. The above identity will be shown by induction on L in [Lemma 2.14](#) below. The meaning of the identity reads as follows: the last term $\prod_{l=0}^L (1 - \sigma_{l,N(n)}\chi_R)$ means the characteristic function supported on $\mathbf{R}^N \setminus \mathcal{B}_{n,R,L}$, and $\sigma_{l,N(n)}\chi_R \prod_{l' \in \mathcal{J}_{l,L}^+} (1 - \sigma_{l',N(n)}\chi_R)$ restricts our perspective onto $B(y_{l,N(n)}, 2^{-j_{l,N(n)}}R)$ excluding balls $B(y_{l',N(n)}, 2^{-j_{l',N(n)}}R)$ for $l' \in \mathcal{J}_{l,L}^+$ which stand for the (essential) supports of the profiles concentrating faster than the l -th profile w^l . Roughly speaking, the identity splits \mathbf{R}^N into disjoint supports of profiles w^0, \dots, w^L and the residual term $r_{N(n)}^L$.

With these devices, one sees that for large n and for a fixed multi-index $\alpha \in (\mathbf{Z}_{\geq 0})^N$ with $|\alpha| = m$,

$$(2.15) \quad \begin{aligned}\int_{\mathbf{R}^N} |\partial^\alpha u_{N(n)}|^p dx &= \sum_{l=0}^L \int_{\mathbf{R}^N} |\partial^\alpha u_{N(n)}|^p \sigma_{l,N(n)}\chi_R \prod_{l' \in \mathcal{J}_{l,L}^+} (1 - \sigma_{l',N(n)}\chi_R) dx \\ &\quad + \int_{\mathbf{R}^N} |\partial^\alpha u_{N(n)}|^p \prod_{l=0}^L (1 - \sigma_{l,N(n)}\chi_R) dx\end{aligned}$$

$$=: \sum_{l=0}^L J_1^l + J_2.$$

Due to the convexity of $|\cdot|^p$, one sees that

$$\begin{aligned}
(2.16) \quad J_2 &\geq \int_{\mathbf{R}^N} \left[|\partial^\alpha r_{N(n)}^L|^p + p |\partial^\alpha r_{N(n)}^L|^{p-2} \partial^\alpha r_{N(n)}^L \partial^\alpha \left(\sum_{l=0}^L g_{l,N(n)} w^l \right) \right] \\
&\quad \times \prod_{l=0}^L (1 - \sigma_{l,N(n)} \chi_R) \, dx \\
&= \int_{\mathbf{R}^N} |\partial^\alpha r_{N(n)}^L|^p \prod_{l=0}^L (1 - \sigma_{l,N(n)} \chi_R) \, dx \\
&\quad + p \sum_{l=0}^L \int_{\mathbf{R}^N} |\partial^\alpha r_{N(n)}^L|^{p-2} \partial^\alpha r_{N(n)}^L \partial^\alpha (g_{l,N(n)} w^l) \\
&\quad \quad \quad \times \prod_{l=0}^L (1 - \sigma_{l,N(n)} \chi_R) \, dx \\
&=: J_3 + p \sum_{l=0}^L J_4^l.
\end{aligned}$$

As is observed before, one has

$$\prod_{l=0}^L (1 - \sigma_{l,N(n)} \chi_R) = \chi_{\mathbf{R}^N \setminus [\cup_{l=0}^L B(y_{l,N(n)}, 2^{-j_{l,N(n)}} R)]} = \chi_{\mathbf{R}^N \setminus \mathcal{B}_{n,R,L}},$$

and hence, we find that

$$(2.17) \quad J_3 = \int_{\mathbf{R}^N \setminus \mathcal{B}_{n,R,L}} |\partial^\alpha r_{N(n)}^L|^p \, dx.$$

From the Hölder inequality, the change of variables and (2.11), one also gets

$$\begin{aligned}
(2.18) \quad |J_4^l| &\leq \|\partial^\alpha r_{N(n)}^L\|_{L^p(\mathbf{R}^N)}^{p-1} \times \left(\int_{\mathbf{R}^N} |\partial^\alpha (g_{l,N(n)} w^l)|^p \prod_{l=0}^L (1 - \sigma_{l,N(n)} \chi_R) \, dx \right)^{1/p} \\
&\leq C_L \left(\int_{\mathbf{R}^N} |\partial^\alpha w^l|^p (1 - \chi_{B(0,R)}) \, dx \right)^{1/p} \\
&= C_L \left(\int_{\mathbf{R}^N \setminus B(0,R)} |\partial^\alpha w^l|^p \, dx \right)^{1/p} \rightarrow 0, \quad \text{as } R \rightarrow \infty.
\end{aligned}$$

From (2.16)–(2.18), one obtains

$$(2.19) \quad J_2 \geq \int_{\mathbf{R}^N \setminus \mathcal{B}_{n,R,L}} |\partial^\alpha r_{N(n)}^L|^p \, dx - C_L \sum_{l=0}^L \left(\int_{\mathbf{R}^N \setminus B(0,R)} |\partial^\alpha w^l|^p \, dx \right)^{1/p}.$$

Now we go back to the first term in the last line of (2.15). By changing variables, we get

$$(2.20) \quad J_1^l = \int_{\mathbf{R}^N} |\partial^\alpha u_{N(n)}|^p \sigma_{l,N(n)} \chi_R \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - \sigma_{\nu,N(n)} \chi_R) \, dx$$

$$= \int_{\mathbf{R}^N} |\partial^\alpha ([g_{l,N(n)}]^{-1} u_{N(n)})|^p \chi_R \times \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) dx,$$

where $[\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R$ denotes the characteristic function supported on the ball $B(2^{j_{l,N(n)}}(y_{\nu,N(n)} - y_{l,N(n)}), 2^{-j_{l,N(n)}} R)$. Due to the convexity again, one has

$$(2.21) \quad \begin{aligned} & \int_{\mathbf{R}^N} |\partial^\alpha ([g_{l,N(n)}]^{-1} u_{N(n)})|^p \chi_R \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) dx \\ & \geq \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) dx \\ & \quad + p \int_{\mathbf{R}^N} |\partial^\alpha w^l|^{p-2} \partial^\alpha w^l (\partial^\alpha ([g_{l,N(n)}]^{-1} u_{N(n)}) - \partial^\alpha w^l) \\ & \quad \quad \quad \times \chi_R \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) dx. \end{aligned}$$

Moreover, one can show that

$$(2.22) \quad \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) \rightarrow 1 \text{ in } L^q(\mathbf{R}^N) \text{ (} q \in [1, +\infty[),$$

$$(2.23) \quad \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) dx \rightarrow \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R dx,$$

$$(2.24) \quad \begin{aligned} & \int_{\mathbf{R}^N} |\partial^\alpha w^l|^{p-2} \partial^\alpha w^l (\partial^\alpha ([g_{l,N(n)}]^{-1} u_{N(n)}) - \partial^\alpha w^l) \\ & \quad \quad \quad \times \chi_R \prod_{\nu \in \mathcal{J}_{l,L}^+} (1 - [\sigma_{l,N(n)}]^{-1} \sigma_{\nu,N(n)} \chi_R) dx \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$ (these facts will be proved in [Lemma 2.15](#) below). Hence from (2.20)–(2.24), one gets

$$(2.25) \quad J_1^l \geq \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R dx + o(1)$$

as $n \rightarrow \infty$. Combining (2.15), (2.19) and (2.25), one obtains that

$$(2.26) \quad \begin{aligned} \int_{\mathbf{R}^N} |\partial^\alpha u_{N(n)}|^p dx & \geq \sum_{l=0}^L \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R dx + \int_{\mathbf{R}^N \setminus \mathcal{B}_{n,R,L}} |\partial^\alpha r_{N(n)}^L|^p dx \\ & \quad - C_L \sum_{k=0}^L \int_{\mathbf{R}^N \setminus B(0,R)} |\partial^\alpha w^k|^p dx + o(1) \end{aligned}$$

as $n \rightarrow \infty$. Summing up (2.26) over all multi-indices α with $|\alpha| = m$ and passing to the limits as $n \rightarrow \infty$, $R \rightarrow \infty$ and then $L \rightarrow \Lambda$, we see that

$$\overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p \geq \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p + \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{R \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N \setminus \mathcal{B}_{n,R,L})}^p.$$

Hence it remains to prove (2.22)–(2.24). □

Lemma 2.12 (Boundedness of the residual term (2.4)). *It holds that*

$$\overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \leq 2 \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)} < +\infty.$$

Proof. Set $S_{N(n)}^L := \sum_{l=0}^L g_{l,N(n)} w^l$. We shall employ the following elementary inequality for the Euclidean norm: for all $\alpha_l \in \mathbf{R}^d$ ($l = 1, \dots, L$, $d, L \in \mathbf{N}$),

$$\left| \left| \sum_{l=1}^L \alpha_l \right|^p - \sum_{l=1}^L |\alpha_l|^p \right| \leq C_L \sum_{l \neq m} |\alpha_l| |\alpha_m|^{p-1}$$

for some constant $C_L > 0$. This inequality will be shown similarly to Lemma 3.4 below. From this, we see that, for any multi-index α with $|\alpha| = m$,

$$\begin{aligned} & \left| \int_{\mathbf{R}^N} |\partial^\alpha S_{N(n)}^L|^p dx - \sum_{l=0}^L \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p dx \right| \\ & \leq \int_{\mathbf{R}^N} \left| \left| \sum_{l=0}^L \partial^\alpha g_{l,N(n)} w^l \right|^p - \sum_{l=0}^L |\partial^\alpha g_{l,N(n)} w^l|^p \right| dx \\ & \leq C_L \sum_{l \neq l'} \int_{\mathbf{R}^N} |\partial^\alpha g_{l,N(n)} w^l| |\partial^\alpha g_{l',N(n)} w^{l'}|^{p-1} dx. \end{aligned}$$

The mutual orthogonality condition (2.9) implies that for any $0 \leq l \neq l' \leq L$,

$$(2.27) \quad \int_{\mathbf{R}^N} |\partial^\alpha g_{l,N(n)} w^l| |\partial^\alpha g_{l',N(n)} w^{l'}|^{p-1} dx \rightarrow 0$$

as $n \rightarrow \infty$. Hence we get

$$\lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} |\partial^\alpha S_{N(n)}^L|^p dx = \sum_{l=0}^L \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p dx.$$

Adding up the above over all multi-indices α with $|\alpha| = m$, we obtain

$$(2.28) \quad \|S_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p = \sum_{l=0}^L \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p + o(1) \quad (n \rightarrow \infty),$$

and from (2.12) and (2.28), one sees that

$$(2.29) \quad \begin{aligned} \lim_{L \rightarrow \Lambda} \lim_{n \rightarrow \infty} \|S_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p &= \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p \\ &\leq \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p. \end{aligned}$$

So we observe that

$$\begin{aligned} & \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \\ & \leq \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)} + \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|S_{N(n)}^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \\ & \leq 2 \overline{\lim}_{n \rightarrow \infty} \|u_{N(n)}\|_{\dot{W}^{m,p}(\mathbf{R}^N)}, \end{aligned}$$

which implies the boundedness of the double sequence $(r_{N(n)}^L)$ in $\dot{W}^{m,p}(\mathbf{R}^N)$. \square

Step 3 (Vanishing of the residual term).

Lemma 2.13 (Vanishing of the residual term (2.5)). *It holds that*

$$\begin{aligned} \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{L^{p_m}(\mathbf{R}^N)} &= 0, \\ \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_{N(n)}^L\|_{\dot{W}^{k,p_{m-k}}(\mathbf{R}^N)} &= 0, \quad k \in \mathbf{Z}_{\geq 0}, k < m. \end{aligned}$$

Proof. This lemma is readily checked by Theorem A.3 and Lemmas 2.5 and 2.6 together with (2.13). \square

The following two lemmas will provide supplementary calculations for proofs of the above lemmas. For the purpose of the proof of (2.14), we generalize the identity as follows:

Lemma 2.14 (Generalization of (2.14)). *For any $L \in \mathbf{N}$ with $L \leq \Lambda + 1$ and for all $l_\nu \in \mathbf{Z}_{\geq 0}^{\leq \Lambda+1}$ ($1 \leq \nu \leq L$) with $l_\nu \neq l_{\nu'}$ ($\nu \neq \nu'$), it holds that*

$$1 \equiv \sum_{\nu=1}^L \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) + \prod_{\nu=1}^L (1 - \sigma_{l_\nu, N(n)} \chi_R)$$

for sufficiently large $n \in \mathbf{Z}_{\geq 0}$, where $\sigma_{l_\nu, N(n)} \chi_R$ denotes the characteristic function supported on the ball $B(y_{l_\nu, N(n)}, 2^{-j_{l_\nu, N(n)}} R)$, and

$$J_{l_\nu, l_L}^+ := \{l_\mu \in \mathbf{Z}_{\geq 0}; 1 \leq \nu \leq L, j_{l_\mu, N(n)} - j_{l_\nu, N(n)} \rightarrow +\infty (n \rightarrow \infty)\}.$$

Proof. We prove the identity by induction on L .

(I) Base step: $L = 1$. In this case, the identity is trivial.

(II) Inductive step. Let $L \in \mathbf{N}$ satisfy $L + 1 \leq \Lambda + 1$ and $L + 1 < +\infty$. Suppose that the identity holds true for all positive integers up to $L \in \mathbf{N}$. Let $l_\nu \in \mathbf{Z}_{\geq 0}^{\leq \Lambda+1}$ ($1 \leq \nu \leq L + 1$) with $l_\nu \neq l_{\nu'}$ ($\nu \neq \nu'$). We use the following partition:

$$\begin{aligned} \{l_1, \dots, l_L\} &= A_1 \sqcup A_2 \sqcup A_3, \\ A_1 &= \{l_\nu \in \mathbf{Z}_{\geq 0}; 1 \leq \nu \leq L, j_{l_\nu, N(n)} - j_{l_{L+1}, N(n)} \rightarrow +\infty (n \rightarrow \infty)\} \\ &= J_{l_{L+1}, l_{L+1}}^+, \\ A_2 &= \{l_\nu \in \mathbf{Z}_{\geq 0}; 1 \leq \nu \leq L, j_{l_\nu, N(n)} - j_{l_{L+1}, N(n)} \rightarrow -\infty (n \rightarrow \infty)\}, \\ A_3 &= \{l_\nu \in \mathbf{Z}_{\geq 0}; 1 \leq \nu \leq L, j_{l_\nu, N(n)} - j_{l_{L+1}, N(n)} \text{ is convergent}\}. \end{aligned}$$

Then we see that

$$\begin{aligned} J_{l_\nu, l_{L+1}}^+ &= J_{l_\nu, l_L}^+ && \text{if } l_\nu \in A_1, \\ J_{l_\nu, l_{L+1}}^+ &= J_{l_\nu, l_L}^+ \sqcup \{l_{L+1}\} && \text{if } l_\nu \in A_2, \\ J_{l_\nu, l_{L+1}}^+ &= J_{l_\nu, l_L}^+ && \text{if } l_\nu \in A_3. \end{aligned}$$

It follows that

$$\begin{aligned} &\sum_{\nu=1}^{L+1} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_{L+1}}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\ &= \sum_{\nu=1}^L \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_{L+1}}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \end{aligned}$$

$$\begin{aligned}
& + \sigma_{l_{L+1}, N(n)} \chi_R \prod_{l_\mu \in J_{l_{L+1}, l_{L+1}}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
= & \sum_{l_\nu \in A_1} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& + \sum_{l_\nu \in A_2} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) (1 - \sigma_{l_{L+1}, N(n)} \chi_R) \\
& + \sum_{l_\nu \in A_3} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& + \sigma_{l_{L+1}, N(n)} \chi_R \prod_{l_\mu \in A_1} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
= & \sum_{\nu=1}^L \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& - \sigma_{l_{L+1}, N(n)} \chi_R \sum_{l_\nu \in A_2} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& + \sigma_{l_{L+1}, N(n)} \chi_R \prod_{l_\mu \in A_1} (1 - \sigma_{l_\mu, N(n)} \chi_R).
\end{aligned}$$

By the induction hypothesis, the last line turns to:

$$\begin{aligned}
& = \sum_{\nu=1}^L \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& - \sigma_{l_{L+1}, N(n)} \chi_R \sum_{l_\nu \in A_2} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& + \sigma_{l_{L+1}, N(n)} \chi_R \left(1 - \sum_{l_\nu \in A_1} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+ \cap A_1} (1 - \sigma_{l_\mu, N(n)} \chi_R) \right).
\end{aligned}$$

It is readily seen that $J_{l_\nu, l_L}^+ \cap A_1 = J_{l_\nu, l_L}^+$ when $l_\nu \in A_1$, and hence, the last line turns to:

$$\begin{aligned}
(2.30) \quad & = \sum_{\nu=1}^L \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \\
& + \sigma_{l_{L+1}, N(n)} \chi_R \left(1 - \sum_{l_\nu \in A_1 \sqcup A_2 \sqcup A_3} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R) \right) \\
& + \sigma_{l_{L+1}, N(n)} \chi_R \sum_{l_\nu \in A_3} \sigma_{l_\nu, N(n)} \chi_R \prod_{l_\mu \in J_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu, N(n)} \chi_R).
\end{aligned}$$

Whenever $l_\nu \in A_3$, the mutual orthogonality condition (2.9) implies that the supports of $\sigma_{l_{L+1},N(n)}\chi_R$ and $\sigma_{l_\nu,N(n)}\chi_R$ are mutually disjoint for sufficiently large n , and thus

$$\sigma_{l_{L+1},N(n)}\chi_R \sum_{l_\nu \in A_3} \sigma_{l_\nu,N(n)}\chi_R \prod_{l_\mu \in \mathcal{J}_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu,N(n)}\chi_R) = 0$$

for sufficiently large n .

Therefore, from the above and the induction hypothesis again, (2.30) leads to:

$$\begin{aligned} &= \sum_{\nu=1}^L \sigma_{l_\nu,N(n)}\chi_R \prod_{l_\mu \in \mathcal{J}_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu,N(n)}\chi_R) \\ &\quad + \sigma_{l_{L+1},N(n)}\chi_R \left(1 - \sum_{\nu=1}^L \sigma_{l_\nu,N(n)}\chi_R \prod_{l_\mu \in \mathcal{J}_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu,N(n)}\chi_R) \right) \\ &= (1 - \sigma_{l_{L+1},N(n)}\chi_R) \sum_{\nu=1}^L \sigma_{l_\nu,N(n)}\chi_R \prod_{l_\mu \in \mathcal{J}_{l_\nu, l_L}^+} (1 - \sigma_{l_\mu,N(n)}\chi_R) \\ &\quad + \sigma_{l_{L+1},N(n)}\chi_R \\ &= (1 - \sigma_{l_{L+1},N(n)}\chi_R) \left(1 - \prod_{\nu=1}^L (1 - \sigma_{l_\nu,N(n)}\chi_R) \right) + \sigma_{l_{L+1},N(n)}\chi_R \\ &= 1 - \prod_{\nu=1}^{L+1} (1 - \sigma_{l_\nu,N(n)}\chi_R) \end{aligned}$$

for sufficiently large n . This completes the proof. \square

Lemma 2.15 (Proofs of (2.22), (2.23) and (2.24)). *Under the same settings as in the proof of Lemma 2.11, it holds that*

$$(2.31) \quad \prod_{l' \in \mathcal{J}_{l, L}^+} (1 - [\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R) \rightarrow 1 \text{ in } L^q(\mathbf{R}^N) \text{ (} q \in [1, +\infty[),$$

$$(2.32) \quad \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R \prod_{l' \in \mathcal{J}_{l, L}^+} (1 - [\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R) dx \rightarrow \int_{\mathbf{R}^N} |\partial^\alpha w^l|^p \chi_R dx,$$

$$(2.33) \quad \int_{\mathbf{R}^N} |\partial^\alpha w^l|^{p-2} \partial^\alpha w^l (\partial^\alpha ([g_{l, N(n)}]^{-1} u_{N(n)}) - \partial^\alpha w^l) \\ \times \chi_R \prod_{l' \in \mathcal{J}_{l, L}^+} (1 - [\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R) dx \rightarrow 0,$$

as $n \rightarrow \infty$.

Proof. When $\mathcal{J}_{l, L}^+ = \emptyset$, we always assume that $\prod_{l' \in \mathcal{J}_{l, L}^+} (1 - [\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R) \equiv 1$, and so the assertions are all trivial. Hence we assume $\mathcal{J}_{l, L}^+ \neq \emptyset$.

We shall prove (2.31). Expanding the products and calculating $1 - \prod_{l' \in \mathcal{J}_{l, L}^+} (1 - [\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R)$, one sees that each term of the summation contains at least one $[\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R$ for some $l' \in \mathcal{J}_{l, L}^+$. Hence the L^q -norm of each term is majorized by the L^q -norm of such $[\sigma_{l, N(n)}]^{-1} \sigma_{l', N(n)} \chi_R$

that is infinitesimal as $n \rightarrow \infty$, thanks to $l' \in \mathcal{J}_{l,L}^+$ and the mutual orthogonality condition. Hence (2.31) follows.

We next show (2.32). By approximation, we may assume that $\partial^\alpha w^l$ is smooth. Then thanks to (2.31), one gets (2.32). Regarding (2.33), we shall approximate $|\partial^\alpha w^l|^{p-2} \partial^\alpha w^l$ by a smooth function. Then (2.33) follows from

$$\partial^\alpha [g_{l,N(n)}]^{-1} u_{N(n)} \rightarrow \partial^\alpha w^l \text{ weakly in } L^p(\mathbf{R}^N)$$

and (2.31). \square

Eventually, the proof of Theorem 2.7 has been complete. \square

2.4. Relations among profile decompositions in inhomogeneous and homogeneous Sobolev spaces

Fix $1 < p < N/m$, and recall that $W^{m,p}(\mathbf{R}^N)$ is continuously embedded into $\dot{W}^{m,p}(\mathbf{R}^N)$ and that $W^{m,p}(\mathbf{R}^N) \cong \dot{W}^{m,p}(\mathbf{R}^N) \cap L^p(\mathbf{R}^N)$. Now consider that

- (u_n) is a bounded sequence in the *inhomogeneous* Sobolev space $W^{m,p}(\mathbf{R}^N)$.

Then there are two types of profile decomposition of (u_n) :

- the profile decomposition in $W^{m,p}(\mathbf{R}^N)$ given by [13, Theorem 3.8];
- the profile decomposition in $\dot{W}^{m,p}(\mathbf{R}^N)$ given by Theorem 2.7.

In this section, we shall discuss the relations between the above two.

Recall that, by Theorem 2.7, there exist:

- a subsequence of (n) , still denoted by n ,
- a number $\Lambda \in \mathbf{N} \cup \{0, +\infty\}$,
- profiles $w^l \in \dot{W}^{m,p}(\mathbf{R}^N)$, $l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$,
- dislocations $(y_{l,n}, j_{l,n}) \in \mathbf{R}^N \times \mathbf{R}$, $l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$,
- residual terms $r_n^L \in \dot{W}^{m,p}(\mathbf{R}^N)$, $L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq L}$,

with the relation of a double-suffix profile decomposition

$$u_n = \sum_{l=0}^L 2^{j_{l,n}N/p_m^*} w^l(2^{j_{l,n}}(\cdot - y_{l,n})) + r_n^L, \quad L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}, \quad n \in \mathbf{Z}_{\geq L},$$

such that all relations in Theorem 2.7 hold true. Furthermore, we may assume that the index set $\mathbf{Z}_{\geq 0}^{<\Lambda+1}$ is decomposed into disjoint sets as

$$\begin{aligned} \mathbf{Z}_{\geq 0}^{<\Lambda+1} &= J_+ \sqcup J_0 \sqcup J_-, \\ J_+ &:= \left\{ l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}; j_{l,n} \rightarrow +\infty \right\}, \\ J_0 &:= \left\{ l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}; j_{l,n} \rightarrow j_{l,\infty} \right\}, \\ J_- &:= \left\{ l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}; j_{l,n} \rightarrow -\infty \right\}. \end{aligned}$$

Under the above situation, we can show the following

Lemma 2.16. (i) $J_- = \emptyset$.

(ii) If $l \in J_0$, then $w^l \in W^{m,p}(\mathbf{R}^N)$ and moreover, without loss of generality, one may assume that $j_{l,\infty} = j_{l,n} = 0$ for all $n \in \mathbf{Z}_{\geq l}$.

Proof. We shall first show the assertion (i). Suppose that $J_- \neq \emptyset$ and let $l \in J_-$. It then follows that for every $e \in C_c^\infty(\mathbf{R}^N)$,

$$\int_{\mathbf{R}^N} 2^{-j_{l,n}N/p_m^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) e \, dx \rightarrow \int_{\mathbf{R}^N} w^l e \, dx \quad (n \rightarrow \infty).$$

On the other hand, one sees that

$$\begin{aligned} \left| \int_{\mathbf{R}^N} 2^{-j_{l,n}N/p_m^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) e \, dx \right| &\leq 2^{-j_{l,n}} \int_{\mathbf{R}^N} \left| 2^{-j_{l,n}N/p} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) e \right| dx \\ &\leq 2^{-j_{l,n}} \|u_n\|_{L^p(\mathbf{R}^N)} \|e\|_{L^{p'}(\mathbf{R}^N)} \\ &= o(1) \end{aligned}$$

as $n \rightarrow \infty$. Thus from the density argument we conclude that $w^l = 0$. But this contradicts the fact that $w^l \neq 0$ for every $l \in \mathbf{Z}_{\geq 0}^{\leq \Lambda+1}$, hence the conclusion.

We then prove (ii). Let $l \in J_0$. Recall that

$$2^{-j_{l,n}N/p_m^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) \rightarrow w^l \text{ weakly in } \dot{W}^{m,p}(\mathbf{R}^N).$$

Since $j_{l,n} \rightarrow j_{l,\infty}$, one sees from changing variables that

$$\begin{aligned} 2^{-(j_{l,n}-j_{l,\infty})N/p_m^*} u_n(2^{-(j_{l,n}-j_{l,\infty})} \cdot + y_{l,n}) &\rightarrow 2^{j_{l,\infty}M/p_m^*} w^l(2^{j_{l,\infty}} \cdot) \\ &\text{weakly in } \dot{W}^{m,p}(\mathbf{R}^N). \end{aligned}$$

Thus with no loss of generality, we assume that $j_{l,\infty} = 0$ for each $l \in J_0$ and $j_{l,n} \rightarrow 0$.

Take $\phi \in [\dot{W}^{m,p}(\mathbf{R}^N)]^*$ arbitrarily, and let $(\phi_\alpha)_\alpha \subset L^p(\mathbf{R}^N)$ be its Riesz representation given by [Lemma C.1](#). It then follows that

$$(2.34) \quad \sum_{|\alpha|=m} \int_{\mathbf{R}^N} \partial_\alpha \left(2^{-j_{l,n}N/p_m^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) \right) \phi_\alpha \, dx \rightarrow \sum_{|\alpha|=m} \int_{\mathbf{R}^N} \partial_\alpha w^l \phi_\alpha \, dx$$

as $n \rightarrow \infty$. On the other hand, it follows that

$$\begin{aligned} (2.35) \quad &\left| \int_{\mathbf{R}^N} \partial_\alpha \left(2^{-j_{l,n}N/p_m^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) \right) \phi_\alpha \, dx - \int_{\mathbf{R}^N} \partial_\alpha u_n(\cdot + y_{l,n}) \phi_\alpha \, dx \right| \\ &\leq \int_{\mathbf{R}^N} |\partial_\alpha u_n(\cdot + y_{l,n})| \left| 2^{j_{l,n}N/p_m^*} \phi_\alpha(2^{j_{l,n}} \cdot) \phi_\alpha - \phi_\alpha \right| dx \\ &\leq \|u_n\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \left\| 2^{j_{l,n}N/p_m^*} \phi_\alpha(2^{j_{l,n}} \cdot) \phi_\alpha - \phi_\alpha \right\|_{L^{p'}(\mathbf{R}^N)} \\ &= o(1) \end{aligned}$$

as $n \rightarrow \infty$.

From (2.34) and (2.35), we infer that

$$\sum_{|\alpha|=m} \int_{\mathbf{R}^N} \partial_\alpha u_n(2^{-j_{l,n}} \cdot + y_{l,n}) \phi_\alpha \, dx \rightarrow \sum_{|\alpha|=m} \int_{\mathbf{R}^N} \partial_\alpha w^l \phi_\alpha \, dx$$

as $n \rightarrow \infty$, which implies that

$$u_n(\cdot + y_{l,n}) \rightarrow w^l \text{ weakly in } \dot{W}^{m,p}(\mathbf{R}^N).$$

Also, from the following lemma, one finds that $w^l \in W^{m,p}(\mathbf{R}^N)$ (not just $\dot{W}^{m,p}(\mathbf{R}^N)$) for all $l \in J_0$, and thus the proof is complete. \square

Lemma 2.17. *Let $u_n \in W^{m,p}(\mathbf{R}^N)$ ($n \in \mathbf{N}$) be such that $u_n \rightarrow u$ weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$ (not $W^{m,p}(\mathbf{R}^N)$) for some $u \in \dot{W}^{m,p}(\mathbf{R}^N)$. If $\sup_{n \in \mathbf{N}} \|u_n\|_{W^{m,p}(\mathbf{R}^N)} < \infty$, then $u \in W^{m,p}(\mathbf{R}^N)$.*

Proof. It suffices to show that $u_n \rightarrow u$ weakly in $L^p(\mathbf{R}^N)$. This follows from the boundedness of (u_n) in $L^p(\mathbf{R}^N)$ and the pointwise convergence $u_n \rightarrow u$ a.e. in \mathbf{R}^N (up to a subsequence) which follows from the local compact embeddings for Sobolev spaces. \square

By the same calculations as in the proof of the energy decomposition for the inhomogeneous Sobolev norm (see also [13]), one can show that

$$\overline{\lim}_{n \rightarrow \infty} \|u_n\|_{W^{m,p}(\mathbf{R}^N)}^p \geq \sum_{l \in J_0} \|w^l\|_{W^{m,p}(\mathbf{R}^N)}^p.$$

Also, we find that

$$\lim_{L \rightarrow \Lambda} \sup_{\phi \in B_{[W^{m,p}(\mathbf{R}^N)]^*}(1)} \overline{\lim}_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^N} \left| \langle \phi, \rho_n^L(\cdot + y) \rangle_{W^{m,p}(\mathbf{R}^N)} \right| = 0,$$

where $\rho_n^L = u_n - \sum_{l=0}^L w^{i_l}(\cdot - y_{i_l,n})$ and $J_0 = \{0 < i_1 < i_2 < \dots\}$. This can be proved by the same argument as in [13, Lemma 3.12].

Therefore, w^l and $y_{l,n}$ ($l \in J_0$, $n \in \mathbf{N}_{\geq l}$) are profile elements for the profile decomposition of (u_n) in $W^{m,p}(\mathbf{R}^N)$, and assertions as in [13, Theorem 3.8] hold. In short, the profile decompositions in $\dot{W}^{m,p}(\mathbf{R}^N)$ of bounded sequences in $W^{m,p}(\mathbf{R}^N)$ include the profile decompositions in $W^{m,p}(\mathbf{R}^N)$ of (u_n) .

Summing up the above arguments, we obtain the following

Theorem 2.18. *Let (u_n) be a bounded sequence in $W^{m,p}(\mathbf{R}^N)$. Then there exist a subsequence of (n) , still denoted by n , a number $\Lambda \in \mathbf{N} \cup \{0, +\infty\}$, profiles $w^l \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$), dislocations $(y_{l,n}, j_{l,n}) \in \mathbf{R}^N \times \mathbf{R}$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$), and residual terms $r_n^L \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq L}$), with the relation of a double-suffix profile decomposition*

$$u_n = \sum_{l=0}^L 2^{j_{l,n}N/p_m^*} w^l \left(2^{k_{l,n}}(\cdot - y_{l,n}) \right), \quad L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}, \quad n \in \mathbf{Z}_{\geq L},$$

such that assertions as in [Theorem 2.7](#) hold true. Moreover, set

$$\begin{aligned} J_+ &:= \{l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}; j_{l,n} \rightarrow +\infty\}, \\ J_0 &:= \{l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}; j_{l,n} \rightarrow j_{l,\infty}\}, \\ J_- &:= \{l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}; j_{l,n} \rightarrow -\infty\}, \\ \mathbf{Z}_{\geq 0}^{<\Lambda+1} &= J_+ \sqcup J_0 \sqcup J_-. \end{aligned}$$

Then,

$$\begin{aligned} J_- &= \emptyset, \\ j_{l,n} &= j_{l,\infty} = 0 \quad (l \in J_0), \\ w^l &\in W^{m,p}(\mathbf{R}^N) \quad (l \in J_0), \\ u_n(\cdot + y_{l,n}) &\rightarrow w^l \quad \text{weakly in } W^{m,p}(\mathbf{R}^N) \quad (l \in J_0), \end{aligned}$$

$$\begin{aligned} \overline{\lim}_{n \rightarrow \infty} \|u_n\|_{W^{m,p}(\mathbf{R}^N)}^p &\geq \sum_{l \in J_0} \|w^l\|_{W^{m,p}(\mathbf{R}^N)}^p, \\ \lim_{L \rightarrow \Lambda} \sup_{\phi \in B_{[W^{m,p}(\mathbf{R}^N)]^*}(1)} \overline{\lim}_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^N} \left| \langle \phi, \rho_n^L(\cdot + y) \rangle_{W^{m,p}(\mathbf{R}^N)} \right| &= 0, \\ \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|\rho_n^L\|_{L^q(\mathbf{R}^N)} &= 0, \quad q \in]p, p_m^*[, \end{aligned}$$

where $\rho_n^L := u_n - \sum_{l=0}^L w^{i_l}(\cdot - y_{i_l,n})$ and $J_0 = \{0 < i_1 < i_2 < \dots\}$.

Summary. The above arguments are summarized as follows: For a bounded sequence in $\dot{W}^{m,p}(\mathbf{R}^N)$ has three types of profiles:

- ones which are translated by vectors $y_{l,n} \in \mathbf{R}^N$,
- ones which are concentrating at some points,
- ones which are anti-concentrating, i.e., vanishing locally.

However, the above arguments indicate that if (u_n) is bounded in $L^p(\mathbf{R}^N)$, then the anti-concentrating profiles do not exist because the L^p -norm of those profiles is increasing to infinity, contradicting the L^p -boundedness. Conversely, the concentrating profiles exist because the L^p -norm of those profiles is converging to zero.

3. Decomposition of integral functionals in critical cases

In this section, we always assume $1 < p < N/m$ for $m, N \in \mathbf{N}$.

As is intended in [13], we shall investigate the results of decomposition of integral functionals of critical order (in other words, the iterated Brezis-Lieb lemma). The author pointed out in [13] that the results of decomposition of integral functionals will be obtained in Lebesgue or Sobolev spaces into which the dislocation Sobolev spaces considered are embedded G -completely continuously. Hence, along with Lemmas 2.5 and 2.6, we shall approach the results of decomposition of integral functionals in lower order homogeneous Sobolev spaces with critical exponents. This result has been obtained by some precursors on profile decomposition, e.g., [18, 19].

3.1. Brief Summary

Firstly, we shall briefly review our results, choosing typical and simple examples in order to describe the essence of the results. Assume that (u_n) is a bounded sequence in $\dot{W}^{m,p}(\mathbf{R}^N)$ and take profile elements $(w^l, y_{l,n}, j_{l,n}\Lambda)$ (on a subsequence still denoted by n) given by Theorem 2.7. According to Lemmas 2.5 and 2.6 and the exactness condition (2.3), the residual term r_n^L and its derivatives are vanishing (as $n \rightarrow \infty$ and then $L \rightarrow \Lambda$) in suitable $L^{p_m^*}(\mathbf{R}^N)$ or $\dot{W}^{k,p_{m-k}^*}(\mathbf{R}^N)$.

Typically, our main results read:

$$(3.1) \quad \lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} |u_n|^{p_m^*} dx = \sum_{l=0}^{\Lambda} \int_{\mathbf{R}^N} |w^l|^{p_m^*} dx,$$

$$(3.2) \quad \lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} |\partial^\alpha u_n|^{p_{m-|\alpha|}^*} dx = \sum_{l=0}^{\Lambda} \int_{\mathbf{R}^N} |\partial^\alpha w^l|^{p_{m-|\alpha|}^*} dx, \quad |\alpha| < m,$$

$$(3.3) \quad \lim_{n \rightarrow \infty} \|u_n\|_{W^{k, p_m^*}(\mathbf{R}^N)}^{p_m^* - k} = \sum_{l=0}^{\Lambda} \|w^l\|_{W^{k, p_m^*}(\mathbf{R}^N)}^{p_m^* - k}, \quad 0 \leq k < m.$$

These formulas are shown as follows. As for Euclidean norms, one can show:

$$(3.4) \quad \left| \left| \sum_{l=1}^L s_l \right|^q - \sum_{l=1}^L |s_l|^q \right| \leq C_L \sum_{l \neq k} |s_l| |s_k|^{q-1}, \quad s_l \in \mathbf{R},$$

$$(3.5) \quad ||a + b|^q - |a|^q| \leq \varepsilon |a|^q + C_\varepsilon |b|^q, \quad \varepsilon > 0, a, b \in \mathbf{R}.$$

Combining the above, we see:

$$\begin{aligned} & \left| \int_{\mathbf{R}^N} |u_n|^{p_m^*} dx - \sum_{l=0}^L \int_{\mathbf{R}^N} |w^l|^{p_m^*} dx \right| \\ &= \left| \int_{\mathbf{R}^N} |u_n|^{p_m^*} dx - \sum_{l=0}^L \int_{\mathbf{R}^N} |w^l(\cdot - y_{l,n})|^{p_m^*} dx \right| \\ &\leq \varepsilon \int_{\mathbf{R}^N} \left| \sum_{l=0}^L w^l(\cdot - y_{l,n}) \right|^{p_m^*} dx + C_\varepsilon \int_{\mathbf{R}^N} |r_n^L|^{p_m^*} dx \\ &\quad + C_L \sum_{l \neq k} \int_{\mathbf{R}^N} |w^l(\cdot - y_{l,n})| |w^k(\cdot - y_{k,n})|^{p_m^* - 1} dx. \end{aligned}$$

Since (u_n) and (r_n^L) are bounded in $\dot{W}^{m,p}(\mathbf{R}^N)$, $\int_{\mathbf{R}^N} |\sum_{l=0}^L w^l(\cdot - y_{l,n})|^{p_m^*} dx$ is bounded. By the mutual orthogonality condition, $\int_{\mathbf{R}^N} |w^l(\cdot - y_{l,n})| |w^k(\cdot - y_{k,n})|^{p_m^* - 1} dx$ is converging to zero as $n \rightarrow \infty$. Also, the residual term $\int_{\mathbf{R}^N} |r_n^L|^{p_m^*} dx$ is vanishing as $n \rightarrow \infty$ and then $L \rightarrow \Lambda$. Therefore, passing to the limits as $n \rightarrow \infty$, $L \rightarrow \Lambda$ and then $\varepsilon \rightarrow 0$, one can conclude (3.1). Similarly, one can show (3.2) and (3.3).

In the above observation, the inequality (3.4) is employed together with the mutual orthogonality condition, yielding the degeneracy of cross terms. The inequality (3.5) is used with $b = r_n^L$ which is vanishing in the suitable space. Hence, one can obtain other variants of the above decompositions of integral functionals, by considering the Lebesgue or Sobolev spaces where $\dot{W}^{m,p}(\mathbf{R}^N)$ is embedded $G[\mathbf{R}^N, \mathbf{Z}; p_m^*]$ -completely continuously. Along the above strategy, in what follows, we shall investigate more general results.

3.2. Main theorems

Now we shall discuss the general results of decomposition of integral functionals of $(\int_{\mathbf{R}^N} F(\partial^\alpha u_n) dx)$ with continuous functions F of critical growth, based on the profile decomposition in *homogeneous* Sobolev spaces. We consider two types of continuous functions here: locally Lipschitz continuous functions with homogeneity of critical order; smooth functions with asymptotic growth of critical power near zero and infinity. We refer the reader to [19, Section 5.2.] as a reference of conditions of continuous functions.

In what follows, $\text{Lip}_{\text{loc}}(\mathbf{R})$ denotes the set of locally Lipschitz continuous functions on \mathbf{R} . Let $\alpha \in (\mathbf{Z}_{\geq 0})^N$ be a multi-index such that $|\alpha| < m$. Firstly, consider a function $F_\alpha \in \text{Lip}_{\text{loc}}(\mathbf{R})$ such that

$$(3.6) \quad F_\alpha(2^{jN/p_m^* - |\alpha|} s) = 2^{jN} F_\alpha(s), \quad s \in \mathbf{R}, j \in \mathbf{Z}.$$

From this condition immediately follows

$$|F(s)| \leq C|s|^{p_m^* - |\alpha|}, \quad s \in \mathbf{R},$$

for some constant $C > 0$. For those functions F , an invariance property follows from the change of variables:

$$\int_{\mathbf{R}^N} F\left(\partial^\alpha \left[2^{jN/p_m^*} u(2^j x)\right]\right) dx = \int_{\mathbf{R}^N} F(\partial^\alpha u(x)) dx$$

for all $u \in L^{p_m^*}(\mathbf{R}^N)$ and $j \in \mathbf{Z}$.

Now one form of the results of decomposition of integral functionals in the critical case reads:

Theorem 3.1 (Decomposition of integral functionals 1). *Let (u_n) be a bounded sequence in $\dot{W}^{m,p}(\mathbf{R}^N)$. Assume that on a subsequence, still denoted by n , (u_n) has a profile decomposition with profile elements $(w^l, y_{l,n}, j_{l,n}, \Lambda) \in \dot{W}^{m,p}(\mathbf{R}^N) \times \mathbf{R}^N \times \mathbf{Z} \times (\mathbf{N} \cup \{0, \infty\})$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$) as in [Theorem 2.7](#). Let $\alpha \in (\mathbf{Z}_{\geq 0})^N$ be a multi-index such that $|\alpha| < m$, and let $F_\alpha \in \text{Lip}_{\text{loc}}(\mathbf{R})$ satisfy [\(3.6\)](#). Then the following holds true:*

$$\lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} F_\alpha(\partial^\alpha u_n(x)) dx = \sum_{l=0}^{\Lambda} \int_{\mathbf{R}^N} F_\alpha(\partial^\alpha w^l(x)) dx.$$

We secondly consider a continuous function F_α as follows. Let $f_\alpha \in C(\mathbf{R})$ satisfy

$$(3.7) \quad |f_\alpha(s)| \leq C|s|^{(p_m^* - |\alpha|)^{-1}}, \quad s \in \mathbf{R},$$

for some $C > 0$. Let $F_\alpha \in C^1(\mathbf{R})$ be the primitive function of f_α given by $F_\alpha(s) = \int_0^s f_\alpha(t) dt$, $s \in \mathbf{R}$. From [\(3.7\)](#), it follows that

$$|F_\alpha(s)| \leq C|s|^{p_m^* - |\alpha|}, \quad s \in \mathbf{R}.$$

Now assume that the following limits exist:

$$(3.8) \quad \begin{aligned} a^+ &:= \lim_{s \rightarrow 0^+} |s|^{-p_m^* - |\alpha|} F_\alpha(s), & A^+ &:= \lim_{s \rightarrow \infty} |s|^{-p_m^* - |\alpha|} F_\alpha(s), \\ a^- &:= \lim_{s \rightarrow 0^-} |s|^{-p_m^* - |\alpha|} F_\alpha(s), & A^- &:= \lim_{s \rightarrow -\infty} |s|^{-p_m^* - |\alpha|} F_\alpha(s). \end{aligned}$$

Set limit functions of F by

$$(3.9) \quad F_{\alpha,0}(s) := \begin{cases} a^+ |s|^{p_m^* - |\alpha|} & \text{if } s \geq 0, \\ a^- |s|^{p_m^* - |\alpha|} & \text{if } s < 0, \end{cases}$$

$$(3.10) \quad F_{\alpha,\infty}(s) := \begin{cases} A^+ |s|^{p_m^* - |\alpha|} & \text{if } s \geq 0, \\ A^- |s|^{p_m^* - |\alpha|} & \text{if } s < 0. \end{cases}$$

With these functions, a decomposition of integral functionals also holds true, that is,

Theorem 3.2 (Decomposition of integral functionals 2). *Let (u_n) be a bounded sequence in $\dot{W}^{m,p}(\mathbf{R}^N)$. Assume that on a subsequence, still denoted by n , (u_n) has a profile decomposition with profile elements $(w^l, y_{l,n}, j_{l,n}, \Lambda) \in \dot{W}^{m,p}(\mathbf{R}^N) \times \mathbf{R}^N \times \mathbf{Z} \times (\mathbf{N} \cup \{0, \infty\})$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$) as in [Theorem 2.7](#). Let $\alpha \in (\mathbf{Z}_{\geq 0})^N$ be a multi-index such that $|\alpha| < m$, and let $f_\alpha \in C(\mathbf{R})$ satisfy [\(3.7\)](#) and let $F_\alpha(s) := \int_0^s f_\alpha(t) dt$, $s \in \mathbf{R}$, admit the limits [\(3.8\)](#). Define $F_{\alpha,0}$ and $F_{\alpha,\infty}$*

by (3.9) and (3.10). Then, on a subsequence again if necessary, still denoted by n , the following holds true:

$$(3.11) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} F_\alpha(\partial^\alpha u_n(x)) \, dx \\ &= \sum_{l \in \mathbf{N}_1} \int_{\mathbf{R}^N} F_\alpha\left(\partial^\alpha \left[2^{j^l N/p_m^*} w^l(2^{j^l} x)\right]\right) \, dx \\ & \quad + \sum_{l \in \mathbf{N}_2} \int_{\mathbf{R}^N} F_{\alpha, \infty}(\partial^\alpha w^l(x)) \, dx + \sum_{l \in \mathbf{N}_3} \int_{\mathbf{R}^N} F_{\alpha, 0}(\partial^\alpha w^l(x)) \, dx, \end{aligned}$$

where $\mathbf{N}_1 \cup \mathbf{N}_2 \cup \mathbf{N}_3 = \mathbf{Z}_{\geq 0}^{< \Lambda + 1}$, $j_{l,n} \rightarrow j^l$ if $l \in \mathbf{N}_1$, $j_{l,n} \rightarrow \infty$ if $l \in \mathbf{N}_2$, $j_{l,n} \rightarrow -\infty$ if $l \in \mathbf{N}_3$.

One can prove other variants of the above results in the same way as follows:

Proposition 3.3. *Along the profile decomposition in $\dot{W}^{m,p}(\mathbf{R}^N)$ ($1 < m < N/p$), there holds*

$$\lim_{n \rightarrow \infty} \|u_n\|_{\dot{W}^{k,q}(\mathbf{R}^N)}^q = \sum_{l=0}^{\Lambda} \|w^l\|_{\dot{W}^{k,q}(\mathbf{R}^N)}^q, \quad q = p_{m-k}^*, \quad k \in \mathbf{Z}_{\geq 0}, \quad k < m.$$

3.3. Proof of Theorem 3.1

We shall prove the above results in the special case $|\alpha| = 0$ and we abbreviate f_α to f and so on; the other cases will be readily shown in the same way.

Basic lemmas. We need important inequalities which play crucial roles, corresponding to (3.4) and (3.5).

Lemma 3.4. *Let $F \in \text{Lip}_{\text{loc}}(\mathbf{R})$ be as in Theorem 3.1. Then for any $L \in \mathbf{N}$, there exists a constant $C_L > 0$ such that for all $s_l \in \mathbf{R}$, $1 \leq l \leq L$,*

$$(3.12) \quad \left| F\left(\sum_{l=1}^L s_l\right) - \sum_{l=1}^L F(s_l) \right| \leq C_L \sum_{1 \leq l \neq k \leq L} |s_l| |s_k|^{p_m^* - 1}.$$

Proof. We argue by induction on L . For the sake of convenience, we set $d = 2^{N/p_m^*} > 1$, $M := [-d^2, -d^{-1}] \cup [d^{-1}, d^2] \subset \mathbf{R}$ and $L_M := \text{Lip}(M)$ which is the Lipschitz constant of F on M . Fix $\alpha > 0$ sufficiently large so that

$$1 + d^{-\alpha} \leq d, \quad 1 - d^{-\alpha} \geq d^{-1}.$$

Note that this is equivalent to

$$\alpha > \max \left\{ 0, -\frac{\log(d-1)}{\log d}, -\frac{\log(1-1/d)}{\log d} \right\}.$$

(I) Base step: $L = 2$. We shall show that there exists $C > 0$ such that for any $a, b \in \mathbf{R}$,

$$(3.13) \quad |F(a+b) - F(a) - F(b)| \leq C(|a|^{p_m^* - 1}|b| + |a||b|^{p_m^* - 1}).$$

If $a = 0$ or $b = 0$, then (3.13) is immediate, so we assume $a, b \neq 0$. Firstly, we consider the case where the ratio $t = |b|/|a|$ is so small that $t \leq d^{-\alpha}$. Also we suppose $|a| \in [1, d]$. Then one has $a+b, a \in M$ since

$$\begin{aligned} |a+b| &\leq |a| + |b| \leq d(1 + d^{-\alpha}) \leq d^2, \\ |a+b| &\geq |a| - |b| \geq 1(1 - d^{-\alpha}) \geq d^{-1}. \end{aligned}$$

It follows that

$$\begin{aligned}
 (3.14) \quad & |F(a+b) - F(a) - F(b)| \\
 & \leq |F(a+b) - F(a)| + |F(b)| \leq L_M|b| + C|b|^{p_m^*} \\
 & \leq L_M|a|^{p_m^*-1}|b| + C|a|^{p_m^*-1}|b| \leq (L_M + C)|a|^{p_m^*-1}|b|,
 \end{aligned}$$

where we used the assumptions $|a| \in [1, d]$ and $|b|/|a| \leq d^{-\alpha}$.

Then we remove the restriction $|a| \in [1, d]$, and fix arbitrary $a \in \mathbf{R}$. Then there is a unique $j = j_a \in \mathbf{Z}$ such that $|a| \in [d^j, d^{j+1}]$, and so one sees $|d^{-j}a| \in [1, d]$. Due to $t \leq d^{-\alpha}$, it follows that $d^{-j}a + d^{-j}b, d^{-j}a \in M$ since

$$\begin{aligned}
 |d^{-j}a + d^{-j}b| & \leq d^{-j}|a| + d^{-j}|b| \leq d^{-j}|a|(1 + d^{-\alpha}) \leq d(1 + d^{-\alpha}) \leq d^2, \\
 |d^{-j}a + d^{-j}b| & \geq d^{-j}|a| - d^{-j}|b| \geq d^{-j}|a|(1 - d^{-\alpha}) \geq 1(1 - d^{-\alpha}) \geq d^{-1}.
 \end{aligned}$$

Hence from (3.6) and (3.14) one sees that

$$\begin{aligned}
 & |F(a+b) - F(a) - F(b)| \\
 & \leq |F(d^j(d^{-j}a + d^{-j}b)) - F(d^j(d^{-j}a))| + |F(d^j(d^{-j}b))| \\
 & = 2^{jN}|F(d^{-j}a + d^{-j}b) - F(d^{-j}a)| + 2^{jN}|F(d^{-j}b)| \\
 & \leq 2^{jN}(L_M + C)|d^{-j}a|^{p_m^*-1}|d^{-j}b| \\
 & = (L_M + C)|a|^{p_m^*-1}|b|.
 \end{aligned}$$

Thus (3.13) holds true if $a \in \mathbf{R}$ and $t \leq d^{-\alpha}$. In the same way, one can show that

$$|F(a+b) - F(a) - F(b)| \leq (L_M + C)|a||b|^{p_m^*-1}$$

if $b \in \mathbf{R}$ and $t \geq d^\alpha$.

We finally have to verify (3.13) in the case of $a, b \in \mathbf{R}$ with $d^{-\alpha} \leq t \leq d^\alpha$. It follows that

$$\begin{aligned}
 |F(a+b) - F(a) - F(b)| & \leq |F(a+b)| + |F(a)| + |F(b)| \\
 & \leq C_p(|a|^{p_m^*} + |b|^{p_m^*}) \\
 & = C_p|a|^{p_m^*}(1 + t^{p_m^*}).
 \end{aligned}$$

There exists a constant $C > 0$ only depending on d^α such that

$$1 + t^{p_m^*} \leq C(t + t^{p_m^*-1}), \quad t \in [d^{-\alpha}, d^\alpha],$$

and thus, we get

$$\begin{aligned}
 |F(a+b) - F(a) - F(b)| & \leq C|a|^{p_m^*}(t + t^{p_m^*-1}) \\
 & = C(|a|^{p_m^*-1}|b| + |a||b|^{p_m^*-1})
 \end{aligned}$$

for all $a, b \in \mathbf{R}$ such that $t \in [d^{-\alpha}, d^\alpha]$. Eventually, (3.13) is verified for all $a, b \in \mathbf{R}$.

(II) Inductive step. Assume that (3.12) holds true for some $L \in \mathbf{N}$. Set $a = \sum_{l=1}^L s_l$. Then by the induction hypothesis and (3.13), one sees that

$$\left| F\left(\sum_{l=1}^{L+1} s_l\right) - \sum_{l=1}^{L+1} F(s_l) \right|$$

$$\begin{aligned}
&\leq |F(a + s_{L+1}) - F(a) - F(s_{L+1})| + \left| F\left(\sum_{l=1}^L s_l\right) - \sum_{l=1}^L F(s_l) \right| \\
&\leq C(|a|^{p_m^*-1}|s_{L+1}| + |a||s_{L+1}|^{p_m^*-1}) + C_L \sum_{1 \leq l \neq k \leq L} |s_l||s_k|^{p_m^*-1} \\
&\leq C_{L+1} \sum_{1 \leq l \neq k \leq L+1} |s_l||s_k|^{p_m^*-1}.
\end{aligned}$$

Hence (3.12) with L being replaced with $L + 1$ is verified. Hence (3.12) is proved for the case $L \geq 2$. The case $L = 1$ is obviously verified. Thus the proof is complete. \square

Lemma 3.5. *Let $F \in \text{Lip}_{\text{loc}}(\mathbf{R})$ be as in Theorem 3.1. Then for any $\varepsilon > 0$, there exists a constant $C = C_\varepsilon > 0$ such that for all $a, b \in \mathbf{R}$,*

$$|F(a + b) - F(a)| \leq \varepsilon |a|^{p_m^*} + C |b|^{p_m^*}.$$

Proof. Fix $\varepsilon > 0$ arbitrarily. The preceding lemma and the Young inequality yield

$$\begin{aligned}
|F(a + b) - F(a)| &\leq |F(a + b) - F(a) - F(b)| + |F(b)| \\
&\leq C(|a|^{p_m^*-1}|b| + |a||b|^{p_m^*-1}) + C|b|^{p_m^*} \\
&\leq \varepsilon |a|^{p_m^*} + C_\varepsilon |b|^{p_m^*},
\end{aligned}$$

hence the conclusion. \square

Combining the above two lemmas, one gets the following

Lemma 3.6. *Let $F \in \text{Lip}_{\text{loc}}(\mathbf{R})$ be as in Theorem 3.1 and let $L \in \mathbf{N}$. Then for any $\varepsilon > 0$, there exists $C_\varepsilon, C_L > 0$ such that for all $s_l, r \in \mathbf{R}$ ($l = 1, \dots, L$),*

$$\begin{aligned}
(3.15) \quad &\left| F\left(\sum_{l=1}^L s_l + r\right) - \sum_{l=1}^L F(s_l) \right| \\
&\leq \varepsilon \left| \sum_{l=1}^L s_l \right|^{p_m^*} + C_\varepsilon |r|^{p_m^*} + C_L \sum_{1 \leq l \neq l' \leq L} |s_l||s_{l'}|^{p_m^*-1}.
\end{aligned}$$

Main body. Now we are ready to prove the first result of decomposition of integral functionals.

Proof of Theorem 3.1. Set

$$E := \sup_{n \geq 0} \|u_n\|_{\dot{W}^{m,p}(\mathbf{R}^N)}, \quad S_n^L := \sum_{l=0}^L g_{l,n} w^l, \quad L \in \mathbf{Z}_{\geq 0}^{\leq \Lambda+1}, \quad n \in \mathbf{Z}_{\geq L},$$

and one has $u_n = S_n^L + r_n^L$. Fix $\varepsilon > 0$ arbitrarily. From (3.15), one sees that

$$\begin{aligned}
(3.16) \quad &\int_{\mathbf{R}^N} \left| F(u_n) - \sum_{l=0}^L F(g_{l,n} w^l) \right| dx \\
&\leq \varepsilon \|S_n^L\|_{L^{p_m^*}(\mathbf{R}^N)}^{p_m^*} + C_\varepsilon \|r_n^L\|_{L^{p_m^*}(\mathbf{R}^N)}^{p_m^*} \\
&\quad + C_L \sum_{1 \leq l \neq l' \leq L} \int_{\mathbf{R}^N} |g_{l,n} w^l| |g_{l',n} w^{l'}|^{p_m^*-1} dx.
\end{aligned}$$

By (2.4), (2.29) and the Sobolev inequality, we have

$$(3.17) \quad \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|S_n^L\|_{L^{p_m^*}(\mathbf{R}^N)} \leq C \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|S_n^L\|_{\dot{W}^{m,p}(\mathbf{R}^N)} \leq CE.$$

The mutual orthogonality condition (the assertion (ii) in Theorem 2.7) implies that for all $l \neq l' \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$,

$$(3.18) \quad \int_{\mathbf{R}^N} |g_{l,n} w^l| |g_{l',n} w^{l'}|^{p_m^*-1} dx \rightarrow 0$$

as $n \rightarrow \infty$. We also have

$$(3.19) \quad \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \|r_n^L\|_{L^{p_m^*}(\mathbf{R}^N)} = 0.$$

Combining (3.16)–(3.19), and passing to the limits as $n \rightarrow \infty$, $L \rightarrow \Lambda$ and then $\varepsilon \rightarrow 0$, one gets

$$\overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \int_{\mathbf{R}^N} \left| F(u_n) - \sum_{l=0}^L F(g_{l,n} w^l) \right| dx = 0.$$

By (3.6) and the change of variables, we get

$$\begin{aligned} & \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \left| \int_{\mathbf{R}^N} F(u_n) dx - \sum_{l=0}^L \int_{\mathbf{R}^N} F(w^l) dx \right| \\ & \leq \overline{\lim}_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \int_{\mathbf{R}^N} \left| F(u_n) - \sum_{l=0}^L F(g_{l,n} w^l) \right| dx \\ & = 0, \end{aligned}$$

which yields

$$\lim_{n \rightarrow \infty} \int_{\mathbf{R}^N} F(u_n) dx = \sum_{l=0}^{\Lambda} \int_{\mathbf{R}^N} F(w^l) dx.$$

This completes the proof. \square

3.4. Proof of Theorem 3.2

We now move on to the proof of Theorem 3.2. In this case, too, we shall prove the result in the special case $|\alpha| = 0$ and we abbreviate f_α to f and so on.

Basic lemma. Firstly, we provide important inequalities corresponding to Lemmas 3.4 and 3.5.

Lemma 3.7. *Let f and F be as in Theorem 3.2. Then for any $L \in \mathbf{N}$, there exists a constant $C = C_L > 0$ such that for any $s_1, \dots, s_L \in \mathbf{R}$,*

$$(3.20) \quad \left| F\left(\sum_{l=1}^L s_l\right) - \sum_{l=1}^L F(s_l) \right| \leq C \sum_{1 \leq l \neq k \leq L} |s_l| |s_k|^{p_m^*-1}.$$

Moreover, for any $\varepsilon > 0$, there exists a constant $C = C_\varepsilon > 0$ such that for all $a, b \in \mathbf{R}$,

$$(3.21) \quad |F(a+b) - F(a)| \leq \varepsilon |a|^{p_m^*} + C |b|^{p_m^*}.$$

Furthermore, for any $\varepsilon > 0$, there exists $C_\varepsilon, C_L > 0$ such that for all $s_l, r \in \mathbf{R}$ ($l = 1, \dots, L$),

$$(3.22) \quad \left| F\left(\sum_{l=1}^L s_l + r\right) - \sum_{l=1}^L F(s_l) \right|$$

$$\leq \varepsilon \left| \sum_{l=1}^L s_l \right|^{p_m^*} + C_\varepsilon |r|^{p_m^*} + C_L \sum_{1 \leq l \neq l' \leq L} |s_l| |s_{l'}|^{p_m^* - 1}.$$

Proof. The inequality (3.20) will be proved in the same way as the proofs of [13, Lemmas 4.4–4.7]. The inequality (3.21) follows from the first inequality with $L = 2$ and the Young inequality. The inequality (3.22) is proved by the above two. \square

Main body. Now we are in a position to prove [Theorem 3.2](#).

Proof of Theorem 3.2. From a similar argument to the preceding proof of [Theorem 3.1](#) together with [Lemma 3.7](#), one obtains the following:

$$(3.23) \quad \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \int_{\mathbf{R}^N} \left| F(u_n) - \sum_{l=0}^L F(g_{l,n} w^l) \right| dx = 0.$$

When $l \in \mathbf{N}_1$, there exists a limit $j^l \in \mathbf{Z}$ such that $j_{l,n} \rightarrow j^l$ as $n \rightarrow \infty$. So if $l \in \mathbf{N}_1$, then by the dominated convergence theorem, one sees that

$$(3.24) \quad \begin{aligned} & \int_{\mathbf{R}^N} F(g_{l,n} w^l) dx \\ &= \int_{\mathbf{R}^N} F(2^{j_{l,n} N / p_m^*} w^l(2^{j_{l,n}}(x))) dx \\ &= \int_{\mathbf{R}^N} F(2^{j^l N / p_m^*} w^l(2^{j^l}(x))) dx + o(1) \end{aligned}$$

as $n \rightarrow \infty$. Similarly, the dominated convergence theorem implies that

$$(3.25) \quad \begin{aligned} & \int_{\mathbf{R}^N} F(2^{j_{l,n} N / p_m^*} w^l(2^{j_{l,n}}(x - y_{l,n}))) dx \\ &= \int_{\mathbf{R}^N} F_\infty(w^l) dx + o(1) \quad (l \in \mathbf{N}_2), \end{aligned}$$

$$(3.26) \quad \begin{aligned} & \int_{\mathbf{R}^N} F(2^{j_{l,n} N / p_m^*} w^l(2^{j_{l,n}}(x - y_{l,n}))) dx \\ &= \int_{\mathbf{R}^N} F_0(w^l) dx + o(1) \quad (l \in \mathbf{N}_3), \end{aligned}$$

as $n \rightarrow \infty$.

Combining (3.24), (3.25) and (3.26), one obtains

$$\begin{aligned} & \left| \int_{\mathbf{R}^N} F(u_n) dx - \sum_{l \in \mathbf{N}_1, l \leq L} \int_{\mathbf{R}^N} F(2^{j^l N / p_m^*} w^l(2^{j^l}(x))) dx \right. \\ & \quad \left. - \sum_{l \in \mathbf{N}_2, l \leq L} \int_{\mathbf{R}^N} F_\infty(w^l) dx - \sum_{l \in \mathbf{N}_3, l \leq L} \int_{\mathbf{R}^N} F_0(w^l) dx \right| \\ & \leq \left| \int_{\mathbf{R}^N} F(u_n) dx - \sum_{l=0}^L \int_{\mathbf{R}^N} F(g_{l,n} w^l) dx \right| + o(1) \end{aligned}$$

as $n \rightarrow \infty$. From the above and (3.23), one concludes that

$$\begin{aligned} & \lim_{L \rightarrow \Lambda} \overline{\lim}_{n \rightarrow \infty} \left| \int_{\mathbf{R}^N} F(u_n) \, dx - \sum_{l \in \mathbf{N}_1, l \leq L} \int_{\mathbf{R}^N} F(2^{j^l N/p_m^*} w^l(2^{j^l}(x))) \, dx \right. \\ & \quad \left. - \sum_{l \in \mathbf{N}_2, l \leq L} \int_{\mathbf{R}^N} F_\infty(w^l) \, dx - \sum_{l \in \mathbf{N}_3, l \leq L} \int_{\mathbf{R}^N} F_0(w^l) \, dx \right| \\ & = 0, \end{aligned}$$

which leads us to (3.11). This completes the proof. \square

Acknowledgment

The author would like to thank his supervisor Goro Akagi for his great support and advice during the preparation of the paper. The author also wishes to express his gratitude to Michinori Ishiwata and Norihisa Ikoma for a lot of valuable comments on proofs and the organization of the paper. The author also wishes to thank editors and reviewers who gave many valuable comments to improve this paper.

Appendix A. Abstract theory of profile decomposition

We here recall two fundamental theorems of profile decomposition in general reflexive Banach spaces provided in the author's previous paper [13]. The following theorem is employed to prove Theorem 2.7.

Theorem A.1 ([13, Theorem 2.1]). *Let $(X, \|\cdot\|_X)$ be a reflexive Banach space, let (X, G) be a dislocation space with a dislocation group G , and let (u_n) be a bounded sequence in X . Then there exist $\Lambda \in \mathbf{N} \cup \{0, +\infty\}$, a subsequence $(N(n)) \preceq (n)$, $w^l \in X$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$) and $g_{l, N(n)} \in G$ ($l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$, $n \in \mathbf{Z}_{\geq l}$) such that*

$$u_{N(n)} = \sum_{l=0}^L g_{l, N(n)} w^l + r_{N(n)}^L, \quad L \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}, \quad n \in \mathbf{Z}_{\geq L},$$

and the following hold:

- (i) $g_{0, N(n)} = \text{Id}_X$ ($n \geq 0$), $w^l \neq 0$ ($1 \leq l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$).
- (ii) $g_{l, N(n)}^{-1} g_{k, N(n)} \rightarrow 0$ operator-weakly as $n \rightarrow \infty$ whenever $l \neq k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$.
- (iii) $g_{l, N(n)}^{-1} u_{N(n)} \rightarrow w^l$ weakly in X ($n \rightarrow \infty$, $l \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$).
- (iv) For $k \in \mathbf{Z}_{\geq 0}^{<\Lambda+1}$,

$$g_{k, N(n)}^{-1} r_{N(n)}^L \rightarrow \begin{cases} 0 & \text{if } k = 0, \dots, L, \\ w^k & \text{if } k \geq L+1, \end{cases}$$

weakly in X as $n \rightarrow \infty$.

Furthermore, if either $\Lambda = \infty$ and $\|w^l\|_X \rightarrow 0$ as $l \rightarrow \Lambda$, or else $\Lambda < \infty$, then the following exactness condition holds:

$$(A.1) \quad \lim_{L \rightarrow \Lambda} \sup_{\phi \in U} \overline{\lim}_{n \rightarrow \infty} \sup_{g \in G} \left| \left\langle \phi, g^{-1} r_{N(n)}^L \right\rangle \right| = 0,$$

where $U := B_{X^*}(1)$.

As is remarked in [13], we put the same remark here:

- Remark A.2.**
- The above theorem gives *qualitative* assertions of profile decomposition, and *quantitative* one (A.1) is obtained under a further assumption that either $\Lambda = \infty$ and $\|w^l\|_X \rightarrow 0$ as $l \rightarrow \Lambda$, or else $\Lambda < \infty$. However, this further assumption will be ensured in theorems of profile decomposition below in the present paper and in [13], by virtue of the direct calculations for the decompositions in energy like (2.2), which shows that $\|w^l\|_X \rightarrow 0$ as $l \rightarrow \infty$ if $\Lambda = \infty$.
 - As for the assumption “either $\Lambda = \infty$ and $\|w^l\|_X \rightarrow 0$ as $l \rightarrow \Lambda$, or else $\Lambda < \infty$ ”, Solimini-Tintarev [15] generally verified this further assumption, by reformulating the profile decomposition theory by means of the so-called “ Δ -convergence”. They established a Δ -convergence-version of the profile decomposition theory for uniformly convex and uniformly smooth Banach spaces, where the above further assumption always holds true, and they also showed that if the Banach space satisfies Opial’s condition, the Δ -limits coincide with the weak limits, so that the above further assumption are also true with respect to the weak-topological profile decomposition theory. It is noteworthy that Hilbert spaces, Besov spaces and Triebel-Lizorkin spaces (including Sobolev spaces) enjoys Opial’s condition, so that our assumption is satisfied in those cases.

The following theorem implies that the residual term satisfying the exactness condition (A.1) becomes arbitrarily small in a normed space Y , where the embedding $X \hookrightarrow Y$ is G -completely continuous. From the following theorem, one can prove (2.5) and (2.6).

Theorem A.3 ([13, Theorem 2.5]). *Let (X, G) be as in Theorem A.1 and let $(Y, \|\cdot\|_Y)$ be a normed space. Suppose that the embedding $X \hookrightarrow Y$ is G -completely continuous. Also assume that a double-suffix sequence (u_n^L) in X satisfies that*

$$\begin{aligned} \sup_{n, L \in \mathbf{N}} \|u_n^L\|_X &< \infty, \\ \lim_{L \rightarrow \infty} \sup_{\phi \in B_{X^*}(1)} \overline{\lim}_{n \rightarrow \infty} \sup_{g \in G} |\langle \phi, g^{-1} u_n^L \rangle| &= 0. \end{aligned}$$

Then it holds that

$$\lim_{L \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|u_n^L\|_Y = 0.$$

Appendix B. Abbreviated form of profile decomposition

In the profile decomposition theorem as before, we explicitly denoted the number of non-trivial profiles by $\Lambda \in \mathbf{Z} \cup \{0, \infty\}$, so that one can distinguish the infinite-profiles case from the finite-profiles case.

In practice, however, many profile decomposition theorems use the abbreviated form as in the introduction. Here, we provide the abbreviated version of the profile decomposition theorem.

Theorem B.1 (Abbreviated form of Theorem 2.7). *Let (u_n) be a bounded sequence in the dislocation Sobolev space $(\dot{W}^{m,p}(\mathbf{R}^N), G[\mathbf{R}^N, \mathbf{R}; p_m^*])$. Then, there exist a subsequence of (n) , still denoted by n , profiles $w^l \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($l \in \mathbf{Z}_{\geq 0}$), dislocations $(y_{l,n}, j_{l,n}) \in \mathbf{R}^N \times \mathbf{R}$ ($l, n \in$*

$\mathbf{Z}_{\geq 0}$), and residual terms $r_n^L \in \dot{W}^{m,p}(\mathbf{R}^N)$ ($L, n \in \mathbf{Z}_{\geq 0}$), with the relation of a double-suffix profile decomposition

$$u_n = \sum_{l=0}^L 2^{j_{l,n}N/p_m^*} w^l(2^{j_{l,n}}(\cdot - y_{l,n})) + r_n^L, \quad L, n \in \mathbf{Z}_{\geq 0},$$

such that the following holds true:

- (i) For all $n \in \mathbf{Z}_{\geq 0}$, $y_{0,n} = 0$ and $j_{0,n} = 0$.
- (ii) If $l \neq k$, then $|j_{l,n} - j_{k,n}| + 2^{j_{k,n}}|y_{l,n} - y_{k,n}| \rightarrow \infty$.
- (iii) For every $l \in \mathbf{Z}_{\geq 0}$, $2^{-j_{l,n}N/p_m^*} u_n(2^{-j_{l,n}} \cdot + y_{l,n}) \rightarrow w^l$ as $n \rightarrow \infty$ weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$ and a.e. on \mathbf{R}^N ($l \in \mathbf{Z}_{\geq 0}$).
- (iv) For every $k \in \mathbf{Z}_{\geq 0}$,

$$2^{-j_{k,n}N/p_m^*} r_n^L(2^{-j_{k,n}} \cdot + y_{k,n}) \rightarrow \begin{cases} 0 & \text{if } k = 0, \dots, L, \\ w^k & \text{if } k \geq L + 1, \end{cases}$$

weakly in $\dot{W}^{m,p}(\mathbf{R}^N)$ as $n \rightarrow \infty$.

- (v) There holds

$$\overline{\lim}_{n \rightarrow \infty} \|u_n\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p \geq \sum_{l=0}^{\infty} \|w^l\|_{\dot{W}^{m,p}(\mathbf{R}^N)}^p + \overline{\lim}_{L \rightarrow \infty} \overline{\lim}_{R \rightarrow \infty} \overline{\lim}_{n \rightarrow \infty} \|r_n^L\|_{\dot{W}^{m,p}(\mathbf{R}^N \setminus \mathcal{B}_{n,R,L})}^p,$$

where $\mathcal{B}_{n,R,L} := \bigcup_{l=0}^L B(y_{l,n}, 2^{-j_{l,n}} R)$.

- (vi) There holds

$$\lim_{L \rightarrow \infty} \sup_{\phi \in U} \overline{\lim}_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^N, j \in \mathbf{R}} \left| \left\langle \phi, 2^{-jN/p_m^*} r_n^L(2^{-j} \cdot + y) \right\rangle_{\dot{W}^{m,p}(\mathbf{R}^N)} \right| = 0,$$

where $U := B_{[\dot{W}^{m,p}(\mathbf{R}^N)]^*}(1)$.

Proof. Take a double-suffix profile decomposition given by [Theorem 2.7](#). At this time we renumber the subsequence and use the index n .

If $\Lambda < +\infty$, then we set

$$\begin{aligned} w^l &= 0 \quad \text{for all } l \geq \Lambda + 1, \\ y_{l,n} &= 0, \quad j_{l,n} = 0 \quad \text{for all } l \geq \Lambda + 1, \quad n \in \mathbf{Z}_{\geq 0}, \\ y_{l,n} &= 0, \quad j_{l,n} = 0 \quad \text{if } n < l. \end{aligned}$$

Then we get the above theorem.

If $\Lambda = \infty$, then we set

$$y_{l,n} = 0, \quad j_{l,n} = 0 \quad \text{if } n < l.$$

Then we get the above theorem. □

Corollary B.2. Let $u_n = \sum_{l=0}^L 2^{j_{l,n}N/p_m^*} w^l(2^{j_{l,n}}(\cdot - y_{l,n})) + r_n^L$ ($n, L \in \mathbf{Z}_{\geq 0}$) be a double-suffix profile decomposition as in the above theorem. Then this profile decomposition is the finite profile decomposition if and only if there exists $L \in \mathbf{Z}_{\geq 0}$ such that $w^l = 0$ for all $l \geq L + 1$.

Appendix C. Preliminaries

We here provide some preliminary facts which are used in the paper.

Dual spaces of Sobolev spaces. We recall that duality structures of Sobolev spaces are well characterized via the Riesz representation theorem.

Lemma C.1 (Dual space of $\dot{W}^{m,p}(\mathbf{R}^N)$). *Let $1 < p < N/m$ with $m, N \in \mathbf{N}$. Then for any $F \in [\dot{W}^{m,p}(\mathbf{R}^N)]^*$, there exist $f_\alpha \in L^{p'}(\mathbf{R}^N)$ ($\alpha \in (\mathbf{Z}_{\geq 0})^N$ with $|\alpha| = m$) such that*

$$\langle F, u \rangle = \sum_{|\alpha|=m} \int_{\mathbf{R}^N} f_\alpha \partial^\alpha u \, dx \quad \text{for all } u \in \dot{W}^{m,p}(\mathbf{R}^N).$$

Proof. Let M be the number of all multi-indices whose length is m , i.e., $M = \#\{\alpha \in (\mathbf{Z}_{\geq 0})^N; |\alpha| = m\}$. Consider the product space $X := L^p(\mathbf{R}^N)^M$ equipped with the norm $\|u\|_X = (\sum_{|\alpha|=m} \|u_\alpha\|_{L^p(\mathbf{R}^N)}^p)^{1/p}$ for $u = (u_\alpha)_{|\alpha|=m} \in X$. Define an isometric injection $\kappa : \dot{W}^{m,p}(\mathbf{R}^N) \rightarrow X$ by $\kappa(u) = (\partial^\alpha u)_{|\alpha|=m} \in X$, and set a closed subspace $Y := \kappa(\dot{W}^{m,p}(\mathbf{R}^N))$ of X .

Let $F \in [\dot{W}^{m,p}(\mathbf{R}^N)]^*$. Then a bounded linear functional $\Phi \in Y^*$ is induced by F as follows: for any $u \in Y$, $\langle \Phi, u \rangle = \langle F, \kappa^{-1}(u) \rangle$, which is bounded and $\|\Phi\|_{Y^*} \leq \|F\|_{[\dot{W}^{m,p}(\mathbf{R}^N)]^*}$. By the Hahn-Banach theorem, there is an extension $\hat{\Phi}$ of Φ onto whole X such that $\|\hat{\Phi}\|_{X^*} = \|\Phi\|_{Y^*}$, i.e., $\hat{\Phi} \in X^*$. Then by the Riesz representation theorem, there exist $\phi_\alpha \in L^{p'}(\mathbf{R}^N)$, $|\alpha| = m$, such that for all $u = (u_\alpha)_{|\alpha|=m} \in X$,

$$\langle \hat{\Phi}, u \rangle = \sum_{|\alpha|=m} \int_{\mathbf{R}^N} \phi_\alpha u_\alpha \, dx.$$

Therefore, we observe that for all $u \in \dot{W}^{m,p}(\mathbf{R}^N)$,

$$\langle F, u \rangle = \langle \Phi, \kappa(u) \rangle = \langle \hat{\Phi}, \kappa(u) \rangle = \sum_{|\alpha|=m} \int_{\mathbf{R}^N} \phi_\alpha \partial^\alpha u \, dx,$$

hence the conclusion. \square

Besov spaces. Here we recall homogeneous Besov spaces for the sake of the reader's convenience. For more details, we refer the reader to [7]. In what follows, we denote by \mathcal{F} and \mathcal{F}^{-1} the Fourier transformation and the inverse Fourier transformation, respectively, defined on $L^1(\mathbf{R}^N)$, $L^2(\mathbf{R}^N)$ or $\mathcal{S}'(\mathbf{R}^N)$, where $\mathcal{S}'(\mathbf{R}^N)$ denotes the set of tempered distributions on \mathbf{R}^N .

Let us recall the Littlewood-Paley decomposition of tempered distributions. Let $\{\phi_j\}_{j \in \mathbf{Z}}$ be a family in the Schwartz class satisfying the following properties:

$$\begin{aligned} \text{supp } \phi_j &\subset \{\xi \in \mathbf{R}^N; 2^{j-1} \leq |\xi| \leq 2^{j+1}\}; \\ \sum_{j \in \mathbf{Z}} \phi_j(\xi) &= 1, \quad \xi \neq 0; \\ \phi_j(\xi) &= \phi_0(2^{-j}\xi), \quad \xi \in \mathbf{R}^N, \quad j \in \mathbf{Z}; \\ \phi_{j-1}(\xi) + \phi_j(\xi) + \phi_{j+1}(\xi) &= 1, \quad \xi \in \text{supp } \phi_j, \quad j \in \mathbf{Z}. \end{aligned}$$

A family of operators $\{P_j\}$ is defined as follows:

$$P_j u = \mathcal{F}^{-1} \phi_j \mathcal{F} u, \quad u \in \mathcal{S}'(\mathbf{R}^N), \quad j \in \mathbf{Z}.$$

Then the Littlewood-Paley decomposition of $u \in \mathcal{S}'(\mathbf{R}^N)$ is

$$u = \sum_{j \in \mathbf{Z}} P_j u, \quad u \in \mathcal{S}'(\mathbf{R}^N).$$

Under these settings, the homogeneous Besov space $\dot{B}_{p,q}^s(\mathbf{R}^N)$ is defined as the set of all tempered distributions whose homogeneous Besov norm is finite; here the homogeneous Besov norm is defined as

$$\|u\|_{\dot{B}_{p,q}^s(\mathbf{R}^N)} := \|(\|2^{js} P_j u\|_{L^p(\mathbf{R}^N)})_{j \in \mathbf{Z}}\|_{\ell^q(\mathbf{Z})}$$

for $s \in \mathbf{R}$, $p \in [1, \infty]$, $q \in [1, \infty]$ and $u \in \mathcal{S}'(\mathbf{R}^N)$.

References

- [1] H. Bahouri, A. Cohen, G. Koch, A general wavelet-based profile decomposition in the critical embedding of function spaces, *Confluentes Math.* **3** (2011), no. 3, 387–411.
- [2] A. Bahri, J.M. Coron, On a nonlinear elliptic equation involving the critical Sobolev exponent: the effect of the topology of the domain, *Comm. Pure Appl. Math.* **41** (1988), no. 3, 253–294.
- [3] H. Brezis, J.M. Coron, Convergence of solutions of H-systems or how to blow bubbles, *Arch. Rational Mech. Anal.* **89** (1985), no. 1, 21–56.
- [4] H. Brezis, E. Lieb, A relation between pointwise convergence of functions and convergence of functionals, *Proc. Amer. Math. Soc.* **88** (1983), no. 3, 486–490.
- [5] G. Devillanova, S. Solimini, Some remarks on profile decomposition theorems, *Adv. Nonlinear Stud.* **16** (2016), no. 4, 795–805.
- [6] P. Gérard, Description du défaut de compacité de l'injection de Sobolev, *ESAIM Control Optim. Calc. Var.* **3** (1998), 213–233.
- [7] L. Grafakos, *Modern Fourier analysis. Third edition*, Graduate Texts in Mathematics, 250, Springer, New York, 2014.
- [8] S. Jaffard, Analysis of the lack of compactness in the critical Sobolev embeddings, *J. Funct. Anal.* **161** (1999), no. 2, 384–396.
- [9] P.L. Lions, The concentration-compactness principle in the calculus of variations. The locally compact case. I, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1** (1984), no. 2, 109–145.
- [10] P.L. Lions, The concentration-compactness principle in the calculus of variations. The locally compact case. II, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1** (1984), no. 4, 223–283.
- [11] P.L. Lions, The concentration-compactness principle in the calculus of variations. The limit case. I, *Rev. Mat. Iberoamericana* **1** (1985), no. 1, 145–201.
- [12] P.L. Lions, The concentration-compactness principle in the calculus of variations. The limit case. II, *Rev. Mat. Iberoamericana* **1** (1985), no. 2, 45–121.
- [13] M. Okumura, Profile decomposition in Sobolev spaces and decomposition of integral functionals I: inhomogeneous case, preprint submitted, [arXiv:2109.08176](https://arxiv.org/abs/2109.08176).
- [14] S. Solimini, A note on compactness-type properties with respect to Lorentz norms of bounded subsets of a Sobolev space, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **12** (1995), no. 3, 319–337.
- [15] S. Solimini, C. Tintarev, Concentration analysis in Banach spaces, *Commun. Contemp. Math.* **18** (2016), no. 3, 1550038, 33 pp.
- [16] M. Struwe, A global compactness result for elliptic boundary value problems involving limiting nonlinearities, *Math. Z.* **187** (1984), no. 4, 511–517.
- [17] T. Tao, Compactness and contradiction, <https://terrytao.files.wordpress.com/2011/06/blog-book.pdf>.
- [18] C. Tintarev, *Concentration compactness. Functional-analytic theory of concentration phenomena*, De Gruyter Series in Nonlinear Analysis and Applications, vol. 33, Walter de Gruyter GmbH, Berlin/Boston, 2020.
- [19] K. Tintarev, K. H. Fieseler, *Concentration compactness. Functional-analytic grounds and applications*, Imperial College Press, London, 2007.

GRADUATE SCHOOL OF SCIENCE

TOHOKU UNIVERSITY

SENDAI 980-8578

JAPAN

Email address: `okumura.mizuho.p3@dc.tohoku.ac.jp`