

Fuglede's conjecture holds in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$

Tao Zhang*

School of Mathematics and Information Science, Guangzhou University, Guangzhou 510006, China.

Abstract

Fuglede's conjecture states that for a subset Ω of a locally compact abelian group G with positive and finite Haar measure, there exists a subset of the dual group of G which is an orthogonal basis of $L^2(\Omega)$ if and only if it tiles the group by translation. In this paper, we prove a divisibility property for a set in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$. Then using the divisibility property and equi-distributed property, we prove that Fuglede's conjecture holds in the group $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$.

Key words and phrases: Fuglede's conjecture, tile, spectral set.

AMS subject classifications: 05B25, 52C20, 11P99, 42B05, 43A40.

1 Introduction

Fourier analysis can be viewed as a tool to study an arbitrary function f on (say) the reals \mathbb{R} , by looking at how such a function decomposes into a series of orthogonal basis. In this paper, we consider a generalization of this concept.

Definition 1.1. *A bounded measurable subset $\Omega \subseteq \mathbb{R}^n$ with $\mu(\Omega) > 0$ is called spectral, if there is a subset $\Lambda \subseteq \mathbb{R}^n$ such that the set of exponential functions $\{e_\lambda(x)\}_{\lambda \in \Lambda}$ is a complete orthogonal basis, where $e_\lambda(x) = e^{2\pi i \langle \lambda, x \rangle}$, that is*

$$\langle e_\lambda, e_{\lambda'} \rangle = \int_{\Omega} e_{\lambda - \lambda'}(x) dx = \delta_{\lambda, \lambda'} \mu(\Omega),$$

where $\delta_{\lambda, \lambda'} = 1$ if $\lambda = \lambda'$ and 0 otherwise, and every $f \in L^2(\Omega)$ can be expressed as

$$f(x) = \sum_{\lambda \in \Lambda} a_\lambda e_\lambda(x),$$

*Research supported by the National Natural Science Foundation of China under Grant No. 11801109.

for some $a_\lambda \in \mathbb{C}$. Λ is called the spectrum of Ω , and (Ω, Λ) is called a spectral pair in \mathbb{R}^n .

The other class of sets under study is that of tiling sets. Tiling is one of the most diverse and ubiquitous concepts of modern mathematics.

Definition 1.2. A subset $A \subseteq \mathbb{R}^n$ tiles \mathbb{R}^n by translation, if there is a translation set $T \subseteq \mathbb{R}^n$ such that almost all elements of \mathbb{R}^n have a unique representation as a sum $a + t$, where $a \in A$, $t \in T$. We will denote this by $A \oplus T = \mathbb{R}^n$. T is called the tiling complement of A , and (A, T) is called a tiling pair in \mathbb{R}^n .

In [9], Fuglede proposed the following conjecture, which connected these two notions.

Conjecture 1.3. A subset $\Omega \subseteq \mathbb{R}^n$ of positive and finite Lebesgue measure is a spectral set if and only if it tiles \mathbb{R}^n by translation.

Fuglede [9] proved this conjecture under the additional assumption that the tiling set or the spectrum is a lattice subset of \mathbb{R}^n . After some positive results, Tao [28] disproved this conjecture by constructing a spectral set in \mathbb{R}^5 which is not a tile. Later, tiles with no spectrum in \mathbb{R}^5 had been found by Kolountzakis and Matolcsi [18]. Currently, Fuglede's conjecture was disproved in \mathbb{R}^n for $n \geq 3$ in both directions [1, 6, 7, 8, 16, 17, 18, 23]. Nonetheless, some important cases had been proven to be true. Iosevich, Katz and Tao [11] proved that Fuglede's conjecture holds for convex sets in \mathbb{R}^2 , then a similar result in dimension 3 was proved by Greenfeld and Lev [10]. Recently, Lev and Matolsci [20] showed that Fuglede's conjecture holds for convex domains in \mathbb{R}^n for all n .

It is natural to formulate the following conjecture, which is in a more general setting, the locally compact abelian group.

Conjecture 1.4. Let G be a locally compact abelian group. A subset $\Omega \subseteq G$ of positive and finite Haar measure is a spectral set if and only if it is a tile.

In this general setting, the conjecture is far from being solved and is false for some specific groups as Tao [28] showed. Hence, the question becomes for which abelian group G , Fuglede's conjecture holds. In [4], Fan et al. proved that Fuglede's conjecture holds in \mathbb{Q}_p , the field of p -adic numbers. In the following of this paper, we focus on finite abelian groups. Let G be a finite abelian group, by the Fundamental Theorem of Finite Abelian Groups, $G \cong \mathbb{Z}_{m_1} \times \mathbb{Z}_{m_2} \times \cdots \times \mathbb{Z}_{m_t}$, where t is the number of generators, \mathbb{Z}_{m_i} ($i = 1, \dots, t$) is a cyclic group of order m_i , and $m_1 \mid m_2 \mid \cdots \mid m_t$, $m_1 > 1$. If the abelian group with at least 3 generators, then there exists a finite abelian group such that the Fuglede's conjecture fails for both directions [1, 6, 7, 16, 17, 18, 23]. For abelian groups with 2 generators, Fuglede's conjecture holds in \mathbb{Z}_p^2 [12], $\mathbb{Z}_p \times \mathbb{Z}_{p^2}$ [25], $\mathbb{Z}_p \times \mathbb{Z}_{pq}$ [15] and $\mathbb{Z}_{pq} \times \mathbb{Z}_{pq}$ [3], where p, q are distinct primes. For abelian groups with 1 generator, Fuglede's conjecture holds in cyclic group \mathbb{Z}_N ,

where $N = p^n, p^n q, p^n q^2, pqr, p^2 qr, pqr s$ [5, 13, 14, 19, 22, 21, 24, 26], and p, q, r, s are distinct primes.

The primary aim of this paper is to continue this investigation and our main result is the following.

Theorem 1.5. *Let p be a prime and n be a positive integer. A subset in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$ is a spectral set if and only if it is a tile of $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$.*

When we consider the Fuglede's conjecture in cyclic group \mathbb{Z}_N , one of the most important tools is the so called (T1) and (T2) conditions, which was introduced by Coven and Meyerowitz [2]. In this paper, we will show a divisibility property in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$ in Lemma 3.2, which is a weaker property than (T1) condition. However, it is very useful for bounding the size of a set. Another useful tool to investigate the Fuglede's conjecture in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$ is the equi-distributed property, which has been proven in [25]. The divisibility property and equi-distributed property are the main tools to prove Theorem 1.5.

This paper is organized as follows. In Section 2, we give some basic results about spectral sets and tiles in finite abelian groups. In Section 3, we show some properties of sets in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$. In Section 4, we prove the Tile \Rightarrow Spectral direction of Theorem 1.5, and in Section 5, we prove the Spectral \Rightarrow Tile direction of Theorem 1.5. Section 6 concludes the paper.

2 Properties of spectral sets and tiles

In this section, we give some properties of spectral sets and tiles in finite abelian groups. The following notations are fixed throughout this paper.

- Let a, b be integers such that $a \leq b$, denote $[a, b] = \{a, a + 1, \dots, b\}$.
- For any positive integer m , \mathbb{Z}_m is the ring of integers modulo m , and $\mathbb{Z}_m^* = \{a \in \mathbb{Z}_m : \gcd(a, m) = 1\}$.
- Let p be a prime, for any $t \in [0, p^m - 1]$, let $t_p[i]$ be the i -th coefficient of p -adic representation of t , i.e., $t = \sum_{i=0}^{m-1} t_p[i] p^i$, where $t_p[i] \in [0, p - 1]$. We will simply use $t[i]$ instead of $t_p[i]$ when there is no misunderstanding. Let $M(t)$ be the minimal i such that $t[i] \neq 0$, i.e.,

$$M(t) := \min\{i \in [0, m - 1] : t[i] \neq 0\}.$$

Let G be a finite abelian group, then $G \cong \mathbb{Z}_{m_1} \times \mathbb{Z}_{m_2} \times \dots \times \mathbb{Z}_{m_t}$, for some m_i with $m_1 \mid m_2 \mid \dots \mid m_t$ and $m_1 > 1$. For any $g \in G$, we will represent g as (g_1, \dots, g_t) , where $g_i \in \mathbb{Z}_{m_i}$. For any $g, h \in G$, we define their inner product by

$$\langle g, h \rangle := \sum_{i=1}^t g_i h_i \frac{m_t}{m_i}.$$

Let

$$\chi_g(h) = e^{\frac{2\pi i \cdot \langle g, h \rangle}{m_t}},$$

and $\chi_g \circ \chi_h = \chi_{g+h}$. Then the set $\widehat{G} = \{\chi_g : g \in G\}$ with \circ forms a group which is isomorphic to G .

Let $\mathbb{Z}[G]$ denote the group ring of G over \mathbb{Z} . For any $A \in \mathbb{Z}[G]$, A can be written as $A = \sum_{g \in G} a_g g$, where $a_g \in \mathbb{Z}$. For any $\chi \in \widehat{G}$, define

$$\chi(A) = \sum_{g \in G} a_g \chi(g).$$

The following inversion formula shows that A is completely determined by its character value $\chi(A)$, where χ ranges over \widehat{G} .

Lemma 2.1 ((Fourier inversion formula)). *Let G be an abelian group. If $A = \sum_{g \in G} a_g g \in \mathbb{Z}[G]$, then*

$$a_g = \frac{1}{|G|} \sum_{\chi \in \widehat{G}} \chi(A) \chi(g^{-1}),$$

for all $g \in G$.

Since we mainly work on the finite abelian group, we restate the definition of spectral set.

Definition 2.2. *Let G be a finite abelian group. A subset $A \subseteq G$ is said to be spectral if there is a subset $B \subseteq G$ such that*

$$\{\chi_b : b \in B\}$$

forms an orthogonal basis in $L^2(A)$, the vector space of complex valued functions on A with Hermitian inner product $\langle f, g \rangle = \sum_{a \in A} f(a) \overline{g(a)}$. In such case, the set B is called a spectrum of A , and (A, B) is called a spectral pair.

Since the dimension of $L^2(A)$ is $|A|$, the pair (A, B) being a spectral pair is equivalent to

$$|A| = |B| \text{ and } \chi_{b-b'}(A) = 0 \text{ for all } b \neq b' \in B.$$

The set of zeros of A is defined by

$$\mathcal{Z}_A = \{g \in G : \chi_g(A) = 0\}.$$

Lemma 2.3. *If $g \in \mathcal{Z}_A$, then $rg \in \mathcal{Z}_A$ for all $r \in \mathbb{Z}_{m_t}^*$.*

Proof. Since $g \in \mathcal{Z}_A$, then we have

$$\chi_g(A) = \sum_{h \in A} e^{\frac{2\pi i \cdot \langle g, h \rangle}{m_t}} = 0.$$

By Galois theory,

$$\chi_{rg}(A) = \sum_{h \in A} e^{\frac{2\pi i \cdot \langle rg, h \rangle}{m_t}} = \sum_{h \in A} e^{\frac{2\pi i \cdot r \cdot \langle g, h \rangle}{m_t}} = 0$$

for all $r \in \mathbb{Z}_{m_t}^*$, which means that $rg \in \mathcal{Z}_A$. □

Now we give some equivalent conditions of spectral pair.

Lemma 2.4. *Let $A, B \subseteq G$. Then the following statements are equivalent.*

- (a) (A, B) is a spectral pair.
- (b) (B, A) is a spectral pair.
- (c) $|A| = |B|$ and $(B - B) \setminus \{0\} \subseteq \mathcal{Z}_A$.
- (d) The pair $(aA + g, bB + h)$ is a spectral pair for all $a, b \in \mathbb{Z}_{m_t}^*$ and $g, h \in G$.

Proof. The equivalence of (a), (b) and (c) can be found in [25].

For (a) \Rightarrow (d), note that $|B| = |bB + h|$ and $(bB + h) - (bB + h) = b(B - B)$, by (c) and Lemma 2.3, $(A, bB + h)$ is a spectral pair. By (b), $(bB + h, A)$ is a spectral pair, then a similar discussion as above, $(aA + g, bB + h)$ is a spectral pair. For (d) \Rightarrow (a), we only need to take $a = b = 1$ and $g = h = 0$. □

Definition 2.5. *Let G be a finite abelian group. A subset $A \subseteq G$ is said to be a tile if there is a subset $T \subseteq G$ such that each element $g \in G$ can be expressed uniquely in the form*

$$g = a + t, \quad a \in A, \quad t \in T.$$

We will denote this by $G = A \oplus T$. The set T is called a tiling complement of A , and (A, T) is called a tiling pair.

The following result can be found in [25], [27, Lemma 2.1].

Lemma 2.6. *Let A, T be subsets in G . Then the following statements are equivalent.*

- (a) (A, T) is a tiling pair.
- (b) (T, A) is a tiling pair.

- (c) $(A + g, T + h)$ is a tiling pair.
- (d) $|A| \cdot |T| = |G|$ and $(A - A) \cap (T - T) = \{0\}$.
- (e) $|A| \cdot |T| = |G|$ and $\mathcal{Z}_A \cup \mathcal{Z}_T = G \setminus \{0\}$.

If $|A| = 1$ or $A = G$, then the set A is called trivial. It is easy to see that a trivial set is a spectral set and also a tiling set. In the following of this paper, we will only consider nontrivial set.

Lemma 2.7. *If (A, T) is a nontrivial tiling pair of G , then $\mathcal{Z}_A \neq \emptyset$ and $\mathcal{Z}_T \neq \emptyset$.*

Proof. If $\mathcal{Z}_A = \emptyset$, then by Lemma 2.6, $\mathcal{Z}_T = G \setminus \{0\}$. By the Fourier inversion formula (Lemma 2.1), we have $T = G$, then $|A| = 1$, which is a contradiction. Hence $\mathcal{Z}_A \neq \emptyset$. Similarly, we can get $\mathcal{Z}_T \neq \emptyset$. \square

3 Sets in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$

Recall that for $u, v \in \mathbb{Z}_p \times \mathbb{Z}_{p^n}$, their inner product is

$$\langle u, v \rangle = p^{n-1}u_1v_1 + u_2v_2 \in \mathbb{Z}_{p^n}.$$

Define

$$H(u, t) := \{x \in \mathbb{Z}_p \times \mathbb{Z}_{p^n} : \langle x, u \rangle = t\},$$

and

$$H_A(u, t) := H(u, t) \cap A,$$

for $u \in \mathbb{Z}_p \times \mathbb{Z}_{p^n}$, $t \in \mathbb{Z}_{p^n}$ and $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$. Then the following lemma, which is called equi-distributed property, can be found in [25].

Lemma 3.1. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ and $u \in \mathbb{Z}_p \times \mathbb{Z}_{p^n}$. Then $u \in \mathcal{Z}_A$ if and only if $|H_A(u, t)| = |H_A(u, t')|$ for all $t' \equiv t \pmod{p^{n-1}}$.*

Now we prove the divisibility property for a set in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$, which is useful in the following sections.

Lemma 3.2. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$. If $(a, p^{i_1}), (0, p^{i_2}), \dots, (0, p^{i_s}) \in \mathcal{Z}_A$ for some $a \in \mathbb{Z}_p$ and $0 \leq i_1 < i_2 < \dots < i_s \leq n-1$, then $p^s \mid |A|$.*

Proof. For any $2 \leq r \leq s$, $v \in [0, p^{n-1-i_r} - 1]$ and $k \in [0, p - 1]$, define

$$S(r, v, k) = \{u \in [0, p^{n-1-i_{r-1}} - 1] : u[i] = v[i] \text{ for } i \in [0, n-2-i_r] \text{ and } u[n-1-i_r] = k\}.$$

Claim: For any $a \in \mathbb{Z}_p$, $v \in [0, p^{n-1-i_r} - 1]$ and $k \in [0, p - 1]$, we have

$$\cup_{u \in S(r, v, k)} \cup_{j=0}^{p-1} H_A((a, p^{i_{r-1}}), up^{i_{r-1}} + jp^{n-1}) = H_A((0, p^{i_r}), vp^{i_r} + kp^{n-1}).$$

If $(x, y) \in \cup_{u \in S(r, v, k)} \cup_{j=0}^{p-1} H_A((a, p^{i_{r-1}}), up^{i_{r-1}} + jp^{n-1})$, then we have

$$axp^{n-1} + yp^{i_{r-1}} \equiv up^{i_{r-1}} + jp^{n-1} \pmod{p^n},$$

which leads to $y \equiv u \pmod{p^{n-1-i_{r-1}}}$. By the definition of $S(r, v, k)$,

$$yp^{i_r} \equiv vp^{i_r} + kp^{n-1} \pmod{p^n},$$

then $(x, y) \in H_A((0, p^{i_r}), vp^{i_r} + kp^{n-1})$. Hence,

$$\cup_{u \in S(r, v, k)} \cup_{j=0}^{p-1} H_A((a, p^{i_{r-1}}), up^{i_{r-1}} + jp^{n-1}) \subseteq H_A((0, p^{i_r}), vp^{i_r} + kp^{n-1}). \quad (1)$$

If $(x, y) \in H_A((0, p^{i_r}), vp^{i_r} + kp^{n-1})$, then we have

$$yp^{i_r} \equiv vp^{i_r} + kp^{n-1} \pmod{p^n},$$

which leads to $y \equiv v + kp^{n-1-i_r} \pmod{p^{n-i_r}}$. Then

$$axp^{n-1} + yp^{i_{r-1}} \equiv up^{i_{r-1}} + jp^{n-1} \pmod{p^n}$$

for some $u \in S(r, v, k)$ and $j \in [0, p - 1]$. Hence $(x, y) \in \cup_{u \in S(r, v, k)} \cup_{j=0}^{p-1} H_A((a, p^{i_{r-1}}), up^{i_{r-1}} + jp^{n-1})$. Therefore

$$\cup_{u \in S(r, v, k)} \cup_{j=0}^{p-1} H_A((a, p^{i_{r-1}}), up^{i_{r-1}} + jp^{n-1}) \supseteq H_A((0, p^{i_r}), vp^{i_r} + kp^{n-1}). \quad (2)$$

Combining (1) and (2), we have

$$\cup_{u \in S(r, v, k)} \cup_{j=0}^{p-1} H_A((a, p^{i_{r-1}}), up^{i_{r-1}} + jp^{n-1}) = H_A((0, p^{i_r}), vp^{i_r} + kp^{n-1}).$$

This ends the proof of the claim.

Since $(a, p^{i_1}), (0, p^{i_2}), \dots, (0, p^{i_s}) \in \mathcal{Z}_A$, by Lemma 3.1, we have

$$\begin{aligned} |H_A((a, p^{i_1}), up^{i_1} + jp^{n-1})| &= |H_A((a, p^{i_1}), up^{i_1} + j'p^{n-1})|, \\ |H_A((0, p^{i_r}), vp^{i_r} + jp^{n-1})| &= |H_A((0, p^{i_r}), vp^{i_r} + j'p^{n-1})| \end{aligned}$$

for all $u \in [0, p^{n-1-i_1} - 1]$, $2 \leq r \leq s$, $v \in [0, p^{n-1-i_r} - 1]$ and $0 \leq j \neq j' \leq p-1$. For any $(x, y) \in A$, we have $yp^{i_s} \equiv v_s p^{i_s} + jp^{n-1}$ for some $v_s \in [0, p^{n-1-i_s} - 1]$, $j \in [0, p-1]$. Hence

$$A = \cup_{j=0}^{p-1} \cup_{v_s \in [0, p^{n-1-i_s}-1]} H_A((0, p^{i_s}), v_s p^{i_s} + jp^{n-1}).$$

Then we have

$$\begin{aligned} |A| &= |\cup_{j=0}^{p-1} \cup_{v_s \in [0, p^{n-1-i_s}-1]} H_A((0, p^{i_s}), v_s p^{i_s} + jp^{n-1})| \\ &= p |\cup_{v_s \in [0, p^{n-1-i_s}-1]} H_A((0, p^{i_s}), v_s p^{i_s})| \\ &= p |\cup_{j=0}^{p-1} \cup_{v_{s-1} \in S(s, v_s, 0)} \cup_{v_s \in [0, p^{n-1-i_s}-1]} H_A((0, p^{i_{s-1}}), v_{s-1} p^{i_{s-1}} + jp^{n-1})| \\ &= p^2 |\cup_{v_{s-1} \in S(s, v_s, 0)} \cup_{v_s \in [0, p^{n-1-i_s}-1]} H_A((0, p^{i_{s-1}}), v_{s-1} p^{i_{s-1}})| \\ &= \dots \\ &= p^s |\cup_{v_1 \in S(2, v_2, 0)} \dots \cup_{v_{s-1} \in S(s, v_s, 0)} \cup_{v_s \in [0, p^{n-1-i_s}-1]} H_A((a, p^{i_1}), v_1 p^{i_1})|. \end{aligned}$$

Hence $p^s \mid |A|$. □

For $u, v \in \mathbb{Z}_p \times \mathbb{Z}_{p^n}$, we define the relation $u \sim v$ if there exists $r \in \mathbb{Z}_{p^n}^*$ such that $u = rv$. Then the equivalent classes in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$ by \sim are

$$(1, 0), (c, p^i) \text{ for all } c \in \mathbb{Z}_p \text{ and } i \in [0, n-1].$$

By Lemma 2.3, when we study the zero set \mathcal{Z}_A , we only need to consider the above equivalent classes.

4 Tile \Rightarrow Spectral

In this section, we will prove that if A tiles $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$, then A is a spectral set in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$. By Lemma 2.6, if A is a nontrivial tiling set in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$, then $|A| \mid p^{n+1}$, hence $|A| = p^i$ for some $1 \leq i \leq n$.

Theorem 4.1. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ be a tiling set. If $|A| = p$, then A is a spectral set.*

Proof. By Lemma 2.7, $\mathcal{Z}_A \neq \emptyset$. Let $(x, y) \in \mathcal{Z}_A$, define the set

$$B = \{r(x, y) : r \in [0, p-1]\}.$$

Note that $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$ and $|A| = |B| = p$, by Lemma 2.4, (A, B) is a spectral pair. □

Theorem 4.2. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ be a tiling set. If $|A| = p^t$, $2 \leq t \leq n$, then A is a spectral set.*

Proof. Suppose (A, T) forms a tiling pair in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$, then $|T| = p^{n-t+1}$. Let $I = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_A\}$ and $J = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_T\}$. By Lemma 3.2, we have $|I| \leq t$ and $|J| \leq n - t + 1$. On the other hand, $\mathcal{Z}_A \cup \mathcal{Z}_T = \mathbb{Z}_p \times \mathbb{Z}_{p^n} \setminus \{(0, 0)\}$, then $|I| + |J| \geq n$. Hence $t - 1 \leq |I| \leq t$.

If $|I| = t$, define

$$B = \left\{ \sum_{i \in I} s_i(0, p^i) : s_i \in [0, p-1] \right\}.$$

For any $b \neq b' \in B$, $b - b' = \sum_{i \in I} r_i(0, p^i)$, where $r_i \in [-p+1, p-1]$. Note that $\sum_{i \in I} r_i(0, p^i) \sim (0, p^j)$, where $j = \min\{i \in I : r_i \neq 0\}$, then $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$. Since $|B| = p^t = |A|$, then we have (A, B) is a spectral pair.

If $|I| = t-1$, then $|J| = n-t+1$. Suppose $I = \{a_1, a_2, \dots, a_{t-1}\}$ and $J = \{b_1, b_2, \dots, b_{n-t+1}\}$, where $0 \leq a_1 < \dots < a_{t-1} \leq n-1$, $0 \leq b_1 < \dots < b_{n-t+1} \leq n-1$ and $a_i \neq b_j$. Now we divide our discussion into three cases.

Case 1: There exist $d \in \mathbb{Z}_p$ and a_k such that $(d, p^{a_k}) \in \mathcal{Z}_T$ and $(c, p^{b_j}) \in \mathcal{Z}_T$ for all $c \in \mathbb{Z}_p$, $b_j < a_k$.

Define

$$B = \{s_0(d, p^{a_k}) + \sum_{i=1}^{n-t+1} s_i(0, p^{b_i}) : s_i \in [0, p-1]\}.$$

For any $b \neq b' \in B$, $b - b' = r_0(d, p^{a_k}) + \sum_{i=1}^{n-t+1} r_i(0, p^{b_i})$, where $r_i \in [-p+1, p-1]$. Note that

$$r_0(d, p^{a_k}) + \sum_{i=1}^{n-t+1} r_i(0, p^{b_i}) \sim \begin{cases} (d, p^{a_k}), & \text{if } r_0 \neq 0 \text{ and } r_i = 0 \text{ for all } b_i < a_k; \\ (r_l^{-1} r_0 d, p^{b_l}), & \text{if } r_0 \neq 0, b_l < a_k, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l; \\ (0, p^{b_l}), & \text{if } r_0 = 0, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l. \end{cases}$$

Then $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_T$, hence B is a spectrum for T , but $|B| = p^{n-t+2} > |T|$, which is a contradiction.

Case 2: There exist $d \in \mathbb{Z}_p$ and b_k such that $(d, p^{b_k}) \in \mathcal{Z}_A$ and $(c, p^{a_j}) \in \mathcal{Z}_A$ for all $c \in \mathbb{Z}_p$, $a_j < b_k$.

Define

$$B = \{s_0(d, p^{b_k}) + \sum_{i=1}^{t-1} s_i(0, p^{a_i}) : s_i \in [0, p-1]\}.$$

For any $b \neq b' \in B$, $b - b' = r_0(d, p^{b_k}) + \sum_{i=1}^{t-1} r_i(0, p^{a_i})$, where $r_i \in [-p+1, p-1]$. Note that

$$r_0(d, p^{b_k}) + \sum_{i=1}^{t-1} r_i(0, p^{a_i}) \sim \begin{cases} (d, p^{b_k}), & \text{if } r_0 \neq 0 \text{ and } r_i = 0 \text{ for all } a_i < b_k; \\ (r_l^{-1} r_0 d, p^{a_l}), & \text{if } r_0 \neq 0, a_l < b_k, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l; \\ (0, p^{a_l}), & \text{if } r_0 = 0, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } i < l. \end{cases}$$

Then $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$. Since $|B| = p^t = |A|$, then we have (A, B) is a spectral pair.

Case 3: $\{(c, p^{a_i}) : c \in \mathbb{Z}_p, i \in [1, t]\} \subseteq \mathcal{Z}_A$, and $\{(c, p^{b_i}) : c \in \mathbb{Z}_p, i \in [1, n - t + 1]\} \subseteq \mathcal{Z}_T$.

If $(1, 0) \in \mathcal{Z}_T$, define

$$B = \{s_0(1, 0) + \sum_{i=1}^{n-t+1} s_i(0, p^{b_i}) : s_i \in [0, p - 1]\}.$$

For any $b \neq b' \in B$, $b - b' = r_0(1, 0) + \sum_{i=1}^{n-t+1} r_i(0, p^{b_i})$, where $r_i \in [-p + 1, p - 1]$. Note that

$$r_0(1, 0) + \sum_{i=1}^{n-t+1} r_i(0, p^{b_i}) \sim \begin{cases} (1, 0), & \text{if } r_0 \neq 0 \text{ and } r_i = 0 \text{ for all } i \geq 1; \\ (r_l^{-1}r_0, p^{b_l}), & \text{if } r_0 \neq 0, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l; \\ (0, p^{b_l}), & \text{if } r_0 = 0, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l. \end{cases}$$

Then $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_T$, B is a spectrum for T , but $|B| = p^{n-t+2} > |T|$, which is a contradiction. Hence $(1, 0) \in \mathcal{Z}_A$, define

$$C = \{s_0(1, 0) + \sum_{i=1}^{t-1} s_i(0, p^{a_i}) : s_i \in [0, p - 1]\}.$$

For any $c \neq c' \in C$, $c - c' = r_0(1, 0) + \sum_{i=1}^{t-1} r_i(0, p^{a_i})$, where $r_i \in [-p + 1, p - 1]$. Note that

$$r_0(1, 0) + \sum_{i=1}^{t-1} r_i(0, p^{a_i}) \sim \begin{cases} (1, 0), & \text{if } r_0 \neq 0 \text{ and } r_i = 0 \text{ for all } i \geq 1; \\ (r_l^{-1}r_0, p^{a_l}), & \text{if } r_0 \neq 0, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l; \\ (0, p^{a_l}), & \text{if } r_0 = 0, r_l \neq 0 \text{ and } r_i = 0 \text{ for all } 1 \leq i < l. \end{cases}$$

Then $(C - C) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$. Since $|C| = p^t = |A|$, we have (A, C) is a spectral pair.

We claim that the tiling pair (A, T) satisfies at least one of Case 1, Case 2 and Case 3. Assume that Case 1 and Case 2 fail. By Lemmas 2.6 and 3.2, $\{(c, p^{a_i}) : c \in [0, p - 1], a_i < b_1\} \subseteq \mathcal{Z}_A$. Since Case 2 fails, then $\{(c, p^{b_1}) : c \in [0, p - 1]\} \subseteq \mathcal{Z}_T$. Repeating above argument, we get that Case 3 holds. This completes the proof. \square

5 Spectral \Rightarrow Tile

In this section, we will prove that if A is a spectral set in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$, then A tiles $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$. Note that $\mathcal{Z}_A \neq \emptyset$, then $p \mid |A|$.

Theorem 5.1. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ be a spectral set. If $|A| > p^n$, then $A = \mathbb{Z}_p \times \mathbb{Z}_{p^n}$.*

Proof. Suppose (A, B) is a spectral pair in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$, then $|A| = |B| > p^n$. By pigeonhole principle, there exist $(b, x), (b', x) \in B$ with $b \neq b'$. Then

$$(b, x) - (b', x) \sim (1, 0) \in \mathcal{Z}_A.$$

For any $i_0 \in [0, n-1]$ and $c_0 \in \mathbb{Z}_p$, $x \in \mathbb{Z}_p \times \mathbb{Z}_{p^n}$, x can be represented as

$$x = \sum_{i \in [0, n-1] \setminus \{i_0\}} x_i(0, p^i) + x_n(1, 0) + x_{i_0}(c_0, p^{i_0}),$$

where $x_i \in [0, p-1]$. Since $|B| > p^n$, by pigeonhole principle, there exist $u, v \in B$ such that

$$\begin{aligned} u &= \sum_{i \in [0, n-1] \setminus \{i_0\}} u_i(0, p^i) + u_n(1, 0) + u_{i_0}(c_0, p^{i_0}), \\ v &= \sum_{i \in [0, n-1] \setminus \{i_0\}} v_i(0, p^i) + v_n(1, 0) + v_{i_0}(c_0, p^{i_0}), \end{aligned}$$

with $u_{i_0} \neq v_{i_0}$, then we obtain

$$u - v \sim (c_0, p^{i_0}) \in \mathcal{Z}_A.$$

Thus, $\mathcal{Z}_A = \mathbb{Z}_p \times \mathbb{Z}_{p^n} \setminus \{(0, 0)\}$. By the Fourier inversion formula, $A = \mathbb{Z}_p \times \mathbb{Z}_{p^n}$. \square

It is easy to see that the set $A = \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ is a spectral set. Now we consider the spectral set $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ with $|A| \leq p^n$.

Theorem 5.2. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ be a spectral set. If $|A| = p$, then A is a tiling set.*

Proof. Note that $\mathcal{Z}_A \neq \emptyset$, we divide our discussion into three cases.

Case 1: $(1, 0) \in \mathcal{Z}_A$.

Define

$$T = \{(0, x) : x \in \mathbb{Z}_{p^n}\}.$$

For any $i \in [0, n-1]$, $c \in \mathbb{Z}_p$, $t \in [0, p^{n-1-i}-1]$ and $j \in [0, p-1]$, if $(0, x) \in H_T((c, p^i), tp^i + jp^{n-1})$, then $xp^i \equiv tp^i + jp^{n-1} \pmod{p^{n-i}}$, which leads to $x \equiv t + jp^{n-1-i} \pmod{p^{n-i}}$. Hence

$$|H_T((c, p^i), tp^i + jp^{n-1})| = p^i$$

for any $i \in [0, n-1]$, $c \in \mathbb{Z}_p$, $t \in [0, p^{n-1-i}-1]$ and $j \in [0, p-1]$. By Lemma 3.1,

$$(c, p^i) \in \mathcal{Z}_T \text{ for all } i \in [0, n-1], c \in \mathbb{Z}_p.$$

Then we have

$$\mathcal{Z}_A \cup \mathcal{Z}_T = (\mathbb{Z}_p \times \mathbb{Z}_{p^n}) \setminus \{(0, 0)\}.$$

Note that $|A||T| = p^{n+1}$, by Lemma 2.6, (A, T) is a tiling pair.

Case 2: there exists $s \in [0, n-1]$ such that $(0, p^s) \in \mathcal{Z}_A$.

Define

$$T = \{(x, \sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i : x, y_i \in [0, p-1]\}.$$

For any $j \in [0, p-1]$ and $t \in [0, p^{n-1}-1]$, if $(x, \sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i) \in H_T((1, 0), t + jp^{n-1})$, then

$$\begin{aligned} x &\equiv j \pmod{p}, \\ t &= 0. \end{aligned}$$

Hence

$$|H_T((1, 0), t + jp^{n-1})| = \begin{cases} p^{n-1}, & \text{if } t = 0; \\ 0, & \text{if } t \neq 0, \end{cases}$$

for any $j \in [0, p-1]$ and $t \in [0, p^{n-1}-1]$. By Lemma 3.1,

$$(1, 0) \in \mathcal{Z}_T. \quad (3)$$

For any $k \in [0, n-1] \setminus \{s\}$, $c \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k}-1]$ and $j \in [0, p-1]$, if $(x, \sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i) \in H_T((c, p^k), tp^k + jp^{n-1})$, then

$$cxp^{n-1} + (\sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i)p^k \equiv tp^k + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} cx + y_{n-1-k} &\equiv j \pmod{p}, \\ \sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i &\equiv t \pmod{p^{n-1-k}}. \end{aligned}$$

Then we have

$$|H_T((c, p^k), tp^k + jp^{n-1})| = \begin{cases} p^{k+1}, & \text{if } k < s \text{ and } t[n-s-1] = 0; \\ 0, & \text{if } k < s \text{ and } t[n-s-1] \neq 0; \\ p^k, & \text{if } k > s, \end{cases}$$

for $k \in [0, n-1] \setminus \{s\}$, $c \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k}-1]$ and $j \in [0, p-1]$. By Lemma 3.1,

$$(c, p^k) \in \mathcal{Z}_T \text{ for all } k \in [0, n-1] \setminus \{s\}, c \in \mathbb{Z}_p. \quad (4)$$

For any $c \in \mathbb{Z}_p^*$, $t \in [0, p^{n-1-s}-1]$ and $j \in [0, p-1]$, if $(x, \sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i) \in H_T((c, p^s), tp^s + jp^{n-1})$, then

$$cxp^{n-1} + \left(\sum_{i \in [0, n-1] \setminus \{n-s-1\}} y_i p^i \right) p^s \equiv tp^s + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} \sum_{i \in [0, n-s-2]} y_i p^i &\equiv t \pmod{p^{n-s-1}}, \\ cx &\equiv j \pmod{p}. \end{aligned}$$

Then we have

$$|H_T((c, p^s), tp^s + jp^{n-1})| = p^s$$

for all $c \in \mathbb{Z}_p^*$, $t \in [0, p^{n-1-s}-1]$ and $j \in [0, p-1]$. By Lemma 3.1,

$$(c, p^s) \in \mathcal{Z}_T \text{ for all } c \in \mathbb{Z}_p^*. \quad (5)$$

Combining (3), (4) and (5), we have

$$\mathcal{Z}_A \cup \mathcal{Z}_T = (\mathbb{Z}_p \times \mathbb{Z}_{p^n}) \setminus \{(0, 0)\}.$$

Note that $|A||T| = p^{n+1}$, by Lemma 2.6, (A, T) is a tiling pair.

Case 3: there exist $c \in \mathbb{Z}_p^*$ and $s \in [0, n-1]$ such that $(c, p^s) \in \mathcal{Z}_A$.

Define

$$T = \{(-c^{-1}y_{n-s-1}, \sum_{i=0}^{n-1} y_i p^i) : y_i \in [0, p-1]\}.$$

For any $t \in [0, p^{n-1}-1]$ and $j \in [0, p-1]$, if $(-c^{-1}y_{n-s-1}, \sum_{i=0}^{n-1} y_i p^i) \in H_T((1, 0), t + jp^{n-1})$, then

$$\begin{aligned} -c^{-1}y_{n-s-1} &\equiv j \pmod{p}, \\ t &= 0. \end{aligned}$$

Hence

$$|H_T((1, 0), t + jp^{n-1})| = \begin{cases} p^{n-1}, & \text{if } t = 0; \\ 0, & \text{if } t \neq 0, \end{cases}$$

for any $t \in [0, p^{n-1}-1]$ and $j \in [0, p-1]$. By Lemma 3.1,

$$(1, 0) \in \mathcal{Z}_T. \quad (6)$$

For any $k \in [0, n-1] \setminus \{s\}$, $d \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k}-1]$ and $j \in [0, p-1]$, if $(-c^{-1}y_{n-s-1}, \sum_{i=0}^{n-1} y_i p^i) \in H_T((d, p^k), tp^k + jp^{n-1})$, then

$$-dc^{-1}y_{n-s-1}p^{n-1} + \left(\sum_{i=0}^{n-1} y_i p^i\right)p^k \equiv tp^k + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} -dc^{-1}y_{n-s-1} + y_{n-1-k} &\equiv j \pmod{p}, \\ \sum_{i=0}^{n-2-k} y_i p^i &\equiv t \pmod{p^{n-1-k}}. \end{aligned}$$

Then we have

$$|H_T((d, p^k), tp^k + jp^{n-1})| = p^k$$

for $k \in [0, n-1] \setminus \{s\}$, $d \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k}-1]$ and $j \in [0, p-1]$. By Lemma 3.1,

$$(d, p^k) \in \mathcal{Z}_T \text{ for all } k \in [0, n-1] \setminus \{s\}, d \in \mathbb{Z}_p. \quad (7)$$

For any $d \in \mathbb{Z}_p$, $d \neq c$, $t \in [0, p^{n-1-s}-1]$ and $j \in [0, p-1]$, if $(-c^{-1}y_{n-s-1}, \sum_{i=0}^{n-1} y_i p^i) \in H_T((d, p^s), tp^s + jp^{n-1})$, then

$$-dc^{-1}y_{n-s-1}p^{n-1} + \left(\sum_{i=0}^{n-1} y_i p^i\right)p^s \equiv tp^s + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} \sum_{i=0}^{n-s-2} y_i p^i &\equiv t \pmod{p^{n-s-1}}, \\ -dc^{-1}y_{n-s-1} + y_{n-s-1} &\equiv (1 - dc^{-1})y_{n-s-1} \equiv j \pmod{p}. \end{aligned}$$

Then we have

$$|H_T((d, p^s), tp^s + jp^{n-1})| = p^s,$$

for all $d \in \mathbb{Z}_p$ with $d \neq c$, $t \in [0, p^{n-1-s}-1]$ and $j \in [0, p-1]$. By Lemma 3.1,

$$(d, p^s) \in \mathcal{Z}_T \text{ for all } d \in \mathbb{Z}_p, d \neq c. \quad (8)$$

Combining (6), (7) and (8), we have

$$\mathcal{Z}_A \cup \mathcal{Z}_T = (\mathbb{Z}_p \times \mathbb{Z}_{p^n}) \setminus \{(0, 0)\}.$$

Note that $|A||T| = p^{n+1}$, by Lemma 2.6, (A, T) is a tiling pair. \square

Theorem 5.3. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$ be a spectral set. If $|A| = p^s$ with $2 \leq s \leq n$, then A is a tiling set.*

Proof. Suppose (A, B) is a spectral pair, then $|A| = |B| = p^s$. Let $I = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_A\}$ and $J = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_B\}$. Since $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$ and $(A - A) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_B$, then $|I| \geq s-1$ and $|J| \geq s-1$. By Lemma 3.2, we have $s-1 \leq |I| \leq s$ and $s-1 \leq |J| \leq s$.

Case 1: $|I| = s$.

Define

$$T = \{(x, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) : x, y_i \in [0, p-1]\}.$$

For any $t \in [0, p^{n-1} - 1]$ and $j \in [0, p-1]$, if $(x, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((1, 0), t + jp^{n-1})$, then

$$\begin{aligned} x &\equiv j \pmod{p}, \\ t &= 0. \end{aligned}$$

Hence

$$|H_T((1, 0), t + jp^{n-1})| = \begin{cases} p^{n-s}, & \text{if } t = 0; \\ 0, & \text{if } t \neq 0, \end{cases}$$

for any $j \in [0, p-1]$ and $t \in [0, p^{n-1} - 1]$. By Lemma 3.1,

$$(1, 0) \in \mathcal{Z}_T. \tag{9}$$

For any $k \in I$, $c \in \mathbb{Z}_p^*$, $t \in [0, p^{n-1-k} - 1]$ and $j \in [0, p-1]$, if $(x, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((c, p^k), tp^k + jp^{n-1})$, then

$$xcp^{n-1} + (\sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i})p^k \equiv tp^k + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} xc &\equiv j \pmod{p}, \\ \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} &\equiv t \pmod{p^{n-1-k}}. \end{aligned}$$

Then we have

$$|H_T((c, p^k), tp^k + jp^{n-1})| = \begin{cases} p^{t_0}, & \text{if } t[i] = 0 \text{ for } i < n-1-k, n-1-i \in I; \\ 0, & \text{otherwise,} \end{cases}$$

for $k \in I$, $c \in \mathbb{Z}_p^*$, $t \in [0, p^{n-1-k} - 1]$ and $j \in [0, p - 1]$, where $t_0 = |\{i \in [0, n - 1] \setminus I : i < k\}|$. By Lemma 3.1,

$$(c, p^k) \in \mathcal{Z}_T \text{ for all } k \in I, c \in \mathbb{Z}_p^*. \quad (10)$$

For any $k \in [0, n - 1] \setminus I$, $c \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k} - 1]$ and $j \in [0, p - 1]$, if $(x, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((c, p^k), tp^k + jp^{n-1})$, then

$$cxp^{n-1} + \left(\sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} \right) p^k \equiv tp^k + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} &\equiv t \pmod{p^{n-1-k}}, \\ cx + y_k &\equiv j \pmod{p}. \end{aligned}$$

Then we have

$$|H_T((c, p^k), tp^k + jp^{n-1})| = \begin{cases} p^{t_0}, & \text{if } t[i] = 0 \text{ for } i < n - 1 - k, n - 1 - i \in I; \\ 0, & \text{otherwise,} \end{cases}$$

for all $k \in [0, n - 1] \setminus I$, $c \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k} - 1]$ and $j \in [0, p - 1]$, where $t_0 = 1 + |\{i \in [0, n - 1] \setminus I : i < k\}|$. By Lemma 3.1,

$$(c, p^k) \in \mathcal{Z}_T \text{ for all } k \in [0, n - 1] \setminus I, c \in \mathbb{Z}_p. \quad (11)$$

Combining (9), (10) and (11), we have

$$\mathcal{Z}_A \cup \mathcal{Z}_T = (\mathbb{Z}_p \times \mathbb{Z}_{p^n}) \setminus \{(0, 0)\}.$$

Note that $|A||T| = p^{n+1}$, by Lemma 2.6, (A, T) is a tiling pair.

Case 2: $|J| = s - 1$.

Define

$$T = \{(0, \sum_{i \in [0, n-1] \setminus J} y_i p^i) : y_i \in [0, p - 1]\}.$$

For any $(a_1, x_1) \neq (a_2, x_2) \in A$ and $(0, \sum_{i \in [0, n-1] \setminus J} y_i p^i) \neq (0, \sum_{i \in [0, n-1] \setminus J} z_i p^i) \in T$, if

$$(a_1, x_1) - (a_2, x_2) = (0, \sum_{i \in [0, n-1] \setminus J} y_i p^i) - (0, \sum_{i \in [0, n-1] \setminus J} z_i p^i),$$

then

$$a_1 = a_2,$$

$$x_1 - x_2 = \sum_{i \in [0, n-1] \setminus J} (y_i - z_i) p^i.$$

Since $J = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_B\}$ and $(A - A) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_B$, then $M(x_1 - x_2) \in J$, which is contradicting to $M(\sum_{i \in [0, n-1] \setminus J} (y_i - z_i) p^i) \in [0, n-1] \setminus J$. Hence $(A - A) \cap (T - T) = \{(0, 0)\}$. Note that $|A||T| = p^{n+1}$, by Lemma 2.6, (A, T) is a tiling pair.

Case 3: $|I| = s - 1$ and $|J| = s$.

Since $|B| = p^s$, $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$ and $|I| = s - 1$. Let

$$B_t = \{(x, y) \in B : x = t\},$$

then for any $(t, y), (t, y') \in B_t$, we have $M(y - y') \in I$. Then $|B_t| \leq p^{s-1}$. But $p^s = |B| = |\cup_{t=0}^{p-1} B_t| \leq p^s$, hence $|B_t| = p^{s-1}$ for all $t \in [0, p-1]$. Thus, for any $i \in I$, there exist $(t, y), (t, y') \in B_t$ such that $M(y - y') = i$. Since $|J| > |I|$, then there exists $j_0 \in J$ such that $n-1-j_0 \notin I$. Recall that $n-1-j_0 \notin I$, then for any $(t, x), (t, x') \in B$, if $x \equiv x' \pmod{p^{n-1-j_0}}$, then $x \equiv x' \pmod{p^{n-j_0}}$. Note that $(0, p^{j_0}) \in \mathcal{Z}_B$ and

$$\langle (t, x), (0, p^{j_0}) \rangle = xp^{j_0},$$

then for any $(0, x_0) \in B$, there exist $(1, x_1), \dots, (p-1, x_{p-1}) \in B$ such that

$$\{x_i - x_0 : i \in [0, p-1]\} \equiv \{i \cdot p^{n-1-j_0} : i \in [0, p-1]\} \pmod{p^{n-j_0}}.$$

Therefore, for any $i \in I$ with $i < n-1-j_0$, there exist $(0, x), (0, x_0), (1, x_1), \dots, (p-1, x_{p-1}) \in B$ such that

$$M(x - x_0) = i,$$

$$\{x_j - x_0 : j \in [0, p-1]\} \equiv \{i \cdot p^{n-1-j_0} : i \in [0, p-1]\} \pmod{p^{n-j_0}}.$$

Then $(j, x_j - x) \in B - B$ and

$$\{(j, x_j - x) : j \in [0, p-1]\} \sim \{(d, p^i) : d \in \mathbb{Z}_p\}.$$

Hence we have

$$(d, p^i) \in \mathcal{Z}_A \text{ for all } d \in \mathbb{Z}_p, i \in I \text{ with } i < n-1-j_0. \quad (12)$$

As we have proved, there exist $(t, x), (t', x') \in B$ such that $t \neq t'$ and $x - x' \equiv c'p^{n-1-j_0} \pmod{p^{n-j_0}}$ for some $c' \in [1, p-1]$. Denote $c := c' \cdot (t - t')^{-1}$, then $(t, x) - (t', x') \sim (c^{-1}, p^{n-1-j_0}) \in \mathcal{Z}_A$. Define

$$T = \{(-cy_{n-1-j_0}, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) : y_i \in [0, p-1]\}.$$

For any $t \in [0, p^{n-1} - 1]$ and $j \in [0, p-1]$, if $(-cy_{n-1-j_0}, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((1, 0), t + jp^{n-1})$, then

$$\begin{aligned} -cy_{n-1-j_0} &\equiv j \pmod{p}, \\ t &= 0. \end{aligned}$$

Hence

$$|H_T((1, 0), t + jp^{n-1})| = \begin{cases} p^{n-s}, & \text{if } t = 0; \\ 0, & \text{if } t \neq 0, \end{cases}$$

for any $j \in [0, p-1]$ and $t \in [0, p^{n-1} - 1]$. By Lemma 3.1,

$$(1, 0) \in \mathcal{Z}_T. \quad (13)$$

For any $k \in I$ with $k > n-1-j_0$, $d \in \mathbb{Z}_p^*$, $t \in [0, p^{n-1-k} - 1]$ and $j \in [0, p-1]$, if $(-cy_{n-1-j_0}, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((d, p^k), tp^k + jp^{n-1})$, then

$$-dcy_{n-1-j_0}p^{n-1} + \left(\sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} \right) p^k \equiv tp^k + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} -dcy_{n-1-j_0} &\equiv j \pmod{p}, \\ \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} &\equiv t \pmod{p^{n-1-k}}. \end{aligned}$$

Then we have

$$|H_T((d, p^k), tp^k + jp^{n-1})| = \begin{cases} p^{t_0}, & \text{if } t[i] = 0 \text{ for } i < n-1-k, n-1-i \in I; \\ 0, & \text{otherwise,} \end{cases}$$

for $k \in I$ with $k > n-1-j_0$, $d \in \mathbb{Z}_p^*$, $t \in [0, p^{n-1-k} - 1]$ and $j \in [0, p-1]$, where $t_0 = |\{i \in [0, n-1] \setminus I : i < k\}| - 1$. By Lemma 3.1,

$$(d, p^k) \in \mathcal{Z}_T \text{ for all } k \in I \text{ with } k > n-1-j_0, d \in \mathbb{Z}_p^*. \quad (14)$$

For any $k \in [0, n-1] \setminus I$ with $k \neq n-1-j_0$, $d \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k}-1]$ and $j \in [0, p-1]$, if $(-cy_{n-1-j_0}, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((d, p^k), tp^k + jp^{n-1})$, then

$$-dcy_{n-1-j_0}p^{n-1} + \left(\sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} \right) p^k \equiv tp^k + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} &\equiv t \pmod{p^{n-1-k}}, \\ -dcy_{n-1-j_0} + y_k &\equiv j \pmod{p}. \end{aligned}$$

Then we have

$$|H_T((d, p^k), tp^k + jp^{n-1})| = \begin{cases} p^{t_0}, & \text{if } t[i] = 0 \text{ for } i < n-1-k, n-1-i \in I; \\ 0, & \text{otherwise,} \end{cases}$$

for all $k \in [0, n-1] \setminus I$ with $k \neq n-1-j_0$, $d \in \mathbb{Z}_p$, $t \in [0, p^{n-1-k}-1]$ and $j \in [0, p-1]$, where $t_0 = |\{i \in [0, n-1] \setminus I : i < k\}|$. By Lemma 3.1,

$$(d, p^k) \in \mathcal{Z}_T \text{ for all } k \in [0, n-1] \setminus I \text{ with } k \neq n-1-j_0, d \in \mathbb{Z}_p. \quad (15)$$

For any $d \in \mathbb{Z}_p$ with $d \neq c^{-1}$, $t \in [0, p^{j_0}-1]$ and $j \in [0, p-1]$, if $(-cy_{n-1-j_0}, \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i}) \in H_T((d, p^{n-1-j_0}), tp^{n-1-j_0} + jp^{n-1})$, then

$$-dcy_{n-1-j_0}p^{n-1} + \left(\sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} \right) p^{n-1-j_0} \equiv tp^{n-1-j_0} + jp^{n-1} \pmod{p^n},$$

which leads to

$$\begin{aligned} \sum_{i \in [0, n-1] \setminus I} y_i p^{n-1-i} &\equiv t \pmod{p^{j_0}}, \\ (1-dc)y_{n-1-j_0} &\equiv j \pmod{p}. \end{aligned}$$

Then we have

$$|H_T((d, p^{n-1-j_0}), tp^{n-1-j_0} + jp^{n-1})| = \begin{cases} p^{t_0}, & \text{if } t[i] = 0 \text{ for } i < j_0, n-1-i \in I; \\ 0, & \text{otherwise,} \end{cases}$$

for all $d \in \mathbb{Z}_p$ with $d \neq c^{-1}$, $t \in [0, p^{j_0}-1]$ and $j \in [0, p-1]$, where $t_0 = |\{i \in [0, n-1] \setminus I : i < n-1-j_0\}|$. By Lemma 3.1,

$$(d, p^{n-1-j_0}) \in \mathcal{Z}_T \text{ for all } d \in \mathbb{Z}_p \text{ with } d \neq c^{-1}. \quad (16)$$

Combining (12), (13), (14), (15) and (16), we have

$$\mathcal{Z}_A \cup \mathcal{Z}_T = (\mathbb{Z}_p \times \mathbb{Z}_{p^n}) \setminus \{(0, 0)\}.$$

Note that $|A||T| = p^{n+1}$, by Lemma 2.6, (A, T) is a tiling pair. \square

Theorem 5.4. *Let $A \subseteq \mathbb{Z}_p \times \mathbb{Z}_{p^n}$. If $|A| = mp^s$ with $2 \leq m \leq p-1$, $1 \leq s \leq n-1$, then A is not a spectral set.*

Proof. Suppose (A, B) is a spectral pair, then $|A| = |B| = mp^s$. Let $I = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_A\}$ and $J = \{i \in [0, n-1] : (0, p^i) \in \mathcal{Z}_B\}$. Since $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$ and $(A - A) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_B$, then $|I| = |J| = s$. Now we divide our discussion into two cases.

Case 1: There exists $r \in [0, n-1]$ such that $n-1-r \notin I$ and $r \notin J$.

We prove this case by induction. It has been proven that Fuglede's conjecture holds in $\mathbb{Z}_p \times \mathbb{Z}_p$ and $\mathbb{Z}_p \times \mathbb{Z}_{p^2}$ [12, 25]. We assume that Fuglede's conjecture holds in $\mathbb{Z}_p \times \mathbb{Z}_{p^i}$ for all $i \leq n-1$, now we consider the Fuglede's conjecture in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$.

Let $\phi_1, \phi_2 : \mathbb{Z}_{p^n} \mapsto \mathbb{Z}_{p^{n-1}}$ be defined by

$$\begin{aligned} \phi_1\left(\sum_{i=0}^{n-1} c_i p^i\right) &= \sum_{i=0}^{r-1} c_i p^i + \sum_{i=r}^{n-2} c_{i+1} p^i, \\ \phi_2\left(\sum_{i=0}^{n-1} c_i p^i\right) &= \sum_{i=0}^{n-2-r} c_i p^i + \sum_{i=n-1-r}^{n-2} c_{i+1} p^i. \end{aligned}$$

Let

$$A' = \{(a, \phi_1(x)) : (a, x) \in A\},$$

and

$$B' = \{(b, \phi_2(y)) : (b, y) \in B\}.$$

Since $(0, p^{n-1-r}) \notin \mathcal{Z}_A$, $(0, p^r) \notin \mathcal{Z}_B$ and $(B - B) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$, $(A - A) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_B$, then $|A'| = |A|$ and $|B'| = |B|$. In the following, we will show that (A', B') forms a spectral pair in $\mathbb{Z}_p \times \mathbb{Z}_{p^{n-1}}$.

Let

$$i' = \begin{cases} i, & \text{if } i < n-1-r; \\ i-1, & \text{if } i > n-1-r. \end{cases}$$

Note that for any $(b_1, y), (b_2, w) \in B$, we have

$$(b_1, \phi_2(y)) - (b_2, \phi_2(w)) \sim \begin{cases} (1, 0), & \text{if } (b_1 - b_2, y - w) \sim (1, 0); \\ (0, p^{i'}), & \text{if } (b_1 - b_2, y - w) \sim (0, p^i); \\ (c, p^{i'}), & \text{if } (b_1 - b_2, y - w) \sim (c, p^i) \text{ and } i \neq n - 1 - r; \\ (d, p^j), & \text{if } (b_1 - b_2, y - w) \sim (c, p^{n-1-r}), y - w \equiv c_1 p^{n-1-r} \pmod{p^{n-r}} \\ & \text{and } M(y - w - c_1 p^{n-1-r}) = j + 1; \\ (1, 0), & \text{if } (b_1 - b_2, y - w) \sim (c, p^{n-1-r}) \text{ and } y - w = c_1 p^{n-1-r}, \end{cases}$$

for some $c, c_1, d \in \mathbb{Z}_p$.

If $(1, 0) \in \mathcal{Z}_A$, then for any $t \in [0, p - 1]$, we have $|\{(a, x) \in A : a = t\}| = \frac{|A|}{p}$, hence $|\{(a, \phi_1(x)) \in A' : a = t\}| = \frac{|A'|}{p}$. Therefore,

$$(1, 0) \in \mathcal{Z}_{A'} \text{ if } (1, 0) \in \mathcal{Z}_A. \quad (17)$$

If $i \in I$ and $(a, x) \in H_A((0, p^i), up^i + jp^{n-1})$, then $xp^i \equiv up^i + jp^{n-1} \pmod{p^n}$. It follows that $\phi_1(x)p^{i'} \equiv \phi_1(u)p^{i'} + jp^{n-2} \pmod{p^{n-1}}$, hence

$$(0, p^{i'}) \in \mathcal{Z}_{A'} \text{ if } (0, p^i) \in \mathcal{Z}_A. \quad (18)$$

If $i \neq n - 1 - r$ and $(a, x) \in H_A((c, p^i), up^i + jp^{n-1})$, then $acp^{n-1} + xp^i \equiv up^i + jp^{n-1} \pmod{p^n}$. It follows that $acp^{n-2} + \phi_1(x)p^{i'} \equiv \phi_1(u)p^{i'} + jp^{n-2} \pmod{p^{n-1}}$, hence

$$(c, p^{i'}) \in \mathcal{Z}_{A'} \text{ if } (c, p^i) \in \mathcal{Z}_A, i \neq n - 1 - r. \quad (19)$$

If $(c, p^{n-1-r}) \in \mathcal{Z}_A$ for some $c \in \mathbb{Z}_p^*$, note that

$$\langle (a, x), (c, p^{n-1-r}) \rangle_n = acp^{n-1} + xp^{n-1-r}$$

and for any $(a, x_1), (a, x_2) \in A$, we have $M(x_1 - x_2) \neq r$, then for any $(a, x_a) \in A$, there exist $(i, x_i), i \in [0, p - 1] \setminus \{a\}$ such that

$$x_i \equiv x_a \pmod{p^r} \text{ for } i \in [0, p - 1].$$

Then A can be represented as

$$A = \cup_{t=0}^{mp^{s-1}-1} A_t,$$

where $A_t = \{(a, x_{a,t}) : a \in [0, p - 1]\}$ and for any $(a, x_{a,t}), (b, x_{b,t}) \in A_t, x_{a,t} \equiv x_{b,t} \pmod{p^r}$. For any $t \in [0, mp^{s-1} - 1]$, we can compute to get that

$$\begin{aligned} \chi_{(1,0)}(A_t) &= 0, \\ \chi_{(c,p^i)}(A_t) &= 0 \text{ for all } c \in \mathbb{Z}_p^* \text{ and } i \in [n - r, n - 1]. \end{aligned}$$

It follows that

$$\begin{aligned}\chi_{(1,0)}(A) &= 0, \\ \chi_{(d,p^i)}(A) &= 0 \text{ for all } d \in \mathbb{Z}_p^* \text{ and } i \in [n-r, n-1].\end{aligned}$$

Hence

$$(1, 0), (d, p^i) \in \mathcal{Z}_A \text{ for all } d \in \mathbb{Z}_p^* \text{ and } i \in [n-r, n-1], \text{ if } (c, p^{n-1-r}) \in \mathcal{Z}_A \text{ for some } c \in \mathbb{Z}_p^*. \quad (20)$$

Combining (17), (18), (19) and (20), we have (A', B') forms a spectral pair in $\mathbb{Z}_p \times \mathbb{Z}_{p^{n-1}}$, which is a contradiction.

Case 2: There does not exist $r \in [0, n-1]$ such that $n-1-r \notin I$ and $r \notin J$.

Claim: there is at most one i such that $i \in I$ and $n-1-i \notin J$. Similarly, there is at most one j such that $j \in J$ and $n-1-j \notin I$.

Assume that there exist $i_1, i_2 \in I$ and $n-1-i_1, n-1-i_2 \notin J$. Since $n-1-i_1 \notin J$ and $(A-A) \setminus \{(0,0)\} \subseteq \mathcal{Z}_B$, then for any $(t, x), (t, x') \in A$, if $x \equiv x' \pmod{p^{n-1-i_1}}$, then $x \equiv x' \pmod{p^{n-i_1}}$. Note that $(0, p^{i_1}) \in \mathcal{Z}_A$ and

$$\langle (t, x), (0, p^{i_1}) \rangle = xp^{i_1},$$

then for any $(0, x_0) \in A$, there exist $(1, x_1), \dots, (p-1, x_{p-1}) \in A$ such that

$$\{x_i - x_0 : i \in [0, p-1]\} \equiv \{i \cdot p^{n-1-i_1} : i \in [0, p-1]\} \pmod{p^{n-i_1}}.$$

Similarly, there exist $(1, y_1), \dots, (p-1, y_{p-1}) \in A$ such that

$$\{y_i - x_0 : i \in [0, p-1]\} \equiv \{i \cdot p^{n-1-i_2} : i \in [0, p-1]\} \pmod{p^{n-i_2}}.$$

If $i_1 > i_2$, then

$$(0, y_1 - x_1) \sim (0, p^{n-1-i_1}) \in \mathcal{Z}_B,$$

which is a contradiction. Similarly, we can get a contradiction for the case $i_1 < i_2$. This ends the proof of the claim.

By the claim, we have $s = n-1$, i.e., $|A| = |B| = mp^{n-1}$. We may assume $I = [0, n-1] \setminus \{i_0\}$ and $J = [0, n-1] \setminus \{j_0\}$, then $n-1-i_0 \in J$ and $n-1-j_0 \in I$.

Since $i_0 \notin I$, then for any $(b, y), (b, y') \in B$, if $y \equiv y' \pmod{p^{i_0}}$, then $y \equiv y' \pmod{p^{i_0+1}}$. Note that $(0, p^{n-1-i_0}) \in \mathcal{Z}_B$ and

$$\langle (b, y), (0, p^{n-1-i_0}) \rangle = yp^{n-1-i_0},$$

then for any $(0, y_0) \in B$, there exist $(1, y_1), \dots, (p-1, y_{p-1}) \in B$ such that

$$\{y_i - y_0 : i \in [0, p-1]\} \equiv \{i \cdot p^{i_0} : i \in [0, p-1]\} \pmod{p^{i_0+1}}.$$

Hence

$$(1, y_1 - y_0) \sim (c_0, p^{i_0}) \in \mathcal{Z}_A \text{ for some } c_0 \in \mathbb{Z}_p^*. \quad (21)$$

Since $|B| = mp^{n-1}$, then for any $k < i_0$, there exist $(t, y), (t, y_t) \in B$ such that $M(y_t - y) = k$, and there exist $(j, y_j) \in B, j \in [0, p-1] \setminus \{t\}$ such that

$$\{y_j - y_t : j \in [0, p-1]\} \equiv \{i \cdot p^{i_0} : i \in [0, p-1]\} \pmod{p^{i_0+1}}.$$

Then $(j - t, y_j - y) \in B - B$ and

$$\{(j - t, y_j - y) : j \in [0, p-1]\} \sim \{(c, p^k) : c \in \mathbb{Z}_p\}.$$

Hence we have

$$(c, p^k) \in \mathcal{Z}_A \text{ for all } c \in \mathbb{Z}_p, k < i_0. \quad (22)$$

Define

$$D = \left\{ \sum_{i \in I} a_i(0, p^i) + a_{i_0}(c_0, p^{i_0}) : a_i \in [0, p-1] \right\}.$$

For any $d \neq d' \in D, d - d' = \sum_{i \in I} d_i(0, p^i) + d_{i_0}(c_0, p^{i_0})$, where $d_i \in [-p+1, p-1]$. Note that

$$\sum_{i \in I} d_i(0, p^i) + d_{i_0}(c_0, p^{i_0}) \sim \begin{cases} (c_0, p^{i_0}), & \text{if } d_{i_0} \neq 0 \text{ and } d_i = 0 \text{ for all } i < i_0; \\ (d_k^{-1} d_{i_0} c_0, p^k), & \text{if } d_{i_0} \neq 0, d_k \neq 0, k < i_0 \text{ and } d_i = 0 \text{ for all } i < k; \\ (0, p^k), & \text{if } d_{i_0} = 0, d_k \neq 0 \text{ and } d_i = 0 \text{ for all } i < k. \end{cases}$$

By (21) and (22), we have $(D - D) \setminus \{(0, 0)\} \subseteq \mathcal{Z}_A$, then D is a spectrum for A , but $|D| = p^n > |A|$, which is a contradiction. This completes the proof. \square

6 Conclusion

From the proof of Theorem 1.5, we can see that the divisibility property and equi-distributed property in $\mathbb{Z}_p \times \mathbb{Z}_{p^n}$ play an important role. In [25], the author had proved that the equi-distributed property holds in $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$. Actually, the divisibility property can also be generalized to the group $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$.

Lemma 6.1. *Let m, n be integers with $m \leq n$. Let $A \subseteq \mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$. If $(a, p^{i_1}), (0, p^{i_2}), \dots, (0, p^{i_s}) \in \mathcal{Z}_A$ for some $a \in p^{m-1}\mathbb{Z}_{p^m}$ and $0 \leq i_1 < i_2 < \dots < i_s \leq n-1$, then $p^s \mid |A|$.*

The proof of Lemma 6.1 is similar with the proof of Lemma 3.2, we omit the details. In order to study the zero set in $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$, we define an equivalence relation. For $u, v \in \mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$, we define the relation $u \sim v$ if there exists $r \in \mathbb{Z}_{p^n}^*$ such that $u = rv$. Then the equivalent classes in $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$ by \sim are

$$(p^i, 0), (c_i p^i, p^j) \text{ for all } i \in [0, m-1], j \in [0, n-1] \text{ and } c_i \in [0, p^{m-i} - 1] \text{ with } \gcd(c_i, p) = 1.$$

Note that the first zero in Lemma 6.1 is (a, p^{i_1}) , where $a \in p^{m-1}\mathbb{Z}_{p^m}$, and there are much more equivalent classes in $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$, it seems that our method does not work for the general case $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$. It would be interesting to consider the Fuglede's conjecture in $\mathbb{Z}_{p^m} \times \mathbb{Z}_{p^n}$.

References

- [1] C. Aten, B. Ayachi, E. Bau, D. FitzPatrick, A. Iosevich, H. Liu, A. Lott, I. MacKinnon, S. Maimon, S. Nan, J. Pakianathan, G. Petridis, C. Rojas Mena, A. Sheikh, T. Tribone, J. Weill, and C. Yu. Tiling sets and spectral sets over finite fields. *J. Funct. Anal.*, 273(8):2547–2577, 2017.
- [2] E. M. Coven, and A. Meyerowitz. Tiling the integers with translates of one finite set. *J. Algebra*, 212(1):161–174, 1999.
- [3] T. Fallon, G. Kiss and G. Somlai. Spectral sets and tiles in $\mathbb{Z}_p^2 \times \mathbb{Z}_q^2$. arXiv: 2105.10575.
- [4] A. Fan, S. Fan, L. Liao, and R. Shi. Fuglede's conjecture holds in \mathbb{Q}_p . *Math. Ann.*, 375(1-2):315–341, 2019.
- [5] A. Fan, S. Fan, and R. Shi. Compact open spectral sets in \mathbb{Q}_p . *J. Funct. Anal.*, 271(12):3628–3661, 2016.
- [6] B. Farkas, M. Matolcsi, and P. Móra. On Fuglede's conjecture and the existence of universal spectra. *J. Fourier Anal. Appl.*, 12(5):483–494, 2006.
- [7] B. Farkas and S. G. Révész. Tiles with no spectra in dimension 4. *Math. Scand.*, 98(1):44–52, 2006.
- [8] S. J. Ferguson and N. Sothanaphan. Fuglede's conjecture fails in 4 dimensions over odd prime fields. *Discrete Math.*, 343(1):111507, 7, 2020.
- [9] B. Fuglede. Commuting self-adjoint partial differential operators and a group theoretic problem. *J. Functional Analysis*, 16:101–121, 1974.

- [10] R. Greenfeld and N. Lev. Fuglede’s spectral set conjecture for convex polytopes. *Anal. PDE*, 10(6):1497–1538, 2017.
- [11] A. Iosevich, N. Katz, and T. Tao. The Fuglede spectral conjecture holds for convex planar domains. *Math. Res. Lett.*, 10(5-6):559–569, 2003.
- [12] A. Iosevich, A. Mayeli, and J. Pakianathan. The Fuglede conjecture holds in $\mathbb{Z}_p \times \mathbb{Z}_p$. *Anal. PDE*, 10(4):757–764, 2017.
- [13] G. Kiss, R. D. Malikiosis, G. Somlai, and M. Vizer. Fuglede’s conjecture holds for cyclic groups of order $pqrs$. arXiv: 2011.09578.
- [14] G. Kiss, R. D. Malikiosis, G. Somlai, and M. Vizer. On the discrete Fuglede and Pompeiu problems. *Anal. PDE*, 13(3):765–788, 2020.
- [15] G. Kiss and G. Somlai. Fuglede’s conjecture holds on $\mathbb{Z}_p^2 \times \mathbb{Z}_q$. *Proc. Amer. Math. Soc.*, 149(10):4181–4188, 2021.
- [16] M. N. Kolountzakis. Non-symmetric convex domains have no basis of exponentials. *Illinois J. Math.*, 44(3):542–550, 2000.
- [17] M. N. Kolountzakis and M. Matolcsi. Complex Hadamard matrices and the spectral set conjecture. *Collect. Math.*, (Vol. Extra):281–291, 2006.
- [18] M. N. Kolountzakis and M. Matolcsi. Tiles with no spectra. *Forum Math.*, 18(3):519–528, 2006.
- [19] I. Łaba. The spectral set conjecture and multiplicative properties of roots of polynomials. *J. London Math. Soc. (2)*, 65(3):661–671, 2002.
- [20] N. Lev and M. Matolcsi. The Fuglede conjecture for convex domains is true in all dimensions. *Acta Math.* to appear.
- [21] R. D. Malikiosis. On the structure of spectral and tiling subsets of cyclic groups. arXiv: 2005.05800.
- [22] R. D. Malikiosis and M. N. Kolountzakis. Fuglede’s conjecture on cyclic groups of order $p^n q$. *Discrete Anal.*, pages Paper No. 12, 16, 2017.
- [23] M. Matolcsi. Fuglede’s conjecture fails in dimension 4. *Proc. Amer. Math. Soc.*, 133(10):3021–3026, 2005.
- [24] R. Shi. Fuglede’s conjecture holds on cyclic groups \mathbb{Z}_{pqr} . *Discrete Anal.*, pages Paper No. 14, 14, 2019.

- [25] R. Shi. Equi-distributed property and spectral set conjecture on $\mathbb{Z}_{p^2} \times \mathbb{Z}_p$. *J. Lond. Math. Soc. (2)*, 102(3):1030–1046, 2020.
- [26] G. Somlai. Spectral sets in \mathbb{Z}_{p^2qr} tile. arXiv: 1907.04398.
- [27] S. Szabó and A. D. Sands. *Factoring groups into subsets*, volume 257 of *Lecture Notes in Pure and Applied Mathematics*. CRC Press, Boca Raton, FL, 2009.
- [28] T. Tao. Fuglede’s conjecture is false in 5 and higher dimensions. *Math. Res. Lett.*, 11(2-3):251–258, 2004.