

PRINCIPAL EIGENVALUE PROBLEM FOR INFINITY LAPLACIAN IN METRIC SPACES

QING LIU AND AYATO MITSUISHI

ABSTRACT. This paper is concerned with the Dirichlet eigenvalue problem associated to the ∞ -Laplacian in metric spaces. We establish a direct PDE approach to find the principal eigenvalue and eigenfunctions in a proper geodesic space without assuming any measure structure. We provide an appropriate notion of solutions to the ∞ -eigenvalue problem and show the existence of solutions by adapting Perron's method. Our method is different from the standard limit process via the variational eigenvalue formulation for p -Laplacian in the Euclidean space. Several concrete examples are given in the case of finite metric graphs. We also prove the uniqueness of eigenfunctions up to a multiplicative factor in this special case.

1. INTRODUCTION

1.1. Background and motivation. In this paper, we consider the principal eigenvalue and eigenfunctions associated to the ∞ -Laplacian with homogeneous Dirichlet boundary condition in metric spaces. One of our major contributions is a general framework that can be applied to study this eigenvalue problem in a large variety of metric spaces. Throughout this paper, the metric space (\mathbf{X}, d) is assumed to satisfy the following two conditions:

- (\mathbf{X}, d) is a geodesic space, namely, for any $x, y \in \mathbf{X}$, there exists a Lipschitz curve $\gamma : [a, b] \rightarrow \mathbf{X}$ such that $\gamma(a) = x, \gamma(b) = y$ and $d(x, y) = \ell(\gamma)$, where $\ell(\gamma)$ stands for the length of γ . (Throughout this paper, by an abuse of notation, we also use γ to denote the image set of the curve.)
- (\mathbf{X}, d) is proper, that is, for any $x \in \mathbf{X}$ and $r > 0$, the closed metric ball $\overline{B_r(x)}$ is compact. Here and in the sequel, we denote by $B_r(x)$ the open metric ball centered at x with radius $r > 0$.

Before stating our main results, let us first go over the background on the topic and describe our motivation of this work.

The study on the eigenvalue problem for the ∞ -Laplacian is initiated by the work [30] (and also [21]), where the limits of eigenvalue and eigenfunctions for the p -Laplacian as $p \rightarrow \infty$ are investigated in the Euclidean space. More precisely, for any given $1 < p < \infty$ and a bounded domain $\Omega \subset \mathbb{R}^n$, we can obtain the first p -eigenvalue, written as Λ_p^p (the p -th power of Λ_p), via the Rayleigh quotient

$$\Lambda_p^p = \min \left\{ \frac{\int_{\Omega} |\nabla u|^p dx}{\int_{\Omega} |u|^p dx} : u \in W_0^{1,p}(\Omega) \setminus \{0\} \right\}. \quad (1.1)$$

Date: September 22, 2021.

2010 Mathematics Subject Classification. 35R15, 35P30, 35J60, 31C05.

Key words and phrases. eigenvalue problems, metric spaces, ∞ -Laplacian, eikonal equation, viscosity solutions, metric graphs.

The minimizers in (1.1) are called the p -eigenfunctions or p -ground states.

It is shown [30, 21] that Λ_p converges as $p \rightarrow \infty$ to

$$\Lambda_\infty = \frac{1}{R_\infty}, \quad (1.2)$$

where $R_\infty > 0$ is the radius of the maximum ball inscribed in Ω :

$$R_\infty = \max_{x \in \overline{\Omega}} \min_{y \in \partial\Omega} |x - y|.$$

Thus, the value Λ_∞ is considered as the principal eigenvalue of the ∞ -Laplacian and is called ∞ -eigenvalue.

Moreover, it is also proved in [30, 21] that any uniform limit of appropriately normalized p -eigenfunctions is a viscosity solutions of the following obstacle problem

$$\min\{|\nabla u| - \Lambda_\infty u, -\Delta_\infty u\} = 0 \quad \text{in} \quad (1.3)$$

with

$$u = 0 \quad \text{on} \quad \partial\Omega, \quad (1.4)$$

where $\Delta_\infty u$ denotes the ∞ -Laplacian of a function $u \in C^2(\Omega)$, i.e.,

$$\Delta_\infty u = \langle \nabla^2 u \nabla u, \nabla u \rangle.$$

The positive solutions obtained from the limit process are called ∞ -ground states in the literature. The higher eigenvalues and eigenfunctions arising in this limit are discussed in [29].

The above results on the principal ∞ -eigenvalue problem are later developed for more general nonlinear elliptic operators [5, 14, 44, 13, 32, 10, 19] and for more general boundary conditions [24, 42, 20]. However, on the other hand, less is known about this problem in general geometric settings. We refer to [6] for generalization under the Finsler metrics. In metric measure spaces, the convergence of Λ_p to Λ_∞ is addressed in [2, 26] and results related to the limit for higher p -eigenvalues are recently provided by the second author [40]. In these results the equipped measure structure plays a fundamental role to invoke the variational setting.

It is worth pointing out that the full convergence of normalized p -eigenfunctions as $p \rightarrow \infty$ is still unclear in general even in the Euclidean space. This is related to the long standing open question on the uniqueness of ∞ -ground states up to a multiplicative factor, or in other words, the simplicity of the ∞ -eigenvalue Λ_∞ . An affirmative result is given by Yu [48] for $\Omega \subset \mathbb{R}^n$ in certain particular shapes when the distance to $\partial\Omega$ is a viscosity solution of (1.3) and (1.4).

Rather than investigating the convergence of p -eigenfunctions, one may try to directly prove the uniqueness of viscosity solutions of (1.3) and (1.4) up to a constant multiple. However, counterexamples have been constructed in [27] to show that it fails in general. At least three linear independent solutions are built in a dumbbell-shaped domain in \mathbb{R}^2 that is symmetric with respect to the axes. However, only one of them is considered to be variational, since the others are not symmetric about the axes and certainly cannot be the uniform limits of the symmetric p -eigenfunctions.

In addition to [27], much progress has been made to tackle this uniqueness problem. For instance, the papers [17, 18] develop the observations in [48] and give new results on the geometry of the domains for which the uniqueness holds. In [19], a variational formulation more general than (1.1) is introduced to allow different normalizations and a maximal

solution to (1.3) and (1.4) is obtained through the limiting process. More recently, in relation to the uniqueness issue in convex domains in \mathbb{R}^2 , Lindgren and Lindqvist [35] discuss the properties of certain specific curves, called attracting streamlines, for the normalized ∞ -eigenfunctions; see also [33, 34] for related results.

In this work, we study the ∞ -eigenvalue problem from a more geometric perspective. Our purpose is twofold.

- (a) We intend to study the principal ∞ -eigenvalue problem in general metric spaces with minimal structure assumptions. In particular, it is of our interest to find an ∞ -eigenvalue consistent with (1.2) and to generalize the known results on Euclidean ∞ -eigenfunctions in a proper geodesic space without any measure structure.
- (b) As an application of our generalized framework, we look into the uniqueness of ∞ -eigenfunctions in finite metric graphs, which is a special class of geodesic spaces with a particularly simple structure. We revisit the uniqueness problem in this special case and provide new insights into this challenging open question.

Our study, especially for the purpose (a), is motivated by the following observations. First, the ∞ -eigenvalue in (1.2) is a completely geometric quantity and requires nothing more than the space metric. We naturally expect that in a general metric space (\mathbf{X}, d) , the expression of the ∞ -eigenvalue turns into

$$\Lambda_\infty = \frac{1}{\max_{x \in \overline{\Omega}} d(x, \partial\Omega)}. \quad (1.5)$$

The denominator represents the radius of the maximum inscribed metric ball, which we still denote by R_∞ , i.e.,

$$R_\infty = \max_{x \in \overline{\Omega}} d(x, \partial\Omega). \quad (1.6)$$

Here and in the sequel, $d(x, E)$ denotes the distance from a point $x \in \mathbf{X}$ to a compact set $E \subset \mathbf{X}$, namely, $d(x, E) = \min_{y \in E} d(x, y)$.

Second, both the eikonal operator and the ∞ -Laplacian appearing in the nonlinear obstacle problem (1.3) can be understood under merely the length structure. In recent years, several notions of solutions to the eikonal equation in geodesic or length spaces are proposed [25, 23, 38]. Concerning the ∞ -Laplace equation, we refer to [43] for a tug-of-war game interpretation in length spaces. Also, it is well known that ∞ -harmonic functions in the Euclidean space can be characterized via comparison with cones; see for example [15, 4, 3]. This geometric characterization is extended to general metric spaces in [31]. None of these results essentially require measures on the spaces.

Besides, the eigenvalue problem can be set up without relying on the limit process via p -Laplacian. Recall that in the Euclidean space the fully nonlinear principal eigenvalue problem

$$F(\nabla u, \nabla^2 u) = \lambda u \quad \text{in } \Omega \subset \mathbb{R}^n \quad (1.7)$$

with the Dirichlet condition (1.4) is treated in [8, 9] based on the maximum principle. More precisely, the eigenvalue is obtained as the maximum value of $\lambda \in \mathbb{R}$ that admits the existence of positive viscosity supersolutions of (1.7).

Based on the observations above, we establish a new approach to the eigenvalue problem that is applicable to general geodesic spaces. Our strategy is as follows. Instead of passing to the limit for the p -eigenvalue problem as $p \rightarrow \infty$ as in [30], we follow [8, 9] to investigate,

in a more straightforward manner, the maximum value λ that guarantees existence of positive supersolutions of

$$\min\{|\nabla u| - \lambda u, -\Delta_\infty u\} = 0 \quad \text{in } \Omega. \quad (1.8)$$

We adopt this method to avoid the use of measures that are required to formulate the variational p -eigenvalue problem. It turns out that such a critical value $\lambda > 0$ does coincide with Λ_∞ as in (1.5).

Once the eigenvalue Λ_∞ is justified, we can discuss positive solutions of (1.3) that also satisfy the boundary condition (1.4). Such solutions are regarded as ∞ -eigenfunctions in our general setting. As in the Euclidean case, we do not know whether they are unique up to a constant multiple in general. But the existence of ∞ -eigenfunctions can be obtained by adapting Perron's method. We construct solutions of (1.3) by taking the infimum of all supersolutions satisfying appropriate conditions that essentially play the role of normalization. In order to make our PDE-based arguments above work, it is crucial to find an appropriate notion of solutions of (1.8), which we will clarify in a moment.

As mentioned above, our general framework facilitates our understanding of the ∞ -eigenvalue and ∞ -eigenfunctions in various specific geodesic spaces. We are particularly interested in the case when \mathbf{X} is a finite metric graph. In fact, thanks to the finiteness and one-dimensional structure, we are able to have a clear view about the uniqueness of solutions of (1.3) and (1.4). Not only can we construct concrete examples, but we also manage to prove the desired uniqueness of ∞ -eigenfunctions in this special case under certain natural constraints. More detailed explanations on this will be given later.

1.2. Main results in general geodesics spaces. In order to present our main results, let us first introduce the notion of solutions to (1.3) in geodesic spaces. Assume that (\mathbf{X}, d) is a proper geodesic space. Let $\Omega \subsetneq \mathbf{X}$ be a bounded domain. We seek solutions of (1.8) in the class of locally Lipschitz functions in Ω . Let us briefly clarify our definitions of solutions below; see Section 2 for more precise descriptions.

As usual, our definition can be divided into a supersolution part and a subsolution part. When defining a supersolution u , we require it to fulfill the supersolution properties for both the eikonal and the infinity Laplace equation. The former can be simply defined by

$$|\nabla^- u| \geq \lambda u \quad (1.9)$$

in Ω , where $|\nabla^- u|$ denotes the subslope of u , given by

$$|\nabla^- u|(x) = \limsup_{y \rightarrow x} \frac{\max\{u(x) - u(y), 0\}}{d(x, y)}.$$

See also the definitions of slope $|\nabla u|$ and superslope $|\nabla^+ u|$ respectively in (2.4) and (2.5).

One can analogously use the subslope to define subsolutions and solutions of the eikonal equation. This type of solutions of eikonal equations is called Monge solutions. Such a notion is studied in the Euclidean space [41, 11] and is introduced in [38] for general metric spaces. See [38] also for the equivalence with other notions of metric viscosity solutions proposed in [1, 25, 22, 23]. For the reader's convenience, we include several basic results on the eikonal equation in metric spaces in Appendix A.

Regarding the ∞ -Laplacian supersolution property (or ∞ -superharmonicity), we adopt the characterization of comparison with cones from below. This requires that an ∞ -superharmonic function u satisfy

$$\min_{\overline{\mathcal{O}}} (u - \phi) \geq \min_{\partial \mathcal{O}} (u - \phi)$$

for any open subset $\mathcal{O} \subset \subset \Omega$ (i.e., $\overline{\mathcal{O}} \subset \Omega$) and any cone function

$$\phi = a + \kappa d(\hat{x}, \cdot) \quad \text{in } \Omega \quad (1.10)$$

with $a \in \mathbb{R}$, $\kappa \leq 0$ and $\hat{x} \in \Omega \setminus \mathcal{O}$. Consult [31] for more details on the properties of comparison with cones in connection with the absolute minimizing Lipschitz extensions in metric spaces.

We consequently call u a supersolution to (1.8) if it satisfies both (1.9) and comparison with cones from below in Ω . This notion looks very different from the usual viscosity supersolutions, since it is not defined pointwise by means of test functions. In contrast, we need to define subsolutions of (1.8) pointwise and choose an appropriate class of test functions. While in the Euclidean case one can test a subsolution at $x_0 \in \Omega$ by a C^2 function v satisfying $-\Delta_\infty v(x_0) > 0$ so as to obtain

$$|\nabla v(x_0)| \leq \lambda u(x_0), \quad (1.11)$$

finding the corresponding test class in general metric spaces is however not straightforward. We overcome the difficulty by adopting the class of ∞ -superharmonic functions introduced above to test the candidate function in a strict manner.

We say that u is a subsolution of (1.8) if whenever there exist an ∞ -superharmonic function v and $x_0 \in \Omega$ such that $u - v$ attains a strict local maximum at x_0 , we have

$$\liminf_{r \rightarrow 0^+} \inf_{B_r(x_0)} |\nabla^- v| \leq \lambda u(x_0). \quad (1.12)$$

The left hand side above looks slightly complicated. Actually (1.12) reduces to (1.11) in \mathbb{R}^n for a test function $v \in C^2(\Omega)$. In our current setting, the lower semicontinuous envelope of the subslope needs to be utilized due to the lack of smoothness of v .

Our definitions of supersolutions and subsolutions of (1.8) prove to be an appropriate generalization of those in the Euclidean case. Indeed, we can show the equivalence between both types of definitions in the case when $\mathbf{X} = \mathbb{R}^n$ based on the results in [15, 4, 31]; see Section 4 for details.

Moreover, our new notions enable us to solve the ∞ -eigenvalue problem in general geodesic spaces. Using our notion of supersolutions of (1.8), we define the ∞ -eigenvalue in Ω by

$$\Lambda = \sup \{ \lambda \in \mathbb{R} : \text{there exists a locally Lipschitz positive supersolution of (1.8)} \}. \quad (1.13)$$

It turns out that this value coincides with (1.5), which is consistent with the Euclidean result.

Theorem 1.1 (∞ -eigenvalue). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $\Lambda_\infty > 0$ be given by (1.5). Let Λ be defined by (1.13). Then $\Lambda = \Lambda_\infty$ holds.*

Our proof of Theorem 1.1, which is elaborated in Section 3.1, consists of two steps. We first verify that

$$u_{dist}(x) = \Lambda_\infty d(x, \partial\Omega), \quad x \in \overline{\Omega}, \quad (1.14)$$

is a supersolution of (1.3). This implies immediately that $\Lambda \geq \Lambda_\infty$. The reverse inequality is shown by proving the non-existence of supersolutions of (1.8) when $\lambda > \Lambda_\infty$. It actually follows from a fundamental property of supersolutions at any $x_0 \in \Omega$ satisfying

$$d(x_0, \partial\Omega) = R_\infty. \quad (1.15)$$

Here and in the sequel we call such a point an incenter of Ω . In fact, for an incenter point x_0 , by comparing any positive supersolution u with the cone function ϕ in (1.10) with $a = u(x_0)$ and any $\kappa < -|\nabla^- u|(x_0)$, one can prove that

$$|\nabla^- u|(x_0) \leq u(x_0)/R_\infty = \Lambda_\infty u(x_0),$$

which yields $\Lambda \leq \Lambda_\infty$ by (1.13).

Our second main result is on the existence of positive solutions of (1.3) satisfying the boundary condition (1.4). As usual, by solutions of (1.3), we mean locally Lipschitz functions that are both supersolutions and subsolutions. The positive solutions of (1.3) satisfying (1.4) are called ∞ -eigenfunctions in our current setting. In order to obtain the existence, we adapt Perron's method by taking the pointwise infimum of all supersolutions under the constraint

$$u = \max_{\overline{\Omega}} u = 1 \quad \text{on } \mathcal{M}(\Omega), \quad (1.16)$$

where $\mathcal{M}(\Omega)$ denotes the set of all incenters in Ω , i.e.,

$$\mathcal{M}(\Omega) := \left\{ x \in \Omega : d(x, \partial\Omega) = \max_{\overline{\Omega}} d(\cdot, \partial\Omega) = R_\infty \right\}. \quad (1.17)$$

In other words, we set, for any $x \in \overline{\Omega}$,

$$u_\infty(x) = \inf \left\{ u(x) : u \in C(\overline{\Omega}) \text{ is a positive supersolution of (1.3) satisfying (1.16)} \right\}. \quad (1.18)$$

We then obtain the following result.

Theorem 1.2 (Existence of ∞ -eigenfunctions). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u_\infty : \overline{\Omega} \rightarrow \mathbb{R}$ be defined by (1.18). Then u_∞ is continuous in $\overline{\Omega}$ and is a positive solution of (1.3) satisfying (1.4) and (1.16).*

The proof of Theorem 1.2 streamlines Perron's method but some arguments need to be adapted. One noteworthy issue is about the regularity of solutions. Recall that Perron's method for classical viscosity solutions is usually carried out in the class of semicontinuous sub- or supersolutions. One can later apply the comparison principle to obtain the continuity of solutions. We refer the reader to [16] for a detailed introduction on this method.

Our method here is slightly different. As shown in (1.18), we choose to construct solutions of (1.3) directly in the class of locally Lipschitz functions. This is possible because the property of comparison with cones implies local Lipschitz regularity. This result is presented in Lemma 2.4. In fact, in Lemma 2.5 we prove more for an ∞ -superharmonic function u in Ω : it satisfies

$$|\nabla u| = |\nabla^- u| \quad (1.19)$$

and the slope $|\nabla u|$ is upper semicontinuous in Ω ; see also [38] for remarks on (1.19) in relation to the semiconcavity regularity. We use the regularity result to obtain the local Lipschitz continuity of u_∞ after showing its ∞ -superharmonicity. We also include a result (Proposition 2.6) on Harnack's inequality for ∞ -superharmonic functions to show that $u_\infty > 0$ in Ω . Our new version of Harnack's inequality, which applies to general metric spaces, generalizes the results in [37, 7] in the Euclidean case.

In view of Theorem 1.2 and (1.18), we see that u_∞ is the minimal positive solution of (1.3) satisfying the boundary condition (1.4) and the constraint (1.16). The constraint (1.16) essentially normalizes any eigenfunction u so that

$$\|u\|_{L^\infty(\Omega)} = 1. \quad (1.20)$$

One may wonder whether u_∞ is the only solution satisfying (1.4) and (1.20). In general it fails to hold. We can generalize our method to find a solution under a partial constraint

$$u = \max_{\overline{\Omega}} u = 1 \quad \text{in } Y \quad (1.21)$$

instead of (1.16), where Y is a compact subset of $\mathcal{M}(\Omega)$. Letting

$$u_\infty^Y(x) := \inf \{u(x) : u \text{ is a positive supersolution of (1.3) satisfying (1.21)}\}, \quad (1.22)$$

we can follow the same proof to show that u_∞^Y is also a solution of (1.3)–(1.4). In general, it happens that $u_\infty^Y \neq u_\infty$ when $\mathcal{M}(\Omega)$ is not a singleton and $Y \subsetneq \mathcal{M}(\Omega)$. In Section 5.4, we present a concrete example in a finite metric graph, which can be regarded as a refined counterpart of the Euclidean example in [27].

It is thus natural to ask whether the positive solutions of (1.3) satisfying both (1.4) and (1.16) are unique. This question is closely related to the open problem on the simplicity of ∞ -eigenvalue in the Euclidean space. We establish a comparison principle, Theorem 3.9, which generalizes [30, Theorem 2.3]. Unfortunately this comparison result does not imply the desired uniqueness, since it requires uniform positivity of subsolutions and supersolutions in $\overline{\Omega}$. However, thanks to this comparison result, the uniqueness problem can be reduced from the whole domain to a neighborhood of $\partial\Omega$, which helps us deal with special geodesic spaces such as metric graphs.

1.3. Application to metric graphs. Let us now discuss an application of our general approach to the case when \mathbf{X} is a finite connected metric graph. A finite metric graph is the result of gluing a finite collection of vertices and edges of finite length equipped with the intrinsic metric; see Section 5.1 as well as [12] for preliminaries on metric graphs.

The one-dimensional structure of metric graphs enables us to better understand the behavior of ∞ -eigenfunctions in a domain $\Omega \subsetneq \mathbf{X}$. A key observation is that any positive solution u of (1.3) is piecewise linear, attains no local minima in Ω , and achieves at most one local maximum on each edge. Moreover, one can show that the subslope $|\nabla^- u|$ and superslope $|\nabla^+ u|$ agree at any point $x \in \Omega$ unless x is a local maximizer of u . This suggests the existence of a curve connecting any x to a local maximizer of u , along which $|\nabla^- u|$ is nondecreasing; we name such a curve an upstream curve of u from x . On the other hand, one can also define a downstream curve from x toward the boundary $\partial\Omega$, along which $|\nabla^- u|$ is also nondecreasing. This property is due to the general regularity (1.19) of ∞ -superharmonic functions. All of these results are presented in detail in Section 5.2 and Section 5.5.

Taking into consideration the properties described above, we can easily build many concrete examples of ∞ -eigenfunctions in metric graphs. Besides the aforementioned example related to [27], in Section 5.3 we also examine the case when \mathbf{X} is a tripod, that is, the graph formed by three edges sharing one common endpoint. A variant of the example containing a loop is also studied.

More importantly, we can prove the uniqueness of ∞ -eigenfunctions satisfying the condition (1.16).

Theorem 1.3 (Unique existence of eigenfunctions in finite metric graphs). *Suppose that $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite connected metric graph equipped with intrinsic metric d . Let $\Omega \subsetneq \mathbf{X}$ be a domain with $\partial\Omega \subset \mathcal{V}$. Let Λ_∞ be as in (1.5). Then the positive solutions of (1.3) satisfying (1.4) and (1.16) are unique. In particular, u_∞ given by (1.18) is the unique positive solution of (1.3) satisfying (1.4) and (1.16).*

Our proof looks very different from the standard uniqueness arguments for elliptic problems. We provide an iterative method to construct solutions satisfying (1.4) and (1.16) and show that the value of any solution in a neighborhood of $\partial\Omega$ is completely determined by this method. Now that the uniqueness near the boundary has been proved, we can use the comparison principle for uniformly positive sub- and supersolutions to conclude the uniqueness of solutions in the whole domain.

There is a PDE interpretation on how we determine the value of solutions near $\partial\Omega$. Essentially, we iteratively solve a free boundary problem for the eikonal equation that admits the steepest constant subslope. Under the Dirichlet boundary condition (1.4), we first look for a solution U of

$$|\nabla^- U| = \lambda \tag{1.23}$$

with a maximum possible $\lambda > 0$ in an open subset $\Omega' \subset \Omega$ with $\partial\Omega' \subset \partial\Omega$. Denote by $L(U, \Omega)$ the largest possible value λ . Let Z_0 denote the largest region Ω' for the existence of solution U of the boundary value problem with $\lambda = L(U, \Omega)$. We can show that U restricted to Z_0 is piecewise linear and has subslope $L(U, \mathcal{O}_0)$ everywhere. It turns out that $L(U, \Omega) = \Lambda_\infty$. We then take $\mathcal{S}_0 = \overline{Z_0}$, which consists of all closed geodesics joining points in $\mathcal{M}(\Omega)$ and on $\partial\Omega$. Also, for any $y \in \mathcal{S}_0$, we see that

$$U(x) = \Lambda_\infty d(x, \partial\Omega)$$

for all $x \in \mathcal{S}_0$.

Our goal is achieved if \mathcal{S}_0 contains a neighborhood of $\partial\Omega$. Otherwise, we take $\mathcal{O}_1 = \Omega \setminus \mathcal{S}_0$ and study the existence of the same free boundary value problem in \mathcal{O}_1 again, for which we denote by $L(U, \mathcal{O}_1)$ the largest possible λ this time. Note that the values of U on \mathcal{S}_0 now becomes a part of the prescribed boundary condition. Once we solve the problem on a subregion Z_1 of \mathcal{O}_1 , we obtain U on the set $\mathcal{S}_1 = \mathcal{S}_0 \cup \overline{Z_1}$. If \mathcal{S}_1 still fails to include a neighborhood of $\partial\Omega$, we repeat this operation to get the next subslope $L(U, \mathcal{O}_2)$ and U on \mathcal{S}_2 . Such a process is continued to extend the value of U in succession to $\mathcal{S}_2, \mathcal{S}_3, \dots$, until \mathcal{S}_N contains a neighborhood of $\partial\Omega$ for some $N \in \mathbb{N}$. This process is stopped in finite steps, since \mathbf{X} contains only finitely many edges and at least one edge is newly included into \mathcal{S}_k for each $k = 1, 2, \dots$ before the process ends.

In practice, we iteratively construct U in a direct manner based on the property that any solution is piecewise linear. For any $k = 1, 2, \dots, N - 1$, we search all possible curves in $\mathcal{O}_k = \Omega \setminus \mathcal{S}_{k-1}$ and all piecewise linear functions along the curves that admit a constant subslope and agree with the existing values of U on $\partial\mathcal{O}_k$. Maximizing the subslope over all these candidates, we obtain the maximum slope $L(U, \mathcal{O}_k)$ and the optimized function on Z_k , by which we make the extension of U to \mathcal{S}_k . The uniqueness result as in Theorem 1.3 is also proved via the process. One can show that any solution u admits the subslope $L(U, \mathcal{O}_k)$ in Z_k for each $k = 1, 2, \dots, N - 1$. If it is not the case, then there exist points in Z_k where u admits a subslope larger than $L(U, \mathcal{O}_k)$. Using the upstream and downstream curves introduced above, we end up with a subregion of \mathcal{O}_k in which u has a constant subslope greater than $L(U, \mathcal{O}_k)$. This contradicts our constructive method to find $L(U, \mathcal{O}_k)$. See Section 5.5 and Section 5.6 for more details.

Our uniqueness arguments seem to be restricted to the graph case or other one dimensional spaces. It would be interesting if this method can be developed for more general metric spaces including \mathbb{R}^n with $n \geq 2$.

The rest of the paper is organized in the following way. In Section 2, we provide precise definitions of supersolutions and subsolutions of (1.3) in geodesic spaces. Several important properties of ∞ -superharmonic functions will also be studied including the Lipschitz regularity and Harnack's inequality. In Section 3, we prove our main results, Theorem 1.1, Theorem 1.2 and a comparison principle for the ∞ -eigenvalue problem. Section 5 is devoted to the special case when \mathbf{X} is a metric graph. For the reader's convenience, we include Appendix A to recall preliminaries on the eikonal equation in metric spaces.

Acknowledgements. The work of the first author is supported by JSPS Grant-in-Aid for Scientific Research (No. 19K03574). The work of the second author is supported by JSPS Grant-in-Aid for Scientific Research (No. 20K03598).

2. DEFINITION AND PROPERTIES OF SOLUTIONS

In this section we give a generalized notion of solutions to (1.8) in a proper geodesic space for $\lambda \in \mathbb{R}$.

It is well known that the comparison with cones (cf. [4]) can be employed to characterize the ∞ -harmonic functions; we refer to [31] for generalization in general metric spaces. We recall the definition of super- and subsolutions of

$$-\Delta_\infty u = 0 \quad \text{in } \Omega \tag{2.1}$$

based on this property.

Definition 2.1 (∞ -superharmonic functions). Let (\mathbf{X}, d) be a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ be a bounded domain. A function $u : \Omega \rightarrow \mathbb{R}$ that is bounded from below is said to be ∞ -superharmonic in Ω (or a supersolution of (2.1)) if it satisfies the following property of comparison with cones from below in Ω : for any $\hat{x} \in \Omega$, any $a \in \mathbb{R}$, $\kappa \leq 0$ and any bounded open set $\mathcal{O} \subset\subset \Omega$ with $\hat{x} \in \Omega \setminus \mathcal{O}$, the condition

$$u \geq \phi \quad \text{on } \partial\mathcal{O} \tag{2.2}$$

for ϕ given by (1.10) implies that

$$u \geq \phi \quad \text{in } \overline{\mathcal{O}}. \tag{2.3}$$

Remark 2.2 (Definition of ∞ -(sub)harmonic functions). We can also define ∞ -subharmonic in a symmetric way. More precisely, we say that any $u : \Omega \rightarrow \mathbb{R}$ bounded from above is ∞ -subharmonic if $-u$ is ∞ -superharmonic. In addition, u is said to be ∞ -harmonic if it is both ∞ -superharmonic and ∞ -subharmonic.

We next provide an immediate consequence of Definition 2.1, which will be used later. Let us recall that for any $x \in \mathbf{X}$ and a locally Lipschitz function u , the local slope of u at x is given by

$$|\nabla u|(x) = \limsup_{y \rightarrow x} \frac{|u(y) - u(x)|}{d(x, y)} \tag{2.4}$$

and the sub- and superslopes of u at x are defined to be

$$|\nabla^\pm u|(x) = \limsup_{y \rightarrow x} \frac{[u(y) - u(x)]_\pm}{d(x, y)}, \tag{2.5}$$

where $[a]_\pm = \max\{\pm a, 0\}$ for any $a \in \mathbb{R}$.

Lemma 2.3 (Comparison with special cone functions). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u : \Omega \rightarrow \mathbb{R}$ be ∞ -superharmonic in Ω , that is, u is bounded from below and obeys the comparison with cones from below in Ω . Let $\Omega' \subset\subset \Omega$ be an open bounded set and $x_0 \in \Omega'$. For $\kappa \leq 0$, if $u \geq \phi$ on $\partial\Omega'$, where*

$$\phi(x) := u(x_0) + \kappa d(x_0, x), \quad x \in \Omega. \quad (2.6)$$

then $u \geq \phi$ in Ω' . In addition, if u is locally Lipschitz in Ω , then for any $r > 0$ satisfying $B_r(x_0) \subset\subset \Omega$, there exists $x_r \in \partial B_r(x_0)$ such that

$$u(x_0) - u(x_r) \geq |\nabla^- u|(x_0)r. \quad (2.7)$$

Proof. This result follows directly from Definition 2.1 with $\mathcal{O} = \Omega' \setminus \{x_0\}$ and the cone function taken to be ϕ as in (2.6). We then can use the comparison with cones to get $u \geq \phi$ in Ω' .

To show the second statement, we assume by contradiction that it fails to hold, which yields existence of $\sigma > 0$ small such that

$$u(x_0) - u(x) \leq (|\nabla^- u|(x_0) - \sigma)r$$

for all $x \in \partial B_r(x_0)$. We may apply the previous result to show

$$u(x_0) - u(x) \leq (|\nabla^- u|(x_0) - \sigma)d(x_0, x)$$

for all $x \in B_r(x_0)$. This is clearly a contradiction by the definition of $|\nabla^- u|$. \square

We next show that ∞ -superharmonic functions are actually locally Lipschitz continuous. Related regularity results in the Euclidean case based on Harnack's inequality can be found in [37, 7]. Our proof for general geodesic spaces is more direct, using the comparison with cones only.

Lemma 2.4 (Local Lipschitz continuity). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u : \Omega \rightarrow \mathbb{R}$ be ∞ -superharmonic in Ω , that is, u is bounded from below and obeys the comparison with cones from below in Ω . Then, u is locally Lipschitz in Ω . More precisely, for any $x \in \Omega$, $y \in B_r(x)$ with $0 < 2r < d(x, \partial\Omega)$,*

$$|u(x) - u(y)| \leq \frac{\max\{u(x), u(y)\} - \inf_{\Omega} u}{r} d(x, y). \quad (2.8)$$

A symmetric result for an ∞ -subharmonic function u in Ω ; in this case, u is still locally Lipschitz in Ω and satisfies (2.8) with u replaced by $-u$.

Proof. Suppose that u is a supersolution that is bounded from below in Ω . Let us take $y \in \Omega$ and any $s > 0$ satisfying $d(y, \partial\Omega) > s$. We consider a cone function defined by

$$\phi(z) := u(x) - \frac{u(x) - \inf_{\Omega} u}{s} d(x, z).$$

Then, $u \geq \inf_{\Omega} u \equiv \phi$ holds on $\partial B_s(x)$. Therefore, by Lemma 2.3 with $x_0 = x$ and $\Omega' = B_s(x)$, we obtain

$$u(x) - u(y) \geq -\frac{u(x) - \inf_{\Omega} u}{s} d(x, y) \quad (2.9)$$

whenever $y \in B_s(x)$ and $s < d(x, \partial\Omega)$.

Let us take $r > 0$ small with $2r < d(x, \partial\Omega)$ and $y \in B_r(x)$. Then (2.9) immediately implies that

$$u(x) - u(y) \geq -\frac{u(x) - \inf_{\Omega} u}{r} d(y, x). \quad (2.10)$$

On the other hand, since

$$d(y, \partial\Omega) \geq d(x, \partial\Omega) - d(x, y) > r,$$

it follows from (2.9) again that

$$u(y) - u(x) \geq -\frac{u(y) - \inf_{\Omega} u}{r} d(x, y). \quad (2.11)$$

We conclude the proof by combining (2.10) and (2.11). \square

A further regularity property of ∞ -harmonic functions is as follows.

Lemma 2.5 (Slope regularity of ∞ -harmonic functions). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u : \Omega \rightarrow \mathbb{R}$ be ∞ -superharmonic in Ω . Then, in Ω the slope and subslope of u coincide, i.e., (1.19) holds, and they are both upper semicontinuous. A symmetric result for an ∞ -subharmonic function u in Ω , that is, $|\nabla u|$ and $|\nabla^+ u|$ coincide and are upper semicontinuous in Ω .*

Proof. We only consider the case when u is ∞ -superharmonic, since the argument is symmetric for ∞ -subharmonic functions.

By Lemma 2.4, u is locally Lipschitz in Ω . Fix $x_0 \in \Omega$ arbitrarily. It is clear that

$$|\nabla u|(x_0) = \max\{|\nabla^+ u|(x_0), |\nabla^- u|(x_0)\}. \quad (2.12)$$

In what follows let us show that

$$|\nabla^- u|(x_0) \geq |\nabla^+ u|(x_0). \quad (2.13)$$

For any $\varepsilon > 0$ small, we can find $r > 0$ such that

$$u(x) \geq u(x_0) - (|\nabla^- u|(x_0) + \varepsilon)d(x, x_0)$$

for any $x \in B_r(x_0)$. Let us take

$$m_{\varepsilon, r} = u(x_0) - (|\nabla^- u|(x_0) + \varepsilon)r.$$

For any $z \in B_r(x_0)$ close to x_0 , let

$$\phi_z(x) := u(z) - \frac{u(z) - m_{\varepsilon, r}}{r_z} d(x, z)$$

for $x \in B_r(x_0)$, where we set

$$r_z = r - d(x_0, z).$$

It is not difficult to see that $u \geq \phi_z$ on $\partial B_{r_z}(z)$. We apply Lemma 2.3 to get

$$u \geq \phi_z \quad \text{in } B_{r_z}(z). \quad (2.14)$$

In particular, for any z sufficiently close to x_0 we have

$$u(x_0) \geq \phi_z(x_0),$$

which yields

$$u(z) - u(x_0) \leq \frac{u(z) - m_{\varepsilon, r}}{r_z} d(x_0, z)$$

and therefore

$$\frac{[u(z) - u(x_0)]_+}{d(x_0, z)} \leq \left| \frac{u(z) - m_{\varepsilon, r}}{r_z} \right| \leq \frac{|u(z) - u(x_0)|}{r - d(x_0, z)} + (|\nabla^- u|(x_0) + \varepsilon) \frac{r}{r - d(x_0, z)}.$$

Sending $d(z, x_0) \rightarrow 0$ and then $\varepsilon \rightarrow 0$, we are led to (2.13). We immediately obtain $|\nabla^- u| = |\nabla u|$ in Ω due to (2.12) and the arbitrariness of $x_0 \in \Omega$.

We now show the upper semicontinuity of $|\nabla^- u|$. It follows from (2.14) again that

$$u(z) - u(x) \leq u(z) - \phi_z(x) = \frac{u(z) - m_{\varepsilon,r}}{r_z} d(x, z)$$

for all x near z , which implies that

$$|\nabla^- u|(z) \leq \frac{u(z) - m_{\varepsilon,r}}{r - d(x_0, z)} = \frac{u(z) - u(x_0) + (|\nabla^- u|(x_0) + \varepsilon)r}{r - d(x_0, z)}.$$

Letting $z \rightarrow x_0$, we have

$$\limsup_{z \rightarrow x_0} |\nabla^- u|(z) \leq |\nabla^- u|(x_0) + \varepsilon.$$

We complete our proof of the upper semicontinuity of $|\nabla^- u|$ in Ω by noticing that $\varepsilon > 0$ and $x_0 \in \Omega$ are arbitrary. \square

In addition, we generalize, in the context of general geodesic spaces, a result in [37, 7, 36] on Harnack's inequality for the ∞ -Laplace equation.

Proposition 2.6 (Harnack's inequality). *Suppose that (\mathbf{X}, d) is a proper geodesic space and Ω is an open subset of \mathbf{X} . Assume that $u : \Omega \rightarrow \mathbb{R}$ satisfies the comparison with cones from below. Assume that $u \geq 0$ in $B_R(x_0)$ with $B_R(x_0) \subset \Omega$. Then,*

$$u(y) \leq 3u(x) \tag{2.15}$$

for any $x, y \in B_r(x_0)$ and $r > 0$ with $4r < R$. In addition, if Ω is connected, u is lower semicontinuous, nonnegative on Ω , and $\sup_{\Omega} u > 0$, then $u > 0$ in Ω .

Proof. Fix $y \in B_r(x_0)$ arbitrarily. Let us define

$$\kappa := \min_{z \in \partial B_{3r}(y)} u(z) - u(y)$$

and consider a cone function

$$\phi(\cdot) := u(y) + \frac{\min\{\kappa, 0\}}{3r} d(y, \cdot).$$

It is clear that $u \geq \phi$ holds on $\partial B_{3r}(y)$. By Lemma 2.3, we then have $u \geq \phi$ in $B_{3r}(y)$.

If $\kappa \geq 0$, then for any $x \in B_r(x_0)$, we have $\phi(x) = u(y)$ and thus $u(x) \geq u(y)$, which immediately implies (2.15).

If $\kappa < 0$, for any $x \in B_r(x_0)$, we have

$$\begin{aligned} u(x) &\geq \phi(x) = u(y) + \frac{\kappa}{3r} d(y, x) \\ &= \left(1 - \frac{1}{3r} d(y, x)\right) u(y) + \min_{\xi \in \partial B_{3r}(y)} u(\xi) \geq \frac{u(y)}{3}. \end{aligned}$$

Hence, we obtain (2.15) again.

Let us prove the second statement. Since u is lower semicontinuous, the set

$$\Omega' := \{x \in \Omega : u(x) > 0\}$$

is open in Ω . If $\Omega \neq \Omega'$, then since Ω is connected, there is a point $x \in \partial\Omega' \cap \Omega$. Consequently, we have $u(x) = 0$, which is a contradiction to the first statement. We thus have completed the proof. \square

We next turn to the definition of supersolutions of (1.8).

Definition 2.7 (Supersolutions of ∞ -eigenvalue problem). Let Ω be a domain in a proper geodesic space (\mathbf{X}, d) . A locally Lipschitz function u in Ω is called a supersolution of (1.8) if u is ∞ -superharmonic in Ω and (1.9) holds everywhere in Ω .

Remark 2.8. Thanks to Lemma 2.4, we may drop the local Lipschitz condition in the definition above provided that u is known to be bounded from below in Ω . In particular, any nonnegative ∞ -superharmonic function in Ω is locally Lipschitz .

The idea of adopting the subslope rather than the entire slope to define the so-called Monge solutions of eikonal-type equations stems from the work [41] in the Euclidean space and is recently applied to general complete length spaces in [38]. We refer to [1, 22, 25, 23] etc. for alternative viscosity approaches to Hamilton-Jacobi equations in general metric spaces.

In contrast to the notion of supersolutions, it is less straightforward to define subsolutions in a general metric space. It turns out that the class of ∞ -superharmonic functions itself becomes a natural option if we use it locally to test the candidate function in a strict manner.

Definition 2.9 (Subsolutions of ∞ -eigenvalue problem). Let Ω be a domain in a proper geodesic space (\mathbf{X}, d) . A locally Lipschitz function u in Ω is called a subsolution of (1.8) if whenever there exist $x_0 \in \Omega$, $r_0 > 0$ small and an ∞ -superharmonic function v in $B_{r_0}(x_0) \subset \Omega$ such that $u - v$ attains a strict local maximum at x_0 , the inequality (1.12) holds.

A locally Lipschitz function u in Ω is called a solution of (1.8) if it is both a supersolution and a subsolution.

If $|\nabla^- v|$ is known to be lower semicontinuous at x_0 , then (1.12) can be rewritten as $|\nabla^- v|(x_0) \leq \lambda u(x_0)$. However, in general we only have upper semicontinuity of $|\nabla^- v|$ due to Lemma 2.5 and the lower semicontinuity of solutions may fail to hold, as shown in Case 3 of Example 5.6, Section 5.3.

Let us construct more ∞ -superharmonic functions for our later use.

Lemma 2.10 (∞ -superharmonic functions by composition). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let v be an ∞ -superharmonic function in Ω . Let $h \in C^2(\mathbb{R})$ satisfy*

$$h'(v(x)) > 0, \quad h''(v(x)) < 0 \quad \text{for all } x \in \Omega. \quad (2.16)$$

Then, for any cone function given by (1.10) with $a \in \mathbb{R}$, $\kappa \leq 0$ and $\hat{x} \in \Omega$, $h(v) - \phi$ cannot attain a local minimum in $\Omega \setminus \{\hat{x}\}$. In particular, $h(v)$ is ∞ -superharmonic in Ω .

Proof. Suppose by contradiction that there exists a bounded open set $\mathcal{O} \subset \subset \Omega$ such that $h(v) - \phi$ attains a minimum at $x_0 \in \mathcal{O}$ for a cone function given in (1.10) with $a \in \mathbb{R}$, $\kappa \leq 0$ and $\hat{x} \in \Omega \setminus \mathcal{O}$. By changing the value of a , we may assume that $h(v(x_0)) = \phi(x_0)$ and $h(v) \geq \phi$ in $B_r(x_0)$ with $r > 0$ small such that $\hat{x} \notin B_r(x_0)$.

Then by assumptions, h admits an inverse function h^{-1} , of C^2 class, near $v(x_0)$. It follows that $v(x_0) = h^{-1}(\phi(x_0))$ and $v \geq h^{-1}(\phi)$ in $B_r(x_0)$. In addition, noticing that h^{-1} is strictly convex near $\phi(x_0)$, by letting $r > 0$ further small if necessary, we get

$$v(x) \geq h^{-1}(\phi(x)) > \psi_0(x)$$

for all $x \in B_r(x_0) \setminus \{x_0\}$, where we define

$$\psi_0(x) := v(x_0) + \frac{\kappa}{h'(v(x_0))} (d(x_0, \hat{x}) - d(x, \hat{x})), \quad x \in B_r(x_0).$$

On the other hand, we have $v(x_0) = \psi_0(x_0)$. We therefore can take a cone function $\psi_\varepsilon = \psi_0 + \varepsilon$ with apex at $\hat{x} \notin B_r(x_0)$ and $\varepsilon > 0$ sufficiently small so that $\psi_\varepsilon \leq v$ on $\partial B_r(x_0)$ but $\psi_\varepsilon(x_0) > v(x_0)$. This means that u fails to obey the comparison with cones from below, which is clearly a contradiction to the assumption that v is ∞ -superharmonic. \square

Since $|\nabla^- h(v)| = h'(v)|\nabla^- v|$ holds for any $h \in C^2(\mathbb{R})$, the result above amounts to saying that any composite function $h(v)$ serves as a test function for subsolutions in Ω provided that v is ∞ -superharmonic, $|\nabla^- v| > 0$ in Ω and h satisfies (2.16).

3. EIGENVALUE AND EIGENFUNCTIONS

In this section, we study the eigenvalue problem (1.3) associated to the infinity Laplacian. We generalize the notion of the radius of the maximal inscribed metric ball in Ω ; namely, we take R_∞ as in (1.6). This generalized radius R_∞ can be used to introduce the value Λ_∞ as in (1.2).

It is not difficult to see that $R_\infty > 0$ in (1.6) can be expressed by

$$R_\infty = \max_{x \in \overline{\Omega}} d(x, \partial\Omega).$$

We introduce a notion of the principal eigenvalue and show that it is indeed the value Λ_∞ given in (1.2). We next discuss a definition and properties of the corresponding eigenfunction.

3.1. The eigenvalue. Let us begin with our notion of the principal eigenvalue associated to the ∞ -Laplacian in geodesic spaces.

Definition 3.1 (∞ -eigenvalue). Let (\mathbf{X}, d) be a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ be a bounded domain. The value $\Lambda \in \mathbb{R}$ given by (1.13) is called the principal eigenvalue for the ∞ -Laplacian in Ω with the Dirichlet condition (1.4).

Our first main result, Theorem 1.1, states that the ∞ -eigenvalue Λ as in (1.13) coincides with Λ_∞ , the reciprocal of $R_\infty > 0$ in (1.6).

In order to prove Theorem 1.1, we present the following result.

Proposition 3.2 (Existence of typical supersolutions). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Assume that $g \in C(\partial\Omega)$. For $\lambda > 0$, set*

$$u(x) = \min_{y \in \partial\Omega} \{g(y) + \lambda d(x, y)\} \quad \text{for } x \in \overline{\Omega}. \quad (3.1)$$

Then u is ∞ -superharmonic and $|\nabla^- u| = \lambda$ holds in Ω .

Proof. The function u given by (3.1) is known as the McShane-Whitney Lipschitz extension. By Theorem A.1, we see that u is Lipschitz in $\overline{\Omega}$ and $|\nabla^- u| = \lambda$ in Ω . It thus suffices to prove that u is ∞ -superharmonic in Ω .

Let $\mathcal{O} \subset\subset \Omega$ be a bounded open set and fix $\hat{x} \in \Omega \setminus \mathcal{O}$. For any $a \in \mathbb{R}$ and $\kappa \leq 0$, let ϕ be given by (1.10). Suppose that (2.2) holds. We aim to show that (2.3) holds.

Assume by contradiction that this fails to hold. Then there exists $x_0 \in \mathcal{O}$ such that

$$\max_{x \in \overline{\mathcal{O}}} (\phi - u)(x) = (\phi - u)(x_0) = \mu \quad (3.2)$$

for some $\mu > 0$. Due to the maximality at x_0 , it is not difficult to see that

$$\lambda = |\nabla^- u|(x_0) \leq |\nabla^- \phi|(x_0) \leq |\nabla \phi|(x_0) = -\kappa. \quad (3.3)$$

Note that there exists a Lipschitz curve $\bar{\gamma}$ with $\bar{\gamma}(0) = x_0$, $\bar{\gamma}(1) = \hat{x}$ and $\ell(\bar{\gamma}) = d(x_0, \hat{x})$. Let

$$t_0 = \inf\{t > 0 : \bar{\gamma}(t) \notin \mathcal{O}\}.$$

Then $y_0 = \bar{\gamma}(t_0) \in \partial\mathcal{O}$. It is clear that

$$d(\hat{x}, x_0) = d(\hat{x}, y_0) + d(y_0, x_0). \quad (3.4)$$

Moreover, we have

$$d(y_0, y) \leq d(y_0, x_0) + d(x_0, y) \quad \text{for all } y \in \partial\mathcal{O},$$

which, by (3.2), implies that

$$\begin{aligned} u(y_0) &= \min_{y \in \partial\Omega} \{g(y) + \lambda d(y_0, y)\} \\ &\leq \lambda d(y_0, x_0) + \min_{y \in \partial\Omega} \{g(y) + \lambda d(x_0, y)\} \\ &\leq \lambda d(y_0, x_0) + a + \kappa d(\hat{x}, x_0) - \mu. \end{aligned}$$

It follows from (3.3) and (3.4) that

$$u(y_0) \leq \kappa d(\hat{x}, x_0) - \kappa d(y_0, x_0) + a - \mu = a + \kappa d(\hat{x}, y_0) - \mu,$$

which is clearly a contradiction to (2.2). \square

Remark 3.3 (A distance-type supersolution). Applying Proposition 3.2 with $g \equiv 0$ on $\partial\Omega$, we can see that

$$u(x) = \lambda d(x, \partial\Omega)$$

is ∞ -superharmonic and satisfies $|\nabla^- u| \geq \lambda$ in Ω . In particular, we see that u_{dist} given by (1.14) is a supersolution of (1.3), since $u_{dist} \leq 1$ holds in Ω .

We now proceed to the proof of Theorem 1.1.

Proof of Theorem 1.1. In view of Remark 3.3, it is clear that $\Lambda_\infty \leq \Lambda$. Let us now prove $\Lambda \leq \Lambda_\infty$. Suppose that there exists a locally Lipschitz positive supersolution u of (1.8) for some $\lambda > 0$. Let $x_0 \in \Omega$ be an incenter, which satisfies (1.15).

Fix $\varepsilon > 0$ arbitrarily small. Noticing that for $s = R_\infty - \varepsilon$, $u \geq \phi$ holds on $\partial B_s(x_0)$, where

$$\phi(x) := u(x_0) - \frac{u(x_0)}{s} d(x, x_0) \quad \text{for } x \in \Omega,$$

by Lemma 2.3, we have

$$|\nabla^- u|(x_0) \leq \frac{u(x_0)}{R_\infty - \varepsilon}.$$

Letting $\varepsilon \rightarrow 0$, we end up with

$$|\nabla^- u|(x_0) \leq \Lambda_\infty u(x_0), \quad (3.5)$$

which implies that $\lambda \leq \Lambda_\infty$. It then follows from (1.13) that $\Lambda \leq \Lambda_\infty$. \square

3.2. Existence of eigenfunctions. Let us now investigate the existence of eigenfunctions. In this section, we aim to prove Theorem 1.2, which states that u_∞ defined by (1.18) is a solution of (1.3) and (1.4). We remark that u_∞ is well defined, since by 3.3, the function class for the infimum in (1.18) is non-empty.

Let us now use Perron's method to prove Theorem 1.2. We begin with the supersolution property of u_∞ .

Theorem 3.4 (Supersolution property of infimum). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u_\infty : \overline{\Omega} \rightarrow \mathbb{R}$ be defined by (1.18). Then u_∞ is a positive supersolution of (1.3) satisfying (1.16).*

To prove Theorem 3.4, we prepare two results regarding the pointwise infima of supersolutions to the ∞ -Laplace equation and to the eikonal equation respectively.

Proposition 3.5 (Supersolution preserving of infimum for ∞ -Laplacian). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let \mathcal{S}_I be a family of nonnegative ∞ -superharmonic functions in Ω . Then $w(x) = \inf\{u(x) : u \in \mathcal{S}_I\}$ is also ∞ -superharmonic in Ω .*

Proof. We only need to show that w enjoys the property of comparison with cones from below. To see this, fix a bounded open set $\mathcal{O} \subset\subset \Omega$ and $\hat{x} \in \Omega \setminus \mathcal{O}$ and take a cone function ϕ as in (1.10). If $w \geq \phi$ on $\partial\mathcal{O}$, then by definition $u \geq \phi$ on $\partial\mathcal{O}$ for all $u \in \mathcal{S}_I$. It follows that $u \geq \phi$ in \mathcal{O} for all $u \in \mathcal{S}_I$, which in turn implies that $w \geq \phi$ in \mathcal{O} . \square

Proposition 3.6 (Supersolution preserving of infimum for eikonal equation). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let \mathcal{S}_E be a family of nonnegative locally Lipschitz functions satisfying (1.9) in Ω for some $\lambda > 0$. Assume that $w(x) = \inf\{u(x) : u \in \mathcal{S}_E\}$ is locally Lipschitz in Ω . Then w is also a nonnegative function satisfying the same inequality in Ω .*

Proof. It suffices to prove that $W := \log w$

$$|\nabla^- v| \geq \lambda \quad \text{in } \Omega.$$

Fix any $x_0 \in \Omega$ and take $r > 0$ arbitrarily small. For any $u \in \mathcal{S}_E$ letting $U = \log u$, we see that U is a locally Lipschitz and satisfies $|\nabla^- U| \geq \lambda$ in $B_r(x_0)$. It is clear that W is the pointwise infimum over all such U .

As shown in Proposition 3.2 (and in Theorem A.1), the McShane-Whitney Lipschitz extension for W

$$\underline{W}(x) = \min_{y \in \partial B_r(x_0)} \{W(y) + \lambda d(x, y)\}, \quad x \in \overline{B}_r(x_0),$$

satisfies $|\nabla^- \underline{W}| = \lambda$ in $B_r(x_0)$. Besides, it is easily seen that $\underline{W} \leq W \leq U$ on $\partial\Omega$. We then can adopt the comparison principle, Theorem A.2 (or [38, Theorem 4.2]), to deduce that $\underline{W} \leq U$ in $B_r(x_0)$, where we recall that $U = \log u$ for each $u \in \mathcal{S}_E$.

By taking the infimum over all such u , we obtain

$$\underline{W} \leq W \quad \text{in } B_r(x_0).$$

In particular, there exists $y_r \in \partial B_r(x_0)$ such that

$$W(x_0) \geq \underline{W}(x_0) \geq W(y_r) + \lambda d(x_0, y_r)$$

and therefore

$$\frac{W(x_0) - W(y_r)}{d(x_0, y_r)} \geq \lambda.$$

Sending $r \rightarrow 0$, we get

$$|\nabla^- W|(x_0) \geq \limsup_{r \rightarrow 0} \frac{W(x_0) - W(y_r)}{d(x_0, y_r)} \geq \lambda.$$

We complete the proof due to the arbitrariness of x_0 in Ω . \square

We are now in a position to prove Theorem 3.4.

Proof of Theorem 3.4. By definition, it is clear that u_∞ satisfies (1.16). In view of Proposition 3.5, we see that u_∞ is ∞ -superharmonic. Using Lemma 2.4, we obtain local Lipschitz continuity of u_∞ . By Proposition 2.6 and the fact that $u_\infty = 1$ on $\mathcal{M}(\Omega)$, we further deduce that $u_\infty > 0$ in Ω . Using Proposition 3.6, we have $|\nabla^- u_\infty| \geq \Lambda_\infty u_\infty$ in Ω . Our proof is thus complete. \square

We complete the proof of Theorem 1.2 by combining Theorem 3.4 with Theorem (3.7) below, which states that u_∞ is also a subsolution of (1.3) in the sense of Definition 2.9.

Theorem 3.7 (Subsolution property of infimum of supersolutions). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u_\infty : \overline{\Omega} \rightarrow \mathbb{R}$ be defined by (1.18). Then u_∞ is continuous in $\overline{\Omega}$ and is a subsolution of (1.3) satisfying (1.4).*

Proof. We have shown in Theorem 3.4 that u_∞ is a positive supersolution of (1.3). Note that the definition of u_∞ , together with Remark 3.3, yields

$$u_\infty \leq u_{dist} = \Lambda_\infty d(\cdot, \partial\Omega) \quad \text{in } \overline{\Omega}.$$

It is then easily seen that $u_\infty \in C(\overline{\Omega})$ and $u_\infty = 0$ on $\partial\Omega$.

Let us focus on the subsolution property of u_∞ . Suppose by contradiction that u_∞ is not a subsolution of (1.3). This means that there exist $r, \sigma > 0$ small and $x_0 \in \Omega$ with $B_r(x_0) \subset\subset \Omega$ such that

$$u_\infty(x) - u(x) < u_\infty(x_0) - u(x_0) = 0$$

for all $x \in B_r(x_0) \setminus \{x_0\}$, where u is an ∞ -superharmonic function satisfying

$$|\nabla^- u| \geq \Lambda_\infty u_\infty + \sigma \quad \text{in } B_r(x_0). \quad (3.6)$$

As shown in the proof of Theorem 1.1, we obtain (3.5) when x_0 is an incenter, i.e., $x_0 \in \mathcal{M}(\Omega)$. This yields $x_0 \notin \mathcal{M}(\Omega)$. We thus can take $r > 0$ small such that $B_r(x_0) \cap \mathcal{M}(\Omega) = \emptyset$.

We next take $\varepsilon > 0$ small such that

$$u - \varepsilon \geq u_\infty \quad \text{on } \Omega \setminus B_{r-\varepsilon}(x_0). \quad (3.7)$$

We further take

$$\tilde{u}(x) = \begin{cases} \min\{u_\infty(x), u(x) - \varepsilon\} & \text{if } x \in B_r(x_0), \\ u_\infty(x) & \text{if } x \in \overline{\Omega} \setminus B_r(x_0). \end{cases}$$

It is clear that \tilde{u} is continuous and positive in Ω . One can also easily observe that u_∞ satisfies (1.16) and

$$\tilde{u}(x_0) \leq u_\infty(x_0) - \varepsilon. \quad (3.8)$$

Let us below prove that \tilde{u} is a supersolution of (1.3). By Proposition 3.5, we deduce that \tilde{u} is ∞ -superharmonic in $B_r(x_0)$. Moreover, for any $x, y \in B_r(x_0)$ we have

$$\tilde{u}(x) - \tilde{u}(y) \geq \min\{u_\infty(x) - u_\infty(y), u(x) - u(y)\},$$

which yields

$$|\nabla^- \tilde{u}|(x) \geq \min\{|\nabla^- u_\infty|(x), |\nabla^- u|(x)\}$$

for all $x \in B_r(x_0)$. Since $|\nabla^- u_\infty| \geq \Lambda_\infty u_\infty$ and (3.6) holds, it follows that

$$|\nabla^- \tilde{u}| \geq \Lambda_\infty \tilde{u}. \quad (3.9)$$

in $B_r(x_0)$. Noticing that $\tilde{u} = u_\infty$ in $\Omega \setminus B_{r-\varepsilon}(x_0)$ due to (3.7), we thus see that (3.9) holds in Ω .

It remains to verify that \tilde{u} is ∞ -superharmonic in Ω . Suppose that there exist a bounded open set $\mathcal{O} \subset\subset \Omega$, $\hat{x} \in \Omega \setminus \mathcal{O}$ and a cone function as in (1.10) such that $\tilde{u} \geq \phi$ on $\partial\mathcal{O}$. Since $\tilde{u} \leq u_\infty$, we have $u_\infty \geq \phi$ on $\partial\mathcal{O}$. Noticing that u_∞ is ∞ -superharmonic, we obtain $u_\infty \geq \phi$ in $\overline{\mathcal{O}}$ and in particular

$$\tilde{u} \geq \phi \quad \text{in } \overline{\mathcal{O}} \setminus B_{r-\varepsilon}(x_0). \quad (3.10)$$

It follows that

$$u - \varepsilon \geq \phi \quad \text{on } \partial\mathcal{O} \setminus B_{r-\varepsilon}(x_0),$$

which implies that $u - \varepsilon \geq \phi$ on $\partial(B_{r-\varepsilon}(x_0) \cap \mathcal{O})$.

Since u is ∞ -superharmonic in $B_r(x_0)$, we obtain $u - \varepsilon \geq \phi$ in $B_{r-\varepsilon}(x_0) \cap \mathcal{O}$. Combining this with (3.10), we are led to $\tilde{u} \geq \phi$ in $\overline{\mathcal{O}}$. Hence, we conclude that \tilde{u} is a positive supersolution of (1.3).

Since \tilde{u} also satisfies (1.16) and (3.8), we reach a contradiction to the definition of u_∞ as in (1.18). Our proof is now complete. \square

Remark 3.8 (Partial incenter constraints). Our argument above can be used to construct more solutions of (1.3) when $\mathcal{M}(\Omega)$ is not a singleton. Recall that $\mathcal{M}(\Omega)$, defined by (1.17), denotes the set of incenters. In fact, for any given compact subset $(\emptyset \neq) Y \subset \mathcal{M}(\Omega)$, replacing the condition (1.16) by (1.21) in (1.18), we can take u_∞^Y as in (1.22). (It is clear that $u_\infty^Y = u_\infty$ when $Y = \mathcal{M}(\Omega)$.) Then, following the proof of Theorem 3.4 and Theorem 3.7, we can prove in a similar way that u_∞^Y is also a positive solution of (1.3). In Section 5.4, we present a concrete example on metric graphs to show that u_∞^Y and u_∞ are really different in general.

3.3. A weak comparison principle. We next turn to discuss the uniqueness of linearly independent eigenfunctions in general geodesic spaces. Although we believe that u_∞ is the only correct eigenfunction to the eigenvalue problem (1.3), the uniqueness of eigenfunctions up to a constant multiple is however still unclear. The uniqueness issue remains to be a challenging open question even for a convex domain $\Omega \subset \mathbb{R}^n$. We refer to [28, 48, 27, 18] for some partial results on this problem.

We present a comparison principle, assuming the subsolution and supersolution are both positive in $\overline{\Omega}$, which extends the result [30, Theorem 2.3] in the Euclidean case to our general setting. Unfortunately, due to the uniform positivity assumption, this comparison principle is weaker than the usual type and is not sufficient for us to obtain the desired uniqueness of eigenfunctions.

Theorem 3.9 (Comparison principle for eigenvalue problem with uniform positivity). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let $u, v \in C(\overline{\Omega})$ satisfy*

$$\min_{x \in \overline{\Omega}} \min\{u(x), v(x)\} > 0 \quad (3.11)$$

and be respectively a subsolution and a supersolution of (1.3). If $u \leq v$ on $\partial\Omega$, then $u \leq v$ in $\overline{\Omega}$.

Proof. Assume by contradiction that there exists $\bar{x} \in \Omega$ such that $u(\bar{x}) - v(\bar{x}) > \sigma$ for some $\sigma > 0$. By appropriate constant multiplication for both u and v , we may assume that there exist $\tau_1, \tau_2 > 0$ such that

$$0 < \tau_1 \leq u, v \leq \tau_2 < 1 \quad \text{in } \overline{\Omega}. \quad (3.12)$$

In view of (3.12), we can find $A > 1$ and $\alpha < 1$ close to 1 such that for all $\tau_1 \leq t \leq \tau_2$,

$$At^{\frac{1}{\alpha}} + 1 - A > 0 \quad (3.13)$$

and

$$\log u \leq \log h(v) - \sigma \quad \text{on } \partial\Omega \quad (3.14)$$

hold, where h is given by

$$h(t) = (At^{\frac{1}{\alpha}} + 1 - A)^{\alpha}, \quad t > 0.$$

The relation (3.13) yields

$$\inf_{\overline{\Omega}} h(v) \geq h(\tau_1) > 0.$$

By direct computation, we see that

$$h(t)^{\frac{1}{\alpha}} - t^{\frac{1}{\alpha}} = (A - 1)(t^{\frac{1}{\alpha}} - 1) < 0$$

and thus

$$\max_{\tau_1 \leq t \leq \tau_2} (h(t) - t) < 0. \quad (3.15)$$

Moreover, by direct calculation, we get

$$h'(t) = A(At^{\frac{1}{\alpha}} + 1 - A)^{\alpha-1} t^{\frac{1-\alpha}{\alpha}} > A^{\alpha} > 1$$

and

$$h''(t) = \frac{1}{\alpha}(\alpha - 1)A(A - 1)(At^{\frac{1}{\alpha}} + 1 - A)^{\alpha-2} t^{\frac{1-2\alpha}{2}} < 0$$

for any $\tau_1 \leq t \leq \tau_2$. In other words, we have

$$\min_{\tau_1 \leq t \leq \tau_2} h'(t) > 1, \quad \max_{\tau_1 \leq t \leq \tau_2} h''(t) < 0. \quad (3.16)$$

Since $u(\bar{x}) - h(v(\bar{x})) > \sigma$ and (3.14) holds, there exist $x_0 \in \Omega$ and $\mu > 0$ such that

$$\max_{\overline{\Omega}} (\log u - \log(h(v))) = \log u(x_0) - \log h(v(x_0)) = \mu,$$

which yields

$$u(x) - e^{\mu} h(v)(x) \leq u(x_0) - e^{\mu} h(v)(x_0) = 0$$

for all $x \in \Omega$.

Let us slightly change h by taking

$$h_{\varepsilon}(t) = h(t) + \varepsilon(t - h(v(x_0)))^2, \quad t > 0$$

for $\varepsilon > 0$ small. By (3.15) and (3.16), we can take $\varepsilon > 0$ sufficiently small such that, for any $t > 0$ close to $v(x_0)$,

$$h_\varepsilon(t) < t \quad (3.17)$$

and

$$h'_\varepsilon(t) > 1, \quad h''_\varepsilon(t) < 0. \quad (3.18)$$

It is also clear that $h_\varepsilon(v(x_0)) = h(v(x_0))$.

Since v is a supersolution of (1.3), using the estimate for h'_ε in (3.18), we deduce that

$$|\nabla^- h_\varepsilon(v)| - \Lambda_\infty h_\varepsilon(v) = h'_\varepsilon |\nabla^- v| - \Lambda_\infty h_\varepsilon(v) > \Lambda_\infty v - \Lambda_\infty h_\varepsilon(v)$$

in $B_r(x_0)$ with $r > 0$ sufficiently small. Together with (3.17), this means that, when $r > 0$ is small,

$$\inf_{B_r(x_0)} (|\nabla^- h_\varepsilon(v)| - \Lambda_\infty h_\varepsilon(v)) > 0. \quad (3.19)$$

In addition, by Lemma 2.10, $h_\varepsilon(v)$ is ∞ -superharmonic in $B_r(x_0)$ due to (3.18) again.

On the other hand, we see that $u(x) - e^\mu h_\varepsilon(v)(x)$ attains a strict maximum in $B_r(x_0)$ at $x = x_0$ with $u(x_0) = e^\mu h_\varepsilon(v)(x_0)$. We then get a contradiction by taking into account that u is a subsolution of (1.3). In fact, noticing that $e^\mu h_\varepsilon(v)$ is also ∞ -superharmonic in $B_r(x_0)$, by Definition 2.9 we have

$$\lim_{r \rightarrow 0^+} \inf_{B_r(x_0)} (e^\mu |\nabla^- h_\varepsilon(v)| - \Lambda_\infty e^\mu h_\varepsilon(v)) \leq 0.$$

This is a contradiction to (3.19). \square

The strong assumption (3.11) demands that the positive functions u and v be uniformly bounded away from 0. Therefore Theorem 3.9 alone does not lead us to general uniqueness of solutions of (1.3) and (1.4), even among those satisfying (1.16). In order to adopt this comparison result to prove the uniqueness, we somehow need to first show that the uniqueness holds near the boundary.

Corollary 3.10 (Uniqueness with positive data near boundary). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let Ω_δ be a subdomain of Ω given by*

$$\Omega_\delta = \{x \in \Omega : d(x, \partial\Omega) > \delta\} \quad (3.20)$$

for $\delta > 0$ small and g be a positive locally Lipschitz function in $\Omega \setminus \Omega_\delta$. Then the positive solutions of (1.3) satisfying

$$u = g \quad \text{in } \Omega \setminus \Omega_\delta$$

are unique if they exist.

This corollary can be obtained immediately by adopting Theorem 3.9. In fact, one can take the domain Ω in Theorem 3.9 to be Ω_δ here.

4. CONSISTENCY WITH THE EUCLIDEAN CASE

By Theorem 1.1, we have seen that our ∞ -eigenvalue is consistent with that in the Euclidean case. In this section we further show that our definition of eigenfunctions in geodesic spaces, as given in Definition 2.9, is also a generalization of the notion proposed in [30] in the Euclidean space.

Let us recall the definition of viscosity supersolutions of (1.8) in the Euclidean space, which is implicitly given in [30] as follows. See [16] for definitions of viscosity solutions to general nonlinear elliptic equations. Let Ω be a bounded domain in \mathbb{R}^n and $\lambda > 0$.

Definition 4.1 (Definition of Euclidean viscosity solutions). A locally bounded lower semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is called a viscosity subsolution of (1.8) if whenever there exist $x_0 \in \Omega$ and $\varphi \in C^2(\Omega)$ such that $u - \varphi$ attains a strict local minimum in Ω at x_0 , both

$$|\nabla\varphi(x_0)| \geq \lambda u(x_0) \quad (4.1)$$

and

$$-\Delta_\infty\varphi(x_0) \geq 0 \quad (4.2)$$

hold. A locally bounded upper semicontinuous function $u : \Omega \rightarrow \mathbb{R}$ is called a viscosity subsolution of (1.8) if whenever there exist $x_0 \in \Omega$ and $\varphi \in C^2(\Omega)$ such that $u - \varphi$ attains a strict local maximum in Ω at x_0 , either

$$|\nabla\varphi(x_0)| \leq \lambda u(x_0) \quad (4.3)$$

or

$$-\Delta_\infty\varphi(x_0) \leq 0$$

holds. A function $u \in C(\Omega)$ is called a viscosity solution of (1.8) if it is both a viscosity supersolution and a viscosity subsolution.

Remark 4.2. As is well known in the theory of viscosity solutions, one may use the semijets instead of the test functions to define super- and subsolutions. More precisely, $u \in C(\Omega)$ is a viscosity supersolution (resp., subsolution) if for any $x_0 \in \Omega$, we have

$$\begin{aligned} |p| \geq \lambda u(x_0) \quad \text{and} \quad -\langle Xp, p \rangle \geq 0 \\ (\text{resp., } |p| \leq \lambda u(x_0) \quad \text{or} \quad -\langle Xp, p \rangle \leq 0) \end{aligned}$$

for every $(p, X) \in \overline{\mathcal{J}}^{2,-}u(x_0)$ (resp., $(p, X) \in \overline{\mathcal{J}}^{2,+}u(x_0)$). We refer to [16] for a detailed introduction on the semijets $\overline{\mathcal{J}}^{2,\pm}u(x_0) \subset \mathbb{R}^n \times \mathcal{S}^n$, where \mathcal{S}^n denotes the set of all $n \times n$ real-valued symmetric matrices.

The definition above is indeed consistent with the standard framework of viscosity solutions. For a general fully nonlinear elliptic equation

$$F(x, u, \nabla u(x), \nabla^2 u) = 0 \quad \text{in } \Omega \subset \mathbb{R}^n,$$

where $F : \Omega \times \mathbb{R} \times \mathbb{R}^n \times \mathcal{S}^n \rightarrow \mathbb{R}$ is continuous, we define a viscosity supersolution (resp., subsolution) u by asking

$$F(x, u(x), p, X) \geq 0 \quad (\text{resp., } F(x, u(x), p, X) \leq 0)$$

for all $(p, X) \in \overline{\mathcal{J}}^{2,-}u(x)$ (resp., $(p, X) \in \overline{\mathcal{J}}^{2,+}u(x)$). See again [16] for details.

The main result of this section is as follows.

Theorem 4.3 (Equivalence of solutions). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and $\lambda > 0$. Let u be a nonnegative continuous function in Ω . Then u is a solution of (1.8) in the sense of Definition 2.7 and Definition 2.9 if and only if u is a viscosity supersolution of (1.8) as defined in Definition 4.1.*

Before starting to prove this theorem, we first present the following equivalence result on ∞ -superharmonic functions, which is essentially obtained in [4].

Lemma 4.4 (Equivalence of ∞ -superharmonic functions). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. Then u is ∞ -superharmonic in $\Omega \subset \mathbb{R}^n$ in the sense of Definition 2.1 if and only if u is a viscosity supersolution of (2.1), that is, whenever there exist $x_0 \in \Omega$ and $\varphi \in C^2(\Omega)$ such that $u - \phi$ attains a strict local minimum at x_0 , the inequality (4.2) holds.*

Proof. In [15, Theorem 3.1] and [4, Theorem 4.13], it is proved that u is a viscosity supersolution of (2.1) if and only if u satisfies the property of comparison with cones from below. Note that in these results each test cone ϕ in an open subset \mathcal{O} is in the form of (1.10) with $a, \kappa \in \mathbb{R}$ and $\hat{x} \in \Omega \setminus \mathcal{O}$.

In order to complete the proof, it suffices to show that in the Euclidean space this more restrictive condition (with $\kappa \in \mathbb{R}$) of cone comparison is equivalent to the version with $\kappa \geq 0$ as stated in Definition 2.1. This can be found in [31]; see the remarks after Definition 2.3 in [31], where a sufficient condition is given for the equivalence of both types of comparison with cones in more general metric spaces. \square

We first show the equivalence between the notions of supersolutions.

Proposition 4.5 (Equivalence of supersolutions). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and $\lambda > 0$. Let u be a nonnegative continuous function in Ω . Then u is a supersolution of (1.8) in the sense of Definition 2.7 if and only if u is a viscosity supersolution of (1.8) as defined in Definition 4.1.*

Proof. Let us first show “ \Rightarrow ”. Suppose that u is locally Lipschitz and ∞ -superharmonic, and (1.9) holds in Ω . Assume that $u - \varphi$ attains a strict local minimum at $x_0 \in \Omega$ for some $\varphi \in C^2(\Omega)$. By Lemma 4.4, we have (4.2). Also, it is not difficult to see that

$$|\nabla \varphi|(x_0) = |\nabla^- \varphi|(x_0) \geq |\nabla^- u|(x_0),$$

which, together with (1.9), immediately yields (4.1). Hence, we conclude that u is a viscosity supersolution of (1.8).

We next prove “ \Leftarrow ”. Suppose that u is viscosity supersolution of (1.8). Applying Lemma 4.4 again, we deduce that u satisfies the property of comparison with cones in Ω . In particular, u is locally Lipschitz by Lemma 2.4. It remains to show (1.9) in Ω .

Fix $x_0 \in \Omega$ arbitrarily. Thanks to the ∞ -superharmonicity of u , for any $\sigma > 0$ small, we can find $r > 0$ small such that

$$u(x) > u(x_0) - (|\nabla^- u|(x_0) + \sigma)d(x_0, x)$$

for all $x \in \overline{B_r(x_0)} \setminus \{x_0\}$. It amounts to saying that $\phi_\sigma - u$ attains a strict maximum in $\overline{B_r(x_0)}$ at x_0 , where

$$\phi_\sigma(x) := u(x_0) - (|\nabla^- u|(x_0) + \sigma)d(x_0, x), \quad x \in \overline{B_r(x_0)}.$$

Let us consider

$$\Phi_\varepsilon(x, y) = \phi_\sigma(x) - u(y) - \frac{|x - y|^2}{\varepsilon}$$

for $\varepsilon > 0$ and $x, y \in \overline{B_r(x_0)}$. Suppose that $(x_\varepsilon, y_\varepsilon)$ is a maximizer of Φ_ε in $\overline{B_r(x_0)} \times \overline{B_r(x_0)}$. By a standard argument of viscosity solutions, we have $x_\varepsilon, y_\varepsilon \rightarrow x_0$ as $\varepsilon \rightarrow 0$. In particular, we get $x_\varepsilon, y_\varepsilon \in B_r(x_0)$ when $\varepsilon > 0$ is taken small.

Since

$$y \mapsto u(y) + \frac{|x_\varepsilon - y|^2}{\varepsilon} - \phi_\sigma(x_\varepsilon)$$

attains a local minimum at $y = y_\varepsilon$, we can apply the supersolution part of Definition 4.1 to deduce that

$$\frac{2|x_\varepsilon - y_\varepsilon|}{\varepsilon} \geq \lambda u(y_\varepsilon). \quad (4.4)$$

On the other hand, the maximality at $x = x_\varepsilon$ of

$$x \mapsto \phi_\sigma(x) - u(y_\varepsilon) - \frac{|x - y_\varepsilon|^2}{\varepsilon}$$

yields

$$\frac{2|x_\varepsilon - y_\varepsilon|}{\varepsilon} \leq |\nabla^- \phi_\sigma|(x_\varepsilon) = |\nabla^- u|(x_0) + \sigma. \quad (4.5)$$

Combining (4.4) and (4.5), we are led to

$$|\nabla^- u|(x_0) + \sigma \geq \lambda u(y_\varepsilon).$$

Our proof is thus complete if we send $\varepsilon \rightarrow 0$ and then $\sigma \rightarrow 0$ in the relation above. \square

We next show the equivalence for subsolutions. Our proof is based on a comparison-type argument.

Proposition 4.6 (Equivalence of subsolutions). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain and $\lambda > 0$. Let u be a nonnegative continuous function in Ω . Then u is a supersolution of (1.8) in the sense of Definition 2.9 if and only if u is a viscosity subsolution of (1.8) as defined in Definition 4.1.*

Proof. We again begin with the proof of “ \Rightarrow ”. Assume that u is locally Lipschitz and there exist x_0 and $\varphi \in C^2(\Omega)$ such that $u - \varphi$ attains a strict local maximum at x_0 . If $-\Delta_\infty \varphi(x_0) > 0$, then $-\Delta_\infty \varphi > 0$ in $B_r(x_0)$ for some $r > 0$ small. In view of Lemma 4.4, we know that φ is ∞ -superharmonic. It thus follows from Definition 2.9 that

$$\lim_{r \rightarrow 0^+} \inf_{B_r(x_0)} (|\nabla^- \varphi| - \lambda u) \leq 0. \quad (4.6)$$

This implies (4.3), since $|\nabla^- \varphi| = |\nabla \varphi|$ is continuous.

Let us show the reverse implication “ \Leftarrow ”. Let u be locally Lipschitz in Ω . Assume that there exist $x_0 \in \Omega$, $r > 0$ small and a function φ that is ∞ -superharmonic in $B_r(x_0)$ such that $u - \varphi$ attains a strict maximum in $B_r(x_0)$ at x_0 . By Lemma 4.4, we see that $-\Delta_\infty \varphi \geq 0$ in $B_r(x_0)$ holds in the viscosity sense. We aim to get (4.6).

By contradiction we may assume that there exists $\sigma > 0$ small such that

$$|\nabla^- \varphi| - \lambda u \geq \sigma \quad \text{in } \overline{B_r(x_0)}. \quad (4.7)$$

In particular, we see that $|\nabla \varphi| \geq \sigma$ holds in the viscosity sense in $B_r(x_0)$. Therefore, by subtracting a quadratic function $a(x - x_0)^2$ with $a > 0$ small, we may also assume that φ satisfies in the viscosity sense

$$-\Delta_\infty \varphi \geq \sigma \quad \text{in } B_r(x_0) \quad (4.8)$$

without affecting other conditions on φ . For each $\varepsilon > 0$, let us consider

$$\Psi_\varepsilon(x, y) = u(x) - \varphi(y) - \frac{|x - y|^2}{\varepsilon}$$

for $x, y \in \overline{B_r(x_0)}$. We can find a maximizer $(x_\varepsilon, y_\varepsilon)$ of Ψ_ε in $\overline{B_r(x_0)} \times \overline{B_r(x_0)}$. As in the proof of Proposition 4.5, we apply a standard argument for comparison principle of viscosity solutions to deduce that $x_\varepsilon, y_\varepsilon \rightarrow x_0$ as $\varepsilon \rightarrow 0$ and therefore $x_\varepsilon, y_\varepsilon \in B_r(x_0)$ for $\varepsilon > 0$ sufficiently small.

Noticing that

$$y \mapsto u(x_\varepsilon) - \varphi(y) - \frac{|x - y|^2}{\varepsilon}$$

attains a maximum at $y = y_\varepsilon$, we apply (4.7) to get

$$\frac{2|x_\varepsilon - y_\varepsilon|}{\varepsilon} \geq |\nabla^- \varphi|(y_\varepsilon) > \lambda u(y_\varepsilon) + \sigma. \quad (4.9)$$

We next adopt the Crandall-Ishii lemma [16] to obtain $(p, X) \in \overline{\mathcal{J}}^{2,+} u(x_\varepsilon)$ and $(q, Y) \in \overline{\mathcal{J}}^{2,-} \varphi(y_\varepsilon)$ satisfying

$$p = q = \frac{2(x_\varepsilon - y_\varepsilon)}{\varepsilon}, \quad (4.10)$$

and

$$\begin{pmatrix} X & 0 \\ 0 & -Y \end{pmatrix} \leq \frac{C}{\varepsilon} \begin{pmatrix} I & -I \\ -I & I \end{pmatrix}.$$

for some $C > 0$. It follows that

$$X \leq Y. \quad (4.11)$$

Using the viscosity inequality (4.8) at y_ε in the form of semijets as in Remark 4.2, we are led to

$$-\langle Yq, q \rangle \geq \sigma,$$

which by (4.10) and (4.11) implies

$$-\langle Xp, p \rangle \geq \sigma > 0. \quad (4.12)$$

Now we apply the alternative definition of subsolutions of (1.8) with semijets on u . We get

$$\frac{2|x_\varepsilon - y_\varepsilon|}{\varepsilon} = |p| \leq \lambda u(x_\varepsilon), \quad (4.13)$$

since (4.12) holds. Combining (4.13) and (4.9), we end up with

$$\lambda u(x_\varepsilon) \geq \lambda u(y_\varepsilon) + \sigma,$$

We reach a contradiction by letting $\varepsilon \rightarrow 0$. □

5. EIGENFUNCTIONS ON METRIC GRAPHS

Let us examine the case when \mathbf{X} is a finite metric graph. In this special case, we present several properties and concrete examples and later discuss the uniqueness issue.

5.1. Preliminaries on metric graphs. Let us first recall the definition of a metric graph. We refer the reader to [12, Section 3.2.2] for a detailed introduction on metric graphs. Hereafter by a metric segment of length $s > 0$ we mean a metric space that is isometric to the interval $[0, s] \subset \mathbb{R}$.

Definition 5.1 (Definition 3.2.9 in [12]). A metric graph is the result of gluing of a disjoint collection of metric segments $\{e_i\}$ and points $\{V_j\}$ (regarded with the length metric of disjoint union) along an equivalence relation defined on the union of the set $\{V_j\}$ and the set of the endpoints of the segments. The segments e_i are called edges, and the equivalence classes of the endpoints are called vertices of the metric graph.

The length of an edge is the length of the segment. The degree of a vertex is defined as the number of endpoints of segments in the corresponding equivalence class. It is thus possible to consider rectifiable curves on a metric graph. Below we always adopt the intrinsic metric for a connected metric graph so that it is a geodesic space.

For a finite topological graph $\mathbf{X} = (\mathcal{V}, \mathcal{E})$, where \mathcal{V} and \mathcal{E} denote respectively the sets of vertices and edges, if the length of each edge is specified, then one can turn it into a finite metric graph by letting the collection of segments $\{e_i\}$ be \mathcal{E} with the given lengths and the common endpoints of edges naturally generates an equivalence relation to define vertices.

A simplest example of metric graphs is the so-called n -multipod with $n \in \mathbb{N}$. More precisely, for metric segments with endpoints O_i and V_i , denoted by $[O_i, V_i]$, for $i = 1, 2, \dots, n$, we consider the endpoints O_i to be equivalent for all $i = 1, 2, \dots, n$. Then the quotient yields a metric space

$$P_n := \bigcup_{i=1,2,\dots,n} [O_i, V_i] / (\text{all } O_i \text{ are identified}),$$

which is called an n -multipod. We denote by O the equivalence class for O_i ($i = 1, 2, \dots, n$) and call it the origin of P_n . In general, a geodesic space (\mathbf{X}, d) is a metric graph if for any $x \in \mathbf{X}$, there exists $n \in \mathbb{N}$ such that $B_r(x) \cap \mathbf{X}$ is isometric to $B_r(O) \cap P_n$ for all $r > 0$ small. This formulation enables us to extend the notion of degree, which is originally for vertices only. Suppose that a metric graph is locally isometric to P_n at x . In this case, we say that x has n directions. We also call each segment isometric to $(O, P_i) \cap B_r(O)$ a direction from x . It is clear that the number of directions from x is constant for all $r > 0$ small.

We emphasize that the presence of loops and multiple edges in metric graphs will be allowed in our analysis. In other words, the metric graphs we consider may contain an edge that connects a vertex to itself, and two or more edges may be incident to the same pair of vertices.

For our convenience in analysis on graphs, we utilize the following notations. For an edge $e = [V, V']$ connecting vertices V and V' , we denote by (x, y) the open connected portion of e between $x, y \in e$. Similarly, we extend the notations $[x, y]$, $[x, y)$ and $(x, y]$ for intervals to denote the portions between $x, y \in e$ with or without endpoints.

5.2. Properties of eigenfunctions on metric graphs. We now study the ∞ -eigenvalue problem on a finite metric graph. Nonlinear PDEs on graphs recently attract much attention; see for example [46, 47, 39] for analysis of finite difference equations associated to various nonlinear operators. Our space setting here is different. We consider directly the continuum equation, which is set up on the whole graph rather than its vertices only.

The one-dimensional structure of metric graphs yields some additional results on geometric properties and regularity of solutions of (1.3).

Proposition 5.2 (Properties of eigenfunctions on metric graphs). *Suppose that $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite connected graph equipped with intrinsic metric d . Let $\Omega \subsetneq \mathbf{X}$ be a domain with $\partial\Omega \subset \mathcal{V}$. Let u be a Lipschitz positive solution of (1.3). Then the following properties hold:*

- (i) *On each edge u is concave and cannot be constant in any open subinterval.*
- (ii) *For any edge $[V_i, V_j]$, if there exists $x \in [V_i, V_j)$ such that $u(y) < u(x)$ for any $y \in (x, V_j)$ close to x , then u is linearly decreasing in (x, V_j) .*

(iii) For any $x \in \Omega$,

$$|\nabla^- u|(x) = \Lambda_\infty u(x) \quad (5.1)$$

holds provided that $|\nabla^- u|(x) > |\nabla^+ u|(x)$.

(iv) For any $x \in \Omega$,

$$|\nabla^- u|(x) > \Lambda_\infty u(x) \quad (5.2)$$

and

$$|\nabla^- u|(x) = |\nabla^+ u|(x) \quad (5.3)$$

hold provided that $|\nabla^+ u|(x) > 0$.

Remark 5.3. The statements (i), (ii), (iii) enable us to understand the behavior of u at a local maximizer in Ω . Let $\hat{x} \in \Omega$ be a local maximizer. Since

$$|\nabla^- u|(x) \geq \Lambda_\infty u(x)$$

holds for any x near \hat{x} , it follows from (i), (ii) and (5.1) that u is linearly decreasing with $|\nabla^- u|(\hat{x})$ along all directions from \hat{x} till the adjacent vertices.

Remark 5.4. The statements (i) and(ii) amount to saying that on each edge the function graph of u is either a line segment or a broken line segment having a unique local maximum. In the latter case, by Remark 5.3, u must be decreasing from the maximizer \hat{x} linearly towards both endpoints of the edge with the same slope $|\nabla^- u|(\hat{x}) = \Lambda_\infty u(\hat{x})$. As a result, we can combine both cases to see that u is piecewise linear on any curve in $\bar{\Omega}$.

As explained in Remark 5.4, there is at most one local maximizer of a solution on each edge. Then the number of local maximizers in the whole domain is also finite, since there are only finitely many edges.

Corollary 5.5 (Finite number of local maximizers). *Suppose that $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite connected graph equipped with intrinsic metric d . Let $\Omega \subsetneq \mathbf{X}$ be a domain with $\partial\Omega \subset \mathcal{V}$. Let u be a Lipschitz positive solution of (1.3). Then there are only finitely many local maximizers of u in Ω .*

Let us now prove Proposition 5.2.

Proof of Proposition 5.2. (i) Since $|\nabla^- u| \geq \Lambda_\infty u$ in Ω , it is clear that at any point in Ω , there is always a direction along which u is decreasing. This implies that u cannot attain a local minimum in Ω . In particular, u cannot be constant in any connected portion of an edge.

In addition, since u is ∞ -superharmonic, parametrizing each edge by $t \mapsto x(t)$ with its length t and applying Lemma 4.4, we get $-v'' \geq 0$ in the interior in the viscosity sense, where $v(t) = u(x(t))$. As an immediate consequence, u is concave on all edges.

(ii) Suppose that u is decreasing near a fixed point $\bar{x} \in [V_i, V_j)$ along a direction toward V_j . Note first that in this case u must be decreasing on (\bar{x}, V_j) , since there are no local minima. In view of (i), we see that u is concave in (\bar{x}, V_j) .

It suffices to show that $v(t)$ is convex in (\bar{x}, V_j) . Let us parametrize (\bar{x}, V_j) by $t \mapsto x(t)$ in terms of the distance $t = d(\cdot, \bar{x})$. Then $v(t) = u(x(t))$ for $t \in [0, T]$, where $T = d(\bar{x}, V_j)$. Assume by contradiction that v is not convex. Then there exist $t_0 \in (0, T)$ and $\psi \in C^2((0, T))$ with $\psi''(t_0) \leq 0$ such that $v - \psi$ attains a strict local maximum at t_0 . By adding appropriate linear functions to ψ , for any for any $\varepsilon > 0$ small, we can construct

$\psi_\varepsilon \in C^2((0, T))$ such that $v - \psi_\varepsilon$ attains a strict local maximum at some $t_0 \in (0, T)$ satisfying $\psi_\varepsilon''(t_0) \leq 0$ and

$$\psi_\varepsilon'(t_0) - \varepsilon \leq \lim_{t \rightarrow t_\varepsilon^+} \frac{v(t) - v(t_0)}{t - t_0} = -|\nabla^- u|(x_0), \quad (5.4)$$

where $x_0 = x(t_0) \in (\bar{x}, V_j)$. Adopting Definition 2.9 on the subsolution u , we deduce that

$$-\psi_\varepsilon'(t_0) \leq \Lambda_\infty v(t_0) = \Lambda_\infty u(x_0).$$

It thus follows from (5.4) that

$$|\nabla^- u|(x_0) \leq \Lambda_\infty u(x_0) + \varepsilon. \quad (5.5)$$

Passing to the limit as $\varepsilon \rightarrow 0$, we obtain

$$|\nabla^- u|(x_0) \leq \Lambda_\infty u(x_0). \quad (5.6)$$

On the other hand, since $v(t)$ is decreasing and concave on $(0, T)$, we can take $t_* \in (0, t_0)$ such that $v(t_*) > v(t_0)$, v is differentiable at t_* and

$$|\nabla^- u|(x_*) = -v'(t_*) \leq |\nabla^- u|(x_0),$$

where $x_* = x(t_*) \in (\bar{x}, x_0)$. We then can use (5.6) to get

$$|\nabla^- u|(x_*) \leq \Lambda_\infty u(x_0) < \Lambda_\infty u(x_*),$$

which contradicts the assumption that u is a supersolution of (1.3).

(iii) Let $x_0 \in \Omega$ be a point such that

$$|\nabla^- u|(x_0) > |\nabla^+ u|(x_0) \quad (5.7)$$

holds. If x_0 is an interior point of some edge, then we may apply an argument analogous to the proof of (ii). In fact, we can still parametrize the edge by the parameter t and find a C^2 function ψ such that $v - \psi$ attains a maximum at $t = t_0$ corresponding to x_0 . Using the modified test function ψ_ε and applying Definition 2.9, we obtain (5.6) again, which yields (5.1) at $x = x_0$.

When $x_0 \in \mathcal{V}$, our proof is similar but the test function looks slightly different. In view of Lemma 2.3, for any $r > 0$ small we can find x_r on an edge incident to x_0 such that (2.7) holds. By (ii) and the condition (5.7), it is not difficult to see that for any $\varepsilon > 0$ small $u - \phi$ attains a strict local maximum at x_0 , where

$$\phi_\varepsilon(x) = u(x_0) + (|\nabla^- u|(x_0) - \varepsilon)(d(x_r, x) - d(x_r, x_0)).$$

Letting $C > 0$ sufficiently large and

$$g(x) = \begin{cases} \phi_\varepsilon(x_r) & \text{for } x = x_r, \\ C & \text{for } x \in \partial B_r(x_0) \setminus \{x_r\}, \end{cases}$$

we have, for all $x \in B_r(x_0)$,

$$\phi_\varepsilon(x) = \min_{\partial y \in \partial B_r(x_0)} \{g(y) + (|\nabla^- u|(x_0) - \varepsilon)d(x, y)\}.$$

It then follows from Proposition 3.2 that ϕ_ε is ∞ -superharmonic in $B_r(x_0)$. Applying the definition of subsolutions, we are again led to (5.5) and consequently (5.6). The desired relation (5.1) at $x = x_0$ thus follows immediately.

(iv) Suppose that $|\nabla^+ u|(x_0) > 0$ at some $x_0 \in \Omega$. Let us show that (5.2) holds at $x = x_0$. Assume by contradiction that (5.1) holds instead. Since u is piecewise linear in Ω as explained in Remark 5.4, we can take a point x_* near x_0 on an incident edge satisfying

$$|\nabla^- u|(x_*) = |\nabla^+ u|(x_0).$$

It is clear that $u(x_*) > u(x_0)$ and therefore

$$|\nabla^- u|(x_*) \leq \Lambda_\infty u(x_0) < \Lambda_\infty u(x_*),$$

which obviously contradicts the fact that u is a supersolution of (1.3). We therefore have (5.2). The relation (5.3) is an immediate consequence of (5.2) and (iii). \square

5.3. Eigenvalue and eigenfunctions on a tripod. Let us examine the solution u_∞ of (1.3) on a tripod, a simple special case of metric graphs. We will later prove that u_∞ is in fact the unique solution of (1.3) satisfying (1.4) and (1.16), as stated in Theorem 1.3.

Example 5.6. Suppose that \mathbf{X} is a tripod, i.e., $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ is a finite graph with $\mathcal{V} = \{O, V_1, V_2, V_3\}$ and $\mathcal{E} = \{e_1, e_2, e_3\}$ such that $e_j = [O, V_j]$ for $j = 1, 2, 3$. Suppose that e_j has length ℓ_j and is parametrized by its length starting from the junction O for each $j = 1, 2, 3$. Let Ω and $\partial\Omega$ be respectively the interior and boundary of (\mathbf{X}, d) , that is, $\Omega = \mathbf{X} \setminus \{V_1, V_2, V_3\}$ and $\partial\Omega = \{V_1, V_2, V_3\}$.

Let us discuss three different cases in detail.

Case 1 ($\ell_1 = \ell_2 \leq \ell_3$). This is the case when

$$R_\infty = \frac{\ell_1 + \ell_3}{2} = \frac{\ell_2 + \ell_3}{2}, \quad (5.8)$$

$\Lambda_\infty = 1/R_\infty$ and the maximizer (incenter) set $\mathcal{M}(\Omega)$ is a singleton. The unique incenter lies on the edge e_3 and u_∞ coincides with u_{dist} given in (1.14), i.e.,

$$u_\infty(x) = \begin{cases} -\Lambda_\infty(t - \ell_1) & \text{for } x \text{ on either } e_1 \text{ or } e_2, \\ -\Lambda_\infty \left| t - \frac{\ell_3 - \ell_1}{2} \right| + 1 & \text{for } x \text{ on } e_3. \end{cases}$$

In the expression above, we used the parametrization of e_i by $t \in [0, t_i]$ ($i = 1, 2, 3$) with $t = 0$ corresponding to the junction point.

Case 2 ($\ell_1 = \ell_2 > \ell_3$). Then the maximum inscribed radius R_∞ is the same as that in (5.8) and $\Lambda_\infty = 1/R_\infty$, but there are two different incenter points, symmetric about the junction O , respectively on e_1 and e_2 . Let us denote them by P_1 and P_2 . The function u_∞ is again consistent with u_{dist} in (1.14). More precisely, we can compute it as

$$u_\infty(x) = \begin{cases} -\Lambda_\infty \left| t - \frac{\ell_3 - \ell_1}{2} \right| + 1 & \text{for } x \text{ on either } e_1 \text{ or } e_2, \\ -\Lambda_\infty(t - \ell_3) & \text{for } x \text{ on } e_3. \end{cases}$$

One can also discuss u_∞^Y in (1.22), the infimum of supersolutions with $Y = \{P_1\}$ (or symmetrically $Y = \{P_2\}$) in the partial constraint (1.21). We can show that $u_\infty^Y = u_\infty$ if $Y = \{P_1\}$. Indeed, assuming that the unique maximizer of u_∞^Y on e_2 appears at $t = \tau \in [0, \ell_2]$ and the slope in $[0, \tau]$ equals to κ , we can use the supersolution property of the eikonal part of (1.3) near $t = \tau$ to obtain the following two inequalities

$$\begin{aligned} \Lambda_\infty \left(\frac{\ell_3}{R_\infty} + \kappa\tau \right) &\leq \kappa, \\ \Lambda_\infty \left(\frac{\ell_3}{R_\infty} + \kappa\tau \right) &\leq \frac{1}{\ell_2 - \tau} \left(\frac{\ell_3}{R_\infty} + \kappa\tau \right). \end{aligned}$$

It follows that

$$R_\infty - \tau \geq \frac{\ell_3}{\kappa R_\infty}, \quad R_\infty + \tau \geq \ell_2. \quad (5.9)$$

We therefore deduce that

$$\ell_2 + \ell_3 = 2R_\infty \geq \ell_2 + \frac{\ell_3}{\kappa R_\infty},$$

which yields $\kappa R_\infty \geq 1$, i.e., $\kappa \geq \Lambda_\infty$. Since by definition $u_\infty^Y \leq u_\infty$, we get $\kappa = \Lambda_\infty$. In view of (5.9), we have

$$\tau = (\ell_2 - \ell_3)/2,$$

which amounts to saying that the unique maximizer of u_∞^Y on e_3 coincides with P_2 . This implies that $u_\infty^Y = u_\infty$. The same relation also holds when $Y = \{P_2\}$.

Case 3 ($\ell_1 < \ell_2 < \ell_3$). In this case,

$$R_\infty = \frac{\ell_1 + \ell_3}{2}$$

and the unique incenter lies on e_3 . Hence, $\Lambda_\infty = 1/R_\infty = 2/(\ell_1 + \ell_3)$. To investigate u_∞ , we need to further discuss two distinct cases.

When $\ell_2 \leq R_\infty$, we have

$$u_\infty(x) = -\Lambda_\infty(t - \ell_1) \quad \text{for } x \text{ on } e_1, \quad (5.10)$$

$$u_\infty(x) = -\frac{\Lambda_\infty \ell_1}{\ell_2}(t - \ell_2) \quad \text{for } x \text{ on } e_2, \quad (5.11)$$

$$u_\infty(x) = -\Lambda_\infty \left| t - \frac{\ell_3 - \ell_1}{2} \right| + 1 \quad \text{for } x \text{ on } e_3. \quad (5.12)$$

One can verify that this is really a positive solution of (1.3). One can also see that $u_\infty(x)$ is not lower semicontinuous at $x = O$. Indeed, $|\nabla^- u|(O) = \Lambda_\infty$ but $|\nabla^- u|(x) = \Lambda \ell_1 / \ell_2 < \Lambda_\infty$ for any $x \in (O, V_2)$.

When $\ell_2 > R_\infty$, the function u_∞ given by (5.10)–(5.12) is no longer a supersolution; indeed, at any point $x \in (O, V_2)$ sufficiently close to the junction, (5.11) implies that

$$\Lambda_\infty u_\infty(x) > \frac{\Lambda_\infty \ell_1}{\ell_2} = |\nabla^- u|(x).$$

In this case, we actually still get (5.10) and (5.12) but on e_2 instead of (5.11) we now have

$$u_\infty(x) = -\frac{\ell_1 \Lambda_\infty}{\ell_1 + \ell_3 - \ell_2} |t - \ell_2 + R_\infty| + \frac{\ell_1}{\ell_1 + \ell_3 - \ell_2}.$$

In either case, we can see that u_∞ satisfies the definition of subsolutions in Definition 2.9 and therefore it is a solution.

Let us make a brief remark on the concavity of ∞ -eigenfunction. As pointed out in [30], u_∞ is log-concave in a bounded convex domain $\Omega \subset \mathbb{R}^n$. This can be proved by passing to the limit as $p \rightarrow \infty$ of the log-concavity of the p -eigenfunctions. The log-concavity results for p -Laplacians are due to [45]. In order to obtain analogous results for the general eigenvalue problem in metric spaces, it seems crucial to introduce an appropriate notion of log-concave functions and a replacement assumption for the convexity of the domain.

In general, one cannot expect the so-called geodesic log-concavity to hold without additional assumptions on the geometry of Ω . Note that in Case 2 of the example above, u_∞ restricted to the geodesic $e_1 \cup e_2$ is not log-concave. The function value along this geodesic joining V_1 and V_2 yields an M-shaped graph. Also, in Case 3 when $\ell_2 > R_\infty$, it is easily seen that u_∞ is not log-concave along the geodesic $e_2 \cup e_3$.

Below we include a simple example of metric graphs containing loops. It is a variant of the tripod example above with e_3 changed to a loop.

Example 5.7. Let $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite graph with $\mathcal{V} = \{O, V_1, V_2\}$ and $\mathcal{E} = \{e_1, e_2, e_3\}$, where $e_j = [O, V_j]$ for $j = 1, 2$ and e_3 represents a loop on the vertex O . Again, for $j = 1, 2, 3$, we denote by ℓ_j the length of e_j and parametrize e_j by its length t from O . Let $\Omega = \mathbf{X} \setminus \{V_1, V_2\}$ and therefore $\partial\Omega = \{V_1, V_2\}$.

Assume that $\ell_1 \leq \ell_2$. We divide our discussion into several cases in terms of the length of the loop.

Case 1 ($\ell_3 < \ell_2 - \ell_1$). We have

$$R_\infty = \frac{\ell_1 + \ell_2}{2}, \quad \Lambda_\infty = \frac{1}{R_\infty} = \frac{2}{\ell_1 + \ell_2}.$$

The unique incenter lies on e_2 and the solution u_∞ of (1.3) is

$$u_\infty(x) = \begin{cases} -\Lambda_\infty(t - \ell_1) & \text{for } x \text{ on } e_1, \\ -\Lambda_\infty \left| t - \frac{\ell_2 - \ell_1}{2} \right| + 1 & \text{for } x \text{ on } e_2, \\ -\frac{4\ell_1}{(\ell_1 + \ell_2)(\ell_1 + \ell_2 - \ell_3)} \left| t - \frac{\ell_3}{2} \right| + \frac{2\ell_1}{\ell_1 + \ell_2 - \ell_3} & \text{for } x \text{ on } e_3. \end{cases} \quad (5.13)$$

Note that the value of u_∞ on e_3 does not depend on the orientation of its arc length parametrization.

Case 2 ($\ell_3 \geq \ell_2 - \ell_1$). In this case, it is easily seen that

$$R_\infty = \ell_1 + \frac{\ell_3}{2}, \quad \Lambda_\infty = \frac{1}{R_\infty} = \frac{2}{2\ell_1 + \ell_3}.$$

Also, u_∞ takes the same value on e_1 as in (5.13). On the loop e_3 , we have

$$u_\infty(x) = -\Lambda_\infty \left| t - \frac{\ell_3}{2} \right| + 1 \quad \text{for } x \text{ on } e_3.$$

To determine u_∞ on e_2 , we need to look into two subcases. If $\ell_3 \geq 2(\ell_2 - \ell_1)$, then we get

$$u_\infty(x) = \Lambda_\infty \ell_1 - \frac{\Lambda_\infty \ell_1}{\ell_2} t \quad \text{for } x \text{ on } e_2.$$

On the other hand, if $\ell_2 - \ell_1 \leq \ell_3 < 2(\ell_2 - \ell_1)$, then we have, for x on e_2 ,

$$u_\infty(x) = \frac{\ell_1}{2\ell_1 + \ell_3 - \ell_2} - \frac{2\ell_1}{(2\ell_1 + \ell_3)(2\ell_1 + \ell_3 - \ell_2)} \left| t + \ell_1 - \ell_2 + \frac{\ell_3}{2} \right|.$$

5.4. Non-uniqueness under partial incenter constraints. In the Euclidean space an example is built [27] in a dumbbell-shaped domain showing that in general there may be multiple linearly independent solutions to (1.3) and (1.4). In our general setting, this observation corresponds to the existence of solutions under partial incenter constraints as described in Remark 3.8.

In a similar manner to [27], for some particular domain Ω one can obtain at least one more solution to (1.3) if the condition (1.16) is weakened in the definition of u_∞ . We below present, on a metric graph, an analogue of the example in [27] for non-uniqueness of solutions.

Example 5.8. Let $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite graph with $\mathcal{V} = \{O\} \cup \{V_j\}_{j=0,\pm 1,\pm 2,\pm 3}$ and $\mathcal{E} = \{e_j\}_{j=0,\pm 1,\pm 2,\pm 3}$ satisfying

$$e_j = \begin{cases} [O, V_j] & \text{for } j = 0, \pm 1, \\ [V_{+1}, V_j] & \text{for } j = +2, +3, \\ [V_{-1}, V_j] & \text{for } j = -2, -3. \end{cases} \quad (5.14)$$

Then, equipped with the intrinsic metric d , (\mathbf{X}, d) is clearly a geodesic space. Let Ω be the interior of \mathbf{X} ; namely,

$$\Omega = \mathbf{X} \setminus \{V_j\}_{j=0,\pm 1,\pm 2,\pm 3}, \quad \partial\Omega = \{V_j\}_{j=0,\pm 1,\pm 2,\pm 3}.$$

Assume that the length ℓ_j of each edge e_j is

$$\ell_0 = \ell_{\pm 1} = \ell_{\pm 2} = 1, \quad \ell_{\pm 3} = 3.$$

We parametrize each e_j , using its length, by $[0, \ell_j]$ with $t = 0$ and $t = \ell_j$ respectively corresponding to the left and right endpoints in the expression (5.14).

One can show that

$$R_\infty = 2, \quad \Lambda_\infty = \frac{1}{2}$$

and there are two incenters P_\pm lying respectively on $e_{\pm 3}$ with $d(P_\pm, V_{\pm 3}) = 2$. We can also prove that u_∞ given below is a solution of (1.3):



$$u_\infty(x) = \begin{cases} -\frac{1}{4}t + \frac{1}{4} & \text{for } x \text{ on } e_0, \\ \frac{1}{4}t + \frac{1}{4} & \text{for } x \text{ on } e_{\pm 1}, \\ -\frac{1}{2}t + \frac{1}{2} & \text{for } x \text{ on } e_{\pm 2}, \\ -\frac{1}{2}|t - 1| + 1 & \text{for } x \text{ on } e_{\pm 3}, \end{cases} \quad (5.15)$$

where t represents the parameter for x on each e_j according to the parametrization given previously. See Figure 1 for an illustration of the function graph of u_∞ . Note that u_{dist} in (1.14) is not a solution and we have $u_\infty < u_{dist}$ on $e_{\pm 1}$.

On the other hand, if we take $Y = \{P_+\}$, then we can construct another solution u_∞^Y :

$$u_\infty^Y(x) = \begin{cases} u_\infty(x) & \text{for } x \text{ on } e_j \text{ with } j = 0, +1, +2, +3, \\ -\frac{1}{8}t + \frac{1}{4} & \text{for } x \text{ on } e_{-1}, \\ -\frac{1}{8}t + \frac{1}{8} & \text{for } x \text{ on } e_{-2}, \\ -\frac{1}{8}|t - 1| + \frac{1}{4} & \text{for } x \text{ on } e_{-3}. \end{cases}$$

See Figure 2 for the graph of u_∞^Y .

FIGURE 1. The graph of u_∞ FIGURE 2. The graph of u_∞^Y

Hence, in general we cannot expect uniqueness of solutions of (1.3) up to a constant multiple. In the Euclidean spaces, the ∞ -ground states (as the limits of p -eigenfunctions) are supposed to be symmetric in space. The function u_∞^Y above, which is not symmetric, thus corresponds to a non-variational solution on the metric graph.

It is not clear whether the solutions fulfilling the constraint (1.16) are unique in general geodesic spaces. In what follows, we attempt to provide partial results, tackling this problem on metric graphs.

5.5. Special curves and maximum slope. As explained in the introduction, we can construct ∞ -eigenfunctions in metric graphs by solving iteratively a free boundary problem associated to the eikonal equation. We actually build these solutions directly. To this end, we adopt the properties described in Proposition 5.2 to find some special curves for any solution u of (1.3).

The first result states that starting from any point in Ω we can find a curve passing through several vertices to reach a local maximizer of u .

Proposition 5.9 (Upstream curves). *Suppose that $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite connected graph equipped with intrinsic metric d . Let $\Omega \subsetneq \mathbf{X}$ be a domain with $\partial\Omega \subset \mathcal{V}$. Let u be a Lipschitz positive solution of (1.3). Then, from any $x_0 \in \Omega$, there exists a non-self-intersecting curve in Ω passing through $x_1, x_2, \dots, x_{m-1} \in \Omega \cap \mathcal{V}$ and $x_m \in \Omega$ with $m \geq 1$ such that*

$$|\nabla^+ u|(x_m) = 0 \quad (5.16)$$

and

$$|\nabla^- u|(x_0) \leq |\nabla^- u|(x_1) \leq \dots \leq |\nabla^- u|(x_m) = \Lambda_\infty u(x_m). \quad (5.17)$$

Proof. Suppose that

$$|\nabla^- u|(x_0) > \Lambda_\infty u(x_0)$$

to avoid the trivial case $m = 0$.

By Proposition 5.2(iii) and (iv), we then have

$$|\nabla^+ u|(x_0) = |\nabla^- u|(x_0) > 0.$$

Using Proposition 5.2(i) and (ii), we see that there exists an interval (x_0, x_*) on an edge such that

$$u(x(t)) - u(x_0) = |\nabla^+ u|(x_0)t \quad (5.18)$$

for $t > 0$ small, where $x(t)$ denotes the length parametrization of $[x_0, x_*]$ with $x(0) = x_0$. If u is not linear on the interval, then due to Remark 5.4 there exists a local maximizer $\hat{x} \in (x_0, x_*)$ of u such that

$$|\nabla^+ u|(x_0) = |\nabla^- u|(\hat{x}) = \Lambda_\infty u(\hat{x}).$$

We then take $x_1 = \hat{x}$. On the other hand, if (5.18) holds for the whole interval, then we take $x_1 = x_*$.

If $|\nabla^- u|(x_1) = \Lambda_\infty u(x_1)$, then we complete this process with $m = 1$. Otherwise, noticing that

$$|\nabla^+ u|(x_1) = |\nabla^- u|(x_1) \geq |\nabla^- u|(x_0),$$

we can continue to find $x_2 \in \Omega$ such that u is linear on $[x_1, x_2]$ with slope $|\nabla^- u|(x_1)$. Repeat this process finitely times until we find the final point x_m , where (5.16) holds. Then (5.17) follows by Proposition 5.2(iii). \square

It is clear that such finite sequences $\{x_1, x_2, \dots, x_m\}$ and the resulting curves are not unique in general. We call each of the curves an upstream curve from x_0 .

In contrast, we can find another type of curves, along which the function value is decreasing. We call them downstream curves.

Proposition 5.10 (Downstream curves). *Suppose that $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite connected metric graph equipped with intrinsic metric d . Let $\Omega \subsetneq \mathbf{X}$ be a domain with $\partial\Omega \subset \mathcal{V}$. Let u be a Lipschitz positive solution of (1.3). Then, from any $x_0 \in \Omega$, there exists a non-self-intersecting curve in $\overline{\Omega}$ passing through $y_1, y_2, \dots, y_{l-1} \in \Omega \cap \mathcal{V}$ and $y_l \in \partial\Omega$ with $l \geq 1$ such that*

$$|\nabla^- u|(x_0) \leq |\nabla^- u|(y_1) \leq \dots \leq |\nabla^- u|(y_{l-1}) \leq |\nabla^+ u|(y_l).$$

We omit the proof of this proposition, since it is a simple result of Definition 2.7 and the piecewise linearity of u as shown in Proposition 5.2. Note that, similar to the upstream curves, in general the sequences $\{y_1, y_2, \dots, y_l\}$ and the corresponding downstream curves are not unique either.

We next intend to present a direct constructive method to determine the value in $\Omega \setminus \Omega_\delta$ of a solution u of (1.3)–(1.4) satisfying (1.16), where Ω_δ is given as in (3.20) and $\delta > 0$ is chosen small. For this purpose, we make some preparations.

For any open subset $\mathcal{O} \subset \Omega$, let $d_{\mathcal{O}}$ denote the distance induced by geodesics restricted in $\overline{\mathcal{O}}$, that is, for any $x, y \in \overline{\mathcal{O}}$, we take

$$d_{\mathcal{O}}(x, y) = \min\{\ell(\gamma) : \gamma \text{ is a Lipschitz curve in } \overline{\mathcal{O}} \text{ joining } x \text{ and } y \text{ with } \gamma \setminus \{x, y\} \subset \mathcal{O}\}$$

Definition 5.11 (Peak points and downhill curves). Let \mathcal{O} be a subdomain of Ω . Let f be a Lipschitz function on $\overline{\Omega} \setminus \mathcal{O}$. A point $x \in \mathcal{O}$ (having n directions) is said to be a peak point in \mathcal{O} with respect to f if there exist $0 < K \leq \Lambda_\infty$ and a set of Lipschitz curves $\gamma_1, \gamma_2, \dots, \gamma_n$ in $\overline{\mathcal{O}}$, along each of n directions of x respectively, such that for any $j = 1, 2, \dots, n$, γ_j joins x to a point $z_j \in \partial\mathcal{O}$ satisfying $\gamma \setminus \{x, z_j\} \subset \Omega$,

$$\ell(\gamma_j) = d_{\mathcal{O}}(x, z_j) \leq R_\infty$$

and

$$KR_\infty = f(z_j) + K\ell(\gamma_j) = \min_{z \in \partial\mathcal{O}} \{f(z) + Kd_{\mathcal{O}}(x, z)\}. \quad (5.19)$$

We call $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ a set of downhill curves associated to the peak point $x \in \mathcal{O}$, f and slope K .

It is not difficult to see that there are only finitely many downhill curve sets for each peak point $x \in \mathcal{O}$ due to the finiteness of the number of distinct curves joining $x \in \mathcal{O}$ to $\partial\mathcal{O}$. For the same reason, there are only finitely many peak points in \mathcal{O} if they exist, which can be observed in the following way. Since the numbers of vertices and of edges are

finite, we only need to show that there are only finitely many peak points on each edge. Noticing that for any point $x \in \mathcal{O}$ in the interior of an edge $[V_i, V_j]$, there are only two directions of x toward V_i, V_j respectively, we see that the total number of peak points in (V_i, V_j) cannot exceed the number of distinct sets of curves from V_i and V_j to $\partial\mathcal{O}$.

Based on the finiteness shown above, we define the maximum slope at any peak point $x \in \Omega$. Suppose that x has n directions in $B_r(x)$ for $r > 0$ small. Then we define

$$A(x; f, \mathcal{O}) = \max \left\{ K \in (0, \Lambda_\infty] : \{\gamma_1, \dots, \gamma_n\} \text{ is a set of downhill curves} \right. \\ \left. \text{associated to } x \in \mathcal{O}, f \text{ and slope } K \right\}. \quad (5.20)$$

In what follows, the downhill curves $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ that attain the maximum slope will be called steepest downhill curves associated to $x \in \mathcal{O}$ and f . We denote by $\Gamma_A(x, f, \mathcal{O})$ the union of all steepest downhill curves associated to $x \in \mathcal{O}$ and f .

Let us now further maximize $A(x; f, \mathcal{O})$ over all peak points x . Set

$$A(f, \mathcal{O}) = \max\{A(x; f, \mathcal{O}) : x \text{ is a peak point in } \mathcal{O} \text{ with respect to } f\} \quad (5.21)$$

and

$$\mathcal{P}(f, \mathcal{O}) = \{x \in \mathcal{O} : x \text{ is a peak point in } \mathcal{O} \text{ w.r.t } f \text{ such that } A(x; f, \mathcal{O}) = A(f, \mathcal{O})\}.$$

Here we adopt the convention $A(f, \mathcal{O}) = -\infty$ if there are no peak points in \mathcal{O} . In this case, $\mathcal{P}(f, \mathcal{O}) = \emptyset$.

Also, we take the region consisting of all steepest downhill curves associated to the points in $\mathcal{P}(f, \mathcal{O})$ by setting

$$\Gamma_A(f, \mathcal{O}) = \bigcup \{\Gamma_A(x, f, \mathcal{O}) : x \in \mathcal{P}(f, \mathcal{O})\}.$$

By the definition above, for any Lipschitz function u in $\overline{\Omega}$ satisfying (1.4), we have $\mathcal{P}(u, \Omega) = \mathcal{M}(\Omega)$, i.e., any point $x \in \mathcal{M}(\Omega)$ is a peak point in $\mathcal{O} = \Omega$ with respect to u with $K = \Lambda_\infty$. In addition, $A(u, \Omega) = \Lambda_\infty$ holds.

Let us next introduce a boundary-point version of the maximizing curves by taking

$$B(f, \mathcal{O}) = \max \left\{ \frac{f(x) - f(z)}{\ell(\gamma_0)} : \gamma_0 \text{ is a Lipschitz curve in } \overline{\mathcal{O}} \text{ joining } x, z \in \partial\mathcal{O} \right. \\ \left. \text{satisfying } f(x) \geq f(z), \gamma_0 \setminus \{x, z\} \subset \mathcal{O}, \ell(\gamma_0) = d_{\mathcal{O}}(x, z) \right\}. \quad (5.22)$$

We call γ_0 a boundary-slope maximizing curve associated to f if γ_0 attains the maximum $B(f, \mathcal{O})$ above. Let $\Gamma_B(f, \mathcal{O})$ denote the union of all such curves.

Let us combine the interior and boundary cases and determine the overall maximum slope. We set

$$L(f, \mathcal{O}) = \max\{A(f, \mathcal{O}), B(f, \mathcal{O})\} \quad (5.23)$$

and

$$\Gamma(f, \mathcal{O}) = \begin{cases} \Gamma_A(f, \mathcal{O}) & \text{if } A(f, \mathcal{O}) > B(f, \mathcal{O}), \\ \Gamma_B(f, \mathcal{O}) & \text{if } A(f, \mathcal{O}) < B(f, \mathcal{O}), \\ \Gamma_A(f, \mathcal{O}) \cup \Gamma_B(f, \mathcal{O}) & \text{if } A(f, \mathcal{O}) = B(f, \mathcal{O}). \end{cases}$$

It is clear that $\Gamma(f, \mathcal{O}) \neq \emptyset$.

5.6. Uniqueness under full incenter constraints. Let us now directly construct a function U in a region containing $\Omega \setminus \Omega_\delta$, where $\delta > 0$ is taken small enough so that $\Omega \setminus \Omega_\delta$ contains no vertices. We shall show later that any solution coincides with U in $\Omega \setminus \Omega_\delta$.

We first assign the boundary condition $U = 0$ on $\partial\Omega$.

(1) Let $\mathcal{P}_0 = \mathcal{M}(\Omega)$. For any $x \in \mathcal{P}_0$ and each $\gamma_j \in \Gamma_A(x, U, \Omega)$, we set

$$U(y) = \Lambda_\infty d(y, z_j) \quad \text{for all } y \in \gamma_j.$$

Take

$$\mathcal{S}_0 = \Gamma(U, \Omega) = \Gamma_A(U, \Omega) = \bigcup_{x \in \mathcal{P}_0} \Gamma_A(x; U, \Omega).$$

In other words, \mathcal{S}_0 consists of all geodesics joining $x \in \mathcal{M}(\Omega)$ and $\partial\Omega$. It is clear that \mathcal{S}_0 is a closed subset of $\overline{\Omega}$.

(2) Suppose that, for $k = 1, 2, \dots$,

$$\overline{\Omega} \setminus \Omega_\delta \not\subset \mathcal{S}_{k-1}. \quad (5.24)$$

Let $\mathcal{O}_k = \Omega \setminus \mathcal{S}_{k-1}$ and $\mathcal{P}_k = \mathcal{P}(U, \mathcal{O}_k)$. For each γ in $\Gamma(U, \mathcal{O}_k)$, which is either from $\Gamma_A(U, \mathcal{O}_k)$ connecting $x \in \mathcal{P}_k$ and $z \in \partial\mathcal{O}$ or from $\Gamma_B(U, \mathcal{O}_k)$ connecting $x, z \in \partial\mathcal{O}$ with $U(x) \geq U(z)$, we set

$$U(y) = U(z) + L(U, \mathcal{O}_k) \ell(\gamma|_{[y,z]}) \quad \text{for all } y \in \gamma. \quad (5.25)$$

Here and in the sequel, $\gamma|_{[y,z]}$ denotes the closed portion of γ between y and z .

In addition, we take $\mathcal{S}_k = \mathcal{S}_{k-1} \cup \Gamma(U, \mathcal{O}_k)$, which is again a closed set.

(3) We stop repeating the above process after N times when $\overline{\Omega} \setminus \Omega_\delta \subset \mathcal{S}_N$.

Remark 5.12. We stress that U is well defined by (5.25). In general, the maximizing curves in $\Gamma(U, \mathcal{O}_k)$ may have nonempty intersections. In fact, when $\mathcal{P}_k \neq \emptyset$ and $L(U, \mathcal{O}_k) = A(U, \mathcal{O}_k)$, there are multiple steepest downhill curves joining at any $x \in \mathcal{P}_k \neq \emptyset$. Noticing that

$$\ell(\gamma|_{[y,z]}) = d_{\mathcal{O}_k}(y, z) = d_{\mathcal{O}_k}(x, z) - d_{\mathcal{O}_k}(x, y),$$

by (5.19) we can verify that, for any γ in $\Gamma_A(U, \mathcal{O}_k)$ with endpoint $z \in \partial\mathcal{O}_k$,

$$\begin{aligned} U(x) &= U(z) + L(U, \mathcal{O}_k) d_{\mathcal{O}_k}(x, z) \\ &= L(U, \mathcal{O}_k) (R_\infty - d_{\mathcal{O}_k}(x, z)) + L(U, \mathcal{O}_k) d_{\mathcal{O}_k}(x, z) = L(U, \mathcal{O}_k) R_\infty. \end{aligned}$$

In other words, U has the same value at any $x \in \mathcal{P}_k$.

In general, we can show that $U(y)$ given by (5.25) does not depend on the curve γ in $\Gamma(U, \mathcal{O}_k)$ that passes through y . Suppose that $\gamma, \tilde{\gamma}$ are two slope maximizing curves in $\Gamma(U, \mathcal{O}_k)$ intersecting at $y \in \overline{\mathcal{O}_k}$. Assume that $\gamma, \tilde{\gamma}$ respectively have endpoints $\{x, z\}$ and $\{\tilde{x}, \tilde{z}\}$ satisfying $U(x) \geq U(z)$ and $U(\tilde{x}) \geq U(\tilde{z})$. We claim that

$$U(z) + L(U, \mathcal{O}_k) \ell(\gamma|_{[y,z]}) = U(\tilde{z}) + L(U, \mathcal{O}_k) \ell(\tilde{\gamma}|_{[y,\tilde{z}]}) \quad (5.26)$$

To see this, let us build a curve γ_* by combining $\gamma|_{[x,y]}$ and $\tilde{\gamma}|_{[y,\tilde{z}]}$. Since γ is a slope maximizing curve, we can use (5.19) (if x is a peak point) or (5.22) (if $x \in \partial\mathcal{O}_k$) to obtain

$$U(z) + L(U, \mathcal{O}_k) \ell(\gamma) = U(x) \leq U(\tilde{z}) + L(U, \mathcal{O}_k) \ell(\gamma_*),$$

which implies that

$$U(z) + L(U, \mathcal{O}_k) \ell(\gamma|_{[y,z]}) \leq U(\tilde{z}) + L(U, \mathcal{O}_k) \ell(\tilde{\gamma}|_{[y,\tilde{z}]})$$

Interchanging the roles of γ and $\tilde{\gamma}$, we can use the same argument to deduce the reverse inequality and conclude the proof of (5.26).

Remark 5.13. It is worth pointing out that the process above is completed in finite steps. For each $k = 1, 2, \dots$, at least one curve in $\overline{\mathcal{O}}_k$ joining two different boundary points is contained in $\Gamma(U, \mathcal{O}_k)$, which means that \mathcal{O}_{k+1} has fewer edges than \mathcal{O}_k .

It turns out that any solution of (1.3) satisfying (1.4) and (1.16) agrees with the construction above in the δ -neighborhood of the boundary.

Theorem 5.14 (Uniqueness near boundary). *Suppose that $\mathbf{X} = (\mathcal{V}, \mathcal{E})$ be a finite connected one dimensional graph equipped with intrinsic metric d . Let $\Omega \subsetneq \mathbf{X}$ be a domain with $\partial\Omega \subset \mathcal{V}$. Choose $\delta > 0$ small such that $\Omega \setminus \Omega_\delta$ contains no vertices, where Ω_δ is given by (3.20). Let u be a Lipschitz positive solution of (1.3) satisfying (1.4) and (1.16). Then the value of u in $\Omega \setminus \Omega_\delta$ coincides with U defined in the construction process (1)–(3) above. In particular, the Lipschitz solutions of (1.3) satisfying (1.4) and (1.16) are unique in $\Omega \setminus \Omega_\delta$.*

Proof. Let us first show $u = U$ in Γ_0 . It suffices to prove that $|\nabla^- u| = \Lambda_\infty$ in Γ_0 . Suppose that it fails to hold. Then, in view of the Lipschitz continuity of u , the boundary condition (1.4) and the fact that $u = 1$ on $\mathcal{P}_0 = \mathcal{M}(\Omega)$, we can find a point x_0 on a certain geodesic of length R_∞ joining \mathcal{P}_0 and $\partial\Omega$ such that $|\nabla^- u|(x_0) > \Lambda_\infty$. Using Proposition 5.9, we can get a finite sequence of points with a final point $\eta_0 \in \Omega$ satisfying

$$|\nabla^- u|(x_0) \leq |\nabla^- u|(\eta_0) = \Lambda_\infty u(\eta_0).$$

which implies that $u(\eta_0) > 1$. This is clearly a contradiction to (1.16).

Assuming that $\overline{\Omega} \setminus \Omega_\delta \not\subset \mathcal{S}_0$, we next show $u = U$ on \mathcal{S}_1 . In view of (5.25), it suffices to show that for any $\gamma \in \Gamma(U, \mathcal{O}_1)$, there holds

$$u(y) = u(z) + L(U, \mathcal{O}_1)\ell(\gamma|_{[y,z]}),$$

where $z \in \partial\mathcal{O}_1$ is connected by γ to a peak point $x \in \mathcal{P}_1$ or a boundary point $x \in \partial\mathcal{O}_1$ satisfying $U(x) \geq U(z)$.

Since u is piecewise linear, we only need to show that

$$|\nabla^- u| = L(U, \mathcal{O}_1) \quad \text{on } \gamma. \tag{5.27}$$

Assume by contradiction that it is not the case. Then, due to the piecewise linearity of u on γ , there must exist $\xi_1 \in \gamma$ such that

$$|\nabla^- u|(\xi_1) = \max_{x \in \gamma} |\nabla^- u| > L(U, \mathcal{O}_1).$$

By Proposition 5.9, we can follow an upstream curve to reach a final point $\eta_1 \in \Omega$, where

$$|\nabla^- u|(\eta_1) = \Lambda_\infty u(\eta_1) \geq |\nabla^- u|(\xi_1) > L(U, \mathcal{O}_1).$$

Let γ^1 denote an upstream curve in Ω joining ξ_1 and η_1 .

Case 1. If $\gamma^1 \cap \mathcal{S}_0 \neq \emptyset$, then there exists $x_* \in \mathcal{S}_0 \cap \gamma_0$ such that we can construct a new curve $\tilde{\gamma}$ by combining the piece of γ from y to ξ_1 and the piece of γ^1 from ξ_1 to x_* such that $\tilde{\gamma} \setminus \{x_*, y\} \subset \mathcal{O}_1$. We easily see that

$$u(x_*) - u(y) \geq |\nabla^- u|(\xi_1)\ell(\tilde{\gamma}) > L(U, \mathcal{O}_1)\ell(\tilde{\gamma}).$$

This is a contradiction, since $L(U, \mathcal{O}_1) \geq B(U, \mathcal{O}_1)$ and $B(U, \mathcal{O}_1)$, as defined in (5.22), maximizes the slope of boundary values of U on $\partial\mathcal{O}_1$.

Case 2. If $\gamma^1 \cap \mathcal{S}_0 = \emptyset$, then η_1 is a local maximizer of u . By Remark 5.3, we have

$$|\nabla^- u|(x) = |\nabla^- u|(\eta_1) = \Lambda_\infty u(\eta_1)$$

for any $x \in \Omega \setminus \gamma^1$ close to η_1 . Hence, using Proposition 5.10, we can find downstream curves starting from η_1 along all directions, reaching $\partial\mathcal{S}_0 \cup \partial\Omega$, and satisfying $|\nabla^- u| \geq |\nabla^- u|(\eta_1)$ along any such curve.

We further claim that $|\nabla^- u| = |\nabla^- u|(\eta_1)$ on these curves. Suppose by contradiction that $|\nabla^- u|(\xi_2) > |\nabla^- u|(\eta_1)$ at some point ξ_2 on one of these curves. We then follow another upstream curve γ^2 from ξ_2 to reach a final point $\eta_2 \in \Omega$. If $\gamma^2 \cap \mathcal{S}_0 \neq \emptyset$, we can reduce the situation to Case 1 with γ^1 replaced by γ^2 and end up with a contradiction again. If $\gamma^2 \cap \mathcal{S}_0 = \emptyset$, then we repeat the process from the beginning of Case 2 with η_1 replaced by η_2 .

Noticing that there are only finitely many local maximizers of u , as stated in Corollary 5.5, if no contradictions as in Case 1 are obtained to stop the argument, we can repeat the process finitely many times to finally find $\eta \in \mathcal{O}_1 = \Omega \setminus \mathcal{S}_0$, a local maximizer of u , and $K > L(U, \mathcal{O}_1)$ such that $|\nabla^- u| = K$ along any curve connecting η and $\partial\mathcal{O}_1$. This amounts to saying that $\eta \in \mathcal{P}(U, \mathcal{O}_1)$ and $A(U, \mathcal{O}_1) \geq A(\eta, U, \mathcal{O}_1) \geq K$; we recall the definition of maximum slopes in (5.20) and (5.21). This however contradicts the maximality of $L(U, \mathcal{O}_1)$, as defined in (5.23). We therefore obtain (5.27).

It follows immediately from (5.27) that (5.25) holds for $k = 1$. Obtaining the value of u in \mathcal{S}_1 , we then continue our discussion provided that (5.24) holds with $k = 2$. We can conclude the proof by iterating the same argument finitely many times. \square

The proof of Theorem 1.3 can be completed now.

Proof of Theorem 1.3. We have shown in Theorem 1.2 that in general proper geodesic spaces u_∞ is a solution of (1.3) satisfying (1.4) and (1.16). The uniqueness of such solutions on a finite metric graph follows from Theorem 5.14 and Corollary 3.10. \square

APPENDIX A. EIKONAL EQUATION IN METRIC SPACES

In this appendix, we present several results on the eikonal equation in metric spaces that are used in the proof of Theorem 3.4.

Let (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Consider the eikonal equation

$$|\nabla u| = \lambda \quad \text{in } \Omega \tag{A.1}$$

with the Dirichlet boundary condition

$$u = g \quad \text{on } \partial\Omega$$

where $\lambda > 0$ and $g \in C(\partial\Omega)$ are given.

In [38], the notion of Monge solutions of the Hamilton-Jacobi equations in the Euclidean space [41, 11] is also generalized for general length spaces. A locally Lipschitz function in Ω is called a Monge solution (resp., Monge supersolution, Monge subsolution) of (A.1) if $|\nabla^- u| = \lambda$ (resp., $\geq \lambda$, $\leq \lambda$) in Ω .

An optimal control interpretation is provided in [25, Theorem 4.2] to construct solutions of general eikonal equations in metric spaces. It is shown in [38] that such solutions are actually Monge solutions. For our particular purpose in this work, we below build a slightly different Monge solution u satisfying $u \leq g$ on $\partial\Omega$, which is simply the celebrated McShane-Whitney extension in $\overline{\Omega}$ with Lipschitz constant λ .

Theorem A.1 (Construction of a Monge solution). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Assume that $\lambda > 0$ and $g \in C(\partial\Omega)$. Let u be given by (3.1). Then u is Lipschitz in $\overline{\Omega}$ and $|\nabla^- u| = |\nabla u| = \lambda$ holds in Ω .*

Proof. We first claim that, for any subdomain \mathcal{O} with $\overline{\mathcal{O}} \subset \Omega$ and any $x \in \mathcal{O}$,

$$u(x) = \min_{z \in \overline{\mathcal{O}}} \{u(z) + \lambda d(x, z)\}. \quad (\text{A.2})$$

For any $z \in \overline{\mathcal{O}}$, by (3.1) there exists $y_z \in \partial\Omega$ such that

$$u(z) \geq g(y_z) + \lambda d(z, y_z). \quad (\text{A.3})$$

It follows from (3.1) again that

$$u(x) \leq g(y_z) + \lambda d(x, y_z) \leq g(y_z) + \lambda d(z, y_z) + \lambda d(x, z) \leq u(z) + \lambda d(x, z), \quad (\text{A.4})$$

which, due to the arbitrariness of $z \in \overline{\mathcal{O}}$ yields

$$u(x) \leq \min_{z \in \overline{\mathcal{O}}} \{u(z) + \lambda d(x, z)\}.$$

On the other hand, we can find $y_x \in \partial\Omega$ such that

$$u(x) \geq g(y_x) + \lambda d(x, y_x). \quad (\text{A.5})$$

Take a geodesic γ connecting x and y_x , i.e., $\gamma(0) = x$, $\gamma(1) = y_x$ and $\ell(\gamma) = d(x, y_x)$. There must exist a point of intersection z_x of γ and $\partial\mathcal{O}$, which satisfies

$$d(x, y_x) = d(x, z_x) + d(z_x, y_x).$$

In view of (A.5), we thus can apply (3.1) once again to get

$$u(x) \geq g(y_x) + \lambda d(z_x, y_x) + \lambda d(x, z_x) \geq u(z_x) + \lambda d(x, z_x) \geq \min_{z \in \overline{\mathcal{O}}} \{u(z) + \lambda d(x, z)\}.$$

Our proof of (A.2) is now complete.

As mentioned before, (3.1) is just the McShane-Whitney Lipschitz extension. We can certainly obtain the Lipschitz regularity:

$$|u(x) - u(z)| \leq \lambda d(x, z) \quad \text{for any } x, z \in \overline{\Omega}. \quad (\text{A.6})$$

In fact, the argument resulting in (A.4) applies to all points $x, z \in \overline{\Omega}$. Even if x or z appears on $\partial\Omega$, we can still get $y_z \in \partial\Omega$ satisfying (A.3) and thus obtain (A.4). Interchanging the roles of x and z in $\overline{\Omega}$, we get (A.6) immediately.

It is then clear that $|\nabla u| \leq \lambda$ in Ω . Let us now take any $x \in \Omega$ and any $r > 0$ small such that $B_r(x) \subset \Omega$. Using (A.2) with $\mathcal{O} = B_r(x)$ we have

$$\sup_{z \in B_r(x)} (u(x) - u(z)) \geq \lambda d(x, z),$$

which implies that

$$\sup_{z \in B_r(x) \setminus \{x\}} \frac{u(x) - u(z)}{d(x, z)} \geq \lambda.$$

Passing to the limit as $r \rightarrow 0$, we end up with

$$|\nabla^- u|(x) \geq \lambda$$

for any $x \in \Omega$. Since $|\nabla^- u| \leq |\nabla u|$, we obtain $|\nabla^- u| = |\nabla u| = \lambda$ in Ω , as desired. \square

The function u defined by (3.1) obviously satisfies $u \leq g$ on $\partial\Omega$. Note that in general one cannot expect the $u = g$ holds on $\partial\Omega$ even in the Euclidean space. A simple example is as follows. Let \mathbf{X} be the closed interval $[0, 1]$ equipped with the standard Euclidean metric and $\Omega = (0, 1)$. Assume that $0 < \lambda < 1$, $g(0) = 0$ and $g(1) = 1$. Then the function u as in (3.1) can be directly computed:

$$u(x) = \lambda x, \quad x \in \overline{\Omega} = \mathbf{X} = [0, 1].$$

In particular, we have $u(1) = \lambda < g(1)$. In general, one needs additional assumptions to guarantee the Dirichlet boundary condition in general metric spaces such as the λ -Lipschitz continuity of g on $\partial\Omega$. See more details in [38, Section 3.3].

For our application in this work, we next present a comparison theorem for the Lipschitz Monge sub- and supersolutions in a proper geodesic space.

Theorem A.2 (Comparison principle for eikonal equation). *Suppose that (\mathbf{X}, d) is a proper geodesic space and $\Omega \subsetneq \mathbf{X}$ is a bounded domain. Let u and v be respectively a Monge subsolution and a Monge supersolution of (A.1) with $\lambda > 0$. Assume in addition that u and v are continuous in $\overline{\Omega}$. If $u \leq v$ on $\partial\Omega$, then $u \leq v$ in $\overline{\Omega}$.*

Proof. It is clear that u and v are bounded, since $u, v \in C(\overline{\Omega})$ and Ω is bounded. We therefore may assume that $u, v \geq 0$ by adding a positive constant to them. It suffices to show that $\mu u \leq v$ in Ω for all $\mu \in (0, 1)$.

Assume by contradiction that there exists $\mu \in (0, 1)$ such that $\sup_{\Omega}(\mu u - v) > 0$. Due to the assumption that $u \leq v$ on $\partial\Omega$, we can find $x \in \Omega$ such that

$$\sup_{\Omega}(\mu u - v) = (\mu u - v)(x). \quad (\text{A.7})$$

Since v is a Monge supersolution of (A.1), there exists a sequence $\{y_n\} \subset \Omega$ converging to x as $n \rightarrow \infty$ such that

$$\lim_{n \rightarrow \infty} \frac{[v(y_n) - v(x)]_-}{d(x, y_n)} \geq \lambda > 0.$$

Hence, by (A.7) we have

$$\lambda \leq \limsup_{n \rightarrow \infty} \frac{\mu(u(x) - u(y_n))}{d(x, y_n)} \leq \limsup_{y \rightarrow x} \frac{\mu(u(x) - u(y))}{d(x, y)} \leq \mu |\nabla^- u|(x).$$

Noticing that u is a Monge subsolution, we thus get

$$\lambda \leq \mu \lambda,$$

which is clearly a contradiction. \square

Similar comparison results also hold in the case of general length spaces; see [38, Theorem 4.2]. In the general case, the assumptions are slightly more complicated and the proof, involving Ekeland's variational principle is more technical due to the lack of local compactness of the metric space.

REFERENCES

- [1] L. Ambrosio and J. Feng. On a class of first order Hamilton-Jacobi equations in metric spaces. *J. Differential Equations*, 256(7):2194–2245, 2014.
- [2] L. Ambrosio and S. Honda. New stability results for sequences of metric measure spaces with uniform Ricci bounds from below. In *Measure theory in non-smooth spaces*, Partial Differ. Equ. Meas. Theory, pages 1–51. De Gruyter Open, Warsaw, 2017.

- [3] S. N. Armstrong and C. K. Smart. A finite difference approach to the infinity Laplace equation and tug-of-war games. *Trans. Amer. Math. Soc.*, 364(2):595–636, 2012.
- [4] G. Aronsson, M. G. Crandall, and P. Juutinen. A tour of the theory of absolutely minimizing functions. *Bull. Amer. Math. Soc. (N.S.)*, 41(4):439–505, 2004.
- [5] M. Belloni and B. Kawohl. The pseudo- p -Laplace eigenvalue problem and viscosity solutions as $p \rightarrow \infty$. *ESAIM Control Optim. Calc. Var.*, 10(1):28–52, 2004.
- [6] M. Belloni, B. Kawohl, and P. Juutinen. The p -Laplace eigenvalue problem as $p \rightarrow \infty$ in a Finsler metric. *J. Eur. Math. Soc. (JEMS)*, 8(1):123–138, 2006.
- [7] T. Bhattacharya. An elementary proof of the Harnack inequality for non-negative infinity-superharmonic functions. *Electron. J. Differential Equations*, pages No. 44, 8, 2001.
- [8] I. Birindelli and F. Demengel. First eigenvalue and maximum principle for fully nonlinear singular operators. *Adv. Differential Equations*, 11(1):91–119, 2006.
- [9] I. Birindelli and F. Demengel. Eigenvalue, maximum principle and regularity for fully non linear homogeneous operators. *Commun. Pure Appl. Anal.*, 6(2):335–366, 2007.
- [10] D. Bonheure, J. D. Rossi, and N. Saintier. The limit as $p \rightarrow \infty$ in the eigenvalue problem for a system of p -Laplacians. *Ann. Mat. Pura Appl. (4)*, 195(5):1771–1785, 2016.
- [11] A. Briani and A. Davini. Monge solutions for discontinuous Hamiltonians. *ESAIM Control Optim. Calc. Var.*, 11(2):229–251 (electronic), 2005.
- [12] D. Burago, Y. Burago, and S. Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [13] F. Charro and E. Parini. Limits as $p \rightarrow \infty$ of p -Laplacian eigenvalue problems perturbed with a concave or convex term. *Calc. Var. Partial Differential Equations*, 46(1-2):403–425, 2013.
- [14] F. Charro and I. Peral. Limit branch of solutions as $p \rightarrow \infty$ for a family of sub-diffusive problems related to the p -Laplacian. *Comm. Partial Differential Equations*, 32(10-12):1965–1981, 2007.
- [15] M. G. Crandall, L. C. Evans, and R. F. Gariepy. Optimal Lipschitz extensions and the infinity Laplacian. *Calc. Var. Partial Differential Equations*, 13(2):123–139, 2001.
- [16] M. G. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [17] G. Crasta and I. Fragalà. On the characterization of some classes of proximally smooth sets. *ESAIM Control Optim. Calc. Var.*, 22(3):710–727, 2016.
- [18] G. Crasta and I. Fragalà. Rigidity results for variational infinity ground states. *Indiana Univ. Math. J.*, 68(2):353–367, 2019.
- [19] J. V. da Silva, J. D. Rossi, and A. M. Salort. Maximal solutions for the ∞ -eigenvalue problem. *Adv. Calc. Var.*, 12(2):181–191, 2019.
- [20] L. Esposito, B. Kawohl, C. Nitsch, and C. Trombetti. The Neumann eigenvalue problem for the ∞ -Laplacian. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 26(2):119–134, 2015.
- [21] N. Fukagai, M. Ito, and K. Narukawa. Limit as $p \rightarrow \infty$ of p -Laplace eigenvalue problems and L^∞ -inequality of the Poincaré type. *Differential Integral Equations*, 12(2):183–206, 1999.
- [22] W. Gangbo and A. Świąch. Optimal transport and large number of particles. *Discrete Contin. Dyn. Syst.*, 34(4):1397–1441, 2014.
- [23] W. Gangbo and A. Świąch. Metric viscosity solutions of Hamilton-Jacobi equations depending on local slopes. *Calc. Var. Partial Differential Equations*, 54(1):1183–1218, 2015.
- [24] J. Garcia-Azorero, J. J. Manfredi, I. Peral, and J. D. Rossi. Partial differential equations—the limit as $p \rightarrow \infty$ for the p -Laplacian with mixed boundary conditions and the mass transport problem through a given window. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 20(2):111–126, 2009.
- [25] Y. Giga, N. Hamamuki, and A. Nakayasu. Eikonal equations in metric spaces. *Trans. Amer. Math. Soc.*, 367(1):49–66, 2015.
- [26] S. Honda. Cheeger constant, p -Laplacian, and Gromov-Hausdorff convergence. *preprint*, 2013.
- [27] R. Hynd, C. K. Smart, and Y. Yu. Nonuniqueness of infinity ground states. *Calc. Var. Partial Differential Equations*, 48(3-4):545–554, 2013.
- [28] P. Juutinen. Minimization problems for Lipschitz functions via viscosity solutions. *Ann. Acad. Sci. Fenn. Math. Diss.*, (115):53, 1998. Dissertation, University of Jyväskylä, Jyväskylä, 1998.
- [29] P. Juutinen and P. Lindqvist. On the higher eigenvalues for the ∞ -eigenvalue problem. *Calc. Var. Partial Differential Equations*, 23(2):169–192, 2005.
- [30] P. Juutinen, P. Lindqvist, and J. J. Manfredi. The ∞ -eigenvalue problem. *Arch. Ration. Mech. Anal.*, 148(2):89–105, 1999.
- [31] P. Juutinen and N. Shanmugalingam. Equivalence of AMLE, strong AMLE, and comparison with cones in metric measure spaces. *Math. Nachr.*, 279(9-10):1083–1098, 2006.

- [32] E. Lindgren and P. Lindqvist. Fractional eigenvalues. *Calc. Var. Partial Differential Equations*, 49(1-2):795–826, 2014.
- [33] E. Lindgren and P. Lindqvist. Infinity-harmonic potentials and their streamlines. *Discrete Contin. Dyn. Syst.*, 39(8):4731–4746, 2019.
- [34] E. Lindgren and P. Lindqvist. The gradient flow of infinity-harmonic potentials. *Adv. Math.*, 378:107526, 24, 2021.
- [35] E. Lindgren and P. Lindqvist. On ∞ -ground states in the plane. *preprint*, 2021.
- [36] P. Lindqvist. *Notes on the infinity Laplace equation*. SpringerBriefs in Mathematics. BCAM Basque Center for Applied Mathematics, Bilbao; Springer, [Cham], 2016.
- [37] P. Lindqvist and J. J. Manfredi. The Harnack inequality for ∞ -harmonic functions. *Electron. J. Differential Equations*, pages No. 04, approx. 5 pp. 1995.
- [38] Q. Liu, N. Shanmugalingam, and X. Zhou. Equivalence of solutions of eikonal equation in metric spaces. *J. Differential Equations*, 272:979–1014, 2021.
- [39] J. J. Manfredi, A. M. Oberman, and A. P. Sviridov. Nonlinear elliptic partial differential equations and p -harmonic functions on graphs. *Differential Integral Equations*, 28(1-2):79–102, 2015.
- [40] A. Mitsuishi. Certain min-max values related to the p -energy and packing radii of Riemannian manifolds and metric measure spaces. *preprint*, 2019.
- [41] R. T. Newcomb, II and J. Su. Eikonal equations with discontinuities. *Differential Integral Equations*, 8(8):1947–1960, 1995.
- [42] S. Patrizi. The principal eigenvalue of the ∞ -Laplacian with the Neumann boundary condition. *ESAIM Control Optim. Calc. Var.*, 17(2):575–601, 2011.
- [43] Y. Peres, O. Schramm, S. Sheffield, and D. B. Wilson. Tug-of-war and the infinity Laplacian. *J. Amer. Math. Soc.*, 22(1):167–210, 2009.
- [44] M. Pérez-Llanos and J. D. Rossi. The behaviour of the $p(x)$ -Laplacian eigenvalue problem as $p(x) \rightarrow \infty$. *J. Math. Anal. Appl.*, 363(2):502–511, 2010.
- [45] S. Sakaguchi. Concavity properties of solutions to some degenerate quasilinear elliptic Dirichlet problems. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 14(3):403–421 (1988), 1987.
- [46] A. P. Sviridov. *Elliptic equations in graphs via stochastic games*. ProQuest LLC, Ann Arbor, MI, 2011. Thesis (Ph.D.)—University of Pittsburgh.
- [47] Y. van Gennip, N. Guillen, B. Osting, and A. L. Bertozzi. Mean curvature, threshold dynamics, and phase field theory on finite graphs. *Milan J. Math.*, 82(1):3–65, 2014.
- [48] Y. Yu. Some properties of the ground states of the infinity Laplacian. *Indiana Univ. Math. J.*, 56(2):947–964, 2007.

(Qing Liu) DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCE, FUKUOKA UNIVERSITY, JAPAN

Email address: qingliu@fukuoka-u.ac.jp

(Ayato Mitsuishi) DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCE, FUKUOKA UNIVERSITY, JAPAN

Email address: mitsuishi@fukuoka-u.ac.jp