

Kaniadakis holographic dark energy and cosmology

Niki Drepanou^a Andreas Lymperis^b Emmanuel N. Saridakis^{c,d,e} Kuralay Yesmakhanova^{e,f}

^a*Department of Physics, National Technical University of Athens, Zografou Campus GR 157 73, Athens, Greece*

^b*Department of Physics, University of Patras, 26500 Patras, Greece*

^c*National Observatory of Athens, Lofos Nymfon, 11852 Athens, Greece*

^d*CAS Key Laboratory for Researches in Galaxies and Cosmology, Department of Astronomy, University of Science and Technology of China, Hefei, Anhui 230026, P.R. China*

^e*Eurasian National University, Nur-Sultan Astana 010008, Kazakhstan*

^f*Ratbay Myrzakulov Eurasian International Centre for Theoretical Physics, Nur-Sultan 010009, Kazakhstan*

E-mail: alymperis@upatras.gr, msaridak@phys.uoa.gr

ABSTRACT: We construct a holographic dark energy scenario based on Kaniadakis entropy, which is a generalization of Boltzmann-Gibbs entropy that arises from relativistic statistical theory and is characterized by a single parameter K which quantifies the deviations from standard expressions, and we use the future event horizon as the Infrared cutoff. We extract the differential equation that determines the evolution of the effective dark energy density parameter, and we provide analytical expressions for the corresponding equation-of-state and deceleration parameters. We show that the universe exhibits the standard thermal history, with the sequence of matter and dark-energy eras, while the transition to acceleration takes place at $z \approx 0.6$. Concerning the dark-energy equation-of-state parameter we show that it can have a rich behavior, being quintessence-like, phantom-like, or experience the phantom-divide crossing in the past or in the future. Finally, in the far future dark energy dominates completely, and the asymptotic value of its equation of state depends on the values of the two model parameters.

KEYWORDS: Holographic dark energy, Kaniadakis entropy, phantom crossing

Contents

1	Introduction	1
2	Kaniadakis holographic dark energy	2
3	Cosmological evolution	7
4	Conclusions	10

1 Introduction

It is now well established that the Universe at late times experienced the transition from the matter era to the accelerated expansion phase. Although the simplest explanation would be the consideration of the cosmological constant, the corresponding problem related to the quantum-field-theoretical calculation of its value, as well as the possibility of a dynamical nature, led to two main paths of constructing extended scenarios. The first is to maintain general relativity as the underlying theory of gravity, and consider new, exotic forms of matter that constitute the concept of dark energy [1–3]. The second is to construct extended or modified theories of gravity, that possess general relativity as a low-energy limit, but which in general provide the extra degrees of freedom that can drive the dynamical universe acceleration [4–7].

Nevertheless, one can acquire an alternative explanation of the dark energy origin, through the cosmological application [8–10] of the holographic principle [11–13]. The corresponding framework is based on the thermodynamics of black holes and the connection of the Ultraviolet cutoff of a quantum field theory (related to the vacuum energy), with the largest distance of the theory (which in turn is a requirement in order for the theory to be applicable at large distances) [14]. In particular, in a given system whose entropy is proportional to its volume, the total energy should not be larger than the mass of a black hole with the same size, whose entropy is proportional to its area, since in such a case the system would collapse to a black hole. If one considers the whole Universe as the system one extracts a vacuum energy of holographic origin, namely a form of holographic dark energy with dynamical nature [15, 16].

The cosmological implications of holographic dark energy proves to be very interesting [15–26] and it proves to be in agreement with observations [27–32]. In this scenario one is free of the naturalness problem of the cosmological constant [33], as well as from pathologies that may arise in various modified gravity constructions [7]. Additionally, it has been shown to be able to alleviate the H_0 and growth tensions between Λ CDM scenario and some direct measurements [34]. That is why a large amount of research has been devoted to these investigations, and the basic models have been extended in various ways [35–71].

The basic expression in the construction of holographic dark energy is the one that connects the entropy of a system with geometrical quantities such as its radius. The standard one is the Bekenstein-Hawking entropy, which arises as the black-hole and cosmological application of the standard Boltzmann-Gibbs entropy. However, Kaniadakis has proposed a one-parameter generalization of the Boltzmann-Gibbs entropy, called Kaniadakis entropy [72, 73]. This results from a self-consistent and coherent relativistic statistical theory, in which the basic features of standard statistical theory are maintained. In such an extended statistical theory, the distribution functions are a one-parameter continuous deformation of the usual Maxwell-Boltzmann ones, and hence standard statistical theory is recovered in a particular limit.

In the present work we will use Kaniadakis entropy in order to formulate Kaniadakis holographic dark energy, and study its cosmological implications. Although in the literature there were some first attempts towards this direction [74–76] the resulting models were not correct. The reason for this failure was the fact that the authors used the Hubble horizon instead of the future event horizon in the basic holographic expression. Therefore, not only the resulting models could not recover usual holographic dark energy in the limit where Kaniadakis entropy becomes standard entropy, as it should, but in order to be able to describe the universe evolution one needs unacceptably large values of the Kaniadakis parameter, namely unacceptably large deviations from standard entropy. Hence, in this work we proceed to the consistent formulation of Kaniadakis holographic dark energy, which is indeed a well-defined extension of standard holographic dark energy, recovering it as a particular limit in the case where Kaniadakis entropy becomes standard Bekenstein-Hawking entropy.

The plan of the manuscript is the following: In Section 2 we formulate Kaniadakis holographic dark energy, we present the corresponding cosmological equations and we extract analytical relations for the dark energy density and equation-of-state parameters. Then in Section 3 we proceed to the study of the resulting cosmological behavior. Finally, in Section 4 we discuss our results and we summarize.

2 Kaniadakis holographic dark energy

In this section we proceed to the formulation of Kaniadakis holographic dark energy. The basic idea behind holographic dark energy is the inequality $\rho_{DE} L^4 \leq S$, where L is the largest distance of the theory (the Infrared cutoff) and S the entropy relation applied in a black hole of radius L [15, 16]. In the case of standard Bekenstein-Hawking entropy $S_{BH} \propto A/(4G) = \pi L^2/G$, with G the Newton’s constant, the saturation of the above inequality gives standard holographic dark energy, namely $\rho_{DE} = 3c^2 M_p^2 L^{-2}$, with M_p the Planck mass and c the model parameter. Hence, we can see that if instead of standard entropy we use a modified one, we will obtain a modified holographic dark energy.

As we mentioned in the Introduction, Kaniadakis entropy is a one-parameter generalization of the classical entropy. It is given by [72, 73]

$$S_K = -k_B \sum_i n_i \ln_{\{K\}} n_i, \quad (2.1)$$

with k_B the Boltzmann constant, and where we have defined $\ln_{\{K\}} x = (x^K - x^{-K})/2K$. Kaniadakis entropy is characterized by the single dimensionless parameter K , which quantifies the deviation from the case of standard statistical mechanics. Hence, standard entropy is recovered in the limit $K \rightarrow 0$, while K can vary in the range $-1 < K < 1$. Additionally, in such a generalized statistical theory the distribution function reads as $n_i = \alpha \exp_{\{K\}}[-\beta(E_i - \mu)]$, where $\exp_{\{K\}}(x) = \left(\sqrt{1 + K^2 x^2} + Kx\right)^{1/K}$, $\alpha = [(1 - K)/(1 + K)]^{1/2K}$, $1/\beta = \sqrt{1 - K^2} k_B T$, and the chemical potential μ can be fixed by normalization [72, 73]. Kaniadakis entropy can be expressed as [77–82]

$$S_K = -k_B \sum_{i=1}^W \frac{P_i^{1+K} - P_i^{1-K}}{2K}, \quad (2.2)$$

where P_i is the probability the system to be in a specific microstate and W the total number of configurations.

Let us apply Kaniadakis entropy in the black-hole framework, which will then be needed for the holographic application. Assuming that $P_i = 1/W$, using the fact that Boltzmann-Gibbs entropy is $S \propto \ln(W)$, while the Bekenstein-Hawking entropy is given by $S_{BH} = A/(4G)$, we acquire $W = \exp[A/(4G)]$ [74], where from now on we impose units in which the Boltzmann constant, the light speed, and the reduced Planck constant are set to $k_B = c = \hbar = 1$. Hence, inserting these into (2.2) we find

$$S_K = \frac{1}{K} \sinh(K S_{BH}). \quad (2.3)$$

As expected in the limit $K \rightarrow 0$ one recovers standard Bekenstein-Hawking entropy, i.e. $S_{K \rightarrow 0} = S_{BH}$. Since in reality one expects the above modified entropy to be close to the standard Bekenstein-Hawking value, we expect that $K \ll 1$ (we remind that $-1 < K < 1$). Thus, it is justified to expand the above Kaniadakis entropy for small K , obtaining

$$S_K = S_{BH} + \frac{K^2}{6} S_{BH}^3 + \mathcal{O}(K^4). \quad (2.4)$$

As one can see, the first term is the usual entropy, while the second term is the lowest-order Kaniadakis correction.

It is now easy to extract the relation of Kaniadakis holographic dark energy. In particular, inserting (2.4) into the inequality $\rho_{DE} L^4 \leq S$, we obtain

$$\rho_{DE} = 3c^2 M_p^2 L^{-2} + 3\tilde{c}^2 K^2 M_p^6 L^2, \quad (2.5)$$

with c and \tilde{c} constants. As mentioned above, for $K = 0$ the above expression gives the usual holographic dark energy $\rho_{DE} = 3c^2 M_p^2 L^{-2}$. In the following we absorb the constant \tilde{c} inside the parameter K , by setting $3\tilde{c}^2 K^2 \equiv \tilde{K}^2$ and we drop the tildes for simplicity.

We proceed by considering a flat homogeneous and isotropic Friedmann-Robertson-Walker (FRW) geometry with metric

$$ds^2 = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j, \quad (2.6)$$

with $a(t)$ the scale factor. As a next step, in any holographic dark energy scenario, one needs to determine the length L that appears in the corresponding relations. In the case of standard holographic dark energy models it is well known that L cannot be the Hubble horizon H^{-1} (where $H \equiv \dot{a}/a$ is the Hubble function), since this choice leads to obvious inconsistencies [83], such as no acceleration. Thus, one must use the future event horizon [15]

$$R_h \equiv a \int_t^\infty \frac{dt}{a} = a \int_a^\infty \frac{da}{Ha^2}. \quad (2.7)$$

As we mentioned in the Introduction, in some recent attempts to construct Kaniadakis holographic dark energy the authors used (2.3) but then they considered the Hubble horizon to be L [74–76]. Thus, the obtained models do not have standard holographic dark energy and standard thermodynamics as a sub-case, and this is a serious disadvantage. One can verify that in a clear way by observing that in order to have reasonable observational results the authors demand K values of the order of 10^3 , namely a huge deviation from standard Bekenstein-Hawking entropy, which is not observed (not mentioning the fact that the initial K parameter of Kaniadakis entropy is bounded in $-1 < K < 1$).

In the present work we desire to formulate Kaniadakis holographic dark energy in a consistent way, and hence we use as L the future event horizon (2.7). In this way, as we will see, standard holographic dark energy is included as a sub-case, and can be obtained for $K \rightarrow 0$. However, let us comment here that using the future event horizon does have the disadvantage that the dark-energy density at present depends on the future expansion of the Universe [16], in a similar way that the use of the Hubble horizon has the disadvantage that the dark-energy density, which is a local concept, depends on the global picture of the whole Universe, namely on the global expansion scale. Nevertheless, we mention that the use of both horizons, as well as other horizons, such as the Granda-Oliveros cutoff [84], is in principle justified exactly by the concept of holography and the dualities known from string theory, that relate the very small with the very large in space and time.

According to the above discussion, and using (2.5) with L the R_h , the energy density of Kaniadakis holographic dark energy writes as

$$\rho_{DE} = 3c^2 M_p^2 R_h^{-2} + K^2 M_p^6 R_h^2. \quad (2.8)$$

The Friedmann equations in a universe containing the dark energy and matter perfect fluids are

$$3M_p^2 H^2 = \rho_m + \rho_{DE} \quad (2.9)$$

$$-2M_p^2 \dot{H} = \rho_m + p_m + \rho_{DE} + p_{DE}, \quad (2.10)$$

with p_{DE} the pressure of Kaniadakis holographic dark energy, and ρ_m and p_m respectively the energy density and pressure of the matter sector. The equations close by considering the matter conservation equation

$$\dot{\rho}_m + 3H(\rho_m + p_m) = 0. \quad (2.11)$$

It proves convenient to introduce the dark energy and matter density parameters through

$$\Omega_m \equiv \frac{1}{3M_p^2 H^2} \rho_m \quad (2.12)$$

$$\Omega_{DE} \equiv \frac{1}{3M_p^2 H^2} \rho_{DE}. \quad (2.13)$$

Using these definitions, relations (2.7),(2.8),(2.13) lead to

$$\int_x^\infty \frac{dx}{Ha} = \frac{1}{a} \left(\frac{3H^2 \Omega_{DE} - \sqrt{9H^4 \Omega_{DE}^2 - 12c^2 K^2 M_p^4}}{2K^2 M_p^4} \right)^{\frac{1}{2}}, \quad (2.14)$$

where $x \equiv \ln a$. Note that solving the fourth-degree algebraic equation (2.8) we have kept only the solutions that give positive R_h and moreover with the usual limiting result for $K \rightarrow 0$. Indeed, as one can see, in the limit $K \rightarrow 0$ the above relation gives the standard holographic dark energy result $\int_x^\infty \frac{dx}{Ha} = \frac{c}{aH\sqrt{\Omega_{DE}}}$.

We focus on the physically interesting dust matter case, where the matter equation-of-state parameter is set to zero. Therefore, (2.11) leads to $\rho_m = \rho_{m0}/a^3$, where ρ_{m0} is the matter energy density at the current scale factor $a_0 = 1$ (we use the subscript “0” to denote the present value of a quantity). Hence, substituting into (2.12) leads to $\Omega_m = \Omega_{m0} H_0^2 / (a^3 H^2)$, and then, using the Friedmann equation $\Omega_m + \Omega_{DE} = 1$, we find

$$\frac{1}{Ha} = \frac{\sqrt{a(1 - \Omega_{DE})}}{H_0 \sqrt{\Omega_{m0}}}. \quad (2.15)$$

Inserting (2.15) into (2.14) leads to

$$\int_x^\infty \frac{dx}{H_0 \sqrt{\Omega_{m0}}} \sqrt{a(1 - \Omega_{DE})} = \frac{1}{a} \left(\frac{3H^2 \Omega_{DE} - \sqrt{9H^4 \Omega_{DE}^2 - 12c^2 K^2 M_p^4}}{2K^2 M_p^4} \right)^{\frac{1}{2}}. \quad (2.16)$$

In the following we use $x = \ln a$ as the independent variable, and therefore for a quantity f we acquire $\dot{f} = f' H$, with primes denoting derivatives with respect to x . Differentiating (2.16) in terms of x we obtain

$$\Omega'_{DE} = \Omega_{DE}(1 - \Omega_{DE}) \left\{ 3 - \frac{2(\mathcal{A} - 2K^2 M_p^4 \mathcal{B})}{\mathcal{A}} \left[1 - \sqrt{3} \left(\frac{\Omega_{DE}}{\mathcal{A}\mathcal{B}} \right)^{\frac{1}{2}} \right] \right\}, \quad (2.17)$$

with

$$\mathcal{A} = \frac{3e^{-3x} H_0^2 \Omega_{m0} \Omega_{DE}}{1 - \Omega_{DE}},$$

$$\mathcal{B} = \frac{\mathcal{A} - \sqrt{\mathcal{A}^2 - 12c^2 K^2 M_p^4}}{2K^2 M_p^4}.$$

Differential equation (2.17) determines the evolution of Kaniadakis holographic dark energy as a function of $x = \ln a$, in the case of flat spatial geometry and for dust matter. We mention that in the limit $K \rightarrow 0$ we have $\mathcal{B}|_{K \rightarrow 0} = \frac{3c^2}{\mathcal{A}}$, and hence (2.17) recovers the corresponding differential equation of usual holographic dark energy [15], i.e. $\Omega'_{DE}|_{K \rightarrow 0} = \Omega_{DE}(1 - \Omega_{DE}) \left(1 + 2\sqrt{\frac{3M_p^2 \Omega_{DE}}{3c^2 M_p^2}}\right)$, which, since the x -dependence is absent, accepts an analytic solution in an implicit form [15].

We proceed by examining the behavior of the equation-of-state parameter $w_{DE} \equiv p_{DE}/\rho_{DE}$ of Kaniadakis holographic dark energy. From the conservation of the matter sector (2.11), and using the two Friedmann equations (2.9),(2.10), we deduce that the dark energy sector is conserved too, i.e.

$$\dot{\rho}_{DE} + 3H\rho_{DE}(1 + w_{DE}) = 0. \quad (2.18)$$

Differentiating (2.8) gives $\dot{\rho}_{DE} = 2M_p^2 (-3c^2 R_h^{-4} + K^2 M_p^4) R_h \dot{R}_h$. In this expression we have that $\dot{R}_h = H R_h - 1$, as it is found from (2.7), where R_h can be further eliminated in terms of ρ_{DE} according to (2.8) as

$$R_h = \left(\frac{\rho_{DE} - \sqrt{\rho_{DE}^2 - 12c^2 K^2 M_p^8}}{2K^2 M_p^6} \right)^{1/2} \equiv \mathcal{C}. \quad (2.19)$$

Substituting all the above into (2.18) we acquire

$$2M_p^2(H\mathcal{C} - 1) \left(\frac{-3c^2 + K^2 M_p^4 \mathcal{C}^4}{\mathcal{C}^3} \right) + 3H\rho_{DE}(1 + w_{DE}) = 0. \quad (2.20)$$

Therefore, inserting H from (2.15), and using definition (2.13) after some algebra we find

$$w_{DE} = -1 - 2 \left(\frac{\Omega_{DE}}{3\mathcal{A}^3} \right)^{\frac{1}{2}} \left(\frac{-3c^2 + K^2 M_p^4 \mathcal{B}^2}{\mathcal{B}^{\frac{3}{2}}} \right) \left[-1 + \frac{\sqrt{3}}{3} \left(\frac{\mathcal{A}\mathcal{B}}{\Omega_{DE}} \right)^{\frac{1}{2}} \right]. \quad (2.21)$$

Hence, w_{DE} as a function of $\ln a$ is known, as long as Ω_{DE} is known from (2.17). Note that for $K \rightarrow 0$ the above expression provides the standard holographic dark energy result, i.e. $w_{DE}|_{K \rightarrow 0} = -\frac{1}{3} - \frac{2}{3} \frac{\sqrt{\Omega_{DE}}}{c}$ [16], as expected. Additionally, we mention that in general w_{DE} can be either quintessence-like or phantom-like, which is an advantage revealing the rich capabilities of the scenario at hand.

Lastly, for convenience we can introduce the deceleration parameter

$$q \equiv -1 - \frac{\dot{H}}{H^2} = \frac{1}{2} + \frac{3}{2} (w_m \Omega_m + w_{DE} \Omega_{DE}), \quad (2.22)$$

which in the case of dust matter is straightforwardly known as long as Ω_{DE} (and thus w_{DE} from (2.21)) is known.

We close this section by discussing the relation of Kaniadakis entropy with other extended entropies, and in particular with Tsallis one. As it is known, the non-extensive

Tsallis entropy S_q^T , with q the parameter which quantifies the deviation from Bekenstein-Hawking entropy [85, 86], is related to Kaniadakis one through [74, 79, 87]

$$S_K = \frac{S_{1+K}^T + S_{1-K}^T}{2}. \quad (2.23)$$

Concerning the other recently proposed generalized entropy by Barrow, namely S_Δ^B , which arises from the intricate structure of the black-hole surface due to quantum-gravitational effects, with Δ the parameter that quantifies the deviation from usual entropy [88], we mention that although mathematically one can extract the relation $S_K = \frac{S_\Delta^B + S_{-\Delta}^B}{2}$, it cannot have a physical application since in Barrow entropy $0 \leq \Delta \leq 1$.

3 Cosmological evolution

In the previous section we formulated Kaniadakis holographic dark energy, and we provided the equations that determine the evolution of the corresponding dark energy density, equation-of-state and deceleration parameters. Hence, we can now proceed to a detailed investigation of the resulting cosmological behavior. Since equation (2.17) can be solved analytically only for $K = 0$, in the general case we should resort to numerical elaboration. As long as we have the solution for $\Omega_{DE}(x)$ we can obtain its behavior in terms of the redshift z through the simple relation $x \equiv \ln a = -\ln(1+z)$. Finally, we mention that Kaniadakis entropy is an even function, namely $S_K = S_{-K}$, and that is why all the above expressions of Kaniadakis holographic dark energy depend only on K^2 . Thus, in the following we focus on the $K \geq 0$ region.

We solve equation (2.17) numerically, imposing $\Omega_{DE}(x = -\ln(1+z) = 0) \equiv \Omega_{DE0} \approx 0.7$ and therefore $\Omega_m(x = -\ln(1+z) = 0) \equiv \Omega_{m0} \approx 0.3$ in agreement with observations [89]. In the upper graph of Fig. 1 we depict the evolution of the dark energy and matter density parameters in terms of the redshift. Additionally, in the middle graph we present the corresponding behavior of the dark-energy equation-of-state parameter as it arises from (2.21). Finally, in the lower graph we show the deceleration parameter as it is given from (2.22). We mention that for reader's convenience we have extended the evolution up to the far future, namely for $z \rightarrow -1$.

As we observe, the scenario at hand can provide the required thermal history of the universe, i.e. the sequence of matter and dark energy epochs, and the universe results asymptotically to a complete dark-energy dominated phase. Moreover, from the middle graph of Fig. 1 we can see that the value of w_{DE} at present is around -1 in agreement with observational data. Note that in this specific example w_{DE} in the future enters slightly inside the phantom regime, which as mentioned above is allowed by (2.21) and shows the capabilities of the model. Finally, from the lower graph of Fig. 1 we deduce that the transition from deceleration to acceleration is realized at $z \approx 0.6$, in agreement with observations.

Let us now study the effect of the model parameters c and K on the dark-energy equation-of-state parameter w_{DE} . In Fig. 2 we depict $w_{DE}(z)$ for fixed $K = 0.1$ and various values of c . As we can see, with c decreasing $w_{DE}(z)$, as well as its present value




Figure 1. **Upper graph:** The Kaniadakis holographic dark energy density parameter Ω_{DE} (blue-solid) and the matter density parameter Ω_m (red-dashed), as a function of the redshift z , for $K = 0.1$ and $c = 0.9$. **Middle graph:** The corresponding dark-energy equation-of-state parameter w_{DE} . **Lower graph:** The corresponding deceleration parameter q . In all graphs we have set $\Omega_{DE}(x = -\ln(1+z) = 0) \equiv \Omega_{DE0} \approx 0.7$ in agreement with observations, and for convenience we have added a vertical dotted line marking the present time $z = 0$.

$w_{DE}(z = 0)$, acquire algebraically lower values, experiencing the phantom-divide crossing during the evolution. Note that for $c < 0.9$ the value of $w_{DE}(z = 0)$ lies in the phantom regime. Furthermore, in Fig. 3 we present $w_{DE}(z)$ for fixed $c = 1$ and various values of K . Here we observe the interesting behavior that for increasing K , at earlier times w_{DE} slightly decreases, in future times it increases, however at times around the present ones it remains almost unaltered. Concerning the asymptotic value of w_{DE} in the far future, namely for $z \rightarrow -1$, as can be deduced from the figures, as well as from (2.21), it depends on the combination of K and c . In summary, we can see that the scenario of Kaniadakis holographic dark energy can lead to very interesting cosmological phenomenology, in which




Figure 2. The redshift-evolution of the equation-of-state parameter w_{DE} of Kaniadakis holographic dark energy, for fixed $K = 0.1$ and various values of c . We have imposed $\Omega_{DE0} \approx 0.7$ and we have added a vertical dotted line marking the present time $z = 0$.

w_{DE} can be quintessence-like, phantom-like, or cross the phantom divide before or after the present time. Note that these results are in agreement with other researches on Kaniadakis holographic dark energy, that appeared after the present work, especially with observational confrontation [90–95].




Figure 3. The redshift-evolution of the equation-of-state parameter w_{DE} of Kaniadakis holographic dark energy, for fixed $c = 1$ and various values of K . We have imposed $\Omega_{DE0} \approx 0.7$ we have added a vertical dotted line marking the present time $z = 0$.

4 Conclusions

In this work we formulated a holographic dark energy scenario based on Kaniadakis entropy. The latter is a generalization of Boltzmann-Gibbs entropy, arising from a coherent relativistic statistical theory and characterized by a single parameter K that quantifies the deviations from standard expressions. Hence, by applying the usual steps of holographic dark energy, imposing the future event horizon as the Infrared cutoff, and using Kaniadakis extended entropy, we obtained Kaniadakis holographic dark energy in a consistent way, namely a one-parameter extension of usual holographic dark energy, possessing it as a particular limit, namely for $K \rightarrow 0$.

In order to investigate the cosmological application of Kaniadakis holographic dark energy we extracted the differential equation that determines the evolution of the effective dark energy density parameter Ω_{DE} . Moreover, we provided analytical expressions for the corresponding equation-of-state parameter w_{DE} , as well as for the deceleration parameter.

The scenario of Kaniadakis holographic dark energy proves to lead to interesting cosmological behavior. In particular, the universe exhibits the standard thermal history, i.e. the sequence of matter and dark-energy eras, while the transition to acceleration takes place at $z \approx 0.6$. Concerning the dark-energy equation-of-state parameter we saw that it can have a rich behavior, being quintessence-like, phantom-like, or experience the phantom-divide crossing in the past or in the future, depending on the values of the two model parameters c and K . In particular, for fixed K decreasing c leads to algebraically smaller w_{DE} values, while for fixed c by increasing K we acquire smaller w_{DE} values at higher redshifts, larger w_{DE} values in the future, and almost unaltered values at present. Finally, in the far future dark energy dominates completely, and the asymptotic w_{DE} value depends on c and K .

We comment here that, as we mentioned in the Introduction, it has been recently shown that holographic dark energy constructions may alleviate the H_0 tension (see the corresponding sections in the recent review [34]), and the reason is that they may lead to $w_{DE} < -1$ which seems to be a requirement if one desires to provide a solution based on late-time modifications [66, 96]. As we saw, the scenario at hand can fulfill this requirement and that is why it is a candidate to be able to alleviate the H_0 tension too. Definitely the phantom regime may have potential disadvantages, however this is not in general the case in models which present phantom behavior in an effective way, among which is holographic dark energy [16].

In conclusion, Kaniadakis holographic dark energy exhibits richer and more interesting behavior in comparison to usual holographic dark energy. Additionally, due to the consistent formulation, it possesses the latter as a limiting sub-case. Definitely, before one considers it as a successful candidate for the description of dark energy, there are necessary investigations that should be performed. In particular, one should confront the scenario with observational data from Supernova type Ia (SNIa), Baryon Acoustic Oscillation (BAO), Cosmic Microwave Background (CMB), and Hubble parameter observations, and extract constraints on the model parameters. Additionally, one should analyze in detail the phase-space behavior, in order to examine the global dynamics and the asymptotic, late-time evolution of the scenario. These investigations will be performed in separate

projects.

Acknowledgments

The authors would like to acknowledge the contribution of the COST Action CA18108 “Quantum Gravity Phenomenology in the multi-messenger approach”. The work is partially supported by the Ministry of Education and Science of the Republic of Kazakhstan, Grant AP08856912.

References

- [1] E. J. Copeland, M. Sami and S. Tsujikawa, *Dynamics of dark energy*, Int. J. Mod. Phys. D **15**, 1753 (2006) [[arXiv:hep-th/0603057](#)].
- [2] Y. -F. Cai, E. N. Saridakis, M. R. Setare and J. -Q. Xia, *Quintom Cosmology: Theoretical implications and observations*, Phys. Rept. **493**, 1 (2010) [[arXiv:0909.2776](#)].
- [3] K. Bamba, S. Capozziello, S. Nojiri and S. D. Odintsov, *Dark energy cosmology: the equivalent description via different theoretical models and cosmography tests*, Astrophys. Space Sci. **342**, 155 (2012) [[arXiv:1205.3421](#)].
- [4] S. Nojiri and S. D. Odintsov, *Unified cosmic history in modified gravity: from $F(R)$ theory to Lorentz non-invariant models*, Phys. Rept. **505**, 59 (2011) [[arXiv:1011.0544](#)].
- [5] S. Capozziello and M. De Laurentis, *Extended Theories of Gravity*, Phys. Rept. **509**, 167 (2011) [[arXiv:1108.6266](#)].
- [6] Y. F. Cai, S. Capozziello, M. De Laurentis and E. N. Saridakis, *$f(T)$ teleparallel gravity and cosmology*, Rept. Prog. Phys. **79**, 106901 (2016) [[arXiv:1511.07586](#)].
- [7] E. N. Saridakis *et al.* [CANTATA], *Modified Gravity and Cosmology: An Update by the CANTATA Network*, [[arXiv:2105.12582](#)].
- [8] W. Fischler and L. Susskind, *Holography and cosmology*, [[arXiv:hep-th/9806039](#)].
- [9] D. Bak and S. J. Rey, *Cosmic holography*, Class. Quant. Grav. **17**, L83 (2000) [[arXiv:hep-th/9902173](#)].
- [10] P. Horava and D. Minic, *Probable values of the cosmological constant in a holographic theory*, Phys. Rev. Lett. **85**, 1610 (2000) [[arXiv:hep-th/0001145](#)].
- [11] G. 't Hooft, *Dimensional reduction in quantum gravity*, Salamfest 1993: 0284-296 [[arXiv:gr-qc/9310026](#)].
- [12] L. Susskind, *The World as a hologram*, J. Math. Phys. **36**, 6377 (1995) [[arXiv:hep-th/9409089](#)].
- [13] R. Bousso, *The Holographic principle*, Rev. Mod. Phys. **74**, 825 (2002) [[arXiv:hep-th/0203101](#)].
- [14] A. G. Cohen, D. B. Kaplan and A. E. Nelson, *Effective field theory, black holes, and the cosmological constant*, Phys. Rev. Lett. **82**, 4971 (1999) [[arXiv:hep-th/9803132](#)].
- [15] M. Li, *A Model of holographic dark energy*, Phys. Lett. B **603**, 1 (2004) [[arXiv:hep-th/0403127](#)].

- [16] S. Wang, Y. Wang and M. Li, *Holographic Dark Energy*, hys. Rept. **696**, 1 (2017) [[arXiv:1612.00345](#)].
- [17] R. Horvat, *Holography and variable cosmological constant*, Phys. Rev. D **70**, 087301 (2004) [[arXiv:astro-ph/0404204](#)].
- [18] Q. G. Huang and M. Li, *The Holographic dark energy in a non-flat universe*, JCAP **0408**, 013 (2004) [[arXiv:astro-ph/0404229](#)].
- [19] D. Pavon and W. Zimdahl, *Holographic dark energy and cosmic coincidence*, Phys. Lett. B **628**, 206 (2005) [[arXiv:gr-qc/0505020](#)].
- [20] B. Wang, Y. g. Gong and E. Abdalla, *Transition of the dark energy equation of state in an interacting holographic dark energy model*, Phys. Lett. B **624**, 141 (2005) [[arXiv:hep-th/0506069](#)].
- [21] S. Nojiri and S. D. Odintsov, *Unifying phantom inflation with late-time acceleration: Scalar phantom-non-phantom transition model and generalized holographic dark energy*, Gen. Rel. Grav. **38**, 1285 (2006) [[arXiv:hep-th/0506212](#)].
- [22] H. Kim, H. W. Lee and Y. S. Myung, *Equation of state for an interacting holographic dark energy model*, Phys. Lett. B **632**, 605 (2006) [[arXiv:gr-qc/0509040](#)].
- [23] B. Wang, C. Y. Lin and E. Abdalla, *Constraints on the interacting holographic dark energy model*, Phys. Lett. B **637**, 357 (2006) [[arXiv:hep-th/0509107](#)].
- [24] M. R. Setare, *Interacting holographic dark energy model in non-flat universe*, Phys. Lett. B **642**, 1 (2006) [[arXiv:hep-th/0609069](#)].
- [25] M. R. Setare and E. N. Saridakis, *Non-minimally coupled canonical, phantom and quintom models of holographic dark energy*, Phys. Lett. B **671**, 331 (2009) [[arXiv:0810.0645](#)].
- [26] M. R. Setare and E. N. Saridakis, *Correspondence between Holographic and Gauss-Bonnet dark energy models*, Phys. Lett. B **670**, 1 (2008) [[arXiv:0810.3296](#)].
- [27] X. Zhang and F. Q. Wu, *Constraints on holographic dark energy from Type Ia supernova observations*, Phys. Rev. D **72**, 043524 (2005) [[arXiv:0506310](#)].
- [28] M. Li, X. D. Li, S. Wang and X. Zhang, *Holographic dark energy models: A comparison from the latest observational data*, JCAP **0906**, 036 (2009) [[arXiv:0904.0928](#)].
- [29] C. Feng, B. Wang, Y. Gong and R. K. Su, *Testing the viability of the interacting holographic dark energy model by using combined observational constraints*, JCAP **0709**, 005 (2007) [[arXiv:0706.4033](#)].
- [30] X. Zhang, *Holographic Ricci dark energy: Current observational constraints, quintom feature, and the reconstruction of scalar-field dark energy*, Phys. Rev. D **79**, 103509 (2009) [[arXiv:0901.2262](#)].
- [31] J. Lu, E. N. Saridakis, M. R. Setare and L. Xu, *Observational constraints on holographic dark energy with varying gravitational constant*, JCAP **1003**, 031 (2010) [[arXiv:0912.0923](#)].
- [32] S. M. R. Micheletti, *Observational constraints on holographic tachyonic dark energy in interaction with dark matter*, JCAP **1005**, 009 (2010) [[arXiv:0912.3992](#)].
- [33] S. Weinberg, *The Cosmological Constant Problem*, Rev. Mod. Phys. **61**, 1-23 (1989).
- [34] E. Abdalla, G. F. Abellán, A. Aboubrahim, A. Agnello, O. Akarsu, Y. Akrami, G. Alestas,

- D. Aloni, L. Amendola and L. A. Anchordoqui, *et al. Cosmology Intertwined: A Review of the Particle Physics, Astrophysics, and Cosmology Associated with the Cosmological Tensions and Anomalies*, [[arXiv:2203.06142](#)].
- [35] Y. g. Gong, *Extended holographic dark energy*, Phys. Rev. D **70**, 064029 (2004) [[arXiv:0404030](#)].
- [36] E. N. Saridakis, *Restoring holographic dark energy in brane cosmology*, Phys. Lett. B **660**, 138 (2008) [[arXiv:0712.2228](#)].
- [37] M. R. Setare and E. C. Vagenas, *The Cosmological dynamics of interacting holographic dark energy model*, Int. J. Mod. Phys. D **18**, 147 (2009) [[arXiv:0704.2070](#)].
- [38] R. G. Cai, *A Dark Energy Model Characterized by the Age of the Universe*, Phys. Lett. B **657**, 228 (2007) [[arXiv:0707.4049](#)].
- [39] E. N. Saridakis, *Holographic Dark Energy in Braneworld Models with Moving Branes and the $w=-1$ Crossing*, JCAP **0804**, 020 (2008) [[arXiv:0712.2672](#)].
- [40] E. N. Saridakis, *Holographic Dark Energy in Braneworld Models with a Gauss-Bonnet Term in the Bulk. Interacting Behavior and the $w=-1$ Crossing*, Phys. Lett. B **661**, 335 (2008) [[arXiv:0712.3806](#)].
- [41] M. R. Setare and E. C. Vagenas, *Thermodynamical Interpretation of the Interacting Holographic Dark Energy Model in a non-flat Universe*, Phys. Lett. B **666**, 111 (2008) [[arXiv:0801.4478](#)].
- [42] M. Jamil, E. N. Saridakis and M. R. Setare, *Holographic dark energy with varying gravitational constant*, Phys. Lett. B **679**, 172 (2009) [[arXiv:0906.2847](#)].
- [43] Y. Gong and T. Li, *A Modified Holographic Dark Energy Model with Infrared Infinite Extra Dimension(s)*, Phys. Lett. B **683**, 241 (2010) [[arXiv:0907.0860](#)].
- [44] M. Suwa and T. Nihei, *Observational constraints on the interacting Ricci dark energy model*, Phys. Rev. D **81**, 023519 (2010) [[arXiv:0911.4810](#)].
- [45] M. Jamil and E. N. Saridakis, *New agegraphic dark energy in Horava-Lifshitz cosmology*, JCAP **1007**, 028 (2010) [[arXiv:1003.5637](#)].
- [46] M. Bouhmadi-Lopez, A. Errahmani and T. Ouali, *The cosmology of an holographic induced gravity model with curvature effects*, Phys. Rev. D **84**, 083508 (2011) [[arXiv:1104.1181](#)].
- [47] M. Malekjani, *Generalized holographic dark energy model described at the Hubble length*, Astrophys. Space Sci. **347**, 405 (2013) [[arXiv:1209.5512](#)].
- [48] M. Khurshudyan, J. Sadeghi, R. Myrzakulov, A. Pasqua and H. Farahani, *Interacting quintessence dark energy models in Lyra manifold*, Adv. High Energy Phys. **2014**, 878092 (2014) [[arXiv:1404.2141](#)].
- [49] R. C. G. Landim, *Holographic dark energy from minimal supergravity*, Int. J. Mod. Phys. D **25**, no. 04, 1650050 (2016) [[arXiv:1508.07248](#)].
- [50] A. Pasqua, S. Chattopadhyay and R. Myrzakulov, *Power-law entropy-corrected holographic dark energy in Hořava-Lifshitz cosmology with Granda-Oliveros cut-off*, Eur. Phys. J. Plus **131**, no. 11, 408 (2016) [[arXiv:1511.00611](#)].
- [51] A. Jawad, N. Azhar and S. Rani, *Entropy corrected holographic dark energy models in modified gravity*, Int. J. Mod. Phys. D **26**, no. 04, 1750040 (2016).

- [52] B. Pourhassan, A. Bonilla, M. Faizal and E. M. C. Abreu, *Holographic Dark Energy from Fluid/Gravity Duality Constraint by Cosmological Observations*, Phys. Dark Univ. **20**, 41 (2018) [[arXiv:1704.03281](#)].
- [53] E. N. Saridakis, *Ricci-Gauss-Bonnet holographic dark energy*, Phys. Rev. D **97**, no. 6, 064035 (2018) [[arXiv:1707.09331](#)].
- [54] S. Nojiri and S. D. Odintsov, *Covariant Generalized Holographic Dark Energy and Accelerating Universe*, Eur. Phys. J. C **77**, no. 8, 528 (2017) [[arXiv:1703.06372](#)].
- [55] E. N. Saridakis, K. Bamba, R. Myrzakulov and F. K. Anagnostopoulos, *Holographic dark energy through Tsallis entropy*, JCAP **12**, 012 (2018) [[arXiv:1806.01301](#)].
- [56] A. Oliveros and M. A. Acero, *Inflation driven by a holographic energy density*, EPL **128**, no.5, 59001 (2019) [[arXiv:1911.04482](#)].
- [57] C. Kritpetch, C. Muhammad and B. Gumjudpai, *Holographic dark energy with non-minimal derivative coupling to gravity effects*, Phys. Dark Univ. **30**, 100712 (2020) [[arXiv:2004.06214](#)].
- [58] E. N. Saridakis, *Barrow holographic dark energy*, Phys. Rev. D **102**, no.12, 123525 (2020) [[arXiv:2005.04115](#)].
- [59] M. P. Dabrowski and V. Salzano, *Geometrical observational bounds on a fractal horizon holographic dark energy*, Phys. Rev. D **102**, no.6, 064047 (2020) [[arXiv:2009.08306](#)].
- [60] W. J. C. da Silva and R. Silva, *Cosmological Perturbations in the Tsallis Holographic Dark Energy Scenarios*, Eur. Phys. J. Plus **136**, no.5, 543 (2021) [[arXiv:2011.09520](#)].
- [61] F. K. Anagnostopoulos, S. Basilakos and E. N. Saridakis, *Observational constraints on Barrow holographic dark energy*, Eur. Phys. J. C **80**, no.9, 826 (2020) [[arXiv:2005.10302](#)].
- [62] A. A. Mamon, A. Paliathanasis and S. Saha, *Dynamics of an Interacting Barrow Holographic Dark Energy Model and its Thermodynamic Implications*, Eur. Phys. J. Plus **136**, no.1, 134 (2021) [[arXiv:2007.16020](#)].
- [63] S. Bhattacharjee, *Growth Rate and Configurational Entropy in Tsallis Holographic Dark Energy*, Eur. Phys. J. C **81**, no.3, 217 (2021) [[arXiv:2011.13135](#)].
- [64] Q. Huang, H. Huang, B. Xu, F. Tu and J. Chen, *Dynamical analysis and statefinder of Barrow holographic dark energy*, Eur. Phys. J. C **81**, no.8, 686 (2021).
- [65] C. Lin, *An effective field theory of holographic dark energy*, JCAP **07**, 003 (2021) [[arXiv:2101.08092](#)].
- [66] E. Ó. Colgáin and M. M. Sheikh-Jabbari, *A critique of holographic dark energy*, Class. Quant. Grav. **38**, no.17, 177001 (2021) [[arXiv:2102.09816](#)].
- [67] H. Hossienkhani, N. Azimi and H. Yousefi, *Constraints on the Ricci dark energy cosmologies in Bianchi type I model*, Int. J. Geom. Meth. Mod. Phys. **18**, no.06, 2150095 (2021).
- [68] S. Nojiri, S. D. Odintsov and T. Paul, *Different Faces of Generalized Holographic Dark Energy*, Symmetry **13**, no.6, 928 (2021) [[arXiv:2105.08438](#)].
- [69] S. H. Shekh, *Models of holographic dark energy in $f(Q)$ gravity*, Phys. Dark Univ. **33**, 100850 (2021).
- [70] E. C. Telali and E. N. Saridakis, *Power-law holographic dark energy and cosmology*, [[arXiv:2112.06821](#)].

- [71] A. Y. Shaikh, *Panorama behaviors of general relativistic hydrodynamics and holographic dark energy in $f(R, T)$ gravity*, New Astron. **91**, 101676 (2022).
- [72] G. Kaniadakis, *Statistical mechanics in the context of special relativity*, Phys. Rev. E **66**, 056125 (2002) [[arXiv:cond-mat/0210467](#)].
- [73] G. Kaniadakis, *Statistical mechanics in the context of special relativity. II*, Phys. Rev. E **72**, 036108 (2005) [[arXiv:cond-mat/0507311](#)].
- [74] H. Moradpour, A. H. Ziaie and M. Kord Zangeneh, *Generalized entropies and corresponding holographic dark energy models*, Eur. Phys. J. C **80**, no.8, 732 (2020) [[arXiv:2005.06271](#)].
- [75] A. Jawad and A. M. Sultan, *Cosmic Consequences of Kaniadakis and Generalized Tsallis Holographic Dark Energy Models in the Fractal Universe*, Adv. High Energy Phys. **2021**, 5519028 (2021).
- [76] U. K. Sharma, V. C. Dubey, A. H. Ziaie and H. Moradpour, *Kaniadakis Holographic Dark Energy in non-flat Universe*, Int. J. Mod. Phys. D **31**, no.03, 2250013 (2022) [[arXiv:2106.08139](#)].
- [77] E. M. C. Abreu, J. Ananias Neto, E. M. Barboza and R. C. Nunes, *Jeans instability criterion from the viewpoint of Kaniadakis' statistics*, EPL **114**, no.5, 55001 (2016) [[arXiv:1603.00296](#)].
- [78] E. M. C. Abreu, J. A. Neto, E. M. Barboza and R. C. Nunes, *Tsallis and Kaniadakis statistics from the viewpoint of entropic gravity formalism*, Int. J. Mod. Phys. A **32**, no.05, 1750028 (2017) [[arXiv:1701.06898](#)].
- [79] E. M. C. Abreu, J. A. Neto, A. C. R. Mendes and A. Bonilla, *Tsallis and Kaniadakis statistics from a point of view of the holographic equipartition law*, EPL **121**, no.4, 45002 (2018) [[arXiv:1711.06513](#)].
- [80] E. M. C. Abreu, J. A. Neto, A. C. R. Mendes and R. M. de Paula, *Loop Quantum Gravity Immirzi parameter and the Kaniadakis statistics*, Chaos, Solitons and Fractals **118**, 307-310 (2019) [[arXiv:1808.01891](#)].
- [81] W. H. Yang, Y. Z. Xiong, H. Chen and S. Q. Liu, *Jeans gravitational instability with κ -deformed Kaniadakis distribution in Eddington-inspired Born Infield gravity*, Chin. Phys. B **29**, no.11, 110401 (2020).
- [82] E. M. C. Abreu and J. Ananias Neto, *Black holes thermodynamics from a dual Kaniadakis entropy*, EPL **133**, no.4, 49001 (2021).
- [83] S. D. H. Hsu, *Entropy bounds and dark energy*, Phys. Lett. B **594**, 13 (2004) [[arXiv:hep-th/0403052](#)].
- [84] L. N. Granda and A. Oliveros, *Infrared cut-off proposal for the Holographic density*, Phys. Lett. B **669**, 275-277 (2008) [[arXiv:0810.3149](#)].
- [85] C. Tsallis, *Possible Generalization of Boltzmann-Gibbs Statistics*, J. Statist. Phys. **52**, 479 (1988).
- [86] C. Tsallis and L. J. L. Cirto, *Black hole thermodynamical entropy*, Eur. Phys. J. C **73**, 2487 (2013) [[arXiv:1202.2154](#)].
- [87] R. C. Nunes, E. M. Barboza, Jr., E. M. C. Abreu and J. A. Neto, *Probing the cosmological viability of non-gaussian statistics*, JCAP **08**, 051 (2016) [[arXiv:1509.05059](#)].
- [88] J. D. Barrow, *The Area of a Rough Black Hole*, Phys. Lett. B **808**, 135643 (2020)

- [[arXiv:2004.09444](#)].
- [89] N. Aghanim *et al.* [Planck], *Planck 2018 results. VI. Cosmological parameters*, Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] [[arXiv:1807.06209](#)].
 - [90] A. Hernández-Almada, G. Leon, J. Magaña, M. A. García-Aspeitia, V. Motta, E. N. Saridakis, K. Yesmakhanova and A. D. Millano, *Observational constraints and dynamical analysis of Kaniadakis horizon-entropy cosmology*, [[arXiv:2112.04615](#)].
 - [91] A. Lymperis, S. Basilakos and E. N. Saridakis, *Modified cosmology through Kaniadakis horizon entropy*, Eur. Phys. J. C **81**, no.11, 1037 (2021) [[arXiv:2108.12366](#)].
 - [92] G. G. Luciano, *Modified Friedmann equations from Kaniadakis entropy and cosmological implications on baryogenesis and ${}^7\text{Li}$ -abundance*, Eur. Phys. J. C **82**, no.4, 314 (2022).
 - [93] A. Hernández-Almada, G. Leon, J. Magaña, M. A. García-Aspeitia, V. Motta, E. N. Saridakis and K. Yesmakhanova, *Kaniadakis-holographic dark energy: observational constraints and global dynamics*, Mon. Not. Roy. Astron. Soc. **511**, no.3, 4147-4158 (2022) [[arXiv:2111.00558](#)].
 - [94] S. Ghaffari, *Kaniadakis holographic dark energy in Brans-Dicke cosmology*, [[arXiv:2112.05813](#)].
 - [95] J. Sadeghi, S. N. Gashti and T. Azizi, *Tsallis and Kaniadakis holographic dark energy with Complex Quintessence theory in Brans-Dicke cosmology*, [[arXiv:2203.04375](#)].
 - [96] S. Vagnozzi, *New physics in light of the H_0 tension: An alternative view*, Phys. Rev. D **102**, no.2, 023518 (2020) [[arXiv:1907.07569](#)].