

SOME RESULTS ON VARIABLE GAUSSIAN BESOV-LIPSCHITZ AND VARIABLE GAUSSIAN TRIEBEL-LIZORKIN SPACES.

EBNER PINEDA, LUZ RODRIGUEZ, AND WILFREDO O. URBINA

ABSTRACT. In a previous paper [6] two of the authors introduced and study Gaussian Besov-Lipschitz spaces $B_{p,q}^\alpha(\gamma_d)$ and Gaussian Triebel-Lizorkin spaces $F_{p,q}^\alpha(\gamma_d)$. Now, in this paper we introduce the variable Gaussian Besov-Lipschitz spaces $B_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$ and the variable Gaussian Triebel-Lizorkin spaces $F_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$, that is to say, Gaussian Besov-Lipschitz and Triebel-Lizorkin spaces with variable exponents, following [7] and [6] under certain additional regularity conditions on the exponents $p(\cdot)$ and $q(\cdot)$ introduced by Dalmasso and Scotto in [7]. Trivially, they include the Gaussian Besov-Lipschitz spaces $B_{p,q}^\alpha(\gamma_d)$ and Gaussian Triebel-Lizorkin spaces $F_{p,q}^\alpha(\gamma_d)$. We consider some inclusion relations of those spaces and finally we also prove some interpolation results for them.

1. INTRODUCTION AND PRELIMINARIES

Let us consider the Gaussian measure

$$(1.1) \quad \gamma_d(dx) = \frac{e^{-\|x\|^2}}{\pi^{d/2}} dx, \quad x \in \mathbb{R}^d$$

on $\mathbb{R}^d$ and the Ornstein-Uhlenbeck differential operator

$$(1.2) \quad L = \frac{1}{2} \Delta_x - \langle x, \nabla_x \rangle.$$

Let $\nu = (\nu_1, \dots, \nu_d)$ be a multi-index such that $\nu_i \geq 0, i = 1, \dots, d$, let $\nu! = \prod_{i=1}^d \nu_i!$, $|\nu| = \sum_{i=1}^d \nu_i$, $\partial_i = \frac{\partial}{\partial x_i}$, for each $1 \leq i \leq d$ and $\partial^\nu = \partial_1^{\nu_1} \dots \partial_d^{\nu_d}$. Consider the normalized Hermite polynomials of order ν in d variables,

$$(1.3) \quad h_\nu(x) = \frac{1}{(2^{|\nu|} \nu!)^{1/2}} \prod_{i=1}^d (-1)^{\nu_i} e^{x_i^2} \partial_i^{\nu_i} (e^{-x_i^2}).$$

The *Ornstein-Uhlenbeck semigroup* on $\mathbb{R}^d$ is defined by

$$T_t f(x) = \frac{1}{(1 - e^{-2t})^{d/2}} \int_{\mathbb{R}^d} e^{-\frac{e^{-2t}(|x|^2 + |y|^2) - 2e^{-t}\langle x, y \rangle}{1 - e^{-2t}}} f(y) \gamma_d(dy) \quad .$$

Using the *Bochner subordination formula*

$$(1.4) \quad e^{-\lambda} = \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{e^{-u}}{\sqrt{u}} e^{-\lambda^2/4u} du,$$

2010 *Mathematics Subject Classification.* Primary 42B25, 42B35; Secondary 46E30, 47G10.

Key words and phrases. Hermite expansions, variable exponent, Besov-Lipschitz, Triebel-Lizorkin, Gaussian measure.

we introduce the *Poisson-Hermite semigroup* by

$$(1.5) \quad P_t f(x) = \frac{1}{\sqrt{\pi}} \int_0^\infty \frac{e^{-u}}{\sqrt{u}} T_{t^2/4u} f(x) du.$$

Now, taking the change of variables $s = \frac{t^2}{4u}$, $P_t f(x)$ can be written as

$$(1.6) \quad P_t f(x) = \int_0^\infty T_s f(x) \mu_t^{(1/2)}(ds),$$

where

$$\mu_t^{(1/2)}(ds) = \frac{t}{2\sqrt{\pi}} e^{-t^2/4s} s^{-3/2} ds,$$

is the one-sided stable measure on $(0, \infty)$ of order $1/2$.

It is easy to see that $\mu_t^{(1/2)}$ is a probability measure on $(0, \infty)$.

It is well known, that the Hermite polynomials are eigenfunctions of the operator L ,

$$(1.7) \quad Lh_\nu(x) = -|\nu| h_\nu(x).$$

In consequence

$$(1.8) \quad T_t h_\nu(x) = e^{-t|\nu|} h_\nu(x),$$

and

$$(1.9) \quad P_t h_\nu(x) = e^{-t\sqrt{|\nu|}} h_\nu(x),$$

i.e. the Hermite polynomials are also eigenfunctions of T_t and P_t for any $t \geq 0$, for more details, see [10].

Next, we present some technical results for the measure $\mu_t^{(1/2)}$ needed in what follows. First, as $\mu_t^{(1/2)}(ds) = \frac{t}{2\sqrt{\pi}} \frac{e^{-t^2/4s}}{s^{3/2}} ds = g(t, s) ds$, for any $k \in \mathbb{N}$, we use the notation $\frac{\partial^k}{\partial t^k} \mu_t^{(1/2)}(ds)$ for

$$(1.10) \quad \frac{\partial^k}{\partial t^k} \mu_t^{(1/2)}(ds) := \frac{\partial^k g(t, s)}{\partial t^k} ds.$$

Lemma 1.1. *Given $k \in \mathbb{N}$*

$$(1.11) \quad \frac{\partial^k \mu_t^{(1/2)}}{\partial t^k}(ds) = \left(\sum_{i,j} a_{i,j} \frac{t^i}{s^j} \right) \mu_t^{(1/2)}(ds)$$

where $\{a_{i,j}\}$ is a finite set of constants and the indexes $i \in \mathbb{Z}$, $j \in \mathbb{N}$ verifies the equation $2j - i = k$.

Lemma 1.2. *Given $k \in \mathbb{N}$ and $t > 0$*

$$\int_0^{+\infty} \frac{1}{s^k} \mu_t^{(1/2)}(ds) = \frac{C_k}{t^{2k}},$$

where $C_k = \frac{2^{2k} \Gamma(k + \frac{1}{2})}{\pi^{\frac{1}{2}}}$.

Corollary 1.1. *Given $k \in \mathbb{N}$ and $t > 0$*

$$(1.12) \quad \int_0^{+\infty} \left| \frac{\partial^k \mu_t^{(1/2)}}{\partial t^k} \right| (ds) \leq \frac{C_k}{t^k}.$$

On the other hand, by considering the *maximal function of the Ornstein-Uhlenbeck semigroup*

$$T^* f(x) = \sup_{t>0} |T_t f(x)|,$$

we obtain the inequality:

Lemma 1.3. *Let $f \in L^1(\gamma_d)$, $x \in \mathbb{R}^d$ and $k \in \mathbb{N}$*

$$(1.13) \quad \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \leq C_k T^* f(x) t^{-k}, \quad \forall t > 0.$$

For the proofs of these technical results, see [6] or [10].

Now, for completeness, we need some background on variable Lebesgue spaces with respect to a Borel measure μ .

A μ -measurable function $p(\cdot) : \Omega \subset \mathbb{R}^d \rightarrow [1, \infty]$ is said to be an *exponent function*; the set of all the exponent functions will be denoted by $\mathcal{P}(\Omega, \mu)$. For $E \subset \Omega$ we set

$$p_-(E) = \operatorname{ess\,inf}_{x \in E} p(x) \text{ and } p_+(E) = \operatorname{ess\,sup}_{x \in E} p(x).$$

$$\Omega_\infty = \{x \in \Omega : p(x) = \infty\}.$$

We use the abbreviations $p_+ = p_+(\Omega)$ and $p_- = p_-(\Omega)$.

Definition 1.1. *Let $E \subset \mathbb{R}^d$. we say that $p(\cdot) : E \rightarrow \mathbb{R}$ is locally log-Hölder continuous, denote by $p(\cdot) \in LH_0(E)$, if there exists a constant $C_1 > 0$ such that*

$$|p(x) - p(y)| \leq \frac{C_1}{\log(e + \frac{1}{|x-y|})}$$

for all $x, y \in E$.

We say that $p(\cdot)$ is log-Hölder continuous at infinity with base point at $x_0 \in \mathbb{R}^d$, and denote this by $p(\cdot) \in LH_\infty(E)$, if there exist constants $p_\infty \in \mathbb{R}$ and $C_2 > 0$ such that

$$|p(x) - p_\infty| \leq \frac{C_2}{\log(e + |x - x_0|)}$$

for all $x \in E$.

We say that $p(\cdot)$ is log-Hölder continuous, and denote this by $p(\cdot) \in LH(E)$ if both conditions are satisfied. The maximum, $\max\{C_1, C_2\}$ is called the log-Hölder constant of $p(\cdot)$.

Definition 1.2. *Let $E \subset \mathbb{R}^d$, we say that $p(\cdot) \in \mathcal{P}_d^{\log}(E)$, if $\frac{1}{p(\cdot)}$ is log-Hölder continuous and denote by $C_{\log}(p)$ or $C_{\log}$ the log-Hölder constant of $\frac{1}{p(\cdot)}$.*

Definition 1.3. Let $\Omega \subset \mathbb{R}^d$ and $p(\cdot) \in \mathcal{P}(\Omega, \mu)$. For a μ -measurable function $f : \Omega \rightarrow \overline{\mathbb{R}}$, we define the modular

$$(1.14) \quad \rho_{p(\cdot), \mu}(f) = \int_{\Omega \setminus \Omega_\infty} |f(x)|^{p(x)} \mu(dx) + \|f\|_{L^\infty(\Omega_\infty, \mu)},$$

and the norm

$$(1.15) \quad \|f\|_{L^{p(\cdot)}(\Omega, \mu)} = \inf \left\{ \lambda > 0 : \rho_{p(\cdot), \mu}(f/\lambda) \leq 1 \right\}.$$

Definition 1.4. The variable exponent Lebesgue space on $\Omega \subset \mathbb{R}^d$, $L^{p(\cdot)}(\Omega, \mu)$ consists on those μ -measurable functions f for which there exists $\lambda > 0$ such that $\rho_{p(\cdot), \mu}\left(\frac{f}{\lambda}\right) < \infty$, i.e.

$$L^{p(\cdot)}(\Omega, \mu) = \left\{ f : \Omega \rightarrow \overline{\mathbb{R}} : f \text{ is measurable and } \rho_{p(\cdot), \mu}\left(\frac{f}{\lambda}\right) < \infty, \text{ for some } \lambda > 0 \right\}.$$

Observation 1.1. When μ is the Lebesgue measure, we write $\rho_{p(\cdot)}$ and $\|f\|_{p(\cdot)}$ instead of $\rho_{p(\cdot), \mu}$ and $\|f\|_{p(\cdot), \mu}$.

Theorem 1.1. (Norm conjugate formula) Let ν a complete, σ -finite measure on Ω . $p(\cdot) \in \mathcal{P}(\Omega, \nu)$, then

$$(1.16) \quad \frac{1}{2} \|f\|_{p(\cdot), \nu} \leq \|f\|'_{p(\cdot), \nu} \leq 2 \|f\|_{p(\cdot), \nu},$$

for all f ν -measurable on Ω ,

$$\text{donde } \|f\|'_{p(\cdot), \nu} = \sup \left\{ \int_{\Omega} |f|g d\mu : g \in L^{p'(\cdot)}(\Omega, \nu), \|g\|_{p'(\cdot), \nu} \leq 1 \right\}.$$

Proof. See Corollary 3.2.14 in [3] □

Theorem 1.2. (Hölder's inequality) Let ν a complete, σ -finite measure on Ω . $r(\cdot), q(\cdot) \in \mathcal{P}(\Omega, \nu)$, define $p(\cdot) \in \mathcal{P}(\Omega, \nu)$ by $\frac{1}{p(x)} = \frac{1}{q(x)} + \frac{1}{r(x)}$ ν a.e. $x \in \Omega$. Then for all $f \in L^{q(\cdot)}(\Omega, \nu)$ and $g \in L^{r(\cdot)}(\Omega, \nu)$, $fg \in L^{p(\cdot)}(\Omega, \nu)$ and

$$(1.17) \quad \|fg\|_{p(\cdot), \nu} \leq 2 \|f\|_{q(\cdot), \nu} \|g\|_{r(\cdot), \nu}$$

Proof. See Lemma 3.2.20 in [3] □

Theorem 1.3. (Minkowski's integral inequality for variable Lebesgue spaces) Given μ and ν complete σ -finite measures on X and Y respectively, $p \in \mathcal{P}(X, \mu)$. Let $f : X \times Y \rightarrow \overline{\mathbb{R}}$ measurable with respect to the product measure on $X \times Y$, such that for almost every $y \in Y$, $f(\cdot, y) \in L^{p(\cdot)}(X, \mu)$. Then

$$(1.18) \quad \left\| \int_Y f(\cdot, y) d\nu(y) \right\|_{p(\cdot), \mu} \leq 4 \int_Y \|f(\cdot, y)\|_{p(\cdot), \mu} d\nu(y)$$

Proof. It is completely analogous to the proof of Corollary 2.38 in [1] by interchanging the Lebesgue measure for complete σ -finite measures μ and ν on X and Y respectively, and by using (1.17), Fubini's theorem and then (1.16). □

In what follows μ represents the measure $\mu(dt) = \frac{dt}{t}$ on $\mathbb{R}^+$.

Observation 1.2. For a μ -measurable function $f : \mathbb{R}^+ \rightarrow \overline{\mathbb{R}}$, $q(\cdot) \in \mathcal{P}(\mathbb{R}^+, \mu)$, and any $\lambda > 0$

$$\begin{aligned} \rho_{q(\cdot), \mu}\left(\frac{f}{\lambda}\right) &= \int_0^\infty \left| \frac{f(t)}{\lambda} \right|^{q(t)} \mu(dt) = \int_0^\infty \left| \frac{t^{-1/q(t)} f(t)}{\lambda} \right|^{q(t)} dt \\ &= \rho_{q(\cdot)}\left(\frac{t^{-1/q(\cdot)} f}{\lambda}\right) \end{aligned}$$

Thus,

$$(1.19) \quad \|f\|_{q(\cdot), \mu} = \|t^{-1/q(\cdot)} f\|_{q(\cdot)}$$

Next, we present an useful technical results for the measure μ .

Lemma 1.4. For $q(\cdot) \in \mathcal{P}(\mathbb{R}^+, \mu)$

- i) For any $\alpha, \beta > 0$ and $q_+ < \infty$, $\|t^\alpha e^{-t\beta}\|_{q(\cdot), \mu} < \infty$.
- ii) For any $\alpha > 0$, $\|t^\alpha \chi_{(0,1]}\|_{q(\cdot), \mu} < \infty$.
- iii) For any $\alpha > 0$, $\|t^{-\alpha} \chi_{(1,\infty)}\|_{q(\cdot), \mu} < \infty$.
- iv) For any $t_0 > 0$, $(\ln 2)^{\frac{1}{q_-}} \leq \|\chi_{[t_0/2, t_0]}\|_{q(\cdot), \mu} \leq 1$.

Proof. Let us prove i). Set $f = t^\alpha e^{-t\beta}$

$$\rho_{q(\cdot), \mu}(f) = \int_0^\infty |f(t)|^{q(t)} \mu(dt) = \int_0^1 |t^\alpha e^{-t\beta}|^{q(t)} \frac{dt}{t} + \int_1^\infty |t^\alpha e^{-t\beta}|^{q(t)} \frac{dt}{t}$$

Now,

$$\int_0^1 |t^\alpha e^{-t\beta}|^{q(t)} \frac{dt}{t} = \int_0^1 t^{\alpha q(t)-1} e^{-t\beta q(t)} dt \leq \int_0^1 t^{\alpha-1} dt < \infty,$$

since $\alpha, \beta > 0$ and $0 \leq t \leq 1$. On the other hand, by making the change of variables $u = t\beta q_-$

$$\begin{aligned} \int_1^\infty |t^\alpha e^{-t\beta}|^{q(t)} \frac{dt}{t} &= \int_1^\infty t^{\alpha q(t)} e^{-t\beta q(t)} \frac{dt}{t} \\ &\leq \int_1^\infty t^{\alpha q_+} e^{-t\beta q_-} \frac{dt}{t} \leq \int_0^\infty t^{\alpha q_+} e^{-t\beta q_-} \frac{dt}{t} \\ &= \int_0^\infty \left(\frac{u}{\beta q_-}\right)^{\alpha q_+} e^{-u} \frac{du}{u} = \frac{1}{(\beta q_-)^{\alpha q_+}} \int_0^\infty u^{\alpha q_+-1} e^{-u} du \\ &= \frac{1}{(\beta q_-)^{\alpha q_+}} \Gamma(\alpha q_+) < \infty. \end{aligned}$$

since $\alpha, \beta > 0$ and $q_+ < \infty$. Thus, $\rho_{q(\cdot), \mu}(f) < \infty$, and therefore

$$\|t^\alpha e^{-t\beta}\|_{q(\cdot), \mu} < \infty.$$

The proof of *ii*) and *iii*) are immediate. Now, in order to prove *iv*), set $g = \chi_{[t_0/2, t_0]}$

$$\rho_{q(\cdot), \mu}(g) = \int_0^\infty |g(t)|^{q(t)} \mu(dt) = \int_{t_0/2}^{t_0} \frac{dt}{t} = \ln 2 < 1.$$

Then, $\lambda \geq 1$ implies $\rho_{q(\cdot), \mu}(\frac{g}{\lambda}) \leq \rho_{q(\cdot), \mu}(g) \leq 1$. Thus, $\|g\|_{q(\cdot), \mu} \leq 1$.

On the other hand, taking $0 < \lambda < 1$

$$\rho_{q(\cdot), \mu}(\frac{g}{\lambda}) = \int_{t_0/2}^{t_0} \lambda^{-q(t)} \frac{dt}{t} \geq \int_{t_0/2}^{t_0} \lambda^{-q_-} \frac{dt}{t} = \lambda^{-q_-} (\ln 2)$$

then $\lambda < (\ln 2)^{1/q_-}$ implies $\rho_{q(\cdot), \mu}(\frac{g}{\lambda}) > 1$. Therefore, $\rho_{q(\cdot), \mu}(\frac{g}{\lambda}) \leq 1$ implies $\lambda \geq (\ln 2)^{1/q_-}$ and then

$$\|g\|_{q(\cdot), \mu} \geq (\ln 2)^{1/q_-}.$$

□

In the case $\Omega = \mathbb{R}^+$, we denote $\mathcal{M}_{0, \infty}$ the set of all measurable functions $p(\cdot) : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ which satisfy the following conditions:

i) $0 \leq p_- \leq p_+ < \infty$,

ii) there exists $p(0) = \lim_{x \rightarrow 0} p(x)$ and $|p(x) - p(0)| \leq \frac{A}{\ln(1/x)}$, $0 < x \leq 1/2$

iii) there exists $p(\infty) = \lim_{x \rightarrow \infty} p(x)$ and $|p(x) - p(\infty)| \leq \frac{A}{\ln(x)}$, $x > 2$.

we denote $\mathcal{P}_{0, \infty}$ the subset of functions $p(\cdot)$ such that $p_- \geq 1$.

Let $\alpha(\cdot), \beta(\cdot) \in LH(\mathbb{R}^+)$, bounded with

$$(1.20) \quad \alpha(0) < \frac{1}{p'(0)}, \alpha(\infty) < \frac{1}{p'(\infty)}$$

and

$$(1.21) \quad \beta(0) > -\frac{1}{p(0)}, \beta(\infty) > -\frac{1}{p(\infty)}$$

Theorem 1.4. *Let $p(\cdot) \in \mathcal{P}_{0, \infty}$, $\alpha(\cdot), \beta(\cdot) \in LH(\mathbb{R}^+)$, bounded. Then the Hardy-type inequalities*

$$(1.22) \quad \left\| x^{\alpha(x)-1} \int_0^x \frac{f(y)}{y^{\alpha(y)}} dy \right\|_{p(\cdot)} \leq C_{\alpha(\cdot), p(\cdot)} \|f\|_{p(\cdot)}$$

$$(1.23) \quad \left\| x^{\beta(x)} \int_x^\infty \frac{f(y)}{y^{\beta(y)+1}} dy \right\|_{p(\cdot)} \leq C_{\beta(\cdot), p(\cdot)} \|f\|_{p(\cdot)}$$

are valid, if and only if, $\alpha(\cdot), \beta(\cdot)$ satisfy conditions (1.20) and (1.21)

Proof. For the proof see Theorem 3.1 and Remark 3.2 in [4].

□

As a consequence, we obtain the Hardy's inequalities associated to the exponent $q(\cdot) \in \mathcal{P}_{0, \infty}$ and the measure μ .

Corollary 1.2. *Let $q(\cdot) \in \mathcal{P}_{0,\infty}$ and $r > 0$, then*

$$(1.24) \quad \left\| t^{-r} \int_0^t g(y) dy \right\|_{q(\cdot), \mu} \leq C_{r,q(\cdot)} \|y^{-r+1} g\|_{q(\cdot), \mu}$$

and

$$(1.25) \quad \left\| t^r \int_t^\infty g(y) dy \right\|_{q(\cdot), \mu} \leq C_{r,q(\cdot)} \|y^{r+1} g\|_{q(\cdot), \mu}$$

Proof. Let $\alpha(t) = -r + \frac{1}{q'(t)} = -r + 1 - \frac{1}{q(t)}$, for any $t \in \mathbb{R}^+$, $f(y) = y^{\alpha(y)} g(y)$, for any $y \in \mathbb{R}^+$ then $\alpha(\cdot) \in LH(\mathbb{R}^+)$ and bounded, $\alpha(0) = -r + \frac{1}{q'(0)} < \frac{1}{q'(0)}$ and $\alpha(\infty) = -r + \frac{1}{q'(\infty)} < \frac{1}{q'(\infty)}$. Then, using (1.19) and (1.22)

$$\begin{aligned} \left\| t^{-r} \int_0^t g(y) dy \right\|_{q(\cdot), \mu} &= \left\| t^{-r - \frac{1}{q(t)}} \int_0^t g(y) dy \right\|_{q(\cdot)} = \left\| t^{\alpha(t)-1} \int_0^t g(y) dy \right\|_{q(\cdot)} \\ &\leq C_{r,q(\cdot)} \|y^{\alpha(y)} g\|_{q(\cdot)} = C_{r,q(\cdot)} \|y^{-r+1 - \frac{1}{q(y)}} g\|_{q(\cdot)} \\ &= C_{r,q(\cdot)} \|y^{-r+1} g\|_{q(\cdot), \mu}. \end{aligned}$$

On the other hand, by taking $\beta(t) = r - \frac{1}{q(t)}$, $\forall t \in \mathbb{R}^+$, $f(y) = y^{\beta(y)+1} g(y)$, $\forall y \in \mathbb{R}^+$ then $\beta(\cdot) \in LH(\mathbb{R}^+)$ and the proof of (1.25) is completely analogous. $\square$

In what follows we will consider only Lebesgue variable spaces with respect to the Gaussian measure γ_d , $L^{p(\cdot)}(\mathbb{R}^d, \gamma_d)$. The next condition was introduced by E. Dalmasso and R. Scotto in [2] and it is crucial to deal with the Gaussian measure.

Definition 1.5. *Let $p(\cdot) \in \mathcal{P}(\mathbb{R}^d, \gamma_d)$, we say that $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d)$ if there exist constants $C_{\gamma_d} > 0$ and $p_\infty \geq 1$ such that*

$$(1.26) \quad |p(x) - p_\infty| \leq \frac{C_{\gamma_d}}{|x|^2},$$

for $x \in \mathbb{R}^d \setminus \{(0, 0, \dots, 0)\}$.

Observation 1.3. *It can be proved that if $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d)$, then $p(\cdot) \in LH_\infty(\mathbb{R}^d)$.*

2. THE MAIN RESULTS

In this section we are going to define the variable Gaussian Besov-Lipschitz spaces and the variable Gaussian Triebel-Lizorkin spaces, which are the main goal of the paper.

The following two technical results are needed for defining variable Gaussian Besov-Lipschitz spaces.

Lemma 2.1. *Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $f \in L^{p(\cdot)}(\gamma_d)$, $\alpha \geq 0$ y k, l integers greater than α , then*

$$\left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \leq A_k t^{-k+\alpha} \text{ if and only if } \left\| \frac{\partial^l u(\cdot, t)}{\partial t^l} \right\|_{p(\cdot), \gamma_d} \leq A_l t^{-l+\alpha}.$$

Moreover, if $A_k(f), A_l(f)$ are the smallest constants in the inequalities above then there exist constants $A_{k,l,\alpha,p(\cdot)}$ and $D_{k,l,\alpha}$ such that

$$A_{k,l,\alpha,p(\cdot)} A_k(f) \leq A_l(f) \leq D_{k,l,\alpha} A_k(f),$$

for all $f \in L^{p(\cdot)}(\gamma_d)$.

Proof. Let us suppose without loss of generality that $k \geq l$. We start by proving the direct implication. For this we use the representation of the Poisson-Hermite semigroup (1.6), this is,

$$P_t f(x) = \int_0^{+\infty} T_s f(x) \mu_t^{(1/2)}(ds).$$

Then, by differentiating k -times with respect to t and by using the dominated convergence theorem, we get

$$\frac{\partial^k P_t f(x)}{\partial t^k} = \int_0^{+\infty} T_s f(x) \frac{\partial^k \mu_t^{(1/2)}}{\partial t^k}(ds).$$

By using Lemma 1.3, it's easy to prove that for all $m \in \mathbb{N}$

$$\lim_{t \rightarrow +\infty} \frac{\partial^m P_t f(x)}{\partial t^m} = 0.$$

Now, given $n \in \mathbb{N}, n > \alpha$

$$\begin{aligned} - \int_t^{+\infty} \frac{\partial^{n+1} P_s f(x)}{\partial s^{n+1}} ds &= - \lim_{s \rightarrow +\infty} \frac{\partial^n P_s f(x)}{\partial s^n} + \frac{\partial^n P_t f(x)}{\partial t^n} \\ &= \frac{\partial^n P_t f(x)}{\partial t^n}. \end{aligned}$$

Thus, for Minkowski's integral inequality (1.18)

$$\begin{aligned} \left\| \frac{\partial^n u(\cdot, t)}{\partial t^n} \right\|_{p(\cdot), \gamma_d} &\leq 4 \int_t^{+\infty} \left\| \frac{\partial^{n+1} u(\cdot, s)}{\partial s^{n+1}} \right\|_{p(\cdot), \gamma_d} ds \leq 4 \int_t^{+\infty} A_{n+1}(f) s^{-(n+1)+\alpha} ds \\ &= 4 \frac{A_{n+1}(f)}{n - \alpha} t^{-n+\alpha}. \end{aligned}$$

Therefore

$$A_n(f) \leq 4 \frac{A_{n+1}(f)}{n - \alpha},$$

and, since $n > \alpha$ is arbitrary, then, by using the above result $k - l$ times, we obtain

$$\begin{aligned} A_l(f) &\leq 4 \frac{A_{l+1}(f)}{l - \alpha} \leq 4^2 \frac{A_{l+2}(f)}{(l - \alpha)(l + 1 - \alpha)} \\ &\leq \dots \leq 4^{k-l} \frac{A_k(f)}{(l - \alpha)(l + 1 - \alpha) \dots (k - 1 - \alpha)} = D_{k,l,\alpha} A_k(f). \end{aligned}$$

To prove the converse, we use again the representation (1.6) and we obtain that

$$u(x, t_1 + t_2) = P_{t_1}(P_{t_2} f)(x) = \int_0^{+\infty} T_s(P_{t_2} f)(x) \mu_{t_1}^{\frac{1}{2}}(ds).$$

Thus, taking $t = t_1 + t_2$ and differentiating l times with respect to t_2 and $k - l$ times with respect to t_1 , we get

$$(2.1) \quad \frac{\partial^k u(x, t)}{\partial t^k} = \int_0^{+\infty} T_s \left(\frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right) \frac{\partial^{k-l} \mu_{t_1}^{\frac{1}{2}}}{\partial t_1^{k-l}}(ds).$$

Then, by Corollary 1.1, Minkowski's integral inequality (1.18) and the $L^{p(\cdot)}$ -boundedness of the Ornstein-Uhlenbeck semigroup (see [5]), we get

$$\begin{aligned} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} &\leq 4 \int_0^{+\infty} \left\| T_s \left(\frac{\partial^l P_{t_2} f}{\partial t_2^l} \right) \right\|_{p(\cdot), \gamma_d} \left| \frac{\partial^{k-l} \mu_{t_1}^{\frac{1}{2}}}{\partial t_1^{k-l}}(ds) \right| \\ &\leq 4C_{p(\cdot)} \left\| \frac{\partial^l P_{t_2} f}{\partial t_2^l} \right\|_{p(\cdot), \gamma_d} \int_0^{+\infty} \left| \frac{\partial^{k-l} \mu_{t_1}^{\frac{1}{2}}}{\partial t_1^{k-l}}(ds) \right| \\ &\leq 4C_{p(\cdot)} \left\| \frac{\partial^l}{\partial t_2^l} P_{t_2} f \right\|_{p(\cdot), \gamma_d} C_{k-l} t_1^{l-k} \leq 4C_{p(\cdot)} A_l(f) C_{k-l} t_2^{-l+\alpha} t_1^{l-k}. \end{aligned}$$

Therefore, taking $t_1 = t_2 = \frac{t}{2}$,

$$\left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \leq 4C_{p(\cdot)} A_l(f) C_{k-l} \left(\frac{t}{2} \right)^{-k+\alpha}.$$

Thus

$$A_k(f) \leq 4C_{p(\cdot)} \frac{C_{k-l}}{2^{-k+\alpha}} A_l(f).$$

□

Lemma 2.2. *Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $q(\cdot) \in \mathcal{P}_{0,\infty}$. Let $\alpha \geq 0$ and k, l integers greater than α . Then*

$$\left\| t^{k-\alpha} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} < \infty$$

if and only if

$$\left\| t^{l-\alpha} \left\| \frac{\partial^l u(\cdot, t)}{\partial t^l} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} < \infty.$$

Moreover, there exist constants $A_{k,l,\alpha,p(\cdot)}$ and $D_{k,l,\alpha,q(\cdot)}$ such that

$$\begin{aligned} D_{k,l,\alpha,q(\cdot)} \left\| t^{l-\alpha} \left\| \frac{\partial^l u(\cdot, t)}{\partial t^l} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} &\leq \left\| t^{k-\alpha} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \\ &\leq A_{k,l,\alpha,p(\cdot)} \left\| t^{l-\alpha} \left\| \frac{\partial^l u(\cdot, t)}{\partial t^l} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu}, \end{aligned}$$

for all $f \in L^{p(\cdot)}(\gamma_d)$.

Proof. Suppose without loss of generality that $k \geq l$. We prove first the converse implication; by proceeding as in lemma 2.1, taking $t_1 = t_2 = \frac{t}{2}$, we have

$$\begin{aligned} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} &\leq 4C_{p(\cdot)} \left\| \frac{\partial^l P_{t_2} f}{\partial t_2^l} \right\|_{p(\cdot), \gamma_d} C_{k-l} t_1^{l-k}, \\ &= 4C_{p(\cdot)} \left\| \frac{\partial^l P_{\frac{t}{2}} f}{\partial (\frac{t}{2})^l} \right\|_{p(\cdot), \gamma_d} C_{k-l} \left(\frac{t}{2}\right)^{l-k}. \end{aligned}$$

Thus

$$\begin{aligned} \left\| t^{k-\alpha} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} &\leq 4C_{p(\cdot)} \frac{C_{k-l}}{2^{l-k}} \left\| t^{l-\alpha} \left\| \frac{\partial^l u(\cdot, \frac{t}{2})}{\partial (\frac{t}{2})^l} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \\ &= A_{k,l,\alpha,p(\cdot)} \left\| s^{l-\alpha} \left\| \frac{\partial^l u(\cdot, s)}{\partial s^l} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu}. \end{aligned}$$

with $A_{k,l,\alpha,p(\cdot)} = 4C_{p(\cdot)} C_{k-l} 2^{k-\alpha}$.

For the direct implication, given $n \in \mathbb{N}$, $n > \alpha$, again, as in the above lemma

$$\left\| \frac{\partial^n u(\cdot, t)}{\partial t^n} \right\|_{p(\cdot), \gamma_d} \leq 4 \int_t^{+\infty} \left\| \frac{\partial^{n+1} u(\cdot, s)}{\partial s^{n+1}} \right\|_{p(\cdot), \gamma_d} ds.$$

Therefore, by the Hardy's inequality (1.25)

$$\begin{aligned} \left\| t^{n-\alpha} \left\| \frac{\partial^n u(\cdot, t)}{\partial t^n} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} &\leq 4 \left\| t^{n-\alpha} \int_t^{+\infty} \left\| \frac{\partial^{n+1} u(\cdot, s)}{\partial s^{n+1}} \right\|_{p(\cdot), \gamma_d} ds \right\|_{q(\cdot), \mu} \\ &\leq 4C_{n,\alpha,q(\cdot)} \left\| s^{n+1-\alpha} \left\| \frac{\partial^{n+1} u(\cdot, s)}{\partial s^{n+1}} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \end{aligned}$$

Now, since $n > \alpha$ is arbitrary, by using the previous result $k - l$ times, we obtain

$$\begin{aligned} \left\| t^{l-\alpha} \left\| \frac{\partial^l u(\cdot, t)}{\partial t^l} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} &\leq 4C_{l,\alpha,q(\cdot)} \left\| t^{l+1-\alpha} \left\| \frac{\partial^{l+1} u(\cdot, t)}{\partial t^{l+1}} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \\ &\leq 4^2 C_{l,\alpha,q(\cdot)} C_{l+1,\alpha,q(\cdot)} \left\| t^{l+2-\alpha} \left\| \frac{\partial^{l+2} u(\cdot, t)}{\partial t^{l+2}} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \\ &\vdots \\ &\leq D_{k,l,\alpha,q(\cdot)} \left\| t^{k-\alpha} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \end{aligned}$$

where $D_{k,l,\alpha,q(\cdot)} = 4^{k-l} C_{l,\alpha,q(\cdot)} \cdots C_{k-1,\alpha,q(\cdot)}$. $\square$

The next technical result will be the key to define the variable Gaussian Triebel-Lizorkin spaces.

Lemma 2.3. *Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $q(\cdot) \in \mathcal{P}_{0,\infty}$. Let $\alpha \geq 0$ and k, l integers greater than α . Then*

$$\left\| \left\| t^{k-\alpha} \left| \frac{\partial^k}{\partial t^k} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} < \infty$$

if and only if

$$\left\| \left\| t^{l-\alpha} \left| \frac{\partial^l}{\partial t^l} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} < \infty.$$

Moreover, there exist constants $A_{k,l,\alpha,p(\cdot)}, D_{k,l,\alpha,q(\cdot)}$ such that

$$\begin{aligned} D_{k,l,\alpha,q(\cdot)} \left\| \left\| t^{l-\alpha} \left| \frac{\partial^l}{\partial t^l} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} &\leq \left\| \left\| t^{k-\alpha} \left| \frac{\partial^k}{\partial t^k} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} \\ &\leq A_{k,l,\alpha,p(\cdot)} \left\| \left\| t^{l-\alpha} \left| \frac{\partial^l}{\partial t^l} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d}, \end{aligned}$$

for all $f \in L^{p(\cdot)}(\gamma_d)$.

Proof. Suppose without loss of generality that $k \geq l$. Let $n \in \mathbb{N}$ such that $n > \alpha$, we can prove that

$$\left| \frac{\partial^n}{\partial t^n} P_t f(x) \right| \leq \int_t^{+\infty} \left| \frac{\partial^{n+1}}{\partial s^{n+1}} P_s f(x) \right| ds.$$

Then, by the Hardy's inequality (1.25),

$$\begin{aligned} \left\| t^{n-\alpha} \left| \frac{\partial^n}{\partial t^n} P_t f(x) \right| \right\|_{q(\cdot), \mu} &\leq \left\| t^{n-\alpha} \int_t^{+\infty} \left| \frac{\partial^{n+1}}{\partial s^{n+1}} P_s f(x) \right| ds \right\|_{q(\cdot), \mu} \\ &\leq C_{n,\alpha,q(\cdot)} \left\| s^{n+1-\alpha} \left| \frac{\partial^{n+1}}{\partial s^{n+1}} P_s f(x) \right| \right\|_{q(\cdot), \mu} \end{aligned}$$

Now, since $n > \alpha$ is arbitrary, by iterating the previous argument $k - l$ times, we obtain

$$\begin{aligned} \left\| t^{l-\alpha} \left| \frac{\partial^l}{\partial t^l} P_t f(x) \right| \right\|_{q(\cdot), \mu} &\leq C_{l,\alpha,q(\cdot)} \left\| t^{l+1-\alpha} \left| \frac{\partial^{l+1}}{\partial t^{l+1}} P_t f(x) \right| \right\|_{q(\cdot), \mu} \\ &\leq C_{l,\alpha,q(\cdot)} C_{l+1,\alpha,q(\cdot)} \left\| t^{l+2-\alpha} \left| \frac{\partial^{l+2}}{\partial t^{l+2}} P_t f(x) \right| \right\|_{q(\cdot), \mu} \\ &\vdots \\ &\leq C_{k,l,\alpha,q(\cdot)} \left\| t^{k-\alpha} \left| \frac{\partial^k}{\partial t^k} P_t f(x) \right| \right\|_{q(\cdot), \mu} \end{aligned}$$

where $C_{k,l,\alpha,q(\cdot)} = C_{l,\alpha,q(\cdot)} C_{l+1,\alpha,q(\cdot)} \cdots C_{k-1,\alpha,q(\cdot)}$. Thus,

$$D_{k,l,\alpha,q(\cdot)} \left\| \left\| t^{l-\alpha} \left| \frac{\partial^l}{\partial t^l} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} \leq \left\| \left\| t^{k-\alpha} \left| \frac{\partial^k}{\partial t^k} P_t f \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d},$$

where $D_{k,l,\alpha,q(\cdot)} = 1/C_{k,l,\alpha,q(\cdot)}$.

The other inequality is obtain from the case $k = l + 1$ by an inductive argument. Let $t_1, t_2 > 0$ and take $t = t_1 + t_2$, from (2.1) we get

$$\frac{\partial^k u(x, t)}{\partial t^k} = \int_0^{+\infty} T_s \left(\frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right) \frac{\partial^{k-l}}{\partial t_1^{k-l}} \mu_{t_1}^{(1/2)}(ds),$$

and since, $\frac{\partial}{\partial t_1} \mu_{t_1}^{(1/2)}(ds) = (t_1^{-1} - \frac{t_1}{2s}) \mu_{t_1}^{(1/2)}(ds)$ we obtain

$$\begin{aligned} \left| \frac{\partial^k u(x, t)}{\partial t^k} \right| &\leq \int_0^{+\infty} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) |t_1^{-1} - \frac{t_1}{2s}| \mu_{t_1}^{(1/2)}(ds) \\ &\leq t_1^{-1} \int_0^{+\infty} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) \mu_{t_1}^{(1/2)}(ds) \\ &\quad + \frac{t_1}{2} \int_0^{+\infty} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) \frac{1}{s} \mu_{t_1}^{(1/2)}(ds). \end{aligned}$$

Therefore

$$\begin{aligned} \left\| t_2^{k-\alpha} \left| \frac{\partial^k u(x, t)}{\partial t^k} \right| \right\|_{q(\cdot), \mu} &\leq \left\| t_2^{k-\alpha} t_1^{-1} \int_0^{+\infty} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) \mu_{t_1}^{(1/2)}(ds) \right\|_{q(\cdot), \mu} \\ &\quad + \left\| t_2^{k-\alpha} \frac{t_1}{2} \int_0^{+\infty} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) \frac{1}{s} \mu_{t_1}^{(1/2)}(ds) \right\|_{q(\cdot), \mu}. \\ &= (I) + (II). \end{aligned}$$

Now, by using Minkowski's integral inequality twice (1.18) (since T_s is an integral transformation with positive kernel) and the fact that $\mu_{t_1}^{(1/2)}(ds)$ is a probability measure, we get

$$\begin{aligned} (I) &= \left\| (t_2^{k-\alpha} t_1^{-1} \int_0^{+\infty} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) \mu_{t_1}^{(1/2)}(ds) \right\|_{q(\cdot), \mu} \\ &\leq 4 \int_0^{+\infty} \left\| t_2^{k-\alpha} t_1^{-1} T_s \left(\left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right) \right\|_{q(\cdot), \mu} \mu_{t_1}^{(1/2)}(ds) \\ &\leq 16 \int_0^{+\infty} T_s \left(\left\| t_2^{k-\alpha} t_1^{-1} \left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right\|_{q(\cdot), \mu} \right) \mu_{t_1}^{(1/2)}(ds) \\ &\leq 16 T^* \left(\left\| t_2^{k-\alpha} t_1^{-1} \left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right\|_{q(\cdot), \mu} \right). \end{aligned}$$

For (II) we proceed in analogous way, and by using Lemma 1.2 we get

$$\begin{aligned} (II) &\leq \frac{16}{2} T^* \left(\left\| t_2^{k-\alpha} t_1 \left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right\|_{q(\cdot), \mu} \right) \int_0^{+\infty} \frac{1}{s} \mu_{t_1}^{(1/2)}(ds) \\ &= 8 T^* \left(\left\| t_2^{k-\alpha} t_1 \left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right\|_{q(\cdot), \mu} \right) C_1 \frac{1}{t_1^2}. \end{aligned}$$

Now, since T^* is defined as a supremum, we get

$$(II) \leq 8 C_1 T^* \left(\left\| t_2^{k-\alpha} t_1^{-1} \left| \frac{\partial^l P_{t_2} f(x)}{\partial t_2^l} \right| \right\|_{q(\cdot), \mu} \right).$$

Then, taking $t_1 = t_2 = \frac{t}{2}$ and the change of variable $s = \frac{t}{2}$, we have

$$(I) \leq 16 T^* \left(\left\| s^{l-\alpha} \left| \frac{\partial^l P_s f(x)}{\partial s^l} \right| \right\|_{q(\cdot), \mu} \right)$$

and

$$(II) \leq 8 C_1 T^* \left(\left\| s^{l-\alpha} \left| \frac{\partial^l P_s f(x)}{\partial s^l} \right| \right\|_{q(\cdot), \mu} \right)$$

Therefore, by the $L^{p(\cdot)}(\gamma_d)$ -boundedness of T^* (see [5]),

$$\begin{aligned} \left\| \left\| t^{k-\alpha} \left| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} &\leq 2^{k-\alpha} 16 \left\| T^* \left(\left\| s^{l-\alpha} \left| \frac{\partial^l P_s f}{\partial s^l} \right| \right\|_{q(\cdot), \mu} \right) \right\|_{p(\cdot), \gamma_d} \\ &\quad + 2^{k-\alpha} 8 C_1 \left\| T^* \left(\left\| s^{l-\alpha} \left| \frac{\partial^l P_s f}{\partial s^l} \right| \right\|_{q(\cdot), \mu} \right) \right\|_{p(\cdot), \gamma_d} \\ &\leq 2^{k-\alpha} C_{p(\cdot)} (16 + 8 C_1) \left\| \left\| s^{l-\alpha} \left| \frac{\partial^l P_s f}{\partial s^l} \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d}. \end{aligned}$$

□

Next, we need the following technical result for the $L^{p(\cdot)}(\gamma_d)$ -norms of the derivatives of the Poisson-Hermite semigroup:

Lemma 2.4. *Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$. Suppose that $f \in L^{p(\cdot)}(\gamma_d)$, then for any integer k , $\left\| \frac{\partial^k}{\partial t^k} P_t f \right\|_{p(\cdot), \gamma_d} \leq C_{p(\cdot)} \left\| \frac{\partial^k}{\partial s^k} P_s f \right\|_{p(\cdot), \gamma_d}$, for whatever $0 < s < t < +\infty$. Moreover,*

$$(2.2) \quad \left\| \frac{\partial^k}{\partial t^k} P_t f \right\|_{p(\cdot), \gamma_d} \leq \frac{C_{k, p(\cdot)}}{t^k} \|f\|_{p(\cdot), \gamma_d}, \quad t > 0.$$

Proof. First, let us consider the case $k = 0$. Fixed $t_1, t_2 > 0$, by using the semigroup property of $\{P_t\}$, we get

$$P_{t_1+t_2} f(x) = P_{t_1}(P_{t_2} f(x))$$

Thus, by the $L^{p(\cdot)}$ -boundedness of $\{P_t\}$ (see [5]),

$$\|P_{t_1+t_2}f\|_{p(\cdot),\gamma_d} \leq C_{p(\cdot)}\|P_{t_2}f\|_{p(\cdot),\gamma_d}.$$

In order to prove the general case, $k > 0$, using the dominated convergence theorem and differentiating the identity $u(x, t_1 + t_2) = P_{t_1}(u(x, t_2))$ k -times with respect to t_2 we obtain

$$\frac{\partial^k u(x, t_1 + t_2)}{\partial(t_1 + t_2)^k} = P_{t_1} \left(\frac{\partial^k u(x, t_2)}{\partial t_2^k} \right)$$

and then we proceed as in the previous argument. In other to prove (2.2) we use again the representation (1.6) of the Poisson-Hermite semigroup and differentiating k -times with respect to t to obtain

$$\frac{\partial^k}{\partial t^k} u(x, t) = \int_0^{+\infty} T_s f(x) \frac{\partial^k}{\partial t^k} \mu_t^{(1/2)}(ds).$$

Thus, by the Minkowski's integral inequality, the $L^{p(\cdot)}$ -boundedness of the Ornstein-Uhlenbeck semigroup (see [5]) and the Corollary 1.1, for $t > 0$

$$\begin{aligned} \left\| \frac{\partial^k u(\cdot, t)}{\partial t^k} \right\|_{p(\cdot),\gamma_d} &\leq 4 \int_0^{+\infty} \left\| T_s f \frac{\partial^k \mu_t^{(1/2)}}{\partial t^k}(ds) \right\|_{p(\cdot),\gamma_d} \\ &= 4 \int_0^{+\infty} \|T_s f\|_{p(\cdot),\gamma_d} \left| \frac{\partial^k \mu_t^{(1/2)}}{\partial t^k}(ds) \right| \\ &\leq 4C_{p(\cdot)}\|f\|_{p(\cdot),\gamma_d} \int_0^{+\infty} \left| \frac{\partial^k \mu_t^{(1/2)}}{\partial t^k}(ds) \right| \leq \frac{C_{k,p(\cdot)}}{t^k} \|f\|_{p(\cdot),\gamma_d}. \end{aligned}$$

□

The Lipschitz spaces can be generalized of the following way (see, for example [6], [7],[8],[9]), using the Poisson-Hermite semigroup.

We are ready to define the variable Gaussian Besov-Lipschitz spaces $B_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$, also called Gaussian Besov-Lipschitz spaces with variable exponents or variable Besov-Lipschitz spaces for expansions in Hermite polynomials.

Definition 2.1. Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $q(\cdot) \in \mathcal{P}_{0,\infty}$. Let $\alpha \geq 0$, k the smallest integer greater than α . The variable Gaussian Besov-Lipschitz space $B_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$ is defined as the set of functions $f \in L^{p(\cdot)}(\gamma_d)$ such that

$$(2.3) \quad \left\| t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \right\|_{q(\cdot),\mu} < \infty.$$

The norm of $f \in B_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$ is defined as

$$(2.4) \quad \|f\|_{B_{p(\cdot),q(\cdot)}^\alpha} := \|f\|_{p(\cdot),\gamma_d} + \left\| t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \right\|_{q(\cdot),\mu}.$$

The variable Gaussian Besov-Lipschitz space $B_{p(\cdot),\infty}^\alpha(\gamma_d)$ is defined as the set of functions $f \in L^{p(\cdot)}(\gamma_d)$ for which there exists a constant A such that

$$\left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \leq A t^{-k+\alpha}$$

and then the norm of $f \in B_{p(\cdot),\infty}^\alpha(\gamma_d)$ is defined as

$$(2.5) \quad \|f\|_{B_{p(\cdot),\infty}^\alpha} := \|f\|_{p(\cdot),\gamma_d} + A_k(f),$$

where $A_k(f)$ is the smallest constant A in the above inequality.

Lemmas 2.1 and 2.2 show that we could have replaced k with any other integer l greater than α and the resulting norms are equivalents.

Now, let us study some inclusion relations between variable Gaussian Besov-Lipschitz spaces. The next result is analogous to Proposition 10, page 153 in [7] (see also [6] or Proposition 7.36 in [10]).

Proposition 2.1. *Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $q_1(\cdot), q_2(\cdot) \in \mathcal{P}_{0,\infty}$. The inclusion $B_{p(\cdot),q_1(\cdot)}^{\alpha_1}(\gamma_d) \subset B_{p(\cdot),q_2(\cdot)}^{\alpha_2}(\gamma_d)$ holds if:*

- i) $\alpha_1 > \alpha_2 > 0$ ($q_1(\cdot)$ y $q_2(\cdot)$ not need to be related), or
- ii) If $\alpha_1 = \alpha_2$ and $q_1(t) \leq q_2(t)$ a.e.

Proof. To prove part ii), let us take α the common value of α_1 and α_2 .

Let $f \in B_{p(\cdot),q_1(\cdot)}^\alpha$ and set $A = \left\| t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \right\|_{q_1(\cdot),\mu}$.

Fixed $t_0 > 0$

$$\left\| \chi_{[\frac{t_0}{2}, t_0]} t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \right\|_{q_1(\cdot),\mu} \leq A.$$

However, by Lemma 2.4,

$$\left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot),\gamma_d} \leq C_{p(\cdot)} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \quad t \in [\frac{t_0}{2}, t_0].$$

Thus, we obtain

$$\begin{aligned} \left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot),\gamma_d} \left\| \chi_{[\frac{t_0}{2}, t_0]} t^{k-\alpha} \right\|_{q_1(\cdot),\mu} &\leq C_{p(\cdot)} \left\| \chi_{[\frac{t_0}{2}, t_0]} t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot),\gamma_d} \right\|_{q_1(\cdot),\mu} \\ &\leq C_{p(\cdot)} A. \end{aligned}$$

Therefore,

$$\begin{aligned} \left(\frac{t_0}{2} \right)^{k-\alpha} \left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot),\gamma_d} \left\| \chi_{[\frac{t_0}{2}, t_0]} \right\|_{q_1(\cdot),\mu} &\leq \left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot),\gamma_d} \left\| \chi_{[\frac{t_0}{2}, t_0]} t^{k-\alpha} \right\|_{q_1(\cdot),\mu} \\ &\leq C_{p(\cdot)} A \end{aligned}$$

and by Lemma 1.4

$$\begin{aligned} \left(\frac{t_0}{2}\right)^{k-\alpha} \left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot), \gamma_d} (\ln 2)^{1/q_1^-} &\leq \left(\frac{t_0}{2}\right)^{k-\alpha} \left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot), \gamma_d} \left\| \chi_{[\frac{t_0}{2}, t_0]} \right\|_{q_1(\cdot), \mu} \\ &\leq C_{p(\cdot)} A \end{aligned}$$

Then,

$$\left\| \frac{\partial^k P_{t_0} f}{\partial t_0^k} \right\|_{p(\cdot), \gamma_d} \leq \frac{C_{p(\cdot)} 2^{k-\alpha}}{(\ln 2)^{1/q_1^-}} A t_0^{-k+\alpha},$$

and since t_0 is arbitrary

$$\left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \leq C_{k, \alpha, p(\cdot) q_1(\cdot)} A t^{-k+\alpha},$$

for all $t > 0$. In other words, $f \in B_{p(\cdot), q_1(\cdot)}^\alpha$ implies that $f \in B_{p(\cdot), \infty}^\alpha$.

Now, let us take $g(t) = t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d}$, then $\rho_{q_1(\cdot), \mu}(g) < \infty$, since, $f \in B_{p(\cdot), q_1(\cdot)}^\alpha$.

Thus, as $q_2(t) \geq q_1(t)$ a.e.

$$\begin{aligned} \rho_{q_2(\cdot), \mu}(g) &= \int_0^{+\infty} \left(t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right)^{q_2(t)} \frac{dt}{t} \\ &= \int_0^{+\infty} \left(t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right)^{q_2(t)-q_1(t)} \left(t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right)^{q_1(t)} \frac{dt}{t} \\ &\leq (C_{k, \alpha, p(\cdot) q_1(\cdot)} A)^{q_2^+ - q_1^-} \int_0^{+\infty} \left(t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right)^{q_1(t)} \frac{dt}{t} \\ &= (C_{k, \alpha, p(\cdot) q_1(\cdot)} A)^{q_2^+ - q_1^-} \rho_{q_1(\cdot), \mu}(g) < +\infty. \end{aligned}$$

Hence, $f \in B_{p(\cdot), q_2(\cdot)}^\alpha$. In order to prove part i), by Lemma 2.4, we obtain

$$\left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \leq C_{k, p(\cdot)} t^{-k}, \quad t > 0.$$

Now, given $f \in B_{p(\cdot), q_1(\cdot)}^{\alpha_1}$, again by setting

$$A = \left\| t^{k-\alpha_1} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_1(\cdot), \mu},$$

we obtain, as in part ii),

$$\left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \leq C_{k, \alpha_1, p(\cdot) q_1(\cdot)} A t^{-k+\alpha_1},$$

for all $t > 0$. Therefore,

$$\begin{aligned} \left\| t^{k-\alpha_2} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_2(\cdot), \mu} &\leq \left\| \chi_{(0,1]} t^{k-\alpha_2} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_2(\cdot), \mu} \\ &\quad + \left\| \chi_{(1,\infty)} t^{k-\alpha_2} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_2(\cdot), \mu} \\ &= (I) + (II). \end{aligned}$$

Now, again by Lemma 1.4 we get,

$$\begin{aligned} (I) &= \left\| \chi_{(0,1]} t^{k-\alpha_2} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_2(\cdot), \mu} \leq \left\| \chi_{(0,1]} t^{k-\alpha_2} C_{k, \alpha_1, p(\cdot)q_1(\cdot)} A t^{-k+\alpha_1} \right\|_{q_2(\cdot), \mu} \\ &= C_{k, \alpha_1, p(\cdot)q_1(\cdot)} A \left\| \chi_{(0,1]} t^{\alpha_1-\alpha_2} \right\|_{q_2(\cdot), \mu} < \infty, \end{aligned}$$

and also by Lemma 1.4,

$$\begin{aligned} (II) &= \left\| \chi_{(1,\infty)} t^{k-\alpha_2} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_2(\cdot), \mu} \leq \left\| \chi_{(1,\infty)} t^{k-\alpha_2} C_{k, p(\cdot)} t^{-k} \right\|_{q_2(\cdot), \mu} \\ &= C_{k, p(\cdot)} \left\| \chi_{(1,\infty)} t^{-\alpha_2} \right\|_{q_2(\cdot), \mu} < \infty. \end{aligned}$$

Hence,

$$\left\| t^{k-\alpha_2} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q_2(\cdot), \mu} < +\infty,$$

and then $f \in B_{p(\cdot), q_2(\cdot)}^{\alpha_2}$. $\square$

Let us now define the variable Gaussian Triebel-Lizorkin spaces $F_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$, which represent another way to measure regularity of functions, proceeding as in [6], [8] or [9].

Definition 2.2. Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $q(\cdot) \in \mathcal{P}_{0, \infty}$. Let $\alpha \geq 0$ and k the smallest integer greater than α . The variable Gaussian Triebel-Lizorkin space $F_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$ is the set of functions $f \in L^{p(\cdot)}(\gamma_d)$ such that

$$(2.6) \quad \left\| \left\| t^{k-\alpha} \frac{\partial^k P_t f}{\partial t^k} \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} < \infty,$$

The norm of $f \in F_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$ is defined as

$$(2.7) \quad \|f\|_{F_{p(\cdot), q(\cdot)}^\alpha} := \|f\|_{p(\cdot), \gamma_d} + \left\| \left\| t^{k-\alpha} \frac{\partial^k P_t f}{\partial t^k} \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d}.$$

By Lemma 2.3, the definition of $F_{p(\cdot), q(\cdot)}^\alpha$ is independent of the integer $k > \alpha$ chosen and the resulting norms are equivalents.

Observation 2.1. *The variable Gaussian Besov-Lipschitz and variable Gaussian Triebel-Lizorkin spaces are, by construction, subspaces of $L^{p(\cdot)}(\gamma_d)$. Moreover, since trivially $\|f\|_{p(\cdot),\gamma_d} \leq \|f\|_{B_{p(\cdot),q(\cdot)}^\alpha}$ and $\|f\|_{p(\cdot),\gamma_d} \leq \|f\|_{F_{p(\cdot),q(\cdot)}^\alpha}$, the inclusions are continuous. On the other hand, from (1.9) it is clear that for all $t > 0$ and $k \in \mathbb{N}$,*

$$\frac{\partial^k}{\partial t^k} P_t h_\beta(x) = (-1)^k |\beta|^{k/2} e^{-t\sqrt{|\beta|}} h_\beta(x),$$

and again by Lemma 1.4,

$$\begin{aligned} \left\| t^{k-\alpha} \left\| \frac{\partial^k}{\partial t^k} P_t h_\beta \right\|_{p(\cdot),\gamma_d} \right\|_{q(\cdot),\mu} &= \left\| t^{k-\alpha} \left\| (-|\beta|^{1/2})^k e^{-t\sqrt{|\beta|}} h_\beta \right\|_{p(\cdot),\gamma_d} \right\|_{q(\cdot),\mu} \\ &= |\beta|^{k/2} \|h_\beta\|_{p(\cdot),\gamma_d} \left\| t^{k-\alpha} e^{-t\sqrt{|\beta|}} \right\|_{q(\cdot),\mu} \\ &= C_{k,\alpha,\beta,q(\cdot)} \|h_\beta\|_{p(\cdot),\gamma_d} < \infty. \end{aligned}$$

Thus, $h_\beta \in B_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$ and

$$\|h_\beta\|_{B_{p(\cdot),q(\cdot)}^\alpha} = (1 + C_{k,\alpha,\beta,q(\cdot)}) \|h_\beta\|_{p(\cdot),\gamma_d}.$$

In a similar way, $h_\beta \in F_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$ and

$$\begin{aligned} \|h_\beta\|_{F_{p(\cdot),q(\cdot)}^\alpha} &= \|h_\beta\|_{p(\cdot),\gamma_d} + \left\| \left\| t^{k-\alpha} \left\| \frac{\partial^k}{\partial t^k} P_t h_\beta \right\|_{q(\cdot),\mu} \right\|_{p(\cdot),\gamma_d} \right\| \\ &= \|h_\beta\|_{p(\cdot),\gamma_d} + |\beta|^{k/2} \left\| t^{k-\alpha} e^{-t\sqrt{|\beta|}} \right\|_{q(\cdot),\mu} \|h_\beta\|_{p(\cdot),\gamma_d} \\ &= (1 + C_{k,\alpha,\beta,q(\cdot)}) \|h_\beta\|_{p(\cdot),\gamma_d} = \|h_\beta\|_{B_{p(\cdot),q(\cdot)}^\alpha}. \end{aligned}$$

Hence, the polinomials $\mathcal{P}$ is contained in $B_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$ and in $F_{p(\cdot),q(\cdot)}^\alpha(\gamma_d)$.

Also, we have an inclusion result for the variable Gaussian Triebel-Lizorkin spaces, which is analogous to Proposition 2.1, see also [6] or Proposition 7.40 in [10].

Proposition 2.2. *Let $p(\cdot) \in \mathcal{P}_{\gamma_d}^\infty(\mathbb{R}^d) \cap LH_0(\mathbb{R}^d)$ and $q_1(\cdot), q_2(\cdot) \in \mathcal{P}_{0,\infty}$. The inclusion $F_{p(\cdot),q_1(\cdot)}^{\alpha_1}(\gamma_d) \subset F_{p(\cdot),q_2(\cdot)}^{\alpha_2}(\gamma_d)$ holds for $\alpha_1 > \alpha_2 > 0$ and $q_1(t) > q_2(t)$ a.e.*

Proof. Let us consider $f \in F_{p(\cdot),q_1(\cdot)}^{\alpha_1}$, then

$$\begin{aligned} \left\| t^{k-\alpha_2} \left\| \frac{\partial^k P_t f(x)}{\partial t^k} \right\|_{q_2(\cdot),\mu} \right\| &\leq \left\| t^{k-\alpha_2} \left\| \frac{\partial^k P_t f(x)}{\partial t^k} \right\|_{\chi_{(0,1]}} \right\|_{q_2(\cdot),\mu} \\ &\quad + \left\| t^{k-\alpha_2} \left\| \frac{\partial^k P_t f(x)}{\partial t^k} \right\|_{\chi_{(1,\infty)}} \right\|_{q_2(\cdot),\mu} \\ &= (I) + (II). \end{aligned}$$

Now, since $q_1(t) > q_2(t)$ a.e., by taking $r(t) = \frac{q_1(t)q_2(t)}{q_1(t) - q_2(t)}$, we obtain that $r(\cdot) \geq 1$ and $\frac{1}{r(\cdot)} + \frac{1}{q_1(\cdot)} = \frac{1}{q_2(\cdot)}$, thus, by Hölder's inequality (1.17) and Lemma 1.4

$$\begin{aligned} (I) &= \left\| t^{\alpha_1 - \alpha_2} \chi_{(0,1]} t^{k - \alpha_1} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right\|_{q_2(\cdot), \mu} \\ &\leq 2 \left\| t^{\alpha_1 - \alpha_2} \chi_{(0,1]} \right\|_{r(\cdot), \mu} \left\| t^{k - \alpha_1} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu} \\ &= C_{\alpha_1, \alpha_2, q_1(\cdot), q_2(\cdot)} \left\| t^{k - \alpha_1} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu}. \end{aligned}$$

Now, for the second term (II), by using Lemmas 1.4 and 1.3, we get

$$\begin{aligned} (II) &= \left\| t^{k - \alpha_2} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \chi_{(1, \infty)} \right\|_{q_2(\cdot), \mu} \leq C_k T^* f(x) \left\| \chi_{(1, \infty)} t^{k - \alpha_2} t^{-k} \right\|_{q_2(\cdot), \mu} \\ &= C_k T^* f(x) \left\| \chi_{(1, \infty)} t^{-\alpha_2} \right\|_{q_2(\cdot), \mu} = C_{k, \alpha_2, q_2(\cdot)} T^* f(x). \end{aligned}$$

Then, by using the $L^{p(\cdot)}(\gamma_d)$ boundedness of T^* (see [5]),

$$\begin{aligned} \left\| \left\| t^{k - \alpha_2} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_2(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} &\leq C_{\alpha_1, \alpha_2, q_1(\cdot), q_2(\cdot)} \left\| \left\| t^{k - \alpha_1} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} \\ &\quad + C_{k, \alpha_2, q_2(\cdot)} \|T^* f\|_{p(\cdot), \gamma_d} \\ &\leq C_{\alpha_1, \alpha_2, q_1(\cdot), q_2(\cdot)} \left\| \left\| t^{k - \alpha_1} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} \\ &\quad + C_{k, \alpha_2, p(\cdot), q_2(\cdot)} \|f\|_{p(\cdot), \gamma_d} < +\infty. \end{aligned}$$

Therefore, $f \in F_{p(\cdot), q_2(\cdot)}^{\alpha_2}$. □

3. INTERPOLATION RESULTS

Finally, we are going to consider some interpolation results for the Gaussian variable Besov-Lipschitz and the variable Triebel-Lizorkin spaces.

We will use the following results for general variable Lebesgue spaces $L^{p(\cdot)}(X, \nu)$.

Lemma 3.1. *Let $p(\cdot) \in \mathcal{P}(\Omega, \nu)$ and $s > 0$ such that $sp^- \geq 1$. Then*

$$\| |f|^s \|_{p(\cdot), \nu} = \|f\|_{sp(\cdot), \nu}^s.$$

Proof. It is the same proof of Lemma 3.2.6 in [3]. □

Lemma 3.2. *Let ν a complete σ -finite measure on X . $r_j(\cdot) \in \mathcal{P}(X, \nu)$, $1 < r_j^-, r_j^+ < \infty$, $j = 0, 1$. For all $0 < \lambda < 1$, if $f \in L^{r_j(\cdot)}(X, \nu)$, $j = 0, 1$ then $f \in L^{r(\cdot)}(X, \nu)$ where $\frac{1}{r(y)} = \frac{1 - \lambda}{r_0(y)} + \frac{\lambda}{r_1(y)}$, a.e. $y \in X$ and*

$$(3.1) \quad \|f\|_{r(\cdot), \nu} \leq 2 \|f\|_{r_0(\cdot), \nu}^{1 - \lambda} \|f\|_{r_1(\cdot), \nu}^{\lambda}.$$

Proof. It is a consequence of Hölder's inequality (1.17) and Lemma 3.1. $\square$

Now, we present the interpolation result.

Theorem 3.1. *Let $p_j(\cdot) \in \mathcal{P}(\mathbb{R}^d, \gamma_d)$ and $q_j \in \mathcal{P}(\mathbb{R}^+, \mu)$, $j = 0, 1$*

i) For $1 < p_j^-, q_j^-, p_j^+, q_j^+ < +\infty$ and $\alpha_j \geq 0$, if $f \in B_{p_j(\cdot), q_j(\cdot)}^{\alpha_j}(\gamma_d)$, $j = 0, 1$, then for all $0 < \theta < 1$, $f \in B_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$, where

$$\alpha = \alpha_0(1 - \theta) + \alpha_1\theta$$

and

$$\begin{aligned} \frac{1}{p(x)} &= \frac{1 - \theta}{p_0(x)} + \frac{\theta}{p_1(x)}, \text{ a.e. } x \in \mathbb{R}^d, \\ \frac{1}{q(t)} &= \frac{1 - \theta}{q_0(t)} + \frac{\theta}{q_1(t)}, \text{ a.e. } t \in \mathbb{R}^+. \end{aligned}$$

ii) For $1 < p_j^-, q_j^-, p_j^+, q_j^+ < +\infty$ and $\alpha_j \geq 0$, if $f \in F_{p_j(\cdot), q_j(\cdot)}^{\alpha_j}(\gamma_d)$, $j = 0, 1$, then for all $0 < \theta < 1$, $f \in F_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$, where

$$\alpha = \alpha_0(1 - \theta) + \alpha_1\theta,$$

and

$$\begin{aligned} \frac{1}{p(x)} &= \frac{1 - \theta}{p_0(x)} + \frac{\theta}{p_1(x)}, \text{ a.e. } x \in \mathbb{R}^d, \\ \frac{1}{q(t)} &= \frac{1 - \theta}{q_0(t)} + \frac{\theta}{q_1(t)}, \text{ a.e. } t \in \mathbb{R}^+. \end{aligned}$$

Proof. i) Let k be any integer greater than α_0 and α_1 , by using Lemma 3.2, we obtain for $\alpha = \alpha_0(1 - \theta) + \alpha_1\theta$,

$$\begin{aligned} & \left\| t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \\ & \leq \left\| t^{k-(\alpha_0(1-\theta)+\alpha_1\theta)} 2 \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_0(\cdot), \gamma_d}^{1-\theta} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_1(\cdot), \gamma_d}^\theta \right\|_{q(\cdot), \mu} \\ & = 2 \left\| t^{(1-\theta)(k-\alpha_0)+\theta(k-\alpha_1)} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_0(\cdot), \gamma_d}^{1-\theta} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_1(\cdot), \gamma_d}^\theta \right\|_{q(\cdot), \mu} \\ & = 2 \left\| \left(t^{k-\alpha_0} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_0(\cdot), \gamma_d} \right)^{1-\theta} \left(t^{k-\alpha_1} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_1(\cdot), \gamma_d} \right)^\theta \right\|_{q(\cdot), \mu}. \end{aligned}$$

Thus, by Hölder's inequality (1.17) and Lemma 3.1,

$$\begin{aligned} & \left\| t^{k-\alpha} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p(\cdot), \gamma_d} \right\|_{q(\cdot), \mu} \\ & \leq 4 \left\| t^{k-\alpha_0} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_0(\cdot), \gamma_d} \right\|_{q_0(\cdot), \mu}^{1-\theta} \left\| t^{k-\alpha_1} \left\| \frac{\partial^k P_t f}{\partial t^k} \right\|_{p_1(\cdot), \gamma_d} \right\|_{q_1(\cdot), \mu}^\theta < +\infty, \end{aligned}$$

that is, $f \in B_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$.

ii) Analogously, by Hölder's inequality (1.17) and Lemma 3.1, we obtain for $\alpha = \alpha_0(1 - \theta) + \alpha_1\theta$,

$$\begin{aligned} \left\| t^{k-\alpha} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right\|_{q(\cdot), \mu} &= \left\| \left(t^{k-\alpha_0} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right)^{1-\theta} \left(t^{k-\alpha_1} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right)^\theta \right\|_{q(\cdot), \mu} \\ &\leq 2 \left\| t^{k-\alpha_0} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right\|_{q_0(\cdot), \mu}^{1-\theta} \left\| t^{k-\alpha_1} \left| \frac{\partial^k P_t f(x)}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu}^\theta, \end{aligned}$$

a.e. $x \in \mathbb{R}^d$. Therefore

$$\left\| \left\| t^{k-\alpha} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} \leq 2 \left\| \left\| t^{k-\alpha_0} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_0(\cdot), \mu}^{1-\theta} \left\| t^{k-\alpha_1} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu}^\theta \right\|_{p(\cdot), \gamma_d},$$

and again by Hölder's inequality and Lemma 3.1,

$$\begin{aligned} &\left\| \left\| t^{k-\alpha} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q(\cdot), \mu} \right\|_{p(\cdot), \gamma_d} \\ &\leq 4 \left\| \left\| t^{k-\alpha_0} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_0(\cdot), \mu} \right\|_{p_0(\cdot), \gamma_d}^{1-\theta} \left\| \left\| t^{k-\alpha_1} \left| \frac{\partial^k P_t f}{\partial t^k} \right| \right\|_{q_1(\cdot), \mu} \right\|_{p_1(\cdot), \gamma_d}^\theta < +\infty. \end{aligned}$$

That is, $f \in F_{p(\cdot), q(\cdot)}^\alpha(\gamma_d)$.

□

REFERENCES

- [1] Cruz-Urbe, D. & Fiorenza, A. *Variable Lebesgue Spaces Foundations and Harmonic Analysis*, Applied and Numerical Harmonic Analysis Birkhäuser-Springer, Basel, (2013)
- [2] Dalmaso, E. & Scotto, R. (2017) *Riesz transforms on variable Lebesgue spaces with Gaussian measure*, Integral Transforms and Special Functions, 28:5, 403-420, DOI: 10.1080/10652469.2017.1296835
- [3] Diening, L., Harjulehto, P., Hästö, P. and Růžička, M. *Lebesgue and Sobolev spaces with variable exponents*. Lecture Notes in Mathematics, 2017. Springer, Heidelberg, 2011.
- [4] Diening, L., Samko, S. *Hardy inequality in variable exponent Lebesgue spaces*. Fractional Calculus & Applied Analysis, Volume 10, Number 1 (2007).
- [5] Moreno, J., Pineda, E., & Urbina, W. *Boundedness of the maximal function of the Ornstein-Uhlenbeck semigroup on variable Lebesgue spaces with respect to the Gaussian measure and consequences*. Revista Colombiana de Matemáticas (2021) (to Appear)
- [6] Pineda, E., Urbina, W. *Some results on Gaussian Besov-Lipschitz and Gaussian Triebel-Lizorkin spaces*. J. Approx. Theor. 161(2), 529-564 (2009)
- [7] Stein E. *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press. Princeton, New Jersey. (1970).
- [8] Triebel, H. *Theory of function spaces*, Birkhäuser Verlag, Basel. (1983).
- [9] Triebel, H. *Theory of function spaces II*, Birkhäuser Verlag, Basel. (1992).
- [10] Urbina W. *Gaussian Harmonic Analysis*, Springer Monographs in Math. Springer Verlag, Switzerland AG (2019).

ESCUELA SUPERIOR POLITÉCNICA DEL LITORAL. ESPOL, FCNM, CAMPUS GUSTAVO GALINDO KM. 30.5
VÍA PERIMETRAL, P.O. Box 09-01-5863, GUAYAQUIL, ECUADOR.

Email address: epineda@espol.edu.ec

ESCUELA SUPERIOR POLITÉCNICA DEL LITORAL. ESPOL, FCNM, CAMPUS GUSTAVO GALINDO KM. 30.5
VÍA PERIMETRAL, P.O. Box 09-01-5863, GUAYAQUIL, ECUADOR.

Email address: luzeurod@espol.edu.ec

DEPARTMENT OF MATHEMATICS, ACTUARIAL SCIENCES AND ECONOMICS, ROOSEVELT UNIVERSITY, CHICAGO,
IL, 60605, USA.

Email address: wurbinaromero@roosevelt.edu