On the valuation of multiple reset options: integral equation approach

Nazym Azimbayev*† & Yerkin Kitapbayev*

February 10, 2022

In this paper, we study a pricing problem of the multiple reset put option, which allows the holder to reset several times a current strike price to obtain an at-the-money European put option. We formulate the pricing problem as a multiple optimal stopping problem, then reduce it to a sequence of single optimal stopping problems and study the associated free-boundary problems. We solve this sequence of problems by induction in the number of remaining reset rights and exploit probabilistic arguments such as local time-space calculus on curves. As a result, we characterize each optimal reset boundary as the unique solution to a nonlinear integral equation and derive the reset premium representations for the option prices. We propose that the multiple reset options can be used as cryptocurrency derivatives and an attractive alternative to standard European options due to the extreme volatility of underlying cryptocurrencies.

1. Introduction

Let us imagine an investor who holds a standard European call or put option with strike price K and maturity date T. Then there are at least two possible scenarios when the holder can feel a regret: 1) there is a period before T when underlying asset price movements are favorable for him; however, he cannot early exercise his option, and then the asset price will turn back and at time T he gets small or zero payoff; 2) the option becomes deep out-of-themoney, and the asset price is below K for the call (above K for the put) option, and most likely he gets zero payoff at time T. In the last two decades, the options with shouting and reset features have been introduced and studied, addressing these unfavorable scenarios. They can be divided into the two groups of options: 1) shout (call or put) option, which allows the holder to lock the profit at some endogenous time τ and then at time T get the maximum between two intrinsic values at τ and T; 2) reset (call or put) option gives to the investor the right to reset the strike K to the current price, i.e., to substitute the current out-of-the-money option to the at-the-money one. The first group, i.e., shout options, allows the investor to lock

^{*}Department of Computational Mathematics and Cybernetics, Lomonosov Moscow State University, Moscow, Russia; nazym.azimbayev@cs.msu.ru

[†]The National Bank of Kazakhstan, Nur-Sultan, Kazakhstan

[‡]Department of Mathematics, NC State University, Raleigh NC, USA; ykitapb@ncsu.edu

Mathematics Subject Classification 2010. Primary 91G20, 60G40. Secondary 60J60, 35R35, 45G10.

Key words and phrases: multiple reset option, shout option, multiple optimal stopping, geometric Brownian motion, free-boundary problem, local time, integral equation, cryptocurrencies, bitcoin.

the profit while having the opportunity of payoff increase at T. In contrast, the second one increases the chances to get a positive payoff at T. The pricing problem for both options can be formulated as optimal stopping problems where stopping times represent shouting or reset times, respectively. They have both European (since the payoff is known at T only) and American features (due to early 'shouting' or 'reset' opportunities).

Below we provide a literature review on the reset and shout options. These options might have both single and multiple opportunities to shout or reset. However, so far, the research has been focused primarily on single shout and reset options. The origin of the shout option goes to the paper [11] and brief analysis can be found in textbooks [7] and [12] where the binomial tree method is offered for pricing the option. There are several papers where these options were thoroughly studied from both theoretical and numerical points of view. In series of works [13]-[15] several sophisticated numerical schemes have been developed to price the reset and shout options. The comprehensive analysis has been performed in [4], where the reset put option was formulated as a free-boundary problem, shouting premium for the option price and recursive integral equation for the optimal shouting boundary were obtained. In [3], authors studied the multiple reset put option problem, i.e., the buyer has several rights to reset a strike. The numerical experiments and analysis have been provided using the binomial method approach. Several monotonicity properties for option prices and optimal reset boundaries were established through theoretical arguments and numerical experiments.

Later, a thorough theoretical analysis was provided in [16], where PDE and variational inequality approaches were applied to show the existence and uniqueness of a solution to the free-boundary problem associated with the reset put option problem. Also, monotonicity and regularity of the optimal shouting boundary have been shown in some cases. Then in [1], using a Laplace transform, the Fredholm integro-differential equations for optimal shouting boundaries of shout call and put options were obtained. Finally, in [6] the formal series expansions have been discovered for the option price and optimal shouting boundaries of the reset put and shout call options.

This paper is devoted to the pricing of the multiple reset put option. The contribution of the paper is twofold. Firstly, we study the single reset put option and reduce it to a freeboundary problem, and then we tackle the latter using mostly probabilistic arguments such as Markov property and local time-space calculus (see [8]). We provide the probabilistic proof of the monotonicity of optimal reset boundary in some situations. We then obtain the same reset premium representation for the option price and recursive integral equation for optimal reset boundary as in [4]. Secondly and most importantly, we complement the work [3] and study the multiple reset option using local time-space formula and by induction in number nof reset rights. To the best of our knowledge, this is the first paper where the reset premium representations for the option prices with multiple rights and the recursive integral equations for each optimal reset boundaries were obtained. This result extends the results for the single reset option. Thirdly, we describe the numerical algorithm for pricing multiple reset options and plotting optimal reset boundaries. Fourthly, the paper also recalls the put-call reset-shout parity, i.e., the reset put option problem is equivalent to shout call one in the sense that their optimal strategies coincide and their prices have a simple relationship. The same fact is true for reset call and shout put options, which was also observed in [3] and [6].

Finally, we also note that these contracts can be attractive for investors in cryptocurrency markets. The high volatility of cryptocurrencies calls for more flexible option contracts, which

could give new opportunities to investors even if they see sudden jumps in the price of the underlying cryptocurrency. It then seems natural to consider reset strike or shout options in this market as an alternative to plain vanilla call/put options.

The paper is organised as follows. In Section 2 we formulate the multiple reset put option problem as a multiple optimal stopping problem, which we reduce to the sequence of single optimal stopping problems. In Section 3 we provide a put-call parity between reset and shout options. Section 4 is devoted to the single reset put option. In Section 5 we derive reset premium representation for the price of the multiple reset option and characterize the optimal reset boundaries as the unique solution to nonlinear integral equations. Finally, Section 6 provides numerical algorithms for numerical solutions to the multiple reset option problem.

2. Formulation of the problem

Let us suppose the probability space $(\Omega, \mathcal{F}, \mathbb{Q})$, where \mathbb{Q} is the risk-neutral measure. In this paper, we study the multiple reset put option problem. The underlying asset price X follows geometric Brownian motion

$$(2.1) dX_t = (r - \delta)X_t dt + \sigma X_t dB_t (X_0 = x)$$

where B is a \mathbb{Q} -standard Brownian motion started at zero, r > 0 is the interest rate, $\delta \geq 0$ is the dividend yield and $\sigma > 0$ is the constant volatility coefficient. The solution X to the stochastic differential equation (2.1) is given by

(2.2)
$$X_t = x \exp\left((r - \delta - \sigma^2/2)t + \sigma B_t\right)$$

for t > 0, x > 0.

By definition of the multiple reset put option with n rights the payoff at maturity time T is the following: if the buyer "resets" at the sequence of endogenous times $\tau_1 < \leq \tau_n \leq T$ (i.e. at least $\tau_1 < T$, others may equal T), he obtains $(X_{\tau_1} \vee \vee X_{\tau_n} - X_T)^+$ and if the buyer does not reset until T (i.e. $\tau_1 = T$), his payoff equals $(K - X_T)^+$, where K > 0 is the original strike price. Hence the multiple reset put option allows to obtain sequentially at-the-money put options by reseting the strike at times τ_i , i = 1, ..., n. Clearly, one should reset her/his i-th right only when $X_t > K_{i-1}$ with $K_i = X_{\tau_i}$, i = 1, ..., n-1 and $K_0 = K$. This option of an European type since the payoff is delivered at T, however it has an "American"-style feature of an early action - reseting.

Now let us turn to the valuation of multiple reset put option. If the holder resets (or does not reset and $\tau_1 = T$) at the sequence of stopping times $\tau := (\tau_1, ..., \tau_n)$ with respect to the natural filtration of X, then the payoff at maturity T is given by

$$(2.3) (X_{\tau_1} \vee ... \vee X_{\tau_n} \vee K - X_T)^+.$$

Thus the arbitrage-free price of the multiple reset put option at time $t \in [0,T)$ as a value function of the following optimal stopping problem

(2.4)
$$V_n^{RP}(t, x; K) = e^{-r(T-t)} \sup_{t \le \tau \le T} \mathbb{E}_{t, x} \left[(X_{\tau_1} \lor \dots \lor X_{\tau_n} \lor K - X_T)^+ \right]$$

where the supremum is taken over the set of sequences of stopping times $\{\tau = (\tau_1, ..., \tau_n) : 0 \le \tau_1 \le ... \le \tau_n \le T\}$, the expectation $\mathbb{E}_{t,x}$ is taken under \mathbb{Q} and $X_t = x$, and we include the discount factor e^{-rT} since the payoff is delivered at T. We also emphasize the dependence of V_n^{RP} on the current strike level K. It is important to note that since the terminal payoff function (2.3) is not known at reset times τ_i , the multiple optimal stopping problem (2.4) falls into the class of optimal prediction problems. Hence we need to reduce this problem to a standard optimal stopping problem with an adapted payoff function and also to reformulate it as the sequence of single stopping problems. For this, let us now rewrite the payoff in the following way

$$(2.5) \qquad \max(X_{\tau_1} \vee ... \vee X_{\tau_n} \vee K - X_T, 0) = \max(X_{\tau_2} \vee ... \vee X_{\tau_n} \vee (X_{\tau_1} \vee K) - X_T, 0).$$

The equality (2.5) shows that the payoff of the option with remaining n rights and strike K is the same as the payoff of the option with remaining n-1 rights left and the strike $X_{\tau_1} \vee K$. Now by conditioning on X_{τ_1} we reduce (2.4) to

(2.6)
$$V_n^{RP}(t, x; K) = \sup_{t < \tau_1 < T} \mathbb{E}_{t, x} \left[e^{-r(\tau_1 - t)} V_{n-1}^{RP}(\tau_1, X_{\tau_1}; X_{\tau_1} \vee K) \right].$$

Therefore to solve the problem (2.4) we need to tackle n single optimal stopping problems sequentially starting from n=1 and V_0^{RP} being the European put option price. This reduction is very natural as it says that the price of the reset option with n rights can be obtained as the expected discounted value at the first optimal reset time τ_1^* of the reset option with n-1 remaining rights. The reduction (2.6) was also stated in [3]. The reduction of multiple optimal stopping problems to the sequence of single optimal stopping problems is quite standard in the literature of swing options, see e.g. [2] and the recent work on finite horizon [5]. The pricing of the swing options on finite horizon are different from reset/shout options due to the presence of so-called refracting period, which is the minimal period between two exercise times.

Let us now write the problem as

(2.7)
$$V_n^{RP}(t, x; K) = \sup_{t < \tau < T} \mathbb{E}_{t, x} \left[e^{-r(\tau - t)} G_n(\tau, X_\tau) \right]$$

for $n \ge 1$, $0 \le t \le T$ and x, K > 0 where the payoff function G_n reads inductively

(2.8)
$$G_n(t,x) := V_{n-1}^{RP}(t,x;x \vee K)$$

and

(2.9)
$$V_0^{RP}(t, x; K) = V^e(t, x; x \vee K) = \mathbb{E}_{t,x} \left[e^{-r(T-t)} (x \vee K - X_T)^+ \right]$$

is the price of European put option at time $t \in [0,T)$ and asset price x > 0, with strike $x \vee K$ and expiry date T.

3. Parity between shout and reset options

In this section we recall definitions of reset call option, shout put and call options with multiple rights, and show the parity between them.

Definition 3.1. Reset call option has the following payoff at maturity time T: if the buyer 'resets' at sequence of times $\tau_1 < \ldots \leq \tau_n < T$ he gets $(X_T - X_{\tau_1} \wedge \ldots \wedge X_{\tau_n})^+$ and if the buyer does not 'reset' until T his payoff equals $(X_T - K)^+$, where K > 0 is the original strike price.

Definition 3.2. Shout call option has the following payoff at maturity time T: if the buyer 'shouts' at sequence of times $\tau_1 < \leq \tau_n < T$ he gets $\max(X_{\tau_1} - K, ..., X_{\tau_n} - K)$ and if the buyer does not 'shout' until T his payoff equals $(X_T - K)^+$, where K > 0 is the strike price.

Definition 3.3. Shout put option has the following payoff at maturity time T: if the buyer 'shouts' at sequence of times $\tau_1 < \leq \tau_n < T$ he gets $\max(K - X_{\tau_1}, ..., K - X_{\tau_n})$ and if the buyer does not 'shout' until T his payoff equals $(K - X_T)^+$, where K > 0 is the strike price.

If the holder resets (or shouts) at sequence of times $\tau_1 \leq \ldots \leq \tau_n < T$, then the payoff for reset call option, and shout call and put options with n rights at maturity T under risk-neutral measure \mathbb{Q} is given by, respectively

$$(3.1) (X_T - X_{\tau_1} \wedge \ldots \wedge X_{\tau_n} \wedge K)^+$$

(3.2)
$$\max(X_{\tau_1} - K, \dots, X_{\tau_n} - K, X_T - K, 0)$$

(3.3)
$$\max(K - X_{\tau_1}, \dots, K - X_{\tau_n}, K - X_T, 0)$$

and their arbitrage-free prices

$$(3.4) V_n^{RC} = e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E}\left[(X_T - X_{\tau_1} \land \dots \land X_{\tau_n} \land K)^+ \right]$$

(3.5)
$$V_n^{SC} = e^{-rT} \sup_{0 < \tau < T} \mathbb{E} \left[\max(X_{\tau_1} - K, \dots, X_{\tau_n} - K, X_T - K, 0) \right]$$

(3.4)
$$V_n^{RC} = e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[(X_T - X_{\tau_1} \land \dots \land X_{\tau_n} \land K)^+ \right]$$
(3.5)
$$V_n^{SC} = e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[\max(X_{\tau_1} - K, \dots, X_{\tau_n} - K, X_T - K, 0) \right]$$
(3.6)
$$V_n^{SP} = e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[\max(K - X_{\tau_1}, \dots, K - X_{\tau_n}, K - X_T, 0) \right].$$

Straightforward calculations show the parity between the prices of reset call and shout put options

(3.7)
$$V_n^{RC} = e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[\max(X_T - X_{\tau_1}, \dots, X_T - X_{\tau_n}, X_T - K, 0) \right]$$
$$= e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[X_T - K + \max(K - X_{\tau_1}, \dots, K - X_{\tau_n}, K - X_T, 0) \right]$$
$$= X_0 e^{-\delta T} - K e^{-rT} + V_n^{SP}$$

and between the prices of reset put and shout call options

(3.8)
$$V_n^{RP} = e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[\max(X_{\tau_1} - X_T, \dots, X_{\tau_n} - X_T, K - X_T, 0) \right]$$
$$= e^{-rT} \sup_{0 \le \tau \le T} \mathbb{E} \left[K - X_T + \max(X_{\tau_1} - K, \dots, X_{\tau_n} - K, X_T - K, 0) \right]$$
$$= Ke^{-rT} - X_0e^{-\delta T} + V_n^{SC}$$

and the relationship between all of them

$$(3.9) V_n^{RP} - V_n^{SC} = V_n^{SP} - V_n^{RC} = Ke^{-rT} - X_0e^{-\delta T}.$$

Finally, we note that the optimal stopping times are the same for reset call and shout put as well as for reset put and shout call options.

4. Single reset put option

Throughout the rest of the paper we focus on the reset put option and for notational convenience we use V_n instead of V_n^{RP} . In this section we study the pricing of single reset put option, i.e., n = 1 in (2.7)

(4.1)
$$V_1(t, x; K) = \sup_{t < \tau < T} \mathbb{E}_{t, x} \left[e^{-r(\tau - t)} G_1(\tau, X_\tau) \right]$$

for $t \in [0, T)$ and x, K > 0, and where

(4.2)
$$G_1(t,x) = \mathbb{E}_{t,x} \left[e^{-r(T-t)} (x \vee K - X_T)^+ \right].$$

We solve the problem for fixed K so we will use notation $V_1(t,x)$ instead of $V_1(t,x;K)$ in this section.

We will reduce the problem into a free-boundary problem and the latter will be tackled in the next section using local time-space calculus ([8]). First using that the gain function G_1 is continuous and standard arguments (see e.g. Corollary 2.9 (Finite horizon) with Remark 2.10 in [10]) we have that continuation and reset sets read

$$(4.3) C_1 = \{ (t, x) \in [0, T) \times [0, \infty) : V_1(t, x) > G_1(t, x) \}$$

$$(4.4) D_1 = \{ (t, x) \in [0, T) \times [0, \infty) : V_1(t, x) = G_1(t, x) \}$$

and the optimal stopping time in (4.1) is given by

(4.5)
$$\tau_1^* = \inf \{ t \le s \le T : (s, X_s) \in D_1 \}.$$

1. We will now discuss some properties of the functions G_1 and V_1 . The gain function G_1 can be simplified

(4.6)
$$G_1(t,x) = xV^e(t,1;1) =: xV^e(t)$$

$$= x \left[e^{-r(T-t)} \Phi\left(\left(\frac{\delta - r}{\sigma} + \frac{\sigma}{2} \right) \sqrt{T - t} \right) - e^{-\delta(T-t)} \Phi\left(\left(\frac{\delta - r}{\sigma} - \frac{\sigma}{2} \right) \sqrt{T - t} \right) \right]$$

for x > K, where $\Phi(\cdot)$ is the cumulative distribution function of the standard normal law and $V^e(t)$ is the price of European put option at t with x = 1 and strike 1. On the other hand, G_1 equals to

(4.7)
$$G_1(t,x) = Ke^{-r(T-t)}\Phi\left(\frac{1}{\sigma\sqrt{T-t}}\left[\log(\frac{K}{x}) - (r-\delta - \frac{\sigma^2}{2})(T-t)\right]\right) - xe^{-\delta(T-t)}\Phi\left(\frac{1}{\sigma\sqrt{T-t}}\left[\log(\frac{K}{x}) - (r-\delta + \frac{\sigma^2}{2})(T-t)\right]\right)$$

for $0 < x \le K$ and this is exactly the European put option price $V^e(t,x;K)$ with strike K. We know that $x \mapsto V^e(t,x)$ is convex on $(0,\infty)$ for any $0 \le t < T$ fixed so that $x \mapsto G_1(t,x)$ is convex on (0,K) for every t fixed. Since G_1 is linear in x on (K,∞) in order to prove that $x \mapsto G_1(t,x)$ is convex on $(0,\infty)$ for every t fixed it is enough to show that $(G_1)_x(t,K+) \ge (G_1)_x(t,K-)$. Using (4.6), (4.7) and well-known expression for 'delta'

coefficient of the European put option $\Delta_p = \frac{\partial V^e}{\partial x} = -\Phi\left(\frac{1}{\sigma\sqrt{T-t}}\left[\log(\frac{K}{x}) - (r-\delta + \frac{\sigma^2}{2})(T-t)\right]\right)$ we have that

$$(4.8) (G_1)_x(t,K+) = e^{-r(T-t)}\Phi\left(\left(\frac{\delta-r}{\sigma} + \frac{\sigma}{2}\right)\sqrt{T-t}\right) - \Phi\left(\left(\frac{\delta-r}{\sigma} - \frac{\sigma}{2}\right)\sqrt{T-t}\right)$$

$$(4.9) (G_1)_x(t,K-) = -\Phi\left(\left(\frac{\delta-r}{\sigma} - \frac{\sigma}{2}\right)\sqrt{T-t}\right)$$

and thus it is clear that $(G_1)_x(t,K+) \geq (G_1)_x(t,K-)$. Hence the function $x \mapsto G_1(t,x)$ is convex on $(0,\infty)$ for every t fixed and thus using standard arguments we obtain that $x \mapsto V_1(t,x)$ is convex on $(0,\infty)$ as well.

2. It will be useful for further analysis to calculate the expression $H_1:=\frac{\partial G_1}{\partial t}+\mathbb{L}_XG_1-rG_1$ for $(t,x)\in[0,T)\times(0,\infty)$ where $\mathbb{L}_X=(r-\delta)xd/dx+(\sigma^2/2)x^2d^2/dx^2$ is the infinitesimal operator of X. Since we have that $G_1(t,x)=V^e(t,x)$ for $(t,x)\in[0,T)\times(0,K)$ and it is well-known that $V_t^e+\mathbb{L}_XV^e-rV^e=0$ for all $(t,x)\in[0,T)\times(0,\infty)$ then

(4.10)
$$H_1(t,x) = 0$$
 on $[0,T) \times (0,K)$.

Now we consider set $\{(t,x) \in [0,T) \times (K,\infty)\}$ and there G_1 is given by (4.6) so that we have

$$(4.11) \quad H_1(t,x) = x \left[(r-\delta)e^{-r(T-t)}\Phi\left(\left(\frac{\delta-r}{\sigma} + \frac{\sigma}{2}\right)\sqrt{T-t}\right) - \frac{\sigma}{2\sqrt{T-t}}e^{-r(T-t)}\varphi\left(\left(\frac{\delta-r}{\sigma} + \frac{\sigma}{2}\right)\sqrt{T-t}\right) \right]$$

for $(t,x) \in [0,T) \times (K,\infty)$ and where $\varphi(\cdot)$ is the probability density function of the standard normal law. For convenience we define

$$(4.12) h_1(t) = (V^e(t))' - \delta V^e(t)$$

$$= (r - \delta)e^{-r(T-t)}\Phi\left(\left(\frac{\delta - r}{\sigma} + \frac{\sigma}{2}\right)\sqrt{T - t}\right) - \frac{\sigma}{2\sqrt{T - t}}e^{-r(T-t)}\varphi\left(\left(\frac{\delta - r}{\sigma} + \frac{\sigma}{2}\right)\sqrt{T - t}\right)$$

for $t \in [0,T)$ so that $H_1(t,x) = xh_1(t)$ for $(t,x) \in [0,T) \times (K,\infty)$.

By straightforward calculations it can be seen from (4.12) that $h_1(t) \downarrow -\infty$ as $t \to T$ and $h_1(0) > 0$ as $T \to \infty$. Moreover, there exists large enough T > 0 and $t_* < T$ such that h_1 is increasing on $[0,t_*)$ and is decreasing on (t_*,T) . Then for this size of T there is a point $t^* > t_*$ with $h_1 > 0$ on $[0,t^*)$ and $h_1 \le 0$ on $[t^*,T)$. We note that for typical values of parameters (T,r,δ,σ) we have that h_1 is decreasing and negative on [0,T) and thus positive points t_* and t^* do not exist.

3. Below we derive some insights into the structure of reset set D_1 . Using Ito-Tanaka's formula and (5.21) with (4.8)-(4.9) we have

$$(4.13) \qquad \mathbb{E}_{t,x}\left[e^{-r(\tau-t)}G_1(\tau,X_\tau)\right] = G_1(t,x) + \mathbb{E}_{t,x}\left[\int_t^\tau e^{-r(s-t)}h_1(s)X_sI(X_s \ge K)ds\right] + \frac{1}{2}\mathbb{E}_{t,x}\left[\int_t^\tau e^{-r(s-t)}\Phi\left(-\left(\frac{r}{\sigma} - \frac{\sigma}{2}\right)\sqrt{T-s}\right)d\ell_s^K(X)\right]$$

for $(t,x) \in [0,T) \times (0,\infty)$ and any stopping time τ of X, where $(\ell_s^K(X))_{s \geq t}$ is the local time process of X at level K. The formula (4.13) shows that for x < K the first integrand

on the right-hand side vanishes and the buyer should continue due to the non-negative integral with respect to local time at K. When x>K and the integrand h_1 is positive, i.e. $t< t^*$, then it is again optimal to wait. Now if x>K and $h_1(t)\leq 0$, i.e. $t\geq t^*$, the only incentive to wait is to accumulate a positive local time at K in future up to T. However this makes sense only for asset prices above but close to K and for large x one should reset the strike immediately due to very negative h_1X . To conclude, we obtain that it is optimal to reset only when $t\geq t^*$ (if t^* exists as described above) and large enough x above K.

Now we can describe the structure of the reset set D_1 . Namely, from the fact that it is not optimal to exercise the reset put option below K, definitions (4.3) and (4.4), convexity of V_1 and linearity of G_1 above K it follows that there exists an optimal reset boundary $b_1:[0,T]\to \mathbb{R}$ such that

$$\tau_1^* = \inf \{ t \le s \le T : X_s \ge b_1(s) \}$$

is optimal in (4.1) and $b_1(t) > K$ for $t \in [0,T)$. Also $b_1 \equiv +\infty$ on $[0,t^*)$ if t^* exists. It is clear that if x > K and t < T is sufficiently close to T then it is optimal to reset immediately (since $h_1 < 0$ near T and the profit obtained from the local time term at K cannot compensate the cost of getting there due to the lack of remaining time). This shows that $b_1(T-) = K$.

4. Standard Markovian arguments (see [10]) lead to the following free-boundary problem (for the value function $V_1 = V_1(t, x)$ and the optimal reset boundary $b_1 = b_1(t)$ to be determined):

(4.15)
$$\frac{\partial V_1}{\partial t} + \mathbb{L}_X V_1 - rV_1 = 0 \qquad \text{in } C_1$$

$$(4.16) V_1(t, b_1(t)) = G_1(t, b_1(t)) \text{for } t \in [0, T)$$

(4.17)
$$\frac{\partial V_1}{\partial x}(t, b_1(t)) = \frac{\partial G_1}{\partial x}(t, b_1(t)) \qquad \text{for } t \in [0, T)$$

(4.18)
$$V_1(T, x) = (K - x)^+ \qquad \text{for } x \in (0, \infty)$$

$$(4.19) V_1(t,x) > G_1(t,x) in C_1$$

$$(4.20) V_1(t,x) = G_1(t,x) in D_1$$

where the continuation set C_1 and the reset set D_1 are given by

(4.21)
$$C_1 = \{ (t, x) \in [0, T) \times (0, \infty) : x < b_1(t) \}$$

(4.22)
$$D_1 = \{ (t, x) \in [0, T) \times (0, \infty) : x \ge b_1(t) \}$$

with $b_1 = +\infty$ on $[0, t^*)$ if $t^* > 0$ exists. It can be shown that this free-boundary problem has a unique solution V_1 and b_1 which coincide with the value function and the optimal reset boundary, respectively.

Completed details of the analysis above go beyond our goals in this paper and for this reason will be omitted. It should be noted however that one of the main issues which makes this analysis quite complicated (in comparison with e.g. the American put option problem) is that the proof of monotonicity in time of b_1 is still an open question for certain values of parameters. This property has been stated in previous papers on reset and shout options but actually have not been proven apart from [16], where PDE arguments were used to prove the

monotonicity for some occasions. We provide the probabilistic proof for the same situations, i.e., when h_1 is decreasing on [0,T) and $r \leq \sigma^2/2$. In this case, we can easily see from (4.13) that both integral terms are decreasing in t and obtain that $\partial V_1/\partial t \leq \partial G_1/\partial t$ and therefore b_1 is decreasing. In the opposite case, when h_1 be increasing on $[0,t_*)$ if such $t_*>0$ exists or $r>\sigma^2/2$, both integrals on (4.13) are not decreasing at the same time and it is not clear how to compare them. However, numerical results clearly support the hypothesis that b_1 is decreasing, and it was also obtained numerically in other papers too.

Another difficulty of the analysis is that the proof of the continuity of the free boundary without having its monotonicity is a challenging problem, which can help to tackle other optimal stopping problems. In the next paragraph we will derive simpler equations which characterize V_1 and b_1 uniquely and can be used for the pricing of the reset put option and computation of the optimal reset boundary.

5. We now provide the reset premium representation formula for the arbitrage-free price V_1 which decomposes it into the sum of the European put option price and the 'reset' premium. The optimal reset boundary b_1 will be obtained as the unique solution to the integral equation.

We will make use of the following function in Theorem 4.1 below:

(4.23)
$$L_{1}(t, u, x, z) = -e^{-r(u-t)} \mathbb{E}_{t,x} \left[h_{1}(u) X_{u} I(X_{u} \geq z) \right]$$
$$= -xe^{-\delta(u-t)} h_{1}(u) \widehat{\mathbb{Q}}(X_{u} \geq z)$$
$$= -xe^{-\delta(u-t)} h_{1}(u) \Phi \left[\left(\log(x/z) + (r - \delta + \sigma^{2}/2)(u - t) \right) / (\sigma \sqrt{u - t}) \right]$$

for $t, u \geq 0$ and x, z > 0 and where the measure $\widehat{\mathbb{Q}}$ is defined as $d\widehat{\mathbb{Q}} = e^{\sigma B_T - \sigma^2 T/2} d\mathbb{Q}$ and we used that $\widehat{B}_t = B_t + \sigma t$ is a standard Brownian motion under $\widehat{\mathbb{Q}}$.

The main result of this section may now be stated as follows.

Theorem 4.1. The optimal reset boundary b_1 can be characterised as the unique solution to the nonlinear integral equation

(4.24)
$$b_1(t)V^e(t) = V^e(t, b_1(t)) + \int_t^T L_1(t, u, b_1(t), b_1(u)) du$$

for $t \in [0,T]$ in the class of continuous functions above K with $b_1(T-) = K$. Then the value function V_1 in (4.1) has the following representation

(4.25)
$$V_1(t,x) = V^e(t,x) + \int_t^T L_1(t,u,x,b_1(u)) du$$

for $t \in [0,T]$ and $x \in (0,\infty)$, where V_1 is decomposed as the sum of the European put price V^e and the reset premium.

Proof. (A) First we clearly have that the following conditions hold: (i) V_1 is $C^{1,2}$ on $C_1 \cup D_1$; (ii) b_1 is of bounded variation; (iii) $(V_1)_t + \mathbb{L}_X V_1 - rV_1$ is locally bounded; (iv) $x \mapsto V_1(t,x)$ is convex (recall paragraph 1 above); (v) $t \mapsto (V_1)_x(t,b_1(t)\pm)$ is continuous (recall (4.17)). Hence we can apply the local time-space formula on curves [8] for $e^{-r(s-t)}V_1(s,X_s)$:

$$(4.26) e^{-r(s-t)}V_1(s, X_s)$$

$$\begin{split} &= V_{1}(t,x) + M_{s} \\ &+ \int_{t}^{s} e^{-r(u-t)} \left(\frac{\partial V_{1}}{\partial t} + \mathbb{L}_{X} V_{1} - r V_{1} \right) (u,X_{u}) I(X_{u} > b_{1}(u)) du \\ &+ \frac{1}{2} \int_{t}^{s} e^{-r(u-t)} \left(\frac{\partial V_{1}}{\partial x} (u,X_{u} +) - \frac{\partial V_{1}}{\partial x} (u,X_{u} -) \right) I(X_{u} = b_{1}(u)) d\ell_{u}^{b_{1}}(X) \\ &= V_{1}(t,x) + M_{s} + \int_{t}^{s} e^{-r(u-t)} \left(\frac{\partial G_{1}}{\partial t} + \mathbb{L}_{X} G_{1} - G_{1} \right) (u,X_{u}) I(X_{u} \ge b_{1}(u)) du \\ &= V_{1}(t,x) + M_{s} + \int_{t}^{s} e^{-r(u-t)} h_{1}(u) X_{u} I(X_{u} \ge b_{1}(u)) du \end{split}$$

where we used (4.15), (5.21), the smooth-fit condition (4.17) and where $M=(M_u)_{u\geq 0}$ is the martingale part, $(\ell_u^{b_1}(X))_{u\geq 0}$ is the local time process of X spending at the boundary b_1 . Now upon letting s=T, taking the expectation $\mathbb{E}_{t,x}$, the optional sampling theorem for M, rearranging terms and noting that $V_1(T,x)=G_1(T,x)=(x\vee K-x)^+=(K-x)^+$ for all x>0, we get (4.25). The integral equation (4.24) one obtains by simply putting $x=b_1(t)$ into (4.25) and using (4.16).

(B) The proof of that b_1 is the unique solution to the equation (4.24) in the class of continuous functions $t \mapsto b_1(t)$ is based on standard arguments originally derived in [9] and omitted here.

5. Multiple reset put option

In this section we turn to the multiple reset put option problem

(5.1)
$$V_n(t,x) = V_n(t,x;K) = \sup_{t < \tau < T} \mathbb{E}\left[e^{-r(\tau - t)}G_n(\tau, X_\tau)\right]$$

for $n \ge 1$, $0 \le t \le T$ and x, K > 0 where the gain function G_n is given inductively

(5.2)
$$G_n(t,x) := V_{n-1}(t,x; x \vee K)$$

for $0 \le t \le T$ and x > 0. Having solved (4.1) for n = 1 in the previous section, we can derive the pricing formulas and integral equations for general $n \ge 2$ by induction.

Theorem 5.1. For any $n \ge 1$ the optimal stopping time in (5.1) is given by

(5.3)
$$\tau_n^* = \inf \{ t \le s \le T : X_s \ge b_n(s) \}$$

where the optimal reset boundary b_n can be characterised as the unique solution to the nonlinear integral equation

(5.4)
$$b_n(t)(V^e(t) + p_{n-1}(t)) = V^e(t, b(t)) + \int_t^T L_n(t, u, b_n(t), b_n(u)) du$$

for $t \in [0,T]$ in the class of continuous functions above K with $b_n(T-) = K$. The value function V_n is characterized by the reset premium representation

(5.5)
$$V_n(t,x) = V^e(t,x) + \int_t^T L_n(t,u,x,b_n(u)) du$$

for $t \in [0,T]$ and $x \in (0,\infty)$, and where

(5.6)
$$L_n(t, u, x, z) = -xe^{-\delta(u-t)}h_n(u)\widehat{\mathbb{Q}}\left(X_{u-t}^x \ge z\right)$$

(5.7)
$$h_n(t) = (V^e(t) + p_{n-1}(t))' - \delta(V^e(t) + p_{n-1}(t))$$

(5.8)
$$p_n(t) = -\int_t^T e^{-\delta(u-t)} h_n(u) \widehat{\mathbb{Q}}_{t,K} (X_u \ge b_n(u;K)) du$$

$$(5.9) p_0(t) \equiv 0.$$

Moreover, V_n and b_n solve the following free-boundary problem

(5.10)
$$\frac{\partial V_n}{\partial t} + \mathbb{L}_X V_n - rV_n = 0 \qquad in C_n$$

(5.11)
$$V_n(t, b_n(t)) = G_n(t, b_n(t)) = b_n(t) \left(V^e(t) + p_{n-1}(t) \right) \qquad \text{for } t \in [0, T)$$

(5.12)
$$\frac{\partial V_n}{\partial x}(t, b_n(t)) = \frac{\partial G_n}{\partial x}(t, b_n(t)) = V^e(t) + p_{n-1}(t) \qquad for \ t \in [0, T)$$

(5.13)
$$V_n(T,x) = (K-x)^+$$
 for $x \in (0,\infty)$

$$(5.14) V_n(t,x) > G_n(t,x) in C_n$$

$$(5.15) V_n(t,x) = G_n(t,x) in D_n$$

where the continuation set C_n and the stopping set D_n are given by

(5.16)
$$C_n = \{ (t, x) \in [T_0, T_n) \times (0, \infty) : x < b_n(t) \}$$

(5.17)
$$D_n = \{ (t, x) \in [T_0, T_n) \times (0, \infty) : x \ge b_n(t) \}.$$

Proof. 1. The proof is followed by induction. In the previous section we verified the statement of theorem for n = 1, so now we assume that the formulas hold for n - 1. In what follows we prove (5.4) and (5.5) for n. The proof will follow along the same lines as for n = 1.

We have that the payoff function G_n is

(5.18)
$$G_n(t,x) = V_{n-1}(t,x;K)$$

for $0 < x \le K$ and thus the local benefits of waiting

(5.19)
$$H_n(t,x) := \left(\frac{\partial G_n}{\partial t} + \mathbb{L}_X G_n - rG_n\right)(t,x) = 0 \quad \text{on} \quad [0,T) \times (0,K)$$

where we used (5.10) for n-1. Now if x>K, using the assumption of induction we obtain

(5.20)
$$G_n(t,x) = V_{n-1}(t,x;x)$$
$$= V^e(t,x;x) + \int_t^T L_{n-1}(t,u,x,b_n(u;x))du$$

$$= x \left(V^{e}(t) - \int_{t}^{T} e^{-\delta(u-t)} h_{n-1}(u) \widehat{\mathbb{Q}}(X_{u-t}^{x} \ge b_{n-1}(u;x)) du \right)$$

= $x \left(V^{e}(t) + p_{n-1}(t) \right)$

so that

(5.21)
$$H_n(t,x) = x \left(V^e(t) + p_{n-1}(t) \right)' - \delta x \left(V^e(t) + p_{n-1}(t) \right) = x h_n(t)$$

for $(t, x) \in [0, T) \times (K, \infty)$.

2. Now by applying Ito-Tanaka's formula for $e^{-r(s-t)}G_n(s,X_s)$ we have

$$(5.22) \qquad \mathbb{E}_{t,x} \left[e^{-r(\tau-t)} G_n(\tau, X_{\tau}) \right]$$

$$= G_n(t,x) + \mathbb{E}_{t,x} \left[\int_t^{\tau} e^{-r(s-t)} H_n(s, X_s) I(X_s \ge K) ds \right]$$

$$+ \frac{1}{2} \mathbb{E}_{t,x} \left[\int_t^{\tau} e^{-r(s-t)} \left(\frac{\partial G_n}{\partial x}(s, K+) - \frac{\partial G_n}{\partial x}(s, K-) \right) d\ell_s^K(X) \right]$$

for $(t,x) \in [0,T) \times (0,\infty)$ and any stopping time τ of X. The formula (5.22) shows that for $X_t < K$ it is not optimal to reset at once.

Using similar arguments as for n=1 we can deduce that there exists an optimal reset boundary $b_n:[0,T]\to \mathbb{R}$ such that

(5.23)
$$\tau_n^* = \inf \{ t \le s \le T : X_s \ge b_n(s) \}$$

is optimal in (5.1) and $b_n(t) > K$ for $t \in [0,T)$ with $b_n(T-) = K$.

3. Using standard Markovian arguments one can show that V_n and the optimal reset boundary b_n uniquely solve the free-boundary problem (5.10)-(5.15). As for the single reset option problem in previous section, the main challenge here is to prove the monotonicity of the boundary b_n . We note that due to complicated expression for h_n , it is even more difficult to prove this property rather than for n=1, where we could show it for some cases. However numerical analysis in the next section and results in [3] supports the claim that all boundaries are monotone.

We now apply the local time-space formula on curves [8] for $e^{-rs}V_n(t+s,X_s^x)$:

$$(5.24) e^{-r(s-t)}V_{n}(s, X_{s})$$

$$= V_{n}(t, x) + M_{s}$$

$$+ \int_{t}^{s} e^{-r(u-t)} \left(\frac{\partial V_{n}}{\partial t} + \mathbb{I}L_{X}V_{n} - rV_{n} \right) (u, X_{u})I(X_{u} \ge b_{n}(u))du$$

$$+ \frac{1}{2} \int_{t}^{s} e^{-r(u-t)} \left(\frac{\partial V_{n}}{\partial x}(u, X_{u} +) - \frac{\partial V_{n}}{\partial x}(u, X_{u} -) \right) I(X_{u} = b_{n}(u))d\ell_{u}^{b_{n}}(X)$$

$$= V_{n}(t, x) + M_{s} + \int_{t}^{s} e^{-r(u-t)} \left(\frac{\partial G_{n}}{\partial t} + \mathbb{I}L_{X}G_{n} - G_{n}(u, X_{u}) I(X_{u} \ge b_{n}(u))du$$

$$= V_{n}(t, x) + M_{s} + \int_{t}^{s} e^{-r(u-t)}h_{n}(u)X_{u}I(X_{u} \ge b_{n}(u))du$$




Figure 1. This figure plots the prices of multiple reset put options (5.1) for n=1 (green line), n=2 (red), n=3 (blue) and n=4 (black). The parameter set is T=1, K=1, r=0.03, $\delta=0.04$, $\sigma=0.4$.

where we used (5.10), (5.21), the smooth-fit condition (5.12) and where $M = (M_u)_{u \geq 0}$ is the martingale part, $(\ell_u^{b_n}(X))_{u \geq 0}$ is the local time process of X spending at the boundary b_n . Now upon letting s = T, taking the expectation $\mathbb{E}_{t,x}$, the optional sampling theorem for M, rearranging terms, noting that $V_n(T,x) = (K-x)^+$ for all x > 0 and recalling (5.6), we finally get (5.5). One obtains The integral equation (5.4) by simply putting $x = b_n(t)$ into (5.5) and using (5.11).

6. Numerical algorithm and implementation

In this section we provide a numerical algorithm to compute prices of reset put option and optimal reset boundaries. Also we explain how to use these results to provide the sequential reset strategies for the buyer, and finally we plot numerically the multiple optimal reset boundaries and the option prices.

1. Now we describe the numerical solution to multiple reset put option problem (5.1) and provide step-by-step algorithm.

Step 1. We calculate the function h_1 from (4.12).

Step 2. Next, we solve numerically the integral equation (4.24). Let us set $t_k = k\Delta$ for k = 0, 1, ..., N and N large enough where $\Delta = T/N$ so that the following discrete approximation of the integral equation (4.24) is valid

(6.1)
$$G_1(t_k, b_1(t_k)) = V^e(t_k, b_1(t_k)) + \Delta \sum_{l=k}^{N-1} L_1(t_k, t_{l+1}, b_1(t_k), b_1(t_{l+1}))$$

for k=0,1,...,N-1. Setting k=N-1 and $b_1(t_N)=K$ we can solve the algebraic equation (6.1) numerically and get number $b_1(t_{N-1})$. Setting k=N-2 and using the values $b_1(t_{N-1})$,




Figure 2. This figure plots the optimal reset boundaries b_n in (5.1) for n=1 (green line), n=2 (red), n=3 (blue) and n=4 (black). The parameter set is $T=1, K=1, r=0.03, \delta=0.04, \sigma=0.4$.

 $b_1(t_N)$, we can solve (6.1) numerically and get number $b_1(t_{N-2})$. Continuing the recursion we obtain $b_1(t_{N-1}), \ldots, b_1(t_1), b_1(t_0)$ as approximations of the optimal boundary b_1 at the points $T - \Delta, \ldots, \Delta, 0$. We can then interpolate $b_1(\cdot)$ between points (t_{k-1}, t_k) , $k = 1, \ldots, N$.

Step 3. The value function (4.25) can be approximated as follows

(6.2)
$$V_1(t_k, x) = V^e(t_k, x) + \Delta \sum_{l=k}^{N-1} L_1(t_k, t_{l+1}, x, b_1(t_{l+1}))$$

for k=0,1,...,N-1 and x>0, then interpolate $V_1(\cdot,x)$ between points (t_{k-1},t_k) , k=1,...,N.

Step 4. We approximate p_1 from (5.8) as

(6.3)
$$p_1(t_k) = -\Delta \sum_{l=k}^{N-1} e^{-\delta(t_{l+1} - t_k)} h_1(t_{l+1}) \widehat{\mathbb{Q}}_{t_k, K} \left(X_{t_{l+1}} \ge b_1(t_{l+1}) \right)$$

for k=0,1,...,N-1 and interpolate between points (t_{k-1},t_k) , k=1,...,N. We can then approximate p_1' .

Step 5. We compute the function h_2 using (5.7).

Step 6. Now we solve (5.4) for b_2 using the same way as in Step 2 above. All components in the equation (5.4) have been computed in previous steps.

Step 7. We calculate V_2 using (5.5).

Step 8. We compute p_2 and p'_2 using (5.8).

Step 9. We then continue the same procedure as above and compute $h_3 \to b_3 \to V_3 \to (p_3, p_3') \to \ldots \to h_n \to b_n \to V_n \to (p_n, p_n')$.

2. We note that the algorithm above is easy to implement, convergent as N increases, and computationally quite fast. In [4] the numerical comparison of recursive integral approach with

binomial method has been provided for the single reset put option, and it was mentioned that for a given numerical accuracy, the former method is faster than the latter. This observation should be more apparent for multiple reset options studied in the current paper.

Figures 1 and 2 display the sequences of the option prices and the optimal reset boundaries, respectively, for n=4. Figure 1 confirms that the reset option prices increase in the number of remaining reset rights n, which is an obvious fact. Also it shows U-shaped option prices for all $n \geq 1$ unlike the European put option price which is decreasing in x. Figure 2 provides an evidence that for fixed n the optimal reset boundaries are decreasing in t and that for fixed $t \in [0,T)$ the boundaries are decreasing in n.

At last, we point out the following issue. When we obtain the sequence of the boundaries $(b_k(\cdot))_{k=1}^n$ for the reset put option with n rights, for the sake of unity we use the same strike level K. However, using the optimal multiple reset policy the buyer will have different strike for each reset right, i.e., for i-th right the strike is $K_i = X_{\tau_{i-1}}$, i = 2, ..., n and $K_1 = K$. Therefore when the holder has n remaining rights and resets at time t, he immediately obtains the reset put option with n-1 rights and the strike X_t , and thus solves the integral equation for next right with $K = X_t$.

References

- [1] Alobaidi, G., Mallier, R. and Mansi, S. (2011). Laplace transforms and shout options. Acta Math. Univ. Comenianae 80 (79–102).
- [2] CARMONA, R. and Touzi, N. (2008). Optimal multiple stopping and valuation of swing options. Math. Finance 18 (239–268).
- [3] Dai, M., Kwok, Y. K. and Wu, L. (2003). Options with multiple reset rights. *International Journal of Theoretical and Applied Finance* 6 (637–653).
- [4] Dai, M., Kwok, Y. K. and Wu, L. (2004). Optimal shouting policies of options with strike reset right. Math. Finance 14 (383–401).
- [5] DE ANGELIS, T. and KITAPBAYEV, Y. (2018). On the optimal exercise boundaries of swing put options. Math. Oper. Res. 43 (252–274).
- [6] Goard, J. (2012). Exact solutions for a strike reset put option and a shout call option. Math and Computer Modelling 55 (1787–1797).
- [7] Hull, J. C. (2009). Options, Futures and Other Derivatives. 7th ed, Pearson Prentice Hall, New Jersey.
- [8] Peskir, G. (2005). A change-of-variable formula with local time on curves. *J. Theoret. Probab.* 18 (499–535).
- [9] Peskir, G. (2005). On the American option problem. Math. Finance 15 (169–181).
- [10] Peskir, G. and Shiryaev, A. N. (2006). Optimal Stopping and Free-Boundary Problems. Lectures in Mathematics, ETH Zürich, Birkhäuser.

- [11] THOMAS, B. (1993). Something to shout about. Risk 6 (56–58).
- [12] WILMOTT, P. (1998). Derivatives: The Theory and Practice of Financial Engineering. John Wiley and Sons, West Sussex UK.
- [13] WINDCLIFF, H., FORSYTH, P. A. and VETZAL, K. R. (2001). Shout options: a framework for pricing contracts which can be modified by the investor. J. Comput. Appl. Math 134 (213–241).
- [14] WINDCLIFF, H., FORSYTH, P. A. and VETZAL, K. R. (2001). Valuation of aggregated funds; Shout options with maturity extensions. *Insurance Math. Econom.* 29 (1–21).
- [15] WINDCLIFF, H., LE ROUX, M. K., FORSYTH, P. A. and VETZAL, K. R. (2002). Understanding the behaviour and hedging of segregated funds offering the reset feature N. Amer. Actuar. J. 6 (107–125).
- [16] Yang, Z., Yi, F. and Dai, M. (2006). A parabolic variational inequality arising from the valuation of strike reset options. J. Differential Equations 230 (481–501).