

Fractional integration with singularity on unit sphere

Zipeng Wang

Abstract

We study a family of convolution operators whose kernels have a singularity on the unit sphere $\mathbb{S}^{n-1}$. As a result, we prove the regarding $\mathbf{L}^p \rightarrow \mathbf{L}^q$ -Sobolev inequalities.

1 Introduction

In this paper, we revisit on a classical problem that has been previously investigated by Strichartz [3].

Let $\lambda(\alpha) \doteq \frac{n+1}{2} \left(1 - \frac{\alpha}{n}\right)$, $\alpha \in \mathbb{C}$. For $\mathbf{Re} \lambda(\alpha) < 1$, define

$$\Omega^\alpha(x) \doteq \begin{cases} \pi^{-\lambda(\alpha)} \Gamma^{-1}(1 - \lambda(\alpha)) \left(\frac{1}{1 - |x|^2} \right)^{\lambda(\alpha)}, & |x| < 1, \\ 0, & |x| \geq 1 \end{cases} \quad (1.1)$$

whose Fourier transform equals

$$\widehat{\Omega}^\alpha(\xi) = \left(\frac{1}{|\xi|} \right)^{\frac{n}{2} - \lambda(\alpha)} \mathbf{J}_{\frac{n}{2} - \lambda(\alpha)}(2\pi|\xi|). \quad (1.2)$$

Γ and $\mathbf{J}$ denote for Gamma and Bessel functions. See chapter IV in the book by Stein and Weiss [8].

Observe that Ω^α in (1.1) has an non-integrable singularity if $\mathbf{Re} \lambda(\alpha) \geq 1$. However, we can extensively define Ω^α for every $\alpha \in \mathbb{C}$ by its Fourier transform in (1.2).

♦ $\mathbf{L}_s^p$ is the generalized Sobolev space for $s > 0$ and $1 < p < \infty$.

♦ Throughout, we regard $\mathfrak{C}$ as a generic constant depending on its sub-indices.

Theorem One *Let Ω^α defined by its Fourier transform in (1.2) for $0 < \alpha < n$. We have*

$$\|f * \Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} \leq \mathfrak{C}_{p,q} \|f\|_{\mathbf{L}_s^p(\mathbb{R}^n)}, \quad s > 0 \quad (1.3)$$


$$\text{if } \frac{\alpha}{n} = \frac{1}{p} - \frac{1}{q} \quad \text{and} \quad (1.4)$$

$$\frac{n-1}{2n-2+4s} + \left(\frac{4s+n-1}{2n-2+4s} \right) \frac{\alpha}{n} < \frac{1}{p} < \frac{n-1+4s}{2n-2+4s} + \left(\frac{n-1}{2n-2+4s} \right) \frac{\alpha}{n}.$$

In particular, at $s = \frac{1}{2}$, we have

$$\|f * \Omega^\alpha\|_{L^q(\mathbb{R}^n)} \leq \mathfrak{C}_{p,q} \|f\|_{L_{1/2}^p(\mathbb{R}^n)} \quad (1.5)$$

if $\frac{\alpha}{n} = \frac{1}{p} - \frac{1}{q'}$, $\frac{n-1}{2n} + \left(\frac{n+1}{2n}\right)\frac{\alpha}{n} < \frac{1}{p} < \frac{n+1}{2n} + \left(\frac{n-1}{2n}\right)\frac{\alpha}{n}$.



One application of **Theorem One** gives an *a priori* estimate of the wave equation

$$\begin{aligned} \partial_t^2 \mathbf{u}(x, t) - \Delta_x \mathbf{u}(x, t) &= f(x, t), \quad (x, t) \in \mathbb{R}^n \times [0, \infty), \\ \mathbf{u}(x, 0) &= \partial_t \mathbf{u}(x, 0) = 0. \end{aligned} \quad (1.6)$$

Corollary One Let $\mathbf{u}$ be the solution of (1.6). We have

$$\|\mathbf{u}(\cdot, t)\|_{L^q(\mathbb{R}^n)} \leq \mathfrak{C}_{p,q} \int_0^t \|f(\cdot, t-r)\|_{L_s^p(\mathbb{R}^n)} r^{-\frac{n-1}{n+1}} dr, \quad s > 0, \quad t \in [0, \infty)$$

if $\frac{2}{n+1} = \frac{1}{p} - \frac{1}{q}$ and $\frac{n-1}{2n-2+4s} + \left(\frac{4s+n-1}{2n-2+4s}\right)\frac{2}{n+1} < \frac{1}{p} < \frac{n-1+4s}{2n-2+4s} + \left(\frac{n-1}{2n-2+4s}\right)\frac{2}{n+1}$. (1.7)

For more discussion of Sobolev inequalities regarding to wave equations, we refer to the references by Sogge [10]-[11].

Proof As shown by Strichartz [4], it is well known that

$$\widehat{\mathbf{u}}(\xi, t) = \int_0^t \frac{\sin[(t-s)|\xi|]}{|\xi|} \widehat{f}(\xi, s) ds. \quad (1.8)$$

From (1.2), we find $\Omega^{\frac{2n}{n+1}}(\xi) = \frac{1}{\pi} \frac{\sin 2\pi|\xi|}{|\xi|}$. Let $f \in \mathbf{L}_s^p(\mathbb{R}^n)$. By applying **Theorem One**, we have

$$\begin{aligned} \|\mathbf{u}(\cdot, t)\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C} \int_0^t \left\| f(\cdot, t-r) * \Omega_r^{\frac{2n}{n+1}} \right\|_{\mathbf{L}^q(\mathbb{R}^n)} r dr \\ &\leq \mathfrak{C}_{p,q} \int_0^t \|f(\cdot, t-r)\|_{\mathbf{L}^p(\mathbb{R}^n)} r^{1-\frac{2n}{n+1}} dr. \end{aligned} \quad (1.9)$$

□

Theorem Two Let Ω^α defined by its Fourier transform in (1.2) for $0 < \alpha < \left(\frac{n-1}{n+1}\right)n$. We have

$$\begin{aligned} \|f * \Omega^\alpha\|_{\mathbf{L}^p(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,\alpha} \|f\|_{\mathbf{L}_s^p(\mathbb{R}^n)}, \quad s > 0 \\ \text{for } \frac{n-1}{2n-2+4s} - \left(\frac{n+1}{2n-2+4s}\right) \frac{\alpha}{n} &< \frac{1}{p} < \frac{n-1+4s}{2n-2+4s} + \left(\frac{n+1}{2n-2+4s}\right) \frac{\alpha}{n}. \end{aligned} \quad (1.10)$$

Remark One For $\left(\frac{n-1}{n+1}\right)n \leq \alpha < n$, we have

$$\|f * \Omega^\alpha\|_{\mathbf{L}^p(\mathbb{R}^n)} \leq \mathfrak{C}_{p,\alpha} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p < \infty. \quad (1.11)$$

For $s = \frac{1}{2}$ in (1.10), we have

$$\begin{aligned} \|f * \Omega^\alpha\|_{\mathbf{L}^p(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,\alpha} \|f\|_{\mathbf{L}_{1/2}^p(\mathbb{R}^n)} \\ \text{for } \frac{n-1}{2n} - \left(\frac{n+1}{2n}\right) \frac{\alpha}{n} &< \frac{1}{p} < \frac{n+1}{2n} + \left(\frac{n+1}{2n}\right) \frac{\alpha}{n}. \end{aligned} \quad (1.12)$$

2 Principal Lemmata

In order to obtain the desired regularity of $f * \Omega^\alpha$, a key estimate is to study

$$\mathfrak{h}\Omega^\alpha(x) \doteq \begin{cases} \left(\frac{1}{1-|x|^2}\right)^{1-\frac{\alpha}{n}}, & |x| < 1, \\ 0, & |x| \geq 1. \end{cases} \quad (2.1)$$

Lemma One Let $\mathfrak{h}\Omega^\alpha$ defined by (2.1) for $0 < \alpha < n$. We have

$$\begin{aligned} \|f * \mathfrak{h}\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{\alpha,p,q} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p \leq q < \infty \\ \text{if } \frac{\alpha}{n} &\geq \frac{1}{p} - \frac{1}{q}. \end{aligned} \quad (2.2)$$

Let $\delta(\alpha) \doteq 1 - \left(\frac{n+1}{2n}\right)\alpha$ for $0 < \mathbf{Re}\alpha < \frac{2n}{n+1}$. Define

$${}^b\Omega^\alpha(x) \doteq \begin{cases} \pi^{-\delta(\alpha)}\Gamma^{-1}(1 - \delta(\alpha))\left(\frac{1}{1 - |x|^2}\right)^{\delta(\alpha)}, & 0 < |x| < 1, \\ 0, & |x| \geq 1 \end{cases} \quad (2.3)$$

whose Fourier transform equals

$$\begin{aligned} {}^b\widehat{\Omega}^\alpha(\xi) &= \left(\frac{1}{|\xi|}\right)^{\frac{n}{2} - \delta(\alpha)} J_{\frac{n}{2} - \delta(\alpha)}(2\pi|\xi|) \\ &= \left(\frac{1}{|\xi|}\right)^{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2} + \frac{n-1}{2}} J_{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2} + \frac{n-1}{2}}(2\pi|\xi|). \end{aligned} \quad (2.4)$$

◊ In the remaining paragraph, we write $\mathbf{c} > 0$ for some large constant.

Observe that

$$\left| {}^b\Omega^\alpha(x) \right| \leq \mathfrak{C}_\alpha e^{\mathbf{c}|\mathbf{Im}\alpha|} {}^b\Omega^{\left(\frac{n+1}{2}\right)\mathbf{Re}\alpha}(x) \quad (2.5)$$

where ${}^b\Omega^\alpha$ is defined in (2.1).

By using (2.5) and applying **Lemma One**, we have

$$\begin{aligned} \|f * {}^b\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{\alpha \, p \, q} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p \leq q < \infty \\ \text{if } \left(\frac{n+1}{2n}\right)\mathbf{Re}\alpha &\geq \frac{1}{p} - \frac{1}{q}. \end{aligned} \quad (2.6)$$

For $s \geq 0$ and $\mathbf{Re}\lambda(\alpha) + s < 1$, define

$${}^s\Omega^\alpha(x) \doteq \begin{cases} \pi^{-\lambda(\alpha)-s}\Gamma^{-1}(1 - \lambda(\alpha) - s)\left(\frac{1}{1 - |x|^2}\right)^{\lambda(\alpha)+s}, & |x| < 1, \\ 0, & |x| \geq 1 \end{cases} \quad (2.7)$$

whose Fourier transform equals

$$\begin{aligned} {}^s\widehat{\Omega}^\alpha(\xi) &= \left(\frac{1}{|\xi|}\right)^{\frac{n}{2} - \lambda(\alpha) - s} J_{\frac{n}{2} - \lambda(\alpha) - s}(2\pi|\xi|) \\ &= \left(\frac{1}{|\xi|}\right)^{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2} - s} J_{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2} - s}(2\pi|\xi|). \end{aligned} \quad (2.8)$$

Let ω_s defined by $\widehat{\omega}_s(\xi) \doteq \left(\frac{1}{1+|\xi|^2}\right)^{\frac{s}{2}}$ for $s \geq 0$. Note that $\widehat{\omega}_s, s \geq 0$ is a $\mathbf{L}^p$ -Fourier multiplier. Namely, we have

$$\|f * \omega_s\|_{\mathbf{L}^p(\mathbb{R}^n)} \leq \mathfrak{C}_p \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p < \infty. \quad (2.9)$$

Regarding estimates can be found in chapter VI of Stein [7].

Lemma Two Let ${}^s\Omega^\alpha$ defined by its Fourier transform in (2. 8) for $0 < \mathbf{Re}\alpha < n$.

For $0 \leq s \leq \frac{1}{2}$, we have

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q,s} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ \text{if } (1-s)\frac{\mathbf{Re}\alpha}{n} &= \frac{1}{p} - \frac{1}{q}, \quad \frac{1}{2} + \frac{\mathbf{Re}\alpha}{n} \left(\frac{1}{2} - s\right) \leq \frac{1}{p} \leq \frac{1}{2} + \frac{\mathbf{Re}\alpha}{2n}. \end{aligned} \quad (2. 10)$$

For $s \geq \frac{1}{2}$, we have

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q,s} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ \text{if } \frac{\mathbf{Re}\alpha}{2n} &= \frac{1}{p} - \frac{1}{q}, \quad \frac{1}{2} \leq \frac{1}{p} \leq \frac{1}{2} + \frac{\mathbf{Re}\alpha}{2n}. \end{aligned} \quad (2. 11)$$

For every $s \geq 0$, we have

$$\|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^2(\mathbb{R}^n)} \leq \mathfrak{C}_s e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^2(\mathbb{R}^n)}. \quad (2. 12)$$

We prove **Lemma One** in section 3 and **Lemma Two** in section 4. We finish the proof of **Theorem One** and **Theorem Two** in section 5.

3 Proof of Lemma One

We prove the lemma in the same sprit of Hedberg [5]. Let $\mathbf{I}_\alpha^{\natural} f \doteq f * {}^{\natural}\Omega^\alpha$ and assume $f \geq 0$. For every $0 \leq \ell \in \mathbb{Z}$, define

$$\begin{aligned} (\Delta_\ell \mathbf{I}_\alpha^{\natural} f)(x) &\doteq \int_{\mathcal{S}_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du, \\ \mathcal{S}_\ell &\doteq \{u \in \mathbb{R}^n : 2^{-\ell-1} \leq 1-|u| < 2^{-\ell}\}. \end{aligned} \quad (3. 1)$$

Momentarily, we consider the partial sum operator

$$\mathbf{I}_\alpha^\rho \doteq \sum_{0 \leq \ell \leq \rho} \Delta_\ell \mathbf{I}_\alpha^{\natural} \quad \text{for } \rho \text{ sufficiently large.} \quad (3. 2)$$

Let $\{u_\rho^{\mathbf{v}}\}_{\mathbf{v}}$ be a collection of points that are equally distributed on $\mathbb{S}^{n-1} \subset \mathbb{R}^n$ with grid length equal to $2^{-\rho}$ multiplied by a suitable constant. Define the *narrow cone*

$$\Gamma_\rho^{\mathbf{v}} \doteq \left\{ u \in \mathbb{R}^n : \left| \frac{u}{|u|} - u_\rho^{\mathbf{v}} \right| \leq 2^{-\rho} \right\}. \quad (3. 3)$$

Let $\mathbf{R}_{\rho,\ell}^{\mathbf{v}}(u_\rho^{\mathbf{v}})$ be the rectangle centered on $u_\rho^{\mathbf{v}}$ with one side parallel to $u_\rho^{\mathbf{v}}$ of side length $5 \times 2^{-\ell}$ and others perpendicular to $u_\rho^{\mathbf{v}}$ equal to $5 \times 2^{-\rho}$.

Observe that

$$\mathcal{S}_\ell \cap \Gamma_\rho^{\mathbf{v}} \subset \mathbf{R}_{\rho,\ell}^{\mathbf{v}}(u_\rho^{\mathbf{v}}), \quad 0 \leq \ell \leq \rho. \quad (3. 4)$$




Figure 1: $\rho = 5$ and $\ell = 3, 5$.

Let

$$\mathbf{R}_{\rho}^v \doteq \left\{ u \in \mathbb{R}^n : u + u_{\rho}^v \in \mathbf{R}_{\rho}^v(u_{\rho}^v) \right\} \quad (3.5)$$

where

$$\left| \mathbf{R}_{\rho}^v(u_{\rho}^v) \right| = 5^n 2^{-\ell} 2^{-\rho(n-1)}. \quad (3.6)$$

We define

$$\begin{aligned} (\mathbf{M}_{\rho}^v f)(x) &\doteq \sup_{0 \leq \ell \leq \rho} \frac{1}{2^{-\ell} 2^{-\rho(n-1)}} \int_{S_{\ell} \cap \Gamma_{\rho}^v} f(x-u) du \\ &\leq \sup_{0 \leq \ell \leq \rho} \frac{5^n}{|\mathbf{R}_{\rho}^v|} \int_{\mathbf{R}_{\rho}^v} f(x - u_{\rho}^v - u) du. \end{aligned} \quad (3.7)$$

Such a maximal operator defined *w.r.t* all rectangles having a same degree is bounded on $\mathbf{L}^p(\mathbb{R}^n)$ for $1 < p < \infty$.

There are at most $\mathfrak{C}2^{\rho(n-1)}$ elements in $\{u_{\rho}^v\}_v$. Consider

$$(\mathbf{M}_{\rho} f)(x) \doteq 2^{-\rho(n-1)} \sum_v (\mathbf{M}_{\rho}^v f)(x). \quad (3.8)$$

Remark Two $\mathbf{M}_{\rho}$ in (3.8) is NOT the strong maximal operator defined *w.r.t* all rectangles with arbitrary directions. The later has unboundedness on every $\mathbf{L}^p$ -space for $1 \leq p < \infty$ due to the Besicovitch construction. See Fefferman [9] or chapter X and XI in the book by Stein [7].

By using (3.8) and applying Minkowski inequality, we have

$$\|\mathbf{M}_{\rho} f\|_{\mathbf{L}^p(\mathbb{R}^n)} \leq \mathfrak{C}_p \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p < \infty. \quad (3.9)$$

In particular, (3. 9) is independent from ρ .

Let $f \in \mathbf{L}^p(\mathbb{R}^n)$. We claim

$$(\mathbf{I}_\alpha^\rho f)(x) \leq \mathfrak{C}_{p,q} (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}^{1-\frac{p}{q}} \quad (3. 10)$$

for

$$\frac{\alpha}{n} = \frac{1}{p} - \frac{1}{q}, \quad 1 < p < q < \infty. \quad (3. 11)$$

By using the $\mathbf{L}^p$ -boundedness of $\mathbf{M}_\rho$ in (3. 9), we find

$$\begin{aligned} \|\mathbf{I}_\alpha^\rho f\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q} \left\{ \int_{\mathbb{R}^n} (\mathbf{M}_\rho f)^p(x) dx \right\}^{\frac{1}{q}} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}^{1-\frac{p}{q}} \\ &\leq \mathfrak{C}_{p,q} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}. \end{aligned} \quad (3. 12)$$

By allowing $\rho \rightarrow \infty$ in (3. 12) and using Fatou's lemma together with the monotone convergence theorem, we conclude

$$\|\mathbf{I}_\alpha^h f\|_{\mathbf{L}^q(\mathbb{R}^n)} \leq \mathfrak{C}_{p,q} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}. \quad (3. 13)$$

On the other hand, from (2. 1), it is clear that $\mathbf{I}_\alpha^h$ is bounded on $\mathbf{L}^p(\mathbb{R}^n)$ for every $1 \leq p \leq \infty$. The desired result in (2. 2) can be obtained by interpolation.

In order to prove (3. 10), we define $\sigma(x) \in \mathbb{R}$ implicitly by

$$2^{\sigma(x)/p} \doteq \frac{(\mathbf{M}_\rho f)(x)}{\|f\|_{\mathbf{L}^p(\mathbb{R}^n)}}. \quad (3. 14)$$

Let $\mathcal{S}_\ell$ and $\Gamma_\rho^\vee$ be given in (3. 1) and (3. 3). From (3. 4)-(3. 6), we have

$$\begin{aligned} \int_{\mathcal{S}_\ell \cap \Gamma_\rho^\vee} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du &= \int_{\mathcal{S}_\ell \cap \Gamma_\rho^\vee} f(x-u) \left(\frac{1}{1+|u|} \right)^{1-\frac{\alpha}{n}} \left(\frac{1}{1-|u|} \right)^{1-\frac{\alpha}{n}} du \\ &\leq \mathfrak{C} 2^{\ell(1-\frac{\alpha}{n})} \int_{\mathcal{S}_\ell \cap \Gamma_\rho^\vee} f(x-u) du \\ &= \mathfrak{C} 2^{\ell(1-\frac{\alpha}{n})} 2^{-\ell} 2^{-\rho(n-1)} \left\{ \frac{1}{2^{-\ell} 2^{-\rho(n-1)}} \int_{\mathcal{S}_\ell \cap \Gamma_\rho^\vee} f(x-u) du \right\} \\ &= \mathfrak{C} 2^{-\ell(\frac{\alpha}{n})} 2^{-\rho(n-1)} \left\{ \frac{1}{2^{-\ell} 2^{-\rho(n-1)}} \int_{\mathcal{S}_\ell \cap \Gamma_\rho^\vee} f(x-u) du \right\} \\ &\leq \mathfrak{C} 2^{-\ell(\frac{\alpha}{n})} 2^{-\rho(n-1)} (\mathbf{M}_\rho^\vee f)(x). \end{aligned} \quad (3. 15)$$

Therefore,

$$\begin{aligned}
\int_{S_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du &\leq \sum_{\mathbf{v}} \int_{S_\ell \cap \Gamma_{\rho}^{\mathbf{v}}} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du \\
&\leq \mathfrak{C} 2^{-\ell(\frac{\alpha}{n})} 2^{-\rho(n-1)} \sum_{\mathbf{v}} (\mathbf{M}_{\rho}^{\mathbf{v}} f)(x) \\
&= \mathfrak{C} 2^{-\ell(\frac{\alpha}{n})} (\mathbf{M}_{\rho} f)(x).
\end{aligned} \tag{3.16}$$

On the other hand, by applying Hölder inequality, we have

$$\int_{S_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du \leq \|f\|_{L^p(\mathbb{R}^n)} \left\{ \int_{S_\ell} \left(\frac{1}{1-|u|^2} \right)^{(1-\frac{\alpha}{n})(\frac{p}{p-1})} du \right\}^{\frac{p-1}{p}} \tag{3.17}$$

where

$$\begin{aligned}
\left\{ \int_{S_\ell} \left(\frac{1}{1-|u|^2} \right)^{(1-\frac{\alpha}{n})(\frac{p}{p-1})} du \right\}^{\frac{p-1}{p}} &= \left\{ \int_{S_\ell} \left(\frac{1}{1+|u|} \right)^{(1-\frac{\alpha}{n})(\frac{p}{p-1})} \left(\frac{1}{1-|u|} \right)^{(1-\frac{\alpha}{n})(\frac{p}{p-1})} du \right\}^{\frac{p-1}{p}} \\
&\leq \mathfrak{C} \left[\frac{1}{2^{-\ell(1-\frac{\alpha}{n})(\frac{p}{p-1})}} 2^{-\ell} \right]^{\frac{p-1}{p}} = \mathfrak{C} 2^{\ell(1-\frac{\alpha}{n})} 2^{-\ell(\frac{p-1}{p})} = \mathfrak{C} 2^{-\ell(\frac{\alpha}{n}-\frac{1}{p})}.
\end{aligned} \tag{3.18}$$

Consider $\sigma(x) \leq \ell \leq \rho$. By inserting (3.14) into (3.16), we find

$$\begin{aligned}
\int_{S_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du &\leq \mathfrak{C} 2^{-\ell(\frac{\alpha}{n})} (\mathbf{M}_\ell f)(x) \\
&= \mathfrak{C} 2^{(\sigma(x)-\ell)(\frac{\alpha}{n})} 2^{-\sigma(x)(\frac{1}{p}-\frac{1}{q})} (\mathbf{M}_\rho f)(x) \\
&= \mathfrak{C} 2^{(\sigma(x)-\ell)(\frac{\alpha}{n})} \left\{ \frac{\|f\|_{L^p(\mathbb{R}^n)}}{(\mathbf{M}_\rho f)(x)} \right\}^{1-\frac{p}{q}} (\mathbf{M}_\rho f)(x) \\
&= \mathfrak{C} 2^{(\sigma(x)-\ell)(\frac{\alpha}{n})} (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{L^p(\mathbb{R}^n)}^{1-\frac{p}{q}}.
\end{aligned} \tag{3.19}$$

By summing over all $\sigma(x) \leq \ell \leq \rho$, we have

$$\begin{aligned}
\sum_{\sigma(x) \leq \ell \leq \rho} \int_{S_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du &\leq \mathfrak{C} \sum_{\sigma(x) \leq \ell \leq \rho} 2^{(\sigma(x)-\ell)(\frac{\alpha}{n})} (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{L^p(\mathbb{R}^n)}^{1-\frac{p}{q}} \\
&\leq \mathfrak{C}_p (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{L^p(\mathbb{R}^n)}^{1-\frac{p}{q}}.
\end{aligned} \tag{3.20}$$

Consider $0 \leq \ell < \sigma(x)$. By inserting (3. 14) into (3. 17)-(3. 18), we find

$$\begin{aligned}
\int_{\mathcal{S}_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du &\leq \mathfrak{C} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} 2^{-\ell(\frac{\alpha}{n}-\frac{1}{p})} \\
&= \mathfrak{C} 2^{(\sigma(x)-\ell)(\frac{\alpha}{n}-\frac{1}{p})} 2^{-\sigma(x)(\frac{\alpha}{n}-\frac{1}{p})} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \\
&= \mathfrak{C} 2^{-(\sigma(x)-\ell)(\frac{1}{q})} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \left\{ \frac{\|f\|_{\mathbf{L}^p(\mathbb{R}^n)}}{(\mathbf{M}_\rho f)(x)} \right\}^{-\frac{p}{q}} \\
&= \mathfrak{C} 2^{-(\sigma(x)-\ell)(\frac{1}{q})} (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}^{1-\frac{p}{q}}.
\end{aligned} \tag{3. 21}$$

By summing over all $0 \leq \ell < \sigma(x)$, we have

$$\begin{aligned}
\sum_{0 \leq \ell < \sigma(x)} \int_{\mathcal{S}_\ell} f(x-u) \left(\frac{1}{1-|u|^2} \right)^{1-\frac{\alpha}{n}} du \\
\leq \mathfrak{C} \sum_{0 \leq \ell < \sigma(x)} 2^{-(\sigma(x)-\ell)(\frac{1}{q})} (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}^{1-\frac{p}{q}} \\
\leq \mathfrak{C}_{p,q} (\mathbf{M}_\rho f)^{\frac{p}{q}}(x) \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}^{1-\frac{p}{q}}.
\end{aligned} \tag{3. 22}$$

By putting together (3. 20) and (3. 22), we obtain (3. 10) as required.

4 Proof of Lemma Two

We begin this section by recalling some well known estimates of Bessel functions. More discussions can be found in the book by Watson [12].

- For $\mu > -\frac{1}{2}, \nu \in \mathbb{R}$ and $\rho > 0$, a Bessel function can be defined by

$$J_{\mu+\mathrm{i}\nu}(\rho) = \frac{(\rho/2)^{\mu+\mathrm{i}\nu}}{\pi^{\frac{1}{2}} \Gamma(\mu + \frac{1}{2} + \mathrm{i}\nu)} \int_{-1}^1 e^{\mathrm{i}\rho s} (1-s^2)^{\mu-\frac{1}{2}+\mathrm{i}\nu} ds. \tag{4. 1}$$

- For $\mu > -\frac{1}{2}, \nu \in \mathbb{R}$ and $\rho > 0$,

$$J_{\mu+\mathrm{i}\nu}(\rho) = \left(\frac{2}{\pi\rho} \right)^{\frac{1}{2}} \cos\left(\rho - \frac{\pi}{2}(\mu + \mathrm{i}\nu) - \frac{\pi}{4}\right) + \mathbf{e}_{\mu+\mathrm{i}\nu}(\rho) \tag{4. 2}$$

where

$$|\mathbf{e}_{\mu+\mathrm{i}\nu}(\rho)| \leq \mathfrak{C}_\mu e^{\mathbf{c}|\nu|} \begin{cases} \rho^{-\frac{1}{2}}, & 0 < \rho \leq 1, \\ \rho^{-\frac{3}{2}}, & \rho > 1. \end{cases} \tag{4. 3}$$

- For every $\mu, \nu \in \mathbb{R}$ and $\rho > 0$, we have the identity

$$\left(\frac{\mu + i\nu}{\rho}\right) \mathbf{J}_{\mu+i\nu}(\rho) = \mathbf{J}_{\mu-1+i\nu}(\rho) + \mathbf{J}_{\mu+1+i\nu}(\rho). \quad (4.4)$$

All together, (4. 1), (4. 2)-(4. 3) and (4. 4) imply:

- For every $\mu, \nu \in \mathbb{R}$ and $\rho > 0$,

$$\left|\rho^{-(\mu+i\nu)} \mathbf{J}_{\mu+i\nu}(\rho)\right| \leq \mathfrak{C}_\mu \left(\frac{1}{1+\rho}\right)^{\frac{1}{2}+\mu} e^{c|\nu|}. \quad (4.5)$$

Let ${}^s\Omega^\alpha$ defined in (2. 7) for $\mathbf{Re}\lambda(\alpha) + s < 1$. We have

$$\left|{}^s\Omega^{n+i\mathbf{Im}\alpha}(x)\right| \leq \mathfrak{C}_s e^{c|\mathbf{Im}\alpha|} \left(\frac{1}{1-|x|^2}\right)^s, \quad 0 \leq s < 1. \quad (4.6)$$

By applying **Lemma One** and then using (2. 9), we obtain

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^{n+i\mathbf{Im}\alpha}\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q,s} e^{c|\mathbf{Im}\alpha|} \|f * \omega_s\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ &\leq \mathfrak{C}_{p,q,s} e^{c|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p < q < \infty \end{aligned} \quad (4.7)$$

$$\text{if} \quad 1-s = \frac{1}{p} - \frac{1}{q}, \quad 0 < s < 1.$$

From (4. 6), it is also clear that

$$\|f * \Omega^{n+i\mathbf{Im}\alpha}\|_{\mathbf{L}^\infty(\mathbb{R}^n)} \leq \mathfrak{C} e^{c|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^1(\mathbb{R}^n)}. \quad (s=0) \quad (4.8)$$

Let ${}^s\Omega^\alpha$ defined by its Fourier transform in (2. 8) for $s \geq 0$ and $\alpha \in \mathbb{C}$. From (4. 5), we have

$$\left|\widehat{{}^s\Omega^\alpha}(\xi)\right| \leq \mathfrak{C}_{\alpha,s} e^{c|\mathbf{Im}\alpha|} \left(\frac{1}{1+|\xi|}\right)^{\left(\frac{n+1}{2n}\right)\mathbf{Re}\alpha-s}. \quad (4.9)$$

By using (4. 9) and applying Plancherel theorem, we find

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^2(\mathbb{R}^n)} &\leq \mathfrak{C}_s e^{c|\mathbf{Im}\alpha|} \|f * \omega_s\|_{\mathbf{L}^2(\mathbb{R}^n)} \\ &\leq \mathfrak{C}_s e^{c|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^2(\mathbb{R}^n)} \quad \text{by (2. 9)}. \end{aligned} \quad (4.10)$$

In particular, we have

$$\|f * \omega_s * {}^s\Omega^{0+i\mathbf{Im}\alpha}\|_{\mathbf{L}^2(\mathbb{R}^n)} \leq \mathfrak{C}_s e^{c|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^2(\mathbb{R}^n)}. \quad (4.11)$$

Let $0 < \mathbf{Re}\alpha < n$ and $0 \leq s \leq \frac{1}{2}$. By using (4. 7)-(4. 8) and (4. 11) then applying Stein interpolation theorem [6], we have

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q,s} e^{c|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ \text{if} \quad (1-s)\frac{\mathbf{Re}\alpha}{n} &= \frac{1}{p} - \frac{1}{q}, \quad \frac{1}{2} + \frac{\mathbf{Re}\alpha}{n} \left(\frac{1}{2} - s\right) \leq \frac{1}{p} \leq \frac{1}{2} + \frac{\mathbf{Re}\alpha}{2n}. \end{aligned} \quad (4.12)$$

Let $0 < \mathbf{Re} \alpha < n$ and $s \geq \frac{1}{2}$. From (4. 9), we find

$$\begin{aligned} \left| \widehat{f}(\xi) \widehat{\omega}_s(\xi) {}^s\widehat{\Omega}^\alpha(\xi) \right| &\leq \mathfrak{C}_{\alpha s} e^{\mathbf{c}|\mathbf{Im} \alpha|} \left| \widehat{f}(\xi) \right| \widehat{\omega}_s(\xi) (1 + |\xi|)^s \left(\frac{1}{1 + |\xi|} \right)^{\left(\frac{n+1}{2n}\right)\mathbf{Re} \alpha} \\ &\leq \mathfrak{C}_{\alpha s} e^{\mathbf{c}|\mathbf{Im} \alpha|} \left| \widehat{f}(\xi) \right| \left(\frac{1}{|\xi|} \right)^{\mathbf{Re} \left(\frac{n}{2}\right)}. \end{aligned} \quad (4. 13)$$

Note that the inverse Fourier transform of $|\xi|^{-\alpha}$ equals $\mathfrak{C}_\alpha |x|^{\alpha-n}$ for $0 < \alpha < n$.

By applying Hardy-Littlewood-Sobolev inequality [1]-[2] together with Plancherel theorem, we simultaneously have

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{q s} e^{\mathbf{c}|\mathbf{Im} \alpha|} \|f\|_{\mathbf{L}^2(\mathbb{R}^n)} \quad \text{if} \quad \frac{\mathbf{Re} \alpha}{2n} = \frac{1}{2} - \frac{1}{q}, \\ \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^2(\mathbb{R}^n)} &\leq \mathfrak{C}_{p s} e^{\mathbf{c}|\mathbf{Im} \alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \quad \text{if} \quad \frac{\mathbf{Re} \alpha}{2n} = \frac{1}{p} - \frac{1}{2}. \end{aligned} \quad (4. 14)$$

From (4. 14), by applying Riesz-Thorin interpolation theorem, we find

$$\begin{aligned} \|f * \omega_s * {}^s\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p q s} e^{\mathbf{c}|\mathbf{Im} \alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ \text{if} \quad \frac{\mathbf{Re} \alpha}{2n} &= \frac{1}{p} - \frac{1}{q}, \quad \frac{1}{2} \leq \frac{1}{p} \leq \frac{1}{2} + \frac{\mathbf{Re} \alpha}{2n}. \end{aligned} \quad (4. 15)$$

5 Interpolation on a family of analytic operators

Consider a family of analytic operators $f * \omega_s * \Theta_z^{\alpha s}$ for $0 < \mathbf{Re} z < 1$ where

$$\widehat{\Theta}_z^{\alpha s}(\xi) \doteq \left(\frac{1}{|\xi|} \right)^{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2} + \left(\frac{n-1}{2}\right)z - s(1-z)} \mathbf{J}_{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2} + \left(\frac{n-1}{2}\right)z - s(1-z)}(2\pi|\xi|). \quad (5. 1)$$

In particular, we have

$$\begin{aligned} \widehat{\Theta}_0^{\alpha s}(\xi) &= {}^b\widehat{\Omega}^\alpha(\xi), \quad \widehat{\Theta}_1^{\alpha s}(\xi) = {}^s\widehat{\Omega}^\alpha(\xi) \\ \text{and} \quad \widehat{\Theta}_{\frac{2s}{n-1+2s}}^{\alpha s}(\xi) &= \left(\frac{1}{|\xi|} \right)^{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2}} \mathbf{J}_{\left(\frac{n+1}{2n}\right)\alpha - \frac{1}{2}}(2\pi|\xi|) = \widehat{\Omega}^\alpha(\xi) \quad \text{by (1. 2)}. \end{aligned} \quad (5. 2)$$

Let $0 < \mathbf{Re} \alpha < \frac{2n}{n+1}$. As a special case of (2. 6), for $0 \leq s \leq \frac{1}{2}$, we have

$$\begin{aligned} \|f * \omega_s * {}^b\Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p q} e^{\mathbf{c}|\mathbf{Im} \alpha|} \|f * \omega_s\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ &\leq \mathfrak{C}_{p q} e^{\mathbf{c}|\mathbf{Im} \alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \quad \text{by (2. 9)} \\ \text{if} \quad \left(\frac{n+1}{2n} \right) \mathbf{Re} \alpha &= \frac{1}{p} - \frac{1}{q}, \quad 1 < p < q < \infty. \end{aligned} \quad (5. 3)$$

On the other hand, for $s \geq \frac{1}{2}$, we have

$$\begin{aligned}
\|f * \omega_s * \Omega^\alpha\|_{L^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q} e^{c|\operatorname{Im}\alpha|} \|f * \omega_s\|_{L^p(\mathbb{R}^n)} \\
&\leq \mathfrak{C}_{p,q} e^{c|\operatorname{Im}\alpha|} \|f\|_{L^p(\mathbb{R}^n)}, \quad \text{by (2.9)} \\
\text{if } \quad \left(\frac{n-1+4s}{4s}\right) \frac{\operatorname{Re}\alpha}{n} &= \frac{1}{p} - \frac{1}{q}, \quad 1 < p < q < \infty.
\end{aligned} \tag{5.4}$$

Recall **Lemma Two**. Let $0 < s \leq \frac{1}{2}$. By using (2.10) and (5.3) respectively for $f * \omega_s * \Theta_z^{\alpha,s}$ defined as (5.1)-(5.2) and applying Stein interpolation theorem [6], we have

$$\begin{aligned}
\|f * \omega_s * \Omega^\alpha\|_{L^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q} e^{c|\operatorname{Im}\alpha|} \|f\|_{L^p(\mathbb{R}^n)}, \quad 1 < p < q < \infty \\
\text{if } \quad \frac{1}{p} - \frac{1}{q} &= \left(\frac{2s}{n-1+2s}\right) \left(\frac{n+1}{2n}\right) \operatorname{Re}\alpha + \left(\frac{n-1}{n-1+2s}\right) \left(\frac{1-s}{n}\right) \operatorname{Re}\alpha = \frac{\operatorname{Re}\alpha}{n} \\
\text{and } \quad \left(\frac{2s}{n-1+2s}\right) \left(\frac{n+1}{2n}\right) \operatorname{Re}\alpha &+ \left(\frac{n-1}{n-1+2s}\right) \left[\frac{1}{2} + \frac{\operatorname{Re}\alpha}{n} \left(\frac{1}{2} - s\right)\right] \\
&= \frac{n-1}{2n-2+4s} + \left(\frac{4s+n-1}{2n-2+4s}\right) \frac{\operatorname{Re}\alpha}{n} < \frac{1}{p} < \\
\left(\frac{2s}{n-1+2s}\right) + \left(\frac{n-1}{n-1+2s}\right) \left[\frac{1}{2} + \frac{\operatorname{Re}\alpha}{2n}\right] &= \frac{n-1+4s}{2n-2+4s} + \left(\frac{n-1}{2n-2+4s}\right) \frac{\operatorname{Re}\alpha}{n}.
\end{aligned} \tag{5.5}$$

Let $s \geq \frac{1}{2}$. By using (2.11) and (5.4) respectively for $f * \omega_s * \Theta_z^{\alpha,s}$ defined as (5.1)-(5.2) and applying Stein interpolation theorem [6], we have

$$\begin{aligned}
\|f * \omega_s * \Omega^\alpha\|_{L^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q} e^{c|\operatorname{Im}\alpha|} \|f\|_{L^p(\mathbb{R}^n)}, \quad 1 < p < q < \infty \\
\text{if } \quad \frac{1}{p} - \frac{1}{q} &= \left(\frac{2s}{n-1+2s}\right) \left(\frac{n-1+4s}{4s}\right) \frac{\operatorname{Re}\alpha}{n} + \left(\frac{n-1}{n-1+2s}\right) \frac{\operatorname{Re}\alpha}{2n} = \frac{\operatorname{Re}\alpha}{n} \\
\text{and } \quad \left(\frac{2s}{n-1+2s}\right) \left(\frac{n-1+4s}{4s}\right) \frac{\operatorname{Re}\alpha}{n} &+ \left(\frac{n-1}{n-1+2s}\right) \frac{1}{2} \\
&= \frac{n-1}{2n-2+4s} + \left(\frac{4s+n-1}{2n-2+4s}\right) \frac{\operatorname{Re}\alpha}{n} < \frac{1}{p} < \\
\left(\frac{2s}{n-1+2s}\right) + \left(\frac{n-1}{n-1+2s}\right) \left[\frac{1}{2} + \frac{\operatorname{Re}\alpha}{2n}\right] &= \frac{n-1+4s}{2n-2+4s} + \left(\frac{n-1}{2n-2+4s}\right) \frac{\operatorname{Re}\alpha}{n}.
\end{aligned} \tag{5.6}$$

By putting together (5.5) and (5.6), we conclude

$$\begin{aligned}
\|f * \omega_s * \Omega^\alpha\|_{L^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p,q} e^{c|\operatorname{Im}\alpha|} \|f\|_{L^p(\mathbb{R}^n)} \\
\text{if } \quad \frac{\operatorname{Re}\alpha}{n} &= \frac{1}{p} - \frac{1}{q} \quad \text{and} \\
\frac{n-1}{2n-2+4s} + \left(\frac{4s+n-1}{2n-2+4s}\right) \frac{\operatorname{Re}\alpha}{n} &< \frac{1}{p} < \frac{n-1+4s}{2n-2+4s} + \left(\frac{n-1}{2n-2+4s}\right) \frac{\operatorname{Re}\alpha}{n}
\end{aligned} \tag{5.7}$$

for $s > 0$ and $0 < \mathbf{Re}\alpha < \frac{2n}{n+1}$.

On the other hand, from (4. 12) and (2. 9), we have

$$\begin{aligned} \|f * \omega_s * \Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p\ q} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f * \omega_s\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ &\leq \mathfrak{C}_{p\ q} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \end{aligned} \quad (5. 8)$$

$$\text{for } \frac{1}{p} = \frac{1}{2} + \frac{\mathbf{Re}\alpha}{2n}, \quad \frac{1}{q} = \frac{1}{2} - \frac{\mathbf{Re}\alpha}{2n}$$

and $0 < \mathbf{Re}\alpha < n$.

By using (5. 7)-(5. 8) and applying Stein interpolation theorem [6], we find

$$\begin{aligned} \|f * \omega_s * \Omega^\alpha\|_{\mathbf{L}^q(\mathbb{R}^n)} &\leq \mathfrak{C}_{p\ q} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad s > 0 \\ \text{if } \frac{\alpha}{n} &= \frac{1}{p} - \frac{1}{q} \quad \text{and} \end{aligned} \quad (5. 9)$$

$$\frac{n-1}{2n-2+4s} + \left(\frac{4s+n-1}{2n-2+4s}\right) \frac{\alpha}{n} < \frac{1}{p} < \frac{n-1+4s}{2n-2+4s} + \left(\frac{n-1}{2n-2+4s}\right) \frac{\alpha}{n}.$$

Note that $f * \Omega^\alpha = \mathfrak{F} * \omega_s * \Omega^\alpha$ of which $\widehat{\mathfrak{F}}(\xi) \doteq \widehat{f}(\xi) (1 + |\xi|^2)^{\frac{s}{2}}$. We have $\mathfrak{F} \in \mathbf{L}^p(\mathbb{R}^n)$ provided that $f \in \mathbf{L}_s^p(\mathbb{R}^n)$. This completes the proof of **Theorem One**.

Let $0 < \mathbf{Re}\alpha < \frac{2n}{n+1}$. As a special case of (2. 6), for every $s \geq 0$, we have

$$\begin{aligned} \|f * \omega_s * {}^b\Omega^\alpha\|_{\mathbf{L}^p(\mathbb{R}^n)} &\leq \mathfrak{C}_{p\ \alpha} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f * \omega_s\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ &\leq \mathfrak{C}_{p\ \alpha} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)}, \quad 1 < p < \infty \quad \text{by (2. 9)}. \end{aligned} \quad (5. 10)$$

Recall **Lemma Two**. Let $s > 0$. By using (2. 12) and (5. 10) respectively for $f * \omega_s * \Theta_z^{\alpha\ s}$ defined as (5. 1)-(5. 2) and applying Stein interpolation theorem [6], we have

$$\begin{aligned} \|f * \omega_s * \Omega^\alpha\|_{\mathbf{L}^p(\mathbb{R}^n)} &\leq \mathfrak{C}_{p\ \alpha} e^{\mathbf{c}|\mathbf{Im}\alpha|} \|f\|_{\mathbf{L}^p(\mathbb{R}^n)} \\ \text{for } \left(\frac{n-1}{n-1+2s}\right) \frac{1}{2} + 0 &= \frac{n-1}{2n-2+4s} < \frac{1}{p} < \\ \left(\frac{n-1}{n-1+2s}\right) \frac{1}{2} + \frac{2s}{n-1+2s} &= \frac{n-1+4s}{2n-2+4s}. \end{aligned} \quad (5. 11)$$

Let Ω^α defined in (1. 1) for $\mathbf{Re}\lambda(\alpha) = \frac{n+1}{2} \left(1 - \frac{\mathbf{Re}\alpha}{n}\right) < 1$. Observe that

$$|\Omega^\alpha(x)| \leq \pi^{-\mathbf{Re}\lambda(\alpha)} |\Gamma^{-1}(1 - \lambda(\alpha))| |\Omega^{\left(\frac{n+1}{2}\right)\mathbf{Re}\alpha - \left(\frac{n-1}{2}\right)n}(x)| \quad (5. 12)$$

for $\left(\frac{n-1}{n+1}\right)n < \mathbf{Re}\alpha < n$ where ${}^b\Omega^\alpha$ is defined in (2. 1) for $0 < \alpha < n$.

By using (5. 12) and applying **Lemma One**, we have

$$\begin{aligned} \|f * \omega_s * \Omega^\alpha\|_{L^p(\mathbb{R}^n)} &\leq \mathfrak{C}_{p-\alpha} e^{c|\operatorname{Im}\alpha|} \|f * \omega_s\|_{L^p(\mathbb{R}^n)} \\ &\leq \mathfrak{C}_{p-\alpha} e^{c|\operatorname{Im}\alpha|} \|f\|_{L^p(\mathbb{R}^n)}, \quad 1 < p < \infty \quad \text{by (2. 9)} \end{aligned} \quad (5. 13)$$

whenever $\left(\frac{n-1}{n+1}\right)n < \operatorname{Re}\alpha < n$. Note that $\widehat{\Omega}^\alpha(\xi)$ at $\alpha = \left(\frac{n-1}{n+1}\right)n$ is the Fourier transform of the surface measure on $\mathbb{S}^{n-1}$. We therefore conclude **Remark One**.

By using (5. 11) with (5. 13) and applying Stein interpolation theorem [6], we have

$$\begin{aligned} \|f * \omega_s * \Omega^\alpha\|_{L^p(\mathbb{R}^n)} &\leq \mathfrak{C}_{p-\alpha} \|f\|_{L^p(\mathbb{R}^n)} \\ \text{for } \left[1 - \frac{\alpha}{n} \left(\frac{n+1}{n-1}\right)\right] \frac{n-1}{2n-2+4s} &= \frac{n-1}{2n-2+4s} - \frac{\alpha}{n} \left(\frac{n+1}{2n-2+4s}\right) < \frac{1}{p} < \\ \left[1 - \frac{\alpha}{n} \left(\frac{n+1}{n-1}\right)\right] \frac{n-1+4s}{2n-2+4s} + \frac{\alpha}{n} \left(\frac{n+1}{n-1}\right) &= \frac{n-1+4s}{2n-2+4s} + \frac{\alpha}{n} \left(\frac{n+1}{2n-2+4s}\right) \end{aligned} \quad (5. 14)$$

and $0 < \alpha < \left(\frac{n-1}{n+1}\right)n$.

Lastly, $f * \Omega^\alpha = \mathfrak{F} * \omega_s * \Omega^\alpha$ where $\widehat{\mathfrak{F}}(\xi) = \widehat{f}(\xi) \left(1 + |\xi|^2\right)^{\frac{s}{2}}$. Moreover, $\mathfrak{F} \in L^p(\mathbb{R}^n)$ because $f \in L_s^p(\mathbb{R}^n)$. We finish the proof of **Theorem Two**.

References

- [1] G. H. Hardy and J. E. Littlewood, *Some Properties of Fractional Integrals*, *Mathematische Zeitschrift* **27**: 565-606, 1928.
- [2] S. L. Sobolev, *On a Theorem of Functional Analysis*, *Matematicheskii Sbornik* **46**: 471-497, 1938.
- [3] R. Strichartz, *Convolutions with kernels having singularity on a sphere*, *Transaction of the American Mathematical Society* **148**: 461-471, 1970.
- [4] R. Strichartz, *A priori estimates for the wave equation and some applications*, *Journal of Functional Analysis* **5**: 218-235, 1970.
- [5] L. Hedberg, *On Certain Convolution Inequalities*, *Proceeding of American Mathematical Society* **36**: 505-510, 1972.
- [6] E. M. Stein, *Interpolation of linear operators*, *Transaction of the American Mathematical Society* **83**: 482-492, 1956.
- [7] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton University Press, 1993.
- [8] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Space*, Princeton University Press, 1971.
- [9] C. Fefferman, *A note on Spherical Summation Multipliers*, *Israel Journal of Mathematics*, **15**: 44-52, 1973.

- [10] C. D. Sogge, *L^p estimates for the wave equation and applications*, Journées Équations aux dérivées partielles, 1-12, 1993.
- [11] C. D. Sogge, *Lectures on Non-linear Wave Equations* (second edition), Johns Hopkins University Press, 2013.
- [12] G. N. Watson, *Theory of Bessel Functions*, Cambridge University Press, Cambridge, 1944.

Department of Mathematics, Westlake University.

email: wangzipeng@westlake.edu.cn