

Liouville-type theorems, radial symmetry and integral representation of solutions to Hardy-Hénon equations involving higher order fractional Laplacians

Hui Yang*

September 20, 2021

Abstract

We study nonnegative solutions to the following Hardy-Hénon type equations involving higher order fractional Laplacians

$$(-\Delta)^\sigma u = |x|^{-\alpha} u^p \quad \text{in } \mathbb{R}^n \setminus \{0\}$$

which have a possible singularity at the origin, where σ is a real number satisfying $0 < \sigma < \frac{n}{2}$, $-\infty < \alpha < 2\sigma$ and $p > 1$. By a more direct approach without using the super poly-harmonic properties of nonnegative solutions, we establish an integral representation for nonnegative solutions to the above higher order fractional equations whether the singularity $\{0\}$ is removable or not. As the first application, we prove Liouville theorems for nonnegative solutions to the above equations with removable singularity for all $\sigma \in (0, \frac{n}{2})$ when

$$1 < p < p_{\sigma, \alpha}^* := \frac{n + 2\sigma - 2\alpha}{n - 2\sigma} \quad \text{and} \quad -\infty < \alpha < 2\sigma.$$

This, in particular, covers a gap occurring for non-integral $\sigma \in (1, \frac{n}{2})$ and $\alpha \in (0, 2\sigma)$ in the current literature. As the second application, we show the radial symmetry of nonnegative solutions to the above equations in the critical case or in the case where the singularity at the origin is not removable. Such radial symmetry would be useful in studying the higher order singular Yamabe problems and characterizing isolated singularities of higher order Hardy-Hénon equations.

Keywords: Higher order fractional Laplacians, integral representation, Liouville-type theorems, radial symmetry, Hardy-Hénon equations

Mathematics Subject Classification (2020): 35J30; 35R11; 35B06; 35C15

*Department of Mathematics, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, China. E-mail address: mahuiyang@ust.hk

1 Introduction and main results

Problems involving the fractional Laplacian operator and other nonlocal diffusion operators have been extensively studied in recent years. Such nonlocal problems arise naturally in mathematical models from various physical phenomena, such as anomalous diffusion and quasi-geostrophic flows, turbulence and water waves, molecular dynamics, and relativistic quantum mechanics of stars (see [6, 11, 22] and the references therein). Moreover, the fractional Laplacian and higher order Laplacian also appear in many areas of mathematics, including conformal geometry [14, 15], probability and finance [3, 23], obstacle problems [8, 55], nonlinear elasticity and crystal dislocation [1, 27].

In this paper, we focus on the nonnegative solutions to the following Hardy-Hénon type equations involving higher order fractional Laplacians

$$(-\Delta)^\sigma u = |x|^{-\alpha} u^p \quad \text{in } \mathbb{R}^n \setminus \{0\} \quad (1.1)$$

which have a possible singularity at the origin, where σ is a real number satisfying $0 < \sigma < \frac{n}{2}$, $-\infty < \alpha < 2\sigma$ and $p > 1$. The fractional Laplacian is defined, for $\sigma \in (0, 1)$, by

$$(-\Delta)^\sigma u(x) = c_{n,\sigma} \text{P.V.} \int_{\mathbb{R}^n} \frac{u(x) - u(z)}{|x - z|^{n+2\sigma}} dz \quad (1.2)$$

while, for higher powers, say $\sigma = k + \sigma'$ with $k \in \mathbb{N}$ and $\sigma' \in (0, 1)$,

$$(-\Delta)^\sigma = (-\Delta)^{\sigma'} \circ (-\Delta)^k. \quad (1.3)$$

When $\sigma = 1$, $0 \leq \alpha < 2$ and $p = 2^*(\alpha) - 1$ with $2^*(\alpha) := \frac{2(n-\alpha)}{n-2}$, Eq. (1.1) arises as an Euler-Lagrange equation of a functional associated with the following Hardy-Sobolev inequality: There exists $C > 0$ such that for all $u \in \mathcal{D}^{1,2}(\mathbb{R}^n)$,

$$\left(\int_{\mathbb{R}^n} \frac{|u|^{2^*(\alpha)}}{|x|^\alpha} dx \right)^{\frac{2}{2^*(\alpha)}} \leq C \int_{\mathbb{R}^n} |\nabla u|^2 dx. \quad (1.4)$$

In particular, when $\alpha = 0$, (1.4) becomes the classical Sobolev inequality whose best constant and extremal functions have been identified by Aubin [4] and Talenti [56]. Furthermore, the best constant in the Hardy-Sobolev inequality (1.4) was first computed by Glaser-Martin-Grosse-Thirring [34] and extremal functions were identified by Lieb [48]. For general $\sigma \in (0, \frac{n}{2})$, Eq. (1.1) with $0 \leq \alpha < 2\sigma$ is also closely related to the fractional Hardy-Sobolev inequality (see, e.g., [51]). On the other hand, Eq. (1.1) has been used in models of astrophysics (see, e.g., [38]) when $\sigma = 1$ and $\alpha < 0$.

Notice also that Eq. (1.1) with $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$ corresponds to the fractional Yamabe problem in conformal geometry [14, 15]. To be more specific, denote $g = u^{\frac{4}{n-2\sigma}} |dx|^2$ the conformal change of the Euclidean metric $|dx|^2$ for some positive function u , then one can construct the fractional order conformal Laplacian associated with g , which satisfies

$$P_\sigma^g(\cdot) = u^{-\frac{n+2\sigma}{n-2\sigma}} (-\Delta)^\sigma(u \cdot). \quad (1.5)$$

The fractional Q -curvature of g is given by

$$Q_\sigma^g := P_\sigma^g(1) = u^{-\frac{n+2\sigma}{n-2\sigma}}(-\Delta)^\sigma u. \quad (1.6)$$

Thus, a solution u of (1.1) with $\alpha = 0$ and $p = (n + 2\sigma)/(n - 2\sigma)$ induces a conformally flat metric $g = u^{\frac{4}{n-2\sigma}}|dx|^2$ which has constant fractional Q -curvature $Q_\sigma^g = 1$. In particular, when $\sigma = 1$, curvature (1.6) is the scalar curvature modulo a positive constant, while for $\sigma = 2$, it coincides with the Branson's Q -curvature corresponding to the Paneitz operator.

Eq. (1.1) that involves the higher order fractional Laplacians has both nonlocal feature and the difficulty of higher order elliptic equations. When $\sigma = m \in (0, n/2)$ is an integer and $\alpha = 0$, Wei and Xu [58] proved the following super poly-harmonic properties for positive classical solutions of (1.1) in \mathbb{R}^n

$$(-\Delta)^i u \geq 0 \quad \text{in } \mathbb{R}^n, \quad i = 1, 2, \dots, m - 1, \quad (1.7)$$

and they further applied these information to classify the entire solutions of (1.1) in \mathbb{R}^n for $1 < p \leq \frac{n+2m}{n-2m}$. See the same results by Lin [49] in the case of $m = 2$. Later, Chen, Li and Ou in [19] used the super poly-harmonic properties (1.7) to establish an integral representation formula for positive entire solutions of (1.1) in \mathbb{R}^n and gave a new proof to the classification results of Lin [49] and Wei-Xu [58] via the corresponding integral equation. We may also see [21, 24, 25] for the similar super poly-harmonic properties and integral representation of nonnegative classical solutions to (1.1) in \mathbb{R}^n when $\sigma = m \in (0, n/2)$ is an integer and $-\infty < \alpha < 2m$. Moreover, an integral representation for nonnegative entire solutions to (1.1) on \mathbb{R}^n in the fractional setting ($0 < \sigma < 1$) has also been obtained in [5, 25, 63] recently to prove Liouville type theorems. Thus, a natural question is whether one can establish an integral representation for nonnegative solutions of (1.1) when $\sigma \in (1, n/2)$ is not an integer? In a recent paper [12], Cao, Dai and Qin showed the following super poly-harmonic properties

$$(-\Delta)^{i+\frac{\nu}{2}} u \geq 0 \quad \text{in } \mathbb{R}^n, \quad i = 0, 1, \dots, m - 1 \quad (1.8)$$

for nonnegative classical solutions of (1.1) in \mathbb{R}^n with $\alpha \leq 0$, where σ is rewritten as $\sigma = m + \frac{\nu}{2}$ for some integer $m \geq 1$ and some $0 < \nu < 2$. Based on the super poly-harmonic properties (1.8), they further proved an integral representation formula for nonnegative classical solutions of (1.1) in \mathbb{R}^n when $1 \leq \sigma < n/2$ and $\alpha \leq 0$. In fact, they studied higher order fractional equations with general continuous nonlinearities which include $|x|^{-\alpha} u^p$ ($\alpha \leq 0$) as a special case.

In this paper, by a more direct approach without using the super poly-harmonic properties (1.8), we will establish an integral representation of nonnegative solutions to the higher order fractional equations (1.1) for all $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Moreover, our results also differ from those in [12] in the following two respects:

1. We consider the higher order fractional equation (1.1) with the Hardy type nonlinearity $|x|^{-\alpha} u^p$ ($0 < \alpha < 2\sigma$), which is not continuous at the origin.
2. Our results apply not only to the nonnegative entire solutions of (1.1) in \mathbb{R}^n , but also to its singular solutions (which have a non-removable singularity at the origin).

For the sake of application, we understand a solution u of (1.1) in the sense of distributions. For $s \in \mathbb{R}$, we define

$$\mathcal{L}_s(\mathbb{R}^n) := \left\{ u \in L_{loc}^1(\mathbb{R}^n) : \int_{\mathbb{R}^n} \frac{|u(x)|}{(1 + |x|)^{n+2s}} dx < \infty \right\}.$$

We say that u is a nonnegative distributional solution of (1.1) in $\mathbb{R}^n \setminus \{0\}$ if $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$, $|\cdot|^{-\alpha}u^p \in L^1_{loc}(\mathbb{R}^n \setminus \{0\})$, $u \geq 0$, and it satisfies (1.1) in the following sense:

$$\int_{\mathbb{R}^n} u(x)(-\Delta)^\sigma \varphi(x) dx = \int_{\mathbb{R}^n} |x|^{-\alpha} u(x)^p \varphi(x) dx \quad \text{for any } \varphi \in C_c^\infty(\mathbb{R}^n \setminus \{0\}). \quad (1.9)$$

Similarly, u is a nonnegative distributional solution of (1.1) in \mathbb{R}^n if $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$, $|\cdot|^{-\alpha}u^p \in L^1_{loc}(\mathbb{R}^n)$, $u \geq 0$, and it satisfies (1.1) in the entire \mathbb{R}^n in the following sense:

$$\int_{\mathbb{R}^n} u(x)(-\Delta)^\sigma \varphi(x) dx = \int_{\mathbb{R}^n} |x|^{-\alpha} u(x)^p \varphi(x) dx \quad \text{for any } \varphi \in C_c^\infty(\mathbb{R}^n). \quad (1.10)$$

Then we have the following integral characterization for nonnegative solutions to the equation (1.1) when $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$.

Theorem 1.1. *Let $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Suppose that one of the following holds:*

- (1) $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ is a nonnegative distributional solution of (1.1) in \mathbb{R}^n with $p > 1$.
- (2) $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n \setminus \{0\})$ is a nonnegative distributional solution of (1.1) in $\mathbb{R}^n \setminus \{0\}$ with $p \geq \frac{n-\alpha}{n-2\sigma}$.

Then u satisfies the integral equation

$$u(x) = C_{n,\sigma} \int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^\alpha |x-y|^{n-2\sigma}} dy \quad \text{for } x \in \mathbb{R}^n \setminus \{0\} \quad (1.11)$$

with some positive constant $C_{n,\sigma}$.

Remark 1.2. *When $\sigma \in (0, n/2)$ is an integer, instead of the assumption $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$ in Theorem 1.1, we only need $u \in L^1_{loc}(\mathbb{R}^n)$ since we have a better bound in this setting by the rescaling method (see Proposition 2.6 in Section 2). Moreover, if the assumption (1) of Theorem 1.1 is satisfied, then u satisfies the integral equation (1.11) for all $x \in \mathbb{R}^n$.*

Note that Theorem 1.1 holds for both entire solutions and singular solutions near the origin, and only requires the growth restriction $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$ near infinity which is slightly weaker than the growth restriction $u \in \mathcal{L}_\nu(\mathbb{R}^n)$ for some $0 < \nu < 2$ satisfying $\sigma = m + \frac{\nu}{2}$ used in [12] when $\sigma > 1$ is not an integer. Moreover, as pointed out in Remark 1.2 above, such a growth restriction is not required in the case where σ is an integer.

For $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$, as a direct result of Theorem 1.1 and the classification results for positive $L^{2n/(n-2\sigma)}_{loc}(\mathbb{R}^n)$ solutions of the integral equation (1.11) in Chen-Li-Ou [19] or Li [44], we give a classification for nonnegative solutions to the following conformally invariant equation with higher order fractional Laplacians

$$(-\Delta)^\sigma = u^{\frac{n+2\sigma}{n-2\sigma}} \quad \text{in } \mathbb{R}^n. \quad (1.12)$$

Theorem 1.3. *Let $0 < \sigma < n/2$. Suppose that $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ is a nonnegative distributional solution of (1.12) in \mathbb{R}^n . Then either $u \equiv 0$ or u must be of the form*

$$u(x) = \gamma_{n,\sigma} \left(\frac{\mu}{\mu^2 + |x - x_0|^2} \right)^{\frac{n-2\sigma}{2}} \quad (1.13)$$

for a fixed constant $\gamma_{n,\sigma} > 0$ and for some $\mu > 0$ and $x_0 \in \mathbb{R}^n$.

It is well-known that such classification results have many important applications in conformal geometry (see, e.g., [40,41] and the references therein). When $\sigma = m \in (0, n/2)$ is an integer, Theorem 1.3 has been proved by Caffarelli-Gidas-Spruck [7] for $m = 1$, Lin [49] for $m = 2$ and Wei-Xu [58] for $m \geq 3$. When $\sigma \in (0, 1)$, Theorem 1.3 was proved by Chen-Li-Li [18] and Jin-Li-Xiong [40]. For general $\sigma \in (0, n/2)$, Chen, Li and Ou in [19] showed Theorem 1.3 for nonnegative weak solutions $u \in H^\sigma(\mathbb{R}^n)$ of (1.12) by using an integral representation. For non-integral $\sigma \in (1, n/2)$, Cao, Dai and Qin in [12] recently proved Theorem 1.3 for nonnegative classical solutions $u \in C_{loc}^{2\sigma+\varepsilon}(\mathbb{R}^n) \cap \mathcal{L}_{\frac{\nu}{2}}(\mathbb{R}^n)$ of (1.12) with $\sigma = m + \frac{\nu}{2}$ for some $0 < \nu < 2$, where $\varepsilon > 0$ is arbitrarily small. Clearly, such weak solutions and classical solutions of (1.12) both satisfy the definition of distributional solutions in this paper. Thus, our Theorem 1.3 in a slightly weaker assumption gives the classification of nonnegative solutions to (1.12) in higher order fractional cases.

Next, as an application of Theorem 1.1, we prove a Liouville type theorem for nonnegative solutions of (1.1) in \mathbb{R}^n for all $\sigma \in (0, n/2)$ and $\alpha \in (-\infty, 2\sigma)$ in the following subcritical range

$$1 < p < p_{\sigma,\alpha}^* := \frac{n + 2\sigma - 2\alpha}{n - 2\sigma}, \quad (1.14)$$

where $p_{\sigma,\alpha}^*$ is a critical exponent.

Theorem 1.4. *Let $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Suppose that $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ is a nonnegative distributional solution of (1.1) in \mathbb{R}^n with $1 < p < p_{\sigma,\alpha}^*$. Then $u \equiv 0$ in \mathbb{R}^n .*

Liouville type theorems for elliptic equations have been extensively studied, which, in particular, are crucial in deriving a priori estimates and then establishing existence of positive solutions to non-variational boundary value problems. When $\alpha = 0$, Liouville-type result in Theorem 1.4 has been proved in [7, 18, 33, 40, 44, 49, 58, 63] for different values of $\sigma \in (0, n/2)$. For the Hardy-Hénon's case $\alpha \neq 0$, Theorem 1.4 was showed in [25, 36, 52, 57] for $\sigma = 1$, in [5, 25] for $0 < \sigma < 1$ and in [25] when $\sigma = m \in (1, n/2)$ is an integer. One may also see [21, 24, 28] and the references therein for some earlier partial results. For the case where $\sigma \in (1, n/2)$ is not an integer, Theorem 1.4 with $\alpha \leq 0$ was recently proved by Cao, Dai and Qin in [12] for nonnegative classical solutions $u \in C_{loc}^{2\sigma+\varepsilon}(\mathbb{R}^n) \cap \mathcal{L}_{\frac{\nu}{2}}(\mathbb{R}^n)$ with $\sigma = m + \frac{\nu}{2}$ for some $0 < \nu < 2$. Hence, Theorem 1.4 not only covers the existing gap when $0 < \alpha < 2\sigma$ and $\sigma \in (1, n/2)$ is not an integer, but also gives a unified approach for both $\alpha < 0$ and $\alpha \geq 0$ in higher order fractional cases.

Finally, applying Theorem 1.1 we study radial symmetry of nonnegative solutions to (1.1) in \mathbb{R}^n with critical exponent or in $\mathbb{R}^n \setminus \{0\}$. Symmetry is a fundamental property in elliptic PDEs which is especially important in searching solutions, classifying solutions, characterizing asymptotic behavior of solutions and so on. In particular, radial symmetry of solutions to an elliptic equation on \mathbb{R}^n plays a crucial role in the classification of entire solutions; see, for example, [7, 16, 17, 32, 49, 58] and the references therein. On the other hand, the importance of studying singular solutions of Yamabe-type equations have been highlighted in the classical works of Schoen and Yau [53, 54] on conformally flat manifolds. In general, radial symmetry of singular solutions of a Yamabe-type equation on $\mathbb{R}^n \setminus \{0\}$ is an important step to understand asymptotic behavior for the corresponding equation near its isolated singularity [7, 37, 43]. The recent works of Yang and Zou [60–62] also showed that radial symmetry of global singular solutions has an important application in characterizing precise asymptotic behavior of solutions to fractional equations

near isolated singularities, which, in particular, helps to overcome the difficulties caused by the lack of ODEs analysis for fractional equations.

We say that the origin $\{0\}$ is a non-removable singularity of the solution u of (1.1) if u cannot be extended to a continuous function near the origin. Based on Theorem 1.1, we show the following radial symmetry of nonnegative solutions to (1.1) with Hardy-type weights ($0 \leq \alpha < 2\sigma$).

Theorem 1.5. *Let $0 < \sigma < n/2$ and $0 \leq \alpha < 2\sigma$. Suppose that $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n \setminus \{0\})$ is a nonnegative distributional solution of (1.1) in $\mathbb{R}^n \setminus \{0\}$, and u has a non-removable singularity at the origin when $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$.*

- (1) *If $\frac{n-\alpha}{n-2\sigma} \leq p \leq \frac{n+2\sigma-2\alpha}{n-2\sigma}$, then u is radially symmetric and monotonically decreasing with respect to the origin.*
- (2) *If $\frac{n+2\sigma-2\alpha}{n-2\sigma} < p \leq \frac{n+2\sigma-\alpha}{n-2\sigma}$, then u is radially symmetric with respect to the origin.*

Clearly, Theorem 1.5 allows a solution u to be singular at the origin. In the case $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$, all of the solutions of (1.1) with removable singularity have been classified in Theorem 1.3. Note also that, for $0 < \alpha < 2\sigma$, the exponent $\frac{n+2\sigma-\alpha}{n-2\sigma}$ in Theorem 1.5 is bigger than the critical exponent $p_{\sigma,\alpha}^* = \frac{n+2\sigma-2\alpha}{n-2\sigma}$. As a special case, by Theorem 1.5 and Remark 1.2 before, we have the following symmetry result for higher order Hardy-Hénon type equations.

Corollary 1.6. *Let $0 < m < n/2$ be an integer and $0 \leq \alpha < 2m$. Suppose that $u \in C(\mathbb{R}^n \setminus \{0\})$ is a nonnegative distributional solution of*

$$(-\Delta)^m u = |x|^{-\alpha} u^p \quad \text{in } \mathbb{R}^n \setminus \{0\}, \quad (1.15)$$

and u has a non-removable singularity at the origin when $\alpha = 0$ and $p = \frac{n+2m}{n-2m}$. If $\frac{n-\alpha}{n-2m} \leq p \leq \frac{n+2m-\alpha}{n-2m}$, then u is radially symmetric with respect to the origin.

As mentioned earlier, these radial symmetry results would be useful in studying the higher order singular Yamabe problems and characterizing isolated singularities of higher order Hardy-Hénon equations. In the case $\alpha = 0$, the radial symmetry in Theorem 1.5 has been showed by Caffarelli-Gidas-Spruck [7] for $\sigma = 1$, by Lin [49] for $\sigma = 2$ and by Caffarelli-Jin-Sire-Xiong [9] for $0 < \sigma < 1$. In the case $0 < \alpha < 2\sigma$, Theorem 1.5 was proved by Li-Bao [47] for positive classical solutions of (1.1) with non-removable singularity at the origin when $0 < \sigma < 1$ and $\frac{n-\alpha}{n-2\sigma} \leq p \leq \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Note that [9, 47] applied the well-known extension formula for the fractional Laplacian introduced by Caffarelli-Silvestre [10] to prove these symmetry results. Moreover, when $0 < \alpha < 2\sigma$, Theorem 1.5 was established by Lu-Zhu [50] for positive weak solutions $u \in H^\sigma(\mathbb{R}^n)$ of (1.1) in \mathbb{R}^n with the critical exponent $p = p_{\sigma,\alpha}^* = \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Hence, on the one hand, instead of Caffarelli-Silvestre's extension, we use a unified approach to show radial symmetry of singular positive solutions of (1.1) for all $\sigma \in (0, n/2)$ and $\alpha \in [0, 2\sigma)$. On the other hand, our Theorem 1.5 in a weaker condition establishes the radial symmetry of entire solutions to (1.1) in \mathbb{R}^n when $0 < \alpha < 2\sigma$ and $p_{\sigma,\alpha}^* \leq p \leq \frac{n+2\sigma-\alpha}{n-2\sigma}$.

In the case $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$, it is well-known that $c|x|^{-\frac{n-2\sigma}{2}}$ is a singular solution of (1.1) with a positive constant c depending only on n and σ , the other singular solutions, e.g., the Fowler solutions, of (1.1) were obtained by Jin-Xiong [42]. In the same case, the existence of Fowler-type singular solutions of (1.1) for $0 < \sigma < 1$ and $\sigma = 2$ was showed earlier by DelaTorre-del

Pino-González-Wei [26] and Guo-Huang-Wang-Wei [35], respectively. We also remark that the classification of singular positive solutions to (1.1) with $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$ was established by Fowler [30] for $\sigma = 1$ and by Frank-König [31] for $\sigma = 2$.

The rest of this paper is organized as follows. In Section 2, we establish the integral representation of nonnegative solutions to (1.1) in Theorem 1.1. Section 3 and Section 4 are devoted to proving Theorem 1.4 and Theorem 1.5, respectively.

In the following, we will use $B_r(x)$ to denote the open ball of radius r in \mathbb{R}^n with center x and sometimes write $B_r(0)$ as B_r for short.

Acknowledgements. The author would like to thank Professor Tianling Jin for his support and encouragement.

2 An integral characterization

In this section, we prove Theorem 1.1 which gives an integral characterization of distributional solutions to (1.1) for all $\sigma \in (0, n/2)$. Firstly, we show, under suitable assumptions, that a distributional solution u of (1.1) is always in $L^p_{loc}(\mathbb{R}^n, |x|^{-\alpha} dx)$ and satisfies the equation (1.1) on the whole \mathbb{R}^n in the sense of distributions. Let $\eta_1, \eta_2 \in C^\infty(\mathbb{R}^n)$ be two cut-off functions satisfying $0 \leq \eta_1 \leq 1$ and

$$\eta_1(x) = \begin{cases} 0 & \text{for } |x| \leq 1, \\ 1 & \text{for } |x| \geq 2, \end{cases} \quad \eta_2(x) := 1 - \eta_1(x).$$

We define

$$\eta_\varepsilon(x) := \eta_{1,\varepsilon}(x)\eta_2(x), \quad \text{where } \eta_{1,\varepsilon}(x) := \eta_1\left(\frac{x}{\varepsilon}\right) \text{ for } \varepsilon > 0. \quad (2.1)$$

Then one has the following estimate for $(-\Delta)^\sigma \eta_\varepsilon$, whose proof can be found in [2, Lemma 2.1].

Lemma 2.1. *Let η_ε be as in (2.1). Then for every $\sigma > 0$ we have*

$$|(-\Delta)^\sigma \eta_\varepsilon(x)| \leq \frac{C_\sigma}{\varepsilon^{2\sigma}} \frac{1}{\left(1 + \frac{|x|}{\varepsilon}\right)^{n+2\sigma}} + \frac{C_\sigma}{(1 + |x|)^{n+2\sigma}} \quad \text{for all } x \in \mathbb{R}^n. \quad (2.2)$$

Using Lemma 2.1 and a dyadic decomposition argument as in Ao-González-Hyder-Wei [2], we prove the following result which has also been obtained in [2] when $\alpha = 0$. See also [7, 42, 59] for the case with $\sigma \in (0, n/2)$ being an integer.

Proposition 2.2. *Suppose that $0 < \sigma < n/2$ and $-\infty < \alpha < +\infty$. Let $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n \setminus \{0\})$ be a nonnegative distributional solution of (1.1) in $\mathbb{R}^n \setminus \{0\}$ with $p > 1$. Suppose in addition that*

$$p \geq \frac{n - \alpha}{n - 2\sigma}.$$

Then $|\cdot|^{-\alpha} u^p \in L^1_{loc}(\mathbb{R}^n)$ and u is a distributional solution to the equation (1.1) in \mathbb{R}^n , i.e.,

$$\int_{\mathbb{R}^n} u(x)(-\Delta)^\sigma \varphi(x) dx = \int_{\mathbb{R}^n} |x|^{-\alpha} u(x)^p \varphi(x) dx \quad \text{for any } \varphi \in C_c^\infty(\mathbb{R}^n). \quad (2.3)$$

Proof. Let η_ε be defined in (2.1) with $\varepsilon \in (0, 1/2)$. Taking η_ε as a test function for the equation (1.1) and using Lemma 2.1 yields

$$\begin{aligned} \int_{\mathbb{R}^n} |x|^{-\alpha} u^p \eta_\varepsilon dx &= \int_{\mathbb{R}^n} u(-\Delta)^\sigma \eta_\varepsilon dx \\ &\leq C + C\varepsilon^{-2\sigma} \int_{\mathbb{R}^n} \frac{u(x)}{\left(1 + \frac{|x|}{\varepsilon}\right)^{n+2\sigma}} dx \\ &\leq C + C\varepsilon^{-2\sigma} \left(\int_{B_{2\varepsilon}} + \int_{B_1 \setminus B_{2\varepsilon}} + \int_{B_1^c} \right) \frac{u(x)}{\left(1 + \frac{|x|}{\varepsilon}\right)^{n+2\sigma}} dx \\ &\leq C + C\varepsilon^{-2\sigma} \left(\int_{B_{2\varepsilon}} u dx + \int_{B_1 \setminus B_{2\varepsilon}} \frac{u(x)}{\left(1 + \frac{|x|}{\varepsilon}\right)^{n+2\sigma}} dx + \varepsilon^{n+2\sigma} \right). \end{aligned}$$

By Hölder inequality with $\frac{1}{p} + \frac{1}{p'} = 1$, we have that for $p > 1$,

$$\begin{aligned} \int_{B_1 \setminus B_{2\varepsilon}} \frac{u(x)}{\left(1 + \frac{|x|}{\varepsilon}\right)^{n+2\sigma}} dx &\leq \varepsilon^{n+2\sigma} \left(\int_{B_1 \setminus B_{2\varepsilon}} |x|^{-\alpha} u^p dx \right)^{1/p} \left(\int_{B_1 \setminus B_{2\varepsilon}} \frac{|x|^{\alpha p'/p}}{(\varepsilon + |x|)^{p'(n+2\sigma)}} dx \right)^{1/p'} \\ &\leq \begin{cases} C\varepsilon^{n+(\alpha-n)/p} \left(\int_{B_1 \setminus B_{2\varepsilon}} |x|^{-\alpha} u^p dx \right)^{1/p} & \text{if } p > (\alpha - n)/2\sigma, \\ -C\varepsilon^{n+2\sigma} \ln \varepsilon \left(\int_{B_1 \setminus B_{2\varepsilon}} |x|^{-\alpha} u^p dx \right)^{1/p} & \text{if } p \leq (\alpha - n)/2\sigma. \end{cases} \end{aligned}$$

Hence, for $p > 1$ and $p \geq \frac{n-\alpha}{n-2\sigma}$ we have

$$\int_{\mathbb{R}^n} |x|^{-\alpha} u^p \eta_\varepsilon dx \leq C + C\varepsilon^{-2\sigma} \int_{B_{2\varepsilon}} u dx + C \left(\int_{B_1 \setminus B_{2\varepsilon}} |x|^{-\alpha} u^p dx \right)^{1/p},$$

and using Young inequality we obtain

$$\int_{\mathbb{R}^n} |x|^{-\alpha} u^p \eta_\varepsilon dx \leq C + C\varepsilon^{-2\sigma} \int_{B_{2\varepsilon}} u dx. \quad (2.4)$$

Now, for every integer $i \geq -1$, we get from Hölder inequality and (2.4) that

$$\begin{aligned} \int_{\{\frac{\varepsilon}{2^{i+1}} \leq |x| \leq \frac{\varepsilon}{2^i}\}} u dx &\leq \left(\frac{\varepsilon}{2^i}\right)^{n-(n-\alpha)/p} \left(\int_{\{\frac{\varepsilon}{2^{i+1}} \leq |x| \leq \frac{\varepsilon}{2^i}\}} |x|^{-\alpha} u^p dx \right)^{1/p} \\ &\leq C \left(\frac{\varepsilon}{2^i}\right)^{n-(n+2\sigma-\alpha)/p} \left(\int_{\{|x| \leq \frac{\varepsilon}{2^{i+1}}\}} u dx \right)^{1/p} + C \left(\frac{\varepsilon}{2^i}\right)^{n-(n-\alpha)/p} \\ &\leq C \left(\frac{\varepsilon}{2^i}\right)^{n-(n+2\sigma-\alpha)/p} \left(\int_{\{|x| \leq \varepsilon\}} u dx \right)^{1/p} + C \left(\frac{\varepsilon}{2^i}\right)^{n-(n-\alpha)/p}. \end{aligned}$$

Note that we always have $n - (n + 2\sigma - \alpha)/p > 0$ since $n - (n + 2\sigma - \alpha)/p > 0 \iff p > (n + 2\sigma - \alpha)/n$ which is true by

$$\begin{cases} p > 1 \geq (n + 2\sigma - \alpha)/n & \text{if } (n + 2\sigma - \alpha)/n \leq 1 \\ p \geq (n - \alpha)/(n - 2\sigma) > (n + 2\sigma - \alpha)/n & \text{if } (n + 2\sigma - \alpha)/n > 1. \end{cases}$$

Thus, we sum the above inequality for all integer $i \geq -1$ to get

$$\int_{B_{2\varepsilon}} u dx \leq C\varepsilon^{n-(n+2\sigma-\alpha)/p} \left(\int_{\{|x| \leq \varepsilon\}} u dx \right)^{1/p} + C\varepsilon^{n-(n-\alpha)/p}.$$

Using Young inequality again, we obtain

$$\int_{B_{2\varepsilon}} u dx \leq C\varepsilon^{n-\frac{2\sigma-\alpha}{p-1}} + C\varepsilon^{n-\frac{n-\alpha}{p}}. \quad (2.5)$$

Since $n - \frac{2\sigma-\alpha}{p-1} \geq 2\sigma$ and $n - \frac{n-\alpha}{p} \geq 2\sigma$ due to $p \geq \frac{n-\alpha}{n-2\sigma}$, combining (2.5) with (2.4) and sending $\varepsilon \rightarrow 0$, we obtain that $\int_{B_1} |x|^{-\alpha} u^p dx < \infty$ and hence, $|\cdot|^{-\alpha} u^p \in L^1_{\text{loc}}(\mathbb{R}^n)$.

To complete the proof, we next prove that u is a distributional solution of (1.1) in the whole space \mathbb{R}^n . Since $|\cdot|^{-\alpha} u^p \in L^1_{\text{loc}}(\mathbb{R}^n)$, we only need to show that for every $\varphi \in C_c^\infty(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n} u(-\Delta)^\sigma(\varphi\eta_{1,\varepsilon}) dx \rightarrow \int_{\mathbb{R}^n} u(-\Delta)^\sigma \varphi dx \quad (2.6)$$

as $\varepsilon \rightarrow 0$, where $\eta_{1,\varepsilon}$ is defined as in (2.1) with $\varepsilon \in (0, 1/4)$. For this purpose, we define

$$I_\varepsilon(x) := (-\Delta)^\sigma(\varphi\eta_{1,\varepsilon})(x) - \eta_{1,\varepsilon}(x)(-\Delta)^\sigma \varphi(x), \quad x \in \mathbb{R}^n.$$

Let k be the integer part of σ , that is, $\sigma := k + \sigma'$ with $\sigma' \in (0, 1)$ and $k \in \{0\} \cup \mathbb{N}$. We consider the two cases $|x| \geq 4\varepsilon$ and $|x| \leq 4\varepsilon$ to estimate $I_\varepsilon(x)$.

Case 1: $|x| \geq 4\varepsilon$. In this case, we note that all derivatives of $\eta_{1,\varepsilon}$ at x are equal to 0. Then we have (for simplicity we omit the constant $c_{n,\sigma}$ in (1.2))

$$\begin{aligned} (-\Delta)^\sigma(\varphi\eta_{1,\varepsilon})(x) &= \int_{\mathbb{R}^n} \frac{\eta_{1,\varepsilon}(x)[(-\Delta)^k \varphi(x) - (-\Delta)^k \varphi(z)] + (-\Delta)^k[\varphi(z)(1 - \eta_{1,\varepsilon}(z))]}{|x-z|^{n+2\sigma'}} dz \\ &= \eta_{1,\varepsilon}(x)(-\Delta)^\sigma \varphi(x) + \int_{\mathbb{R}^n} \frac{(-\Delta)^k[\varphi(z)(1 - \eta_{1,\varepsilon}(z))]}{|x-z|^{n+2\sigma'}} dz \\ &= \eta_{1,\varepsilon}(x)(-\Delta)^\sigma \varphi(x) + C \int_{\mathbb{R}^n} \frac{\varphi(z)(1 - \eta_{1,\varepsilon}(z))}{|x-z|^{n+2\sigma}} dz \end{aligned}$$

where the integration by parts is used in the last equality. Notice also that the integrand is not singular at $z = x$ since the function $1 - \eta_{1,\varepsilon}$ is supported in $B_{2\varepsilon}$. Hence we have

$$|I_\varepsilon(x)| \leq \int_{B_{2\varepsilon}} \frac{|\varphi(z)(1 - \eta_{1,\varepsilon}(z))|}{|x-z|^{n+2\sigma}} dz \leq C \frac{\varepsilon^n}{|x|^{n+2\sigma}} \quad \text{for } |x| \geq 4\varepsilon. \quad (2.7)$$

Case 2: $|x| \leq 4\varepsilon$. In this case, instead of estimating $I_\varepsilon(x)$ we estimate $(-\Delta)^\sigma(\varphi\eta_{1,\varepsilon})$. Denote $\varphi_\varepsilon := \varphi\eta_{1,\varepsilon}$. In fact, we have

$$(-\Delta)^\sigma \varphi_\varepsilon(x) = \frac{1}{2} \int_{\mathbb{R}^n} \frac{2(-\Delta)^k \varphi_\varepsilon(x) - (-\Delta)^k \varphi_\varepsilon(x+y) - (-\Delta)^k \varphi_\varepsilon(x-y)}{|y|^{n+2\sigma'}} dy.$$

Moreover, there exists $C > 0$ depending on φ such that the integrand in the above integral can be controlled by

$$\frac{C}{\varepsilon^{2k}|y|^{n+2\sigma'}} \min \left\{ \frac{|y|^2}{\varepsilon^2}, 1 \right\}.$$

Thus, we obtain

$$|(-\Delta)^\sigma \varphi_\varepsilon(x)| \leq \frac{C}{\varepsilon^{2\sigma}} \quad \text{for } |x| \leq 4\varepsilon.$$

Hence

$$|I_\varepsilon(x)| \leq \frac{C}{\varepsilon^{2\sigma}} \quad \text{for } |x| \leq 4\varepsilon \quad (2.8)$$

due to the boundedness of $\eta_{1,\varepsilon}(-\Delta)^\sigma \varphi$. Combining (2.7) and (2.8), we get

$$|I_\varepsilon(x)| \leq C \frac{1}{\varepsilon^{2\sigma}} \frac{\chi_{B_1}(x)}{(1 + \frac{|x|}{\varepsilon})^{n+2\sigma}} + C\varepsilon^n \frac{\chi_{B_1^c}(x)}{|x|^{n+2\sigma}} \quad \text{for all } x \in \mathbb{R}^n. \quad (2.9)$$

This is obviously true if σ is an integer. For $p' > 1$ with $\frac{1}{p} + \frac{1}{p'} = 1$, we have

$$\left\| |\cdot|^{-\frac{\alpha}{p}} I_\varepsilon \right\|_{L^{p'}(B_1)}^{p'} \leq C\varepsilon^{-2\sigma p'} \int_{B_\varepsilon} |x|^{\frac{\alpha p'}{p}} dx + C\varepsilon^{np'} \int_{\{\varepsilon \leq |x| \leq 1\}} |x|^{\frac{\alpha p'}{p} - (n+2\sigma)p'} \leq C$$

with some $C > 0$ independent of ε . Hence, for each $\delta \in (0, 1)$,

$$\int_{B_\delta} u(x) |I_\varepsilon(x)| dx \leq \left\| |\cdot|^{-\frac{\alpha}{p}} u \right\|_{L^p(B_\delta)} \left\| |\cdot|^{-\frac{\alpha}{p}} I_\varepsilon \right\|_{L^{p'}(B_\delta)} \leq C \left\| |\cdot|^{-\frac{\alpha}{p}} u \right\|_{L^p(B_\delta)}$$

uniformly in ε and δ . On the other hand, by the assumption $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$ and (2.9), we have that for any fixed $\delta \in (0, 1)$,

$$\lim_{\varepsilon \rightarrow 0} \int_{B_\delta^c} u(x) |I_\varepsilon(x)| dx = 0.$$

Therefore, by sending $\delta \rightarrow 0$ we obtain

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^n} u(x) |I_\varepsilon(x)| dx = 0.$$

This implies that (2.6) holds and thus, u is a distributional solution to the equation (1.1) in \mathbb{R}^n . \square

Now we give some growth estimates for solutions of (1.1) near infinity. We start by recalling a result from [2, Lemma 5.2].

Lemma 2.3. *Let $\psi \in C^\infty(\mathbb{R}^n)$ be such that $\psi(x) = \frac{1}{|x|^\tau}$ on B_1^c for some $\tau > 0$. Let $\xi \in C^\infty(\mathbb{R}^n)$ be a cut-off function satisfying*

$$\xi(x) = 1 \quad \text{for } |x| \leq 1 \quad \text{and} \quad \xi(x) = 0 \quad \text{for } |x| \geq 2.$$

For any $\varepsilon > 0$, we define $\xi_\varepsilon := \xi(\varepsilon x)$ and $\psi_\varepsilon(x) := \psi \xi_\varepsilon(x)$. Then for every $\sigma > 0$ we have

$$(-\Delta)^\sigma \psi_\varepsilon \rightarrow (-\Delta)^\sigma \psi \quad \text{locally uniformly in } \mathbb{R}^n \text{ as } \varepsilon \rightarrow 0.$$

Moreover, there exists $C = C(n, \sigma, \varphi) > 0$ (independent of ε) such that

$$|(-\Delta)^\sigma \psi_\varepsilon(x)| \leq C \begin{cases} (1 + |x|)^{-2\sigma - \tau} & \text{if } \tau < n, \\ (1 + |x|)^{-2\sigma - \tau} \log(2 + |x|) & \text{if } \tau = n, \\ (1 + |x|)^{-2\sigma - n} & \text{if } \tau > n. \end{cases}$$

Based on Lemma 2.3, a bootstrap argument allows us to give the following growth estimate of solutions to (1.1) near infinity.

Proposition 2.4. *Suppose that $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Let $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$ be a nonnegative distributional solution of*

$$(-\Delta)^\sigma u = |x|^{-\alpha} u^p \quad \text{in } \mathbb{R}^n \quad (2.10)$$

with $|\cdot|^{-\alpha} u^p \in L^1_{loc}(\mathbb{R}^n)$ for some $p > 1$.

(1) *If $1 < p < \frac{n-\alpha}{n-2\sigma}$, then*

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^\gamma} dx < +\infty \quad \text{for every } \gamma > 0. \quad (2.11)$$

(2) *If $p \geq \frac{n-\alpha}{n-2\sigma}$, then*

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^\gamma} dx < +\infty \quad \text{for every } \gamma > n - 2\sigma + \frac{\alpha - 2\sigma}{p - 1}. \quad (2.12)$$

Remark 2.5. *When $\alpha = 0$, the growth estimate in (2.12) has been proved by Ao-González-Hyder-Wei [2] for $\gamma = n - 2\sigma$. Here we continue to use the bootstrap argument to obtain the sharp estimate (2.12), which is essential in showing the integral characterization of solutions. We also remark that such sharp growth estimate was obtained by Du-Yang [29] when $\alpha = 0$, $p = \frac{n+2\sigma}{n-2\sigma}$ and σ is an integer by using a more direct argument (see also Proposition 2.6 below) that applies only to equations of integer order if $\sigma > 1$.*

Proof. For any $\tau > 0$ we take $\psi \in C^\infty(\mathbb{R}^n)$ satisfying $\psi(x) = \frac{1}{|x|^{n+\tau}}$ on B_1^c . Let ξ_ε be defined as in Lemma 2.3 and let $\psi_\varepsilon := \psi \xi_\varepsilon$. Then, using Lemma 2.3, dominated convergence theorem and monotone convergence theorem we obtain

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^{n+\tau}} dx = \int_{\mathbb{R}^n} u(-\Delta)^\sigma \psi dx < \infty. \quad (2.13)$$

For any $q > n + \frac{\alpha}{p}$ we write $q = q_1 + q_2$ with $q_1 > \frac{n}{p}$ and $q_2 > \frac{n}{p'} + \frac{\alpha}{p}$, where p' is the conjugate exponent of p . By Hölder inequality and (2.13) we have

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{u(x)}{1 + |x|^q} dx &\leq C_1 + \int_{B_1^c} \frac{|x|^{-\alpha/p} u(x)}{|x|^{q_1}} \cdot \frac{|x|^{\alpha/p}}{|x|^{q_2}} dx \\ &\leq C_1 + \left(\int_{B_1^c} \frac{|x|^{-\alpha} u(x)^p}{|x|^{q_1 p}} dx \right)^{1/p} \left(\int_{B_1^c} \frac{|x|^{\alpha p'/p}}{|x|^{q_2 p'}} dx \right)^{1/p'} \\ &< \infty \quad \text{for every } q > n + \frac{\alpha}{p}. \end{aligned}$$

Thus, the integrability of u lifts from $u \in \mathcal{L}_\sigma(\mathbb{R}^n)$ to $u \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{\alpha}{2p}$. Next we consider two cases for $p > 1$ separately.

Case 1: $p \leq \frac{-\alpha}{n-2\sigma}$. In this case we have $n + \frac{\alpha}{p} \leq 2\sigma$. By Lemma 2.3, for any $\tau > 0$ we can take $\psi \in C^\infty(\mathbb{R}^n)$ with $\psi(x) = \frac{1}{|x|^\tau}$ on B_1^c as a test function to (2.10), and consequently we have

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^\tau} dx < \infty \quad \text{for every } \tau > 0.$$

Case 2: $p > \frac{-\alpha}{n-2\sigma}$. In this case we have $0 < n + \frac{\alpha}{p} - 2\sigma < n$. By Lemma 2.3, we use $\psi \in C^\infty(\mathbb{R}^n)$ with $\psi(x) = \frac{1}{|x|^{n-2\sigma+\alpha/p+\tau}}$ on B_1^c for small $\tau > 0$ as a test function to (2.10), and thus we get

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^q} dx \leq C \int_{\mathbb{R}^n} \frac{u(x)}{1 + |x|^{q+2\sigma}} dx < \infty \quad \text{for every } q > n + \frac{\alpha}{p} - 2\sigma.$$

Again by Hölder inequality

$$\int_{\mathbb{R}^n} \frac{u(x)}{1 + |x|^q} dx < \infty \quad \text{for every } q > n + \frac{\alpha}{p} + \frac{\alpha}{p^2} - \frac{2\sigma}{p}.$$

Obviously, this further lifts the integrability of u . As discussed before we are still divided into two cases.

Case 1': $p \leq \frac{-(\alpha-2\sigma)+\sqrt{(\alpha-2\sigma)^2-4\alpha(n-2\sigma)}}{2(n-2\sigma)}$. In this case we know $n + \frac{\alpha}{p} + \frac{\alpha}{p^2} - \frac{2\sigma}{p} \leq 2\sigma$. Using Lemma 2.3 and taking $\psi \in C^\infty(\mathbb{R}^n)$ with $\psi(x) = \frac{1}{|x|^\tau}$ on B_1^c for $\tau > 0$ as a test function yields

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^\tau} dx < \infty \quad \text{for every } \tau > 0.$$

Case 2': $p > \frac{-(\alpha-2\sigma)+\sqrt{(\alpha-2\sigma)^2-4\alpha(n-2\sigma)}}{2(n-2\sigma)}$. In this case we have $0 < n + \frac{\alpha}{p} + \frac{\alpha}{p^2} - \frac{2\sigma}{p} - 2\sigma < n$. Using Lemma 2.3 and taking $\psi \in C^\infty(\mathbb{R}^n)$ with $\psi(x) = \frac{1}{|x|^q}$ on B_1^c for $q > n + \frac{\alpha}{p} + \frac{\alpha}{p^2} - \frac{2\sigma}{p} - 2\sigma$ as a test function yields

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{1 + |x|^q} dx < \infty \quad \text{for every } q > n + \frac{\alpha}{p} + \frac{\alpha}{p^2} - \frac{2\sigma}{p} - 2\sigma.$$

Again by Hölder inequality

$$\int_{\mathbb{R}^n} \frac{u(x)}{1 + |x|^q} dx < \infty \quad \text{for every } q > n + \left(\frac{\alpha}{p} + \frac{\alpha}{p^2} + \frac{\alpha}{p^3} \right) - \left(\frac{2\sigma}{p} + \frac{2\sigma}{p^2} \right).$$

This lifts the integrability of u again.

Continuing this process, after finite many steps we will get the desired conclusion for any fix $p > 1$ and fix γ satisfying the assumption. The proof is completed. \square

When $\sigma \in (0, n/2)$ is an integer, the growth estimates of solutions to (2.10) are much easier to obtain by the standard rescaling method. More precisely, we have

Proposition 2.6. *Suppose that $\sigma \in (0, n/2)$ is an integer and $\alpha \in (-\infty, 2\sigma)$. Let $u \in L_{loc}^1(\mathbb{R}^n)$ be a nonnegative distributional solution of (2.10) with $|\cdot|^{-\alpha}u^p \in L_{loc}^1(\mathbb{R}^n)$ for some $p > 1$. Then*

$$\int_{\mathbb{R}^n} \frac{u(x)}{(1+|x|)^\gamma} dx < \infty \quad \text{for every } \gamma > n + \frac{\alpha - 2\sigma}{p-1} \quad (2.14)$$

and

$$\int_{\mathbb{R}^n} \frac{|x|^{-\alpha}u(x)^p}{(1+|x|)^\gamma} dx < \infty \quad \text{for every } \gamma > n - 2\sigma + \frac{\alpha - 2\sigma}{p-1}. \quad (2.15)$$

Proof. Let $\varphi \in C_c^\infty(\mathbb{R}^n)$ be a nonnegative function satisfying $\varphi = 1$ on B_1 and $\varphi = 0$ on B_2^c . For $R > 0$, set $\varphi_R(x) := \varphi(\frac{x}{R})$. Fix an integer $q > \frac{2\sigma p}{p-1}$ and choose φ_R^q as a test function in (2.10). Note that here $\sigma \in (0, n/2)$ is an integer, we have $|(-\Delta)^\sigma \varphi_R^q(x)| \leq CR^{-2\sigma} \varphi_R(x)^{q-2\sigma}$ for some constant $C > 0$. By Hölder inequality we obtain

$$\begin{aligned} \int_{\mathbb{R}^n} |x|^{-\alpha} u^p \varphi_R^q dx &= \int_{\mathbb{R}^n} u (-\Delta)^\sigma \varphi_R^q dx \leq \frac{C}{R^{2\sigma}} \int_{B_{2R} \setminus B_R} u \varphi_R^{q-2\sigma} dx \\ &= \frac{C}{R^{2\sigma}} \int_{B_{2R} \setminus B_R} |x|^{-\alpha/p} u \varphi_R^{q/p} \cdot |x|^{\alpha/p} \varphi_R^{q-q/p-2\sigma} dx \\ &\leq \frac{C}{R^{2\sigma}} \left(\int_{B_{2R} \setminus B_R} |x|^{-\alpha} u^p \varphi_R^q dx \right)^{1/p} \left(\int_{B_{2R} \setminus B_R} |x|^{\alpha p'/p} dx \right)^{1/p'} \\ &\leq CR^{\frac{\alpha}{p} + \frac{n}{p'} - 2\sigma} \left(\int_{B_{2R} \setminus B_R} |x|^{-\alpha} u^p \varphi_R^q dx \right)^{1/p}. \end{aligned}$$

This gives that

$$\int_{B_R} |x|^{-\alpha} u^p dx \leq CR^{n + \frac{\alpha p'}{p} - 2\sigma p'} = CR^{n-2\sigma + \frac{\alpha-2\sigma}{p-1}} \quad \text{for any } R > 0. \quad (2.16)$$

Here we also point out that the above two estimates imply $u \equiv 0$ if $1 < p \leq \frac{n-\alpha}{n-2\sigma}$. Using Hölder inequality again, we obtain, for any $\gamma \in \mathbb{R}$ and $R \geq 1$, that

$$\begin{aligned} \int_{B_{2R} \setminus B_R} \frac{u(x)}{(1+|x|)^\gamma} dx &\leq \left(\int_{B_{2R} \setminus B_R} |x|^{-\alpha} u^p dx \right)^{1/p} \left(\int_{B_{2R} \setminus B_R} \frac{|x|^{\alpha p'/p}}{(1+|x|)^{\gamma p'}} dx \right)^{1/p'} \\ &\leq CR^{n + \frac{\alpha-2\sigma}{p-1} - \gamma}. \end{aligned}$$

Hence, when $\gamma > n + \frac{\alpha-2\sigma}{p-1}$, a dyadic sum argument leads to that (2.14) holds.

On the other hand, estimate (2.16) also implies that

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{|x|^{-\alpha} u(x)^p}{(1+|x|)^\gamma} dx &= \int_{B_1} \frac{|x|^{-\alpha} u(x)^p}{(1+|x|)^\gamma} dx + \sum_{i=0}^{\infty} \int_{B_{2^{i+1}} \setminus B_{2^i}} \frac{|x|^{-\alpha} u(x)^p}{(1+|x|)^\gamma} dx \\ &\leq C_1 + C \sum_{i=0}^{\infty} (2^i)^{n-2\sigma + \frac{\alpha-2\sigma}{p-1} - \gamma} < \infty \end{aligned}$$

if $\gamma > n - 2\sigma + \frac{\alpha-2\sigma}{p-1}$. This proves (2.15). \square

Suppose that $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Let $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n \setminus \{0\})$ be a nonnegative distributional solution of (1.1) with $p \geq \frac{n-\alpha}{n-2\sigma}$. Then, by Propositions 2.2 and 2.4, the following function

$$v(x) := C_{n,\sigma} \int_{\mathbb{R}^n} \frac{|y|^{-\alpha} u(y)^p}{|x-y|^{n-2\sigma}} dy \quad (2.17)$$

is well-defined for every $x \in \mathbb{R}^n \setminus \{0\}$ and continuous on $\mathbb{R}^n \setminus \{0\}$, where $C_{n,\sigma}$ is a positive constant such that $C_{n,\sigma} |\cdot|^{2\sigma-n}$ is the fundamental solution of the fractional Laplacian. For any $R > 0$, we can write v as $v = v_{1,R} + v_{2,R}$ with

$$v_{1,R}(x) = \int_{B_{2R}} \frac{|y|^{-\alpha} u(y)^p}{|x-y|^{n-2\sigma}} dy \quad \text{and} \quad v_{2,R}(x) = \int_{B_{2R}^c} \frac{|y|^{-\alpha} u(y)^p}{|x-y|^{n-2\sigma}} dy.$$

From Proposition 2.2 we know that $|\cdot|^{-\alpha} u^p \in L^1(B_{2R})$ and hence $v_{1,R} \in L^1(B_R)$. By Proposition 2.4 we easily get $v_{2,R} \in L^\infty(B_R)$. Thus $v \in L^1_{loc}(\mathbb{R}^n)$. Moreover, we also have the following growth estimate for v near infinity.

Proposition 2.7. *Suppose that $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$.*

- (1) *Let $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n \setminus \{0\})$ be a nonnegative distributional solution of (1.1) with $p \geq \frac{n-\alpha}{n-2\sigma}$, and let v be defined by (2.17). Then we have $v \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{\alpha-2\sigma}{2(p-1)}$.*
- (2) *Let $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ be a nonnegative distributional solution of (1.1) in \mathbb{R}^n with $1 < p < \frac{n-\alpha}{n-2\sigma}$, and let v be defined by (2.17). Then we have $v \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{2\sigma-n}{2}$.*

Proof. By Fubini's theorem, for any $q > 2\sigma$ we have

$$\int_{\mathbb{R}^n} \frac{v(x)}{(1+|x|)^q} dx = c_{n,\sigma} \int_{\mathbb{R}^n} |y|^{-\alpha} u(y)^p \left(\int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-2\sigma}} \frac{1}{(1+|x|)^q} dx \right) dy. \quad (2.18)$$

If $|y| \leq 1$, then

$$\begin{aligned} \int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-2\sigma}} \frac{1}{(1+|x|)^q} dx &\leq C \left(\int_{B_3} \frac{1}{|x|^{n-2\sigma}} dx + \int_{B_2^c} \frac{1}{|x|^{n-2\sigma+q}} dx \right) \\ &\leq C < \infty. \end{aligned}$$

If $|y| > 1$, then

$$\int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-2\sigma}} \frac{1}{(1+|x|)^q} dx =: \sum_{i=1}^3 I_i,$$

where

$$I_1 = \int_{\{|x| \leq \frac{|y|}{2}\}} \frac{1}{|x-y|^{n-2\sigma}} \frac{1}{(1+|x|)^q} dx \leq C \begin{cases} |y|^{2\sigma-q}, & \text{if } q < n, \\ \log(1+|y|)|y|^{2\sigma-n}, & \text{if } q = n, \\ |y|^{2\sigma-n}, & \text{if } q > n, \end{cases}$$

$$\begin{aligned}
I_2 &= \int_{\{\frac{|y|}{2} < |x| < 2|y|\}} \frac{1}{|x-y|^{n-2\sigma}} \frac{1}{(1+|x|)^q} dx \leq \frac{C}{|y|^q} \int_{\{\frac{|y|}{2} < |x| < 2|y|\}} \frac{1}{|x-y|^{n-2\sigma}} dx \\
&\leq C|y|^{2\sigma-q}
\end{aligned}$$

and

$$I_3 = \int_{\{|x| \geq 2|y|\}} \frac{1}{|x-y|^{n-2\sigma}} \frac{1}{(1+|x|)^q} dx \leq C \int_{\{|x| \geq 2|y|\}} \frac{1}{|x|^{n-2\sigma+q}} dx \leq C|y|^{2\sigma-q}.$$

Let $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n \setminus \{0\})$ be a nonnegative distributional solution of (1.1) with $p \geq \frac{n-\alpha}{n-2\sigma}$. Then for $q > n + \frac{\alpha-2\sigma}{p-1}$ (note that $n + \frac{\alpha-2\sigma}{p-1} \geq 2\sigma$ due to $p \geq \frac{n-\alpha}{n-2\sigma}$), by (2.18) and (2.12) we get

$$\int_{\mathbb{R}^n} \frac{v(x)}{(1+|x|)^q} dx \leq C \int_{B_1} |y|^{-\alpha} u(y)^p dy + C \int_{B_1^c} \frac{\log(1+|y|) |y|^{-\alpha} u(y)^p}{|y|^{q-2\sigma}} dy < \infty.$$

That is, $v \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{\alpha-2\sigma}{2(p-1)}$.

Similarly, if $u \in \mathcal{L}_\sigma(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ is a nonnegative distributional solution of (1.1) in \mathbb{R}^n with $1 < p < \frac{n-\alpha}{n-2\sigma}$, then by using (2.11) we get that $v \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{2\sigma-n}{2}$. \square

Based on Propositions 2.2, 2.4 and 2.7, we now show the integral representation of solutions to the higher order fractional equation (1.1) stated in Theorem 1.1. Our proof is inspired by that of [2, Theorem 1.8], where the superharmonicity property for fractional Laplacian equations in \mathbb{R}^n was obtained. Similar idea has also been used in [29] for the case when $\alpha = 0$, $p = \frac{n+2\sigma}{n-2\sigma}$ and $\sigma \in (0, n/2)$ is an integer.

Proof of Theorem 1.1. Define v as in (2.17). Then, by Proposition 2.7 we know that $v \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{\alpha-2\sigma}{2(p-1)}$ if $p \geq \frac{n-\alpha}{n-2\sigma}$ and $v \in \mathcal{L}_s(\mathbb{R}^n)$ for any $s > \frac{2\sigma-n}{2}$ if $1 < p < \frac{n-\alpha}{n-2\sigma}$. In particular, $v \in \mathcal{L}_0(\mathbb{R}^n)$ for any case. Moreover, for any $\varphi \in C_c^\infty(\mathbb{R}^n)$ we have

$$\begin{aligned}
\int_{\mathbb{R}^n} v(x) (-\Delta)^\sigma \varphi(x) dx &= \int_{\mathbb{R}^n} \left(C_{n,\sigma} \int_{\mathbb{R}^n} \frac{|y|^{-\alpha} u(y)^p}{|x-y|^{n-2\sigma}} dy \right) (-\Delta)^\sigma \varphi(x) dx \\
&= \int_{\mathbb{R}^n} |y|^{-\alpha} u(y)^p \left(C_{n,\sigma} \int_{\mathbb{R}^n} \frac{(-\Delta)^\sigma \varphi(x)}{|x-y|^{n-2\sigma}} dx \right) dy \\
&= \int_{\mathbb{R}^n} |y|^{-\alpha} u(y)^p \varphi(y) dy,
\end{aligned}$$

where the Fubini's theorem was used in the second equality. Thus v is a nonnegative distributional solution to

$$(-\Delta)^\sigma v = |x|^{-\alpha} u^p \quad \text{in } \mathbb{R}^n. \tag{2.19}$$

Let $w := u - v$. Then, by Proposition 2.2 we have that $(-\Delta)^\sigma w = 0$ on \mathbb{R}^n in the distributional sense. On the other hand, from Propositions 2.4 and 2.7 we obtain that both u and v are in the space $\mathcal{L}_0(\mathbb{R}^n)$ and hence $w \in \mathcal{L}_0(\mathbb{R}^n)$. It follows from a Liouville type theorem (see, e.g., [2]) that $w \equiv 0$ on \mathbb{R}^n . Thus, we have

$$u(x) = C_{n,\sigma} \int_{\mathbb{R}^n} \frac{|y|^{-\alpha} u(y)^p}{|x-y|^{n-2\sigma}} dy \quad \text{for } x \in \mathbb{R}^n \setminus \{0\}.$$

When $u \in C(\mathbb{R}^n)$, the above holds for all $x \in \mathbb{R}^n$. Theorem 1.1 is established. \square

3 Liouville-type theorems

In this section, we first prove a Liouville-type theorem for nonnegative solutions of the following integral equation

$$u(x) = \int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^\alpha |x-y|^{n-2\sigma}} dy, \quad x \in \mathbb{R}^n. \quad (3.1)$$

Theorem 3.1. *Let $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Suppose that $u \in C(\mathbb{R}^n)$ is a nonnegative solution of (3.1) with $0 < p < \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Then $u \equiv 0$ in \mathbb{R}^n .*

Remark 3.2. *Such Liouville-type theorem for the integral equation (3.1) has been proved by Li [44] for $\alpha = 0$, by Dai-Qin [25] for $\sigma = m \in (1, n/2)$ being an integer and $-\infty < \alpha < 2m$, and by Cao-Dai-Qin [12] for non-integral $\sigma \in (1, n/2)$ and $-\infty < \alpha \leq 0$. Here we will prove Theorem 3.1 by using the method of moving spheres in Li, Zhang and Zhu [44–46] and a “Bootstrap” iteration argument in Dai-Qin [25].*

For any $\lambda > 0$ and nonnegative function u , we define the Kelvin transform of u with respect to the ball $B_\lambda(0)$ by

$$u_\lambda(\xi) = \left(\frac{\lambda}{|\xi|} \right)^{n-2\sigma} u(\xi^\lambda) \quad \text{for } \xi \neq 0, \quad (3.2)$$

where $\xi^\lambda := \frac{\lambda^2 \xi}{|\xi|^2}$. It is easy to check that $(\xi^\lambda)^\lambda = \xi$ and $(u_\lambda)_\lambda \equiv u$. By making a change of variables $y = z^\lambda = \frac{\lambda^2 z}{|z|^2}$, we have

$$\begin{aligned} \int_{|y| \geq \lambda} \frac{|y|^{-\alpha} u(y)^p}{|\xi^\lambda - y|^{n-2\sigma}} dy &= \int_{|z| \leq \lambda} \frac{|z^\lambda|^{-\alpha} u(z^\lambda)^p}{|\xi^\lambda - z^\lambda|^{n-2\sigma}} \left(\frac{\lambda}{|z|} \right)^{2n} dz \\ &= \left(\frac{\lambda}{|\xi|} \right)^{-(n-2\sigma)} \int_{|z| \leq \lambda} \frac{|z|^{-\alpha} u_\lambda(z)^p}{|\xi - z|^{n-2\sigma}} \left(\frac{\lambda}{|z|} \right)^{n+2\sigma-2\alpha-p(n-2\sigma)} dz, \end{aligned}$$

where we have used the fact that $\frac{|z|}{\lambda} \frac{|\xi|}{\lambda} |\xi^\lambda - z^\lambda| = |\xi - z|$. Hence, we obtain

$$\left(\frac{\lambda}{|\xi|} \right)^{n-2\sigma} \int_{|y| \geq \lambda} \frac{|y|^{-\alpha} u(y)^p}{|\xi^\lambda - y|^{n-2\sigma}} dy = \int_{|z| \leq \lambda} \frac{|z|^{-\alpha} u_\lambda(z)^p}{|\xi - z|^{n-2\sigma}} \left(\frac{\lambda}{|z|} \right)^{n+2\sigma-2\alpha-p(n-2\sigma)} dz. \quad (3.3)$$

Similarly, we also have

$$\left(\frac{\lambda}{|\xi|} \right)^{n-2\sigma} \int_{|y| \leq \lambda} \frac{|y|^{-\alpha} u(y)^p}{|\xi^\lambda - y|^{n-2\sigma}} dy = \int_{|z| \geq \lambda} \frac{|z|^{-\alpha} u_\lambda(z)^p}{|\xi - z|^{n-2\sigma}} \left(\frac{\lambda}{|z|} \right)^{n+2\sigma-2\alpha-p(n-2\sigma)} dz. \quad (3.4)$$

Thus, if $u \in C(\mathbb{R}^n)$ is a nonnegative solution of (3.1), then by (3.3) and (3.4) we get

$$u_\lambda(\xi) = \int_{\mathbb{R}^n} \frac{|z|^{-\alpha} u_\lambda(z)^p}{|\xi - z|^{n-2\sigma}} \left(\frac{\lambda}{|z|} \right)^{n+2\sigma-2\alpha-p(n-2\sigma)} dz \quad (3.5)$$

and

$$u(\xi) - u_\lambda(\xi) = \int_{|z| \geq \lambda} K(0, \lambda; \xi, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\lambda}{|z|} \right)^{n+2\sigma-2\alpha-p(n-2\sigma)} \frac{u_\lambda(z)^p}{|z|^\alpha} \right] dz, \quad (3.6)$$

where

$$K(0, \lambda; \xi, z) = \frac{1}{|\xi - z|^{n-2\sigma}} - \left(\frac{\lambda}{|\xi|}\right)^{n-2\sigma} \frac{1}{|\xi\lambda - z|^{n-2\sigma}}.$$

It is elementary to check that

$$K(0, \lambda; \xi, z) > 0 \quad \text{for all } |\xi|, |z| > \lambda > 0.$$

To apply the moving sphere method, we first prove the following result which provides us a starting point.

Lemma 3.3. *Let $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Suppose that $u \in C(\mathbb{R}^n)$ is a positive solution of (3.1) with $0 < p < \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Then there exists a small $\lambda_0 > 0$ such that for any $0 < \lambda < \lambda_0$ we have*

$$u_\lambda(y) \leq u(y) \quad \forall |y| \geq \lambda. \quad (3.7)$$

Proof. By Fatou lemma, we have

$$\liminf_{|x| \rightarrow \infty} |x|^{n-2\sigma} u(x) = \liminf_{|x| \rightarrow \infty} \int_{\mathbb{R}^n} \frac{|x|^{n-2\sigma} u(y)^p}{|y|^\alpha |x-y|^{n-2\sigma}} dy \geq \int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^\alpha} dy > 0.$$

This, together with the positivity and continuity of u , yields that for each $r_0 > 0$

$$u(y) \geq \frac{C(r_0)}{|y|^{n-2\sigma}} \quad \forall |y| \geq r_0 \quad (3.8)$$

with some constant $C(r_0) > 0$. Hence, for every $r_0 > 0$, there exists $\lambda_1 = \lambda_1(r_0) \in (0, r_0)$ such that for any $0 < \lambda < \lambda_1$,

$$u_\lambda(y) = \left(\frac{\lambda}{|y|}\right)^{n-2\sigma} u\left(\frac{\lambda^2 y}{|y|^2}\right) \leq \left(\frac{\lambda}{|y|}\right)^{n-2\sigma} \sup_{B_{r_0}} u \leq u(y) \quad \forall |y| \geq r_0. \quad (3.9)$$

Next we will show that $u_\lambda(y) \leq u(y)$ on $B_{r_0} \setminus B_\lambda$ if r_0 is small and $0 < \lambda < \lambda_1(r_0)$. Define $w^\lambda(y) = u(y) - u_\lambda(y)$ for $|y| \geq \lambda$ and denote

$$B_\lambda^- := \{y \in \mathbb{R}^n \setminus B_\lambda : w^\lambda(y) < 0\}.$$

Then by (3.9) we know that $B_\lambda^- \subset B_{r_0} \setminus B_\lambda$ for any fixed $r_0 > 0$ and for all $0 < \lambda < \lambda_1(r_0)$. Using (3.6) we have that for $0 < \lambda < \lambda_1(r_0)$ and $y \in B_\lambda^-$,

$$\begin{aligned} 0 > w^\lambda(y) &= \int_{|z| \geq \lambda} K(0, \lambda; y, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\lambda}{|z|}\right)^{n+2\sigma-2\alpha-p(n-2\sigma)} \frac{u_\lambda(z)^p}{|z|^\alpha} \right] dz \\ &\geq \int_{B_\lambda^-} K(0, \lambda; y, z) |z|^{-\alpha} (u(z)^p - u_\lambda(z)^p) dz \\ &\geq p \int_{B_\lambda^-} \frac{1}{|y-z|^{n-2\sigma}} |z|^{-\alpha} \max\{u(z)^{p-1}, u_\lambda(z)^{p-1}\} w^\lambda(z) dz. \end{aligned}$$

Hence, by Hardy-Littlewood-Sobolev inequality (see, e.g., [48]), we obtain that for any $\frac{n}{n-2\sigma} < q < \infty$,

$$\begin{aligned} \|w^\lambda\|_{L^q(B_\lambda^-)} &\leq C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} w^\lambda \right\|_{L^{\frac{nq}{n+2\sigma q}}(B_\lambda^-)} \\ &\leq C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} \right\|_{L^{\frac{n}{2\sigma}}(B_\lambda^-)} \|w^\lambda\|_{L^q(B_\lambda^-)}. \end{aligned} \quad (3.10)$$

Since $\|\max\{u^{p-1}, u_\lambda^{p-1}\}\|_{L^\infty(B_\lambda^-)} \leq \|u^{p-1}\|_{L^\infty(B_{r_0})}$ and $\alpha < 2\sigma$, we can take $r_0 = \varepsilon_0$ for some $\varepsilon_0 > 0$ small enough such that

$$C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} \right\|_{L^{\frac{n}{2\sigma}}(B_\lambda^-)} \leq C \left\| |\cdot|^{-\alpha} \right\|_{L^{\frac{n}{2\sigma}}(B_{\varepsilon_0})} \|u^{p-1}\|_{L^\infty(B_{\varepsilon_0})} \leq \frac{1}{2}$$

for all $0 < \lambda < \lambda_0 := \lambda_1(\varepsilon_0)$, and thus, (3.10) implies that

$$\|w^\lambda\|_{L^q(B_\lambda^-)} = 0.$$

Therefore, $B_\lambda^- = \emptyset$ for any $0 < \lambda < \lambda_0$. That is, (3.7) holds for any $0 < \lambda < \lambda_0$. \square

We define

$$\bar{\lambda} := \sup\{\mu > 0 \mid u_\lambda(y) \leq u(y) \forall |y| \geq \lambda, \forall 0 < \lambda < \mu\}.$$

By Lemma 3.3, $\bar{\lambda}$ is well-defined and $\bar{\lambda} > 0$. Moreover, we have the following result.

Lemma 3.4. *Let $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Suppose that $u \in C(\mathbb{R}^n)$ is a positive solution of (3.1) with $0 < p < \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Then $\bar{\lambda} = \infty$.*

Proof. Suppose by contradiction that $\bar{\lambda} < \infty$. The definition of $\bar{\lambda}$ implies that

$$u_{\bar{\lambda}}(y) \leq u(y) \quad \forall |y| \geq \bar{\lambda}. \quad (3.11)$$

Since $n + 2\sigma - 2\alpha - p(n - 2\sigma) > 0$, we have $\left(\frac{\bar{\lambda}}{|z|}\right)^{n+2\sigma-2\alpha-p(n-2\sigma)} < 1$ for $|z| > \bar{\lambda}$. Using (3.6) we get, for $|y| > \bar{\lambda}$,

$$\begin{aligned} u(y) - u_{\bar{\lambda}}(y) &= \int_{|z| \geq \bar{\lambda}} K(0, \bar{\lambda}; y, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\bar{\lambda}}{|z|}\right)^{n+2\sigma-2\alpha-p(n-2\sigma)} \frac{u_{\bar{\lambda}}(z)^p}{|z|^\alpha} \right] dz \\ &\geq \int_{|z| \geq \bar{\lambda}} K(0, \bar{\lambda}; y, z) \left[1 - \left(\frac{\bar{\lambda}}{|z|}\right)^{n+2\sigma-2\alpha-p(n-2\sigma)} \right] \frac{u_{\bar{\lambda}}(z)^p}{|z|^\alpha} dz > 0. \end{aligned}$$

This, together with Fatou lemma, yields

$$\begin{aligned} \liminf_{|y| \rightarrow \infty} |y|^{n-2\sigma} (u - u_{\bar{\lambda}})(y) &\geq \liminf_{|y| \rightarrow \infty} \int_{|z| \geq \bar{\lambda}} |y|^{n-2\sigma} K(0, \bar{\lambda}; y, z) \frac{u(z)^p - u_{\bar{\lambda}}(z)^p}{|z|^\alpha} dz \\ &\geq \int_{|z| \geq \bar{\lambda}} \left[1 - \left(\frac{\bar{\lambda}}{|z|}\right)^{n-2\sigma} \right] \frac{u(z)^p - u_{\bar{\lambda}}(z)^p}{|z|^\alpha} dz > 0. \end{aligned}$$

Consequently, for $\delta > 0$ small which will be fixed later, there exists $c_1 = c_1(\delta) > 0$ such that

$$u(y) - u_{\bar{\lambda}}(y) \geq \frac{c_1}{|y|^{n-2\sigma}} \quad \forall |y| \geq \bar{\lambda} + \delta.$$

By the above and explicit formula of u_λ , there exists a small $\varepsilon \in (0, \delta)$ such that for all $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$,

$$u(y) - u_\lambda(y) \geq \frac{c_1}{|y|^{n-2\sigma}} + (u_{\bar{\lambda}}(y) - u_\lambda(y)) \geq \frac{c_1}{2|y|^{n-2\sigma}} \quad \forall |y| \geq \bar{\lambda} + \delta. \quad (3.12)$$

Now, we focus on the narrow region $B_{\bar{\lambda}+\delta} \setminus B_\lambda$. As in the proof of Lemma 3.3, we define $w^\lambda(y) = u(y) - u_\lambda(y)$ for $|y| \geq \lambda$ and denote

$$B_\lambda^- := \{y \in \mathbb{R}^n \setminus B_\lambda : w^\lambda(y) < 0\}.$$

By (3.12) we have $B_\lambda^- \subset B_{\bar{\lambda}+\delta} \setminus B_\lambda$ for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$. Thus, using (3.6) we have, for $y \in B_\lambda^-$ and for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$,

$$\begin{aligned} 0 > w^\lambda(y) &\geq \int_{|z| \geq \lambda} K(0, \lambda; y, z) |z|^{-\alpha} (u(z)^p - u_\lambda(z)^p) dz \\ &\geq p \int_{B_\lambda^-} \frac{1}{|y-z|^{n-2\sigma}} |z|^{-\alpha} \max\{u(z)^{p-1}, u_\lambda(z)^{p-1}\} w^\lambda(z) dz. \end{aligned}$$

By Hardy-Littlewood-Sobolev inequality (see, e.g., [48]), we obtain that for any $\frac{n}{n-2\sigma} < q < \infty$,

$$\begin{aligned} \|w^\lambda\|_{L^q(B_\lambda^-)} &\leq C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} w^\lambda \right\|_{L^{\frac{nq}{n+2\sigma q}}(B_\lambda^-)} \\ &\leq C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} \right\|_{L^{\frac{n}{2\sigma}}(B_\lambda^-)} \|w^\lambda\|_{L^q(B_\lambda^-)}. \end{aligned} \quad (3.13)$$

Note that we can choose $\delta > 0$ small enough such that

$$C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} \right\|_{L^{\frac{n}{2\sigma}}(B_\lambda^-)} \leq C \left\| |\cdot|^{-\alpha} \max\{u^{p-1}, u_\lambda^{p-1}\} \right\|_{L^{\frac{n}{2\sigma}}(B_{\bar{\lambda}+\delta} \setminus B_\lambda)} \leq \frac{1}{2}$$

for all $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$, and thus, (3.13) implies that

$$\|w^\lambda\|_{L^q(B_\lambda^-)} = 0.$$

Therefore, $B_\lambda^- = \emptyset$ for all $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$. That is, there exists $\varepsilon > 0$ such that for all $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$,

$$u_\lambda(y) \leq u(y) \quad \forall |y| \geq \lambda,$$

which contradicts the definition of $\bar{\lambda}$. \square

Proof of Theorem 3.1. Suppose by contrary that $u \in C(\mathbb{R}^n)$ is a nontrivial nonnegative solution of (3.1), then there exists $x_0 \in \mathbb{R}^n$ such that $u(x_0) > 0$. It follows from the equation (3.1) that $u(x) > 0$ for all $x \in \mathbb{R}^n$. Thus, $u \in C(\mathbb{R}^n)$ is a positive solution of (3.1). By Lemma 3.4 we have

$$u(x) \geq \left(\frac{\lambda}{|x|}\right)^{n-2\sigma} u\left(\frac{\lambda^2 x}{|x|^2}\right) \quad \forall |x| \geq \lambda, \quad \forall 0 < \lambda < \infty. \quad (3.14)$$

For any $|x| \geq 1$, taking $\lambda = \sqrt{|x|}$ in (3.14) yields that

$$u(x) \geq |x|^{-\frac{n-2\sigma}{2}} u\left(\frac{x}{|x|}\right). \quad (3.15)$$

Hence, we have the following lower bound:

$$u(x) \geq \left(\min_{\partial B_1} u \right) |x|^{-\frac{n-2\sigma}{2}} =: C_0 |x|^{-\frac{n-2\sigma}{2}} \quad \forall |x| \geq 1. \quad (3.16)$$

Now, we use the ‘‘Bootstrap’’ iteration argument to improve the lower bound (3.16). Let $\mu_0 := \frac{n-2\sigma}{2}$. From (3.1) and (3.16) we have, for $|x| \geq 1$,

$$\begin{aligned} u(x) &\geq C \int_{|x| \leq |y| \leq 2|x|} \frac{1}{|x-y|^{n-2\sigma} |y|^{p\mu_0+\alpha}} dy \\ &\geq C |x|^{-(n-2\sigma)} \int_{|x| \leq |y| \leq 2|x|} \frac{1}{|y|^{p\mu_0+\alpha}} dy \\ &:= C_1 |x|^{-(p\mu_0+\alpha-2\sigma)} \end{aligned}$$

with some constant $C_1 > 0$. Let $\mu_1 := p\mu_0 + \alpha - 2\sigma$. Since $0 < p < \frac{n+2\sigma-2\alpha}{n-2\sigma}$, we have

$$\mu_1 = p\mu_0 + \alpha - 2\sigma < \mu_0. \quad (3.17)$$

This means that we have obtained a better lower bound after one iteration,

$$u(x) \geq C_1 |x|^{-\mu_1} \quad \forall |x| \geq 1. \quad (3.18)$$

Continuing this process, after k iterations we would get the following lower bound

$$u(x) \geq C_k |x|^{-\mu_k} \quad \forall |x| \geq 1, \quad (3.19)$$

where $C_k > 0$ is a constant and the sequence $\{\mu_k\}_{k \geq 1}$ satisfies

$$\mu_k = p\mu_{k-1} + \alpha - 2\sigma, \quad k = 1, 2, \dots. \quad (3.20)$$

It is easy to check that the sequence $\{\mu_k\}_{k \geq 1}$ is monotonically decreasing with respect to k . Moreover, as $k \rightarrow \infty$,

$$\mu_k \rightarrow \begin{cases} \frac{\alpha-2\sigma}{1-p} & \text{if } 0 < p < 1, \\ -\infty & \text{if } 1 \leq p < \frac{n+2\sigma-2\alpha}{n-2\sigma}. \end{cases}$$

Thus, by (3.19) we have the following lower estimates: for $|x| \geq 1$,

$$u(x) \geq \begin{cases} C_\kappa |x|^\kappa & \forall \kappa < \frac{2\sigma-\alpha}{1-p}, & \text{if } 0 < p < 1, \\ C_\kappa |x|^\kappa & \forall \kappa < +\infty, & \text{if } 1 \leq p < \frac{n+2\sigma-2\alpha}{n-2\sigma}. \end{cases}$$

It follows from the equation (3.1) and above estimates that

$$u(0) = \int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^{n+\alpha-2\sigma}} dy \geq \int_{|y| \geq 1} \frac{1}{|y|^{n+\alpha-2\sigma}} dy = +\infty,$$

which is a contradiction. Therefore, we must have $u \equiv 0$ in \mathbb{R}^n . The proof of Theorem 3.1 is completed. \square

Proof of Theorem 1.4. It follows from Theorems 1.1 and 3.1. \square

4 Radial symmetry of solutions

In this section, we first prove the following radial symmetry of nonnegative solutions to the integral equation

$$u(x) = \int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^\alpha |x-y|^{n-2\sigma}} dy, \quad x \in \mathbb{R}^n \setminus \{0\}. \quad (4.1)$$

Then we have

Theorem 4.1. *Let $0 < \sigma < n/2$ and $0 \leq \alpha < 2\sigma$. Suppose that $u \in C(\mathbb{R}^n \setminus \{0\})$ is a nonnegative solution of (4.1) with $p > 0$, and u has a non-removable singularity at the origin when $\alpha = 0$ and $p = \frac{n+2\sigma}{n-2\sigma}$.*

(1) *If $0 < p \leq \frac{n+2\sigma-2\alpha}{n-2\sigma}$, then u is radially symmetric and monotonically decreasing with respect to the origin.*

(2) *If $\frac{n+2\sigma-2\alpha}{n-2\sigma} < p \leq \frac{n+2\sigma-\alpha}{n-2\sigma}$, then u is radially symmetric with respect to the origin.*

Remark 4.2. *When $\alpha = 0$, Theorem 4.1 has been proved by Chen-Li-Ou [20] via the method of moving planes in an integral form. Here we will apply the method of moving spheres in Li, Zhang and Zhu [44–46] to show Theorem 4.1 for all $\alpha \in [0, 2\sigma)$.*

For $x \in \mathbb{R}^n$, $\lambda > 0$ and a nonnegative function u , we define the Kelvin transform of u with respect to the ball $B_\lambda(x)$ by

$$u_{x,\lambda}(\xi) = \left(\frac{\lambda}{|\xi-x|} \right)^{n-2\sigma} u(\xi^{x,\lambda}), \quad (4.2)$$

where

$$\xi^{x,\lambda} = x + \frac{\lambda^2(\xi-x)}{|\xi-x|^2} \quad \text{for } \xi \neq x. \quad (4.3)$$

One can check that $(\xi^{x,\lambda})^{x,\lambda} = \xi$ and $(u_{x,\lambda})_{x,\lambda} \equiv u$. If $x = 0$, we use the notation $u_\lambda = u_{0,\lambda}$ and $\xi^\lambda = \xi^{0,\lambda}$ as in Section 3. Making the following change of variables

$$y = z^{x,\lambda} = x + \frac{\lambda^2(z-x)}{|z-x|^2},$$

we have

$$\begin{aligned} \int_{|y-x| \geq \lambda} \frac{|y|^{-\alpha} u(y)^p}{|\xi^{x,\lambda} - y|^{n-2\sigma}} dy &= \int_{|z-x| \leq \lambda} \frac{|z^{x,\lambda}|^{-\alpha} u(z^{x,\lambda})^p}{|\xi^{x,\lambda} - z^{x,\lambda}|^{n-2\sigma}} \left(\frac{\lambda}{|z-x|} \right)^{2n} dz \\ &= \left(\frac{\lambda}{|\xi-x|} \right)^{-(n-2\sigma)} \int_{|z-x| \leq \lambda} \frac{|z^{x,\lambda}|^{-\alpha} u_{x,\lambda}(z)^p}{|\xi-z|^{n-2\sigma}} \left(\frac{\lambda}{|z-x|} \right)^{n+2\sigma-p(n-2\sigma)} dz, \end{aligned}$$

where we have used the fact

$$\frac{|z-x|}{\lambda} \frac{|\xi-x|}{\lambda} |\xi^{x,\lambda} - z^{x,\lambda}| = |\xi-z|.$$

Thus, we obtain

$$\left(\frac{\lambda}{|\xi-x|}\right)^{n-2\sigma} \int_{|y-x|\geq\lambda} \frac{|y|^{-\alpha}u(y)^p}{|\xi^{x,\lambda}-y|^{n-2\sigma}} dy = \int_{|z-x|\leq\lambda} \frac{|z^{x,\lambda}|^{-\alpha}u_{x,\lambda}(z)^p}{|\xi-z|^{n-2\sigma}} \left(\frac{\lambda}{|z-x|}\right)^{n+2\sigma-p(n-2\sigma)} dz. \quad (4.4)$$

Similarly, we also have

$$\left(\frac{\lambda}{|\xi-x|}\right)^{n-2\sigma} \int_{|y-x|\leq\lambda} \frac{|y|^{-\alpha}u(y)^p}{|\xi^{x,\lambda}-y|^{n-2\sigma}} dy = \int_{|z-x|\geq\lambda} \frac{|z^{x,\lambda}|^{-\alpha}u_{x,\lambda}(z)^p}{|\xi-z|^{n-2\sigma}} \left(\frac{\lambda}{|z-x|}\right)^{n+2\sigma-p(n-2\sigma)} dz. \quad (4.5)$$

Let $u \in C(\mathbb{R}^n \setminus \{0\})$ be a nonnegative solution of (4.1). Then, using (4.4) and (4.5) we get

$$u_{x,\lambda}(\xi) = \int_{\mathbb{R}^n} \frac{|z^{x,\lambda}|^{-\alpha}u_{x,\lambda}(z)^p}{|\xi-z|^{n-2\sigma}} \left(\frac{\lambda}{|z-x|}\right)^{n+2\sigma-p(n-2\sigma)} dz \quad (4.6)$$

and

$$u(\xi) - u_{x,\lambda}(\xi) = \int_{|z-x|\geq\lambda} K(x, \lambda; \xi, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\lambda}{|z-x|}\right)^{n+2\sigma-p(n-2\sigma)} \frac{u_{x,\lambda}(z)^p}{|z^{x,\lambda}|^\alpha} \right] dz, \quad (4.7)$$

where

$$K(x, \lambda; \xi, z) = \frac{1}{|\xi-z|^{n-2\sigma}} - \left(\frac{\lambda}{|\xi-x|}\right)^{n-2\sigma} \frac{1}{|\xi^{x,\lambda}-z|^{n-2\sigma}}.$$

It is elementary to check that

$$K(x, \lambda; \xi, z) > 0 \quad \text{for all } |\xi-x|, |z-x| > \lambda > 0.$$

We first prove the following result, which allows us to move the spheres from a starting point.

Lemma 4.3. *Suppose that $0 < \sigma < n/2$ and $-\infty < \alpha < 2\sigma$. Let $u \in C(\mathbb{R}^n \setminus \{0\})$ be a positive solution of (4.1) with $p > 0$. Then for every $x \in \mathbb{R}^n \setminus \{0\}$, there exists a real number $\lambda_2 \in (0, |x|)$ such that for any $0 < \lambda < \lambda_2$, we have*

$$u_{x,\lambda}(y) \leq u(y) \quad \forall |y-x| \geq \lambda, y \neq 0, \quad (4.8)$$

where $u_{x,\lambda}$ is the Kelvin transform of u as defined in (4.2).

Proof. The proof of Lemma 4.3 consists of two steps.

Step 1. There exists $0 < \lambda_1 < |x|$ such that for any $0 < \lambda < \lambda_1$,

$$u_{x,\lambda}(y) \leq u(y) \quad \forall 0 < \lambda \leq |y-x| \leq \lambda_1. \quad (4.9)$$

By [41, Theorem 2.5], we have $u \in C^1(\mathbb{R}^n \setminus \{0\})$. Suppose that

$$|\nabla \ln u| \leq C_1 \quad \text{in } B_{|x|/2}(x)$$

for some constant $C_1 > 0$. Then we have

$$\begin{aligned} \frac{d}{dr} (r^{\frac{n-2\sigma}{2}} u(x+r\theta)) &= r^{\frac{n-2\sigma}{2}-1} u(x+r\theta) \left(\frac{n-2\sigma}{2} - r \frac{\nabla u \cdot \theta}{u} \right) \\ &\geq r^{\frac{n-2\sigma}{2}-1} u(x+r\theta) \left(\frac{n-2\sigma}{2} - C_1 r \right) > 0 \end{aligned} \quad (4.10)$$

for all $0 < r < \lambda_1 := \min\{\frac{n-2\sigma}{2C_1}, \frac{|x|}{2}\}$ and $\theta \in \mathbb{S}^{n-1}$. For any $0 < \lambda < |y-x| \leq \lambda_1$, let $\theta = \frac{y-x}{|y-x|}$, $r_1 = |y-x|$ and $r_2 = \frac{\lambda^2 r_1}{|y-x|^2}$. Applying (4.10) we obtain

$$r_2^{\frac{n-2\sigma}{2}} u(x+r_2\theta) < r_1^{\frac{n-2\sigma}{2}} u(x+r_1\theta).$$

This is equivalent to

$$u_{x,\lambda}(y) \leq u(y), \quad 0 < \lambda \leq |y-x| \leq \lambda_1.$$

Step 2. There exists $0 < \lambda_2 < \lambda_1 < |x|$ such that (4.8) holds for all $0 < \lambda < \lambda_2$. By Fatou lemma we get

$$\liminf_{|x| \rightarrow \infty} |x|^{n-2\sigma} u(x) = \liminf_{|x| \rightarrow \infty} \int_{\mathbb{R}^n} \frac{|x|^{n-2\sigma} u(y)^p}{|y|^\alpha |x-y|^{n-2\sigma}} dy \geq \int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^\alpha} dy > 0.$$

Hence, there exist two constants $c_0, R_0 > 0$ such that

$$u(y) \geq \frac{c_0}{|y|^{n-2\sigma}} \quad \text{for all } |y| \geq R_0. \quad (4.11)$$

For $y \in B_{R_0} \setminus \{0\}$, by the equation (4.1) and positivity of u we have

$$u(y) \geq \int_{B_{R_0}} \frac{u(z)^p}{|y|^\alpha |y-z|^{n-2\sigma}} dz \geq (2R_0)^{2\sigma-n} \int_{B_{R_0}} \frac{u(z)^p}{|y|^\alpha} dz > 0.$$

This, together with (4.11), implies that there exists $c_1 > 0$ such that

$$u(y) \geq \frac{c_1}{|y-x|^{n-2\sigma}} \quad \forall |y-x| \geq \lambda_1, y \neq 0.$$

Thus, for sufficiently small $\lambda_2 \in (0, \lambda_1)$ and for any $0 < \lambda < \lambda_2$, we have

$$\begin{aligned} u_{x,\lambda}(y) &= \left(\frac{\lambda}{|y-x|} \right)^{n-2\sigma} u \left(x + \frac{\lambda^2(y-x)}{|y-x|^2} \right) \\ &\leq \left(\frac{\lambda_2}{|y-x|} \right)^{n-2\sigma} \sup_{B_{\lambda_1}(x)} u \leq u(y), \quad \forall |y-x| \geq \lambda_1, y \neq 0. \end{aligned}$$

From (4.9) and the above estimate, the proof of Lemma 4.3 is completed. \square

For any $x \in \mathbb{R}^n \setminus \{0\}$, we define

$$\bar{\lambda}(x) := \sup\{0 < \mu \leq |x| \mid u_{x,\lambda}(y) \leq u(y), \forall |y-x| \geq \lambda, y \neq 0, \forall 0 < \lambda < \mu\}.$$

By Lemma 4.3, $\bar{\lambda}(x)$ is well defined and $\bar{\lambda}(x) > 0$. Next we will show that $\bar{\lambda}(x) = |x|$ for all $x \in \mathbb{R}^n \setminus \{0\}$. Before proving this, we need the following lemma which is similar to [13, Lemma 4] and [39, Lemma 2.1]. We include its proof here for completeness.

Lemma 4.4. For $\lambda \in (0, |x|)$, we have

$$\frac{\lambda^2|z|}{|z-x|^2 \left| x + \frac{\lambda^2(z-x)}{|z-x|^2} \right|} < 1 \quad (4.12)$$

for any $z \in \mathbb{R}^n$ satisfying $|z-x| > \lambda$.

Proof. For any $z \in \mathbb{R}^n$ with $|z-x| > \lambda$, we have that

$$\begin{aligned} (4.12) &\Leftrightarrow \lambda^2|z| < |z-x|^2 \left| x + \frac{\lambda^2(z-x)}{|z-x|^2} \right| \\ &\Leftrightarrow \lambda^2|z| < (|z-x|^2 - \lambda^2)x + \lambda^2z \\ &\Leftrightarrow -2\lambda^2\langle z, x \rangle < (|z-x|^2 - \lambda^2)|x|^2 \quad (\text{by taking square}) \\ &\Leftrightarrow -2\lambda^2\langle (z-x), x \rangle < (|z-x|^2 + \lambda^2)|x|^2. \end{aligned}$$

The last inequality holds since

$$-2\lambda^2\langle (z-x), x \rangle \leq 2\lambda^2|z-x||x| < 2\lambda|z-x||x|^2 < (|z-x|^2 + \lambda^2)|x|^2.$$

Lemma 4.4 is established. \square

Lemma 4.5. Suppose that $0 < \sigma < n/2$ and $0 \leq \alpha < 2\sigma$. Let $u \in C(\mathbb{R}^n \setminus \{0\})$ be a positive solution of (4.1) with $0 < p \leq \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Then $\bar{\lambda}(x) = |x|$ for all $x \in \mathbb{R}^n \setminus \{0\}$.

Proof. Suppose $\bar{\lambda}(x) < |x|$ for some $x \in \mathbb{R}^n \setminus \{0\}$. For simplicity we write $\bar{\lambda} = \bar{\lambda}(x)$ and denote $\bar{\delta} := \min\{1, \frac{|x|-\bar{\lambda}}{2}\} > 0$. By the definition of $\bar{\lambda}$,

$$u_{x,\bar{\lambda}}(y) \leq u(y) \quad \text{for all } |y-x| \geq \bar{\lambda}, y \neq 0. \quad (4.13)$$

We next will separate into three cases to show that $u_{x,\bar{\lambda}}(y) < u(y)$ for any $|y-x| \geq \bar{\lambda}$ with $y \neq 0$, and there exists $\varepsilon_1 \in (0, 1)$ such that

$$(u - u_{x,\bar{\lambda}})(y) \geq \frac{\varepsilon_1}{|y-x|^{n-2\sigma}} \quad \text{for all } |y-x| \geq \bar{\lambda} + \bar{\delta}, y \neq 0. \quad (4.14)$$

Case 1: $0 < \alpha < 2\sigma$. Since $n + 2\sigma - p(n - 2\sigma) \geq 2\alpha$, we obtain that for $|z-x| \geq \lambda > 0$,

$$\left(\frac{\lambda}{|z-x|} \right)^{n+2\sigma-p(n-2\sigma)} \leq \left(\frac{\lambda}{|z-x|} \right)^{2\alpha}.$$

Therefore, using (4.7), (4.13), Lemma 4.4 and the positivity of the kernel K , we have, for any $|y-x| \geq \bar{\lambda}$ with $y \neq 0$,

$$\begin{aligned} u(y) - u_{x,\bar{\lambda}}(y) &= \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\bar{\lambda}}{|z-x|} \right)^{n+2\sigma-p(n-2\sigma)} \frac{u_{x,\bar{\lambda}}(z)^p}{|z^{x,\bar{\lambda}}|^\alpha} \right] dz \\ &\geq \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\bar{\lambda}}{|z-x|} \right)^{2\alpha} \frac{u_{x,\bar{\lambda}}(z)^p}{|z^{x,\bar{\lambda}}|^\alpha} \right] dz \\ &\geq \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[1 - \left(\frac{\bar{\lambda}^2|z|}{|z-x|^2|x + \frac{\lambda^2(z-x)}{|z-x|^2}|} \right)^\alpha \right] \frac{u_{x,\bar{\lambda}}(z)^p}{|z|^\alpha} dz \\ &> 0. \end{aligned}$$

This, together with Fatou Lemma, yields

$$\begin{aligned}
& \liminf_{|y| \rightarrow \infty} |y-x|^{n-2\sigma} (u - u_{x,\bar{\lambda}})(y) \\
& \geq \liminf_{|y| \rightarrow \infty} \int_{|z-x| \geq \bar{\lambda}} |y-x|^{n-2\sigma} K(x, \bar{\lambda}; y, z) \left[1 - \left(\frac{\bar{\lambda}^2 |z|}{|z-x|^2 |x + \frac{\bar{\lambda}^2(z-x)}{|z-x|^2}|} \right)^\alpha \right] \frac{u_{x,\bar{\lambda}}(z)^p}{|z|^\alpha} dz \\
& \geq \int_{|z-x| \geq \bar{\lambda}} \left[1 - \left(\frac{\bar{\lambda}}{|z-x|} \right)^{n-2\sigma} \right] \left[1 - \left(\frac{\bar{\lambda}^2 |z|}{|z-x|^2 |x + \frac{\bar{\lambda}^2(z-x)}{|z-x|^2}|} \right)^\alpha \right] \frac{u_{x,\bar{\lambda}}(z)^p}{|z|^\alpha} dz > 0.
\end{aligned}$$

Consequently, there exist $c_2 > 0$ and sufficiently large $R_2 > 0$ such that

$$(u - u_{x,\bar{\lambda}})(y) \geq \frac{c_2}{|y-x|^{n-2\sigma}} \quad \text{for all } |y-x| \geq R_2. \quad (4.15)$$

On the other hand, by the positivity of the kernel K there exists $\beta_1 > 0$ such that for all $\bar{\lambda} + \bar{\delta} \leq |y-x| \leq R_2$ and $2R_2 \leq |z-x| \leq 4R_2$,

$$K(x, \bar{\lambda}; y, z) \geq \beta_1 > 0. \quad (4.16)$$

Thus, for $\bar{\lambda} + \bar{\delta} \leq |y-x| \leq R_2$ and $y \neq 0$, we have

$$\begin{aligned}
u(y) - u_{x,\bar{\lambda}}(y) & \geq \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[1 - \left(\frac{\bar{\lambda}^2 |z|}{|z-x|^2 |x + \frac{\bar{\lambda}^2(z-x)}{|z-x|^2}|} \right)^\alpha \right] \frac{u_{x,\bar{\lambda}}(z)^p}{|z|^\alpha} dz \\
& \geq \int_{2R_2 \leq |z-x| \leq 4R_2} \beta_1 \left[1 - \left(\frac{\bar{\lambda}^2 |z|}{|z-x|^2 |x + \frac{\bar{\lambda}^2(z-x)}{|z-x|^2}|} \right)^\alpha \right] \frac{u_{x,\bar{\lambda}}(z)^p}{|z|^\alpha} dz \\
& =: C_2 > 0.
\end{aligned}$$

Combining this with (4.15), there exists $\varepsilon_1 \in (0, 1)$ such that

$$(u - u_{x,\bar{\lambda}})(y) \geq \frac{\varepsilon_1}{|y-x|^{n-2\sigma}} \quad \text{for all } |y-x| \geq \bar{\lambda} + \bar{\delta}, y \neq 0.$$

Case 2: $\alpha = 0$ and $0 < p < \frac{n+2\sigma}{n-2\sigma}$. Using (4.7), (4.13) and the positivity of the kernel K , we have, for any $|y-x| \geq \bar{\lambda}$ with $y \neq 0$,

$$\begin{aligned}
u(y) - u_{x,\bar{\lambda}}(y) & = \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[u(z)^p - \left(\frac{\bar{\lambda}}{|z-x|} \right)^{n+2\sigma-p(n-2\sigma)} u_{x,\bar{\lambda}}(z)^p \right] dz \\
& \geq \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[1 - \left(\frac{\bar{\lambda}}{|z-x|} \right)^{n+2\sigma-p(n-2\sigma)} \right] u_{x,\bar{\lambda}}(z)^p dz \\
& > 0.
\end{aligned}$$

By a similar argument as in Case 1, we obtain that (4.14) holds.

Case 3: $\alpha = 0$, $p = \frac{n+2\sigma}{n-2\sigma}$ and 0 is a non-removable singularity. In this case, we have $u_{x,\bar{\lambda}}(y) \neq u(y)$. By (4.7), (4.13) and the positivity of the kernel K , we have, for any $|y-x| \geq \bar{\lambda}$ with $y \neq 0$,

$$u(y) - u_{x,\bar{\lambda}}(y) = \int_{|z-x| \geq \bar{\lambda}} K(x, \bar{\lambda}; y, z) \left[u(z)^{\frac{n+2\sigma}{n-2\sigma}} - u_{x,\bar{\lambda}}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right] dz > 0.$$

By Fatou lemma, we have

$$\begin{aligned} & \liminf_{|y| \rightarrow \infty} |y-x|^{n-2\sigma} (u - u_{x,\bar{\lambda}})(y) \\ & \geq \int_{|z-x| \geq \bar{\lambda}} \left[1 - \left(\frac{\bar{\lambda}}{|z-x|} \right)^{n-2\sigma} \right] \left(u(z)^{\frac{n+2\sigma}{n-2\sigma}} - u_{x,\bar{\lambda}}(z)^{\frac{n+2\sigma}{n-2\sigma}} \right) dz > 0. \end{aligned}$$

An argument similar to the Case 1 leads to that (4.14) holds.

Therefore, in any case, we have that (4.14) holds. By (4.14) and the explicit formula of $u_{x,\lambda}$, there exists a small $\varepsilon_2 \in (0, \bar{\delta}/2)$ such that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon_2 < |x|$,

$$\begin{aligned} (u - u_{x,\lambda})(y) & \geq \frac{\varepsilon_1}{|y-x|^{n-2\sigma}} + (u_{x,\bar{\lambda}} - u_{x,\lambda})(y) \\ & \geq \frac{\varepsilon_1}{2|y-x|^{n-2\sigma}} \quad \forall |y-x| \geq \bar{\lambda} + \bar{\delta}, y \neq 0. \end{aligned} \tag{4.17}$$

For $\varepsilon \in (0, \varepsilon_2)$ which we choose below, using (4.7) and Lemma 4.4, we have, for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$ and $\lambda \leq |y-x| \leq \bar{\lambda} + \bar{\delta}$,

$$\begin{aligned} u(y) - u_{x,\lambda}(y) & = \int_{|z-x| \geq \lambda} K(x, \lambda; y, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\lambda}{|z-x|} \right)^{n+2\sigma-p(n-2\sigma)} \frac{u_{x,\lambda}(z)^p}{|z^{x,\lambda}|^\alpha} \right] dz \\ & \geq \int_{|z-x| \geq \lambda} K(x, \lambda; y, z) \left[\frac{u(z)^p}{|z|^\alpha} - \left(\frac{\lambda^2 |z|}{|z-x|^2 |x + \frac{\lambda^2(z-x)}{|z-x|^2}|} \right)^\alpha \frac{u_{x,\lambda}(z)^p}{|z|^\alpha} \right] dz \\ & \geq \int_{|z-x| \geq \lambda} K(x, \lambda; y, z) \frac{u(z)^p - u_{x,\lambda}(z)^p}{|z|^\alpha} dz, \end{aligned}$$

and thus, by (4.17) and (4.13),

$$\begin{aligned} u(y) - u_{x,\lambda}(y) & \geq \int_{\lambda \leq |z-x| \leq \bar{\lambda} + \bar{\delta}} K(x, \lambda; y, z) \frac{u(z)^p - u_{x,\lambda}(z)^p}{|z|^\alpha} dz \\ & \quad + \int_{\bar{\lambda} + 2 \leq |z-x| \leq \bar{\lambda} + 3} K(x, \lambda; y, z) \frac{u(z)^p - u_{x,\lambda}(z)^p}{|z|^\alpha} dz \\ & \geq -C \int_{\lambda \leq |z-x| \leq \lambda + \varepsilon} K(x, \lambda; y, z) \frac{|z-x| - \lambda}{|z|^\alpha} dz \\ & \quad + \int_{\lambda + \varepsilon < |z-x| \leq \bar{\lambda} + \bar{\delta}} K(x, \lambda; y, z) \frac{u_{x,\bar{\lambda}}(z)^p - u_{x,\lambda}(z)^p}{|z|^\alpha} dz \\ & \quad + \int_{\bar{\lambda} + 2 \leq |z-x| \leq \bar{\lambda} + 3} K(x, \lambda; y, z) \frac{u(z)^p - u_{x,\lambda}(z)^p}{|z|^\alpha} dz, \end{aligned}$$

where we have used

$$|u(z)^p - u_{x,\lambda}(z)^p| \leq C(|z-x| - \lambda)$$

in the second inequality. By (4.17), there exists $\delta_1 > 0$ such that for any $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$,

$$u(z)^p - u_{x,\lambda}(z)^p > \delta_1 \quad \forall \bar{\lambda} + 2 \leq |z-x| \leq \bar{\lambda} + 3, \quad z \neq 0.$$

Since $u \in C^1(B_{\bar{\lambda}+\varepsilon_2}(x))$, there exists $C > 0$ (independent of ε) such that for all $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$,

$$|u_{x,\bar{\lambda}}(z)^p - u_{x,\lambda}(z)^p| \leq C(\lambda - \bar{\lambda}) \leq C\varepsilon \quad \forall \bar{\lambda} \leq \lambda \leq |z-x| \leq \bar{\lambda} + \bar{\delta}.$$

Moreover, we have, for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$ and for $\lambda \leq |y-x| \leq \bar{\lambda} + \bar{\delta}$,

$$\begin{aligned} \int_{\lambda+\varepsilon \leq |z-x| \leq \bar{\lambda}+\bar{\delta}} \frac{K(x, \lambda; y, z)}{|z|^\alpha} dz &\leq C \left| \int_{\lambda+\varepsilon \leq |z-x| \leq \bar{\lambda}+\bar{\delta}} \left(\frac{1}{|y-z|^{n-2\sigma}} - \frac{1}{|y^{x,\lambda}-z|^{n-2\sigma}} \right) dz \right| \\ &\quad + C \int_{\lambda+\varepsilon \leq |z-x| \leq \bar{\lambda}+\bar{\delta}} \left| \left(\frac{\lambda}{|y-x|} \right)^{n-2\sigma} - 1 \right| \frac{1}{|y^{x,\lambda}-z|^{n-2\sigma}} dz \\ &\leq C(\varepsilon^{2\sigma-1} + |\ln \varepsilon| + 1)(|y-x| - \lambda) \end{aligned}$$

and

$$\begin{aligned} \int_{\lambda \leq |z-x| \leq \lambda+\varepsilon} K(x, \lambda; y, z) \frac{|z-x| - \lambda}{|z|^\alpha} dz &\leq C \left| \int_{\lambda \leq |z-x| \leq \lambda+\varepsilon} \left(\frac{|z-x| - \lambda}{|y-z|^{n-2\sigma}} - \frac{|z-x| - \lambda}{|y^{x,\lambda}-z|^{n-2\sigma}} \right) dz \right| \\ &\quad + \varepsilon C \int_{\lambda \leq |z-x| \leq \lambda+\varepsilon} \left| \left(\frac{\lambda}{|y-x|} \right)^{n-2\sigma} - 1 \right| \frac{1}{|y^{x,\lambda}-z|^{n-2\sigma}} dz \\ &\leq C(|y-x| - \lambda)\varepsilon^{2\sigma/n} + C\varepsilon(|y-x| - \lambda) \\ &\leq C(|y-x| - \lambda)\varepsilon^{2\sigma/n}. \end{aligned}$$

On the other hand, since $K(x, \lambda; y, z) = 0$ for $y \in \partial B_\lambda(x)$ and

$$\langle \nabla_y K(x, \lambda; y, z), y-x \rangle \Big|_{|y-x|=\lambda} = (n-2\sigma)|y-z|^{2\sigma-n-2}(|z-x|^2 - |y-x|^2) > 0$$

for all $\bar{\lambda} + 2 \leq |z-x| \leq \bar{\lambda} + 3$, and using the positivity of the kernel K , we obtain that, for $\bar{\lambda} \leq \lambda \leq |y-x| \leq \bar{\lambda} + \bar{\delta}$,

$$K(x, \lambda; y, z) \geq \delta_2(|y-x| - \lambda) \quad \forall \bar{\lambda} + 2 \leq |z-x| \leq \bar{\lambda} + 3,$$

where $\delta_2 > 0$ is independent of ε . Thus, it follows from the above that for $\bar{\lambda} \leq \lambda \leq \bar{\lambda} + \varepsilon$ and for $\lambda \leq |y-x| \leq \bar{\lambda} + \bar{\delta}$,

$$u(y) - u_{x,\lambda}(y) \geq \left(-C\varepsilon^{\frac{2\sigma}{n}} + \delta_1\delta_2 \int_{\bar{\lambda}+2 \leq |z-x| \leq \bar{\lambda}+3} \frac{1}{|z|^\alpha} dz \right) (|y-x| - \lambda) \geq 0$$

if ε is chosen sufficiently small. This and (4.17) contradict the definition of $\bar{\lambda}$. Lemma 4.5 is established. \square

Proof of Theorem 4.1. We first assume that $0 < p \leq \frac{n+2\sigma-2\alpha}{n-2\sigma}$. Let $u \in C(\mathbb{R}^n \setminus \{0\})$ be a nonnegative solution of (4.1). If $u(x_0) = 0$ for some $x_0 \in \mathbb{R}^n \setminus \{0\}$, then

$$\int_{\mathbb{R}^n} \frac{u(y)^p}{|y|^\alpha |x_0 - y|^{n-2\sigma}} dy = 0,$$

which implies that $u \equiv 0$ on $\mathbb{R}^n \setminus \{0\}$. Next we suppose that $u \in C(\mathbb{R}^n \setminus \{0\})$ be a positive solution of (4.1). It follows from Lemma 4.5 that for every $x \in \mathbb{R}^n \setminus \{0\}$,

$$u_{x,\lambda}(y) \leq u(y) \quad \forall |y - x| \geq \lambda, y \neq 0, \quad \forall 0 < \lambda < |x|. \quad (4.18)$$

For any unit vector $e \in \mathbb{R}^n$, for any $a > 0$, for any $z \in \mathbb{R}^n \setminus \{0\}$ satisfying $(z - ae) \cdot e < 0$, and for any $R > a$, we have, by (4.18) with $x = Re$ and $\lambda = R - a$,

$$u(z) \geq u_{x,\lambda}(z) = \left(\frac{\lambda}{|z - x|} \right)^{n-2\sigma} u \left(x + \frac{\lambda^2(z - x)}{|z - x|^2} \right).$$

Sending R to infinity in the above, we obtain

$$u(z) \geq u(z - 2(z \cdot e - a)e). \quad (4.19)$$

Since $e \in \mathbb{R}^n$ and $a > 0$ are arbitrary, (4.19) implies that u is radially symmetric about the origin. Moreover, (4.19) also gives

$$u(z) = u(z_1, z_2, \dots, z_n) \geq u_a(z) := u(2a - z_1, z_2, \dots, z_n) \quad \forall z_1 \leq a, a > 0.$$

Thus, u is also monotonically decreasing about the origin.

Next we consider the case $\frac{n+2\sigma-2\alpha}{n-2\sigma} < p \leq \frac{n+2\sigma-\alpha}{n-2\sigma}$ and $0 < \alpha < 2\sigma$. Let $u \in C(\mathbb{R}^n \setminus \{0\})$ be a positive solution of (4.1). Define

$$v(x) = \left(\frac{1}{|x|} \right)^{n-2\sigma} u \left(\frac{x}{|x|^2} \right) \quad \text{for } x \neq 0.$$

By (4.6), v is a positive solution to the following integral equation

$$v(x) = \int_{\mathbb{R}^n} \frac{v(y)^p}{|y|^\theta |x - y|^{n-2\sigma}} dy$$

with $\theta := n + 2\sigma - \alpha - p(n - 2\sigma)$. Note that the new exponent θ satisfies

$$\begin{aligned} 0 \leq \theta &\iff p \leq \frac{n + 2\sigma - \alpha}{n - 2\sigma} \\ \theta < \alpha &\iff \frac{n + 2\sigma - 2\alpha}{n - 2\sigma} < p \quad (\text{hence } \theta < 2\sigma) \\ p < \frac{n + 2\sigma - 2\theta}{n - 2\sigma} &\iff p > \frac{n + 2\sigma - 2\alpha}{n - 2\sigma}. \end{aligned}$$

From the previous case, we know that v is radially symmetric about the origin and hence, u is also radially symmetric about the origin. Theorem 4.1 is established. \square

Proof of Theorem 1.5. It follows from Theorems 1.1 and 4.1. \square

References

- [1] S. Antman, Nonlinear problems of elasticity. Second edition. *Applied Mathematical Sciences*, 107. Springer, New York, 2005.
- [2] W. Ao, M. González, A. Hyder, J. Wei, Removability of singularities and superharmonicity for some fractional Laplacian equations. arXiv:2001.11683v2.
- [3] D. Applebaum, Lévy processes and stochastic calculus. Second edition. *Cambridge Studies in Advanced Mathematics*, 116. Cambridge University Press, Cambridge, 2009.
- [4] T. Aubin, Problemes isopérimétriques et espaces de Sobolev. *J. Differential Geom.*, **11** (4) (1976) 573-598.
- [5] B. Barrios, A. Quaas, The sharp exponent in the study of the nonlocal Hénon equation in \mathbb{R}^N : a Liouville theorem and an existence result. *Calc. Var. Partial Differential Equations*, **59** (2020), no. 4, Paper No. 114, 22 pp.
- [6] J.-P. Bouchaud, A. Georges, Anomalous diffusion in disordered media: statistical mechanisms, models and physical applications, *Phys. Rep.*, **195** (1990) 127-293.
- [7] L. Caffarelli, B. Gidas, J. Spruck, Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth. *Comm. Pure Appl. Math.*, **42** (1989) 271-297.
- [8] L. Caffarelli, S. Salsa, L. Silvestre, Regularity estimates for the solution and the free boundary of the obstacle problem for the fractional Laplacian. *Invent. Math.*, **171** (2008), no. 2, 425-461.
- [9] L. Caffarelli, T. Jin, Y. Sire, J. Xiong, Local analysis of solutions of fractional semi-linear elliptic equations with isolated singularities. *Arch. Ration. Mech. Anal.*, **213** (2014), no.1, 245-268.
- [10] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian. *Comm. Partial. Diff. Equ.*, **32** (2007) 1245-1260.
- [11] L. Caffarelli, A. Vasseur, Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation, *Ann. of Math. (2)* **171** (2010), no. 3, 1903-1930.
- [12] D. Cao, W. Dai, G. Qin, Super poly-harmonic properties, Liouville theorems and classification of nonnegative solutions to equations involving higher-order fractional Laplacians. *Trans. Amer. Math. Soc.*, **374** (2021), no. 7, 4781-4813.
- [13] D. Cao, Y.Y. Li, Results on positive solutions of elliptic equations with a critical Hardy-Sobolev operator. *Methods Appl. Anal.*, **15** (2008), no. 1, 81-95.
- [14] J. Case, S.-Y. A. Chang, On fractional GJMS operators. *Comm. Pure Appl. Math.*, **69** (2016), no. 6, 1017-1061.
- [15] S.-Y. A. Chang, M. González, Fractional Laplacian in conformal geometry. *Adv. Math.*, **226** (2011), no. 2, 1410-1432.
- [16] S.-Y. A. Chang, P. C. Yang, On uniqueness of solutions of n -th order differential equations in conformal geometry. *Math. Res. Lett.*, **4** (1997), no. 1, 91-102.

- [17] W. Chen, C. Li, Classification of solutions of some nonlinear elliptic equations. *Duke Math. J.*, **63** (1991), no. 3, 615-622.
- [18] W. Chen, C. Li, Y. Li, A direct method of moving planes for the fractional Laplacian. *Adv. Math.*, **308** (2017) 404-437.
- [19] W. Chen, C. Li, B. Ou, Classification of solutions for an integral equation. *Comm. Pure Appl. Math.*, **59** (2006) 330-343.
- [20] W. Chen, C. Li, B. Ou, Qualitative properties of solutions for an integral equation. *Discrete Contin. Dyn. Syst.*, **12** (2005), no. 2, 347-354.
- [21] T. Cheng, S. Liu, A Liouville type theorem for higher order Hardy-Hénon equation in \mathbb{R}^n . *J. Math. Anal. Appl.*, **444** (2016) 370-389.
- [22] P. Constantin, Euler equations, Navier-Stokes equations and turbulence. Mathematical foundation of turbulent viscous flows, 1-43, *Lecture Notes in Math.*, 1871, Springer, Berlin, 2006.
- [23] R. Cont, P. Tankov, Financial modelling with jump processes. *Chapman & Hall/CRC Financial Mathematics Series*. Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [24] W. Dai, S. Peng, G. Qin, Liouville type theorems, a priori estimates and existence of solutions for non-critical higher order Lane-Emden-Hardy equations. arXiv:1808.10771 (2018).
- [25] W. Dai, G. Qin, Liouville type theorems for fractional and higher order Hénon-Hardy type equations via the method of scaling spheres. arXiv:1810.02752v7 (2020).
- [26] A. DelaTorre, M. del Pino, M. González, J. Wei, Delaunay-type singular solutions for the fractional Yamabe problem. *Math. Ann.*, **369** (2017) 597-626.
- [27] S. Dipierro, A. Figalli, E. Valdinoci, Strongly nonlocal dislocation dynamics in crystals. *Comm. Partial Differential Equations*, **39** (2014), no. 12, 2351-2387.
- [28] J. Dou, H. Zhou, Liouville theorems for fractional Hénon equation and system on \mathbb{R}^n . *Commun. Pure Appl. Anal.*, **14** (2015), no. 5, 1915-1927.
- [29] X. Du, H. Yang, Local behavior of positive solutions of higher order conformally invariant equations with a singular set. *Calc. Var. Partial Differential Equations*, **60** (2021), no. 6, Paper No. 204.
- [30] R. H. Fowler, Further studies of Emden's and similar differential equations. *Q. J. Math., Oxf. Ser. 2* (1931) 259-288.
- [31] R. Frank, T. König, Classification of positive solutions to a nonlinear biharmonic equation with critical exponent. *Anal. PDE*, **12** (2019) 1101-1113.
- [32] B. Gidas, W. Ni, L. Nirenberg, Symmetry and related properties via the maximum principle. *Comm. Math. Phys.*, **68** (1979), no. 3, 209-243.
- [33] B. Gidas, J. Spruck, Global and local behavior of positive solutions of nonlinear elliptic equations. *Comm. Pure Appl. Math.*, **34** (1981), no. 4, 525-598.
- [34] V. Glaser, A. Martin, H. Grosse, W. Thirring, A family of optimal conditions for the absence of bound states in a potential, in: E.H. Lieb, B. Simon, A.S. Wightman (Eds.), *Studies in Mathematical Physics*, Princeton University Press, 1976, pp. 169-194.

- [35] Z. Guo, X. Huang, L. Wang, J. Wei, On Delaunay solutions of a biharmonic elliptic equation with critical exponent. *J. Anal. Math.*, **140** (2020), no. 1, 371-394.
- [36] Z. Guo, F. Wan, Further study of a weighted elliptic equation. *Sci. China Math.*, **60** (2017), no. 12, 2391-2406.
- [37] Z.-C. Han, Y.Y. Li, E. V. Teixeira, Asymptotic behavior of solutions to the σ_k -Yamabe equation near isolated singularities. *Invent. Math.*, **182** (2010), no. 3, 635-684.
- [38] M. Hénon, Numerical experiments on the stability of spherical stellar systems. *Astron. Astrophys.*, **24** (1973) 229-238.
- [39] T. Jin, Symmetry and nonexistence of positive solutions of elliptic equations and systems with Hardy terms. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **28** (2011), no. 6, 965-981.
- [40] T. Jin, Y.Y. Li, J. Xiong, On a fractional Nirenberg problem, part I: blow up analysis and compactness of solutions. *J. Eur. Math. Soc.*, **16** (2014) 1111-1171.
- [41] T. Jin, Y.Y. Li, J. Xiong, The Nirenberg problem and its generalizations: a unified approach. *Math. Ann.*, **369** (2017), no. 1-2, 109-151.
- [42] T. Jin, J. Xiong, Asymptotic symmetry and local behavior of solutions of higher order conformally invariant equations with isolated singularities. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **38** (2021), no. 4, 1167-1216.
- [43] N. Korevaar, R. Mazzeo, F. Pacard, R. Schoen, Refined asymptotics for constant scalar curvature metrics with isolated singularities. *Invent. Math.*, **135** (2) (1999) 233-272.
- [44] Y.Y. Li, Remark on some conformally invariant integral equations: the method of moving spheres. *J. Eur. Math. Soc. (JEMS)*, **6** (2004) 153-180.
- [45] Y.Y. Li, L. Zhang, Liouville-type theorems and Harnack-type inequalities for semilinear elliptic equations. *J. Anal. Math.*, **90** (2003) 27-87.
- [46] Y.Y. Li, M. Zhu, Uniqueness theorems through the method of moving spheres. *Duke Math. J.*, **80** (1995) 383-418.
- [47] Y. Li, J. Bao, Fractional Hardy-Hénon equations on exterior domains. *J. Differential Equations*, **266** (2019), no. 2-3, 1153-1175.
- [48] E. H. Lieb, Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities. *Ann. of Math. (2)*, **118** (1983), no. 2, 349-374.
- [49] C.-S. Lin, A classification of solutions of a conformally invariant fourth order equation in \mathbb{R}^n . *Comment. Math. Helv.*, **73** (1998) 206-231.
- [50] G. Lu, J. Zhu, Symmetry and regularity of extremals of an integral equation related to the Hardy-Sobolev inequality. *Calc. Var. Partial Differential Equations*, **42** (2011), no. 3-4, 563-577.
- [51] R. Musina, A. I. Nazarov, Complete classification and nondegeneracy of minimizers for the fractional Hardy-Sobolev inequality, and applications. *J. Differential Equations*, **280** (2021), 292-314.
- [52] W. Reichel, H. Zou, Non-existence results for semilinear cooperative elliptic systems via moving spheres. *J. Differential Equations*, **161** (2000), no. 1, 219-243.

- [53] R. Schoen, The existence of weak solutions with prescribed singular behavior for a conformally invariant scalar equation. *Comm. Pure Appl. Math.*, **41** (1988), no. 3, 317-392.
- [54] R. Schoen, S.-T. Yau, Conformally flat manifolds, Kleinian groups and scalar curvature. *Invent. Math.*, **92** (1988) 47-71.
- [55] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator. *Comm. Pure Appl. Math.*, **60** (2007), no. 1, 67-112.
- [56] G. Talenti, Best constant in Sobolev inequality. *Ann. Mat. Pura Appl.*, **110** (1976) 353-372.
- [57] W. Wang, K. Li, L. Hong, Nonexistence of positive solutions of $-\Delta u = K(x)u^p$ in \mathbb{R}^n . *Appl. Math. Lett.*, **18** (2005), no. 3, 345-351.
- [58] J. Wei, X. Xu, Classification of solutions of higher order conformally invariant equation. *Math. Ann.*, **313** (1999) 207-228.
- [59] H. Yang, Asymptotic behavior of positive solutions to a nonlinear biharmonic equation near isolated singularities. *Calc. Var. Partial Differential Equations*, **59** (2020), no. 4, Paper No. 130, 21 pp.
- [60] H. Yang, W. Zou, Exact asymptotic behavior of singular positive solutions of fractional semi-linear elliptic equations. *Proc. Amer. Math. Soc.*, **147** (2019), no. 7, 2999-3009.
- [61] H. Yang, W. Zou, On isolated singularities of fractional semi-linear elliptic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **38** (2021), no. 2, 403-420.
- [62] H. Yang, W. Zou, Sharp blow up estimates and precise asymptotic behavior of singular positive solutions to fractional Hardy-Hénon equations. *J. Differential Equations*, **278** (2021) 393-429.
- [63] R. Zhuo, W. Chen, X. Cui, Z. Yuan, Symmetry and non-existence of solutions for a nonlinear system involving the fractional Laplacian. *Discrete Contin. Dyn. Syst.*, **36** (2016), no. 2, 1125-1141.