

GROUND STATES FOR THE PLANAR NLSE WITH A POINT DEFECT AS MINIMIZERS OF THE CONSTRAINED ENERGY

RICCARDO ADAMI, FILIPPO BONI, RAFFAELE CARLONE, AND LORENZO TENTARELLI

ABSTRACT. We investigate the ground states for the focusing, subcritical nonlinear Schrödinger equation with a point defect in dimension two, defined as the minimizers of the energy functional at fixed mass. We prove that ground states exist for every positive mass and show a logarithmic singularity at the defect. Moreover, up to a multiplication by a constant phase, they are positive, radially symmetric, and decreasing along the radial direction. In order to overcome the obstacles arising from the uncommon structure of the energy space, that complicates the application of standard rearrangement theory, we move to the study of the minimizers of the action functional on the Nehari manifold and then establish a connection with the original problem. An ad hoc result on rearrangements is given to prove qualitative features of the ground states.

AMS Subject Classification: 35Q40, 35Q55, 35B07, 35B09, 35R99, 49J40, 49N15.

Keywords: standing waves, nonlinear Schrödinger, ground states, delta interaction, radially symmetric solutions, rearrangements.

1. INTRODUCTION

The Nonlinear Schrödinger Equation (NLSE) has provided for almost fifty years the effective description of the evolution of the wave function of a Bose-Einstein condensate (BEC) in the Gross-Pitaevskii regime. More recently, interest has grown in the possibility of modeling a BEC in the presence of defects or impurities by means of a NLSE with an additional pointwise interaction located at the defect. If the spatial scale of the impurity is supposed to be much smaller than the dispersion of the wave function, one can try to describe it by means of a Dirac's delta potential (see, e.g., [27, 45, 46]), obtaining the evolution equation

$$i\frac{\partial\psi}{\partial t} = (-\Delta + \beta\delta_0)\psi \pm |\psi|^{p-2}\psi, \quad \beta \in \mathbb{R} \setminus \{0\}, \quad p > 2, \quad (1)$$

where the sign of the nonlinearity establishes the character focusing or defocusing of the equation, and, correspondingly, the attractive or repulsive behaviour of the condensate.

A large part of the available results concerns the one-dimensional case, that models the so-called cigar-shaped condensates. In particular, well-posedness, existence, and stability of standing waves have been studied for several pointwise potentials ([11, 12, 10, 15, 39]), while the well-posedness for the two and three-dimensional models has been addressed in [22]. Here we aim at discussing the existence and the properties of the standing waves for the two-dimensional case, with particular reference to the ground states.

In our intention, the present paper is the first step of a research programme devoted to the study of the standing waves of the NLSE on multi-dimensional structures, that is domains consisting of pieces of different space dimensions glued together through suitable boundary conditions. Such structures are known in the literature as *quantum hybrids* and one of the simplest models is provided by a plane attached to a half-line. It has been shown ([25, 30, 31]) that the conditions to be imposed at the junction between the plane and the half-line prescribe a logarithmic singularity for the restriction of the wave function to the plane, exactly like for the Schrödinger equation with a point interaction. Therefore, the present work is the first step of our research plan towards nonlinear quantum hybrids.

Date: September 21, 2021.

Although all results and proofs hold in space dimension two, we believe that it should be possible to prove analogous results for the three-dimensional case. This will be the subject of a forthcoming paper.

A further research project inspired by the model (1) concerns a different family of singular perturbation of delta, called concentrated nonlinearities, namely

$$i\frac{\partial\psi}{\partial t} = (-\Delta + \beta|\psi|^{p-2}\delta_0)\psi, \quad \beta \in \mathbb{R} \setminus \{0\}, \quad p > 2. \quad (2)$$

Here the nonlinearity affects exclusively the interaction point where the defect is located. Such equation has been studied in one (cfr. [9, 16, 26, 36, 37]), two (cfr. [5, 6, 24]) and three dimensions (cfr. [7, 8, 13, 14]). It is natural to investigate the dynamics of a system in the presence of both types of nonlinearity. The one-dimensional case and the case of the star graphs have been already addressed in [18] and [2], while high-dimensional cases are still unexplored.

Finally, we recall that the issue of PDE's with point singularities is not new in the literature. In particular, the study of singular solutions for elliptic equations traces back to the eighties ([20, 27, 33, 38, 41, 43, 44, 47, 48]).

1.1. Setting and main results. We study equation (1) in space dimension two.

Let us stress preliminarily that writing (1) is formal, as in two dimensions the delta interaction is not a small perturbation of the laplacian. It is well-known that a rigorous definition can be given through the theory of self-adjoint extensions of hermitian operators. Eventually, one finds ([17]) that there exists a family $(H_\alpha)_{\alpha \in \mathbb{R}}$ of self-adjoint operators that realizes a nontrivial point perturbation of $-\Delta$.

The domains of such operators are

$$D(H_\alpha) := \{v \in L^2(\mathbb{R}^2) : \exists q \in \mathbb{C}, \lambda > 0 \text{ s.t.} \\ v - q\mathcal{G}_\lambda =: \phi_\lambda \in H^2(\mathbb{R}^2) \text{ and } \phi_\lambda(0) = (\alpha + \theta_\lambda)q\}, \quad (3)$$

and the action reads

$$H_\alpha v := -\Delta\phi_\lambda - q\lambda\mathcal{G}_\lambda, \quad \forall v \in D(H_\alpha). \quad (4)$$

We denoted

$$\theta_\lambda := \frac{\log\left(\frac{\sqrt{\lambda}}{2}\right) + \gamma}{2\pi}, \quad (5)$$

with γ denoting the Euler-Mascheroni constant, and \mathcal{G}_λ the Green's function of $-\Delta + \lambda$, namely

$$\mathcal{G}_\lambda(\mathbf{x}) := \frac{1}{2\pi}\mathcal{F}^{-1}\left[(|\mathbf{k}|^2 + \lambda)^{-1}\right](\mathbf{x}) = \frac{K_0(\sqrt{\lambda}\mathbf{x})}{2\pi}. \quad (6)$$

Here K_0 denotes the modified Bessel function of the second kind of order 0, also known as Macdonald function (see [1, Section 9.6]), and \mathcal{F} denotes the unitary Fourier transform.

The function \mathcal{G}_λ has a singular behaviour at the origin, namely

$$\mathcal{G}_\lambda(\mathbf{x}) = -\frac{\log|\mathbf{x}|}{2\pi} + o(\log|\mathbf{x}|), \quad \mathbf{x} \rightarrow 0$$

that prevents \mathcal{G}_λ from belonging to the space $H^1(\mathbb{R}^2)$.

Functions in $D(H_\alpha)$ consist then of a *regular part* ϕ_λ , on which the operator acts as the standard Laplacian, and a *singular part* $q\mathcal{G}_\lambda$, on which the operator acts as the multiplication by $-\lambda$. The two components are connected by the boundary condition $\phi_\lambda(0) = (\alpha + \theta_\lambda)q$. The strength q of the singular part is called the *charge*.

We highlight that λ is a dumb parameter as it does not affect the definition of H_α . To check this, consider $v \in D(H_\alpha)$. Therefore, there exist $q \in \mathbb{C}$ and $\lambda > 0$ such that

$v = \phi_\lambda + q\mathcal{G}_\lambda$, $\phi_\lambda \in H^2(\mathbb{R}^2)$ and $\phi_\lambda(0) = (\alpha + \theta_\lambda)q$. Fix $\nu > 0$. By (6)

$$\mathcal{G}_\lambda - \mathcal{G}_\nu \in H^2(\mathbb{R}^2),$$

thus $\phi_\nu := \phi_\lambda + q(\mathcal{G}_\lambda - \mathcal{G}_\nu)$ belongs to $H^2(\mathbb{R}^2)$. Moreover, (6) implies

$$(\mathcal{G}_\lambda - \mathcal{G}_\nu)(\mathbf{0}) = (4\pi)^{-1} \log(\nu/\lambda), \quad (7)$$

so that $\theta_\lambda + (\mathcal{G}_\lambda - \mathcal{G}_\nu)(\mathbf{0}) = \theta_\nu$, whence $\phi_\nu(0) = (\alpha + \theta_\nu)q$. Finally, by (6)

$$-\Delta(\mathcal{G}_\lambda - \mathcal{G}_\nu) = \nu\mathcal{G}_\nu - \lambda\mathcal{G}_\lambda,$$

so $-\Delta\phi_\lambda - q\lambda\mathcal{G}_\lambda = -\Delta\phi_\nu - q\nu\mathcal{G}_\nu$ and thus the decompositions with ν and λ are equivalent.

Remark 1.1. Another way to see the independence of H_α of λ is to rewrite (3)-(4) as follows:

$$D(H_\alpha) := \{v \in L^2(\mathbb{R}^2) : \exists q \in \mathbb{C} \text{ s.t.}$$

$$v - q\mathcal{G}_0 =: \phi \in H_{loc}^1(\mathbb{R}^2) \cap \dot{H}^2(\mathbb{R}^2) \text{ and } \phi(\mathbf{0}) = \alpha q\},$$

$$H_\alpha v := -\Delta\phi, \quad \forall \mathbf{x} \neq \mathbf{0}, \quad \forall v \in D(H_\alpha),$$

with $\mathcal{G}_0 := -\frac{\log|\mathbf{x}|}{2\pi}$ the Green function of $-\Delta$. However, through the paper we prefer to decompose the wave function using $\lambda > 0$, because in such a way the regular part ϕ_λ belongs to $H^1(\mathbb{R}^2)$, that allows a simple expression of the energy functional.

Finally, we recall that the spectrum of H_α is given by

$$\sigma(H_\alpha) = \{\ell_\alpha\} \cup [0, +\infty), \quad \text{with } \ell_\alpha := -4e^{-4\pi\alpha-2\gamma} < 0 \text{ sole eigenvalue.} \quad (8)$$

The rigorous form of the NLSE with a pointwise impurity (δ -NLSE) is therefore

$$i\frac{\partial\psi}{\partial t} = H_\alpha\psi - |\psi|^{p-2}\psi, \quad \alpha \in \mathbb{R}, \quad p > 2. \quad (9)$$

As proven in [22], the flow generated by (9) preserves the mass.

Standing waves are solutions to (9) of the form $\psi = e^{i\omega t}u(\mathbf{x})$, with $\omega \in \mathbb{R}$. An easy computation shows that ψ is a standing wave for (9) whenever u is a bound state, i.e.

$$u \in D(H_\alpha), \quad (10)$$

$$H_\alpha u + \omega u - |u|^{p-2}u = 0. \quad (11)$$

We focus on ground states, rather than generic bound states, as they are orbitally stable via [35, Theorem 3] and then physically observable. The proof that a ground state satisfies (10) and (11) is straightforward and is reported in Appendix A.

In order to give a precise definition of the ground states of (11), we first introduce the quadratic form associated with H_α , which has domain

$$D := \{v \in L^2(\mathbb{R}^2) : \exists q \in \mathbb{C}, \lambda > 0 \text{ s.t. } v - q\mathcal{G}_\lambda =: \phi_\lambda \in H^1(\mathbb{R}^2)\},$$

and action

$$Q(v) := \langle H_\alpha v, v \rangle = \|\nabla\phi_\lambda\|_2^2 + \lambda(\|\phi_\lambda\|_2^2 - \|v\|_2^2) + (\alpha + \theta_\lambda)|q|^2, \quad \forall v \in D, \quad (12)$$

where we denoted by $\langle \cdot, \cdot \rangle$ the hermitian product in $L^2(\mathbb{R}^2)$ and by $\|\cdot\|_p$ the usual norm in $L^p(\mathbb{R}^2)$. As (12) is the quadratic form of the self-adjoint operator H_α , it is independent of the choice of λ . Notice that, as expected when passing from operator to quadratic form, functions in D have a rougher regular part than function in $D(H_\alpha)$ (from $H^2(\mathbb{R}^2)$ to $H^1(\mathbb{R}^2)$), and that in D there is no boundary condition to link regular and singular part.

Let us introduce the main object of our study, the energy functional associated with the δ -NLSE.

Definition 1.2. Given $\alpha \in \mathbb{R}$ and $p > 2$, the δ -NLS energy is the functional $E : D \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} E(v) &:= \frac{1}{2}Q(v) - \frac{1}{p}\|v\|_p^p \\ &= \frac{1}{2}\|\nabla\phi_\lambda\|_2^2 + \frac{\lambda}{2}(\|\phi_\lambda\|_2^2 - \|v\|_2^2) + \frac{(\alpha + \theta_\lambda)}{2}|q|^2 - \frac{1}{p}\|v\|_p^p. \end{aligned} \quad (13)$$

The δ -NLS energy is conserved by the flow generated by (9) ([22]).

Remark 1.3. As a peculiar feature of point interactions in dimensions two and three, the energy space D is strictly larger than $H^1(\mathbb{R}^2)$. Furthermore, if v belongs to $H^1(\mathbb{R}^2)$, i.e. it has no charge, then its energy reduces to the standard NLS energy defined by

$$E^0(v) := \frac{1}{2}\|\nabla v\|_2^2 - \frac{1}{p}\|v\|_p^p,$$

so that the δ -NLS energy is an extension of the NLS energy.

We can now give the following fundamental definition.

Definition 1.4. Given $\mu > 0$, a function u belonging to the space

$$D_\mu := \{v \in D : \|v\|_2^2 = \mu\},$$

and satisfying

$$E(u) = \inf_{v \in D_\mu} E(v) =: \mathcal{E}(\mu),$$

is a *ground state at mass μ* for the NLSE with a point defect.

In other words, a ground state is a minimizer of the energy constrained to a submanifold of constant mass μ .

We can now state the main result of the paper.

Theorem 1.5 (δ -NLS ground states). *Let $p \in (2, 4)$ and $\alpha \in \mathbb{R}$. Therefore, for every $\mu > 0$,*

- (i) *there exists a ground state for the δ -NLS at mass μ ;*
- (ii) *if $u = \phi_\lambda + q\mathcal{G}_\lambda$ is a ground state, then:*
 - (a) *there does not exist $\lambda > 0$ such that ϕ_λ or $q\mathcal{G}_\lambda$ are identically zero,*
 - (b) *u is positive, radially symmetric, and decreasing along the radial direction, up to multiplication by a constant phase; moreover, ϕ_λ is nonnegative as $\lambda = \omega$, and positive as $\lambda > \omega$, with $\omega = \mu^{-1}(\|u\|_p^p - Q(u))$.*

Remark 1.6. One can also see that if u is a ground state for the δ -NLS, then the associated frequency $\omega = \mu^{-1}(\|u\|_p^p - Q(u))$ has to be positive. Indeed, by the Lagrange Multiplier Theorem, one has

$$\langle E'(u), v \rangle + \omega \langle u, v \rangle = 0, \quad \forall v \in D,$$

so that, setting $v = u$ and combining with (64), (12) and (13),

$$2E(u) - \frac{p-2}{p}\|u\|_p^p = -\omega\|u\|_2^2. \quad (14)$$

Then, by remark 1.3,

$$E(u) = \mathcal{E}(\mu) \leq \mathcal{E}^0(\mu) := \inf_{v \in H_\mu^1(\mathbb{R}^2)} E^0(v),$$

with $H_\mu^1(\mathbb{R}^2) := \{v \in H^1(\mathbb{R}^2) : \|v\|_2^2 = \mu\}$, which is negative whenever $p \in (2, 4)$ (see, e.g., [42] or the proof of Proposition 3.3). Thus (14) implies that $\omega > 0$.

We stress that Theorem 1.5 treats power nonlinearities with $p \in (2, 4)$ only, namely the so-called subcritical power nonlinearities of the NLSE, since as in the standard case this is necessary to establish boundedness from below of the constrained energy without prescriptions on the mass μ (see Proposition 3.1). In addition, also regarding existence, positivity, and symmetry, Theorem 1.5 retraces the results on the NLSE. However, point (ii)(a) shows that δ -NLS ground states and NLS ground states cannot coincide as the singular part of the former ones cannot vanish.

Moreover, point (ii)(a) marks a difference with the model (2). Indeed, it was proven in [5, 6] that for any bound state there exists a value of $\lambda > 0$ such that the regular part of the decomposition vanishes.

As a final remark, we highlight that, while point (i) of Theorem 1.5 is proved by minimization of E on D_μ , point (ii) is proved through minimization of another functional, called *Action*, constrained to a set called *Nehari manifold*. More precisely,

Definition 1.7. Fixed $\omega \in \mathbb{R}$, the δ -NLS action at frequency ω is the functional $S_\omega : D \rightarrow \mathbb{R}$ defined

$$S_\omega(v) := E(v) + \frac{\omega}{2} \|v\|_2^2, \quad (15)$$

with the following constraint

Definition 1.8. Fixed $\omega \in \mathbb{R}$, the Nehari manifold at frequency ω associated to the δ -NLS is defined by

$$N_\omega := \{v \in D \setminus \{0\} : I_\omega(v) = 0\}, \quad (16)$$

where $I_\omega : D \rightarrow \mathbb{R}$ denotes the quantity

$$I_\omega(v) := \langle S'_\omega(v), v \rangle = \|\nabla \phi_\lambda\|_2^2 + \lambda \|\phi_\lambda\|_2^2 + (\omega - \lambda) \|v\|_2^2 + (\alpha + \theta_\lambda) |q|^2 - \|v\|_p^p. \quad (17)$$

As a consequence, the minimizers of the δ -NLS action at frequency ω are all functions $u \in N_\omega$ such that

$$S_\omega(u) = d(\omega) := \inf_{v \in N_\omega} S_\omega(v),$$

and, as showed in Appendix A, they are bound states of the δ -NLS.

Remark 1.9. We use the notation

$$Q_\omega(v) := Q(v) + \omega \|v\|_2^2, \quad (18)$$

so that

$$S_\omega(v) = \frac{1}{2} Q_\omega(v) - \frac{1}{p} \|v\|_p^p \quad \text{and} \quad I_\omega(v) = Q_\omega(v) - \|v\|_p^p. \quad (19)$$

The bridge between minimizers of the action and ground states is cast by the following Lemma, whose proof is an adaptation of what was established in [29] for the NLSE. We report it in Appendix B.

Lemma 1.10. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\mu > 0$. If u is a ground state of mass μ , then it is also a minimizer of the action at the frequency $\omega = \mu^{-1}(\|u\|_p^p - Q(u))$.*

We give the following result for the minimizers of the action functional.

Theorem 1.11 (δ -NLS action minimizers). *Let $p > 2$ and $\alpha \in \mathbb{R}$. Then,*

- (i) *a minimizer of the action (15) at frequency ω does exist if and only if $\omega > \omega_0 := -\ell_\alpha$, with ℓ_α defined in (8);*
- (ii) *if $u = \phi_\lambda + q\mathcal{G}_\lambda$ is a minimizer of the action (15) at frequency $\omega > \omega_0$, then:*
 - (a) *there does not exist $\lambda > 0$ such that ϕ_λ or $q\mathcal{G}_\lambda$ are identically zero,*
 - (b) *u is positive, radially symmetric, and decreasing along the radial direction, up to multiplication by a constant phase factor; in particular, ϕ_λ is nonnegative as $\lambda = \omega$, and positive as $\lambda > \omega$.*

First, we note that, in view of Lemma 1.10, point (ii) of Theorem 1.5 is a straightforward consequence of Theorem 1.11. Indeed, if there exists a ground state of mass μ , then by Lemma 1.10 and point (i) of Theorem 1.11 it is also a minimizer of the action at frequency $\omega > \omega_0$. Hence, the conclusion follows by point (ii) of Theorem 1.11.

Furthermore, we mention that in order to establish (ii)(b) we use an equivalent formulation of the problem of minimization of the action consisting in minimizing Q_ω on the functions in D with fixed L^p norm. More details on this point are given in Remark 5.8. The technique is variational and different from those sometimes used for the standard NLSE, such as the moving planes introduced by [34].

Theorem 1.11 establishes that minimizers of the action exist if and only if the frequency exceeds the energy of the unique linear ground state. This fact confirms that nonlinear ground states bifurcate from the linear ones. Moreover, while Theorem 1.5 and Lemma 1.10 allow deducing the existence of minimizers of the action only in the subcritical regime, Theorem 1.11 shows the existence of ground states and of action minimizers in the critical and supercritical regime too.

During the final draft of the present paper we got acquainted that the results of Theorem 1.11 have been proved independently and with a different strategy in a contemporary work ([32]).

Organization of the paper.

- Section 2 introduces some preliminary results that are useful throughout the paper; more precisely:
 - in Section 2.1 we recall some well-known features of the Green's function of $-\Delta + \lambda$,
 - in Section 2.2 we establish two extensions of the Gagliardo-Nirenberg inequality (Proposition 2.1),
 - in Section 2.3 we establish a rearrangement inequality for the L^p -norms of the sum of nonnegative functions (Proposition 2.3);
- Section 3 addresses the existence of ground states (Theorem 1.5–(i));
- Section 4 addresses the existence of action minimizers (Theorem 1.11–(i));
- Section 5 establishes the main features both of the δ -NLS ground states and of the action minimizers (Theorem 1.5 –(ii)/Theorem 1.11–(ii)).

Acknowledgements. The authors wish to thank Simone Dovetta for helpful discussions. Moreover, the work has been partially supported by the MIUR project “Dipartimenti di Eccellenza 2018-2022” (CUP E11G18000350001) and by the INdAM GNAMPA project 2020 “Modelli differenziali alle derivate parziali per fenomeni di interazione”.

2. PRELIMINARY RESULTS

In this section we collect some preliminary results, that will be exploited in the proofs of Theorem 1.5 and Theorem 1.11.

2.1. Further properties of the Green's function. First, we recall that $\mathcal{G}_\lambda \in L^2(\mathbb{R}^2) \setminus H^1(\mathbb{R}^2)$, is positive, radially symmetric, decreasing along the radial direction, has exponential decay at infinity, and is smooth up to the origin, where it satisfies

$$\mathcal{G}_\lambda(\mathbf{x}) = -\frac{1}{2\pi} \log \left(\frac{\sqrt{\lambda}|\mathbf{x}|}{2} \right) + \frac{\gamma}{2\pi} + o(1), \quad \text{as } |\mathbf{x}| \rightarrow \mathbf{0}$$

(see [1, Sec. 9.6] and [23, Eq. (1.5)]). Thus \mathcal{G}_λ belongs to $L^p(\mathbb{R}^2)$, $2 \leq p < \infty$. Moreover,

$$\|\mathcal{G}_\lambda\|_2^2 = \frac{1}{4\pi\lambda} \quad \text{and} \quad \|\mathcal{G}_\lambda\|_p^p = \frac{\|\mathcal{G}_1\|_p^p}{\lambda}. \quad (20)$$

As shown in Section 1.1, $\mathcal{G}_\lambda - \mathcal{G}_\nu \in H^2(\mathbb{R}^2)$. Furthermore, by direct computation,

$$\|\mathcal{G}_\lambda - \mathcal{G}_\nu\|_2^2 = \frac{1}{4\pi} \left(\frac{1}{\lambda} + \frac{1}{\nu} + \frac{2 \log(\nu/\lambda)}{\lambda - \nu} \right), \quad (21)$$

$$\|\nabla(\mathcal{G}_\lambda - \mathcal{G}_\nu)\|_2^2 = \frac{1}{4\pi} \left(\frac{(\lambda + \nu) \log(\nu/\lambda)}{\lambda - \nu} - 2 \right). \quad (22)$$

Finally, we note that if $\nu < \lambda$, then

$$\mathcal{G}_\lambda(\mathbf{x}) = \frac{K_0(\sqrt{\lambda}\mathbf{x})}{2\pi} = \mathcal{G}_\nu \left(\sqrt{\frac{\lambda}{\nu}} \mathbf{x} \right) < \mathcal{G}_\nu(\mathbf{x}), \quad \forall \mathbf{x} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}. \quad (23)$$

2.2. Extensions of the Gagliardo-Nirenberg inequality. We need a generalization of Gagliardo-Nirenberg inequality to the energy space D .

Let us recall the standard two-dimensional Gagliardo-Nirenberg inequality ([28, Theorem 1.3.7]): there exists $C_p > 0$ such that

$$\|v\|_p^p \leq C_p \|\nabla v\|_2^{p-2} \|v\|_2^2, \quad \forall v \in H^1(\mathbb{R}^2). \quad (24)$$

First, notice that the set of function with $q \neq 0$ can be written as

$$D \setminus H^1(\mathbb{R}^2) = \left\{ u \in L^2(\mathbb{R}^2) : \exists q \in \mathbb{C} \setminus \{0\} \text{ s.t. } u - q \mathcal{G}_{\frac{|q|^2}{\|u\|_2^2}} =: \phi \in H^1(\mathbb{R}^2) \right\}. \quad (25)$$

In other words, functions in $D \setminus H^1(\mathbb{R}^2)$ admit the decomposition with $\lambda = \frac{|q|^2}{\|u\|_2^2}$. We can now state the following

Proposition 2.1 (Extended Gagliardo-Nirenberg inequalities). *For every $p > 2$, there exists $K_p > 0$ such that*

$$\|v\|_p^p \leq K_p \left(\|\nabla \phi_\lambda\|_2^{p-2} \|\phi_\lambda\|_2^2 + \frac{|q|^p}{\lambda} \right), \quad \forall v = \phi_\lambda + q \mathcal{G}_\lambda \in D, \quad \forall \lambda > 0. \quad (26)$$

Moreover, there exists $M_p > 0$

$$\|v\|_p^p \leq M_p \left(\|\nabla \phi\|_2^{p-2} + |q|^{p-2} \right) \|v\|_2^2, \quad \forall v = \phi + q \mathcal{G}_{\frac{|q|^2}{\|v\|_2^2}} \in D \setminus H^1(\mathbb{R}^2). \quad (27)$$

Proof. If we fix $v = \phi_\lambda + q \mathcal{G}_\lambda \in D$, for some $\lambda > 0$, then (24) and (20) yield

$$\|v\|_p^p = \|\phi + q \mathcal{G}_\lambda\|_p^p \leq 2^{p-1} (\|\phi\|_p^p + |q|^p \|\mathcal{G}_\lambda\|_p^p) \leq K_p \left(\|\nabla \phi\|_2^{p-2} \|\phi\|_2^2 + \frac{|q|^p}{\lambda} \right),$$

that is (26).

If we suppose, in addition, that $q \neq 0$ and set $\lambda = \lambda_q := \frac{|q|^2}{\|v\|_2^2}$, then by (20), (26) and the triangle inequality there results

$$\|v\|_p^p \leq M_p \left(\|\nabla \phi\|_2^{p-2} \|v\|_2^2 + \|\nabla \phi\|_2^{p-2} \frac{|q|^2}{\lambda_q} + \frac{|q|^p}{\lambda_q} \right) \leq M_p \left(\|\nabla \phi\|_2^{p-2} + |q|^{p-2} \right) \|v\|_2^2$$

possibly redefining M_p , which concludes the proof. \square

Remark 2.2. Note that, whenever $q = 0$, i.e. $v \in H^1(\mathbb{R}^2)$, (26) reduces to (24).

2.3. A rearrangement inequality in L^p -spaces. Let us start by recalling the definition of *radially symmetric nonincreasing rearrangement* of a function in \mathbb{R}^2 and its main features (see e.g. [40, Chapter 3]). First, given a measurable $A \subset \mathbb{R}^2$ with finite Lebesgue measure, we denote by A^* the open ball centred at zero with Lebesgue measure equal to $|A|$, that is

$$A^* := \{x \in \mathbb{R}^2 : \pi|\mathbf{x}|^2 < |A|\}.$$

Now, let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a nonnegative measurable function *vanishing at infinity*, i.e. $|\{f > t\}| := |\{\mathbf{x} \in \mathbb{R}^2 : f(\mathbf{x}) > t\}| < +\infty$, for every $t > 0$. We call the *radially symmetric nonincreasing rearrangement* of f the function $f^* : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f^*(\mathbf{x}) = \int_0^\infty \mathbb{1}_{\{f > t\}^*}(\mathbf{x}) dt, \quad (28)$$

with $\mathbb{1}_{\{f > t\}^*}$ the characteristic function of $\{f > t\}^*$. Definition (28) clearly implies

$$|\{f > t\}| = |\{f^* > t\}| \quad \text{and} \quad \{f > t\}^* = \{f^* > t\} \quad (29)$$

and

$$\mathbb{1}_A^* \equiv \mathbb{1}_{A^*}, \quad \text{for every measurable } A \subset \mathbb{R}^2, \quad |A| < +\infty. \quad (30)$$

One can also check that

$$(\Phi \circ f)^* \equiv \Phi \circ f^*, \quad \text{for every nondecreasing } \Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+ \quad (31)$$

and that (29) yields

$$\|f^*\|_p = \|f\|_p, \quad \forall f \in L^p(\mathbb{R}^2), f \geq 0, \quad \forall p \geq 1. \quad (32)$$

Another well known property of rearrangements is the *Hardy-Littlewood inequality*, which states that, given two nonnegative measurable functions $f, g : \mathbb{R}^2 \rightarrow \mathbb{R}$ vanishing at infinity, there results

$$\int_{\mathbb{R}^2} f(\mathbf{x})g(\mathbf{x}) d\mathbf{x} \leq \int_{\mathbb{R}^2} f^*(\mathbf{x})g^*(\mathbf{x}) d\mathbf{x} \quad (33)$$

and, if f is radially symmetric and decreasing, then the equality holds in (33) if and only if $g = g^*$ a.e. on \mathbb{R}^2 .

The result we need in the sequel compares $\|f + g\|_p$ and $\|f^* + g^*\|_p$. It is stated in the next Proposition, whose proof is an adaptation of that of [40, Theorems 3.4 and 3.5].

Proposition 2.3 (Rearrangement inequality). *For every pair of nonnegative functions $f, g \in L^p(\mathbb{R}^2)$, with $p > 1$, there results*

$$\int_{\mathbb{R}^2} |f + g|^p d\mathbf{x} \leq \int_{\mathbb{R}^2} |f^* + g^*|^p d\mathbf{x}. \quad (34)$$

Moreover, if f is radially symmetric and strictly decreasing along the radial direction, then the equality in (34) implies that $g = g^$ a.e. on \mathbb{R}^2 .*

Proof. First, we introduce the function

$$J_+(t) := \begin{cases} J(t) & \text{if } t \geq 0, \\ 0 & \text{if } t < 0, \end{cases} \quad \text{with} \quad J(t) := |t|^p.$$

It is straightforward that J_+ is of class C^1 and that $J_+(t) = \int_0^t J'_+(s) ds$, where J'_+ denotes the derivative of J_+ . We also note that J'_+ is nonnegative and nondecreasing in \mathbb{R} and, in particular, is positive and increasing in \mathbb{R}^+ . Therefore,

$$|f(\mathbf{x}) + g(\mathbf{x})|^p = J_+(f(\mathbf{x}) + g(\mathbf{x})) = \int_{-f(\mathbf{x})}^{g(\mathbf{x})} J'_+(f(\mathbf{x}) + s) ds = \int_{-\infty}^{+\infty} J'_+(f(\mathbf{x}) + s) \mathbb{1}_{\{g > s\}}(\mathbf{x}) ds,$$

whence, integrating over \mathbb{R}^2 and using Tonelli's theorem, we get

$$\begin{aligned} & \int_{\mathbb{R}^2} |f(\mathbf{x}) + g(\mathbf{x})|^p d\mathbf{x} \\ &= \int_{\mathbb{R}^2} J_+(f(\mathbf{x}) + g(\mathbf{x})) d\mathbf{x} = \int_{-\infty}^{+\infty} \left(\int_{\mathbb{R}^2} J'_+(f(\mathbf{x}) + s) \mathbb{1}_{\{g > s\}}(\mathbf{x}) d\mathbf{x} \right) ds \\ &= \underbrace{\int_0^{+\infty} \int_{\mathbb{R}^2} J'_+(f(\mathbf{x}) - s) d\mathbf{x} ds}_{=: I_1} + \underbrace{\int_0^{+\infty} \int_{\mathbb{R}^2} J'_+(f(\mathbf{x}) + s) \mathbb{1}_{\{g > s\}}(\mathbf{x}) d\mathbf{x} ds}_{=: I_2} \end{aligned} \quad (35)$$

where we used the fact that $\mathbb{1}_{\{g > -s\}} \equiv 1$ for every $s > 0$. Now, combining (29) and (31) with $\Phi(\cdot) = J'_+(\cdot - s)$, one sees that

$$I_1 = \int_0^{+\infty} \int_{\mathbb{R}^2} J'_+(f^*(\mathbf{x}) - s) d\mathbf{x} ds. \quad (36)$$

On the other hand, combining (29), (30), (31) with $\Phi(\cdot) = J'_+(\cdot + s) - J'_+(s)$ and (33), one sees that

$$\begin{aligned} & \int_{\mathbb{R}^2} J'_+(f(\mathbf{x}) + s) \mathbb{1}_{\{g > s\}}(\mathbf{x}) d\mathbf{x} \\ &= \int_{\mathbb{R}^2} (J'_+(f(\mathbf{x}) + s) - J'_+(s)) \mathbb{1}_{\{g > s\}}(\mathbf{x}) d\mathbf{x} + J'_+(s) |\{g > s\}| \\ &\leq \int_{\mathbb{R}^2} (J'_+(f(\mathbf{x}) + s) - J'_+(s))^* \mathbb{1}_{\{g > s\}}(\mathbf{x}) d\mathbf{x} + J'_+(s) |\{g^* > s\}| \\ &= \int_{\mathbb{R}^2} (J'_+(f^*(\mathbf{x}) + s) - J'_+(s)) \mathbb{1}_{\{g^* > s\}}(\mathbf{x}) d\mathbf{x} + J'_+(s) |\{g^* > s\}| \\ &= \int_{\mathbb{R}^2} J'_+(f^*(\mathbf{x}) + s) \mathbb{1}_{\{g^* > s\}}(\mathbf{x}) d\mathbf{x}, \end{aligned} \quad (37)$$

so that

$$I_2 \leq \int_0^{+\infty} \int_{\mathbb{R}^2} J'_+(f^*(\mathbf{x}) + s) \mathbb{1}_{\{g^* > s\}}(\mathbf{x}) d\mathbf{x} ds. \quad (38)$$

Hence, in view of (35), (36) and (38) one easily finds that (34) is satisfied.

It is left to prove that, if f is radially symmetric decreasing and the equality is fulfilled in (34), then $g = g^*$ a.e. on \mathbb{R}^2 . To this aim, fix f radially symmetric and decreasing and assume that the equality in (34) holds. Then, one can check that $f = f^*$ a.e. on \mathbb{R}^2 and that, by (37),

$$\int_{\mathbb{R}^2} J'_+(f(\mathbf{x}) + s) \mathbb{1}_{\{g > s\}}(\mathbf{x}) d\mathbf{x} = \int_{\mathbb{R}^2} J'_+(f(\mathbf{x}) + s) \mathbb{1}_{\{g^* > s\}}(\mathbf{x}) d\mathbf{x}, \quad \text{for a.e. } s \geq 0. \quad (39)$$

Since J'_+ is increasing on \mathbb{R}^+ and f is radially symmetric decreasing, $J'_+(f(\cdot) + s)$ is radially symmetric decreasing too. Hence there exists a continuous bijection $r : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\{\mathbf{x} \in \mathbb{R}^2 : J'_+(f(\mathbf{x}) + s) - J'_+(s) > t\} = B_{r(t)}(\mathbf{0})$, namely the centered ball of radius $r(t)$. In addition, by dominated converge, the function

$$F_C(t) := \int_{\mathbb{R}^2} \mathbb{1}_{B_{r(t)}(\mathbf{0})}(\mathbf{x}) \mathbb{1}_C(\mathbf{x}) d\mathbf{x} = |B_{r(t)}(\mathbf{0}) \cap C|$$

is continuous on \mathbb{R}^+ for any measurable $C \subset \mathbb{R}^2$ fixed.

Now, fix $s > 0$ such that (39) holds and set $C = \{\mathbf{x} \in \mathbb{R}^2 : g(\mathbf{x}) > s\}$. Arguing as before, one can find that $F_C(t) \leq F_{C^*}(t)$. From (37) and (39), using the layer-cake representation, one obtains that $\int_0^\infty F_C(t) dt = \int_0^\infty F_{C^*}(t) dt$ and, hence, $F_C(t) = F_{C^*}(t)$ for every $t > 0$. As C^* is a centered ball too, this implies that for every $r > 0$ either $C, C^* \subset B_r(\mathbf{0})$ or $C, C^* \supset B_r(\mathbf{0})$ up to sets of zero Lebesgue measure, so that $C = C^*$. Finally, as this is

valid for every $s > 0$, using again (29) and the layer-cake representation, there results that $g = g^*$ a.e. on \mathbb{R}^2 . \square

Remark 2.4. Up to some minor modifications, in order to prove the first part of Proposition 2.3 it suffices the simple convexity of J , the strict convexity being necessary for the sole second part. Hence, (34) holds also for $p = 1$.

Before concluding the section, we also mention another well known result on rearrangements that will be used in the sequel: if $f \in H^1(\mathbb{R}^2)$, then $f^* \in H^1(\mathbb{R}^2)$ and in particular

$$\|\nabla f^*\|_2 \leq \|\nabla f\|_2. \quad (40)$$

Equation (40) is usually called the *Pólya-Szegő inequality*.

Remark 2.5. All results in Section 2.3 are valid in every \mathbb{R}^d , with $d \geq 2$.

3. GROUND STATES EXISTENCE: PROOF OF THEOREM 1.5–(I)

In this section, we prove point (i) of Theorem 1.5, that is the existence of δ -NLS ground states of mass μ for every $\mu > 0$.

To this aim, the first step is to establish boundedness from below of $E|_{D_\mu}$ in the $L^2(\mathbb{R}^2)$ -subcritical case. Preliminarily, it is convenient to write the functional E as

$$E(u) = \begin{cases} \frac{1}{2} \|\nabla \phi\|_2^2 + \frac{|q|^2 \|\phi\|_2^2}{2\|u\|_2^2} + \left(\alpha - 1 + \frac{\log\left(\frac{|q|}{2\|u\|_2}\right) + \gamma}{2\pi} \right) \frac{|q|^2}{2} - \frac{\|u\|_p^p}{p}, & \text{if } u \in D \setminus H^1(\mathbb{R}^2), \\ \frac{1}{2} \|\nabla u\|_2^2 - \frac{1}{p} \|u\|_p^p, & \text{if } u \in H^1(\mathbb{R}^2), \end{cases} \quad (41)$$

where we use the decomposition $u = \phi + q\mathcal{G}_{\frac{|q|^2}{\|u\|_2^2}}$ introduced in (25), for every $u \in D \setminus H^1(\mathbb{R}^2)$.

Proposition 3.1. *Let $p \in (2, 4)$ and $\alpha \in \mathbb{R}$. Then $\mathcal{E}(\mu) > -\infty$, for every $\mu > 0$.*

Proof. Let $u \in D_\mu$. Assume, first, that $u \in H_\mu^1(\mathbb{R}^2)$. Therefore, (24) entails

$$E(u) \geq \frac{1}{2} \|\nabla u\|_2^2 - \frac{C_p}{p} \|\nabla u\|_2^{p-2} \mu,$$

so that E is bounded from below on $H_\mu^1(\mathbb{R}^2)$ since $2 < p < 4$.

Then, assume that $u \in D_\mu \setminus H_\mu^1(\mathbb{R}^2)$. By (27)

$$E(u) \geq \left(\frac{1}{2} \|\nabla \phi\|_2^2 - \frac{M_p}{p} \|\nabla \phi\|_2^{p-2} \mu \right) + \frac{|q|^2 \|\phi\|_2^2}{2\mu} + \left[\left(\alpha - 1 + \frac{\log\left(\frac{|q|}{2\sqrt{\mu}}\right) + \gamma}{2\pi} \right) \frac{|q|^2}{2} - \frac{M_p}{p} |q|^{p-2} \mu \right], \quad (42)$$

and here again E is bounded from below in $D_\mu \setminus H_\mu^1(\mathbb{R}^2)$ since $2 < p < 4$ (note that the $\log(|q|)|q|^2$ term balances the negatively diverging $|q|^2$ term). Summing up, E is lower bounded on the whole D_μ . \square

Further than boundedness from below, it is also useful to establish a comparison between the δ -NLS energy infimum and the NLS energy infimum.

Proposition 3.2. *Let $p \in (2, 4)$ and $\alpha \in \mathbb{R}$. Then,*

$$\mathcal{E}(\mu) < \mathcal{E}^0(\mu) < 0, \quad \forall \mu > 0. \quad (43)$$

In order to prove this, we preliminarily recall without proof a well known result about NLS ground states (see [42] for a proof of the existence part, while the proof of the properties satisfied by the ground states is a consequence of [34]).

Proposition 3.3. *Let $p \in (2, 4)$ and $\mu > 0$. Then, there exists an NLS ground state of mass μ , i.e. $u \in H_\mu^1(\mathbb{R}^2)$ such that $E^0(u) = \mathcal{E}^0(\mu)$. Moreover, such minimizer is unique, positive and radially symmetric decreasing, up to multiplication by a constant phase and translation.*

The positive minimizer of the two-dimensional standard NLS functional at mass μ is usually called a two-dimensional *soliton* and we denote it S_μ .

Proof of Proposition 3.2. Fix $\mu > 0$ and let u be the unique NLS ground state of mass μ mentioned in Proposition 3.3. First, note that, as u is positive, it cannot be a δ -NLS ground state of mass μ . Indeed, if u is a δ -NLS ground state, then u has to satisfy (10) and, in particular, $\phi_\lambda(\mathbf{0}) = (\alpha + \theta_\lambda)q$. However, as mentioned in Section 2.2, $u \in H^1(\mathbb{R}^2)$ implies $q = 0$, so that $u \equiv \phi_\lambda$ and $\phi_\lambda(\mathbf{0}) = 0$. Hence, $u(\mathbf{0}) = 0$, which contradicts its positivity. Summing up, u is not a δ -NLS ground state mass μ and, thus, there exists $v \in D_\mu$ such that $E(v) < E(u) = \mathcal{E}^0(\mu)$, which proves the left inequality in (43).

Concerning the right inequality, fix again $\mu > 0$, and consider $v \in H_\mu^1(\mathbb{R}^2)$. Now, using the mass-preserving transformation

$$v_\sigma(x) = \sigma v(\sigma x),$$

There results

$$E^0(v_\sigma) = \frac{\sigma^2}{2} \|\nabla v\|_2^2 - \frac{\sigma^{p-2}}{p} \|v\|_p^p.$$

However, as $p \in (2, 4)$, this immediately entails that $\mathcal{E}^0(\mu) \leq E(u_\sigma) < 0$, for every $\sigma \ll 1$, which completes the proof. \square

The second step of the proof of point (i) in Theorem 1.5 consists in a characterization of the δ -NLS energy minimizing sequences of mass μ , i.e. sequences

$$(u_n)_n \subset D_\mu \quad \text{such that} \quad E(u_n) \rightarrow \mathcal{E}(\mu), \quad \text{as } n \rightarrow +\infty.$$

This is provided by the next two lemmas.

Lemma 3.4. *Let $p \in (2, 4)$, $\alpha \in \mathbb{R}$ and $\mu > 0$. If $u_n = \phi_{\lambda,n} + q_n \mathcal{G}_\lambda$ is a minimizing sequence for the δ -NLS energy, then there exists $\bar{n} \in \mathbb{N}$ and a constant $C > 0$, such that $|q_n| > C$ for every $n \geq \bar{n}$.*

Proof. We proceed by contradiction. Suppose that there exists a subsequence of q_n , that we do not rename, such that $q_n \rightarrow 0$. Then, $\|\phi_{\lambda,n}\|_2^2$ is bounded since it converges to μ . Moreover, applying Gagliardo-Nirenberg inequality (24) to definition (13) one obtains

$$E(u_n) \geq \frac{1}{2} \|\nabla \phi_{\lambda,n}\|_2^2 + \frac{\lambda}{2} (\|\phi_{\lambda,n}\|_2^2 - \mu) + \frac{(\alpha + \theta_\lambda)}{2} |q_n|^2 - \frac{C_p}{p} \|\nabla \phi_{\lambda,n}\|_2^{p-2} \|\phi_{\lambda,n}\|_2^2$$

that, since $E(u_n)$ is bounded from above and $p < 4$, guarantees the boundedness of $\|\nabla \phi_{\lambda,n}\|_2$.

We introduce the sequence $\xi_n = \frac{\sqrt{\mu}}{\|\phi_{\lambda,n}\|_2}$, such that $\|\xi_n\|_2^2 = \mu$ and $\|\nabla \xi_n\|_2^2 = \frac{\mu}{\|\phi_{\lambda,n}\|_2^2} \|\nabla \phi_{\lambda,n}\|_2^2$ is bounded. Then, using $q_n \rightarrow 0$, $\phi_{\lambda,n} - u_n \rightarrow 0$ strongly in every space $L^p(\mathbb{R}^2)$ with $2 \leq p < \infty$, and the listed properties of ξ_n , one obtains

$$\begin{aligned} E(u_n) &= E^0(\phi_{\lambda,n}) + o(1) = E^0(\xi_n) + o(1) \\ &\geq E^0(S_\mu) + o(1), \quad n \rightarrow \infty, \end{aligned}$$

where S_μ is a ground state for E^0 at mass μ . So, passing to the limit,

$$\mathcal{E}(\mu) \geq \mathcal{E}^0(\mu),$$

that contradicts Proposition 3.2 and then q_n cannot converge to zero. This conclusion holds for every subsequence of a minimizing sequence for E , therefore limit points of the complex sequence q_n must be separated from zero, and the proof is complete. \square

Lemma 3.5. *Let $p \in (2, 4)$, $\alpha \in \mathbb{R}$ and $\mu > 0$. Let also $(u_n)_n$ be a δ -NLS energy minimizing sequence of mass μ . Then, it is bounded in $L^r(\mathbb{R}^2)$, for every $r \geq 2$, and there exists $u \in D \setminus H^1(\mathbb{R}^2)$ such that, up to subsequences,*

- $u_n \rightharpoonup u$ in $L^2(\mathbb{R}^2)$,
- $u_n \rightarrow u$ a.e. in \mathbb{R}^2 ,

as $n \rightarrow +\infty$. In particular, if one fixes $\lambda > 0$ and the decomposition $u_n = \phi_{n,\lambda} + q_n \mathcal{G}_\lambda$, then $(\phi_{n,\lambda})_n$ and $(q_n)_n$ are bounded in $H^1(\mathbb{R}^2)$ and \mathbb{C} , respectively, and there exist $\phi_\lambda \in H^1(\mathbb{R}^2)$ and $q \in \mathbb{C} \setminus \{0\}$ such that $u = \phi_\lambda + q \mathcal{G}_\lambda$ and, up to subsequences,

- $\phi_{n,\lambda} \rightharpoonup \phi_\lambda$ in $L^2(\mathbb{R}^2)$,
- $\nabla \phi_{n,\lambda} \rightharpoonup \nabla \phi_\lambda$ in $L^2(\mathbb{R}^2)$,
- $q_n \rightarrow q$ in \mathbb{C} ,

as $n \rightarrow +\infty$.

Proof. Let $(u_n)_n$ be a δ -NLS energy minimizing sequence of mass μ . By Banach-Alaoglu Theorem, $u_n \rightharpoonup u$ in $L^2(\mathbb{R}^2)$ up to subsequences. Moreover, owing to Lemma 3.4 we can suppose without loss of generality that for every n the charge q_n satisfies $|q_n| > C > 0$, then we can rely on the decomposition introduced in (25) and used in (27), namely $u_n = \phi_n + q_n \mathcal{G}_{\nu_n}$ with $\nu_n := \frac{|q_n|^2}{\|u_n\|_2^2}$. This decomposition guarantees

$$\|\phi_n\|_2^2 \leq C \mu$$

for every n , so that the sequence ϕ_n is bounded in $L^2(\mathbb{R}^2)$.

Using (41) and (42), we have

$$E(u_n) \geq \left(\frac{1}{2} \|\nabla \phi_n\|_2^2 - \frac{M_p}{p} \mu \|\nabla \phi_n\|_2^{p-2} \right) + \left(\alpha - 1 + \frac{\log\left(\frac{|q_n|}{2\sqrt{\mu}}\right)}{2\pi} \right) \frac{|q_n|^2}{2} - \frac{M_p}{p} \mu |q_n|^{p-2}, \quad (44)$$

for a suitable $M_p > 0$. First, we note by (44) that $(\nabla \phi_n)_n$ is bounded in $L^2(\mathbb{R}^2)$ and $(q_n)_n$ is bounded in \mathbb{C} , so that, up to subsequences, $q_n \rightarrow q$ and $q \neq 0$ since $|q_n| > C > 0$.

Fix $\lambda \geq \frac{C_2}{\mu}$ with $C_2 = 1 + \sup_n |q_n|$ and consider the decomposition of each u_n according to λ , that is $u_n = \phi_{n,\lambda} + q_n \mathcal{G}_\lambda$ with $\phi_{\lambda,n} := \phi_n + q_n (\mathcal{G}_{\nu_n} - \mathcal{G}_\lambda)$. Exploiting (21) and (22) and the estimates on ϕ_n and q_n , one finds that there exists $M_1, M_2 > 0$ such that for every $n \geq \bar{n}$

$$\|\phi_{n,\lambda}\|_2^2 \leq 2 \left[\|\phi_n\|_2^2 + \frac{1}{4\pi} \left(\frac{|q_n|^2}{\lambda} + \mu + 2|q_n|^2 \frac{\log \lambda + \log(\mu) - 2 \log(|q_n|)}{\nu_n - \lambda} \right) \right] \leq M_1$$

and

$$\|\nabla \phi_{n,\lambda}\|_2^2 \leq 2 \left[\|\nabla \phi_n\|_2^2 + \frac{|q_n|^2}{4\pi} \left(\left(\lambda + \frac{|q_n|^2}{\mu} \right) \frac{\log \lambda + \log(\mu) - 2 \log(|q_n|)}{\lambda - \nu_n} - 2 \right) \right] \leq M_2.$$

Hence $(\phi_{n,\lambda})_n, (\nabla \phi_{n,\lambda})_n$ are bounded in $L^2(\mathbb{R}^2)$, which implies, via the Banach-Alaoglu theorem, that $\phi_{n,\lambda} \rightharpoonup \phi_\lambda, \nabla \phi_{n,\lambda} \rightharpoonup \nabla \phi_\lambda$ in $L^2(\mathbb{R}^2)$, up to subsequences, and that $u = \phi_\lambda + q \mathcal{G}_\lambda$. Furthermore, by Rellich-Kondrakov theorem, $\phi_{n,\lambda} \rightarrow \phi_\lambda$ in $L_{loc}^r(\mathbb{R}^2)$, for every $r > 2$, so that $u_n \rightarrow u$ a.e. in \mathbb{R}^2 .

It is then left to prove that $(\phi_{n,\lambda})_n, (\nabla\phi_{n,\lambda})_n$ are bounded in $L^2(\mathbb{R}^2)$ also when the decomposition parameter is smaller than $\frac{C_2}{\mu}$. To this aim, let $0 < \tilde{\lambda} < \frac{C_2}{\mu}$. We can use the decomposition $u_n = \phi_{n,\tilde{\lambda}} + q_n\mathcal{G}_{\tilde{\lambda}}$, where $\phi_{n,\tilde{\lambda}} = \phi_{n,\lambda} + q_n(\mathcal{G}_\lambda - \mathcal{G}_{\tilde{\lambda}})$, with $\lambda \geq \frac{C_2}{\mu}$. However, arguing as before, one can see that $q_n(\mathcal{G}_\lambda - \mathcal{G}_{\tilde{\lambda}})$ is bounded in $H^1(\mathbb{R}^2)$, which concludes the proof. \square

Finally, we have all the tools to prove the existence part of Theorem 1.5

Proof of Theorem 1.5-(i). Let $(u_n)_n$ be a δ -NLS energy minimizing sequence of mass μ . Assume also, without loss of generality, that it is a subset of $D_\mu \setminus H^1(\mathbb{R}^2)$, so that we can write $u_n = \phi_{n,\lambda} + q_n\mathcal{G}_\lambda$, with $q_n \neq 0$ and $\lambda > 0$. As a consequence, all the results of Lemma 3.5 hold and all the following limits hold up to subsequences.

Set $m := \|u\|_2^2$. By weak lower semicontinuity of the $L^2(\mathbb{R}^2)$ -norm, $m \leq \mu$. Moreover, as $q \neq 0$, $m \neq 0$. Assume, then, by contradiction, that $0 < m < \mu$. Note that, since $u_n \rightharpoonup u$ in $L^2(\mathbb{R}^2)$, $\|u_n - u\|_2^2 = \mu - m + o(1)$, as $n \rightarrow +\infty$. On the one hand, since $p > 2$ and $\frac{\mu}{\|u_n - u\|_2^2} > 1$ for n sufficiently large, there results that

$$\begin{aligned} \mathcal{E}(\mu) &\leq E\left(\sqrt{\frac{\mu}{\|u_n - u\|_2^2}}(u_n - u)\right) = \frac{1}{2}\frac{\mu}{\|u_n - u\|_2^2}Q(u_n - u) - \frac{1}{p}\left(\frac{\mu}{\|u_n - u\|_2^2}\right)^{\frac{p}{2}}\|u_n - u\|_p^p \\ &< \frac{\mu}{\|u_n - u\|_2^2}E(u_n - u) \end{aligned}$$

and thus

$$\liminf_n E(u_n - u) \geq \frac{\mu - m}{\mu}\mathcal{E}(\mu). \quad (45)$$

On the other hand, a similar computation yields

$$\mathcal{E}(\mu) \leq E\left(\sqrt{\frac{\mu}{\|u\|_2^2}}u\right) < \frac{\mu}{\|u\|_2^2}E(u),$$

so that

$$E(u) > \frac{m}{\mu}\mathcal{E}(\mu). \quad (46)$$

In addition, we can also prove that

$$E(u_n) = E(u_n - u) + E(u) + o(1) \quad \text{as } n \rightarrow +\infty \quad (47)$$

Indeed, since, $u_n \rightharpoonup u$, $\phi_{n,\lambda} \rightharpoonup \phi_\lambda$, $\nabla\phi_{n,\lambda} \rightharpoonup \nabla\phi_\lambda$ in $L^2(\mathbb{R}^2)$ and $q_n \rightarrow q$, we have that

$$Q(u_n - u) = Q(u_n) - Q(u) + o(1), \quad n \rightarrow +\infty,$$

while $\|u_n\|_p^p \leq C$ and $u_n \rightarrow u$ a.e. on \mathbb{R}^2 , enable one to use the well known Brezis-Lieb lemma ([19]) in order to get

$$\|u_n\|_p^p = \|u_n - u\|_p^p + \|u\|_p^p + o(1), \quad n \rightarrow +\infty.$$

Combining (45), (46) and (47), one can see that

$$\mathcal{E}(\mu) = \liminf_n E(u_n) = \liminf_n E(u_n - u) + E(u) > \frac{\mu - m}{\mu}\mathcal{E}(\mu) + \frac{m}{\mu}\mathcal{E}(\mu) = \mathcal{E}(\mu),$$

which is a contradiction. Therefore, $m = \mu$, so that $u \in D_\mu$ and, in particular, $u_n \rightarrow u$ in $L^2(\mathbb{R}^2)$ and $\phi_{n,\lambda} \rightarrow \phi_\lambda$ in $L^2(\mathbb{R}^2)$.

It is, then, left to show that

$$E(u) \leq \liminf_n E(u_n) = \mathcal{E}(\mu). \quad (48)$$

However, by all the limits obtained before, it suffices to prove that $u_n \rightarrow u$ in $L^p(\mathbb{R}^2)$, in order to get (48). Now, from (26),

$$\|u_n - u\|_p^p \leq K_p \left(\|\nabla\phi_{n,\lambda} - \nabla\phi_\lambda\|_2^{p-2} \|\phi_{n,\lambda} - \phi_\lambda\|_2^2 + \frac{|q_n - q|^p}{\lambda} \right)$$

and then since $\|\nabla\phi_{n,\lambda} - \nabla\phi_\lambda\|_2$ is bounded, $\phi_{n,\lambda} \rightarrow \phi_\lambda$ in $L^2(\mathbb{R}^2)$ and $q_n \rightarrow q$ in \mathbb{C} , the claim is proved. \square

4. ACTION MINIMIZERS EXISTENCE: PROOF OF THEOREM 1.11 – (I)

The aim of this section is proving point (i) of Theorem 1.11, that is the existence/nonexistence of δ -NLS action minimizers at frequency ω .

Preliminarily, we recall that, in the standard case, NLS-action minimizers are those functions $u \in N_\omega^0$ such that $S_\omega^0(v) = d^0(\omega)$, with

$$\begin{aligned} d^0(\omega) &:= \inf_{v \in N_\omega^0} S_\omega^0(v), \\ S_\omega^0(v) &:= E^0(v) + \frac{\omega}{2} \|v\|_2^2, \\ N_\omega^0 &:= \{v \in H^1(\mathbb{R}^2) \setminus \{0\} : I_\omega^0(v) = 0\}, \quad I_\omega^0(v) := \langle S_\omega^{0'}(v), v \rangle. \end{aligned}$$

We also note that

$$S_\omega(v) = \tilde{S}(v) > 0, \quad \forall v \in N_\omega, \quad (49)$$

with S_ω and N_ω given by (15) and (16), respectively, and

$$\tilde{S}(v) := \frac{p-2}{2p} \|v\|_p^p.$$

Hence, combining with the fact that $S_\omega|_{H^1(\mathbb{R}^2)} = S_\omega^0$ and $N_\omega \cap H^1(\mathbb{R}^2) = N_\omega^0$, it is straightforward that

$$0 \leq d(\omega) \leq d^0(\omega), \quad \forall \omega \in \mathbb{R}. \quad (50)$$

In addition, since $d^0(\omega) = 0$, for every $\omega \leq 0$ (see, e.g., [29, Lemma 2.4 and Remark 2.5]), one immediately sees that $d(\omega) = 0$, for every $\omega \leq 0$, which entails that there cannot be any δ -NLS action minimizer at frequency ω whenever $\omega \leq 0$. In view of this we will focus throughout only on the case $\omega > 0$.

Now, the first step of our discussion is to detect for which $\omega > 0$ the two inequalities in (50) are strict. To this aim let us introduce the set

$$\widehat{N}_\omega := \{q\mathcal{G}_\lambda : \lambda > 0, q \in \mathbb{C} \setminus \{0\}, I_\omega(q\mathcal{G}_\lambda) = 0\},$$

which is the subset of N_ω containing those functions admitting a decomposition with the sole singular part for at least one value of $\lambda > 0$. The next two lemmas characterize the set \widehat{N}_ω on varying $\omega > 0$.

Lemma 4.1. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > 0$. Then, $q\mathcal{G}_\lambda \in \widehat{N}_\omega$ if and only if $\lambda > 0$ and $q \in \mathbb{C} \setminus \{0\}$ satisfy*

$$\frac{\omega - \lambda}{4\pi} + \lambda(\alpha + \theta_\lambda) > 0 \quad (51)$$

(with θ_λ defined by (5)) and

$$|q| = \frac{1}{K_p} \left[\frac{\omega - \lambda}{4\pi} + \lambda(\alpha + \theta_\lambda) \right]^{\frac{1}{p-2}}, \quad (52)$$

with $K_p = \|\mathcal{G}_1\|_p^{\frac{p}{p-2}}$.

Proof. Fix $\psi = q\mathcal{G}_\lambda$ with $q \neq 0$ and $\lambda > 0$. By (20), $I_\omega(\psi) = 0$ if and only if

$$\frac{\omega - \lambda}{4\pi\lambda} |q|^2 + (\alpha + \theta_\lambda) |q|^2 - \frac{K}{\lambda} |q|^p = 0,$$

with $K := \|\mathcal{G}_1\|_p^p$, which entails

$$|q|^{p-2} = \frac{1}{K} \left[\frac{\omega - \lambda}{4\pi} + \lambda(\alpha + \theta_\lambda) \right].$$

Since $|q|^{p-2} > 0$, (51) and (52) follow. \square

Lemma 4.2. *Let $p > 2$, $\alpha \in \mathbb{R}$, $\omega > 0$ and recall that $\omega_0 := -\ell_\alpha$, with ℓ_α defined by (8). Therefore:*

(i) *if $\omega \in (0, \omega_0)$, then there exists $\lambda_1(\omega) \in (0, \omega_0)$ and $\lambda_2(\omega) > \omega_0$ such that*

$$\widehat{N}_\omega = \{q\mathcal{G}_\lambda : \lambda \in (0, \lambda_1(\omega)) \cup (\lambda_2(\omega), +\infty), q \in \mathbb{C} \setminus \{0\} \text{ and satisfies (52)}\};$$

in particular, $\lambda_1(\omega)$ and $\lambda_2(\omega)$ are the sole solutions of the equation

$$\frac{\omega - \lambda}{4\pi} + \lambda(\alpha + \theta_\lambda) = 0;$$

(ii) *if $\omega = \omega_0$, then*

$$\widehat{N}_\omega = \{q\mathcal{G}_\lambda : \lambda > 0, \lambda \neq \omega_0, q \in \mathbb{C} \setminus \{0\} \text{ and satisfies (52)}\};$$

(iii) *if $\omega > \omega_0$, then*

$$\widehat{N}_\omega = \{q\mathcal{G}_\lambda : \lambda > 0, q \in \mathbb{C} \setminus \{0\} \text{ and satisfies (52)}\}$$

Proof. Let $\omega > 0$ and introduce the function

$$g(\lambda) := \frac{\omega - \lambda}{4\pi} + \lambda(\alpha + \theta_\lambda).$$

Recall that, in view of Lemma 4.1, $q\mathcal{G}_\lambda \in \widehat{N}_\omega$ if and only if $g(\lambda) > 0$ and q satisfies (52), namely $|q| = K_p^{-1} g^{\frac{1}{p-2}}(\lambda)$. Now, it is straightforward (by (5)) that

$$\lim_{\lambda \rightarrow 0^+} g(\lambda) = \frac{\omega}{4\pi} > 0, \quad \lim_{\lambda \rightarrow +\infty} g(\lambda) = +\infty$$

and

$$g'(\lambda) = \alpha + \theta_\lambda.$$

Hence, one can easily see that g is decreasing for $\lambda < \omega_0$ and increasing for $\lambda > \omega_0$, has a global minimizer at $\lambda = \omega_0$ and $g(\omega_0) = \frac{\omega - \omega_0}{4\pi}$. Therefore, if $\omega > \omega_0$, then condition (51) can be satisfied for every $\lambda > 0$. On the contrary, if $\omega = \omega_0$, then (51) can be satisfied provided that $\lambda > 0$ and $\lambda \neq \omega_0$. Finally, if $\omega < \omega_0$, then $g(\omega_0) < 0$ and this implies that there exist $\lambda_1(\omega)$, $\lambda_2(\omega) > 0$ such that (51) does not hold if and only if $\lambda \in [\lambda_1(\omega), \lambda_2(\omega)]$. Note that $\lambda_1(\omega)$ and $\lambda_2(\omega)$ are the only values of $\lambda > 0$ for which g vanishes. \square

After this characterization of the set \widehat{N}_ω , we can estimate the value of $d(\omega)$ for $\omega \in (0, \omega_0]$.

Proposition 4.3. *Let $p > 2$, $\alpha \in \mathbb{R}$. Then, $d(\omega) = 0$, for every $\omega \in (0, \omega_0]$.*

Proof. Let us discuss separately the cases $\omega \in (0, \omega_0)$ and $\omega = \omega_0$. If $\omega \in (0, \omega_0)$, then (in view of Lemma 4.2) one can check that

$$\lim_{\substack{\lambda \rightarrow \lambda_1(\omega)^-, \\ q\mathcal{G}_\lambda \in N_\omega}} |q| = \lim_{\lambda \rightarrow \lambda_1(\omega)^-} \frac{1}{K_p} \left[\frac{\omega - \lambda}{4\pi} + \lambda(\alpha + \theta_\lambda) \right]^{\frac{1}{p-2}} = 0.$$

Hence, recalling and (49) and (20),

$$\begin{aligned} 0 \leq d(\omega) &\leq \inf_{q\mathcal{G}_\lambda \in N_\omega} S_\omega(q\mathcal{G}_\lambda) \leq \lim_{\substack{\lambda \rightarrow \lambda_1(\omega)^-, \\ q\mathcal{G}_\lambda \in N_\omega}} S_\omega(q\mathcal{G}_\lambda) \\ &= \lim_{\substack{\lambda \rightarrow \lambda_1(\omega)^-, \\ q\mathcal{G}_\lambda \in N_\omega}} \tilde{S}(q\mathcal{G}_\lambda) = \lim_{\substack{\lambda \rightarrow \lambda_1(\omega)^-, \\ q\mathcal{G}_\lambda \in N_\omega}} \|\mathcal{G}_1\|_p^p \frac{|q|^p}{\lambda} = 0. \end{aligned}$$

If, on the contrary, $\omega = \omega_0$, then one obtains the same result, just arguing as before and replacing the limits for $\lambda \rightarrow \lambda_1(\omega)^-$ with the limits for $\lambda \rightarrow \omega_0$. \square

This result has an immediate consequence on the existence of the δ -NLS action minimizers below ω_0 .

Corollary 4.4. *Let $p > 2$, $\alpha \in \mathbb{R}$. Then, there exists no δ -NLS action minimizer at frequency ω , for every $\omega \in (0, \omega_0]$.*

Proof. The claim follows by Proposition 4.3 and (49). \square

On the other hand, in order to discuss the behavior of $d(\omega)$ when $\omega > \omega_0$, it is preliminarily necessary to further investigate the relation between S_ω and \tilde{S} .

Lemma 4.5. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Then*

$$d(\omega) = \inf_{v \in \tilde{N}_\omega} \tilde{S}(v), \quad (53)$$

with

$$\tilde{N}_\omega := \{v \in D \setminus \{0\} : I_\omega(v) \leq 0\}$$

(and I_ω defined by (17)). Moreover, for any function $u \in D \setminus \{0\}$,

$$\begin{cases} \tilde{S}(u) = d(\omega) \\ I_\omega(u) \leq 0 \end{cases} \iff \begin{cases} S_\omega(u) = d(\omega) \\ I_\omega(u) = 0. \end{cases} \quad (54)$$

Remark 4.6. In view of this lemma, searching for δ -NLS action minimizers is equivalent to searching for

$$u \in \tilde{N}_\omega \quad \text{such that} \quad \tilde{S}(u) = \inf_{v \in \tilde{N}_\omega} \tilde{S}(v) = d(\omega).$$

Proof of Lemma 4.5. We divide in proof in two parts.

Part (i): proof of (53). On the one hand, if $u \in N_\omega$, then $S_\omega(u) = \tilde{S}(u)$, so that

$$\inf_{v \in \tilde{N}_\omega} \tilde{S}(v) \leq d(\omega),$$

as $\tilde{N}_\omega \supset N_\omega$. On the other hand, fix $u \in D \setminus \{0\}$ such that $I_\omega(u) < 0$ (i.e. $u \in \tilde{N}_\omega \setminus N_\omega$). Now, for any fixed $\beta > 0$

$$I_\omega(\beta u) = \beta^2 Q_\omega(u) - \beta^p \|u\|_p^p,$$

(see (18) for the definition of Q_ω), and thus $I_\omega(\beta u) = 0$ (i.e. $u \in N_\omega$) if and only

$$\beta = \beta(u) := \left(\frac{Q_\omega(u)}{\|u\|_p^p} \right)^{\frac{1}{p-2}}$$

(where we also used that $Q_\omega(u) > 0$, for every $u \in D \setminus \{0\}$, whenever $\omega > \omega_0$). Moreover, since $I_\omega(u) < 0$, $\beta(u) < 1$ and hence

$$S_\omega(\beta(u)u) = \tilde{S}(\beta(u)u) = \beta(u)^p \tilde{S}(u) < \tilde{S}(u).$$

As a consequence

$$d(\omega) \leq \inf_{v \in \tilde{N}_\omega} \tilde{S}(v),$$

which completes the proof.

Part (ii): proof of (54). If $u \in N_\omega$ and $S_\omega(u) = d(\omega)$, then clearly $u \in \tilde{N}_\omega$ and (by (49)) $\tilde{S}(u) = d(\omega)$. On the contrary, assume by contradiction that $u \in \tilde{N}_\omega \setminus N_\omega$. If $\tilde{S}(u) = d(\omega)$, then, arguing as before, one obtains that $\beta(u)u \in N_\omega$ and

$$S_\omega(\beta(u)u) < d(\omega),$$

which is impossible. Hence, if $u \in \tilde{N}_\omega$ and $\tilde{S}(u) = d(\omega)$, then $u \in N_\omega$ and $S_\omega(u) = d(\omega)$. \square

We can now prove that the left inequality of (50) is strict.

Proposition 4.7. *Let $p > 2$, $\alpha \in \mathbb{R}$. Then, $d(\omega) > 0$, for every $\omega > \omega_0$.*

Proof. First, let $u \in \tilde{N}_\omega \cap H^1(\mathbb{R}^2)$. By Sobolev inequality, there exists $C_p > 0$, depending only on p , such that

$$0 \geq I_\omega(u) = \|\nabla u\|_2^2 + \omega \|u\|_2^2 - \|u\|_p^p \geq C_p \|u\|_p^2 + \omega \|u\|_2^2 - \|u\|_p^p \geq C_p \|u\|_p^2 - \|u\|_p^p.$$

Hence, $\|u\|_p^{p-2} \geq C_p$ and so

$$\tilde{S}(u) \geq \frac{p-2}{2p} C_p^{\frac{p}{p-2}},$$

whence

$$\inf_{v \in \tilde{N}_\omega \cap H^1(\mathbb{R}^2)} \tilde{S}(v) \geq \frac{p-2}{2p} C_p^{\frac{p}{p-2}} > 0 \quad (55)$$

Consider now a function $u = \phi_\lambda + q\mathcal{G}_\lambda \in \tilde{N}_\omega \setminus H^1(\mathbb{R}^2)$ (so that $q \neq 0$) and fix $\lambda \in (\omega_0, \omega)$. Clearly $(\alpha + \theta_\lambda) > 0$, and thus there exists a constant $C > 0$ such that

$$\|\nabla \phi_\lambda\|_2^2 + \lambda \|\phi_\lambda\|_2^2 + (\omega - \lambda) \|u\|_2^2 + |q|^2 (\alpha + \theta_\lambda) \geq C (\|\phi_\lambda\|_{H^1}^2 + |q|^2). \quad (56)$$

Moreover, by Sobolev inequality we have that

$$\|u\|_p^p \leq C_p (\|\phi_\lambda\|_p^p + |q|^p) \leq C_p (\|\phi_\lambda\|_{H^1}^p + |q|^p) \leq C_p (\|\phi_\lambda\|_{H^1}^2 + |q|^2)^{\frac{p}{2}},$$

which implies

$$\|\phi_\lambda\|_{H^1}^2 + |q|^2 \geq \frac{1}{C_p} \|u\|_p^2. \quad (57)$$

Then, combining (56) and (57),

$$0 \geq I_\omega(u) \geq C (\|\phi_\lambda\|_{H^1}^2 + |q|^2) - \|u\|_p^p \geq \frac{C}{C_p} \|u\|_p^2 - \|u\|_p^p$$

and so, arguing as before, there exists $K_p > 0$, depending only on p , such that

$$\tilde{S}(u) \geq K_p$$

and, consequently,

$$\inf_{v \in \tilde{N}_\omega \setminus H^1(\mathbb{R}^2)} \tilde{S}(v) \geq K_p > 0. \quad (58)$$

Finally, combining (55) and (58), we obtain the claim. \square

For what concerns the right inequality in (50), we need to recall preliminarily some of the main properties of the NLS action minimizers at frequency ω , that is functions $u \in N_\omega^0$ such that $S_\omega^0(u) = d^0(\omega)$ (see Theorem 8.1.5 in [28]).

Proposition 4.8. *Let $p > 2$ and $\omega > 0$. Then, there exists at least an NLS action minimizer at frequency ω . In particular, such minimizer u is unique, positive and radially symmetric decreasing (up to gauge and translations invariances).*

Then, we can prove that also the right inequality of (50) is strict.

Proposition 4.9. *Let $p > 2$, $\alpha \in \mathbb{R}$. Then, $d(\omega) < d^0(\omega)$, for every $\omega > \omega_0$.*

Proof. For a fixed $\omega > \omega_0$, let u be the unique positive NLS action minimizer at frequency ω provided by Proposition 4.8. Then, u cannot be also a δ -NLS action minimizer at frequency ω . Indeed, if u were a δ -NLS action minimizer at frequency ω , then u would have to satisfy (10) and, in particular, $\phi_\lambda(\mathbf{0}) = (\alpha + \theta_\lambda)q$, but this can be proved to be a contradiction with the positivity of u by arguing as in the proof of Proposition 3.2. Hence, there exists $v \in N_\omega \setminus H^1(\mathbb{R}^2)$ such that $S_\omega(v) < S_\omega(u) = d^0(\omega)$, which concludes the proof. \square

Finally, we have all the tools to prove the existence part of Theorem 1.11.

Proof of Theorem 1.11-(i). The case $\omega \leq \omega_0$ has been already proved by the remarks at the beginning of the section and by Corollary 4.4. On the contrary, it is convenient to divide the proof of the case $\omega > \omega_0$ in four steps. We also note that, as in the proof of point (i) of Theorem 1.5, many of the following limits has to be meant as valid up to subsequences. We do not repeat it for the sake of simplicity and since this does not give rise to misunderstandings.

Step 1: weak convergence of the minimizing sequences. Fix $\omega > \omega_0$ and let $(u_n)_n$ be a δ -NLS action minimizing sequence at frequency ω , that is (by Remark 4.6) $(u_n)_n \subset \tilde{N}_\omega$ and $\tilde{S}(u_n) \rightarrow d(\omega)$, as $n \rightarrow +\infty$. In addition, for any fixed $\lambda > 0$ we can use for u_n the decomposition $u_n = \phi_{n,\lambda} + q_n \mathcal{G}_\lambda$. First, we see that, since $\|u_n\|_p^p \rightarrow \frac{2p}{p-2}d(\omega)$, $(u_n)_n$ is bounded in $L^p(\mathbb{R}^2)$. Moreover, as $I_\omega(u_n) \leq 0$, we get

$$\|\nabla \phi_{n,\lambda}\|_2^2 + \lambda \|\phi_{n,\lambda}\|_2^2 + (\omega - \lambda)\|u_n\|_2^2 + (\alpha + \theta_\lambda)|q_n|^2 \leq \|u_n\|_p^p.$$

Now, if one sets $\lambda = \frac{\omega + \omega_0}{2}$, then the three constants in front of $\|\phi_{n,\lambda}\|_2^2$, $\|u_n\|_2^2$ and $|q_n|^2$ are all strictly positive. Hence, $(\nabla \phi_{n,\lambda})_n$, $(\phi_{n,\lambda})_n$ and $(u_n)_n$ are bounded in $L^2(\mathbb{R}^2)$ and $(q_n)_n$ is bounded in \mathbb{C} . Thus, there exists $\phi_\lambda \in H^1(\mathbb{R}^2)$, $q \in \mathbb{C}$ and $u \in D$ such that $u = \phi_\lambda + q \mathcal{G}_\lambda$ and

$$\nabla \phi_{n,\lambda} \rightharpoonup \nabla \phi_\lambda, \quad \phi_{n,\lambda} \rightharpoonup \phi_\lambda, \quad u_n \rightharpoonup u \quad \text{in } L^2(\mathbb{R}^2) \quad \text{and} \quad q_n \rightarrow q \quad \text{in } \mathbb{C}.$$

Step 2: $u \in D \setminus H^1(\mathbb{R}^2)$. Assume, by contradiction, that $u \in H^1(\mathbb{R}^2)$, namely that $q = 0$, and define the sequence $w_n := \sigma_n \phi_{n,\lambda} \in H^1(\mathbb{R}^2)$, with

$$\sigma_n := \left(1 + \frac{I_\omega(u_n) - (\alpha + \theta_\lambda)|q_n|^2 + (\|u_n\|_p^p - \|\phi_{n,\lambda}\|_p^p) + (\omega - \lambda)(\|\phi_{n,\lambda}\|_2^2 - \|u_n\|_2^2)}{\|\phi_{n,\lambda}\|_p^p} \right)^{\frac{1}{p-2}},$$

so that $I_\omega^0(\sigma_n \phi_{n,\lambda}) = 0$. Note that σ_n is well defined since there exists $C > 0$ such that $\|\phi_{n,\lambda}\|_p^p \geq C$ for every $n \in \mathbb{N}$. Indeed, by Proposition 4.7, $\|u_n\|_p^p$ is uniformly bounded away from zero and $q_n \rightarrow 0$. On the other hand, since $|q_n|^2 \rightarrow 0$, it follows that both $\|\phi_{n,\lambda}\|_2^2 - \|u_n\|_2^2 \rightarrow 0$ and

$$\left| \|u_n\|_p^p - \|\phi_{n,\lambda}\|_p^p \right| \leq C_1 \left| \|u_n\|_p - \|\phi_{n,\lambda}\|_p \right| \leq C_2 \|u_n - \phi_{n,\lambda}\|_p \rightarrow 0.$$

As a consequence, since $I_\omega(u_n) \leq 0$, $(\sigma_n^p)_n$ is bounded from above by a sequence $(a_n)_n$ converging to 1. Thus, as $I_\omega^0(w_n) = 0$ and $\tilde{S}(u_n) \rightarrow d(\omega)$,

$$d^0(\omega) + o(1) = \tilde{S}(w_n) = \sigma_n^p \tilde{S}(\phi_{n,\lambda}) \leq a_n \left(\tilde{S}(u_n) + o(1) \right) = \tilde{S}(u_n) + o(1) = d(\omega) + o(1),$$

that implies that $d(\omega) \geq d^0(\omega)$, which contradicts Proposition 4.9.

Step 3: $u \in \tilde{N}_\omega$. In view of Step 2, it is left to prove that $I_\omega(u) \leq 0$. Assume by contradiction that $I_\omega(u) > 0$. From boundedness of $\phi_{n,\lambda}$ in $H^1(\mathbb{R}^2)$ and $q_n \rightarrow q$, one sees that $u_n \rightarrow u$ in $L_{loc}^p(\mathbb{R}^2)$ and hence $u_n \rightarrow u$ a.e. in \mathbb{R}^2 . As $(u_n)_n$ is bounded in $L^p(\mathbb{R}^2)$, one can use Brezis-Lieb lemma to get $\|u_n\|_p^p - \|u_n - u\|_p^p - \|u\|_p^p \rightarrow 0$, and thus

$$\tilde{S}(u_n) - \tilde{S}(u_n - u) - \tilde{S}(u) \rightarrow 0. \tag{59}$$

Since, in addition, $q_n \rightarrow q$, $\nabla\phi_{n,\lambda} \rightharpoonup \nabla\phi_\lambda$, $\phi_{n,\lambda} \rightharpoonup \phi_\lambda$ and $u_n \rightharpoonup u$ in $L^2(\mathbb{R}^2)$ and Q_ω is quadratic, one can also check that

$$I_\omega(u_n) - I_\omega(u_n - u) - I_\omega(u) \rightarrow 0. \quad (60)$$

Let us prove now that $I_\omega(u_n) \rightarrow 0$. Assume by contradiction that $I_\omega(u_n) \not\rightarrow 0$. As $\|u_n\|_p^p \leq C$, for some $C > 0$,

$$-C \leq I_\omega(u_n) \leq 0.$$

Hence, without loss of generality, we can suppose that $I_\omega(u_n) \rightarrow -\beta$, with $\beta > 0$. Consider, then, the sequence $v_n := \theta_n u_n$, with

$$\theta_n := \left(1 + \frac{I_\omega(u_n)}{\|u_n\|_p^p}\right)^{\frac{1}{p-2}},$$

so that $I_\omega(v_n) = 0$. Thus, an easy computation shows that

$$\theta_n \rightarrow l := \left(1 - \frac{\beta(p-2)}{2pd(\omega)}\right)^{\frac{1}{p-2}} < 1.$$

As a consequence,

$$\tilde{S}(v_n) = \tilde{S}(\theta_n u_n) = \theta_n^p \tilde{S}(u_n) \rightarrow l^p d(\omega) < d(\omega),$$

which is a contradiction. Hence $I_\omega(u_n) \rightarrow 0$. Finally, looking back at (60), since $I_\omega(u) > 0$ and $I_\omega(u_n) \rightarrow 0$,

$$I_\omega(u_n - u) = I_\omega(u_n) - I_\omega(u) + o(1) = -I_\omega(u) + o(1),$$

entailing that $I_\omega(u_n - u) \rightarrow -I_\omega(u) < 0$. Choose, then, \bar{n} such that $I_\omega(u_n - u) < 0$ for every $n \geq \bar{n}$. Since $d(\omega) \leq \tilde{S}(u_n - u)$ and $\tilde{S}(u) > 0$, (59) yields

$$d(\omega) \leq \liminf_n \tilde{S}(u_n - u) = d(\omega) - \tilde{S}(u) < d(\omega),$$

which is again a contradiction and entails $I_\omega(u) \leq 0$.

Step 4: conclusion. As boundedness in $L^p(\mathbb{R}^2)$ entails that $u_n \rightharpoonup u$ in $L^p(\mathbb{R}^2)$, by weak lower semicontinuity

$$\tilde{S}(u) \leq \liminf_{n \rightarrow +\infty} \tilde{S}(u_n) = d(\omega),$$

which concludes the proof. \square

5. FURTHER PROPERTIES: PROOF OF POINT (II) OF THEOREMS 1.5 AND 1.11

In this section we address (ii) of both Theorem 1.5 and Theorem 1.11, that is the features of δ -NLS ground states and δ -NLS action minimizers. However, we basically work on Theorem 1.11 since, by Lemma 1.10, this implies the result for Theorem 1.5.

We start by proving (ii)(a), namely the coexistence of the regular and the singular part for a δ -NLS action minimizer.

Proposition 5.1. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Let also u be a δ -NLS action minimizer at frequency ω . Then, $q \neq 0$ and $\phi_\lambda := u - q\mathcal{G}_\lambda \neq 0$, for every $\lambda > 0$.*

Proof. Let $\lambda > 0$ and consider the decomposition $u = \phi_\lambda + q\mathcal{G}_\lambda$. Assume by contradiction that $\phi_\lambda = 0$. Since $u \neq 0$, clearly $q \neq 0$. As u has to satisfy (10), then $(\alpha + \theta_\lambda) = 0$, so that $\lambda = \omega_0$. Since u has to satisfy also (11), with some computations one obtains that q have to satisfy

$$\omega - \omega_0 + |q|^{p-2} |\mathcal{G}_{\omega_0}(\mathbf{x})|^{p-2} = 0, \quad \forall \mathbf{x} \in \mathbb{R}^2 \setminus \{\mathbf{0}\},$$

which is clearly not possible.

On the other hand, assume by contradiction that $q = 0$, or equivalently that $u \in H^1(\mathbb{R}^2)$. This would imply that $d(\omega) = d^0(\omega)$, which contradicts Proposition 4.9. \square

We can move to the proof of point (ii)(b). Preliminarily, we note that, up to the multiplication times a phase factor, a δ -NLS action minimizer $u = \phi_\lambda + q\mathcal{G}_\lambda$ can be assumed to display a charge $q > 0$. Indeed, since $\mathcal{G}_\lambda(\mathbf{x}) > 0$ for every $\mathbf{x} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$ and $q \neq 0$, it is sufficient to multiply u times $e^{i\theta}$ in such a way that $qe^{i\theta} > 0$. In particular, if θ satisfies the equation $e^{i\theta} = \frac{\bar{q}}{|q|}$, then $qe^{i\theta} = |q|$. As a consequence, we will always assume throughout that $q > 0$.

The first key point for the proof of (ii)(b) is the switch to a minimum problem equivalent to that of S_ω constrained on N_ω , which is introduced in the next two results.

Lemma 5.2. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Then, for every fixed $C > 0$, there results*

$$\inf_{v \in D_C^p} I_\omega(v) = \inf_{v \in D_C^p} Q_\omega(v) - C.$$

with

$$D_C^p := \inf\{v \in D : \|v\|_p^p = C\}.$$

Moreover, if any, the minimizers of $I_\omega|_{D_C^p}$ and $Q_\omega|_{D_C^p}$ coincide.

Proof. This is a straightforward consequence of (19). Indeed, if $v \in D$ with $\|v\|_p^p = C$, then

$$I_\omega(v) = Q_\omega(v) - \|v\|_p^p = Q_\omega(v) - C. \quad \square$$

Proposition 5.3. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Let also $C(\omega) := \frac{2p}{p-2}d(\omega)$. Then,*

$$\inf_{v \in D_{C(\omega)}^p} Q_\omega(v) = C(\omega)$$

and there exists a function $u \in D_{C(\omega)}^p$ such that $Q_\omega(u) = C(\omega)$. In particular, there results that

$$\begin{cases} Q_\omega(w) = C(\omega) \\ w \in D_{C(\omega)}^p \end{cases} \iff \begin{cases} S_\omega(w) = d(\omega) \\ w \in N_\omega \end{cases} \quad (61)$$

Remark 5.4. In view of this result, one sees that, in order to study the features of δ -NLS action minimizers at frequency ω , it is sufficient (in fact, equivalent) to study the minimizers of Q_ω on $D_{C(\omega)}^p$, with $C(\omega) = \frac{2p}{p-2}d(\omega)$.

Proof of Proposition 5.3. Let u be δ -NLS action minimizer at frequency ω . Namely, (by the results obtained in Section 4) u is a minimizer of \tilde{S} on \tilde{N}_ω , so that $\|u\|_p^p \leq \|v\|_p^p$ for every $v \in \tilde{N}_\omega$, $\|u\|_p^p = C(\omega)$ and $I_\omega(u) = 0$.

Let $v \in D_{C(\omega)}^p$. First we see that $I_\omega(v) \geq 0 = I_\omega(u)$. Indeed, if we assume by contradiction that there exists $v \in D \setminus \{0\}$ such that $I_\omega(v) < 0$, then by Lemma 4.5 v cannot be a minimizer of \tilde{S} on \tilde{N}_ω , and thus $\|v\|_p^p > C(\omega)$, which contradicts the fact that $v \in D_{C(\omega)}^p$. Therefore, u is a minimizer of I_ω on $D_{C(\omega)}^p$, which yields, by Lemma 5.2, that u is also a minimizer of Q_ω on $D_{C(\omega)}^p$ and that $Q_\omega(u) = C(\omega)$.

This clearly proves the first part of the proposition and the reverse implication in (61). It is, then, to prove that every minimizer of Q_ω on $D_{C(\omega)}^p$ is a δ -NLS action minimizer at frequency ω . To this aim, let w be a minimizer of Q_ω on $D_{C(\omega)}^p$. It is straightforward that $I_\omega(w) = Q_\omega(w) - \|w\|_p^p = C(\omega) - C(\omega) = 0$ and that

$$S_\omega(w) = \tilde{S}(w) = \frac{p-2}{2p}\|w\|_p^p = \frac{p-2}{2p}C(\omega) = d(\omega),$$

which conclude the proof. □

We can now prove the first part of (ii)(b), which is the positivity up to gauge invariance.

Proposition 5.5. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Then, δ -NLS action minimizers at frequency ω are positive, up to gauge invariance.*

Proof. Let u be a δ -NLS action minimizer at frequency ω . Up to gauge invariance, it is not restrictive to assume $q > 0$. In addition, by Proposition 5.5, u is also a minimizer of Q_ω on $D_{C(\omega)}^p$, with $C(\omega) = \frac{2p}{p-2}d(\omega)$. Now, let us choose $\lambda = \omega$ in the decomposition of u and define $\Omega := \{\mathbf{x} \in \mathbb{R}^2 : \phi_\omega(\mathbf{x}) \neq 0\}$. By Proposition 5.1, $|\Omega| > 0$. Then, we can write

$$u(\mathbf{x}) = \phi_\omega(\mathbf{x}) + q\mathcal{G}_\omega(\mathbf{x}) = e^{i\eta(\mathbf{x})}|\phi_\omega(\mathbf{x})| + q\mathcal{G}_\omega(\mathbf{x}), \quad \forall \mathbf{x} \in \Omega \setminus \{\mathbf{0}\},$$

for some $\eta : \Omega \rightarrow [0, 2\pi)$. If one can prove that $\eta(\mathbf{x}) = 0$ for a.e. $\mathbf{x} \in \Omega \setminus \{\mathbf{0}\}$, then the proof is complete as this entails that $\phi_\omega(\mathbf{x}) = |\phi_\omega(\mathbf{x})| \geq 0$ for every $\mathbf{x} \in \mathbb{R}^2$, whence $u(\mathbf{x}) > 0$ for every $\mathbf{x} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$.

To this aim, assume by contradiction that $\eta \neq 0$ on $\Omega_1 \subset (\Omega \setminus \{\mathbf{0}\})$, with $|\Omega_1| > 0$. Letting $\tilde{u} := |\phi_\omega| + q\mathcal{G}_\omega$ (note that $u = \tilde{u}$ in $\mathbb{R}^2 \setminus \Omega_1$), there results that

$$\begin{aligned} |u(\mathbf{x})|^2 &= |\phi_\omega(\mathbf{x})|^2 + q^2\mathcal{G}_\omega^2(\mathbf{x}) + 2\cos(\eta(\mathbf{x}))|\phi_\omega(\mathbf{x})\mathcal{G}_\omega(\mathbf{x}) \\ &< |\phi_\omega(\mathbf{x})|^2 + q^2\mathcal{G}_\omega^2(\mathbf{x}) + 2|\phi_\omega(\mathbf{x})\mathcal{G}_\omega(\mathbf{x})| = |\tilde{u}(\mathbf{x})|^2, \quad \forall \mathbf{x} \in \Omega_1. \end{aligned}$$

Hence, as $|\Omega_1| > 0$,

$$\|u\|_p^p = \int_{\mathbb{R}^2} (|u|^2)^{\frac{p}{2}} d\mathbf{x} < \int_{\mathbb{R}^2} (|\tilde{u}|^2)^{\frac{p}{2}} d\mathbf{x} = \|\tilde{u}\|_p^p. \quad (62)$$

On the other hand, it is straightforward to check that $Q_\omega(\tilde{u}) \leq Q_\omega(u)$. Now, from (62) and the positivity of Q_ω , there exists $\beta \in (0, 1)$ such that $\|\beta\tilde{u}\|_p^p = \|u\|_p^p = C(\omega)$ and

$$Q_\omega(\beta\tilde{u}) = \beta^2 Q_\omega(\tilde{u}) < Q_\omega(u),$$

which contradicts the fact that u minimizes Q_ω on $D_{C(\omega)}^p$. Thus $\eta = 0$ a.e. on $\Omega \setminus \{\mathbf{0}\}$, which concludes the proof. \square

The proof of the previous result also entails that for, $\lambda = \omega$, the regular part ϕ_ω of a δ -NLS action minimizer at frequency ω is nonnegative. The following corollary points out that, whenever $\lambda > \omega$, it is in fact positive.

Corollary 5.6. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Let also u be a δ -NLS action minimizer at frequency ω . Then the regular part $\phi_\lambda := u - q\mathcal{G}_\lambda$ is positive for every $\lambda > \omega$, up to gauge invariance.*

Proof. Let u be a positive δ -NLS action minimizer at frequency ω and consider the decomposition $u = \phi_\lambda + q\mathcal{G}_\lambda$ for a fixed $\lambda > \omega$. First, using (23) and $q > 0$, we see that

$$\phi_\lambda(\mathbf{x}) = \phi_\omega(\mathbf{x}) + q(\mathcal{G}_\omega(\mathbf{x}) - \mathcal{G}_\lambda(\mathbf{x})) > 0, \quad \forall \mathbf{x} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}.$$

Then, one concludes the proof just recalling (7). \square

Finally, we may address the problem of the radially symmetric monotonicity of δ -NLS action minimizers.

Proposition 5.7. *Let $p > 2$, $\alpha \in \mathbb{R}$ and $\omega > \omega_0$. Then, δ -NLS action minimizers at frequency ω are radially symmetric decreasing, up to gauge invariance.*

Proof. Without loss of generality let u be a positive δ -NLS action minimizers at frequency ω . Consider also the decomposition $u = \phi_\omega + q\mathcal{G}_\omega$, corresponding to the choice $\lambda = \omega$. In order to prove the claim is sufficient to show that $\phi_\omega = \phi_\omega^*$, with ϕ_ω^* the radially symmetric nonincreasing rearrangement of ϕ_ω .

Assume, by contradiction, that $\phi_\omega \neq \phi_\omega^*$, that is ϕ_ω is not radially symmetric nonincreasing. Then, define the function $\tilde{u} = \phi_\omega^* + q\mathcal{G}_\omega$. By (40) and (32), we have $\|\nabla\phi_\omega^*\|_2 \leq \|\nabla\phi_\omega\|_2$ and $\|\phi_\omega^*\|_2 = \|\phi_\omega\|_2$, so that

$$Q_\omega(\tilde{u}) \leq Q_\omega(u).$$

Now, applying Proposition 2.3 with $f = q\mathcal{G}_\omega$ and $g = \phi_\omega$, there results that $\|\tilde{u}\|_p^p > \|u\|_p^p$, as $\phi_\omega \neq \phi_\omega^*$. Therefore, (as Q_ω is positive) there exists $\beta < 1$ such that $\|\beta\tilde{u}\|_p^p = \|u\|_p^p$ and

$$Q_\omega(\beta\tilde{u}) = \beta^2 Q_\omega(\tilde{u}) < Q_\omega(\tilde{u}) \leq Q_\omega(u),$$

but, via Proposition 5.3 (arguing as in the proof of Proposition 5.5), this contradicts that u is a δ -NLS action minimizer, thus concluding the proof. \square

We can now sum up all the previous results to prove point (ii) of Theorems 1.5 and 1.11.

Proof of Theorems 1.5 and 1.11-(ii). Let u be a δ -NLS action minimizer at frequency $\omega > \omega_0$. Then, by Proposition 5.1, Proposition 5.5, Corollary 5.6 and Proposition 5.7, u satisfies all the properties stated in item (ii).

Let $p \in (2, 4)$ and u be a δ -NLS ground state of mass μ . Combining Lemma 1.10 and point (i) of Theorem 1.11 one sees that u is also a δ -NLS action minimizer at some frequency $\omega > \omega_0$ (in particular, $\omega = \mu^{-1}(\|u\|_p^p - Q(u))$). Then, one concludes by point (ii) of Theorem 1.11. \square

Remark 5.8. Finally, we want to give some more details on the reasons for which all the features of both δ -NLS ground states and δ -NLS action minimizers are not obtained in a direct way, but by means of the strategy seen before.

Roughly speaking, our strategy mainly relies on the fact that we can trace back our problems to the issue of minimizing Q_ω on $D_{C(\omega)}^p$ with $C(\omega) = \frac{2p}{p-2}d(\omega)$. Then, given a minimizer u of this problem without the required property (i.e., positivity or radially symmetric monotonicity), we can exhibit a function \tilde{u} with such property such that

$$\|\tilde{u}\|_p > \|u\|_p \quad \text{and} \quad Q_\omega(\tilde{u}) \leq Q_\omega(u),$$

Moreover, noting that there exists $\beta < 1$ such that

$$\|\beta\tilde{u}\|_p^p = \|u\|_p^p \quad \text{and} \quad Q_\omega(\beta\tilde{u}) < Q_\omega(\tilde{u}),$$

we find a better competitor with respect to the minimizer, namely a contradiction.

Unfortunately, such a strategy is not applicable when dealing directly with the energy E or the action S_ω . More in detail, trying to repeat the procedure explained above on E , one obtains that

$$\|\tilde{u}\|_r > \|u\|_r \quad \text{for every } r \geq 2 \quad \text{and} \quad E(\tilde{u}) < E(u).$$

However, since the mass constraint is not fulfilled by \tilde{u} , we note that there exists $\beta < 1$ such that $\|\beta\tilde{u}\|_2^2 = \|u\|_2^2$, but here, since $\beta^2 > \beta^p$ and $E(\tilde{u}) < 0$, this yields

$$E(\beta\tilde{u}) = \frac{1}{2}\beta^2 Q(\tilde{u}) - \frac{1}{p}\beta^p \|\tilde{u}\|_p^p > \beta^p E(\tilde{u}) > E(\tilde{u}),$$

providing namely an inequality in the opposite direction with respect to the aimed one.

On the other hand, moving to the action S_ω and repeating again the same procedure, there results that

$$I_\omega(\tilde{u}) < 0 \quad \text{and} \quad S_\omega(\tilde{u}) < S_\omega(u).$$

However, if we set

$$\bar{\beta} := \left(\frac{Q_\omega(\tilde{u})}{\|\tilde{u}\|_p^p} \right)^{\frac{1}{p-2}},$$

then $\bar{\beta} < 1$, $I_\omega(\bar{\beta}\tilde{u}) = 0$ and

$$S_\omega(\bar{\beta}\tilde{u}) = \frac{1}{2}\bar{\beta}^2 Q(\tilde{u}) - \frac{1}{p}\bar{\beta}^p \|\tilde{u}\|_p^p > S_\omega(\tilde{u}). \quad (63)$$

Indeed, computing

$$\frac{d}{d\beta} S_\omega(\beta\tilde{u}) = \beta Q_\omega(\tilde{u}) - \beta^{p-1} \|\tilde{u}\|_p^p,$$

we find that $\frac{d}{d\beta}S_\omega(\beta\tilde{u}) > 0$ if and only if $0 < \beta < \bar{\beta}$, and $\frac{d}{d\beta}S_\omega(\beta\tilde{u})|_{\beta=\bar{\beta}} = 0$, so that $S_\omega(\bar{\beta}\tilde{u}) > S_\omega(\tilde{u})$. In other words, here again (63) is an inequality in the opposite direction with respect to the aimed one.

APPENDIX A. GROUND STATES, ACTION MINIMIZERS AND BOUND STATES

In this section, we show that both δ -NLS ground states and δ -NLS action minimizers are δ -NLS bound states, that is satisfy (10) and (11).

First, we note that (using either the Lagrange Multipliers theorem in the former case or the simple Du Bois-Raymond equation in the latter case), if u is either a δ -NLS ground state of mass μ or a δ -NLS action minimizers at frequency ω , then it satisfies, for any fixed $\lambda > 0$,

$$\begin{aligned} \langle \nabla\chi_\lambda, \nabla\phi_\lambda \rangle + \lambda\langle \chi_\lambda, \phi_\lambda \rangle + (\omega - \lambda)\langle \chi, u \rangle + \bar{\xi}q(\alpha + \theta_\lambda) - \langle \chi, |u|^{p-2}u \rangle &= 0 \\ \forall \chi &= \chi_\lambda + \xi\mathcal{G}_\lambda \in D. \end{aligned} \quad (64)$$

(whenever u is a δ -NLS ground state of mass μ , $\omega = \mu^{-1}(\|u\|_p^p - Q(u))$). Now, letting $\xi = 0$ in (64), so that $\chi = \chi_\lambda \in H^1(\mathbb{R}^2)$, there results

$$\langle \nabla\chi, \nabla\phi_\lambda \rangle + \langle \chi, \omega\phi_\lambda + (\omega - \lambda)q\mathcal{G}_\lambda - |u|^{p-2}u \rangle = 0 \quad \forall \chi \in H^1(\mathbb{R}^2).$$

Hence, as $\omega\phi_\lambda + (\omega - \lambda)q\mathcal{G}_\lambda - |u|^{p-2}u \in L^2(\mathbb{R}^2)$, $\phi_\lambda \in H^2(\mathbb{R}^2)$ and, by density,

$$-\Delta\phi_\lambda + \omega\phi_\lambda + (\omega - \lambda)q\mathcal{G}_\lambda - |u|^{p-2}u = 0 \quad \text{in } L^2(\mathbb{R}^2), \quad (65)$$

which is equivalent to (11). On the other hand, letting $\chi_\lambda = 0$ and $\xi = 1$ in (64), so that $\chi = \mathcal{G}_\lambda$, there results

$$\langle \mathcal{G}_\lambda, (\omega - \lambda)u - |u|^{p-2}u \rangle + q(\alpha + \theta_\lambda) = 0.$$

Finally, using (65), we obtain

$$\langle \mathcal{G}_\lambda, (-\Delta + \lambda)\phi_\lambda \rangle = q(\alpha + \theta_\lambda),$$

which is equivalent to $\phi_\lambda(0) = q(\alpha + \theta_\lambda)$, so that also (10) is satisfied.

APPENDIX B. ENERGY AND ACTION

Proof of Lemma 1.10. Let u be a δ -NLS ground state at mass μ and let $\omega > 0$ be the associated Lagrange multiplier, given by $\omega = \mu^{-1}(\|u\|_p^p - Q(u))$. Assume, by contradiction, that there exists $v = \eta_\lambda + \xi\mathcal{G}_\lambda \in N_\omega$ such that $S_\omega(v) < S_\omega(u)$ and let $\sigma > 0$ be such that $\|\sigma v\|_2^2 = \mu$. Then

$$S_\omega(\sigma v) = \frac{\sigma^2}{2}Q_\omega(v) - \frac{\sigma^p}{p}\|v\|_p^p.$$

Computing the derivative with respect to σ and using that $v \in N_\omega$, we get

$$\frac{d}{d\sigma}S_\omega(\sigma v) = \sigma Q_\omega(v) - \sigma^{p-1}\|v\|_p^p = \sigma I_\omega(v) + (\sigma - \sigma^{p-1})\|v\|_p^p = \sigma(1 - \sigma^{p-2})\|v\|_p^p,$$

which is greater than or equal to zero if and only if $0 < \sigma \leq 1$. Hence $S_\omega(\sigma v) \leq S_\omega(v)$, for every $\sigma > 0$. Therefore, since $S_\omega(\sigma v) \leq S_\omega(v) < S_\omega(u)$,

$$E(\sigma v) + \frac{\omega}{2}\|\sigma v\|_2^2 < E(u) + \frac{\omega}{2}\|u\|_2^2,$$

and using the fact that $\|\sigma v\|_2^2 = \|u\|_2^2 = \mu$, this entails $E(\sigma v) < E(u)$. However, as this contradicts the assumptions on u , we obtain that u is a δ -NLS action minimizer at frequency ω . \square

REFERENCES

- [1] Abramovitz M., Stegun I.A., *Handbook of Mathematical Functions: with Formulas, Graphs, and Mathematical Tables*, Dover, New York, 1965.
- [2] Adami R., Boni F., Dovetta S., Competing nonlinearities in NLS equations as source of threshold phenomena on star graphs, arXiv:2009.06424 [math.AP] (2020).
- [3] Adami R., Cacciapuoti C., Finco D., Noja D., Constrained energy minimization and orbital stability for the NLS equation on a star graph, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **31** (2014), no. 6, 1289–1310.
- [4] Adami R., Cacciapuoti C., Finco D., Noja D., Stable standing waves for a NLS on star graphs as local minimizers of the constrained energy, *J. Differential Equations* **260** (2016), no. 10, 7397–7415.
- [5] Adami R., Carlone R., Correggi M., Tentarelli L., Blow-up for the pointwise NLS in dimension two: absence of critical power, *J. Differential Equations* **269** (2020), no. 1, 1–37.
- [6] Adami R., Carlone R., Correggi M., Tentarelli L., Stability of the standing waves of the concentrated NLSE in dimension two, *Mathematics in Engineering* **3** (2021), no. 2, 1–15.
- [7] Adami R., Dell’Antonio G., Figari R., Teta A., The Cauchy problem for the Schrödinger equation in dimension three with concentrated nonlinearity, *Ann. I. H. Poincaré Anal. Non Linéaire* **20** (2003), no. 3, 477–500.
- [8] Adami R., Dell’Antonio G., Figari R., Teta A., Blow-up solutions for the Schrödinger equation in dimension three with a concentrated nonlinearity, *Ann. I. H. Poincaré Anal. Non Linéaire* **21** (2004), no. 1, 121–137.
- [9] Adami R., Fukuizumi R., Holmer J., Scattering for the L^2 supercritical point NLS, *Trans. Amer. Math. Soc.* **374** (2021), no. 1, 35–60.
- [10] Adami R., Nakamura T., Ruighi A., Discontinuous Ground States for NLSE on \mathbb{R} with a Fülöp-Tsutsui δ interaction, arXiv:2010.00895 [math.AP] (2020).
- [11] Adami R., Noja D., Existence of dynamics for a 1D NLS equation perturbed with a generalized point defect, *J. Phys. A* **42** (2009), no. 49, art. num. 495302, 19 pp.
- [12] Adami R., Noja D., Stability and symmetry-breaking bifurcation for the ground states of a NLS with a δ' interaction, *Commun. Math. Phys.* **318** (2013), no. 1, 247–289.
- [13] Adami R., Noja D., Ortoleva C., Orbital and asymptotic stability for standing waves of a nonlinear Schrödinger equation with concentrated nonlinearity in dimension three, *J. Math. Phys.* **54** (2013), no. 1, art. num. 013501, 33 pp.
- [14] Adami R., Noja D., Ortoleva C., Asymptotic stability for standing waves of a NLS equation with subcritical concentrated nonlinearity in dimension three: neutral modes, *Discrete Contin. Dyn. Syst. Ser. B* **36** (2016), no. 11, 5837–5879.
- [15] Adami R., Noja D., Visciglia N., Constrained energy minimization and ground states for NLS with point defects, *Discrete Contin. Dyn. Syst. Ser. B* **18** (2013), no. 5, 1155–1188.
- [16] Adami R., Teta A., A class of nonlinear Schrödinger equations with concentrated nonlinearity, *J. Funct. Anal.* **180** (2001), no. 1, 148–175.
- [17] Albeverio S., Gesztesy F., Høegh-Krohn R., Holden H., *Solvable Models in Quantum Mechanics*, Texts and Monographs in Physics, Springer-Verlag, New York, 1988.
- [18] Boni F., Dovetta S., Prescribed mass ground states for a doubly nonlinear Schrödinger equation in dimension one, *J. Math. Anal. Appl.* **496** (2021), no. 1, art. num. 124797, 16 pp.
- [19] Brézis H., Lieb E.H., A relation between pointwise convergence of functions and convergence of functionals, *Proc. Amer. Math. Soc.* **88** (1983), no. 3, 486–490.
- [20] Brézis H., Lions P.-L., A note on isolated singularities for linear elliptic equations, *Mathematical analysis and applications*, Part A, pp. 263–266, Adv. in Math. Suppl. Stud. 7a, Academic Press, New York-London, 1981.
- [21] Cacciapuoti C., Finco D., Noja D., Ground state and orbital stability for the NLS equation on a general starlike graph with potentials, *Nonlinearity* **30** (2017), no. 8, 3271–3303.
- [22] Cacciapuoti C., Finco D., Noja D., Well posedness of the nonlinear Schrödinger equation with isolated singularities, arXiv:2012.12391 [math.AP] (2021).
- [23] Carlone R., Correggi M., Figari R., Two-dimensional time-dependent point interactions, *Functional analysis and operator theory for quantum physics*, 189–211, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2017.
- [24] Carlone R., Correggi M., Tentarelli L., Well-posedness of the two-dimensional nonlinear Schrödinger equation with concentrated nonlinearity, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **36** (2019), no. 1, 257–294.
- [25] Carlone R., Exner P., Dynamics of an electron confined to a “hybrid plane” and interacting with a magnetic field, *Rep. Math. Phys.* **67** (2011), no. 2, 211–227.

- [26] Carlone R., Finco D., Tentarelli L., Nonlinear singular perturbations of the fractional Schrödinger equation in dimension one, *Nonlinearity* **32** (2019), no. 8, 3112–3143.
- [27] Caspers W., Clément P., A different approach to singular solutions, *Differential Integral Equations* **7** (1994), no. 5–6, 1227–1240.
- [28] Cazenave T. *Semilinear Schrödinger Equations*, Courant Lecture Notes 10, American Mathematical Society, Providence, RI, 2003.
- [29] Dovetta S., Serra E., Tilli P., Action versus energy ground states in nonlinear Schrödinger equations, arXiv:2107.08655 [math.AP] (2021).
- [30] Exner P., Šeba P., Quantum motion on a half-line connected to a plane, *J. Math. Phys.* **28** (1987), no. 2, 386–391.
- [31] Exner P., Šeba P., Mathematical models for quantum point-contact spectroscopy, *Czechoslovak J. Phys. B* **38** (1988), no. 1, 1–11.
- [32] Fukaya N., Georgiev V., Ikeda M., On stability and instability of standing waves for 2d-nonlinear Schrödinger equations with point interaction, arXiv:2109.04680 [math.AP] (2021).
- [33] Ghergu M., Kim S., Shahgholian H., Isolated singularities for semilinear elliptic systems with power-law nonlinearity, *Anal. PDE* **13** (2020), no. 3, 701–739.
- [34] Gidas B., Ni W.M., Nirenberg L., Symmetry and Related Properties via the Maximum Principle *Comm. Math. Phys.* **68** (1979), no. 3, 209–243.
- [35] Grillakis M., Shatah J., Strauss W., Stability theory of solitary waves in the presence of symmetry. I, *J. Funct. Anal.* **74** (1987), no. 1, 160–197.
- [36] Holmer J., Liu C., Blow-up for the 1D nonlinear Schrödinger equation with point nonlinearity I: Basic theory, *J. Math. Anal. Appl.* **483** (2020), no. 1, art. num. 123522, 20 pp.
- [37] Holmer J., Liu C., Blow-up for the 1D nonlinear Schrödinger equation with point nonlinearity II: Supercritical blow-up profiles, *Commun. Pure Appl. Anal.* **20** (2021), no. 1, 215–242.
- [38] Johnson R.A., Pan X.B., Yi Y., Singular solutions of the elliptic equation $-\Delta u - u + u^p = 0$, *Ann. Mat. Pura Appl. (4)* **166** (1994), 203–225.
- [39] Le Coz S., Fukuizumi R., Fibich G., Kshirim B., Sivan Y., Instability of bound states of a nonlinear Schrödinger equation with a Dirac potential, *Phys. D* **237** (2008), no. 8, 1103–1128.
- [40] Lieb E.H., Loss M., *Analysis. Second edition*, Graduate Studies in Mathematics 14, American Mathematical Society, Providence, RI, 2001.
- [41] Lions P.-L. Isolated singularities in semilinear problems, *J. Differential Equations* **38** (1980), no. 3, 441–450.
- [42] Lions P.-L., The concentration-compactness principle in the Calculus of Variations. The locally compact case. II, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **36** (1984), no. 4, 223–283.
- [43] Ni W.-M., Serrin J., Nonexistence theorems for singular solutions of quasilinear partial differential equations. *Comm. Pure Appl. Math.* **39** (1986), no. 3, 379–399.
- [44] Ni W.-M., Serrin J., Existence and nonexistence theorems for ground states for quasilinear partial differential equations, *Accad. Naz. dei Lincei* **77** (1986), 231–257.
- [45] Sakaguchi H., Malomed B.A., Singular solitons, *Phys. Rev. E* **101** (2020), no. 1, art. num. 012211, 10 pp.
- [46] Shamriz E., Chen Z., Malomed B.A., Sakaguchi H., Singular Mean-Field States: A Brief Review of Recent Results, *Condens. Matter* **5** (2020), no. 1, art. num. 20, 24 pp.
- [47] Véron L., Singular solutions of some nonlinear elliptic equations, *Nonlinear Anal.* **5** (1981), no. 3, 225–242.
- [48] Véron L., *Singularities of solutions of second order quasilinear equations*, Pitman Research Notes in Mathematics Series 353, Longman, Harlow, 1996.

POLITECNICO DI TORINO, DIPARTIMENTO DI SCIENZE MATEMATICHE “G.L. LAGRANGE”, CORSO DUCA DEGLI ABRUZZI, 24, 10129, TORINO, ITALY
 Email address: riccardo.adami@polito.it

POLITECNICO DI TORINO, DIPARTIMENTO DI SCIENZE MATEMATICHE “G.L. LAGRANGE”, CORSO DUCA DEGLI ABRUZZI, 24, 10129, TORINO, ITALY

UNIVERSITÀ DEGLI STUDI DI TORINO, DIPARTIMENTO DI MATEMATICA “G. PEANO”, VIA CARLO ALBERTO, 10, 10123, TORINO, ITALY

Email address: `filippo.boni@polito.it`

UNIVERSITÀ DEGLI STUDI DI NAPOLI FEDERICO II, DIPARTIMENTO DI MATEMATICA E APPLICAZIONI “RENATO CACCIOPOLI”, VIA CINTIA, MONTE S. ANGELO, 80126, NAPOLI, ITALY

Email address: `raffaele.carlone@unina.it`

POLITECNICO DI TORINO, DIPARTIMENTO DI SCIENZE MATEMATICHE “G.L. LAGRANGE”, CORSO DUCA DEGLI ABRUZZI, 24, 10129, TORINO, ITALY

Email address: `lorenzo.tentarelli@polito.it`