

THE STRONGLY QUASI-LOCAL COARSE NOVIKOV CONJECTURE AND BANACH SPACES WITH PROPERTY (H)

XIAOMAN CHEN, KUN GAO, AND JIAWEN ZHANG

ABSTRACT. In this paper, we introduce a strongly quasi-local version of the coarse Novikov conjecture, which states that certain assembly map from the coarse K -homology of a metric space to the K -theory of its strongly quasi-local algebra is injective. We prove that the conjecture holds for metric spaces with bounded geometry which can be coarsely embedded into Banach spaces with Property (H), introduced by Kasparov and Yu. Besides, we also generalise the notion of strong quasi-locality to proper metric spaces and provide a (strongly) quasi-local picture for K -homology.

1. INTRODUCTION

The coarse Novikov conjecture [8, 12, 26, 28] asserts that the coarse Baum-Connes assembly map from the coarse K -homology $KX_*(X)$ of a metric space X to the K -theory of its Roe algebra $C^*(X)$ is injective. Here the Roe algebra $C^*(X)$, encoding the coarse geometry of the space X , was introduced by John Roe [18, 20] in his pioneering work to study index theory on open manifolds.

The coarse Novikov conjecture is a geometric analogue of the strong Novikov conjecture [2] and suggests an approach to detect the higher index of the Dirac operator on a noncompact complete Riemannian manifold, which has many significant applications in geometry and topology. In particular, it implies the Gromov conjecture on non-existence of positive scalar curvature metrics on uniformly contractible Riemannian manifolds, and the zero-in-the-spectrum conjecture concerning the spectrum of the Laplacian operator.

Over the last two decades, there are a lot of remarkable progresses on the coarse Novikov conjecture. For example, Yu proved the coarse Baum-Connes conjecture, and consequently the coarse Novikov conjecture, for metric spaces with bounded geometry which admit a coarse embedding into a Hilbert space [29]. Later on, Kasparov and Yu proved the coarse Novikov conjecture for metric spaces with bounded geometry which can be coarsely embedded into a uniformly convex Banach space [12]. In [7], Gong, Wang and Yu introduced the maximal Roe algebras and studied a maximal version of the coarse Novikov conjecture. So far, no counterexample to the coarse Novikov conjecture has been discovered.

Date: December 20, 2021.

Key words and phrases. Roe algebras, (Strongly) quasi-local algebras, Coarse Novikov conjecture, Coarse embeddability, Property (H).

KG was supported by the National Natural Science Foundation of China under Grant (No.12071183).

Property (H) is a geometric condition to Banach spaces, introduced by Kasparov and Yu in [13] where they proved the strong Novikov conjecture for groups which can be coarsely embedded into a Banach space with Property (H). Later on, Wang, Yu and the first author proved the coarse Novikov conjecture for metric spaces with bounded geometry which can be coarsely embedded into a Banach space with Property (H) [4].

Definition A ([13]). A real Banach space V is said to have Property (H) if there exist increasing sequences of finite dimensional subspaces $\{V_n\}_{n \in \mathbb{N}}$ of V and $\{W_n\}_{n \in \mathbb{N}}$ of a real Hilbert space W such that:

- (1) $\bigcup_{n=1}^{\infty} V_n$ is dense in V ;
- (2) there exists a uniformly continuous map $\psi : S(\bigcup_{n=1}^{\infty} V_n) \rightarrow S(\bigcup_{n=1}^{\infty} W_n)$ such that $\psi|_{S(V_n)}$ is a homeomorphism onto $S(W_n)$ for any $n \in \mathbb{N}$, where $S(\cdot)$ denotes the unit sphere of a subspace of a Banach space.

For example, the Banach space $\ell^p(\mathbb{N})$ and the Banach space of all Schatten- p class operators on a separable Hilbert space have Property (H) for $p \geq 1$ due to the Mazur(-type) map. One can also check that uniformly convex Banach spaces with unconditional basis have Property (H) (see, *e.g.*, [3, Chapter 9]).

In this paper, we introduce a strongly quasi-local version of the coarse Novikov conjecture and prove it for metric spaces with bounded geometry which can be coarsely embedded into a Banach space with Property (H).

Recall that the notion of quasi-locality was introduced by Roe as an intrinsic characterisation for elements in the Roe algebra [18]. It turns out that the quasi-local algebra $C_q^*(X)$ of a metric space X (*i.e.*, the C^* -algebra of all locally compact and quasi-local operators on X) contains the Roe algebra $C^*(X)$ as a C^* -subalgebra, and they coincide when the space X has Yu's property A [15, 23]. On the other hand, Engel [6] discovered that while indices of genuine differential operators on Riemannian manifolds live in the K -theory of (appropriate) Roe algebras, the indices of uniform pseudo-differential operators are only known to be in the K -theory of quasi-local algebras. Hence it is important to study whether the Roe algebra and the quasi-local algebra have the same K -theory.

As a potential approach, the first and third authors together with Bao introduced the following notion of strong quasi-locality in [1] (see Section 3 for more details). Fix an infinite-dimensional separable Hilbert space \mathcal{H} and denote $\mathfrak{K}(\mathcal{H})_1$ the unit ball of the compact operators on \mathcal{H} .

Definition B ([1]). Let X be a discrete metric space with bounded geometry and $T \in \mathfrak{B}(\ell^2(X; \mathcal{H}))$. We say that T is *strongly quasi-local* if for any $\varepsilon > 0$ there exist $\delta, R > 0$ such that for any map $g : X \rightarrow \mathfrak{K}(\mathcal{H})_1$ satisfying that $d(x, y) < R$ implies $\|g(x) - g(y)\| < \delta$, we have

$$\|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| < \varepsilon$$

where $\Lambda(g) \in \mathfrak{B}(\ell^2(X; \mathcal{H} \otimes \mathcal{H}))$ is defined by $\Lambda(g)(\delta_x \otimes \xi \otimes \eta) := \delta_x \otimes \xi \otimes g(x)\eta$ for $\delta_x \otimes \xi \otimes \eta \in \ell^2(X; \mathcal{H} \otimes \mathcal{H}) \cong \ell^2(X) \otimes \mathcal{H} \otimes \mathcal{H}$.

The strongly quasi-local algebra $C_{sq}^*(X)$ for a discrete metric space X is defined to be the C^* -algebra of all locally compact and strongly quasi-local operators on X . The strongly quasi-local algebra $C_{sq}^*(X)$ contains the Roe algebra $C^*(X)$, and the main result of [1] states that they have the same K -theory when X has bounded geometry and can be coarsely embedded into a Hilbert space.

The main focus of this paper is the following strongly quasi-local version of the coarse Novikov conjecture:

The strongly quasi-local coarse Novikov conjecture. If X is a discrete metric space with bounded geometry, then the following *strongly quasi-local coarse assembly map* $\mu_{sq} := i_* \circ \mu$ is injective:

$$\mu_{sq} : KX_*(X) \xrightarrow{\mu} K_*(C^*(X)) \xrightarrow{i_*} K_*(C_{sq}^*(X))$$

where μ is the coarse Baum-Connes assembly map.

One can also consider the strongly quasi-local coarse Baum-Connes conjecture which asserts that the above map μ_{sq} is bijective. Combining with Yu's result [29], the main result of [1] can be restated that the strongly quasi-local coarse Baum-Connes conjecture holds for metric spaces with bounded geometry which can be coarsely embedded into a Hilbert space.

The following is the main result of this paper:

Theorem C. *Let X be a bounded geometry metric space which can be coarsely embedded into a Banach space with Property (H). Then the strongly quasi-local coarse Novikov conjecture holds for X .*

The proof of Theorem C is inspired by that of [4, Theorem 1.1], but is more involved and requires new techniques. Amongst other pieces, the key ingredient in the proof is that we introduce a twisted quasi-local algebra and construct a Bott map from the K -theory of strongly quasi-local algebra to that of the twisted quasi-local algebra. Recall that in the case of Roe algebras, a Bott map was constructed in [4] thanks to the finite propagation approximations for operators in Roe algebras. However for general operators in the strongly quasi-local algebra, it is unclear whether one can still find finite propagation approximations. To overcome this issue, we make use of the hypothesis of strong quasi-locality together with some technical arguments to obtain the required Bott map.

On the other hand, we also provide a (strongly) quasi-local picture for K -homology. Recall that Yu [27] introduced a notion of localisation algebra $C_L^*(X)$ for a proper metric space X and showed that its K -theory is isomorphic to the K -homology of X (see also [17]). Note that operators in $C_L^*(X)$ have smaller and smaller propagation as the parameter tends to infinity. Following the same idea, we introduce a notion of localisation (strongly) quasi-local algebra where the

“extent” of (strong) quasi-locality of operators therein get better and better as the parameter tends to infinity (see Definition 4.3 and 4.5). We prove the following:

Theorem D. *Let X be a proper metric space. The localisation algebra, the localisation quasi-local algebra and the localisation strongly quasi-local algebra have the same K -theory, which coincides with the K -homology of X .*

Consequently, we obtain a (strongly) quasi-local description of the coarse K -homology. Moreover, the strongly quasi-local coarse assembly map μ_{sq} above is induced by an evaluation map (see Corollary 4.12). This is also based on our generalised notion of strong quasi-locality for proper metric spaces (note that the original one introduced in [1] is only for discrete metric spaces).

The paper is organised as follows. In Section 2, we collect notions from coarse geometry and recall the definitions of Roe and quasi-local algebras. In Section 3, we generalise the notion of strong quasi-locality to the case of proper metric spaces and verify their coarse geometric properties, based on which we introduce a (strong) quasi-local picture for K -homology and prove Theorem D in Section 4. The rest of the paper is contributed to the proof of Theorem C. More precisely, in Section 5 we recall the twisted Roe algebras and introduce their quasi-local counterparts for metric spaces which can be coarsely embedded into a Banach space with Property (H). Then in Section 6 we construct Bott maps to link the K -theories of Roe and strongly quasi-local algebras with those of their twisted counterparts, and prove that these twisted algebras have the same K -theory (Theorem 7.1) in Section 7. We finish the proof of Theorem C in Section 8. Finally in Section 9, we outline the case of metric spaces which can be coarsely embedded into a simply-connected complete Riemannian manifold with non-positively sectional curvature.

Convention. Throughout the paper, we fix an infinite-dimensional separable Hilbert space \mathcal{H} . Denote by $\mathfrak{K} := \mathfrak{K}(\mathcal{H})$ the C^* -algebra of compact operators on \mathcal{H} , and \mathfrak{K}_1 its closed unit ball (with respect to the operator norm).

2. PRELIMINARIES

We start with some notions from coarse geometry and higher index theory.

2.1. Notions from coarse geometry. Here we collect several basic notions.

Definition 2.1. Let (X, d) be a metric space.

- (1) A subset $A \subseteq X$ is said to be *bounded* if $\sup\{d(x, y) : x, y \in A\}$ is finite.
- (2) For $x \in X$ and $R > 0$, the *open R -ball* of x is $B(x, R) := \{y \in X : d(x, y) < R\}$.
- (3) (X, d) is said to be *proper* if any bounded closed subset in X is compact.
- (4) For a subset $A \subseteq X$ and $R \geq 0$, the *R -neighbourhood* of A is defined to be $\mathcal{N}_R(A) := \{x \in X : d_X(x, A) \leq R\}$.

- (5) A subset $A \subseteq X$ is called a *net* in X if there exists $C > 0$ such that for any $x \in X$ there exists $y \in A$ with $d(x, y) \leq C$. In this case, we also say that A is a *C-net* in X .
- (6) A subset $A \subseteq X$ is said to be *uniformly discrete* if there exists $C > 0$ such that $d(x, y) \geq C$ for any $x \neq y$ in A . In this case, we also say that A is *C-discrete*.
- (7) (X, d) is said to have *bounded geometry* if X contains a uniformly discrete net Γ with bounded geometry, i.e., for any $R \geq 0$ there exists an $N \in \mathbb{N}$ such that $\#(B(x, R) \cap \Gamma) \leq N$ for any $x \in \Gamma$.

Definition 2.2. Let f be a map between two metric spaces (X, d_X) and (Y, d_Y) .

- (1) f is *uniformly expansive* if there exists a non-decreasing function $\rho_+ : [0, \infty) \rightarrow [0, \infty)$ such that for any $x, y \in X$ we have:

$$d_Y(f(x), f(y)) \leq \rho_+(d_X(x, y)).$$

- (2) f is *proper* if for any bounded $B \subseteq Y$, the preimage $f^{-1}(B)$ is bounded in X .

- (3) f is *coarse* if it is uniformly expansive and proper.

- (4) f is *effectively proper* if there exists a proper non-decreasing function $\rho_- : [0, \infty) \rightarrow [0, \infty)$ such that for any $x, y \in X$ we have:

$$\rho_-(d_X(x, y)) \leq d_Y(f(x), f(y)).$$

- (5) f is a *coarse embedding* if it is uniformly expansive and effectively proper.

Definition 2.3. Let (X, d_X) and (Y, d_Y) be metric spaces.

- (1) Two maps $f, g : (X, d_X) \rightarrow (Y, d_Y)$ are *close* if there exists $R \geq 0$ such that for any $x \in X$, we have $d_Y(f(x), g(x)) \leq R$.
- (2) A coarse map $f : (X, d_X) \rightarrow (Y, d_Y)$ is called a *coarse equivalence* if there exists another coarse map $g : (Y, d_Y) \rightarrow (X, d_X)$ such that $f \circ g$ and $g \circ f$ are close to identities, where g is called a *coarse inverse* of f .
- (3) (X, d_X) and (Y, d_Y) are said to be *coarsely equivalent* if there exists a coarse equivalence from X to Y .

2.2. Roe algebras and quasi-local algebras. For a proper metric space (X, d_X) , recall that an *X-module* is a non-degenerate $*$ -representation $C_0(X) \rightarrow \mathfrak{B}(\mathcal{H}_X)$ where \mathcal{H}_X is some infinite-dimensional separable Hilbert space. We also say that \mathcal{H}_X is an *X-module* if the representation is clear from the context. An *X-module* is called *ample* if no non-zero element of $C_0(X)$ acts as a compact operator on \mathcal{H}_X . Note that every proper metric space X admits an ample *X-module*.

Let \mathcal{H}_X and \mathcal{H}_Y be ample modules of proper metric spaces X and Y , respectively. Given an operator $T \in \mathfrak{B}(\mathcal{H}_X, \mathcal{H}_Y)$, the *support* of T is defined to be

$$\text{supp}(T) := \{(y, x) \in Y \times X : \chi_V T \chi_U \neq 0 \text{ for all neighbourhoods } U \text{ of } x \text{ and } V \text{ of } y\}.$$

When $X = Y$, the *propagation* of $T \in \mathfrak{B}(\mathcal{H}_X)$ is defined to be

$$\text{prop}(T) := \sup\{d_X(y, x) : (y, x) \in \text{supp}(T)\}.$$

We say that an operator $T \in \mathfrak{B}(\mathcal{H}_X)$ has *finite propagation* if $\text{prop}(T)$ is finite, and T is *locally compact* if fT and Tf are compact operators for all $f \in C_0(X)$.

Definition 2.4. For a proper metric space X and an ample X -module \mathcal{H}_X , $\mathbb{C}[\mathcal{H}_X]$ is defined to be the $*$ -algebra of locally compact and finite propagation operators on \mathcal{H}_X , and the *Roe algebra* $C^*(\mathcal{H}_X)$ of \mathcal{H}_X is defined to be the norm-closure of $\mathbb{C}[\mathcal{H}_X]$ in $\mathfrak{B}(\mathcal{H}_X)$.

It is a standard result that the Roe algebra $C^*(\mathcal{H}_X)$ does not depend on the chosen ample module \mathcal{H}_X up to $*$ -isomorphisms, hence denoted by $C^*(X)$ and called the *Roe algebra of X* . Furthermore, $C^*(X)$ is a coarse invariant of the metric space X (up to non-canonical $*$ -isomorphisms), and their K -theories are coarse invariants up to canonical isomorphisms (see, e.g., [19]).

Now we move on to the notion of quasi-locality, introduced by Roe in [18].

Definition 2.5. Let (X, d) be a metric space, and \mathcal{H}_X be an ample X -module. An operator $T \in \mathfrak{B}(\mathcal{H}_X)$ is said to be *quasi-local* if for any $\epsilon > 0$ there exists $R > 0$ such that for any $A, B \subseteq X$ with $d(A, B) > R$, we have $\|\chi_A T \chi_B\| < \epsilon$.

It is clear that the set of all quasi-local operators on \mathcal{H}_X forms a C^* -subalgebra in $\mathfrak{B}(\mathcal{H}_X)$, which leads to the following:

Definition 2.6. For a proper metric space X and an ample X -module \mathcal{H}_X , the set of locally compact quasi-local operators on \mathcal{H}_X is called the *quasi-local algebra of \mathcal{H}_X* , denoted by $C_q^*(\mathcal{H}_X)$.

It follows directly from definitions that finite propagation operators are quasi-local, and hence the Roe algebra $C^*(\mathcal{H}_X)$ is a subalgebra of $C_q^*(\mathcal{H}_X)$. For the converse, it was shown in [23] that these two algebras coincide when X has property A. However, the general case is still widely open.

As in the case of Roe algebras, it was proved in [1, Corollary 2.10] that the quasi-local algebra $C_q^*(\mathcal{H}_X)$ does not depend on the chosen ample module \mathcal{H}_X up to $*$ -isomorphisms. Hence we call it the *quasi-local algebra of X* and denote by $C_q^*(X)$. Furthermore, it was shown that quasi-local algebras are coarse invariants (up to non-canonical $*$ -isomorphisms), and their K -theories are coarse invariants up to canonical isomorphisms.

We would like to recall the following characterisation for quasi-locality from [22], which is also the motivation to introduce strong quasi-locality in [1]. To state the result, we need some more notions:

Definition 2.7. Let (X, d_X) be a metric space and $g : X \rightarrow \mathbb{C}$ be a Borel function. We say that g has (ϵ, R) -variation if for any $x, y \in X$ with $d_X(x, y) < R$, we have $|g(x) - g(y)| < \epsilon$. We say that g is *bounded* if its norm $\|g\|_\infty := \sup_{x \in X} |g(x)|$ is finite.

Proposition 2.8. Let X be a proper metric space, \mathcal{H}_X an ample X -module and $T \in \mathfrak{B}(\mathcal{H}_X)$ be a locally compact operator. Then the following are equivalent:

- (1) T is quasi-local in the sense of Definition 2.5;
- (2) For any $\epsilon > 0$, there exist $\delta, R > 0$ such that for any continuous function $g : X \rightarrow \mathbb{C}$ with norm 1 and (δ, R) -variation then $\|[T, g]\| < \epsilon$;

- (3) For any $\varepsilon > 0$, there exist $\delta, R > 0$ such that for any Borel function $g : X \rightarrow \mathbb{C}$ with norm 1 and (δ, R) -variation then $\| [T, g] \| < \varepsilon$.

Note that the equivalence between (1) and (2) is the “easier” part of [22, Theorem 2.8] (see also [1, Proposition 3.3]). And also note the same argument can deduce the equivalence between (1) and (3), hence omitted.

We remark that condition (3) above is the starting point to introduce the notion of strong quasi-locality for general proper metric spaces in Section 3.

2.3. Coarse K -homology and the assembly map. Recall that the locally finite K -homology groups $K_i(X)$ ($i = 0, 1$) for a proper metric space X are generated by certain cycles modulo certain equivalence relations [10, 11]:

- (1) a cycle for $K_0(X)$ is a pair (\mathcal{H}_X, F) , where \mathcal{H}_X is an X -module and F is a bounded linear operator acting on \mathcal{H}_X such that $F^*F - I$ and $FF^* - I$ are locally compact, and $\phi F - F\phi$ is compact for all $\phi \in C_0(X)$;
- (2) a cycle for $K_1(X)$ is a pair (\mathcal{H}_X, F) , where \mathcal{H}_X is an X -module and F is a self-adjoint operator acting on \mathcal{H}_X such that $F^2 - I$ are locally compact, and $\phi F - F\phi$ is compact for all $\phi \in C_0(X)$;

In both cases, the equivalence relations on cycles are given by homotopy of the operators F , unitary equivalence, and direct sum with “degenerate” cycles, *i.e.*, those cycles for which $F\phi - \phi F$, $\phi(F^*F - I)$ and so on, are actually zero.

Now we recall the definition of the assembly map $\mu : K_*(X) \rightarrow K_*(C^*(X))$.

Definition 2.9. Let (\mathcal{H}_X, F) represent a cycle in $K_0(X)$. For each $R > 0$, we take a locally finite uniformly bounded open cover $\{U_i\}_i$ of X with $\text{diam}(U_i) < R$ for all i , and a continuous partition unity $\{\phi_i\}_i$ subordinate to $\{U_i\}_i$. Define $F = \sum_i \phi_i^{1/2} T \phi_i^{1/2}$, where the sum converges in the strong operator topology. Then (\mathcal{H}_X, F) and (\mathcal{H}_X, T) are equivalent in $K_0(X)$. The operator F has finite propagation less than R , hence $F^*F - I$ and $FF^* - I$ are in $\mathbb{C}[X]$. We set:

$$W = \begin{pmatrix} I & F \\ 0 & F \end{pmatrix} \begin{pmatrix} I & 0 \\ -F^* & I \end{pmatrix} \begin{pmatrix} I & F \\ 0 & F \end{pmatrix} \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix} \in \mathfrak{B}(\mathcal{H}_X \oplus \mathcal{H}_X).$$

Then both W and W^{-1} have finite propagations (at most $3R$), and

$$W \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} W^{-1} - \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \in \mathbb{C}[X] \otimes M_2.$$

We then define

$$\mu([(\mathcal{H}_X, T)]) := \left[W \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} W^{-1} \right] - \left[\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} \right]$$

in $K_0(\mathbb{C}[X])$. Furthermore, $\mu([(\mathcal{H}_X, T)])$ defines an element in $K_0(C^*(X))$ by considering $\mathbb{C}[X]$ as a $*$ -subalgebra of $C^*(X)$. This element is denoted by $\mu([(\mathcal{H}_X, T)]) \in K_0(C^*(X))$. Thus, we obtain the assembly map $\mu : K_0(X) \rightarrow K_0(C^*(X))$. Similarly, we can define $\mu : K_1(X) \rightarrow K_1(C^*(X))$.

To recall the definition of coarse K -homology, we need the notion of *Rips complex*.

Definition 2.10. Let X be a discrete metric space with bounded geometry. For each $d \geq 0$, the *Rips complex* $P_d(X)$ at scale d is defined to be the simplicial complex in which the vertex set is X and a finite subset $\{x_0, x_1, \dots, x_q\} \subseteq X$ spans a simplex if and only if $d(x_i, x_j) \leq d$ for all $0 \leq i, j \leq q$.

We endow $P_d(X)$ with the following *spherical metric*. On each connected component of $P_d(X)$, the spherical metric is the maximal metric whose restriction on each simplex $\Delta := \{\sum_{i=0}^q t_i x_i : t_i \geq 0, \sum_{i=0}^q t_i = 1\}$ is the metric obtained by identifying Δ with S_+^q via the map

$$\sum_{i=0}^q t_i x_i \mapsto \left(\frac{t_0}{\sqrt{\sum_{i=0}^q t_i^2}}, \frac{t_1}{\sqrt{\sum_{i=0}^q t_i^2}}, \dots, \frac{t_q}{\sqrt{\sum_{i=0}^q t_i^2}} \right),$$

where $S_+^q := \{(s_0, s_1, \dots, s_q) \in \mathbb{R}^{q+1} : s_i \geq 0, \sum_{i=0}^q s_i^2 = 1\}$ is endowed with the standard Riemannian metric. If y_0, y_1 belong to two different connected components Y_0, Y_1 of $P_d(X)$, respectively, we define

$$d(y_0, y_1) = \min \{d(y_0, x_0) + d_X(x_0, x_1) + d(x_1, y_1) : x_0 \in X \cap Y_0, x_1 \in X \cap Y_1\}.$$

Note that for any $d \geq 0$, $P_d(X)$ is coarsely equivalent to X via the inclusion map. If $d < d'$, then $P_d(X) \subseteq P_{d'}(X)$ as a subcomplex. Hence we define:

Definition 2.11. The coarse K -homology for a metric space (X, d) , denoted by $KX_*(X)$, is defined to be the direct limit of the K -homology of $P_d(X)$, i.e.,

$$KX_*(X) := \lim_{d \rightarrow \infty} K_*(P_d(X)).$$

Passing the assembly map μ to direct limit as well, we obtain the *coarse assembly map*:

$$(2.1) \quad \mu : \lim_{d \rightarrow \infty} K_*(P_d(X)) \longrightarrow \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) \cong K_*(C^*(X)).$$

The coarse Baum-Connes conjecture. If X is a discrete metric space with bounded geometry, then the coarse assembly map μ defined in (2.1) is bijective.

The coarse Novikov conjecture. If X is a discrete metric space with bounded geometry, then the coarse assembly map μ defined in (2.1) is injective.

The coarse Novikov conjecture provides a method to determine nonvanishing of the higher index of the Dirac operator on a noncompact complete Riemannian manifold, which has many significant applications in geometry and topology. In particular, it implies the Gromov positive scalar curvature conjecture and the zero-in-the-spectrum conjecture (see [9, 19, 20, 26, 29]).

3. STRONGLY QUASI-LOCAL ALGEBRAS

In [1], the authors introduced a new class of operator algebras for discrete metric spaces, called strongly quasi-local algebras. They sit between Roe algebras

and quasi-local algebras, and are shown to be coarse invariants. In this section, we introduce strongly quasi-local algebras for general proper metric spaces and study their coarse geometric properties. This is the preparation to explore the strongly quasi-local perspective on coarse K -homology in the next section.

3.1. Strong quasi-locality. First, let us recall the notion of strong quasi-locality for operators over discrete metric spaces from [1]. Recall that \mathcal{H} is the infinite-dimensional separable Hilbert space fixed at the very beginning, $\mathfrak{K} = \mathfrak{K}(\mathcal{H})$ is the C^* -algebra of compact operators on \mathcal{H} , and \mathfrak{K}_1 is the closed unit ball of \mathfrak{K} .

Let X be a proper *discrete* metric space and \mathcal{H}_X be an ample X -module. For each $x \in X$, denote $\mathcal{H}_x := \chi_{\{x\}} \mathcal{H}_X$. An operator $S \in \mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H})$ can be regarded as an X -by- X matrix $(S_{xy})_{x,y \in X}$, where $S_{xy} \in \mathfrak{B}(\mathcal{H}_y \otimes \mathcal{H}, \mathcal{H}_x \otimes \mathcal{H})$.

Recall that a map $g : X \rightarrow \mathfrak{K}$ has (ε, R) -variation if for any $x, y \in X$ with $d(x, y) < R$, we have $\|g(x) - g(y)\| < \varepsilon$. The map g is *bounded* if $\|g\|_\infty := \sup_{x \in X} \|g(x)\| < \infty$. Given a bounded map $g : X \rightarrow \mathfrak{K}$, we define an operator $\Lambda(g) \in \mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H})$ by setting its matrix entry as follows:

$$(3.1) \quad \Lambda(g)_{xy} := \begin{cases} \text{Id}_{\mathcal{H}_x} \otimes g(x), & y = x; \\ 0, & \text{otherwise.} \end{cases}$$

Note that this is a block-diagonal operator with respect to the decomposition $\mathcal{H}_X = \bigoplus_{x \in X} \mathcal{H}_x$.

Definition 3.1 ([1, Definition 3.4]). Let X be a proper *discrete* metric space and \mathcal{H}_X be an ample X -module. An operator $T \in \mathfrak{B}(\mathcal{H}_X)$ is called *strongly quasi-local* if for any $\varepsilon > 0$ there exist $\delta, R > 0$ such that for any map $g : X \rightarrow \mathfrak{K}_1$ with (δ, R) -variation, we have $\|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\|_{\mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H})} < \varepsilon$. The set of all locally compact strongly quasi-local operators on \mathcal{H}_X forms a C^* -algebra, called the *strongly quasi-local algebra of \mathcal{H}_X* and denoted by $C_{sq}^*(\mathcal{H}_X)$.

It was shown in [1] that $C_{sq}^*(\mathcal{H}_X)$ does not depend on the chosen module \mathcal{H}_X . Note that when \mathcal{H}_X is chosen to be $\ell^2(X; \mathcal{H})$, then the above definition coincides with Definition B.

Now we move to the case of proper metric spaces which are *not* necessarily discrete. Our idea is inspired by Proposition 2.8(3) while we need the following notion of measurability for operator-valued maps, which can be found in standard textbooks on functional analysis (e.g., [25, Section V.4]).

Definition 3.2. Let (X, d) be a proper metric space with the Borel σ -algebra and $g : X \rightarrow \mathfrak{K}$ be a map.

- (1) g is said to be *countably-valued* if there is a countable Borel partition $\{E_n\}_{n \in \mathbb{N}}$ of X such that $g|_{E_n}$ is constant, i.e., $g = \sum_{n \in \mathbb{N}} a_n \chi_{E_n}$ where $a_n \in \mathfrak{K}$;

- (2) g is said to be *strongly measurable*¹ if there is a sequence of countably-valued maps $\{g_n\}_{n \in \mathbb{N}}$ such that $g_n \rightarrow g$ pointwise;
- (3) g is said to be *weakly measurable* if for any bounded linear functional $\phi : \mathfrak{K} \rightarrow \mathbb{C}$, the composition $\phi \circ g : X \rightarrow \mathbb{C}$ is a Borel function on X .

The following result is essentially due to Pettis:

Proposition 3.3 ([16]). *Let (X, d) be a proper metric space with the Borel σ -algebra and $g : X \rightarrow \mathfrak{K}$ be a map. Then the following are equivalent:*

- (1) g is strongly measurable;
- (2) g is weakly measurable;
- (3) there is a sequence of countably-valued maps $\{g_n\}_{n \in \mathbb{N}}$ such that $\|g_n - g\|_\infty \rightarrow 0$.

A key ingredient for strong quasi-locality is to define the operator $\Lambda(g)$ for “well-behaved” operator-valued maps $g : X \rightarrow \mathfrak{K}$ as in (3.1). Let us start with the case of countably-valued maps.

Definition 3.4. Let (X, d) be a proper metric space and \mathcal{H}_X an ample X -module. Let $g : X \rightarrow \mathfrak{K}$ be a bounded countably-valued map, i.e., $g = \sum_{n \in \mathbb{N}} a_n \chi_{E_n}$ where $\{E_n\}_n$ is a Borel partition of X and $a_n \in \mathfrak{K}$ with $\sup_{n \in \mathbb{N}} \|a_n\| < \infty$. We decompose $\mathcal{H}_X = \bigoplus_n \chi_{E_n} \mathcal{H}_X$ and define $\Lambda(g) \in \prod_{n \in \mathbb{N}} \mathfrak{B}(\chi_{E_n} \mathcal{H}_X \otimes \mathcal{H}) \subset \mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H})$ by

$$\Lambda(g) := (\text{SOT}) - \sum_{n \in \mathbb{N}} \chi_{E_n} \otimes a_n.$$

In other words, $\Lambda(g)$ is block-diagonal with respect to the decomposition $\mathcal{H}_X \otimes \mathcal{H} = \bigoplus_n ((\chi_{E_n} \mathcal{H}_X) \otimes \mathcal{H})$ and coincides with $\text{Id}_{\chi_{E_n} \mathcal{H}_X} \otimes a_n$ on each block.

Now we consider a bounded strongly measurable map $g : X \rightarrow \mathfrak{K}$. By Definition 3.2 and Proposition 3.3, there exists a sequence of countably-valued maps g_n such that $\|g_n - g\|_\infty \rightarrow 0$. Note that for each $n, m \in \mathbb{N}$, the map $g_n - g_m$ is again countably-valued, hence we have $\|\Lambda(g_n) - \Lambda(g_m)\| = \|g_n - g_m\|_\infty$. Therefore the sequence $\{\Lambda(g_n)\}_{n \in \mathbb{N}}$ is a Cauchy sequence, which converges to some $T \in \mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H})$. By a standard argument, it is easy to see that the operator T is independent of the approximating sequence $\{g_n\}_{n \in \mathbb{N}}$. This leads to the following:

Definition 3.5. Let (X, d) be a proper metric space and \mathcal{H}_X an ample X -module. Let $g : X \rightarrow \mathfrak{K}$ be a bounded strongly measurable map, and take a sequence of countably-valued maps $\{g_n\}_{n \in \mathbb{N}}$ such that $\|g_n - g\|_\infty \rightarrow 0$. We define $\Lambda(g) \in \mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H}_0)$ by $\Lambda(g) := \lim_{n \rightarrow \infty} \Lambda(g_n)$ (which is well-defined due to the above explanation). Note that we also have $\|\Lambda(g)\| = \|g\|_\infty$.

Now we are in the position to introduce strong quasi-locality for operators on proper metric spaces:

¹Note that the classic notion of strong measurability is defined for maps on a measure space, where the word “pointwise” is replaced by “almost everywhere”.

Definition 3.6. Let (X, d) be a proper metric space and \mathcal{H}_X be an ample X -module. An operator $T \in \mathfrak{B}(\mathcal{H}_X)$ is said to be *strongly quasi-local* if for any $\epsilon > 0$, there exist $\delta, R > 0$ such that for any strongly measurable map $g : X \rightarrow \mathfrak{K}_1$ with (δ, R) -variation (where \mathfrak{K}_1 denotes the closed unit ball of $\mathfrak{K}(\mathcal{H})$), we have:

$$\|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\|_{\mathfrak{B}(\mathcal{H}_X \otimes \mathcal{H})} < \epsilon.$$

It is easy to see that the set of all locally compact strongly quasi-local operators on \mathcal{H}_X forms a C^* -algebra, called the *strongly quasi-local algebra* of \mathcal{H}_X and denoted by $C_{sq}^*(\mathcal{H}_X)$.

It is straightforward to see that when X is discrete, the above definition coincides with [1, Definition 3.4]. Analogous to Roe and quasi-local algebras, it is also easy to see that the algebra $C_{sq}^*(\mathcal{H}_X)$ is *stable* in the sense that $C_{sq}^*(\mathcal{H}_X) \otimes M_n \cong C_{sq}^*(\mathcal{H}_X)$ for any $n \in \mathbb{N}$.

3.2. Properties of strongly quasi-local algebras. Recall that for a proper discrete metric space, it was shown in [1, Proposition 3.7] that the strongly quasi-local algebra sits between the Roe algebra and the quasi-local algebra. Here we prove a similar result for the general case.

Proposition 3.7. *Let X be a proper metric space and \mathcal{H}_X be an ample X -module. Then we have:*

- (1) $C_{sq}^*(\mathcal{H}_X) \subseteq C_q^*(\mathcal{H}_X)$;
- (2) *If X has bounded geometry, then $C^*(\mathcal{H}_X) \subseteq C_{sq}^*(\mathcal{H}_X)$;*
- (3) *If X has bounded geometry and Property A, then $C^*(\mathcal{H}_X) = C_{sq}^*(\mathcal{H}_X) = C_q^*(\mathcal{H}_X)$.*

Proof. To see (1), we fix a rank-one projection $p \in \mathfrak{B}(\mathcal{H})$. For any Borel function $g : X \rightarrow \mathbb{C}$ with $\|g\|_\infty = 1$, define $\tilde{g} : X \rightarrow \mathfrak{K}(\mathcal{H})_1$ by $\tilde{g}(x) := g(x)p$. Note that $[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})] = [T, g] \otimes p$. Hence the conclusion follows from the definition of strong quasi-locality together with Proposition 2.8(3).

It remains to prove (2), since (3) follows from [23, Theorem 3.3] together with (1) and (2). To see (2), we take an operator $T \in C[\mathcal{H}_X]$ with propagation R_0 . Since X has bounded geometry, there exists a Borel partition $\{E_n\}_{n \in \mathbb{N}}$ of X satisfying:

- there exists $C > 0$ such that $\text{diam}(E_n) \leq C$;
- there exists $N \in \mathbb{N}$ such that $\#\{m : d(E_n, E_m) \leq R_0\} \leq N$ for each $n \in \mathbb{N}$.

We take an $x_n \in E_n$ for each $n \in \mathbb{N}$.

Since T has propagation R_0 , we know that for each $n \in \mathbb{N}$ there are at most N -many k 's such that $\chi_{E_k} T \chi_{E_n} \neq 0$. We enumerate such k 's by $k_{1,n}, k_{2,n}, \dots, k_{N,n}$. For each $i = 1, 2, \dots, N$, we denote

$$T_i = \sum_{n=1}^{\infty} \chi_{E_{k_{i,n}}} T \chi_{E_n}.$$

Then it is clear that $T = \sum_{i=1}^N T_i$.

Now for any $\epsilon > 0$, take $\delta := \frac{\epsilon}{4N\|T\|}$ and $R := R_0 + 2C$. Then for any strongly measurable map $g : X \rightarrow \mathfrak{K}_1$ with (δ, R) -variation, we consider a countably-valued map $\tilde{g} : X \rightarrow \mathfrak{K}_1$ defined by $\tilde{g}|_{E_n} := g(x_n)$ for each $n \in \mathbb{N}$. Then $\|g - \tilde{g}\|_\infty \leq \delta$, which implies

$$\|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| \leq 2\|T\|\delta + \|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})]\|.$$

On the other hand for each $i = 1, 2, \dots, N$, we have

$$\|[T_i \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})]\| = \left\| \sum_{n=1}^{\infty} (\chi_{E_{k_{i,n}}} T \chi_{E_n}) \otimes (g(x_n) - g(x_{k_{i,n}})) \right\| \leq \sup_{n \in \mathbb{N}} \|\chi_{E_{k_{i,n}}} T \chi_{E_n}\| \cdot \delta \leq \frac{\epsilon}{2N}.$$

Therefore we obtain:

$$\|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| \leq \frac{\epsilon}{2} + \sum_{i=1}^N \|[T_i \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})]\| \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon,$$

which concludes the proof. \square

Our next goal is to explore the coarse invariance of strongly quasi-local algebras. Note that a similar result hold in the discrete case as shown in [1, Section 3.3]. While the proof is similar but more involved, we include a detailed one here for completeness.

Proposition 3.8. *Let X, Y be proper metric spaces with bounded geometry and $\mathcal{H}_X, \mathcal{H}_Y$ be ample modules for X and Y , respectively. Let $f : X \rightarrow Y$ be a coarse map with a covering isometry $V : \mathcal{H}_X \rightarrow \mathcal{H}_Y$. Then V induces the following $*$ -homomorphism*

$$\text{Ad}_V : C_{sq}^*(\mathcal{H}_X) \longrightarrow C_{sq}^*(\mathcal{H}_Y), T \mapsto VTV^*.$$

Furthermore, the induced K -theoretic map $(\text{Ad}_V)_* : K_*(C_{sq}^*(\mathcal{H}_X)) \rightarrow K_*(C_{sq}^*(\mathcal{H}_Y))$ does not depend on the choice of the covering isometry V , hence denoted by f_* .

Proof. Without loss of generality, we may assume that the coarse map f is a Borel map (see [24, Lemma A.3.12]). We only show that $VTV^* \in C_{sq}^*(\mathcal{H}_Y)$ if $T \in C_{sq}^*(\mathcal{H}_X)$. The “Furthermore” part follows a similar argument as in the case of Roe algebra, hence omitted.

First note that VTV^* is locally compact since T is, which follows the same argument as in the discrete case (see [1, Proposition 2.9]). To see that VTV^* is strongly quasi-local, we assume that $\text{supp}(V) \subseteq \{(y, x) : d_Y(f(x), y) < R_0\}$ for some $R_0 > 0$. Since Y has bounded geometry, there exists a Borel partition $\{E_n\}_{n \in \mathbb{N}}$ of Y satisfying:

- there exists $C > 0$ such that $\text{diam}(E_n) \leq C$;
- there exists $N \in \mathbb{N}$ such that $\#\{m : d_Y(E_n, E_m) \leq R_0\} \leq N$ for each $n \in \mathbb{N}$.

We take an $\gamma_n \in E_n$ for each $n \in \mathbb{N}$. Note that for each $n, m \in \mathbb{N}$, $\chi_{E_m} V \chi_{f^{-1}(E_n)} \neq 0$ implies that $d(E_m, E_n) \leq R_0$. Hence for each $n \in \mathbb{N}$, there are at most N -many m 's such that $\chi_{E_m} V \chi_{f^{-1}(E_n)} \neq 0$. Enumerate such m by $m_{1,n}, m_{2,n}, \dots, m_{N,n}$. Hence V can

be decomposed into:

$$V = W_1 + W_2 + \cdots + W_N \quad \text{where} \quad W_i = \sum_{n \in \mathbb{N}} \chi_{E_{m_{i,n}}} V \chi_{f^{-1}(E_n)}.$$

Let $M = \max_{1 \leq i \leq N} \|W_i\|$ and $\rho_+ : [0, \infty) \rightarrow [0, \infty)$ be the control function of f , i.e., $d_Y(f(x), f(y)) \leq \rho_+(d_X(x, y))$ for any $x, y \in X$. For any $\epsilon > 0$, T being strongly quasi-local implies that there exist δ', R' such that for any strongly measurable function $h : X \rightarrow \mathfrak{K}_1$ with (δ', R') -variation, we have $\|[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(h)]\| < \frac{\epsilon}{2N^2M^2}$.

We take $\delta = \min\{\frac{\epsilon}{8\|T\|}, \frac{\epsilon}{8M^2N^2\|T\|}, \frac{\delta'}{2}\}$ and $R = 2C + R_0 + \rho_+(R')$. Let $g : Y \rightarrow \mathfrak{K}_1$ be a strongly measurable map with (δ, R) -variation. Define a countably-valued map $\tilde{g} : Y \rightarrow \mathfrak{K}_1$ by setting $\tilde{g}|_{E_n} := g(\gamma_n)$, then it is clear that $\|g - \tilde{g}\|_\infty \leq \delta$. Hence:

$$\|[VTV^* \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| \leq 2\|T\|\delta + \|[VTV^* \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})]\|.$$

For each $i = 1, 2, \dots, N$, define a subset $D_i \subseteq X$ by

$$D_i := \{x \in X : \text{there exists } y \in Y \text{ such that } (y, x) \in \text{supp}(W_i)\}$$

and a map $t_i : D_i \rightarrow Y$ by setting $t_i|_{D_i \cap f^{-1}(E_n)} := \gamma_{m_{i,n}}$ for each $n \in \mathbb{N}$. Define a map $\varphi_i : X \rightarrow \mathfrak{K}_1$ by

$$\varphi_i(x) = \begin{cases} \tilde{g}(t_i(x)) & \text{if } x \in D_i; \\ 0 & \text{if } x \notin D_i. \end{cases}$$

It follows that

$$\Lambda(\tilde{g})(W_i \otimes \text{Id}_{\mathcal{H}}) = (W_i \otimes \text{Id}_{\mathcal{H}})\Lambda(\varphi_i) \quad \text{and} \quad (W_i^* \otimes \text{Id}_{\mathcal{H}})\Lambda(g) = \Lambda(\varphi_i)(W_i^* \otimes \text{Id}_{\mathcal{H}}).$$

On the other hand, for any $x \in D_i \cap f^{-1}(E_n)$ we have

$$d_Y(t_i(x), f(x)) \leq d_Y(\gamma_{m_{i,n}}, E_n) + C \leq R_0 + 2C.$$

Hence we obtain

$$\sup_{x \in D_i} \|\varphi_i(x) - \tilde{g}(f(x))\| \leq \delta,$$

which implies

$$\|(W_i \otimes \text{Id}_{\mathcal{H}})\Lambda(\varphi_i - \tilde{g} \circ f)\| \leq \delta M \quad \text{and} \quad \|\Lambda(\varphi_i - \tilde{g} \circ f)(W_i^* \otimes \text{Id}_{\mathcal{H}})\| \leq \delta M.$$

Therefore, for $i, j \in \{1, 2, \dots, N\}$ we have

$$\begin{aligned} & \|[(W_i T W_j^*) \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})]\| \\ &= \|((W_i T) \otimes \text{Id}_{\mathcal{H}})\Lambda(\varphi_j)(W_j^* \otimes \text{Id}_{\mathcal{H}}) - (W_i \otimes \text{Id}_{\mathcal{H}})\Lambda(\varphi_i)((T W_j^*) \otimes \text{Id}_{\mathcal{H}})\| \\ &\leq \|((W_i T) \otimes \text{Id}_{\mathcal{H}})\Lambda(\tilde{g} \circ f)(W_j^* \otimes \text{Id}_{\mathcal{H}}) - (W_i \otimes \text{Id}_{\mathcal{H}})\Lambda(\tilde{g} \circ f)((T W_j^*) \otimes \text{Id}_{\mathcal{H}})\| + 2M^2\|T\|\delta \\ &\leq \|(W_i \otimes \text{Id}_{\mathcal{H}})[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g} \circ f)](W_j^* \otimes \text{Id}_{\mathcal{H}})\| + \frac{\epsilon}{4N^2}. \end{aligned}$$

Note that \tilde{g} has $(2\delta, R)$ -variation on Y , hence $\tilde{g} \circ f$ has (δ', R') -variation. This implies that for $i, j \in \{1, 2, \dots, N\}$, we have

$$\|(W_i \otimes \text{Id}_{\mathcal{H}})[T \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g} \circ f)](W_j^* \otimes \text{Id}_{\mathcal{H}})\| \leq M^2 \cdot \frac{\epsilon}{2N^2M^2} = \frac{\epsilon}{2N^2}.$$

Combining the above together, we obtain

$$\begin{aligned} \|[VTV^* \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| &\leq \frac{\epsilon}{4} + \|[VTV^* \otimes \text{Id}_{\mathcal{H}}, \Lambda(\tilde{g})]\| \\ &\leq \frac{\epsilon}{4} + \sum_{i,j=1}^N \left(\frac{\epsilon}{2N^2} + \frac{\epsilon}{4N^2} \right) = \epsilon. \end{aligned}$$

Hence we conclude the proof. \square

As a direct corollary, we obtain:

Corollary 3.9. *Let \mathcal{H}_X and \mathcal{H}_Y be ample modules for proper metric spaces X and Y of bounded geometry, respectively. If X and Y are coarsely equivalent, then the strongly quasi-local algebra $C_{sq}^*(\mathcal{H}_X)$ is $*$ -isomorphic to $C_{sq}^*(\mathcal{H}_Y)$. In particular, for a proper metric space X of bounded geometry the strongly quasi-local algebra $C_{sq}^*(\mathcal{H}_X)$ does not depend on the chosen ample X -module \mathcal{H}_X up to $*$ -isomorphisms, hence called the strongly quasi-local algebra of X and denoted by $C_{sq}^*(X)$.*

4. LOCALISATION ALGEBRAS FOR (STRONG) QUASI-LOCALITY

In this section, we use the notion of strong quasi-locality for proper metric spaces introduced in Section 3 to provide an alternative version of K -homology. Recall that Yu introduced the localisation algebra for a proper metric space [27] and showed that its K -theory is isomorphic to the K -homology of the underlying space (see also [17]). Our idea is to provide a (strongly) quasi-local version of Yu's localisation algebra and show that they have the same K -theory (Theorem D).

First let us recall Yu's localisation algebra.

Definition 4.1 ([27]). Let (X, d) be a proper metric space and \mathcal{H}_X be an ample X -module. The *localisation algebra* $C_L^*(\mathcal{H}_X)$ is defined to be the norm closure of the $*$ -algebra of all bounded uniformly continuous map $T : [0, \infty) \rightarrow C^*(\mathcal{H}_X)$ satisfying $\lim_{t \rightarrow \infty} \text{prop}(T_t) \rightarrow 0$ (here we write T_t instead of $T(t)$), with respect to the supremum norm $\|T\|_\infty := \sup_{t \in [0, \infty)} \|T_t\|$. It is shown in [27] that $C_L^*(\mathcal{H}_X)$ is independent of the module \mathcal{H}_X , hence denoted by $C_L^*(X)$.

The evaluation map $e : C_L^*(X) \rightarrow C^*(X)$ is defined by $e(g) = g(0)$ for $g \in C_L^*(X)$. One can also define a local assembly map μ_L [27] from the K -homology of X to the K -theory of the localisation algebra:

$$\mu_L : K_*(X) \longrightarrow K_*(C_L^*(X)).$$

Proposition 4.2 ([17, 27]). *Let X be a proper metric space. Then the local assembly map μ_L is an isomorphism.*

Consequently, for a proper metric space X with bounded geometry, we have the following commutative diagram:

$$\begin{array}{ccc}
& \lim_{d \rightarrow \infty} K_*(C_L^*(P_d(X))) & \\
\mu_L \swarrow & \nearrow e_* & \\
\lim_{d \rightarrow \infty} K_*(P_d(X)) & \xrightarrow{\mu} & \lim_{d \rightarrow \infty} K_*(C^*(P_d(X)))
\end{array}$$

Therefore, the coarse Novikov conjecture is equivalent to that e_* is injective and the coarse Baum-Connes conjecture is equivalent to that e_* is bijective.

Now we introduce a (strongly) quasi-local version of Definition 4.1.

Definition 4.3. Let X be a proper metric space and \mathcal{H}_X be an ample X -module. We define the *localisation quasi-local algebra* $C_{L,q}^*(\mathcal{H}_X)$ to be the C^* -algebra consisting of all bounded uniformly continuous map $T : [0, \infty) \rightarrow C_q^*(\mathcal{H}_X)$ satisfying that for any $\epsilon > 0, R > 0$ there is a $t_0 > 0$ such that for any $t > t_0$ and Borel subsets $C, D \subseteq X$ with $d(C, D) > R$, we have $\|\chi_C T_t \chi_D\| < \epsilon$.

The intuition behind the above notion is that the “extent” of quasi-locality of T_t gets better and better as $t \rightarrow \infty$. Hence it is easy to see that $C_L^*(\mathcal{H}_X) \subseteq C_{L,q}^*(\mathcal{H}_X)$. The following lemma follows from the proof of [22, Theorem 2.8, “(i) \Leftrightarrow (ii)”] and Proposition 2.8, hence we omit the proof.

Lemma 4.4. Let X be a proper metric space and \mathcal{H}_X an ample X -module. Then a bounded uniformly continuous map $T : [0, \infty) \rightarrow C_q^*(\mathcal{H}_X)$ belongs to $C_{L,q}^*(\mathcal{H}_X)$ if and only if for any $\epsilon > 0$ there exists $\delta > 0$ such that for any $R > 0$ there exists $t_0 > 0$ satisfying the following: for any $t > t_0$ and Borel function $g : X \rightarrow \mathbb{C}$ with norm 1 and (δ, R) -variation, we have $\|[T_t, g]\| < \epsilon$.

The above lemma inspires us to introduce the following:

Definition 4.5. Let X be a proper metric space and \mathcal{H}_X an ample X -module. The *localisation strongly quasi-local algebra* $C_{L,sq}^*(\mathcal{H}_X)$ is defined to be the C^* -algebra consisting of all bounded uniformly continuous map $T : [0, \infty) \rightarrow C_{sq}^*(\mathcal{H}_X)$ satisfying that for any $\epsilon > 0$ there exists $\delta > 0$ such that for any $R > 0$ there exists $t_0 > 0$ for any $t > t_0$ and strongly measurable function $g : X \rightarrow \mathfrak{K}_1$ with (δ, R) -variation, we have $\|[T_t \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| < \epsilon$.

Following similar arguments as in the proof of [27, Proposition 3.7] with slight changes from those of [1, Proposition 2.9] and Proposition 3.8, we can show that both $C_{L,q}^*(\mathcal{H}_X)$ and $C_{L,sq}^*(\mathcal{H}_X)$ do not depend on the ample module \mathcal{H}_X , hence denoted by $C_{L,q}^*(X)$ and $C_{L,sq}^*(X)$, respectively. We omit the details here. Moreover, from a similar argument as in the proof of Proposition 3.7, we have:

Lemma 4.6. Let X be a proper metric space, then we have:

$$C_L^*(X) \subseteq C_{L,sq}^*(X) \subseteq C_{L,q}^*(X).$$

Note that the above three localisation algebras are closely related to another algebra, which comes from an “almost commutant picture” of K -homology introduced in [5]:

Definition 4.7 ([5]). Let X be a proper metric space and \mathcal{H}_X an ample X -module. We define $C_L(C_0(X))$ to be the C^* -algebra consisting of all bounded uniformly continuous map $T : [0, \infty) \rightarrow \mathfrak{B}(\mathcal{H}_X)$ satisfying

- (1) for any $t \in [0, \infty)$ and $f \in C_0(X)$, we have $T_t f, f T_t \in \mathfrak{K}(\mathcal{H}_X)$;
- (2) for any $f \in C_0(X)$, we have $\lim_{t \rightarrow \infty} \|[T_t, f]\| = 0$.

As pointed out in [5], it is a routine work to check that the C^* -algebra $C_L(C_0(X))$ does not depend on the module \mathcal{H}_X . The following lemma relates our localisation (strongly) quasi-local algebras with $C_L(C_0(X))$:

Lemma 4.8. $C_L^*(X) \subseteq C_{L,sq}^*(X) \subseteq C_{L,q}^*(X) \subseteq C_L(C_0(X))$.

Proof. From Lemma 4.6, it remains to show that $C_{L,q}^*(\mathcal{H}_X) \subseteq C_L(C_0(X))$. Let $T = (T_t) \in C_{L,q}^*(\mathcal{H}_X)$, it suffices to check that $\lim_{t \rightarrow \infty} \|[T_t, f]\| = 0$ for any real-valued non-negative function $f \in C_0(X)$ with $\|f\|_\infty \leq 1$.

Fix such an $f \in C_0(X)$. For any $\epsilon > 0$, it follows from Lemma 4.4 that there exists $\delta > 0$ such that for any $R > 0$ there exists $t_0 > 0$ satisfying the following: for any $t > t_0$ and Borel function $g : X \rightarrow \mathbb{C}$ with norm 1 and (δ, R) -variation, we have $\|[T_t, g]\| < \epsilon$.

Note that f is uniformly continuous, hence for the above δ there exists an $R > 0$ such that f has (δ, R) -variation. Therefore, the above paragraph provides a t_0 such that for any $t > t_0$ we have $\|[T_t, f]\| < \epsilon$. This concludes the proof. \square

As a special case of the main result in [5], we know that $K_*(C_L(C_0(X))) \cong K_*(X)$. This inspires the following theorem, which is the main result of this section and implies Theorem D. It shows that the above three localisation algebras have the same K -theories, hence any of them can serve as the K -homology of X by Proposition 4.2.

Theorem 4.9. *Let X be a proper metric space, then the inclusion maps induce isomorphisms on K -theories:*

$$K_*(C_L^*(X)) \cong K_*(C_{L,sq}^*(X)) \cong K_*(C_{L,q}^*(X)) \cong K_*(C_L(C_0(X))).$$

To achieve, we consider a two-sided $*$ -ideal $C_{L,0}(C_0(X))$ in $C_L(C_0(X))$ consisting of maps $T = (T_t) \in C_L^*(X)$ such that for any compact $K \subseteq X$, we have:

$$\lim_{t \rightarrow \infty} \chi_K T_t = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} T_t \chi_K = 0.$$

Moreover, we consider $C_{L,0}^*(X) := C_{L,0}(C_0(X)) \cap C_L^*(X)$, $C_{L,sq,0}^*(X) := C_{L,0}(C_0(X)) \cap C_{L,sq}^*(X)$ and $C_{L,q,0}^*(X) := C_{L,0}(C_0(X)) \cap C_{L,q}^*(X)$. They are two-sided $*$ -ideals in $C_L^*(X)$, $C_{L,sq}^*(X)$ and $C_{L,q}^*(X)$, respectively.

The following is a crucial step to achieve Theorem 4.9:

Proposition 4.10. *Let X be a proper metric space. Then the inclusions from Lemma 4.8 induces the following isomorphisms:*

$$\frac{C_L^*(X)}{C_{L,0}^*(X)} \cong \frac{C_{L,sq}^*(X)}{C_{L,sq,0}^*(X)} \cong \frac{C_{L,q}^*(X)}{C_{L,q,0}^*(X)} \cong \frac{C_L(C_0(X))}{C_{L,0}(C_0(X))}.$$

Proof. It is clear that the inclusions induce the following injections:

$$\frac{C_L^*(X)}{C_{L,0}^*(X)} \hookrightarrow \frac{C_{L,sq}^*(X)}{C_{L,sq,0}^*(X)} \hookrightarrow \frac{C_{L,q}^*(X)}{C_{L,q,0}^*(X)} \hookrightarrow \frac{C_L(C_0(X))}{C_{L,0}(C_0(X))}.$$

Hence it suffices to show that for any $T = (T_t) \in C_L(C_0(X))$, there exists $S = (S_t) \in C_L^*(X)$ such that $T - S \in C_{L,0}(C_0(X))$.

For each $n \in \mathbb{N}$, let $\{U_i^{(n)}\}_{i \in \mathbb{N}}$ be a locally finite open cover of X such that $\text{diam}(U_i^{(n)}) < 1/n$, and let $\{\phi_i^{(n)}\}_{i \in \mathbb{N}}$ be a partition of unity subordinate to $\{U_i^{(n)}\}_{i \in \mathbb{N}}$. We define

$$S_{n,t} = \sum_{i \in \mathbb{N}} \sqrt{\phi_i^{(n)}} T_t \sqrt{\phi_i^{(n)}},$$

where the sum converges in the strong operator topology. It is clear that $S_{n,t}$ has propagation at most $1/n$.

Since X is proper, we can take a sequence of compact subsets $\{K_m\}_{m \in \mathbb{N}}$ such that each K_m is a closed ball, $K_m \subseteq K_{m+1}$ and $X = \bigcup_{m \in \mathbb{N}} K_m$. For each $m, n \in \mathbb{N}$ we have:

$$\chi_{K_m}(S_{n,t} - T_t) = \sum_{i \in \mathbb{N}} \chi_{K_m} \sqrt{\phi_i^{(n)}} [T_t, \sqrt{\phi_i^{(n)}}].$$

Since K_m is compact, we know that there are only finitely many non-zero items in the above summation. Combining with the assumption that $(T_t) \in C_L(C_0(X))$, we obtain that $\chi_{K_m}(S_{n,t} - T_t)$ tends to zero as $t \rightarrow \infty$ for each $m, n \in \mathbb{N}$. Hence for each $n \in \mathbb{N}$, there is a t_n such that for any $t \geq t_n$ we have

$$\|\chi_{K_n}(S_{n,t} - T_t)\| < 1/n \quad \text{and} \quad \|\chi_{K_n}(S_{n+1,t} - T_t)\| < 1/n.$$

Set $t_0 = 0$ and we assume that t_n is monotonically increasing to infinity. Let $\{\psi_n\}_{n \in \mathbb{N}}$ be a partition of unity of $[0, +\infty)$ such that the $\text{supp}(\psi_n) \subseteq [t_n, t_{n+2}]$. Define

$$S_t := \sum_{m=0}^{\infty} \psi_m(t) S_{m+1,t}$$

for each $t \in [0, \infty)$. It is easy to check that $S := (S_t)$ belongs to $C_L^*(X)$.

Now we claim that $T - S \in C_{L,0}(C_0(X))$. Fixing a compact subset $K \subseteq X$, there is an $m_0 \in \mathbb{N}$ such that $K \subseteq K_n$ for any $n \geq m_0$. Then for any $n > m_0$ and $t \in [t_n, t_{n+1}]$, we have

$$\begin{aligned} \chi_K(T_t - S_t) &= \chi_K \chi_{K_n}(T_t - S_t) \\ &= \chi_K \chi_{K_n} \left(\sum_{m=0}^{\infty} \psi_m(t) T_t - \sum_{m=0}^{\infty} \psi_m(t) S_{m+1,t} \right) \\ &= \chi_K \chi_{K_n} \left(\psi_{n-1}(t) \cdot (T_t - S_{n,t}) + \psi_n(t) \cdot (T_t - S_{n+1,t}) \right). \end{aligned}$$

According to the choice of t_n , we have $\|\chi_K(T_t - S_t)\| \leq 2/n$. Similarly, we obtain that $\|(T_t - S_t)\chi_K\| \leq 2/n$. Hence we conclude the proof. \square

We also need the following lemma:

Lemma 4.11. *Let X be a proper metric space, and \mathcal{H}_X be an ample X -module. Then we have:*

$$K_*(C_{L,0}^*(\mathcal{H}_X)) = K_*(C_{L,sq,0}^*(\mathcal{H}_X)) = K_*(C_{L,q,0}^*(\mathcal{H}_X)) = K_*(C_{L,0}(C_0(X))) = 0.$$

Proof. The proof follows from a standard Eilenberg Swindle argument (see, e.g., [24, Lemma 6.4.11]), hence omitted. \square

Proof of Theorem 4.9. The proof follows easily from Proposition 4.10, Lemma 4.11 together with associated six-term short exact sequences. We omit the details. \square

Now we consider the following commutative diagram:

$$\begin{array}{ccc} \lim_{d \rightarrow \infty} K_*(C_L^*(P_d(X))) & \xrightarrow{i_*} & \lim_{d \rightarrow \infty} K_*(C_{L,sq}^*(P_d(X))) \\ \mu_L \nearrow & e_* & \searrow (e_{sq})_* \\ \lim_{d \rightarrow \infty} K_*(P_d(X)) & \xrightarrow{\mu} & \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) & \xrightarrow{i_*} & \lim_{d \rightarrow \infty} K_*(C_{sq}^*(P_d(X))), \end{array}$$

where i_* are induced by inclusions. Recall that the strongly quasi-local coarse assembly map μ_{sq} is defined to be the composition

$$\mu_{sq} : \lim_{d \rightarrow \infty} K_*(P_d(X)) \xrightarrow{\mu} \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) \xrightarrow{i_*} \lim_{d \rightarrow \infty} K_*(C_{sq}^*(P_d(X))) \cong K_*(C_{sq}^*(X)),$$

where the last isomorphism is from Corollary 3.9. The strongly quasi-local coarse Novikov/Baum-Connes conjecture asserts that μ_{sq} is injective/bijective.

From Proposition 4.2 and Theorem 4.9, we know that the composition

$$\mu_{L,sq} := i_* \circ \mu_L : \lim_{d \rightarrow \infty} K_*(P_d(X)) \longrightarrow \lim_{d \rightarrow \infty} K_*(C_L^*(P_d(X))) \longrightarrow \lim_{d \rightarrow \infty} K_*(C_{L,sq}^*(P_d(X)))$$

is an isomorphism. Consequently, we obtain the following:

Corollary 4.12. *Let X be a discrete metric space with bounded geometry. Then the strongly quasi-local coarse Baum-Connes/Novikov conjecture holds for X if and only if the following homomorphism induced by evaluation at 0:*

$$(e_{sq})_* : \lim_{d \rightarrow \infty} K_*(C_{L,sq}^*(P_d(X))) \longrightarrow \lim_{d \rightarrow \infty} K_*(C_{sq}^*(P_d(X)))$$

is bijective/injective.

We can similarly define a quasi-local coarse assembly map

$$\mu_{L,q} : \lim_{d \rightarrow \infty} K_*(P_d(X)) \longrightarrow K_*(C_q^*(X))$$

and consider the corresponding quasi-local coarse Baum-Connes/Novikov conjecture. Moreover, according to Theorem 4.9 again, we also have an analogous result to Corollary 4.12 for quasi-local algebras.

5. TWISTED ALGEBRAS

The notion of twisted Roe algebras for metric spaces which admit a coarse embedding into a Banach space with Property (H) were introduced in [4]. They play a key role in the proof of the coarse Novikov conjecture for such spaces ([4, Theorem 1.1]). In this section, we study a modified version of the twisted Roe algebras and introduce their quasi-local counterparts. These algebras will provide a bridge to link the K -theories of Roe algebras and strongly quasi-local ones.

Let us fix some notation following [4]. Throughout the section, let X be a discrete metric space with bounded geometry which admits a coarse embedding $f : X \rightarrow V$ into a real Banach space V with Property (H). From Definition A, there exist increasing sequences of finite dimensional subspaces $\{V_n\}_{n \in \mathbb{N}}$ of V and $\{W_n\}_{n \in \mathbb{N}}$ of a real Hilbert space W such that:

- (1) each V_n and W_n are even dimensional, and $\bigcup_{n=1}^{\infty} V_n$ is dense in V ;
- (2) there exists a uniformly continuous map $\psi : S(\bigcup_{n=1}^{\infty} V_n) \rightarrow S(\bigcup_{n=1}^{\infty} W_n)$ such that $\psi|_{S(V_n)}$ is a homeomorphism onto $S(W_n)$ for any $n \in \mathbb{N}$.

By a slight modification, we may assume without loss of generality that

$$f(X) \subset \bigcup_{n=1}^{\infty} V_n.$$

For each $d \geq 0$, the coarse embedding $f : X \rightarrow \bigcup_{n=1}^{\infty} V_n$ can be extended to a continuous coarse embedding $f : P_d(X) \rightarrow \bigcup_{n=1}^{\infty} V_n$ by affine extension (see [4, Section 3.1]). We also take a countable dense subset $Z_d \subset P_d(X)$ such that $Z_d \subseteq Z_{d'}$ whenever $d < d'$.

For each $n \in \mathbb{N}$, let $\text{Cliff}(W_n)$ be the complex Clifford algebra of W_n with respect to the relation $w^2 = \|w\|^2$ for all $w \in W_n$. Denote

$$\mathcal{A}_n := C_0(V_n) \otimes \text{Cliff}(W_n) \cong C_0(V_n, \text{Cliff}(W_n)).$$

We follow the notation from Section 3 that $\mathfrak{K} = \mathfrak{K}(\mathcal{H})$ denotes the C^* -algebra of compact operators on the fixed infinite dimensional separable Hilbert space \mathcal{H} . For a function $h \in \mathcal{A}_n \otimes \mathfrak{K} \cong C_0(V_n, \text{Cliff}(W_n) \otimes \mathfrak{K})$, we define the support of h to be

$$\text{supp}(h) = \overline{\{x \in V_n : h(x) \neq 0\}}.$$

Now we consider two C^* -algebras which serve as coefficients for twisted algebras. Let $\prod_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K})$ be the C^* -algebra direct product and $\bigoplus_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K})$ be the C^* -algebra direct sum. Denote the quotient algebra by

$$\mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_{n \in \mathbb{N}}) := \frac{\prod_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K})}{\bigoplus_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K})},$$

where the norm of $[(h_1, \dots, h_n, \dots)] \in \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_{n \in \mathbb{N}})$ can be calculated by:

$$\|[(h_1, \dots, h_n, \dots)]\| = \limsup_{n \rightarrow \infty} \|h_n\|_{\mathcal{A}_n \otimes \mathfrak{K}}.$$

Also denote the C^* -algebra

$$Q((\mathcal{A}_n)_{n \in \mathbb{N}}) := \frac{\prod_{n=1}^{\infty} \mathcal{A}_n}{\bigoplus_{n=1}^{\infty} \mathcal{A}_n}.$$

Since \mathfrak{K} is nuclear, we have:

$$(5.1) \quad Q((\mathcal{A}_n)_{n \in \mathbb{N}}) \otimes \mathfrak{K} \cong \frac{(\prod_{n=1}^{\infty} \mathcal{A}_n) \otimes \mathfrak{K}}{(\bigoplus_{n=1}^{\infty} \mathcal{A}_n) \otimes \mathfrak{K}} \hookrightarrow \frac{\prod_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K})}{\bigoplus_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K})} = Q((\mathcal{A}_n \otimes \mathfrak{K})_n),$$

where the second inclusion is induced by the natural inclusion

$$\left(\prod_{n=1}^{\infty} \mathcal{A}_n \right) \otimes \mathfrak{K} \hookrightarrow \prod_{n=1}^{\infty} (\mathcal{A}_n \otimes \mathfrak{K}).$$

Hence in the sequel, we regard $Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}$ as a C^* -subalgebra of $Q((\mathcal{A}_n \otimes \mathfrak{K})_n)$ via (5.1). Also for short we write $Q((\mathcal{A}_n \otimes \mathfrak{K})_n)$ and $Q((\mathcal{A}_n)_n)$ instead of $Q((\mathcal{A}_n \otimes \mathfrak{K})_{n \in \mathbb{N}})$ and $Q((\mathcal{A}_n)_{n \in \mathbb{N}})$, respectively.

5.1. Preliminaries on Hilbert modules. To define twisted algebras, we need to recall some facts from the theory of Hilbert module. Readers can refer to standard textbooks (e.g., [14]) for basic notions and more details.

Let Z be a countable set and A be a C^* -algebra. We denote a map $Z \rightarrow A$ which maps $x \mapsto a_x$ by $\sum_{x \in Z} a_x[x]$. Let

$$\ell^2(Z; A) := \left\{ \sum_{x \in Z} a_x[x] : a_x \in A \text{ and } \sum_{x \in Z} a_x^* a_x \text{ converges in norm} \right\}.$$

Then $\ell^2(Z; A)$ is a Hilbert A -module induced by:

$$\begin{aligned} \left\langle \sum_{x \in Z} a_x[x], \sum_{x \in Z} b_x[x] \right\rangle &:= \sum_{x \in Z} a_x^* b_x, \\ \left(\sum_{x \in Z} a_x[x] \right) \cdot a &:= \sum_{x \in Z} (a_x a)[x] \end{aligned}$$

for any $a \in A$. Let $\mathfrak{B}(\ell^2(Z; A))$ be the C^* -algebra of all module homomorphisms from $\ell^2(Z; A)$ to itself for which there exists an adjoint module homomorphism, and $\mathfrak{K}(\ell^2(Z; A))$ be the norm closure of all finite-rank operators in $\mathfrak{B}(\ell^2(Z; A))$. Note that $\mathfrak{K}(\ell^2(Z; A))$ is a $*$ -ideal in $\mathfrak{B}(\ell^2(Z; A))$.

Moreover, note that there is a $*$ -representation $\rho : \ell^\infty(Z) \rightarrow \mathfrak{B}(\ell^2(Z; A))$ by pointwise scalar multiplication. For abbreviation, we will write fT instead of $\rho(f)T$ for $f \in \ell^\infty(Z)$ and $T \in \mathfrak{B}(\ell^2(Z; A))$.

For each $T \in \mathfrak{B}(\ell^2(Z; A))$, we can associate a matrix form $(T_{xy})_{x, y \in Z}$ as follows. For $x, y \in Z$, we define a linear map $T_{xy} : A \rightarrow A$ by

$$T_{xy}(a) := (T(a[y]))(x) \text{ for } a \in A,$$

called the xy -matrix entry of T . It is easy to check that $\|T_{xy}\|_{\mathfrak{B}(A)} = \|\chi_{\{x\}} T \chi_{\{y\}}\|_{\mathfrak{B}(\ell^2(Z; A))}$, and each T_{xy} is indeed in the multiplier algebra $\mathcal{M}(A)$. Furthermore, it follows from $T \in \mathfrak{B}(\ell^2(Z; A))$ that T can be recovered by its matrix form via matrix multiplication. More precisely, for each $\sum_{x \in Z} a_x[x] \in \ell^2(Z; A)$ we have:

- (1) for $x \in Z$, the sum $\sum_{y \in Z} T_{xy}(a_y)$ converges in norm;
- (2) the map $x \mapsto \sum_{y \in Z} T_{xy}(a_y)$ belongs to $\ell^2(Z; A)$;
- (3) $T(\sum_{x \in Z} a_x[x]) = \sum_{x \in Z} (\sum_{y \in Z} T_{xy}(a_y))[x]$.

We need the following auxiliary lemma to define the Bott maps in Section 6.2:

Lemma 5.1. *Let (π, \mathcal{H}) be a faithful representation of a C^* -algebra A (where the Hilbert space \mathcal{H} is not necessarily separable), and Z be a countable set. For any $T \in \mathfrak{B}(\ell^2(Z; \mathcal{H}))$, we write its matrix form as $T = (T_{xy})_{x,y \in Z}$ where $T_{xy} \in \mathfrak{B}(\mathcal{H})$. Assume that each $T_{xy} \in \mathcal{M}(A)$. Define a linear operator $\Psi(T)$ on the Hilbert module $\ell^2(Z; A)$ by:*

$$(5.2) \quad \Psi(T) \left(\sum_{x \in Z} a_x[x] \right) := \sum_{x \in Z} \left(\sum_{y \in Z} T_{xy}(a_y) \right) [x]$$

for any $\sum_{x \in Z} a_x[x] \in \ell^2(Z; A)$. Then $\Psi(T)$ is well-defined and belongs to $\mathfrak{B}(\ell^2(Z; A))$. Moreover, we have $\|\Psi(T)\| \leq \|T\|$ and $T_{xy} = \Psi(T)_{xy}$ for any $x, y \in Z$.

Proof. For any finite subset $F \subset Z$ and $\sum_{x \in F} a_x[x] \in \ell^2(Z; A)$ with $\|\sum_{x \in F} a_x^* a_x\| = 1$, we consider a vector $b \in \ell^2(F, A)$ by $b_x := \sum_{y \in F} T_{xy} a_y$ for $x \in F$. Then we have:

$$\left\| \sum_{x \in F} b_x[x] \right\|^2 = \left\| \sum_{x \in F} b_x^* b_x \right\| = \sup_{\|\xi\|=1} \left\langle \sum_{x \in F} b_x^* b_x \xi, \xi \right\rangle = \sup_{\|\xi\|=1} \sum_{x \in F} \|b_x \xi\|^2.$$

For any $\xi \in \mathcal{H}$ with $\|\xi\| = 1$, we set $\xi' = \sum_{x \in F} (a_x \xi)[x] \in \ell^2(Z; \mathcal{H})$. Then:

$$\|\xi'\|^2 = \sum_{x \in F} \|a_x \xi\|^2 = \left\langle \left(\sum_{x \in F} a_x^* a_x \right) \xi, \xi \right\rangle \leq \left\| \sum_{x \in F} a_x^* a_x \right\| \cdot \|\xi\|^2 = 1,$$

and hence

$$\sum_{x \in F} \|b_x \xi\|^2 = \left\| \sum_{x \in F} (b_x \xi)[x] \right\|^2 = \|T \xi'\|^2 \leq \|T\|^2 \cdot \|\xi'\|^2 \leq \|T\|^2.$$

This concludes the proof. \square

5.2. Twisted Roe algebras. Here we recall the original definition of twisted Roe algebras from [4, Section 3.1]. We also introduce a restricted version and study their relations. This will play a key role in the sequel to study the Bott map for strongly quasi-local algebras and to prove the main theorem. Let us fix a $d \geq 0$, and recall that Z_d is a countable dense subset in the Rips complex $P_d(X)$.

Definition 5.2 ([4, Definition 3.1]). Define $\mathbb{C}[P_d(X), Q((\mathcal{A}_n \otimes \mathcal{R})_n)]$ to be the set of bounded functions $T : Z_d \times Z_d \rightarrow Q((\mathcal{A}_n \otimes \mathcal{R})_n)$ satisfying the following conditions:

- (1) for any bounded subset $B \subseteq X$, the set $\{(x, y) \in (B \times B) \cap (Z_d \times Z_d) : T(x, y) \neq 0\}$ is finite;
- (2) there exists an $L > 0$ such that $\#\{x \in Z_d : T(x, y) \neq 0\} < L$ and $\#\{y \in Z_d : T(x, y) \neq 0\} < L$ for each $x, y \in Z_d$;
- (3) there exists an $R \geq 0$ such that $T(x, y) = 0$ whenever $d(x, y) > R$ for $x, y \in Z_d$;
- (4) there exists an $r > 0$ such that for any $x, y \in Z_d$, $T(x, y)$ is of the form $[(h_1, \dots, h_n, \dots)]$ satisfying $\text{supp}(h_n) \subseteq B_{V_n}(f(x), r)$ for $n \in \mathbb{N}$ large enough such that $f(x) \in V_n$.

The algebraic structure for $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$ is defined by regarding elements T as $Z_d \times Z_d$ -matrices. The $*$ -structure for $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$ is defined by

$$(T^*)(x, y) := [(h_1^*, \dots, h_n^*, \dots)]$$

where $T(y, x) = [(h_1, \dots, h_n, \dots)]$ for all $x, y \in Z_d$.

Denote by E_1 the Hilbert module $\ell^2(Z_d; \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n))$. It is clear that the $*$ -algebra $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$ has an action on E_1 by the formula

$$(5.3) \quad T\left(\sum_{x \in Z_d} a_x[x]\right) = \sum_{x \in Z_d} \left(\sum_{y \in Z_d} T(x, y)a_y\right)[x]$$

for $T \in \mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$ and $\sum_{x \in Z_d} a_x[x] \in \ell^2(Z_d; \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n))$. Also note that T is an adjointable module homomorphism, hence defines an element in $\mathfrak{B}(E_1)$. It is easy to check that the xy -matrix entry of $T \in \mathfrak{B}(E_1)$ coincides with $T(x, y)$.

The following is the twisted Roe algebra introduced in [4, Definition 3.2]:

Definition 5.3 ([4]). The *twisted Roe algebra* $C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n))$ is defined to be the norm closure of $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$ in $\mathfrak{B}(E_1)$.

Now we introduce a modified version of the twisted Roe algebra. The idea is to restrict the coefficient algebra $\mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)$ to the subalgebra $\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$.

Definition 5.4. Define $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ to be the $*$ -algebra consisting of all bounded functions $T : Z_d \times Z_d \rightarrow \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$ satisfying condition (1) ~ (4) in Definition 5.2.

Note that the only difference between the $*$ -algebras $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$ and $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ is the requirement on ranges of their elements. It is also clear that $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ is a $*$ -subalgebra of $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)]$.

We denote by E the Hilbert module $\ell^2(Z_d; \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$. Note that the algebra $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ has an action on E by the same formula (5.3) and similarly, $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ can be regarded as a subalgebra in $\mathfrak{B}(E)$. This leads to the following:

Definition 5.5. The *restricted twisted Roe algebra* $C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ is defined to be the norm closure of $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ in $\mathfrak{B}(E)$.

Remark 5.6. The motivation to introduce the restricted version of the twisted Roe algebra is to pave the way for the Bott maps constructed in Section 6. We will see that the Bott map can be defined for strongly quasi-local algebras using the smaller coefficient algebra $\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$, while it is unclear to us whether this can be done in terms of the original $\mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)$. See Remark 6.2 for more details.

To study the relation between the two twisted Roe algebras, we need an auxiliary algebra which is a subalgebra in $C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n))$. Note that the

$*$ -algebra $\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ also has an action on the module E_1 by the same formula (5.3), hence we can consider its norm closure in $\mathfrak{B}(E_1)$:

$$\mathfrak{A} := \overline{\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]}^{\|\cdot\|_{\mathfrak{B}(E_1)}}.$$

It is clear that \mathfrak{A} is a C^* -subalgebra of $C^*(P_d(X), Q((\mathcal{A}_n \otimes \mathfrak{K})_n))$. The following lemma shows that \mathfrak{A} is isomorphic to the restricted twisted Roe algebra, hence builds a bridge between $C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ and $C^*(P_d(X), Q((\mathcal{A}_n \otimes \mathfrak{K})_n))$.

Lemma 5.7. *The identity map on $\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ can be extended continuously to a $*$ -isomorphism from $C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ to \mathfrak{A} . In particular, we have $\|T\|_{\mathfrak{B}(E)} = \|T\|_{\mathfrak{B}(E_1)}$ for any $T \in \mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$.*

Proof. To prove the lemma, we consider the restriction map as follows. For any $T \in \mathfrak{A}$, we claim that $T(E) \subseteq E$. In fact, assume that there exists a sequence $\{T_n\}_n$ in $\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ converging to T in $\mathfrak{B}(E_1)$. It is clear that $T_n(E) \subseteq E$ for each n . Hence for any $\xi \in E \subseteq E_1$, we have $T\xi = \lim_{n \rightarrow \infty} T_n(\xi) \in E$ since E is closed. This leads to a C^* -homomorphism

$$\Phi : \mathfrak{A} \rightarrow C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}), \quad T \mapsto T|_E.$$

It is clear that Φ is surjective since its image contains the dense subalgebra $\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$, hence it suffices to show that Φ is injective. Assume that $\Phi(T) = 0$ for some $T \in \mathfrak{A}$, and take a sequence $\{T_n\}_n$ in $\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ converging to T in $\mathfrak{B}(E_1)$. For any $x, y \in Z_d$, it is clear that the matrix entry $(T_n)_{xy}$ converges to T_{xy} in $\mathcal{M}(Q((\mathcal{A}_n \otimes \mathfrak{K})_n))$. Since each $(T_n)_{xy} \in Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}$, which is regarded as an element in $\mathcal{M}(Q((\mathcal{A}_n \otimes \mathfrak{K})_n))$ via the inclusion $i : Q((\mathcal{A}_n)_n) \otimes \mathfrak{K} \rightarrow \mathcal{M}(Q((\mathcal{A}_n \otimes \mathfrak{K})_n))$, we obtain that T_{xy} also belongs to the image of i . Hence $T_{xy}(Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}) = 0$ implies that $T_{xy} = 0$, and therefore we obtain that $T = 0$. \square

As a direct corollary, we obtain the following:

Corollary 5.8. *There is a C^* -monomorphism*

$$\iota : C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}) \rightarrow C^*(P_d(X), Q((\mathcal{A}_n \otimes \mathfrak{K})_n))$$

induced by the inclusion $\mathbb{C}[P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}] \hookrightarrow \mathbb{C}[P_d(X), Q((\mathcal{A}_n \otimes \mathfrak{K})_n)]$.

5.3. Twisted quasi-local algebras. Now we introduce a quasi-local version of the twisted algebras. Due to the lack of control on propagations, we need a slightly different approach.

First note that any continuous bounded function $h : V \rightarrow \mathbb{C}$ can be regarded as an element in the multiplier algebra $\mathcal{M}(Q((\mathcal{A}_n)_n))$ by

$$h \cdot [(h_1, \dots, h_n, \dots)] = [(h|_{V_1} \cdot h_1, \dots, h|_{V_n} \cdot h_n, \dots)],$$

where $[(h_1, \dots, h_n, \dots)] \in Q((\mathcal{A}_n)_n)$. Hence we can also regard h as an element in $\mathfrak{B}(E)$ by setting its matrix entry to be:

$$h_{xy} := \begin{cases} h \otimes \text{Id}_{\mathcal{H}} (\in \mathcal{M}(Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})), & x = y; \\ 0, & \text{otherwise.} \end{cases}$$

Definition 5.9. The *twisted quasi-local algebra* $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ is defined to be the subset in $\mathfrak{B}(E)$ consisting of elements T satisfying the following conditions:

- (1) each matrix entry of T belongs to $\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$;
- (2) for any compact $K \subseteq P_d(X)$, we have that $\chi_K T$ and $T \chi_K$ belongs to $\mathfrak{K}(E)$;
- (3) for any $\epsilon > 0$ there is an r_1 such that for any Borel subsets $C, D \subseteq P_d(X)$ with $d(C, D) > r_1$, we have $\|\chi_C T \chi_D\| < \epsilon$ and $\|\chi_D T \chi_C\| < \epsilon$;
- (4) for any $\epsilon > 0$, there is an r_2 such that for any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(V)_1$ with $d(f(C), \text{supp}(h)) > r_2$, we have $\|\chi_C T h\| < \epsilon$.

Remark 5.10. It follows from condition (3) and (4) in Definition 5.9 that for any $T \in C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ and $\epsilon > 0$, there exists $r'_2 > 0$ such that for any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(V)_1$ with $d(f(C), \text{supp}(h)) > r'_2$, we have $\|h T \chi_C\| < \epsilon$. In fact, assume that $\rho_+ : [0, \infty) \rightarrow [0, \infty)$ is a proper function such that $\|f(x) - f(y)\| \leq \rho_+(d(x, y))$ for any $x, y \in P_d(X)$. For the given $\epsilon > 0$, there exist $r_1, r_2 > 0$ satisfying condition (3) and (4) therein for $\epsilon/2$. We take $r'_2 := \rho_+(r_1) + r_2$, then for any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(V)_1$ with $d(f(C), \text{supp}(h)) > r'_2$, we have $d(f(C), \text{supp}(h)) > r_2$ and hence:

$$\|h T \chi_C\| = \|T \chi_C h\| \leq \|\chi_{N_{r_1}^c(C)} T \chi_C h\| + \|\chi_{N_{r_1}(C)} T h \chi_C\| \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

In the sequel, we will use this observation without further explanation.

The following result shows that twisted quasi-local algebras are indeed C^* -algebras:

Lemma 5.11. *The set $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ forms a C^* -subalgebra of $\mathfrak{B}(E)$.*

Proof. It follows from Remark 5.10 that $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ is a complete $*$ -closed linear space in $\mathfrak{B}(E)$. It remains to show that $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ is closed under multiplication. Let $T_1, T_2 \in C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ and fix $x, y \in Z_d$. We claim that the series $\sum_{z \in Z_d} (T_1)_{xz} (T_2)_{zy}$ converges in norm $\|\cdot\|_{\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}}$, which implies that condition (1) holds for $T_1 T_2$. In fact due to condition (2), for any $\epsilon > 0$ there exist finite rank operators $S_1, S_2 \in \mathfrak{K}(E)$ with finitely many non-zero matrix entry such that $\|\chi_{\{x\}} T_1 - S_1\| < \epsilon$ and $\|T_2 \chi_{\{y\}} - S_2\| < \epsilon$, which implies that $\|\chi_{\{x\}} T_1 - \chi_{\{x\}} S_1\| < \epsilon$ and $\|T_2 \chi_{\{y\}} - S_2 \chi_{\{y\}}\| < \epsilon$. Hence there exists a finite $F_0 \subset Z_d$ such that for any $F \subseteq Z_d \setminus F_0$, we have

$$\epsilon > \|\chi_{\{x\}} T_1 \chi_F - \chi_{\{x\}} S_1 \chi_F\| = \|\chi_{\{x\}} T_1 \chi_F\| \quad \text{and} \quad \epsilon > \|\chi_F T_2 \chi_{\{y\}} - \chi_F S_2 \chi_{\{y\}}\| = \|\chi_F T_2 \chi_{\{y\}}\|.$$

Therefore we obtain that

$$\left\| \sum_{z \in F} (T_1)_{xz} (T_2)_{zy} \right\|_{\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}} = \|\chi_{\{x\}} T_1 \chi_F T_2 \chi_{\{y\}}\|_{\mathfrak{B}(E)} < \epsilon^2,$$

which concludes the claim. It is routine to check that condition (2)-(4) hold for $T_1 T_2$, hence we finish the proof. \square

It is straightforward to check that $C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ is a C^* -subalgebra of $C_q^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})$. For later use, we denote the natural inclusion:

$$(5.4) \quad i^{\mathcal{A}} : C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}) \hookrightarrow C_q^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}).$$

In the sequel, we will show that $i^{\mathcal{A}}$ induces isomorphisms on K -theories. This is a crucial step to achieve the main result.

6. CONSTRUCTION OF THE BOTT MAPS

This section is devoted to linking the K -theories of Roe and strongly quasi-local algebras with those of their twisted counterparts. To achieve, we construct the so-called Bott maps β_r, β_{sq} following the idea from [4] and fill them into the following commutative diagram:

$$(6.1) \quad \begin{array}{ccc} K_*(C^*(P_d(X))) & \xrightarrow{\beta_r} & K_*(C^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})) \\ \downarrow i_* & & \downarrow i_*^{\mathcal{A}} \\ K_*(C_{sq}^*(P_d(X))) & \xrightarrow{\beta_{sq}} & K_*(C_q^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})), \end{array}$$

where the vertical lines are induced by inclusions. We follow the same notation introduced in Section 5 and fix $d \geq 0$.

6.1. The Roe algebra case. We start by recalling the Bott map constructed in [4] for Roe algebras. A key observation here is that the image of the Bott map

$$\beta : K_*(P_d(X)) \rightarrow K_*(C^*(P_d(X), Q((M_2(\mathcal{A}_n^+) \otimes \mathfrak{K})_n)))$$

can be made into the K -theory of the restricted twisted algebra introduced above, which makes it possible to relate to the strongly quasi-local case.

The crucial ingredient to construct the Bott map is the existence of the uniformly almost flat Bott generators. For each $n \in \mathbb{N}$, denote by \mathcal{A}_n^+ the C^* -algebra unitization of $\mathcal{A}_n = C_0(V_n, \text{Cliff}(W_n))$ and by $M_2(\mathcal{A}_n^+) = \mathcal{A}_n^+ \otimes M_2(\mathbb{C})$ the algebra of 2×2 matrices over \mathcal{A}_n^+ . Set $b_0 := \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \in M_2(\mathcal{A}_n^+)$. For a matrix $a = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ of functions on V_n , we define the support of a to be

$$\text{supp}(a) := \bigcup_{i,j=1}^2 \text{supp}(a_{ij}).$$

Lemma 6.1 ([4, Lemma 3.6 and Definition 3.7]). *For any $R > 0$ and $\epsilon > 0$, there exist $r > 0$ and a family of idempotents $\{b_{x,r}^{(n)}\}_{n \in \mathbb{N}, x \in V_n}$ in $M_2(\mathcal{A}_n^+)$ which is $(R, \epsilon; r)$ -flat in the following sense: $\text{supp}(b_{x,r}^{(n)} - b_0) \subseteq B_{V_n}(x, r)$ and for any $n \in \mathbb{N}$ and $x, y \in V_n$ with $\|x - y\| < R$, we have*

$$\sup_{v \in V_n} \|b_{x,r}^{(n)} - b_{y,r}^{(n)}\|_{\text{Cliff}(W_n) \otimes M_2(\mathbb{C})} < \epsilon.$$

For convenience to readers, we briefly recall the construction here. Details can be found in [4, Section 3.2]. For each $n \in \mathbb{N}$, $x \in V_n$ and $r > 0$, we define a function $f_{x,r}^{(n)} : V_n \rightarrow W_n \subset \text{Cliff}(W_n)$ by:

$$f_{x,r}^{(n)}(v) = \varphi_r(\|v - x\|) \psi\left(\frac{v - x}{\|v - x\|}\right)$$

where $\psi : S(\bigcup_{n=1}^{\infty} V_n) \rightarrow S(\bigcup_{n=1}^{\infty} W_n)$ is the uniformly continuous function from the definition of Property (H) for V , and $\varphi_r : [0, \infty) \rightarrow [0, \infty)$ is defined by

$$\varphi_r(t) = \begin{cases} 0 & \text{if } 0 \leq t \leq r/2, \\ (2t/r) - 1 & \text{if } r/2 \leq t \leq r, \\ 1 & \text{if } t \geq r. \end{cases}$$

Let

$$W_{x,r} = \begin{pmatrix} 1 & f_{x,r}^{(n)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ f_{x,r}^{(n)} & 1 \end{pmatrix} \begin{pmatrix} 1 & f_{x,r}^{(n)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

and we set:

$$b_{x,r}^{(i)} = W_{x,r} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} W_{x,r}^{-1}.$$

Then it was shown in [4] that $\{b_{x,r}^{(n)}\}$ satisfies the conditions in Lemma 6.1.

We now recall the construction of the Bott map β_r for the Roe algebra $C^*(P_d(X))$ from [4, Section 3.3]. As in [4, Definition 2.4], we choose $\mathcal{H}_{P_d(X)} := \ell^2(Z_d; \mathcal{H})$ as the $P_d(X)$ -module and define $\mathbb{C}_f[P_d(X)]$ to be the $*$ -algebra of all bounded functions $T : Z_d \times Z_d \rightarrow \mathfrak{K}$ such that

- (1) for any bounded subset $B \subseteq X$, the set $\{(x, y) \in (B \times B) \cap (Z_d \times Z_d) : T(x, y) \neq 0\}$ is finite;
- (2) there exists an $L > 0$ such that $\#\{x \in Z_d : T(x, y) \neq 0\} < L$ and $\#\{y \in Z_d : T(x, y) \neq 0\} < L$ for each $x, y \in Z_d$;
- (3) there exists an $R \geq 0$ such that $T(x, y) = 0$ whenever $d(x, y) > R$ for $x, y \in Z_d$.

Note that $\mathbb{C}_f[X]$ can be regarded as a subalgebra in $\mathfrak{B}(\mathcal{H}_{P_d(X)})$, and its norm closure is $*$ -isomorphic to $C^*(P_d(X))$.

Since $C^*(P_d(X))$ is stable, it suffices to define $\beta_r([P])$ for any idempotent $P \in C^*(P_d(X))$. For any $0 < \epsilon_1 < 1/100$, take $Q \in \mathbb{C}_f[P_d(X)]$ such that $\|P - Q\| < \epsilon_1/(4\|P\|)$. Then $\|Q - Q^2\| < \epsilon_1$ and there exists $R_{\epsilon_1} > 0$ such that the propagation of Q does not exceed R_{ϵ_1} .

For any $\epsilon_2 > 0$, it follows from Lemma 6.1 that there exists a family of $(R_{\epsilon_1}, \epsilon_2; r)$ -flat idempotents $\{b_v^{(n)}\}_{n \in \mathbb{N}, v \in V_n}$ in $M_2(\mathcal{A}_n^+)$. Denote:

$$Q((M_2(\mathcal{A}_n^+))_n) := \frac{\prod_{n=1}^{\infty} M_2(\mathcal{A}_n^+)}{\bigoplus_{n=1}^{\infty} M_2(\mathcal{A}_n^+)} \quad \text{and} \quad Q((M_2(\mathcal{A}_n))_n) := \frac{\prod_{n=1}^{\infty} M_2(\mathcal{A}_n)}{\bigoplus_{n=1}^{\infty} M_2(\mathcal{A}_n)}.$$

We define $\widetilde{Q}, \widetilde{Q}_0 : Z_d \times Z_d \rightarrow Q((M_2(\mathcal{A}_n^+))_n) \otimes \mathfrak{K}$ by the formula

$$(6.2) \quad \widetilde{Q}(x, y) := \left[(b_{f(x)}^{(1)}, \dots, b_{f(x)}^{(n)}, \dots) \right] \otimes Q(x, y),$$

$$(6.3) \quad \widetilde{Q}_0(x, y) := \left[(b_0, \dots, b_0, \dots) \right] \otimes Q(x, y),$$

respectively, for all $(x, y) \in Z_d \times Z_d$. Note that here $b_{f(x)}^{(n)}$ is well defined for sufficiently large n such that $f(x) \in V_n$. It is clear that

$$\widetilde{Q}, \widetilde{Q}_0 \in \mathbb{C}[P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n^+))_n) \otimes \mathfrak{K}]$$

and

$$\widetilde{Q} - \widetilde{Q}_0 \in \mathbb{C}[P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K}].$$

Furthermore, since X has bounded geometry and Q has finite propagation, it follows from Lemma 6.1 that $\|\widetilde{Q}^2 - \widetilde{Q}\| < 1/5$ and $\|\widetilde{Q}_0^2 - \widetilde{Q}_0\| < 1/5$ if we choose ϵ_1 and ϵ_2 sufficiently small. Hence the spectrum of either \widetilde{Q} or \widetilde{Q}_0 is contained in disjoint neighbourhoods S_0 of 0 and S_1 of 1. Let $\chi : S_0 \sqcup S_1 \rightarrow \mathbb{C}$ be a continuous function such that $\chi(S_0) = 0$ and $\chi(S_1) = 1$. Define $\Theta := \chi(\widetilde{Q})$ and $\Theta_0 := \chi(\widetilde{Q}_0)$. Then Θ and Θ_0 are idempotents in $C^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n^+))_n) \otimes \mathfrak{K})$, and $\Theta - \Theta_0$ belongs to the closed two-sided ideal $C^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K})$.

Finally we apply the *difference construction* in K -theory of Banach algebras introduced by Kasparov-Yu [12]. More precisely, for a closed two-sided ideal J of a Banach algebra B and two idempotents $p, q \in B$ with $p - q \in J$, we can define a difference element $D(p, q) \in K_0(J)$ associated to p, q . We omit the details here but guide the readers to their original paper. Now we define

$$\beta_r([P]) := D(\Theta, \Theta_0) \in K_0(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})).$$

The correspondence $[P] \mapsto \beta_r([P])$ extends to a homomorphism

$$\beta_r : K_0(C^*(P_d(X))) \longrightarrow K_0(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})),$$

which is called the *Bott map*. By suspension, we define the Bott map for K_1 -groups in a similar way:

$$\beta_r : K_1(C^*(P_d(X))) \longrightarrow K_1(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})).$$

Remark 6.2. Readers might already notice that the above Bott map β_r is slightly different from the one constructed in [4, Section 3.3], which is defined to be

$$\beta = \iota_* \circ \beta_r : K_*(C^*(P_d(X))) \longrightarrow K_*(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)))$$

where ι is the monomorphism from Corollary 5.8. This is due to the observation that the elements \widetilde{Q} and \widetilde{Q}_0 defined in (6.2) and (6.3) indeed belong to the subalgebra $\mathcal{Q}((M_2(\mathcal{A}_n^+))_n) \otimes \mathfrak{K}$ instead of merely the original one $\mathcal{Q}((M_2(\mathcal{A}_n^+))_n \otimes \mathfrak{K})$. This is crucial to construct Diagram (6.1) as revealed in the next subsection.

6.2. The strongly quasi-local case. Now we move to the strongly quasi-local case and define the Bott map $\beta_{sq} : K_*(C_{sq}^*(P_d(X))) \rightarrow K_*(C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}))$. The construction is similar to the Bott map β_r for Roe algebras in Section 6.1, while the argument is more involved. Again we only consider the K_0 -case, since the K_1 -case can be done by suspension.

As in the Roe case, we choose $\mathcal{H}_X := \ell^2(Z_d, \mathcal{H})$ as the fixed $P_d(X)$ -module. Since $C_{sq}^*(\mathcal{H}_X)$ is stable, it suffices to define $\beta_{sq}([P])$ for any idempotent $P \in C_{sq}^*(\mathcal{H}_X)$. In the rest of this subsection, let us fix an idempotent $P \in C_{sq}^*(\mathcal{H}_X)$ and an $\epsilon \in$

$(0, \frac{1}{4})$. It follows from Definition 3.6 that there exist δ, R such that for any strongly measurable map $g : P_d(X) \rightarrow \mathfrak{K}_1$ with (δ, R) -variation, we have $\|[P \otimes \text{Id}_{\mathcal{H}}, \Lambda(g)]\| < \epsilon$. We also fix such parameters δ and R .

Recall that $f : X \rightarrow V$ is the coarse embedding and is extend to a continuous coarse embedding $f : P_d(X) \rightarrow V$ by affine extension (see the beginning of Section 5). We fix two proper functions $\rho_{\pm} : [0, \infty) \rightarrow [0, \infty)$ such that for any $x, y \in P_d(X)$ we have:

$$\rho_-(d(x, y)) \leq \|f(x) - f(y)\| \leq \rho_+(d(x, y)).$$

Moreover, Lemma 6.1 provides an $r > 0$ and a family of idempotents $\{b_x^{(n)}\}_{n \in \mathbb{N}, x \in V_n}$ which is $(\rho_+(R), \delta; r)$ -flat. Again let us fix such r and $\{b_x^{(n)}\}$.

Denote $E_2 := \ell^2(Z_d; \mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K})$ the Hilbert module over $\mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K}$, and fix a faithful representation $\pi : \mathcal{Q}((M_2(\mathcal{A}_n))_n) \rightarrow \mathfrak{B}(\mathcal{H})$. Then $\pi \otimes i_{\mathfrak{K}} : \mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K} \rightarrow \mathfrak{B}(\mathcal{H} \otimes \mathcal{H})$ is also a faithful representation, where $i_{\mathfrak{K}} : \mathfrak{K} \rightarrow \mathfrak{B}(\mathcal{H})$ is the inclusion map. For any $T \in C_{sq}^*(\mathcal{H}_X) \subset \mathfrak{B}(\ell^2(Z_d; \mathcal{H}))$, we consider its amplification $a(T) \in \mathfrak{B}(\ell^2(Z_d; \mathcal{H} \otimes \mathcal{H}))$ defined by $a(T)_{xy} = \text{Id}_{\mathcal{H}} \otimes T_{xy}$ for any $x, y \in Z_d$. It follows from Lemma 5.1 that there is an operator $\Psi(a(T)) \in \mathfrak{B}(E_2)$ defined by (5.2). Note that $\Psi(a(T))_{xy} = \text{Id}_{\mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K}} \otimes T_{xy}$ for any $x, y \in Z_d$.

On the other hand, we define a diagonal operator $B \in \mathfrak{B}(E_2)$ by

$$B_{xy} := \begin{cases} \left[\left(b_{f(x)}^{(1)}, \dots, b_{f(x)}^{(n)}, \dots \right) \right] \otimes \text{Id}_{\mathcal{H}}, & y = x; \\ 0, & \text{otherwise} \end{cases}$$

where $b_{f(x)}^{(n)}$ is well-defined for n large enough such that $f(x) \in V_n$. Note that each $\left[\left(b_{f(x)}^{(1)}, \dots, b_{f(x)}^{(n)}, \dots \right) \right] \in \mathcal{M}(\mathcal{Q}((M_2(\mathcal{A}_n))_n))$, hence B is well-defined. Similarly, we define $B_0 \in \mathfrak{B}(E_2)$ by

$$(B_0)_{xy} := \begin{cases} \left[(b_0, \dots, b_0, \dots) \right] \otimes \text{Id}_{\mathcal{H}}, & y = x; \\ 0, & \text{otherwise.} \end{cases}$$

It is clear that B and B_0 are idempotents in $\mathfrak{B}(E_2)$, and $\|B\| = \|B_0\| = 1$.

Now for the given idempotent $P \in C_{sq}^*(\mathcal{H}_X)$, we define $\widetilde{P} := B \cdot \Psi(a(P)) \in \mathfrak{B}(E_2)$ and $\widetilde{P}_0 := B_0 \cdot \Psi(a(P)) \in \mathfrak{B}(E_2)$. It is clear that for any $x, y \in Z_d$, we have:

$$(6.4) \quad \widetilde{P}(x, y) = \left[\left(b_{f(x)}^{(1)}, \dots, b_{f(x)}^{(n)}, \dots \right) \right] \otimes P(x, y),$$

$$(6.5) \quad \widetilde{P}_0(x, y) = \left[(b_0, \dots, b_0, \dots) \right] \otimes P(x, y).$$

Readers may compare (6.4), (6.5) with (6.2), (6.3).

We define the C^* -algebra $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$ similarly to Definition 5.9 except that we use $\mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K}$ to replace $\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$ therein. It is clear that $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$ is a subalgebra of $\mathfrak{B}(E_2)$. We also define $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n^+))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$ to be the C^* -subalgebra in $\mathfrak{B}(E_2)$ generated by $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$ and $\{B_0 \cdot \Psi(a(Q)) : Q \text{ is an idempotent in } C_{sq}^*(\mathcal{H}_X)\}$. It is easy to check that $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$ is a two-sided $*$ -ideal in $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n^+))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$.

The following lemma is a crucial step to define the Bott map:

Lemma 6.3. *With the same notation as above, we have:*

$$\widetilde{P}, \widetilde{P}_0 \in C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n^+))_{n \in \mathbb{N}}) \otimes \mathfrak{K}) \text{ and } \widetilde{P} - \widetilde{P}_0 \in C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_{n \in \mathbb{N}}) \otimes \mathfrak{K}).$$

Proof. It suffices to prove that $\widetilde{P} - \widetilde{P}_0 \in C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_{n \in \mathbb{N}}) \otimes \mathfrak{K})$. We need to check condition (1)-(4) in Definition 5.9 for $\widetilde{P} - \widetilde{P}_0$. Note that (1) holds trivially.

For (3): Since P is quasi-local, it follows that for any $\epsilon' > 0$ there exists $r_1 > 0$ such that for any Borel subsets $C, D \subseteq P_d(X)$ with $d(C, D) > r_1$, then $\|\chi_C P \chi_D\| < \epsilon' / \|B - B_0\|$. Hence

$$\begin{aligned} \|\chi_C(B - B_0)\Psi(a(P))\chi_D\| &= \|(B - B_0)\chi_C\Psi(a(P))\chi_D\| \\ &\leq \|B - B_0\| \cdot \|\Psi(a(\chi_C P \chi_D))\| \leq \|B - B_0\| \cdot \|\chi_C P \chi_D\| < \epsilon', \end{aligned}$$

where we use Lemma 5.1 in the second last inequality.

For (4): Given $\epsilon' > 0$, we take $r_2 = r$. For any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(V)_1$ with $d(f(C), \text{supp}(h)) > r_2$, we have:

$$\chi_C(\widetilde{P} - \widetilde{P}_0)h = \chi_C(B - B_0)\Psi(a(P))h = h(B - B_0)\chi_C\Psi(a(P)).$$

Since $\text{supp}(b_{f(x)}^{(n)} - b_0) \subseteq B_{V_n}(f(x), r)$ for any $x \in P_d(X)$ and n large enough such that $f(x) \in V_n$, we obtain that $h(B - B_0)\chi_C = 0$ due to the choice of r_2 .

It remains to prove (2). Fixing a compact subset $K \subset P_d(X)$, we aim to show that $\chi_K(\widetilde{P} - \widetilde{P}_0)$ and $(\widetilde{P} - \widetilde{P}_0)\chi_K$ belong to $\mathfrak{K}(E_2)$. Here we only consider $\chi_K(\widetilde{P} - \widetilde{P}_0)$ since the other is similar. Note that

$$\chi_K(\widetilde{P} - \widetilde{P}_0) = (B - B_0)\chi_K\Psi(a(P)) = (B - B_0) \cdot \Psi(a(\chi_K P)).$$

Since $P \in C_{sq}^*(P_d(X))$, then $\chi_K P$ is compact. Without loss of generality, we may assume that $\chi_K P$ is rank-one, i.e., $(\chi_K P)(\zeta) = \xi \langle \eta, \zeta \rangle$ where $\xi, \eta, \zeta \in \ell^2(Z_d; \mathcal{H})$. We fix a unit vector $v_0 \in \mathcal{H}$. Denote the rank one operator $u \mapsto v \langle w, u \rangle$ by $\theta_{v,w} \in \mathfrak{B}(\mathcal{H})$ where $u, v, w \in \mathcal{H}$. Direct calculation shows that for any $x, y \in Z_d$, we have:

$$\left((B - B_0) \cdot \Psi(a(\chi_K P)) \right)_{xy} = \left[(b_{f(x)}^{(1)} - b_0, \dots, b_{f(x)}^{(n)} - b_0, \dots) \right] \otimes \theta_{\xi(x), v_0} \theta_{v_0, \eta(y)}.$$

Let $\{q_i\}_{i \in I}$ be a net of approximating units for the C^* -algebra $\mathcal{Q}((M_2(\mathcal{A}_n))_n)$. We consider two vectors $\Xi_1, \Xi_2 \in E_2$ defined by

$$\Xi_1 := \sum_{x \in Z_d} \left([(b_{f(x)}^{(1)} - b_0, \dots, b_{f(x)}^{(n)} - b_0, \dots)] \otimes \theta_{\xi(x), v_0} \right) [x],$$

and

$$\Xi_2^{(i)} := \sum_{x \in Z_d} (q_i \otimes \theta_{\eta(x), v_0}) [x].$$

For each $i \in I$, denote the operator $\Theta_i : \Xi \mapsto \Xi_1 \langle \Xi_2^{(i)}, \Xi \rangle$ where $\Xi \in E_2$. Then clearly $\Theta \in \mathfrak{K}(E_2)$. Moreover, we obtain that the net $\{\Theta_i\}_{i \in I}$ converges to $\chi_K(\widetilde{P} - \widetilde{P}_0)$ in norm, which concludes the proof. \square

Our last ingredient to construct the Bott map is the following:

Lemma 6.4. *With the same notation as above, we have $\|\widetilde{P}^2 - \widetilde{P}\| \leq \epsilon$ and $\|\widetilde{P}_0^2 - \widetilde{P}_0\| \leq \epsilon$.*

Proof. We only deal with the case of \widetilde{P} while the other is trivial. Recall that $\widetilde{P} = B \cdot \Psi(a(P))$ and B, P are idempotents, hence it suffices to control $\|[B, \Psi(a(P))]\|$.

Recall that $\pi : \mathcal{Q}((M_2(\mathcal{A}_n))_n) \rightarrow \mathfrak{B}(\mathcal{H})$ is the faithful representation fixed above. We consider the following map $g : P_d(X) \rightarrow \mathfrak{B}(\mathcal{H})_1$ defined by

$$g(x) = \pi\left(\left[b_{f(x)}^{(1)}, \dots, b_{f(x)}^{(n)}, \dots\right]\right), \quad x \in P_d(X).$$

Since $\{b_z^{(n)}\}_{n \in \mathbb{N}, z \in V_n}$ is $(\rho_+(R), \delta; r)$ -flat for the R, δ chosen above, we obtain that the map g has (δ, R) -variation on $P_d(X)$. Moreover, it is easy to check that g is continuous. Define a diagonal operator $G \in \mathfrak{B}(\ell^2(Z_d; \mathcal{H} \otimes \mathcal{H}))$ by

$$G_{xy} := \begin{cases} g(x) \otimes \text{Id}_{\mathcal{H}}, & \text{if } x = y; \\ 0, & \text{otherwise.} \end{cases}$$

Then it is clear that $\Psi([G, a(P)]) = [B, \Psi(a(P))]$. Hence from Lemma 5.1, it suffices to control $\|[G, a(P)]\|_{\mathfrak{B}(\ell^2(Z_d; \mathcal{H} \otimes \mathcal{H}))}$.

For any finite rank projection $q \in \mathfrak{B}(\mathcal{H})$, denote $\hat{q} := \text{Id}_{\ell^2(Z_d)} \otimes q \otimes \text{Id}_{\mathcal{H}} \in \mathfrak{B}(\ell^2(Z_d; \mathcal{H} \otimes \mathcal{H}))$. We regard $q(\mathcal{H})$ as a subspace in \mathcal{H} . Consider the continuous map $g_q : P_d(X) \rightarrow \mathfrak{B}(q\mathcal{H})_1 \subseteq \mathfrak{K}(\mathcal{H})_1$ defined by $g_q(x) := q \cdot g(x) \cdot q$, which also has (δ, R) -variation. It is straightforward to check that for any $x, y \in Z_d$, we have:

$$(\Lambda(g_q))_{xy} = \begin{cases} \text{Id}_{\mathcal{H}} \otimes g_q(x), & \text{if } x = y; \\ 0, & \text{otherwise} \end{cases}$$

where $\Lambda(g_q)$ is defined as in Definition 3.5. Hence we obtain:

$$\|\hat{q}[G, a(P)]\hat{q}\| \leq \|[P \otimes \text{Id}_{\mathcal{H}}, \Lambda(g_q)]\| < \epsilon.$$

Finally note that

$$\|[G, a(P)]\| = \sup \left\{ \|\hat{q}[G, a(P)]\hat{q}\| : q \text{ is a finite rank projection in } \mathfrak{B}(\mathcal{H}) \right\},$$

hence we conclude the proof. \square

Consequently, we obtain from Lemma 6.4 that the spectrum of either \widetilde{P} or \widetilde{P}_0 is contained in disjoint neighbourhoods S_0 of 0 and S_1 of 1. Let $\chi : S_0 \sqcup S_1 \rightarrow \mathbb{C}$ be a continuous function such that $\chi(S_0) = 0$ and $\chi(S_1) = 1$. Define $\Theta := \chi(\widetilde{P})$ and $\Theta_0 := \chi(\widetilde{P}_0)$. Then it follows from Lemma 6.3 that Θ and Θ_0 are idempotents in $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n^+))_n) \otimes \mathfrak{K})$, and $\Theta - \Theta_0$ belongs to the closed two-sided ideal $C_q^*(P_d(X), \mathcal{Q}((M_2(\mathcal{A}_n))_n) \otimes \mathfrak{K})$.

Finally we apply the difference construction from [12] as in the Roe case, and define

$$\beta_{sq}([P]) := D(\Theta, \Theta_0) \in K_0(C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})).$$

The correspondence $[P] \mapsto \beta_{sq}([P])$ extends to a homomorphism

$$\beta_{sq} : K_0(C_{sq}^*(P_d(X))) \longrightarrow K_0(C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})),$$

which is also called the *Bott map*. From the constructions above, it is easy to see that Diagram (6.1) commutes.

7. LOCAL ISOMORPHISM

In this section, we relate the K -theories of restricted twisted Roe algebras and their quasi-local counterparts. The main result is the following:

Theorem 7.1. *Let X be a discrete metric space with bounded geometry which admits a coarse embedding into a Banach space with Property (H). Then for each $d \geq 0$, the natural inclusion $i^{\mathcal{A}}$ from (5.4) induces isomorphisms on K -theories, i.e., the following*

$$i_*^{\mathcal{A}} : K_*\left(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})\right) \longrightarrow K_*\left(C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})\right)$$

is an isomorphism for $*$ = 0, 1.

The idea of the proof is to decompose the twisted algebras into smaller subalgebras which are easier to handle and apply the Mayer-Vietoris sequence to paste them together. This follows the outline of [4, Section 4], which originates from [29]. We use the same notation introduced in Section 5 and 6, and fix a $d \geq 0$ throughout the section.

The following notion is inspired by [4, Definition 4.2]:

Definition 7.2. Let $O \subseteq V$ be an open subset of V . Define $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]_O$ to be the $*$ -subalgebra of $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ generated by elements $T \in \mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]$ such that for any $x, y \in Z_d$, we have:

$$\text{supp}(h_n) \subseteq O \cap V_n$$

where $T_{xy} = [(h_1, \dots, h_n, \dots)] \in \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$. We define $C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O$ to be the norm closure of $\mathbb{C}[P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}]_O$ in $\mathfrak{B}(E)$.

Similarly, we introduce the following for quasi-local algebras:

Definition 7.3. Let $O \subseteq V$ be an open subset of V . Define $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O$ to be the $*$ -subalgebra of $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ generated by elements $T \in C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ such that for any $x, y \in Z_d$, we have:

$$\text{supp}(h_n) \subseteq O \cap V_n$$

where $T_{xy} = [(h_1, \dots, h_n, \dots)] \in \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$. For a Borel subset $Y \subseteq P_d(X)$, we also denote by $C_q^*(Y, \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O$ the C^* -algebras consisting of elements in $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ with support in $Y \times Y$.

Note that $C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O$ is a two-sided $*$ -ideal of $C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$. It is also easy to see that $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O$ is norm closed, and is a two-sided $*$ -ideal of $C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$. Moreover, it is clear that we have the inclusion

$$i^{\mathcal{A}} : C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O \hookrightarrow C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O.$$

We need to consider some special open subsets O as follows:

Definition 7.4. Let $\Gamma \subseteq X$ and $r > 0$. An open subset $O \subseteq V$ is said to be (Γ, r) -separate if the following holds:

- (1) $O = \bigsqcup_{\gamma \in \Gamma} O_\gamma$, where $O_\gamma \subset V$ such that $O_\gamma \cap O_{\gamma'} = \emptyset$ whenever $\gamma \neq \gamma'$;
- (2) for any $\gamma \in \Gamma$, we have $O_\gamma \subseteq B_V(f(\gamma), r)$.

The following is the key building block to achieve Theorem 7.1:

Proposition 7.5. *Suppose $\Gamma \subset X$ and $r > 0$. Then for any (Γ, r) -separate open subset $O \subset V$, we have that*

$$i_*^{\mathcal{A}} : K_*(C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O) \cong K_*(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O)$$

is an isomorphism for $*$ = 0, 1.

The proof is divided into several lemmas. We start with some extra notation. Let $O = \bigsqcup_{\gamma \in \Gamma} O_\gamma$ be a (Γ, r) -separate open subset of V . For each $\gamma \in \Gamma$, we define $(\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}$ to be a C^* -subalgebra of $\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$ generated by $[(h_1, \dots, h_n, \dots)] \in \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}$ such that $\text{supp}(h_n) \subseteq O_\gamma \cap V_n$ for all $n \in \mathbb{N}$.

For each $\gamma \in \Gamma$ and $S \geq 0$, we denote $Y_\gamma(S) := \{x \in P_d(X) : d(x, \gamma) \leq S\}$. We also fix two proper functions $\rho_\pm : [0, \infty) \rightarrow [0, \infty)$ such that for any $x, y \in P_d(X)$ we have:

$$\rho_-(d(x, y)) \leq \|f(x) - f(y)\| \leq \rho_+(d(x, y)).$$

The following lemma is actually from [4, Lemma 4.5], hence we omit the proof.

Lemma 7.6. *Let $O = \bigsqcup_{\gamma \in \Gamma} O_\gamma$ be a (Γ, r) -separate open subset of V . Then*

$$C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O \cong \lim_S \prod_{\gamma \in \Gamma} C^*(Y_\gamma(S)) \otimes (\mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}.$$

For the quasi-local case, we have an analogous decomposition. Since the proof is similar, we only provide a sketch here.

Lemma 7.7. *Let $O = \bigsqcup_{\gamma \in \Gamma} O_\gamma$ be a (Γ, r) -separate open subset of V . Then*

$$C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O \cong \lim_S \prod_{\gamma \in \Gamma} C_q^*(Y_\gamma(S), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}.$$

Sketch of proof. For any $T \in C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O$ and $\gamma \in \Gamma$, it is clear that $T\chi_{O_\gamma} \in C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}$. Given $\epsilon > 0$, it follows from Definition 5.9 together with Remark 5.10 that there exist $r_2 > 0$ such that for any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(V)_1$ with $d(f(C), \text{supp}(h)) \geq r_2$, we have $\|\chi_C Th\| < \epsilon/2$ and $\|hT\chi_C\| < \epsilon/2$.

Take $S := \sup\{s \in [0, \infty) : \rho_-(s) \leq r + r_2\}$. Then for every $\gamma \in \Gamma$, we have

$$f(X \setminus Y_\gamma(S)) \subseteq V \setminus B(f(\gamma), r + r_2),$$

which implies that $d(f(X \setminus Y_\gamma(S)), O_\gamma) \geq r_2$. Hence we obtain:

$$\|T\chi_{O_\gamma} - \chi_{Y_\gamma(S)} T\chi_{O_\gamma}\| = \|\chi_{X \setminus Y_\gamma(S)} T\chi_{O_\gamma}\| < \epsilon/2,$$

and

$$\|\chi_{O_\gamma} T - \chi_{O_\gamma} T\chi_{Y_\gamma(S)}\| = \|\chi_{O_\gamma} T\chi_{X \setminus Y_\gamma(S)}\| < \epsilon/2.$$

Note that $T\chi_{O_\gamma} = \chi_{O_\gamma}T$, so we obtain:

$$\begin{aligned} \|T\chi_{O_\gamma} - \chi_{Y_\gamma(S)}(T\chi_{O_\gamma})\chi_{Y_\gamma(S)}\| &\leq \|T\chi_{O_\gamma} - \chi_{Y_\gamma(S)}T\chi_{O_\gamma}\| + \|\chi_{Y_\gamma(S)}T\chi_{O_\gamma} - \chi_{Y_\gamma(S)}(T\chi_{O_\gamma})\chi_{Y_\gamma(S)}\| \\ &\leq \epsilon/2 + \|\chi_{O_\gamma}T - \chi_{O_\gamma}T\chi_{Y_\gamma(S)}\| \leq \epsilon/2 + \epsilon/2 = \epsilon. \end{aligned}$$

Therefore, we obtain a well-defined homomorphism

$$\Phi : C_q^*(P_d(X), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_O \longrightarrow \lim_S \prod_{\gamma \in \Gamma} C_q^*(Y_\gamma(S), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}$$

defined by

$$T \mapsto (T\chi_{O_\gamma})_{\gamma \in \Gamma} = \lim_{S \rightarrow \infty} (\chi_{Y_\gamma(S)}(T\chi_{O_\gamma})\chi_{Y_\gamma(S)})_{\gamma \in \Gamma}.$$

On the other hand, note that $T = \sum_{\gamma \in \Gamma} T\chi_{O_\gamma}$ where the summation converges in the strong $*$ -operator topology. Moreover, the above argument shows that

$$T = \sum_{\gamma \in \Gamma} T\chi_{O_\gamma} = \lim_{S \rightarrow \infty} \sum_{\gamma \in \Gamma} \chi_{Y_\gamma(S)}(T\chi_{O_\gamma})\chi_{Y_\gamma(S)}.$$

Therefore, it is not hard to check that Φ provides a $*$ -isomorphism. Details are omitted here. \square

Furthermore, we have the following:

Lemma 7.8. *For each $S \geq 0$ and $\gamma \in \Gamma$, we have*

$$C_q^*(Y_\gamma(S), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma} \cong C_q^*(Y_\gamma(S)) \otimes (Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}.$$

Proof. For any $T \in C_q^*(Y_\gamma(S), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}$, we have $T = \chi_{Y_\gamma(S)}T \in \mathfrak{K}(E)$. Hence it is easy to see that

$$\begin{aligned} C_q^*(Y_\gamma(S), Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma} &= \mathfrak{K}(\ell^2(Y_\gamma \cap Z_d, Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})) \\ &\cong \mathfrak{K}(\ell^2(Y_\gamma(S) \cap Z_d)) \otimes (Q((\mathcal{A}_n)_n) \otimes \mathfrak{K}) \\ &\cong C_q^*(Y_\gamma(S)) \otimes (Q((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_\gamma}, \end{aligned}$$

which concludes the proof. \square

Proof of Proposition 7.5. Note that for each $\gamma \in \Gamma$, we have:

$$C_q^*(Y_\gamma(S)) \cong \mathfrak{K}(\ell^2(Y_\gamma(S) \cap Z_d)) \cong C^*(Y_\gamma(S)).$$

Hence the proof follows directly from Lemma 7.6, Lemma 7.7 and Lemma 7.8. \square

Having established Proposition 7.5, we now apply a Mayer-Vietoris sequence argument to obtain Theorem 7.1. To achieve, we need two extra lemmas. The first is actually [4, Lemma 4.8] together with its quasi-local version. The proof is similar, hence omitted.

Lemma 7.9. *Let $N \in \mathbb{N}$ and $\Gamma_1, \dots, \Gamma_N$ be N mutually disjoint subsets of X . For any $r > 0$ and $j \in \{1, 2, \dots, N\}$, let*

$$O_{r,j} := \bigcup_{\gamma \in \Gamma_j} B_V(f(\gamma), r).$$

Then for any $r_0 > 0$ and $k \in \{1, 2, \dots, N-1\}$, we have the following:

$$\begin{aligned}
(1) \quad & \lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{r,k}} + \lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{\bigcup_{j=1}^{k-1} O_{r,j}} \\
& = \lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{\bigcup_{j=1}^k O_{r,j}}; \\
(2) \quad & \lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{r,k}} \cap \lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{\bigcup_{j=1}^{k-1} O_{r,j}} \\
& = \lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{r,k} \cap (\bigcup_{j=1}^{k-1} O_{r,j})}; \\
(3) \quad & \lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{r,k}} + \lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{\bigcup_{j=1}^{k-1} O_{r,j}} \\
& = \lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{\bigcup_{j=1}^k O_{r,j}}; \\
(4) \quad & \lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{r,k}} \cap \lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{\bigcup_{j=1}^{k-1} O_{r,j}} \\
& = \lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{r,k} \cap (\bigcup_{j=1}^{k-1} O_{r,j})}.
\end{aligned}$$

Lemma 7.10. For any $r > 0$, denote

$$O_r := \bigcup_{x \in X} B_V(f(x), r) = \mathcal{N}_r(\text{Im} f)$$

where $X \rightarrow V$ is the coarse embedding. Then we have:

$$\begin{aligned}
\lim_{r \rightarrow +\infty} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_r} & \cong C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}); \\
\lim_{r \rightarrow +\infty} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_r} & \cong C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K}).
\end{aligned}$$

Proof. The Roe case is already known from [4], hence we only show the second. Given $T \in C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})$ and $\epsilon > 0$, there exists $r > 0$ such that for any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(V)_1$ with $d(f(C), \text{supp}(h)) > r$, we have $\|\chi_C Th\| < \epsilon$. In particular, taking a function $h \in C_b(V)_1$ such that $h|_{O_r} = 0$ and $h|_{O_{2r}} = 1$, we obtain:

$$\|T - T(1 - h)\| = \|Th\| = \|\chi_X Th\| < \epsilon.$$

Note that $T(1 - h) \in C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_{2r}}$, hence we conclude the proof. \square

Now we are in the position to prove Theorem 7.1:

Proof of Theorem 7.1. For any $r > 0$, denote $O_r := \bigcup_{x \in X} B_V(f(x), r)$. From Lemma 7.10, it suffice to show that for each $r_0 > 0$, the map

$$i_*^{\mathcal{A}} : K_* \left(\lim_{r < r_0, r \rightarrow r_0} C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_r} \right) \longrightarrow K_* \left(\lim_{r < r_0, r \rightarrow r_0} C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})_{O_r} \right)$$

is an isomorphism.

Since X has bounded geometry, there follows (see, e.g., [24, Lemma12.2.3]) that there exists an $N \in \mathbb{N}$ such that

- (1) $X = \bigsqcup_{j=1}^N \Gamma_j$ for some $\Gamma_j \subset X$ with $\Gamma_j \cap \Gamma_{j'} = \emptyset$ whenever $j \neq j'$;
- (2) for each $j \in \{1, 2, \dots, N\}$ and $\gamma \neq \gamma'$ in Γ_j , we have

$$\|f(\gamma) - f(\gamma')\|_V > 2r_0.$$

For any $0 < r < r_0$ and $j \in \{1, 2, \dots, N\}$, let

$$O_{r,j} := \bigcup_{x \in \Gamma_j} B_V(f(x), r) = \mathcal{N}_r(f(\Gamma_j)).$$

Then $O_r = \bigcup_{j=1}^N O_{r,j}$, and each $O_{r,j}$ or $O_{r,j} \cap \left(\bigcup_{i=1}^{j-1} O_{r,i}\right)$ are (Γ_j, r) -separate for any $j \in \{1, 2, \dots, N\}$. Applying the Mayer-Vietoris sequence argument finitely many times based on Lemma 7.9 and using Proposition 7.5, we conclude the proof. \square

8. PROOF OF THE MAIN RESULT

Proof of Theorem C. Combining Diagram (6.1) with the coarse assembly maps, we obtain the following commutative diagram:

$$\begin{array}{ccccc} \lim_{d \rightarrow \infty} K_*(P_d(X)) & \xrightarrow{\mu} & \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) & \xrightarrow{\beta_r} & \lim_{d \rightarrow \infty} K_*(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})) \\ & \searrow \mu_{sq} & \downarrow i_* & & \downarrow i_*^{\mathcal{A}} \\ & & \lim_{d \rightarrow \infty} K_*(C_{sq}^*(P_d(X))) & \xrightarrow{\beta_{sq}} & \lim_{d \rightarrow \infty} K_*(C_q^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})), \end{array}$$

Here we use the same notation to denote the homomorphisms after taking $d \rightarrow \infty$. Recall from [4, Theorem 1.1] together with Remark 6.2 that the composition $\iota_* \circ \beta_r \circ \mu$ is injective, where

$$\iota_* : K_*(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n)_n) \otimes \mathfrak{K})) \rightarrow K_*(C^*(P_d(X), \mathcal{Q}((\mathcal{A}_n \otimes \mathfrak{K})_n)))$$

is induced by the inclusion from Corollary 5.8. Hence we obtain that the composition $\beta_r \circ \mu$ is injective as well. Also note from Theorem 7.1 that $i_*^{\mathcal{A}}$ is an isomorphism, hence we conclude that μ_{sq} is injective via a diagram chasing argument. \square

9. THE CASE OF NON-POSITIVELY CURVED MANIFOLDS

In this final section, we outline the case of metric spaces which can be coarsely embedded into a simply-connected complete Riemannian manifold with non-positively sectional curvature. Recall that the coarse Novikov conjecture was verified for such spaces in [21]. Combining the approach from [21] with the techniques developed in this paper, we are able to prove the following:

Theorem 9.1. *Let X be a bounded geometry metric space which can be coarsely embedded into a simply-connected complete Riemannian manifold of non-positively sectional curvature. Then the strongly quasi-local coarse Novikov conjecture holds for X .*

Since the proof is quite similar to the one for Theorem C, we only provide an outline as follows.

First let us fix some notation. Let M be a simply-connected complete Riemannian manifold of non-positively sectional curvature. Without loss of generality, we may assume the dimension of M is $2n$ for some $n \in \mathbb{N}$. Following the notation from [21], denote the Clifford bundle over M by $\text{Cliff}(TM)$ and let $\mathcal{A} := C_0(M, \text{Cliff}(TM))$ be the C^* -algebra of continuous functions $a : M \rightarrow \text{Cliff}(TM)$ such

that $a(x) \in \text{Cliff}(T_x M)$ and vanish at infinity. Let $C_b(M, \text{Cliff}(TM))$ be the C^* -algebra of bounded continuous functions $a : M \rightarrow \text{Cliff}(TM)$ such that $a(x) \in \text{Cliff}(T_x M)$.

For each $d \geq 0$, we take a countable dense subset Z_d in the Rips complex $P_d(X)$ such that $Z_d \subseteq Z_{d'}$ whenever $d < d'$. Consider the Hilbert module $E := \ell^2(Z_d; \mathcal{A} \otimes \mathfrak{K})$ over $\mathcal{A} \otimes \mathfrak{K}$, where \mathfrak{K} is the C^* -algebra of compact operators on the separable infinite-dimensional Hilbert space \mathcal{H} . Recall that the associated twisted Roe algebra $C^*(P_d(X), \mathcal{A} \otimes \mathfrak{K})$ was defined in [21, Definition 3.1] as a C^* -subalgebra in $\mathfrak{B}(E)$.

Following the idea in Definition 5.9, we introduce the following:

Definition 9.2. The twisted quasi-local algebra $C_q^*(P_d(X), \mathcal{A} \otimes \mathfrak{K})$ is defined to be the C^* -subalgebra in $\mathfrak{B}(E)$ consisting of elements T satisfying the following conditions:

- (1) each matrix entry of T belongs to $\mathcal{A} \otimes \mathfrak{K}$;
- (2) for any compact $K \subseteq P_d(X)$, we have that $\chi_K T$ and $T \chi_K$ belongs to $\mathfrak{K}(E)$;
- (3) for any $\epsilon > 0$ there is an r_1 such that for any Borel subsets $C, D \subseteq P_d(X)$ with $d(C, D) > r_1$, we have $\|\chi_C T \chi_D\| < \epsilon$ and $\|\chi_D T \chi_C\| < \epsilon$;
- (4) for any $\epsilon > 0$, there is an r_2 such that for any Borel subset $C \subseteq P_d(X)$ and $h \in C_b(M)_1$ with $d(f(C), \text{supp}(h)) > r_2$, we have $\|\chi_C T h\| < \epsilon$.

The associated Bott maps β, β_{sq} can be defined similarly to the case of Banach spaces with Property (H) in Section 6, except that we replace Lemma 6.1 by the following:

Lemma 9.3 ([21, Lemma 5.2]). *For any $R > 0$ and $\epsilon > 0$, there exist $r > 0$ and a family of idempotents $\{b_{x,r}\}_{x \in M}$ in $M_2(\mathcal{A}^+)$ which is $(R, \epsilon; r)$ -flat in the following sense: $\text{supp}(b_{x,r} - b_0) \subseteq B_M(x, r)$ and for any $x, y \in M$ with $\|x - y\| < R$, we have*

$$\sup_{z \in M} \|b_{x,r}(z) - b_{y,r}(z)\|_{\text{Cliff}(T_z M) \otimes M_2(\mathbb{C})} < \epsilon,$$

where \mathcal{A}^+ represents the unitization of \mathcal{A} .

We omit the details and simply conclude that we have the following homomorphisms:

$$\beta : K_*(C^*(P_d(X))) \longrightarrow K_*(C^*(P_d(X), \mathcal{A} \otimes \mathfrak{K}))$$

$$\beta_{sq} : K_*(C_{sq}^*(P_d(X))) \longrightarrow K_*(C_q^*(P_d(X), \mathcal{A} \otimes \mathfrak{K})).$$

The last piece is an isomorphism result analogous to Theorem 7.1, which states that the natural inclusion

$$i^{\mathcal{A}} : C^*(P_d(X), \mathcal{A} \otimes \mathfrak{K}) \longrightarrow C_q^*(P_d(X), \mathcal{A} \otimes \mathfrak{K})$$

induces an isomorphism on K -theories. Again the proof is similar hence omitted.

Finally, we consider the following commutative diagram:

$$\begin{array}{ccc}
 \lim_{d \rightarrow \infty} K_*(P_d(X)) & \xrightarrow{\mu} \lim_{d \rightarrow \infty} K_*(C^*(P_d(X))) & \xrightarrow{\beta} \lim_{d \rightarrow \infty} K_*(C^*(P_d(X), \mathcal{A} \otimes \mathfrak{K})) \\
 & \mu_{sq} \searrow \quad i_* \searrow & \quad \quad \quad i_*^{\mathcal{A}} \searrow \\
 & \lim_{d \rightarrow \infty} K_*(C_{sq}^*(P_d(X))) & \xrightarrow{\beta_{sq}} \lim_{d \rightarrow \infty} K_*(C_q^*(P_d(X), \mathcal{A} \otimes \mathfrak{K})),
 \end{array}$$

where $\beta \circ \mu$ is an isomorphism due to the proof for [21, Theorem 1.1] and $i_*^{\mathcal{A}}$ is an isomorphism from above. Hence we obtain that μ_{sq} is injective, which concludes Theorem 9.1.

REFERENCES

- [1] Hengda Bao, Xiaoman Chen, and Jiawen Zhang. Strong quasi-locality and their K -theories. 2021.
- [2] Paul Baum, Alain Connes, and Nigel Higson. Classifying space for proper actions and K -theory of group C^* -algebras. In *C^* -algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 240–291. Amer. Math. Soc., Providence, RI, 1994.
- [3] Yoav Benyamini and Joram Lindenstrauss. *Geometric nonlinear functional analysis. Vol. 1*, volume 48 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2000.
- [4] Xiaoman Chen, Qin Wang, and Guoliang Yu. The coarse Novikov conjecture and Banach spaces with Property (H). *J. Funct. Anal.*, 268(9):2754–2786, 2015.
- [5] Marius Dadarlat, Rufus Willett, and Jianchao Wu. Localization C^* -algebras and K -theoretic duality. *Ann. K-Theory*, 3(4):615–630, 2018.
- [6] Alexander Engel. Rough index theory on spaces of polynomial growth and contractibility. *arXiv preprint arXiv:1505.03988*, 2015.
- [7] Guihua Gong, Qin Wang, and Guoliang Yu. Geometrization of the strong Novikov conjecture for residually finite groups. *J. Reine Angew. Math.*, 621:159–189, 2008.
- [8] Nigel Higson and John Roe. On the coarse Baum-Connes conjecture. In *Novikov conjectures, index theorems and rigidity, Vol. 2 (Oberwolfach, 1993)*, volume 227 of *London Math. Soc. Lecture Note Ser.*, pages 227–254. Cambridge Univ. Press, Cambridge, 1995.
- [9] Nigel Higson, John Roe, and Guoliang Yu. A coarse Mayer-Vietoris principle. *Math. Proc. Cambridge Philos. Soc.*, 114(1):85–97, 1993.
- [10] Gennadi Kasparov. Topological invariants of elliptic operators. I. K -homology. *Izv. Akad. Nauk SSSR Ser. Mat.*, 39(4):796–838, 1975.
- [11] Gennadi Kasparov. Equivariant KK -theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988.
- [12] Gennadi Kasparov and Guoliang Yu. The coarse geometric Novikov conjecture and uniform convexity. *Adv. Math.*, 206(1):1–56, 2006.
- [13] Gennadi Kasparov and Guoliang Yu. The Novikov conjecture and geometry of Banach spaces. *Geom. Topol.*, 16(3):1859–1880, 2012.
- [14] E. Christopher Lance. *Hilbert C^* -modules*, volume 210 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1995. A toolkit for operator algebraists.
- [15] Kang Li, Piotr Nowak, Ján Špakula, and Jiawen Zhang. Quasi-local algebras and asymptotic expanders. *Groups, Geometry, and Dynamics*, 2021.
- [16] Billy J. Pettis. On integration in vector spaces. *Trans. Amer. Math. Soc.*, 44(2):277–304, 1938.
- [17] Yu Qiao and John Roe. On the localization algebra of Guoliang Yu. *Forum Math.*, 22(4):657–665, 2010.
- [18] John Roe. An index theorem on open manifolds. I, II. *J. Differential Geom.*, 27(1):87–113, 115–136, 1988.
- [19] John Roe. *Coarse cohomology and index theory on complete Riemannian manifolds*, volume 497. American Mathematical Soc., 1993.

- [20] John Roe. *Index theory, coarse geometry, and topology of manifolds*, volume 90 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1996.
- [21] Lin Shan and Qin Wang. The coarse geometric Novikov conjecture for subspaces of non-positively curved manifolds. *J. Funct. Anal.*, 248(2):448–471, 2007.
- [22] Ján Špakula and Aaron Tikuisis. Relative Commutant Pictures of Roe Algebras. *Comm. Math. Phys.*, 365(3):1019–1048, 2019.
- [23] Ján Špakula and Jiawen Zhang. Quasi-locality and property A. *Journal of Functional Analysis*, 278(1):108299, 2020.
- [24] Rufus Willett and Guoliang Yu. *Higher index theory*, volume 189. Cambridge University Press, 2020.
- [25] Kôzoku Yosida. *Functional analysis*, volume 123 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin-New York, sixth edition, 1980.
- [26] Guoliang Yu. Coarse Baum-Connes conjecture. *K-Theory*, 9(3):199–221, 1995.
- [27] Guoliang Yu. Localization algebras and the coarse Baum-Connes conjecture. *K-Theory*, 11(4):307–318, 1997.
- [28] Guoliang Yu. The Novikov conjecture for groups with finite asymptotic dimension. *Ann. of Math. (2)*, 147(2):325–355, 1998.
- [29] Guoliang Yu. The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.*, 139(1):201–240, 2000.

SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, 220 HANDAN ROAD, SHANGHAI, 200433, CHINA.

Email address: xchen@fudan.edu.cn, 19110180024@fudan.edu.cn, jiawenzhang@fudan.edu.cn