

A thermodynamical description of third grade fluids mixtures

M. Gorgone and P. Rogolino

Department of Mathematical and Computer Sciences,
Physical Sciences and Earth Sciences, University of Messina
Viale F. Stagno d'Alcontres 31, 98166 Messina, Italy
mgorgone@unime.it; progolino@unime.it

Submitted to *J. Non-Equilib. Thermodyn.* (2021).

Abstract

A complete thermodynamical analysis for a non-reacting binary mixtures exhibiting the features of a third grade fluid is analyzed. The constitutive functions are allowed to depend on the mass density of the mixture and the concentration of one of the constituents, together with their first and second order gradients, on the specific internal energy of the mixture with its first order gradient, as well as on the symmetric part of the gradient of barycentric velocity. Compatibility with second law of thermodynamics is investigated by applying the extended Liu procedure. An explicit solution of the set of thermodynamic restrictions is obtained by postulating a suitable form of the constitutive relations for the diffusional mass flux, the heat flux and the Cauchy stress tensor. Taking a first order expansion in the gradients of the specific entropy, the expression of the entropy flux is determined. It includes an additional contribution due to non non-local effects.

Keywords. Binary mixtures with one temperature; Korteweg fluids; Exploitation of entropy principle; Extended Liu procedure.

1 Introduction

In the context of rational thermodynamics, a fundamental role is played by models of fluid continua; these models, in many applications, need to be represented by mixtures [1, 2, 3, 4], *i.e.*, continua supposed to be microscopically formed by two or more separate components having different specific physical properties.

In the general case, for a N -component mixture, we have to determine the evolution of the N partial mass densities $\rho^{(a)}$ ($a = 1, \dots, N$), the N velocities $\mathbf{v}^{(a)}$, and the N partial temperatures $\theta^{(a)}$ (or, equivalently, partial internal energies $\varepsilon^{(a)}$) of the constituents.

Mixtures can be modeled at various degrees of detail [5, 6, 7, 8, 9]. Here, we focus on a non-reacting model of a binary mixture described by means of the mass density, barycentric velocity and specific internal energy of the mixture, as well as the concentration of one constituent [2, 4]. More complex models can be considered in the case in which the constituents have different velocities, [8, 10, 11], or temperatures [5, 7]; recently, a complete thermodynamical analysis of a binary mixture of viscous Kortweg fluids with two velocities and two temperatures, has been developed in [12]. Moreover, in some physical problems, the description of mixtures requires the introduction of internal state variables as additional fields [10, 11, 13]; this case will not be faced hereafter.

In this paper, we consider a non-reacting mixture exhibiting the features of a third grade fluid, *i.e.*, the constitutive quantities possess second order non-localities [14]. Solving all the thermodynamical restrictions arising from the entropy principle, we prove that the constitutive equation for Cauchy stress tensor involves the second spatial derivatives of the mass density [15], *i.e.*, behaves like a Kortweg fluid.

Kortweg-type fluids have been considered in [16]; they are characterized by the following constitutive equation for the Cauchy stress tensor:

$$T_{ij} = (-p + \alpha_1 \rho_{,k} \rho_{,k} + \alpha_2 \rho_{,kk}) \delta_{ij} + \alpha_3 \rho_{,i} \rho_{,j} + \alpha_4 \rho_{,ij}, \quad (1)$$

where ρ denotes the mass density, p the pressure, δ_{ij} the Kronecker delta, and α_i ($i = 1, \dots, 4$) suitable material functions depending on mass density and temperature ($[\alpha_1] = [\alpha_3] = Kg^{-1}m^7s^{-2}$, $[\alpha_2] = [\alpha_4] = m^4s^{-2}$); moreover, the subscripts denote the partial derivatives and the Einstein convention on sums over repeated indices has been used.

These fluids have been investigated in a pioneeristic paper by Dunn and Serrin [15], in accordance with the principles of rational continuum thermodynamics [17]; moreover, they have been extensively studied by Cimmelli and coworkers [18, 19, 20, 21, 22, 23] through a generalized Liu procedure [24, 25], and by Heida and Málek [26] following a different methodology. In a recent paper [27], an explicit solution of the thermodynamical restrictions imposed by second law of thermodynamics in the general three-dimensional case of a viscous Korteweg fluid has been recovered.

The plan of the paper is as follows. In Section 2, we present the governing equations for the basic fields we choose to describe the mixture as a whole (mixture mass density, concentration of one constituent, barycentric velocity, specific internal energy), together with the entropy inequality; then, we sketch the extended Liu procedure applied for the exploitation of the entropy principle [21, 25], and recover a set of conditions ensuring that second law of thermodynamics to be satisfied for arbitrary thermodynamical processes. In Section 3, we assume some constitutive relations in order to solve all the thermodynamical restrictions and provide an explicit, physically admissible, solution. The lengthy computations necessary to derive and solve the algebraic and differential conditions are handled with the help of some symbolic routines written in the Computer Algebra System Reduce [28]. As a result, it is proved that the constitutive equations are compatible with the second law of thermodynamics; in particular, the Cauchy stress tensor for the mixture has been characterized in a very general form typical of Korteweg fluids [27]. Finally, Section 4 contains our conclusions as well as further developments of the present theory.

2 Basic fields and extended Liu procedure

Let us consider a non-reacting binary mixture of third grade fluids and choose the fundamental fields to describe the mixture as a whole: the mass density $\rho(t, x_j)$ of the mixture, the concentration $c(t, x_j)$ of one constituent [2, 4], the barycentric velocity $\mathbf{v}(t, x_j) \equiv (v_1, v_2, v_3)$, and the specific internal energy $\varepsilon(t, x_j)$ of the mixture. As usual in the study of mixtures, mass density of the whole mixture and the concentration of one constituent are given by

$$\rho = \rho^{(1)} + \rho^{(2)}, \quad c = \frac{\rho^{(1)}}{\rho}, \quad (2)$$

where the superscript (a) ($a = 1, 2$) labels the two components.

Therefore, in the absence of external forces and heat sources, we are led to consider the following governing equations for our mixture, in components:

$$\begin{aligned}
\rho_{,t} + \rho_{,j}v_j + \rho v_{j,j} &= 0, \\
\rho(c_{,t} + v_j c_{,j}) + J_{j,j}^{(m)} &= 0, \\
\rho(v_{i,t} + v_j v_{i,j}) - T_{ij,j} &= 0, \\
\rho(\varepsilon_{,t} + v_j \varepsilon_{,j}) - T_{ij}v_{i,j} + q_{j,j} &= 0,
\end{aligned} \tag{3}$$

where $J_j^{(m)}$, T_{ij} and q_j denote the components of the diffusional mass flux, the symmetric Cauchy stress tensor and the heat flux for the mixture, respectively.

Then, in our general framework, in order to ensure the compatibility of this model with the second law of thermodynamics, we have to impose that the local entropy production for the mixture,

$$\rho(s_{,t} + v_j s_{,j}) + J_{j,j}^{(s)} \geq 0, \tag{4}$$

be non-negative along any admissible thermodynamic process, where s is the specific entropy, and $J_j^{(s)}$ are the components of the entropy flux.

The field equations (3) and the entropy inequality (4), once the variables entering the state space have been assigned, must be closed by the constitutive equations for the diffusional mass flux, Cauchy stress tensor, heat flux, specific entropy and entropy flux. Constitutive theories require that all the constitutive functions must depend on a state variables; consequently, let us assume the state space, containing second order non-localities, to be spanned by

$$\mathcal{Z} = \{\rho, c, \varepsilon, \rho_{,j}, c_{,j}, L_{ij}, \varepsilon_{,j}, \rho_{,ij}, c_{,ij}\}, \tag{5}$$

where $L_{ij} = \frac{v_{i,j} + v_{j,i}}{2}$ is the symmetric part of the barycentric velocity gradient.

Since we need to find a set of conditions which are at least sufficient for the fulfillment of the entropy inequality (4) for arbitrary thermodynamical processes [29, 30], we apply the extended Liu procedure [21, 25], incorporating new restrictions consistent with higher order non-local constitutive theories. According to this procedure, in the entropy inequality, we have to use as constraints the field equations, and their gradient extensions too,

up to the order of the gradients entering the state space, by means of suitable Lagrange multipliers. This methodology is motivated by the fact that thermodynamical processes are solutions of the field equations, and, if these solutions are smooth enough, are trivially solutions of their differential consequences [31]. Since the entropy inequality (4) has to be satisfied in arbitrary smooth processes, then, from a mathematical viewpoint, we have to use also the differential consequences of the equations governing those processes as constraints for such an inequality. On the contrary, if we limit ourselves to consider as constraints only the field equations, we are led straightforwardly to a specific entropy, Lagrange multipliers and Cauchy stress tensor which do not depend upon the gradients of the variables entering the state space [9, 23, 27], and, hence, losing the typical features of Korteweg fluids.

In order to exploit second law of thermodynamics, we take into account the constraints imposed on entropy inequality by the field equations and their gradient extensions; this task is accomplished by introducing the Lagrange multipliers $\lambda^{(1)}$, $\lambda^{(2)}$, $\lambda_i^{(3)}$, $\lambda^{(4)}$, $\Lambda_i^{(1)}$, $\Lambda_i^{(2)}$, $\Lambda_{ij}^{(3)}$, $\Lambda_i^{(4)}$, $\Theta_{ij}^{(1)}$ and $\Theta_{ij}^{(2)}$ ($i, j = 1, 2, 3$), depending on the state space variables. Thus, the entropy inequality writes

$$\begin{aligned}
& \rho(s_{,t} + v_j s_{,j}) + J_{j,j}^{(s)} - \left(\lambda^{(1)} + \Lambda_i^{(1)} \frac{\partial}{\partial x_i} + \Theta_{ik}^{(1)} \frac{\partial^2}{\partial x_i \partial x_k} \right) (\rho_{,t} + \rho_{,j} v_j + \rho v_{j,j}) \\
& - \left(\lambda^{(2)} + \Lambda_i^{(2)} \frac{\partial}{\partial x_i} + \Theta_{ik}^{(2)} \frac{\partial^2}{\partial x_i \partial x_k} \right) \left(\rho(c_{,t} + v_j c_{,j}) + J_{j,j}^{(m)} \right) \\
& - \left(\lambda_i^{(3)} + \Lambda_{ik}^{(3)} \frac{\partial}{\partial x_k} \right) (\rho(v_{i,t} + v_j v_{i,j}) - T_{ij,j}) \\
& - \left(\lambda^{(4)} + \Lambda_k^{(4)} \frac{\partial}{\partial x_k} \right) (\rho(\varepsilon_{,t} + v_j \varepsilon_{,j}) - T_{ij} v_{i,j} + q_{j,j}) \geq 0,
\end{aligned} \tag{6}$$

that needs to be expanded by means of the chain rule; we perform the task by using some routines written in the Computer Algebra System Reduce [28], and, due to the long computations, we omit to report the full form here. The main advantages of using symbolic computation are that we are able to extract the coefficients of a multivariate polynomial in some derivatives of the field variables, and then solve, with the help of the Crack package [32], the set of differential and algebraic conditions for the unknown constitutive functions.

When the inequality (6) is expanded, we can distinguish the *highest derivatives* and the *higher derivatives* [20]. The formers are both the time

derivatives of the field variables and of the elements of the state space, which cannot be expressed through the governing equations as functions of the thermodynamical variables, and the spatial derivatives whose order is the highest one. On the contrary, the latter are the spatial derivatives whose order is not maximal but higher than that of the gradients entering the state space. Due to specific choice of the state space (5), the components of the highest derivatives consist of

$$\mathbf{X} = \{\rho_{,t}, c_{,t}, v_{i,t}, \varepsilon_{,t}, \rho_{,it}, c_{,it}, v_{i,kt}, \varepsilon_{,kt}, \rho_{,ikt}, c_{,ikt}, v_{i,jklm}, \varepsilon_{,ijkl}, \rho_{,ijklm}, c_{,ijklm}\}, \quad (7)$$

whereas the components of the higher ones are

$$\mathbf{Y} = \{\rho_{,jkl}, c_{,jkl}, v_{i,jk}, \varepsilon_{,jk}, \rho_{,ijkl}, c_{,ijkl}, v_{i,jkl}, \varepsilon_{,ijk}\}. \quad (8)$$

By lengthy though straightforward computation, it is easily ascertained that the entropy inequality (6) can be written in the compact form

$$A_p X_p + B_{qrs}^{(3)} Y_q Y_r Y_s + B_{qr}^{(2)} Y_q Y_r + B_q^{(1)} Y_q + C \geq 0, \quad (9)$$

where the functions A_p , $B_q^{(1)}$, $B_{qr}^{(2)}$, $B_{qrs}^{(3)}$ and C depend at most on the field and state variables; this means that the left hand side of the inequality (9) is a polynomial which is linear in the highest derivatives and cubic in the higher ones. Because of the constraints we imposed, the highest and higher derivatives may assume arbitrary values [21].

Since in principle nothing prevents the possibility of a thermodynamical process where $C = 0$, and in order the inequality (9) to be satisfied for arbitrary X_p and Y_q , the conditions

$$A_p = 0, \quad B_{qrs}^{(3)} = 0, \quad B_q^{(1)} = 0, \quad C \geq 0, \quad (10)$$

as well the requirement that the quantities $B_{qr}^{(2)}$ are the entries of a positive semidefinite symmetric matrix, are sufficient for the fulfillment of the entropy inequality.

From $A_p = 0$, we obtain the expressions for the components of Lagrange

multipliers,

$$\begin{aligned}
\lambda^{(1)} &= \rho \frac{\partial s}{\partial \rho}, & \lambda^{(2)} &= \frac{\partial s}{\partial c} - \frac{\rho, k}{\rho} \left(\frac{\partial s}{\partial c, k} - 2 \frac{\rho, i}{\rho} \frac{\partial s}{\partial c, ik} \right) - \frac{\rho, ik}{\rho} \frac{\partial s}{\partial c, ik}, & \lambda_i^{(3)} &= -\frac{\rho, k}{\rho} \frac{\partial s}{\partial v_{i, k}}, \\
\lambda^{(4)} &= \frac{\partial s}{\partial \varepsilon} - \frac{\rho, k}{\rho} \frac{\partial s}{\partial \varepsilon, k}, & \Lambda_k^{(1)} &= \rho \frac{\partial s}{\partial \rho, k}, & \Lambda_k^{(2)} &= \frac{\partial s}{\partial c, k} - 2 \frac{\rho, i}{\rho} \frac{\partial s}{\partial c, ik}, \\
\Lambda_{ik}^{(3)} &= \frac{\partial s}{\partial v_{i, k}}, & \Lambda_k^{(4)} &= \frac{\partial s}{\partial \varepsilon, k}, & \Theta_{ik}^{(1)} &= \rho \frac{\partial s}{\partial \rho, ik}, & \Theta_{ik}^{(2)} &= \frac{\partial s}{\partial c, ik},
\end{aligned} \tag{11}$$

as well as the following restrictions involving the specific entropy and the diffusional mass flux:

$$\begin{aligned}
\left\langle \frac{\partial s}{\partial c, kl} \frac{\partial J_j^{(m)}}{\partial v_{n, i}} \right\rangle_{(ijk\ell)} &= 0, & \left\langle \frac{\partial s}{\partial c, kl} \frac{\partial J_j^{(m)}}{\partial \varepsilon, i} \right\rangle_{(ijk\ell)} &= 0, \\
\left\langle \frac{\partial s}{\partial c, kl} \frac{\partial J_j^{(m)}}{\partial \rho, in} \right\rangle_{(ijk\ell n)} &= 0, & \left\langle \frac{\partial s}{\partial c, kl} \frac{\partial J_j^{(m)}}{\partial c, in} \right\rangle_{(ijk\ell n)} &= 0,
\end{aligned} \tag{12}$$

where the symbol $\langle \mathcal{F} \rangle_{(i_1 \dots i_r)}$ denotes the symmetric part of the tensor function \mathcal{F} with respect to the indices $i_1 \dots i_r$.

The further thermodynamical restrictions, even if their computation is straightforward, are omitted since their expression is rather long. Furthermore, by analyzing the conditions which vanish the linear terms in the higher and highest derivatives, *i.e.*, $B_q^{(1)} = 0$ and $B_{qrs}^{(3)} = 0$, it follows that the entropy flux is no longer given by the ratio between the heat flux and the absolute temperature of the mixture, and extra-terms can be obtained.

In the next section, all the thermodynamical constraints recovered by applying the extended Liu procedure are solved, and explicit constitutive equations for the diffusional mass flux, the Cauchy stress tensor, the heat flux, the specific entropy and the entropy flux are determined. The achieved results provide a complete and physically admissible solution.

3 A solution of thermodynamical restrictions

All the thermodynamical restrictions, derived performing the extended Liu procedure, result too much general for being useful in concrete applications; therefore, a further simplification is necessary in order to exploit the entropy

inequality. Under the hypothesis that the non-reacting binary mixture of third grade fluids acts as a viscous Korteweg fluid, let us choose the following ansatz for the Cauchy stress tensor

$$T_{ij} = (\tau_0 + \tau_1 \rho_{,k} \rho_{,k} + \tau_2 \rho_{,k} c_{,k} + \tau_3 c_{,k} c_{,k} + \tau_4 \rho_{,kk} + \tau_5 c_{,kk} + \tau_6 v_{k,k}) \delta_{ij} \\ + \tau_7 \rho_{,i} \rho_{,j} + \tau_8 \frac{\rho_{,i} c_{,j} + \rho_{,j} c_{,i}}{2} + \tau_9 c_{,i} c_{,j} + \tau_{10} \rho_{,ij} + \tau_{11} c_{,ij} + \tau_{12} L_{ij}; \quad (13)$$

moreover, let us consider the following constitutive relations for the diffusional mass and heat fluxes:

$$J_j^{(m)} = J_\rho \rho_{,j} + J_c c_{,j}, \quad q_j = q \varepsilon_{,j}, \quad (14)$$

where τ_k ($k = 0, \dots, 12$), J_ρ , J_c , and q are suitable scalar material functions depending at most on (ρ, c, ε) .

In addition, let us expand the specific entropy s around the homogeneous equilibrium state (where all gradients vanish), at the first order on the gradients of mass density of the whole mixture and concentration of one constituent, *i.e.*,

$$s = \hat{s}_0 + \hat{s}_1 \rho_{,k} \rho_{,k} + \hat{s}_2 \rho_{,k} c_{,k} + \hat{s}_3 c_{,k} c_{,k}, \quad (15)$$

where $\hat{s}_i \equiv \hat{s}_i(\rho, c, \varepsilon)$ ($i = 0, \dots, 3$) are some scalar functions of the indicated arguments. We are aware that expression (15) is not the most general representation of the entropy density as an isotropic scalar function, but it is enough for our purposes.

On the basis of the assumptions on constitutive relations (13)-(15), we immediately note that restrictions (12) are identically satisfied; the remaining conditions ($B_q^{(1)} = 0$, $B_{qrs}^{(3)} = 0$) provide a large set of partial differential equations that we are able to solve using some routines written in the Computer Algebra System Reduce [28]. Firstly, the coefficients entering the specific entropy and the diffusional mass flux have been recovered:

$$\hat{s}_0 = s_{01} + J_c s_{02} + s_{03}, \quad \hat{s}_1 = \frac{J_\rho^2 - \kappa_2^2}{J_c^2} \hat{s}_3 + \frac{\phi'}{\rho}, \quad \hat{s}_2 = 2 \frac{J_\rho}{J_c} \hat{s}_3, \quad \hat{s}_3 = \frac{s_3}{\rho}, \\ J_\rho = \kappa_1 c + \kappa_2, \quad J_c = -\kappa_1 \rho + \kappa_3, \quad (16)$$

with the prime $'$ denoting differentiation with respect to the argument, along with the functions $s_{01} \equiv s_{01}(\varepsilon)$, $s_{02} \equiv s_{02}\left(\frac{J_\rho}{J_c}\right)$, $s_{03} \equiv s_{03}(\rho)$, $\phi \equiv \phi(\rho)$, κ_i

($i = 1, 2, 3$) and s_3 constant. In order to guarantee the principle of maximum entropy at equilibrium the quadratic part in the gradients of (15) must be negative semidefinite, whence

$$\frac{\kappa_2^2 s_3}{(\kappa_3 - \kappa_1 \rho)^2} - \phi' \geq 0, \quad s_3 \leq 0. \quad (17)$$

As far as the absolute temperature θ of the mixture is concerned, we have

$$\frac{1}{\theta} = \frac{\partial \hat{s}_0}{\partial \varepsilon} = \frac{ds_{01}}{d\varepsilon}; \quad (18)$$

with the hypothesis that $\frac{d^2 s_{01}}{d\varepsilon^2} \neq 0$, by using the implicit function theorem, it follows that the specific internal energy ε depends only on the absolute temperature θ of the whole mixture, thus the heat flux takes the form

$$q_j = q \frac{d\varepsilon}{d\theta} \theta_{,j}, \quad (19)$$

i.e., we recognize the classical Fourier law of heat conduction, wherein the coefficient $q \frac{d\varepsilon}{d\theta}$ is the opposite of the thermal conductivity.

Moreover, we also obtain the following expressions for the material functions entering the Cauchy stress tensor:

$$\begin{aligned} \tau_0 &= \rho^2 \frac{\partial \hat{s}_0}{\partial \rho} \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = \rho^2 \theta \left(-\kappa_1 \left(s_{02} - \frac{J_\rho}{J_c} s'_{02} \right) + s'_{03} \right), \\ \tau_1 &= -\frac{\partial(\rho^2 \hat{s}_1)}{\partial \rho} \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = -\theta \left(\frac{(J_\rho^2 - \kappa_2^2)(J_c + 2\kappa_1 \rho)}{J_c^3} s_3 + \frac{d(\rho \phi')}{d\rho} \right), \\ \tau_2 &= -\left(\frac{\partial(\rho^2 \hat{s}_2)}{\partial \rho} + \rho^2 \frac{\partial \hat{s}_1}{\partial c} \right) \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = -2\theta \frac{J_\rho}{J_c^2} (J_c + 2\kappa_1 \rho) s_3, \\ \tau_3 &= -\rho \left(\rho \frac{\partial \hat{s}_2}{\partial c} + \hat{s}_3 \right) \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = -\theta \frac{J_c + 2\kappa_1 \rho}{J_c} s_3, \\ \tau_4 &= -2\rho^2 \hat{s}_1 \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = -2\rho \theta \left(\frac{J_\rho^2 - \kappa_2^2}{J_c^2} s_3 + \phi' \right), \\ \tau_5 &= -\rho^2 \hat{s}_2 \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = -2\rho \theta \frac{J_\rho}{J_c} s_3, \quad \tau_7 = 2\rho \hat{s}_1 \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = 2\theta \left(\frac{J_\rho^2 - \kappa_2^2}{J_c^2} s_3 + \phi' \right), \\ \tau_8 &= 2\rho \hat{s}_2 \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = 4\theta \frac{J_\rho}{J_c} s_3, \quad \tau_9 = 2\rho \hat{s}_3 \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right)^{-1} = 2\theta s_3, \quad \tau_{10} = \tau_{11} = 0, \end{aligned}$$

and it is recovered for entropy flux the following form

$$J_j^{(s)} = \frac{q_j}{\theta} + \frac{\partial \hat{s}_0}{\partial c} J_j^{(m)} + (\hat{s}_2 (J_\rho \rho_{,kk} + J_c c_{,kk}) + 2\rho^2 \hat{s}_1 v_{k,k}) \rho_{,j} + (2\hat{s}_3 (J_\rho \rho_{,kk} + J_c c_{,kk}) + \rho^2 \hat{s}_2 v_{k,k}) c_{,j}, \quad (20)$$

where the relation (16) has been used.

In the entropy flux (20) we recognize the contribution of the classical term, the ratio between the heat flux and the absolute temperature of the whole mixture, and an entropy extra-flux [33] depending on the divergence of the barycentric velocity, together with the first and second order gradients of the mixture mass density and concentration of one constituent.

The matrix $B_{qr}^{(2)}$, with the above results, identically vanishes, and the entropy inequality reduces to

$$q_j \frac{\partial}{\partial x_j} \left(\frac{\partial \hat{s}_0}{\partial \varepsilon} \right) + (\tau_6 (v_{k,k})^2 + \tau_{12} L_{ij} L_{ij}) \frac{\partial \hat{s}_0}{\partial \varepsilon} + 2\hat{s}_3 \frac{(J_\rho \rho_{,kk} + J_c c_{,kk})^2}{J_c} + J_\rho \frac{\partial^2 \hat{s}_0}{\partial \rho \partial c} \rho_{,k} \rho_{,k} + 2J_\rho \frac{\partial^2 \hat{s}_0}{\partial c^2} \rho_{,k} c_{,k} + J_c \frac{\partial^2 \hat{s}_0}{\partial c^2} c_{,k} c_{,k} \geq 0. \quad (21)$$

The residual entropy inequality (21) turns out to be a homogeneous quadratic polynomial in some gradients entering the state space, whose coefficients depend at most on the field variables; such a relation is fulfilled for all the thermodynamical processes if and only if the following inequalities hold true:

$$q \frac{\partial^2 \hat{s}_0}{\partial \varepsilon^2} \geq 0, \quad J_c \frac{\partial^2 \hat{s}_0}{\partial c^2} \geq 0, \quad J_c \hat{s}_3 \geq 0, \quad \tau_6 \frac{\partial \hat{s}_0}{\partial \varepsilon} \geq 0, \quad \tau_{12} \frac{\partial \hat{s}_0}{\partial \varepsilon} \geq 0. \quad (22)$$

Due to (16)-(17) and

$$\frac{\partial \hat{s}_0}{\partial \varepsilon} = s'_{01} = \frac{1}{\theta} > 0, \quad \frac{\partial^2 \hat{s}_0}{\partial \varepsilon^2} = s''_{01} = -\frac{1}{\theta^2} \theta' < 0, \quad \frac{\partial^2 \hat{s}_0}{\partial c^2} = \frac{\kappa_1^2}{J_c} s''_{02}, \quad (23)$$

it follows that the inequalities (22) are satisfied if

$$q \leq 0, \quad J_c \leq 0, \quad s''_{02} \geq 0, \quad \tau_6 \geq 0, \quad \tau_{12} \geq 0; \quad (24)$$

we notice that the viscosity coefficients, τ_6 and τ_{12} , result non-negative as physics prescribes.

Finally, for the three-dimensional case of a binary non-reacting mixture that acts as a viscous Korteweg fluid, described by the mass density of the

mixture, the concentration of one constituent, the barycentric velocity and the specific internal energy of the whole mixture, all the thermodynamical restrictions arising from the exploitation of the entropy principle are solved. Furthermore, all the results above detailed allow us to characterize the constitutive functions involved in the model.

4 Conclusions

In this paper, we considered a non-reacting binary mixtures of viscous third grade fluids with one temperature, and analyzed the compatibility of the constitutive functions with the second law of thermodynamics. Because of the non-locality of the constitutive equations, we exploited the entropy principle by means of the extended Liu procedure, which consists in taking into account the field equations and their gradient extensions. Firstly, we introduced the basic fields to describe the mixture as a whole, *i.e.*, the mass density of the mixture, the concentration of one constituent, the barycentric velocity, and the specific internal energy of the mixture and, then, the related governing equations. We assumed a state space containing, as well as the first order gradients of the basic fields, also the second order ones of mixture mass density and concentration. Therefore, by applying the procedure, we were able to characterize the non-local constitutive equations guaranteeing the compatibility with second law of thermodynamics. All the straightforward and long computation have been managed by using some symbolic routines written in the Computer Algebra System Reduce [28].

The present model exhibits a smaller level of detail than the case with two temperatures analyzed in [9, 12]; nevertheless, the diffusive effects are described by a diffusional mass flux recovered from the thermodynamic restrictions.

As a consequence of the algorithmic procedure, after expanding around the equilibrium the specific entropy, retaining only first order terms in the gradients of the mixture mass density and concentration of one constituent, the expression of the entropy flux naturally arises. The latter contains the classical part and an extra-flux depending on the first and second order gradients of mixture mass density and concentration, and on the divergence of barycentric velocity.

It is worth of being remarked that we do not assume *a priori* a modified energy equation including additional contributions (*i.e.*, no interstitial

working [15]) and do not postulate the functional form of entropy flux.

Remarkably, we were able to explicitly solve the thermodynamical constraints by obtaining a Cauchy stress tensor that is sufficiently general to include that proposed by Korteweg in [16], and prove that the mixture's behavior is typical of a viscous Korteweg fluid [27].

The theoretical results above derived contain some degrees of freedom and may serve as a basis for experimental and/or numerical investigations. Further application of the theory could be to investigate diffusive binary mixtures of Korteweg-type fluids with one temperature, where the state space includes the diffusion velocity; as a consequence, the diffusional mass flux is not a constitutive function, and the thermodynamical constraints involve a contribution due to the diffusion velocity.

Acknowledgments

Work supported by G.N.F.M. of the Istituto Nazionale di Alta Matematica “F. Severi”. M.G. acknowledges the support through the “Progetto Giovani GNFM 2020”.

References

- [1] I. Müller, A thermodynamic theory of mixtures of fluids, *Arch. Rational Mech. Anal.* **28** (1968), 1–39 .
- [2] M. E. Gurtin and A. S. Vargas, On the classical theory of reacting fluid mixtures, *Arch. Rational Mech. Anal.* **43** (1971), 179–197.
- [3] R. M. Bowen, Theory of mixtures, Part I, in: *A. C. Eringen (ed), Continuum Physics*, Academic Press, New York (2002), 1–127.
- [4] I.-S Liu and I. Müller, Thermodynamics of mixtures of fluids, in: *C. Truesdell, Rational Thermodynamics*, Springer-Verlag (1984), 264–285.
- [5] R. M. Bowen and D. J. Garcia, On the thermodynamics of mixtures with several temperatures, *Int. J. Eng. Sci.* **8** (1970), 63–83.

- [6] A. Palumbo, C. Papenfuss and P. Rogolino, A mesoscopic approach to diffusion phenomena in mixtures, *J. Non-Equilib. Thermodyn.* **30** (2005), 401–419.
- [7] H. Gouin and T. Ruggeri, Identification of an average temperature and a dynamical pressure in a multitemperature mixture of fluids, *Phys. Rev. E* **78** (2008), 016303-1-7.
- [8] D. Bothe and W. Dreyer, Continuum thermodynamics of chemically reacting fluid mixtures, *Acta Mechanica* **226** (2015), 1757–1805.
- [9] V. A. Cimmelli, M. Gorgone, F. Oliveri and A. R. Pace, Weakly non-local thermodynamics of binary mixtures of Korteweg fluids with two velocities and two temperatures, *European J. Mech./B Fluids* **83** (2020), 58–65.
- [10] M. Francaviglia, A. Palumbo and P. Rogolino, Thermodynamics of mixtures as a problem with internal variables. The general theory, *J. Non-Equilib. Thermodyn.* **31** (2006), 419–429.
- [11] M. Francaviglia, A. Palumbo and P. Rogolino, Internal variables thermodynamics of two component mixtures under linear constitutive hypothesis with an application to superfluid helium, *J. Non-Equilib. Thermodyn.* **33** (2008), 149–164.
- [12] M. Gorgone, F. Oliveri and P. Rogolino, Thermodynamical analysis and constitutive equations for a mixture of viscous Korteweg fluids, submitted (2021).
- [13] F. Oliveri, A. Palumbo and P. Rogolino, On a model of mixtures with internal variables: extended Liu procedure for the exploitation of the entropy principle, *Atti Accad. Pelorit. Pericol. Cl. Sci. Fis. Mat. Nat.* **94** (2016), A2-1-17.
- [14] C. Truesdell and K. R. Rajagopal, *An Introduction to the Mechanics of Fluids*, Birkhäuser, Boston–Basel–Berlin, 2000.
- [15] J. E. Dunn and J. Serrin, On the thermomechanics of the interstitial working, *Arch. Rational Mech. Anal.* **88** (1985), 95–133.

- [16] D. J. Korteweg, Sur la forme qui prennent les équations du mouvement des fluids si l'on tient compte des forces capillaires par des variations de densité considérables mais continues et sur la théorie de la capillarité dans l'hypothèse d'une variation continue de la densité, *Archives Néerlandaises des sciences exactes et naturelles* **6** (1901), Ser. II, 1–24.
- [17] C. Truesdell, *Rational thermodynamics*, Springer-Verlag, 1984.
- [18] V. A. Cimmelli, F. Oliveri and A. R. Pace, On the thermodynamics of Korteweg fluids with heat conduction and viscosity, *J. Elast.* **104** (2011), 115–131.
- [19] V. A. Cimmelli, A. Sellitto and V. Triani, A new perspective on the form of the first and second laws in rational thermodynamics: Korteweg fluids as an example, *J. Non-Equilib. Thermodyn.* **35** (2010), 251–265.
- [20] V. A. Cimmelli, A. Sellitto and V. Triani, A new thermodynamic framework for second-grade Korteweg-type fluids, *J. Math. Phys.* **50** (2009), 053101-1-16.
- [21] V. A. Cimmelli, F. Oliveri and V. Triani, Exploitation of the entropy principle: proof of Liu theorem if the gradients of the governing equations are considered as constraints, *J. Math. Phys.* **52** (2011), 023511-1-15.
- [22] V. A. Cimmelli, F. Oliveri and A. R. Pace, A nonlocal phase-field model of Ginzburg–Landau–Korteweg fluids, *Contin. Mech. Thermodyn.* **27** (2015), 367–378.
- [23] M. Gorgone, F. Oliveri and P. Rogolino, Continua with non-local constitutive laws: Exploitation of entropy inequality, *Int. J. Non-Linear Mech.* **126** (2020), 103573.
- [24] I.-S. Liu, Method of Lagrange multipliers for exploitation of the entropy principle, *Arch. Rational Mech. Anal.* **46** (1972), 131–148.
- [25] V. A. Cimmelli, An extension of Liu procedure in weakly nonlocal thermodynamics, *J. Math. Phys.* **48** (2007), 113510-1-13.
- [26] M. Heida and J. Málek, On Korteweg-type compressible fluid-like materials, *Int. J. Eng. Sci.* **48** (2010), 1313–1324.

- [27] M. Gorgone and P. Rogolino, On the characterization of constitutive equations for third-grade viscous Korteweg fluids, *Phys. Fluids* **33** (2021), 043107.
- [28] A. C. Hearn, *REDUCE user's manual, version 3.8* (Technical report, Rand Corporation, Santa Monica, CA, USA, 1995).
- [29] V. A. Cimmelli, D. Jou, T. Ruggeri and P. Ván, Entropy principle and recent results in non-equilibrium theories, *Entropy* **16** (2014), 1756–1807.
- [30] D. Jou, J. Casas-Vázquez and G. Lebon, *Extended irreversible thermodynamics*, fourth revised ed., Springer, 2010.
- [31] P. Rogolino and V. A. Cimmelli, Differential consequences of balance laws in extended irreversible thermodynamics of rigid heat conductors, *Proc. R. Soc. A* **16** (2019), 20180482-1-19.
- [32] T. Wolf, Investigating differential equations with CRACK, LiePDE, Ap-plysym and ConLaw, in: *J. Grabmeier, E. Kaltofen, V. Weispfenning (eds.), Handbook of Computer Algebra, Foundations, Applications, Systems*, Springer, New York (2002), 465–468.
- [33] I. Müller, On the entropy inequality, *Arch. Rational Mech. Anal.* **26** (1967), 118–141.