

Insertion and Lie Bracket Concerning Finite Sets

Zhou Mai *

September 6, 2023

Abstract

In this article we discuss the operations of partitions (sequence of disjoint finite subsets) which are quotient, insertion, composition and Lie bracket. Moreover, we discuss applications of those operations for Feynman diagrams and Kontsevich's graphs.

Keywords : partition, quotient, insertion, Lie bracket, Feynman diagrams, admissible graphs.

Contents

1	Introduction	1
2	Quotient and insertion of the partitions	2
2.1	Notations concerning the partitions	2
2.2	Quotient of the partitions	3
2.3	Insertion of the partitions	8
2.4	Coproduct	12
3	Composition and Lie bracket	16
4	Applications to graphs	19
4.1	The graphs in the sense of graphic theory	20
4.2	Feynman diagrams	23
4.3	Kontsevich graphs	26

1 Introduction

In this article we discuss some operations of partitions, where the partition means the sequence of disjoint finite subsets. The operations what we focus on include quotient, insertion, composition and Lie bracket. All of operation is generalization of ones concerning Feynman diagrams (see A. Connes and D. Kreimer [1, 2], A. Connes and M. Marcolli [3], D. Kreimer [6]) and Kontsevich's graphs (see L. M. Ionescu [4], M. Kontsevich [7]). The construction in this article is suitable for the cases of Feynman diagrams and Kontsevich's graphs, actually, if we consider some structure maps additionally, the operations concerning Feynman diagrams and Kontsevich's graphs can be reduced to our construction. Here the construction of quotient follows the ideas in Zhou Mai [9], but, some modification occurs such that it is more suitable for the cases of Feynman

*address: Colleague of Mathematical Science, Nankai University, Weijin Road, Tianjin City, Republic China;
email address: zhoumai@nankai.edu.cn

diagrams and Kontsevich's graphs. Our construction is suitable for ordinary graphs in the sense of graphic theory as well. Somehow, a ordinary graph can be regarded as a Feynman diagram without external lines, but, the case of subgraph is different. It seems that three types of above graphs can be dealt with in an uniform way.

This paper is organized as follows. In section 2 we discuss the quotient and insertion of partitions in details. Based on the quotient we construct the coproduct which will result in a hopf algebra, but we do not discuss this issue more. In section 3 we construct the composition and Lie bracket. Here two types of composition are considered, both of them will result in well defined Lie bracket. Finally, in section 4 we discuss the cases of Feynman diagrams, Kontsevich's graphs and ordinary graphs starting from our construction.

2 Quotient and insertion of the partitions

2.1 Notations concerning the partitions

Partitions: Firstly, we introduce some notations which will be useful for discussion below.

- For a finite set A , we denote the power set of A by $\mathcal{P}(A)$. The reversion map $\mathcal{R} : \mathcal{P}(\mathcal{P}(A)) \rightarrow \mathcal{P}(A)$ is defined to be:

$$\mathcal{R}(\{I_i\}) = \bigcup_i I_i, \{I_i\} \in \mathcal{P}(\mathcal{P}(A)).$$

Let $\mathcal{P}_{dis}^2(A)$ denote a subset of $\mathcal{P}(\mathcal{P}(A))$,

$$\mathcal{P}_{dis}^2(A) = \{\{I_i\} \in \mathcal{P}(\mathcal{P}(A)) | \{I_i\} \in \mathbf{Part}(\mathcal{R}(\{I_i\}))\}.$$

We call the element $\{I_i\} \in \mathcal{P}_{dis}^2(A)$ the partition in A . If we ignore the order of $\{I_i\}$, then we identify $\{I_i\}$ with $\{I_{\sigma(i)}\}$, where $\sigma \in \mathbb{S}_m$ and $m = |\{I_i\}|$ ($|B|$ denotes the number of the elements in a finite set B).

A partition $\{I_i\}_{i=1}^m \in \mathcal{P}_{dis}^2(A)$ can be decribed by a function f from $I = \mathcal{R}(\{I_i\})$ to \mathbb{N} with $|Im(f)| = m$. Precisely, let $Im(f) = \{i_1, \dots, i_m\}$ ($0 < i_1 < \dots < i_m$), then $\{f^{-1}(i_k)\}_{k=1}^m$ is a partition in A . If $f^{-1}(i_k) = I_k$, then the function f defines the partition $\{I_i\}_{i=1}^m$. For a permutation $\sigma : \{i_1, \dots, i_m\} \rightarrow \{i_{\sigma(1)}, \dots, i_{\sigma(m)}\}$, it is obvious that $\sigma \circ f$ defines a same partition. On the other hand, let τ be a map from $\{i_1, \dots, i_m\}$ to $\underline{m} = \{1, \dots, m\}$, $\tau(i_k) = k$ ($k = 1, \dots, m$), then $\tau \circ f$ defines the same partition also. Without loss of generality, we can always assume $Im(f) = \underline{m}$. We call the function $f : I \rightarrow \underline{m}$ satisfying $f^{-1}(i) = I_i$ ($i = 1, \dots, m$) the defining function of the partition $\{I_i\}_{i=1}^m$ denoted by $f_{\{I_i\}}$.

- Let $\{I_i\}_{i=1}^m, \{J_j\}_{j=1}^n \in \mathcal{P}_{dis}^2(A)$ with defining functions $f_{\{I_i\}}, f_{\{J_j\}}, \mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) = \emptyset$, Then $\{I_1, \dots, I_m, J_1, \dots, J_n\} \in \mathcal{P}_{dis}^2(A)$, the new partition as above is denoted by $\{I_i\} \cup \{J_j\}$. Let $f_{\{I_i\} \cup \{J_j\}}$ be the defining function of $\{I_i\} \cup \{J_j\}$, then, $f_{\{I_i\} \cup \{J_j\}}|_I = f_{\{I_i\}}$, $f_{\{I_i\} \cup \{J_j\}}|_J = \iota \circ f_{\{J_j\}}$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$ and $\iota(j) = j + m$ ($j = 1, \dots, n$). We denote $f_{\{I_i\} \cup \{J_j\}}$ by $f_{\{I_i\}} \cup f_{\{J_j\}}$ also. It is obvious that $f_{\{J_j\}} \cup f_{\{I_i\}}$ defines same partition, therefore, we will identify $f_{\{J_j\}} \cup f_{\{I_i\}}$ with $f_{\{I_i\}} \cup f_{\{J_j\}}$.
- Let $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, we say $\{J_j\} \subset \{I_i\}$, if for any j , there is an i , such that $J_j \subset I_i$. An important situation is $\{J_j\} = \{I_i \cap B\}$, where $B \subset \mathcal{R}(\{I_i\})$. It is obvious that $\mathcal{R}(\{J_j\}) = B$. In this situation we call $\{J_j\}$ the restriction of $\{I_i\}$ on B , denoted by $\{J_j\} = \{I_i\}_{(B)}$. The defining function of $\{J_j\}$ can be determined as follows. Let $\{i_1, \dots, i_n\} = \{i | 1 \leq i \leq m, I_i \cap B \neq \emptyset\}$, where $i_1 < \dots < i_n$ and $n = |\{J_j\}|$. If we take $J_j = I_{i_j} \cap B$, then, $f_{\{J_j\}}$ should be such a function, $f_{\{J_j\}} : B \rightarrow \underline{n}$,

$$(f_{\{J_j\}})^{-1}(j) = (f_{\{I_i\}})^{-1}(i_j) \cap B = J_j.$$

Thus, $f_{\{I_i\}}|_B = \iota \circ f_{\{I_i\}}$, where $\iota : \underline{n} \rightarrow \{i_1, \dots, i_n\}$, $\iota(j) = i_j$ ($j = 1, \dots, n$). If we do not distinguish $f_{\{J_j\}}$ and $\iota \circ f_{\{I_i\}}$, then we have $f_{\{I_i\}}|_B = f_{\{I_i\}}$. In fact the functions to define a partition is an equivalent class under a natural equivalent relation, above formula is exactly valid in the sense of the equivalent relation.

If $D \subset B \subset \mathcal{R}(\{I_i\})$, $\{J_j\} = \{I_i\}_{(B)}$, we have $\{J_j\}_{(D)} = \{I_i\}_{(D)}$.

Map-union of the partitions Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B \subset A$ be a subset, $B \cap \mathcal{R}(\{I_i\}) = \emptyset$, $f : B \rightarrow \mathcal{R}(\{I_i\})$ be a map, we define a new partition $\{f^{-1}(I_i) \cup I_i\}$ which is called the map-union of $\{I_i\}$ and B by f , denoted by $B \sqcup_f \{I_i\}$.

Now we discuss some properties of the map-union.

- Let $\{I_i\}_{i=1}^m \in \mathcal{P}_{dis}^2(A)$ with defining function $f_{\{I_i\}}$, $B \subset A$ be a subset, $B \cap I = \emptyset$ ($I = \mathcal{R}(\{I_i\})$), for two maps $f : B \rightarrow I$, $g : B \rightarrow I$, then

$$B \sqcup_f \{I_i\} = B \sqcup_g \{I_i\} \Leftrightarrow f^{-1}(I_i) = g^{-1}(I_i) \ (1 \leq i \leq m) \Leftrightarrow f_{\{I_i\}} \circ f = f_{\{I_i\}} \circ g.$$

Thus, the map-union $B \sqcup_f \{I_i\}$ depends only on $f_{\{I_i\}} \circ f$. With the help of defining function $f_{\{I_i\}}$, there is a one-one correspondence between I_i and its index i , the map $f_{\{I_i\}} \circ f$ can be regarded as a map $\iota : B \rightarrow \{I_i\}$, and the map-union $B \sqcup_f \{I_i\}$ can be expressed as $\{\iota^{-1}(I_i) \cup I_i\}$ denoted by $B \sqcup_\iota \{I_i\}$.

- For two partitions $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, and two subsets $B, C \subset A$, which satisfy $I \cap J = \emptyset$, $B \cap C = \emptyset$, $(B \cup C) \cap (I \cup J) = \emptyset$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$, it is obvious that

$$(B \cup C) \sqcup_{f \cup g} (\{I_i\} \cup \{J_j\}) = (B \sqcup_f \{I_i\}) \cup (C \sqcup_g \{J_j\}),$$

where $f : B \rightarrow I$, $g : C \rightarrow J$, $f \cup g : B \cup C \rightarrow I \cup J$, $(f \cup g)|_B = f$, $(f \cup g)|_C = g$.

- Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B, C \subset A$, which satisfy $B \cap I = \emptyset$, $C \subset I$, where $I = \mathcal{R}(\{I_i\})$, then

$$B \sqcup_f \{I_i\} = (B_1 \sqcup_{f|_{B_1}} \{I_i\}_{(I \cap C)}) \cup (B_2 \sqcup_{f|_{B_2}} \{I_i\}_{(I \setminus C)}),$$

where $f : B \rightarrow I$, $B_1 = f^{-1}(I \cap C)$, $B_2 = f^{-1}(I \setminus C)$.

- Let $B, C \subset A$, $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B \cap C = \emptyset$, $I \cap (B \cup C) = \emptyset$, $I = \mathcal{R}(\{I_i\})$, then we have

$$(B \cup C) \sqcup_{f \cup g} \{I_i\} = B \sqcup_f (C \sqcup_g \{I_i\}) = C \sqcup_g (B \sqcup_f \{I_i\}),$$

where $f : B \rightarrow I$, $g : C \rightarrow I$, $f \cup g : B \cup C \rightarrow I$, $(f \cup g)|_B = f$, $(f \cup g)|_C = g$.

2.2 Quotient of the partitions

Now we define the quotient between two partitions.

Definition 2.1. Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B \subset \mathcal{R}(\{I_i\})$, we define the quotient of $\{I_i\}$ by $\{I_i\}_{(B)}$, denoted by $\{I_i\} / \{I_i\}_{(B)}$, to be

$$\{I_i\} / \{I_i\}_{(B)} = \{I_i\}_{I_i \cap B = \emptyset} \cup \{\mathcal{R}(\{I_i\}_{I_i \cap B \neq \emptyset}) \setminus B\}. \quad (2.1)$$

We call $\{\mathcal{R}(\{I_i\}_{I_i \cap B \neq \emptyset}) \setminus B\}$ the ideal part of the quotient $\{I_i\} / \{I_i\}_{(B)}$.

Remark 2.1. • It is convenience to denote $\mathcal{R}(\{I_i\}_{I_i \cap B \neq \emptyset})$ by $\mathcal{R}_{\{I_i\}, B}$, or, \mathcal{R}_B for short sometime. From definition 2.1 we can easily see that

$$\mathcal{R}(\{I_i\}) = \mathcal{R}(\{I_i\} / \{I_i\}_{(B)}) \cup B, \mathcal{R}(\{I_i\} / \{I_i\}_{(B)}) \cap B = \emptyset.$$

Therefore

$$\mathcal{R}(\{I_i\} / \{I_i\}_{(B)}) = \mathcal{R}(\{I_i\}) \setminus B.$$

- Particularly, when $\mathcal{R}_B = B$, which means that if $I_i \cap B \neq \emptyset$ we have $I_i \subset B$, or, $\{I_i\}_{(B)} = \{I_i\}_{I_i \cap B \neq \emptyset}$ which is a subset of $\{I_i\}$, then $\{I_i\} / \{I_i\}_{(B)} = (\{I_i\} \setminus \{I_i\}_{(B)}) \cup \{\emptyset\}$. We will identify $(\{I_i\} \setminus \{I_i\}_{(B)}) \cup \{\emptyset\}$ with $\{I_i\} \setminus \{I_i\}_{(B)}$. In this special situation the ideal part of quotient $\{I_i\} / \{I_i\}_{(B)}$ is $\{\emptyset\}$, we call this special situation the travail quotient. Furthermore, $\{I_i\} / \{I_i\}_{(I)} = \{I_i\} \setminus \{I_i\} = \emptyset$.
- Noting that $\{I_i\} = \{I_i\}_{I_i \cap B = \emptyset} \cup \{I_i\}_{I_i \cap B \neq \emptyset}$, and $\{I_i\}_{I_i \cap B \neq \emptyset} / \{I_i\}_{(B)} = \{\emptyset\} \cup \{\mathcal{R}_B \setminus B\}$, thus we can rewrite the formula (2.1) in the following form:

$$\{I_i\} / \{I_i\}_{(B)} = \{I_i\}_{I_i \cap B = \emptyset} \cup (\{I_i\}_{I_i \cap B \neq \emptyset} / \{I_i\}_{(B)}).$$

- Let $J \subset A$, $J \cap I = \emptyset$, $B \subset I$, $I = \mathcal{R}(\{I_i\})$, then the quotient $\{I_i\} / \{I_i\}_{(B)}$ induces the quotient of map-union $J \sqcup_f \{I_i\}$, where $f : J \rightarrow \{I_i\}$ is a map. Actually, let p_B be the projection from $\mathcal{R}(\{I_i\})$ to $\{I_i\} / \{I_i\}_{(B)}$, $p_B(I_i) = I_i$ for $I_i \cap B = \emptyset$, $p_B(I_i) = \mathcal{R}_B \setminus B$ for $I_i \cap B \neq \emptyset$, it is easy to check that

$$(J \sqcup_f \{I_i\}) / \{I_i\}_{(B)} = J \sqcup_{p_B \circ f} (\{I_i\} / \{I_i\}_{(B)}).$$

We will take a look at the properties of the quotient. Here we focus on the case of

$$(\{I_i\} / \{I_i\}_{(B)}) / (\{I_i\} / \{I_i\}_{(B)})_{(C)},$$

where $B, C \subset I = \mathcal{R}(\{I_i\})$, $B \cap C = \emptyset$. The key point is that when $\mathcal{R}_B \cap \mathcal{R}_C = \emptyset$, we have the following decomposition

$$\{I_i\}_{I_i \cap (B \cup C) \neq \emptyset} = \{I_i\}_{I_i \cap B \neq \emptyset} \cup \{I_i\}_{I_i \cap C \neq \emptyset}.$$

Otherwise, above decomposition is not valid.

Lemma 2.1. Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B, C \subset \mathcal{R}(\{I_i\})$, $B \cap C = \emptyset$, then following formulas are valid.

•

$$(\{I_i\} / \{I_i\}_{(B)})_{(C)} = \{I_i\}_{(B \cup C)} / \{I_i\}_{(B)}. \quad (2.2)$$

•

$$\begin{aligned} & (\{I_i\} / \{I_i\}_{(B)}) / (\{I_i\} / \{I_i\}_{(B)})_{(C)} \\ &= (\{I_i\} / \{I_i\}_{(B)}) / (\{I_i\}_{(B \cup C)} / \{I_i\}_{(B)}). \end{aligned} \quad (2.3)$$

- If $\mathcal{R}_B \cap \mathcal{R}_C = \emptyset$, then

$$\{I_i\}_{(B \cup C)} / \{I_i\}_{(B)} = \{I_i\}_{(C)}, \quad (2.4)$$

and

$$(\{I_i\} / \{I_i\}_{(B)})_{(C)} = \{I_i\}_{(C)}.$$

Proof. At first, in order to prove the formula (2.2), we calculate the left side of the formula. From definition 2.1 we have

$$\begin{aligned} & (\{I_i\} / \{I_i\}_{(B)})_{(C)} \\ &= (\{I_i\}_{I_i \cap B = \emptyset} \cup \{\mathcal{R}_B \setminus B\})_{(C)} \\ &= \{I_i \cap C\}_{I_i \cap B = \emptyset} \cup \{\mathcal{R}_B \cap C\}. \end{aligned}$$

On the other hand, for the right side of the formula (2.2) we have

$$\begin{aligned} & \{I_i\}_{(B \cup C)} / \{I_i\}_{(B)} \\ &= \{I_i \cap (B \cup C)\}_{I_i \cap B = \emptyset} \cup \{[\mathcal{R}_B \cap (B \cup C)] \setminus B\} \\ &= \{I_i \cap C\}_{I_i \cap B = \emptyset} \cup \{\mathcal{R}_B \cap C\}. \end{aligned}$$

Therefore, the formula (2.2) is valid. The formula (2.3) is the corollary of the formula (2.2).

We turn to prove the formula (2.4). Because $\mathcal{R}_B \cap \mathcal{R}_C = \emptyset$, we know that

$$I_i \cap B \neq \emptyset \iff I_i \cap C = \emptyset \text{ (or } I_i \cap C \neq \emptyset \iff I_i \cap B = \emptyset), \text{ if } I_i \cap (B \cup C) \neq \emptyset.$$

Thus

$$\{I_i\}_{(B \cup C)} = \{I_i\}_{(B)} \cup \{I_i\}_{(C)},$$

then, in this situation the quotient is trivial, that is

$$\{I_i\}_{(B \cup C)} / (\{I_i\}_{(B \cup C)})_{(B)} = \{I_i\}_{(B \cup C)} / \{I_i\}_{(B)} = \{I_i\}_{(B \cup C)} \setminus \{I_i\}_{(B)} = \{I_i\}_{(C)}.$$

Combining the formulas (2.2), (2.4) we can get the last formula in lemma 2.1. □

Proposition 2.1. Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B, C \subset \mathcal{R}(\{I_i\})$, $B \cap C = \emptyset$, then following formulas are valid.

- When $\mathcal{R}_B \cap \mathcal{R}_C \neq \emptyset$,

$$(\{I_i\} / \{I_i\}_{(B)}) / (\{I_i\}_{(B \cup C)} / \{I_i\}_{(B)}) = \{I_i\} / \{I_i\}_{(B \cup C)}. \quad (2.5)$$

- When $\mathcal{R}_B \cap \mathcal{R}_C = \emptyset$, we have

$$(\{I_i\} / \{I_i\}_{(B)}) / (\{I_i\}_{(B \cup C)} / \{I_i\}_{(B)}) = (\{I_i\} / \{I_i\}_{(B)}) / \{I_i\}_{(C)}. \quad (2.6)$$

Proof. From definition of quotient, we know that

$$\{I_i\} / \{I_i\}_{(B)} = \{I_i\}_{I_i \cap B = \emptyset} \cup \{\mathcal{R}_B \setminus B\},$$

and

$$\{I_i\}_{(B \cup C)} = \{I_i\}_{I_i \cap (B \cup C) = \emptyset} \cup \{\mathcal{R}_{B \cup C} \setminus (B \cup C)\}.$$

The discussions will be divided into two situations.

Case of $\mathcal{R}_B \cap \mathcal{R}_C \neq \emptyset$:

By the definition of the quotient we have

$$\begin{aligned}
& (\{I_i\}/\{I_i\}_{(B)})/(\{I_i\}/\{I_i\}_{(B)})_{(C)} \\
& = \{I_i\}_{I_i \cap (B \cup C) = \emptyset} \cup \{(\mathcal{R}(\{I_i\}_{I_i \cap B = \emptyset, I_i \cap C \neq \emptyset}) \cup (\mathcal{R}_B \setminus B)) \setminus C\}.
\end{aligned}$$

Noting $\mathcal{R}(\{I_i\}_{I_i \cap B = \emptyset, I_i \cap C \neq \emptyset} \cap B = \emptyset$, thus

$$\begin{aligned}
& \mathcal{R}(\{I_i\}_{I_i \cap B = \emptyset, I_i \cap C \neq \emptyset}) \cup (\mathcal{R}_B \setminus B) \\
& = (\mathcal{R}(\{I_i\}_{I_i \cap B = \emptyset, I_i \cap C \neq \emptyset}) \cup \mathcal{R}_B) \setminus B = \mathcal{R}_{B \cup C} \setminus B.
\end{aligned}$$

In summary, we get

$$(\{I_i\}/\{I_i\}_{(B)})/(\{I_i\}/\{I_i\}_{(B)})_{(C)} = \{I_i\}_{I_i \cap (B \cup C) = \emptyset} \cup (\mathcal{R}_{B \cup C} \setminus (B \cup C)).$$

Above formula is (2.5) exactly.

Case of $\mathcal{R}_B \cap \mathcal{R}_C = \emptyset$:

In this situation, we have $(\mathcal{R}_B \setminus B) \cap C = \emptyset$, and $\mathcal{R}(\{I_i\}_{I_i \cap B = \emptyset, I_i \cap C \neq \emptyset}) = \mathcal{R}(\{I_i\}_{I_i \cap C \neq \emptyset})$, thus

$$\begin{aligned}
& (\{I_i\}/\{I_i\}_{(B)})/(\{I_i\}/\{I_i\}_{(B)})_{(C)} \\
& = \{I_i\}_{I_i \cap (B \cup C) = \emptyset} \cup \{\mathcal{R}_B \setminus B\} \cup \{(\mathcal{R}(\{I_i\}_{I_i \cap B = \emptyset, I_i \cap C \neq \emptyset}) \setminus C)\} \\
& = \{I_i\}_{I_i \cap (B \cup C) = \emptyset} \cup \{\mathcal{R}_B \setminus B\} \cup \{\mathcal{R}_C \setminus C\}.
\end{aligned}$$

Now we reach the formula (2.6). □

Corollary 2.1.

$$(\{I_i\}/\{I_i\}_{(B)})/(\{I_i\}/\{I_i\}_{(B)})_{(C)} = (\{I_i\}/\{I_i\}_{(C)})/(\{I_i\}/\{I_i\}_{(C)})_{(B)}. \quad (2.7)$$

$$(\{I_i\}/\{I_i\}_{(B)})/(\{I_i\}/\{I_i\}_{(C)}) = (\{I_i\}/\{I_i\}_{(C)})/(\{I_i\}/\{I_i\}_{(B)}), \quad (2.8)$$

Now we consider more general situation.

Definition 2.2. Let $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{I_i\})$, we define the following quotient inductively,

$$\begin{aligned}
& \{I_i\}/(J_1) \doteq \{I_i\}/\{I_i\}_{(J_1)}, \{I_i\}/(J_1, J_2) \doteq (\{I_i\}/(J_1))/(\{I_i\}/(J_1))_{(J_2)}, \\
& \dots, \{I_i\}/(J_1, \dots, J_{k+1}) \doteq (\{I_i\}/(J_1, \dots, J_k))/(\{I_i\}/(J_1, \dots, J_k))_{(J_{k+1})}, \dots
\end{aligned}$$

Remark 2.2. By the formula (2.7) in corollary 2.1, it is easy to check that the quotient in definition 2.2 does not depend on the order of J_j . Precisely, let $n = |\{J_j\}|$, $\tau \in \mathbb{S}_n$ be a permutation on \underline{n} , then,

$$\{I_i\}/(J_j) = \{I_i\}/(J_{\tau(j)}).$$

Definition 2.3. Let $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, if $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{I_i\})$, and

$$\mathcal{R}_{\{I_i\}, J_j} \cap \mathcal{R}_{\{I_i\}, J_{j'}} = \emptyset, j \neq j', \quad (2.9)$$

we call $\{J_j\}$ admits to $\{I_i\}$, denoted by $\{J_j\} \sqsubset \{I_i\}$.

Remark 2.3. The condition (2.9) is equivalent to the following conditions

$$\mathcal{R}_{\{I_i\}, J_j} \cap J_{j'} = \emptyset \text{ (or, } \mathcal{R}_{\{I_i\}, J_{j'}} \cap J_j = \emptyset), j \neq j'.$$

$\mathcal{R}_{\{I_i\}, J_j}$ will be shortly denoted by \mathcal{R}_{J_j} later.

Actually, the fact $\{J_j\}_{j=1}^p \sqsubset \{I_i\}$ means that $\{I_i\}$ adapts to a decomposition as follows,

$$\{I_i\} = \{I_i\}_{I_i \cap J = \emptyset} \cup \left(\bigcup_j \{I_i\}_{I_i \cap J_j \neq \emptyset} \right),$$

where $J = \mathcal{R}(\{J_j\})$. Therefore, we have

$$\begin{aligned} & \{I_i\} / (J_j) \\ &= \{I_i\}_{I_i \cap J = \emptyset} \cup \left(\bigcup_j \{I_i\}_{I_i \cap J_j \neq \emptyset} / \{I_i\}_{(J_j)} \right) \\ &= \{I_i\}_{I_i \cap J = \emptyset} \cup \left(\bigcup_j \{ \mathcal{R}_{J_j} \setminus J_j \} \right). \end{aligned}$$

Generally, $\{J_j\}$ may do not admit to $\{I_i\}$, even though $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{I_i\})$. But if we consider the quotient, the situation can always be induced to the simple case.

Proposition 2.2. *Let $\{J_j\}, \{I_i\} \in \mathcal{P}_{dis}^2(A)$, $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{I_i\})$. Then, there is a partition $\{L_l\} \in \mathcal{P}_{dis}^2(A)$ satisfying*

- $\{L_l\} \sqsubset \{I_i\}$.
- $\{J_j\} \subset \{L_l\}$, and $\mathcal{R}(\{J_j\}) = \mathcal{R}(\{L_l\})$.
- If $\{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfies $\{J_j\} \subset \{K_k\} \sqsubset \{I_i\}$, then we have $\{L_l\} \subset \{K_k\}$.

If we ignore the order in $\{L_l\}$, $\{L_l\}$ is unique.

Proof. In $\{J_j\}$ we define an equivalent relation as follows. Let $n = |\{J_j\}|$. For any $j, j' \in \underline{n}$, we say $\mathcal{R}_j \sim \mathcal{R}_{j'}$, if there is a subset $\{j_0, j_1, \dots, j_m\}$ of \underline{n} , such that $j_0 = j$, $j_m = j'$, and $\mathcal{R}_{J_{j_k}} \cap \mathcal{R}_{J_{j_{k+1}}} \neq \emptyset$ ($k = 0, \dots, m-1$). It is obvious that \sim is an equivalent relation.

Under the equivalent relation defined above $\{J_j\}$ can be divided into the set of equivalent class, i.e. we have

$$\{J_j\} = \bigcup \{J_j\}_{j \in E_l},$$

where $\{E_l\} \in \mathbf{Part}(\underline{n})$, each $\{J_j\}_{j \in E_l}$ is an equivalent class under \sim . We take L_l to be $L_l = \bigcup_{j \in E_l} J_j$. It is easy to check that $\mathcal{R}(\{L_l\}) = \mathcal{R}(\{J_j\})$ and $\mathcal{R}_{L_l} \cap \mathcal{R}_{L_{l'}} = \emptyset$ for $l \neq l'$. From proposition 2.1 we know that

$$\{I_i\} / (J_j)_{j \in E_l} = \{I_i\} / \{I_i\}_{(\bigcup_{j \in E_l} J_j)} = \{I_i\} / \{I_i\}_{(L_l)}.$$

By the previous discussion, we can reach the formula

$$\{I_i\} / (J_j) = \{I_i\} / (L_l).$$

Let $\{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfy $\{J_j\} \subset \{K_k\} \sqsubset \{I_i\}$, then we can prove that for each l , there is k such that $\mathcal{R}_{\{I_i\}, L_l} \subset \mathcal{R}_{\{I_i\}, K_k}$. On the other hand, $\{J_j\} \subset \{K_k\}$, thus we have $\{L_l\} \subset \{K_k\}$.

Let $\{L'_{l'}\} \in \mathcal{P}_{dis}^2(A)$ satisfy the conditions same as ones of $\{L_l\}$, then both of $\{L_l\} \subset \{L'_{l'}$ and $\{L'_{l'}\} \subset \{L_l\}$ are valid, which implies $\{L_l\} = \{L'_{l'}\}$. □

We denote $\{L_l\}$ by $\{L_l\} = \{J_j\}_{ad, \{I_i\}}$. From proposition 2.2, when we discuss the quotient $\{I_i\} / (J_j)$, we can always assume $\{J_j\} \sqsubset \{I_i\}$.

Corollary 2.2. Let $\{I_i\}, \{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfying $J \cap K = \emptyset$, $J, K \subset I$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$, $K = \mathcal{R}(\{K_k\})$. Then we have

$$\begin{aligned} & (\{J_j\} \cup \{K_k\})_{ad, \{I_i\}} \\ &= (\{J_j\}_{ad, \{I_i\}} \cup \{K_k\}_{ad, \{I_i\}})_{ad, \{I_i\}} \\ &= (\{J_j\} \cup \{K_k\}_{ad, \{I_i\} / \{J_j\}})_{ad, \{I_i\}} \\ &= (\{K_k\} \cup \{J_j\}_{ad, \{I_i\} / \{K_k\}})_{ad, \{I_i\}}. \end{aligned}$$

2.3 Insertion of the partitions

Now we turn to the discussion of insertion. Let $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) = \emptyset$. We hope to define the insertion of $\{J_j\}$ into $\{I_i\}$ at I_a .

Definition 2.4. Let $\{I_i\}_{1 \leq i \leq m}, \{J_j\}_{1 \leq j \leq n} \in \mathcal{P}_{dis}^2(A)$, $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) = \emptyset$, and $\iota : I_a \rightarrow \{J_j\}_{1 \leq j \leq n}$ be a map ($1 \leq a \leq m$). The insertion of $\{J_j\}_{j=1}^n$ into $\{I_i\}_{i=1}^m$ at I_a by ι is a partition $\{I_i\} \circ_a^\iota \{J_j\} \in \mathcal{P}_{dis}^2(A)$, where

$$= \{I_1, \dots, I_{a-1}, J_1 \cup \iota^{-1}(J_1), \dots, J_n \cup \iota^{-1}(J_n), I_{a+1}, \dots, I_m\}. \quad (2.10)$$

I_a is called the position of insertion, and ι is called insertion map.

Remark 2.4.

- We can explain the insertion in terms of map-union. If we ignore the order of the partitions, the insertion $\{I_i\} \circ_a^\iota \{J_j\}$ can be expressed as $\{I_i\}_{i \neq a} \cup (I_a \sqcup_\iota \{J_j\})$. In fact, the map-union is a special situation of insertion, $B \sqcup_\iota \{J_j\} = \{B\} \circ_B^\iota \{J_j\}$ ($B \cap \mathcal{R}(\{J_j\}) = \emptyset$). For simplicity, we can denote $\{I_i\} \circ_a^\iota \{J_j\}$ in the following intuitive way:

$$\begin{array}{c} \{\dots, \hat{I}_a, \dots\} \\ \uparrow \\ I_a \sqcup_\iota \{J_j\}. \end{array}$$

- Particularly, we can always identify $\{I_i\}$ with $\{I_i\} \cup \{\emptyset\}$, then the insertion of $\{J_j\}$ into $\{I_i\}$ at \emptyset is defined as $\{I_i\} \cup \{J_j\}$ denoted by $\{I_i\} \circ_\emptyset \{J_j\}$. We call $\{I_i\} \circ_\emptyset \{J_j\}$ the travail insertion. Let $\{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfying $(I \cup J) \cap \mathcal{R}(\{K_k\}) = \emptyset$, then, it is obvious that $(\{I_i\} \cup \{J_j\}) \circ_{J_b}^\iota \{K_k\} = \{I_i\} \cup (\{J_j\} \circ_{J_b}^\iota \{K_k\})$, or $(\{I_i\} \circ_\emptyset \{J_j\}) \circ_{J_b}^\iota \{K_k\} = \{I_i\} \circ_\emptyset (\{J_j\} \circ_{J_b}^\iota \{K_k\})$.

In the case of non-travail insertion we have:

Proposition 2.3. Let $\{I_i\}, \{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfying $I \cap J = \emptyset$, $I \cup J = K$ ($I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$, $K = \mathcal{R}(\{K_k\})$), $\{J_j\} = \{K_k\}_{(J)}$, $\mathcal{R}_{\{K_k\}, J} \setminus J \neq \emptyset$. Then

$$\{K_k\} / \{J_j\} = \{I_i\},$$

if and only if, there is an insertion map $\iota : I_a \rightarrow \{J_j\}$ for some I_a ($1 \leq a \leq \#\{I_i\}$) such that

$$\{K_k\} = \{I_i\} \circ_{I_a}^\iota \{J_j\}, \text{ (or } \{K_k\} = (\{K_k\} / \{J_j\}) \circ_{I_a}^\iota \{J_j\}).$$

Proof. Let $\{K_k\} = \{I_i\} \circ_{I_a}^{\iota} \{J_j\}$. By proposition 2.2 and remark 2.2 we have

$$\{K_k\} = \{I_i\}_{i \neq a} \cup (I_a \sqcup_{\iota} \{J_j\}).$$

Thus $\{K_k\}_{K_k \cap J = \emptyset} = \{I_i\}_{i \neq a}$, $\{K_k\}_{K_k \cap J \neq \emptyset} = I_a \sqcup_{\iota} \{J_j\}$ and $\{K_k\}_{(J)} = \{J_j\}$. Then we have

$$\{K_k\}_{K_k \cap J \neq \emptyset} / \{J_j\} = (I_a \sqcup_{\iota} \{J_j\}) / \{J_j\} = \{I_a\}.$$

Finally, we get

$$\{K_k\} / \{J_j\} = \{K_k\}_{K_k \cap J = \emptyset} \cup (\{K_k\}_{K_k \cap J \neq \emptyset} / \{J_j\}) = \{I_i\}_{i \neq a} \cup \{I_a\} = \{I_i\},$$

i.e.

$$(\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) / \{J_j\} = \{I_i\}.$$

Conversely, we assume $\{K_k\}$ satisfies $\{K_k\} / \{J_j\} = \{I_i\}$. Let $K = \mathcal{R}(\{K_k\})$, $\mathcal{R}_J = \mathcal{R}(\{K_k\}_{K_k \cap J \neq \emptyset})$, then we have $K = I \cup J$, $\{J_j\} = \{K_k\}_{(J)} = \{K_k \cap J\}_{K_k \cap J \neq \emptyset}$, and

$$\{K_k\}_{K \cap J = \emptyset} \cup \{\mathcal{R}_J \setminus J\} = \{I_i\}.$$

In order to recover $\{K_k\}$ by insertion, the position of the insertion should be taken to be $I_a = \mathcal{R}_J \setminus J$ for some a ($1 \leq a \leq \#\{I_i\}$). Thus $\{K_k\}_{K \cap J = \emptyset} = \{I_i\}_{i \neq a}$. Let $\{K_k\}_{K_k \cap J \neq \emptyset} = \{K_{k_j}\}$, and $K_{k_j} = J_j \cup L_j$, where $J_j \cap L_j = \emptyset$, we get a decomposition of I_a that is $I_a = \bigcup_j L_j$. Above decomposition define a map $\iota : I_a \rightarrow \{J_j\}$, such that $\iota^{-1}(J_j) = L_j$. It is obvious that $\{K_{k_j}\} = I_a \sqcup_{\iota} \{J_j\}$. Up to now, we have proved

$$\{K_k\} = \{I_i\} \circ_{I_a}^{\iota} \{J_j\}.$$

□

Remark 2.5. In the first situation of proposition 2.3, the position of insertion is the ideal part of quotient, and the insertion map is taken in a canonical way, thus we denote this insertion by symbol \circ_{ideal_B} . Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $B \subset \mathcal{R}(\{I_i\})$.

- When $\mathcal{R}(\{I_i\}_{I_i \cap B \neq \emptyset}) \setminus B \neq \emptyset$, we have

$$(\{I_i\} / \{I_i\}_{(B)}) \circ_{\text{ideal}_B} \{I_i\}_{(B)} = \{I_i\},$$

where the position of insertion is at $\mathcal{R}(\{I_i\}_{I_i \cap B \neq \emptyset}) \setminus B$, and the insertion map is taken to be $\iota^{-1}(I_i \cap B) = I_i \setminus B$ if $I_i \setminus B \neq \emptyset$.

- When $\mathcal{R}(\{I_i\}_{I_i \cap B \neq \emptyset}) = B$, we have

$$\{I_i\} = (\{I_i\} / \{I_i\}_{(B)}) \circ_{\emptyset} \{I_i\}_{(B)},$$

where the insertion is travail one.

There is a conclusion about insertion and map-union as follows.

Proposition 2.4. Let $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, $B \subset A$, $I \cap J = \emptyset$, $B \cap (I \cup J) = \emptyset$, $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$. For a pair (f, ι) there is a unique pair (f', ι') such that

$$(B \sqcup_f \{I_i\}) \circ_{f^{-1}(I_a) \cup I_a}^{\iota} \{J_j\} = B \sqcup_{f'} (\{I_i\} \circ_{I_a}^{\iota'} \{J_j\}),$$

and vice-versa, where $1 \leq a \leq |\{I_i\}|$, $f : B \rightarrow \{I_i\}$, $f' : B \rightarrow \{I_i\} \circ_a^{\iota'} \{J_j\}$, $\iota : f^{-1}(I_a) \sqcup I_a \rightarrow \{J_j\}$, $\iota' : I_a \rightarrow \{J_j\}$. On the other hand, we also have

$$(B \sqcup_{f'} (\{I_i\} \circ_a^{\iota'} \{J_j\})) / \{J_j\} = B \sqcup_{f' \circ p} ((\{I_i\} \circ_a^{\iota'} \{J_j\}) / \{J_j\}) = B \sqcup_{f' \circ p} \{I_i\},$$

where $p : \{I_i\} \circ_a^{\iota'} \{J_j\} \rightarrow \{I_i\}$ is a projection satisfying

$$p(I_i) = I_i \ (i \neq a) \quad p((\iota')^{-1}(J_j) \cup J_j) = I_a \ (\forall j), \quad f' \circ p = f.$$

Proof. Let (f, ι) be a given pair, we want to construct the pair (f', ι') based on (f, ι) . By the definition of insertion and map-union we have

$$(B \sqcup_f \{I_i\}) \circ_{f^{-1}(I_a) \cup I_a}^{\iota} \{J_j\} = \{f^{-1}(I_i) \cup I_i\}_{i \neq a} \cup ((f^{-1}(I_a) \cup I_a) \sqcup_{\iota} \{J_j\}),$$

and

$$B \sqcup_{f'} (\{I_i\} \circ_a^{\iota'} \{J_j\}) = \{f'^{-1}(I_i) \cup I_i\}_{i \neq a} \cup (B_1 \sqcup_{f'|_{B_1}} (I_a \sqcup_{\iota'} \{J_j\})),$$

where $B_1 = f'^{-1}(I_a \cup J)$. The previous formulas imply $f^{-1}(I_i) = f'^{-1}(I_i)$ ($i \neq a$), and $f^{-1}(I_a) = f'^{-1}(I_a \cup J)$. Thus we have $B_1 = f^{-1}(I_a)$ and $f|_{B \setminus B_1} = f'|_{B \setminus B_1}$. If we regard $\iota, \iota', f'|_{B_1}$ as maps from some subsets to $\{J_j\}$, then the first formula in proposition 2.4 means that the formula $\iota = f'|_{B_1} \cup \iota'$ should be valid, i.e. we have

$$f'|_{B_1} = \iota|_{B_1}, \quad \iota' = \iota|_{I_a}.$$

Conversely, starting from (f', ι') we can determine (f, ι) in similar way.

The second formula in proposition 2.4 is obviously valid. □

Remark 2.6. Observing the proof of proposition 2.4, in the procedure of from (f', ι') to (f, ι) , f' determines f uniquely, where f' and f depend on “ a ” only. If we fix f' , there is an one-one corresponding between ι and ι' .

There is another version of proposition 2.1 in terms of the insertion.

Proposition 2.5. Let $\{I_i\}, \{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$, $I \cap J = \emptyset$, $K \cap (I \cup J) = \emptyset$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$, $K = \mathcal{R}(\{K_k\})$. Let $\{L_l\} = (\{K_k\} \circ_{K_a}^{\iota} \{I_i\}) \circ_{*}^{\tau} \{J_j\}$, then

•

$$\mathcal{R}_{\{L_l\}, I} \cap \mathcal{R}_{\{L_l\}, J} = \emptyset \Leftrightarrow \{L_l\} = (\{K_k\} \circ_{K_a}^{\iota} \{I_i\}) \circ_{K_b}^{\tau} \{J_j\}, \quad a \neq b.$$

•

$$\mathcal{R}_{\{L_l\}, I} \cap \mathcal{R}_{\{L_l\}, J} \neq \emptyset \Leftrightarrow \{L_l\} = (\{K_k\} \circ_{K_a}^{\iota} \{I_i\}) \circ_{\iota^{-1}(I_c) \cup I_c}^{\tau} \{J_j\}.$$

In this situation, we have

$$\{L_l\} = \{K_k\} \circ_{K_a}^{\iota'} (\{I_i\} \circ_{I_c}^{\kappa} \{J_j\}),$$

and

$$\{L_l\}_{(I \cup J)} = \{I_i\} \circ_{I_c}^{\kappa} \{J_j\},$$

for some ι' and κ .

Proof. **Case of $\mathcal{R}_{\{\mathbf{L}_1\}, \mathbf{I}} \cap \mathcal{R}_{\{\mathbf{L}_1\}, \mathbf{J}} = \emptyset$:**

Let $\mathcal{R}_{\{L_l\}, I} \cap \mathcal{R}_{\{L_l\}, J} = \emptyset$, then

$$\{L_l\} = \{L_l\}_{L_l \cap (I \cup J) = \emptyset} \cup \{L_l\}_{L_l \cap I \neq \emptyset} \cup \{L_l\}_{L_l \cap J \neq \emptyset}.$$

Noting $\{L_l\}_{(J)} = \{J_j\}$, by proposition 2.1 and proposition 2.3 we have

$$\{L_l\} / \{J_j\} = \{L_l\}_{L_l \cap (I \cup J) = \emptyset} \cup \{L_l\}_{L_l \cap I \neq \emptyset} \cup \{\mathcal{R}_{\{L_l\}, J} \setminus J\} = \{K_k\} \circ_{K_a}^l \{I_i\}.$$

Furthermore, we now that $\{L_l\}_{(I)} = \{I_i\}$. Thus

$$(\{L_l\} / \{J_j\}) / \{I_i\} = \{L_l\}_{L_l \cap (I \cup J) = \emptyset} \cup \{\mathcal{R}_{\{L_l\}, I} \setminus I\} \cup \{\mathcal{R}_{\{L_l\}, J} \setminus J\} = \{K_k\}.$$

Above formula means that

$$\{\mathcal{R}_{\{L_l\}, I} \setminus I\} = \{L_l\}_{L_l \cap I \neq \emptyset} / \{I_i\} = K_a \sqcup_l \{I_i\},$$

and

$$\{\mathcal{R}_{\{L_l\}, J} \setminus J\} = \{L_l\}_{L_l \cap J \neq \emptyset} / \{J_j\} = K_b \sqcup_\tau \{J_j\},$$

for some a and b ($a \neq b$).

Conversely, let

$$\{L_l\} = (\{K_k\} \circ_{K_a}^l \{I_i\}) \circ_{K_b}^\tau \{J_j\}, a \neq b.$$

Then, we have

$$\{L_l\} = \{K_k\}_{k \neq a, b} \cup (K_a \sqcup_l \{I_i\}) \cup (K_b \sqcup_\tau \{J_j\}),$$

and $\{L_l\}_{L_l \cap I \neq \emptyset} = K_a \sqcup_l \{I_i\}$, $\{L_l\}_{(I)} = \{I_i\}$, $\{L_l\}_{L_l \cap J \neq \emptyset} = K_b \sqcup_\tau \{J_j\}$, $\{L_l\}_{(J)} = \{J_j\}$. Above facts imply $\mathcal{R}_{\{L_l\}, I} \cap \mathcal{R}_{\{L_l\}, J} = \emptyset$.

Case of $\mathcal{R}_{\{\mathbf{L}_1\}, \mathbf{I}} \cap \mathcal{R}_{\{\mathbf{L}_1\}, \mathbf{J}} \neq \emptyset$:

The conclusion in this situation is the corollary of the one in the case of $\mathcal{R}_{\{L_l\}, I} \cap \mathcal{R}_{\{L_l\}, J} = \emptyset$. Let

$$\{L_l\} = \{K_k\}_{k \neq a} \cup ((K_a \sqcup_l \{I_i\}) \circ_{\ell^{-1}(I_c) \cup I_c}^\tau \{J_j\}).$$

By proposition 2.4 we have

$$(K_a \sqcup_l \{I_i\}) \circ_{\ell^{-1}(I_c) \cup I_c}^\tau \{J_j\} = K_a \sqcup_{\ell'} (\{I_i\} \circ_{I_c}^\kappa \{J_j\}),$$

for some ℓ' and κ . It is natural that we have

$$\{L_l\}_{(I \cup J)} = \{I_i\} \circ_{I_c}^\kappa \{J_j\}.$$

□

Proposition 2.5 can be generalized to more general situation. Let $\{I_i\}, \{J_{j_1}^{(1)}\}, \dots, \{J_{j_n}^{(n)}\} \in \mathcal{P}_{dis}^2(A)$, $J^{(l)} \cap J^{(l')} = \emptyset$ ($l \neq l'$), $I \cap (\bigcup_l J^{(l)}) = \emptyset$, $I = \mathcal{R}(\{I_i\})$, $J^{(l)} = \mathcal{R}(\{J_{j_l}^{(l)}\})$ ($1 \leq l \leq n$),

$$\begin{aligned} \{K_{k_1}^{(1)}\} &= \{I_i\} \circ_{I_a}^{\iota_1} \{J_{j_1}^{(1)}\}, \{K_{k_2}^{(2)}\} = \{K_{k_1}^{(1)}\} \circ_{K_{a_1}}^{\iota_2} \{J_{j_2}^{(2)}\}, \dots, \\ \{K_k\} &= \{K_{k_{n-1}}^{(n-1)}\} \circ_{K_{a_{n-1}}}^{\iota_n} \{J_{j_n}^{(n)}\}. \end{aligned}$$

Corollary 2.3. $\mathcal{R}_{\{K_k\}, J^{(l)}} \cap \mathcal{R}_{\{K_k\}, J^{(l')}} = \emptyset$ ($l \neq l'$) if and only if there are a_1, \dots, a_n such that

$$\{K_k\} = \{I_i\}_{i \neq a_1, \dots, a_n} \cup \left(\bigcup_{l=1}^n I_{a_l} \sqcup_{l_l} \{J_{j_l}^{(l)}\} \right).$$

Remark 2.7. The conclusion of corollary 2.2 can be described by a different way, which is

$$\{J^{(l)}\}_{l=1}^n \subset \{K_k\} \Leftrightarrow \{K_k\} = \{I_i\}_{i \neq a_1, \dots, a_n} \cup \left(\bigcup_{l=1}^n I_{a_l} \sqcup_{l_l} \{J_{j_l}^{(l)}\} \right),$$

for some a_1, \dots, a_n .

2.4 Coproduct

Now we consider the coproduct as an application of quotient. At first we introduce

$$\mathcal{P}_{dis, k}^2(A) \subset \underbrace{\mathcal{P}_{dis}^2(A) \times \dots \times \mathcal{P}_{dis}^2(A)}_{k\text{-times}},$$

$$\begin{aligned} (\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) \in \mathcal{P}_{dis, k}^2(A) &\iff \{I_{i_\lambda}^{(\lambda)}\} \in \mathcal{P}_{dis}^2(A), \lambda = 1, \dots, k, \\ \mathcal{R}(\{I_{i_\lambda}^{(\lambda)}\}) \cap \mathcal{R}(\{I_{i_\mu}^{(\mu)}\}) &= \emptyset, 1 \leq \lambda < \mu \leq k. \end{aligned}$$

Similar to definition 2.3 we have

Definition 2.5. Let $\{J_j\} \in \mathcal{P}_{dis}^2(A)$, $(\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) \in \mathcal{P}_{dis, k}^2(A)$, we say $\{J_j\} \sqsubset (\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\})$, if $\mathcal{R}(\{J_j\}) \subset \bigcup_{\lambda=1}^k \mathcal{R}(\{I_{i_\lambda}^{(\lambda)}\})$, and $\{J_j\}_{J_j \cap \mathcal{R}(\{I_{i_\lambda}^{(\lambda)}\}) \neq \emptyset} \subset \{I_{i_\lambda}^{(\lambda)}\}$, ($\lambda = 1, \dots, k$). In this situation, we define

$$(\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) / (J_j) = (\{I_{i_1}^{(1)}\} / (J_j), \dots, \{I_{i_k}^{(k)}\} / (J_j)),$$

where

$$\{I_{i_\lambda}^{(\lambda)}\} / (J_j) = \{I_{i_\lambda}^{(\lambda)}\} / (J_j)_{J_j \subset \mathcal{R}(\{I_{i_\lambda}^{(\lambda)}\})}, \lambda = 1, \dots, k.$$

Similarly, we can discuss the insertion for the case of $(\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\})$ in a obvious way.

Let \mathbb{K} be a field of characteristic zero, $V_{\mathbb{K}, A} = \bigoplus_{1 \leq k \leq |A|} \text{Span}_{\mathbb{K}}(\mathcal{P}_{dis, k}^2(A))$. Based on the above discussions we now define coproduct on $V_{\mathbb{K}, A}$, $\Delta : V_{\mathbb{K}, A} \rightarrow V_{\mathbb{K}, A} \otimes V_{\mathbb{K}, A}$.

Definition 2.6. Let $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, $(\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) \in \mathcal{P}_{dis, k}^2(A)$, we define

•

$$\begin{aligned} \Delta\{I_i\} &= \emptyset \otimes \{I_i\} + \{I_i\} \otimes \emptyset + \\ &\sum_{\{J_1, \dots, J_l\} \sqsubset \{I_i\}} (\{I_i\}_{(J_1)} \cdots \{I_i\}_{(J_l)}) \otimes \{I_i\} / (J_j). \end{aligned} \quad (2.11)$$

•

$$\begin{aligned} &\Delta(\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) \\ &= \emptyset \otimes (\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) + (\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) \otimes \emptyset + \\ &\sum_{\{J_j\} \sqsubset (\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\})} (\{I_i\}_{(J_1)} \cdots \{I_i\}_{(J_l)}) \otimes (\{I_{i_1}^{(1)}\}, \dots, \{I_{i_k}^{(k)}\}) / (J_j), \end{aligned} \quad (2.12)$$

where $\{I_i\} = \bigcup_{\lambda} \{I_{i_\lambda}^{(\lambda)}\}$.

For associativity of coproduct defined in definition 2.6 we need the following conclusion.

Theorem 2.1. *Let $\{I_i\}, \{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$, $\{J_j\} \sqsubset \{I_i\}$, $\{K_k\} \sqsubset \{I_i\}/(J_j)$. If we take*

$$\{M_\mu\} = (\{J_j\} \cup \{K_k\})_{ad, \{I_i\}},$$

then

$$\{M_\mu\} = \{J_j\}_{\mathcal{R}_{J_j} \cap K = \emptyset} \cup \{N_k\},$$

where $N_k = K_k \cup (\bigcup_{\mathcal{R}_{\{I_i\}, J_j} \cap K_k \neq \emptyset} J_j)$, $K = \mathcal{R}(\{K_k\})$, $k = 1, \dots, |\{K_k\}|$. $\{M_\mu\}$ satisfies the following conditions:

- $\{K_k\} = \{M_\mu\}/(J_j)$.
- $\{M_\mu\} \sqsubset \{I_i\}$.
- $\{J_j\} \sqsubset (\dots, \{I_i\}_{(M_\mu)}, \dots)$.
- $(\dots, (\{I_i\}/(J_j))_{(K_k)}, \dots) = (\dots, \{I_i\}_{(M_\mu)}, \dots)/(J_j)$.
- $(\{I_i\}/(J_j))/(K_k) = \{I_i\}/(M_\mu)$.

Proof. At first, we prove

$$\{M_\mu\} = \{J_j\}_{\mathcal{R}_{J_j} \cap K = \emptyset} \cup \{N_k\}.$$

Let $\{L_\lambda\} = \{I_i\}/(J_j)$, then, from definition of quotient we have

$$\{L_\lambda\} = \{I_i\}_{I_i \cap J = \emptyset} \cup \{\mathcal{R}_{J_j} \setminus J_j\}_{j=1}^p,$$

where $p = |\{J_j\}|$, $J = \mathcal{R}(\{J_j\}_{j=1}^p)$, and $\mathcal{R}_{J_j} = \mathcal{R}_{\{I_i\}, J_j}$. Recalling definition 2.3, $\{K_k\} \sqsubset \{L_\lambda\}$ means that

$$\mathcal{R}_{\{L_i\}, K_k} \cap \mathcal{R}_{\{L_i\}, K_{k'}} = \emptyset, k \neq k',$$

On the other hand, according to $\{K_k\}$ there is the decomposition of $\{L_\lambda\}$ as follows,

$$\{L_\lambda\} = \{L_\lambda\}_{L_\lambda \cap K = \emptyset} \cup \{L_\lambda\}_{L_\lambda \cap K_1 \neq \emptyset} \cup \dots \cup \{L_\lambda\}_{L_\lambda \cap K_q \neq \emptyset},$$

where $K = \mathcal{R}(\{K_k\})$, and $q = |\{K_k\}|$. It is obvious that

$$\{L_\lambda\}_{L_\lambda \cap K = \emptyset} = \{I_i\}_{I_i \cap (J \cup K) = \emptyset} \cup \{\mathcal{R}_{J_j} \setminus J_j\}_{\mathcal{R}_{J_j} \cap K = \emptyset},$$

and

$$\{L_\lambda\}_{L_\lambda \cap K_k \neq \emptyset} = \{I_i\}_{I_i \cap J = \emptyset, I_i \cap K_k \neq \emptyset} \cup \{\mathcal{R}_{J_j} \setminus J_j\}_{\mathcal{R}_{J_j} \cap K_k \neq \emptyset},$$

where $k = 1, \dots, q$. Notong $J \cap K = \emptyset$, thus

$$(\mathcal{R}_{J_j} \setminus J_j) \cap K_k \neq \emptyset \Leftrightarrow \mathcal{R}_{J_j} \cap K_k \neq \emptyset.$$

In summary, we know that for each \mathcal{R}_{J_j} there two possibilities:

- $\mathcal{R}_{J_j} \cap K = \emptyset$,

- or, there is an unique k such that $\mathcal{R}_{J_j} \cap K_k \neq \emptyset$.

Up to now we have proved

$$\{M_\mu\} = \{J_j\}_{\mathcal{R}_{J_j} \cap K = \emptyset} \cup \{N_k\}.$$

We now begin to prove that $\{M_\mu\}$ satisfies the conditions in theorem 2.1 step by step.

$\{\mathbf{K}_k\} = \{\mathbf{M}_\mu\} / (\mathbf{J}_j)$:

Let $\{G_l\} = \{J_j\}_{ad, \{M_\mu\}}$, then we have

$$\{G_l\} = \{J_j\}_{\mathcal{R}_{J_j} \cap K = \emptyset} \cup \left(\bigcup_k \{J_j\}_{\mathcal{R}_{J_j} \cap K_k \neq \emptyset} \right),$$

and $\{M_\mu\} / (J_j) = \{M_\mu\} / (G_l)$. It is easy to check that $\{K_k\} = \{M_\mu\} / (G_l)$.

$\{\mathbf{M}_\mu\} \sqsubset \{\mathbf{I}_i\}$:

Noting

$$\{I_i\}_{I_i \cap N_k \neq \emptyset} = \{I_i\}_{I_i \cap J = \emptyset, I_i \cap K_k \neq \emptyset} \cup \left(\bigcup_j \{I_i\}_{I_i \cap J_j \neq \emptyset, \mathcal{R}_{J_j} \cap K_k \neq \emptyset} \right),$$

thus

$$\mathcal{R}_{\{I_i\}, N_k} = \mathcal{R}_{\{L_\lambda\}, K_k} \cup \left(\bigcup_{\mathcal{R}_{J_j} \cap K_k \neq \emptyset} J_j \right), \quad k = 1, \dots, q.$$

Because $\{K_k\} \sqsubset \{L_\lambda\}$, which means

$$\mathcal{R}_{\{L_\lambda\}, K_k} \cap \mathcal{R}_{\{L_\lambda\}, K_{k'}} = \emptyset, \quad k \neq k',$$

hence, $\{N_k\} \sqsubset \{I_i\}$.

$\{\mathbf{J}_j\} \sqsubset (\dots, \{\mathbf{I}_i\}_{(\mathbf{M}_\mu)}, \dots)$:

It is enough for us to prove $\{J_j\}_{J_j \subset N_k} \sqsubset \{I_i\}_{(N_k)}$, $k = 0, 1, \dots, q$. By the definition of $\{N_k\}$ we know that for each J_j , $J_j \cap K \neq \emptyset$, there is a N_k such that $J_j \subset N_k$. Thus we have

$$\{I_i\}_{(N_k)} = \{I_i \cap K\}_{I_i \cap J = \emptyset, I_i \cap K_k \neq \emptyset} \cup \left(\bigcup_{\mathcal{R}_{J_j} \cap K_k \neq \emptyset} \{I_i\}_{(\mathcal{R}_{J_j} \cap (J \cup K))} \right).$$

Noting $\mathcal{R}(\{I_i\}_{(N_k)}) = N_k$, ($1 \leq k \leq q$), thus

$$J_j \subset N_k \Leftrightarrow \mathcal{R}_{J_j} \cap K_k \neq \emptyset.$$

It is obvious that we have $\{J_j\}_{J_j \subset N_k} \sqsubset \{I_i\}_{(N_k)}$.

$(\dots, (\{\mathbf{I}_i\} / (\mathbf{J}_j))_{(\mathbf{K}_k)}, \dots) = (\dots, \{\mathbf{I}_i\}_{(\mathbf{M}_\mu)}, \dots) / (\mathbf{J}_j)$:

Noting

$$\{I_i\}_{(M_\mu)} = \begin{cases} \{I_i\}_{(J_j)}, & \mathcal{R}_{J_j} \cap K = \emptyset, \\ \{I_i\}_{(N_k)}, & \text{others,} \end{cases}$$

thus, in the situation of $\mathcal{R}_{J_j} \cap K = \emptyset$, $\{I_i\}_{(J_j)} / \{I_i\}_{(J_j)} = \emptyset$, and in the part of $\{I_i\}_{(N_k)}$, by the expressions of $\{I_i\}_{(N_k)}$ and $\{L_\lambda\}_{L_\lambda \cap K_k \neq \emptyset}$, we have

$$\{L_\lambda\}_{(K_k)} = \{I_i\}_{(N_k)} / (J_j)_{J_j \subset N_k}.$$

$(\{\mathbf{I}_i\} / (\mathbf{J}_j)) / (\mathbf{K}_k) = \{\mathbf{I}_i\} / (\mathbf{M}_\mu)$:

We note that

$$\{I_i\}/(M_\mu) = \{I_i\}_{I_i \cap M = \emptyset} \cup \left(\bigcup_{j, \mathcal{R}_{J_j} \cap K = \emptyset} \{I_i\}_{I_i \cap J_j \neq \emptyset} / \{I_i\}_{(J_j)} \right) \cup \left(\bigcup_{k=0}^q \{I_i\}_{I_i \cap M_k \neq \emptyset} / \{I_i\}_{(M_k)} \right),$$

and

$$\{L_\lambda\}/(K_k) = \{L_\lambda\}_{L_\lambda \cap K = \emptyset} \cup \left(\bigcup_{k=1}^q \{L_\lambda\}_{L_\lambda \cap K_k \neq \emptyset} / \{L_\lambda\}_{(K_k)} \right).$$

It is obvious that

$$\{L_\lambda\}_{L_\lambda \cap K = \emptyset} = \{I_i\}_{I_i \cap M = \emptyset} \cup \left(\bigcup_{j, \mathcal{R}_{J_j} \cap K = \emptyset} \{I_i\}_{I_i \cap J_j \neq \emptyset} / \{I_i\}_{(J_j)} \right).$$

Noting $M_k = K_k \cup (\bigcup_{\mathcal{R}_{J_j} \cap K_k \neq \emptyset} J_j)$, hence, we have

$$\mathcal{R}_{\{L_\lambda\}, K_k} \setminus K_k = \mathcal{R}_{\{I_i\}, M_k} \setminus M_k.$$

Above formula implies

$$\{L_\lambda\}_{L_\lambda \cap K_k \neq \emptyset} / \{L_\lambda\}_{(K_k)} = \{I_i\}_{I_i \cap M_k \neq \emptyset} / \{I_i\}_{(M_k)},$$

where $1 \leq k \leq q$.

Up to now we have proved the proposition. □

As the corollary of theorem 2.1 we have:

Proposition 2.6. *The coproduct in definition 2.6 satisfies*

$$(\Delta \otimes id)\Delta = (id \otimes \Delta) \Delta. \quad (2.13)$$

Proof. For simplicity, we will replace the coproduct Δ in (2.13) by the reduce coproduct Δ' , where

$$\Delta'(\cdot) = \Delta(\cdot) - (\cdot) \otimes \emptyset - \emptyset \otimes (\cdot).$$

Here we only give the proof in the situation of

$$(\Delta' \otimes id) \Delta'(\{I_i\}) = (id \otimes \Delta') \Delta'(\{I_i\}), \quad (*)$$

where $\{I_i\} \in \mathcal{P}_{dis}^2(A)$. The general situation is similar.

Recalling the formula (2.11) in definition 2.6, we have

$$\Delta'\{I_i\} = \sum_{\{J_1, \dots, J_l\} \sqsubset \{I_i\}} (\{I_i\}_{(J_1)} \cdots, \{I_i\}_{(J_l)}) \otimes \{I_i\}/(J_j).$$

The right side of (*) should be of the following form,

$$\begin{aligned} & (id \otimes \Delta') \Delta'(\{I_i\}) \\ &= \sum_{\{J_1, \dots, J_l\} \sqsubset \{I_i\}} (\{I_i\}_{(J_1)} \cdots, \{I_i\}_{(J_l)}) \otimes \Delta'(\{I_i\}/(J_j)) \\ &= \sum_{\{J_j\} \sqsubset \{I_i\}, \{K_k\} \sqsubset \{L_\lambda\}} (\{I_i\}_{(J_j)}) \otimes (\{L_\lambda\}_{(K_k)}) \otimes (\{L_\lambda\}/(K_k)), \end{aligned}$$

where $\{L_\lambda\} = \{I_i\}/(J_j)$. Recalling proposition 2.8, we know that there is $\{M_\mu\} \in \mathcal{P}_{dis}^2(A)$ satiafying the following conditions:

$$\begin{aligned} & \{M_\mu\} \sqsubset \{I_i\}, \{J_j\} \sqsubset (\cdots, \{I_i\}_{(M_\mu)}, \cdots), \\ & (\cdots, (\{I_i\}/(J_j))_{(K_k)}, \cdots) = (\cdots, \{I_i\}_{(M_\mu)}, \cdots)/(J_j), \\ & (\{I_i\}/(J_j))/(K_k) = \{I_i\}/(M_\mu), \end{aligned}$$

then, we have

$$= \sum_{\{M_\mu\} \sqsubset \{I_i\}, \{J_j\} \sqsubset (\{I_i\}_{(M_\mu)})} (id \otimes \Delta') \Delta' (\{I_i\}) (\{I_i\}_{(M_\mu \cap J_j)}) \otimes (\{I_i\}_{(M_\mu)}/(J_j) \otimes (\{I_i\}/(M_\mu))).$$

Because $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{M_\mu\})$, we have $(\{I_i\}_{(M_\mu \cap J_j)}) = (\{I_i\}_{(J_j)})$. Finally, we have

$$\begin{aligned} & = \sum_{\{M_\mu\} \sqsubset \{I_i\}, \{J_j\} \sqsubset (\{I_i\}_{(M_\mu)})} (id \otimes \Delta') \Delta' (\{I_i\}) (\{I_i\}_{(J_j)}) \otimes (\{I_i\}_{(M_\mu)}/(J_j) \otimes (\{I_i\}/(M_\mu))) \\ & = (\Delta' \otimes id) \Delta' (\{I_i\}). \end{aligned}$$

□

Remark 2.8. Proposition 2.9 means that $V_{\mathbb{K},A}$ is a coalgebra under the coproduct in definition 2.6. It is easy to check that for each $\{I_i\} \in \mathcal{P}_{dis}^2(A)$ (or $(\{I_{i_1}^{(1)}\}, \cdots, \{I_{i_k}^{(k)}\}) \in \mathcal{P}_{dis,k}^2(A)$) $(\Delta')^m(\{I_i\}) = 0$ (or $(\Delta')^m((\{I_{i_1}^{(1)}\}, \cdots, \{I_{i_k}^{(k)}\})) = 0$) for some positive integer m , thus, from $V_{\mathbb{K},A}$ we can construct a Hopf algebra in a standard way (see D. E. Radford [8]).

3 Composition and Lie bracket

Let \mathbb{K} be a field of characteristic zero, $\mathbb{K}(\mathcal{P}_{dis}^2(A))$ denote the vector space over \mathbb{K} spanned by $\mathcal{P}_{dis}^2(A)$. We now go to the discussion of the composition between $\{I_i\}$ and $\{J_j\}$, where $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) = \emptyset$. When we focus on the composition, we restrict us to consider the non-travail insertion only. Similar to the cases of Feymman diagrams and Kontsevich's graphs, we have:

Definition 3.1. We define

$$\{I_i\} \circ_a \{J_j\} = \sum_{\iota} \{I_i\} \circ_{I_a}^{\iota} \{J_j\}, \quad (3.1)$$

and

$$\{I_i\} \circ \{J_j\} = \sum_a \{I_i\} \circ_a \{J_j\}. \quad (3.2)$$

The composition \circ defined by (3.2) is not associative generally. Let $\{I_i\}, \{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfying $I \cap J = \emptyset$, $K \cap (I \cup J) = \emptyset$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$ and $K = \mathcal{R}(\{K_k\})$. We are interested in the difference between $(\{I_i\} \circ \{J_j\}) \circ \{K_k\}$ and $\{I_i\} \circ (\{J_j\} \circ \{K_k\})$. In general, we have

$$(\{I_i\} \circ \{J_j\}) \circ \{K_k\} \neq \{I_i\} \circ (\{J_j\} \circ \{K_k\}).$$

Observing $(\{I_i\} \circ \{J_j\}) \circ \{K_k\}$, by (3.1) and (3.2) we know that

$$(\{I_i\} \circ \{J_j\}) \circ \{K_k\} = \left(\sum_{a,\iota} \{I_i\} \circ_{I_a}^{\iota} \{J_j\} \right) \circ \{K_k\},$$

and

$$(\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) \circ \{K_k\} = \sum_{b, \tau, a \neq b} (\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) \circ_{I_b}^{\tau} \{K_k\} + \sum_{c, \kappa} (\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) \circ_{J_c \cup \iota^{-1}(J_c)}^{\kappa} \{K_k\}.$$

Recalling proposition 2.5 we know that

$$\{I_i\} \circ_{I_a}^{\iota} \{J_j\} \circ_{J_c \cup \iota^{-1}(J_c)}^{\kappa} \{K_k\} = \{I_i\} \circ_{I_a}^{\iota'} (\{J_j\} \circ_{J_c}^{\kappa'} \{K_k\})$$

for some ι' and κ' . By proposition 2.4 we know that, for fixed I_a and J_c , there is an one-one corresponding between (ι, κ) and (ι', κ') . Summarizing previous discussions, we reach the following formula.

Proposition 3.1.

$$(\{I_i\} \circ \{J_j\}) \circ \{K_k\} = \sum_{a, b, a \neq b} \sum_{\iota, \tau} (\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) \circ_{I_b}^{\tau} \{K_k\} + \{I_i\} \circ (\{J_j\} \circ \{K_k\}). \quad (3.3)$$

We introduce a compact notation

$$<\{I_i\}, \{J_j\}, \{K_k\}> = \sum_{a, b, a \neq b} \sum_{\iota, \tau} (\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) \circ_{I_b}^{\tau} \{K_k\},$$

then the formula (3.3) can be rewritten as

$$(\{I_i\} \circ \{J_j\}) \circ \{K_k\} = \{I_i\} \circ (\{J_j\} \circ \{K_k\}) + <\{I_i\}, \{J_j\}, \{K_k\}>.$$

Now we have

Corollary 3.1.

$$<\{I_i\}, \{J_j\}, \{K_k\}> = <\{I_i\}, \{K_k\}, \{J_j\}>.$$

Proof. Noting

$$(\{I_i\} \circ_{I_a}^{\iota} \{J_j\}) \circ_{I_b}^{\tau} \{K_k\} = \{I_i\}_{i \neq a, b} \cup (I_a \sqcup_{\iota} \{J_j\}) \cup (I_b \sqcup_{\tau} \{K_k\}),$$

the conclusion will be implied immediately. \square

Corollary 3.2.

$$\begin{aligned} & \{I_i\} \circ (\{J_j\} \circ \{K_k\}) - (\{I_i\} \circ \{J_j\}) \circ \{K_k\} \\ &= \{I_i\} \circ (\{K_k\} \circ \{J_j\}) - (\{I_i\} \circ \{K_k\}) \circ \{J_j\}. \end{aligned}$$

Based on the discussion as above we are able to define the Lie bracket for partitions.

Definition 3.2. Let $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$ with $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) = \emptyset$, their Lie bracket is defined to be

$$[\{I_i\}, \{J_j\}] = \{I_i\} \circ \{J_j\} - \{J_j\} \circ \{I_i\}. \quad (3.4)$$

In order to prove the bracket (3.4) is Lie bracket, it is enough for us to check that Jacobi identity is valid.

Theorem 3.1. The bracket (3.4) satisfies Jacobi identity.

Proof. Let $\{I_i\}, \{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$ satisfying $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) \neq \emptyset$, $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{K_k\}) \neq \emptyset$, $\mathcal{R}(\{J_j\}) \cap \mathcal{R}(\{K_k\}) \neq \emptyset$, we want to prove Jacobi identity

$$[\{I_i\}, [\{J_j\}, \{K_k\}]] + cocycle = 0.$$

By a straightforward calculation we get

$$\begin{aligned} & [\{I_i\}, [\{J_j\}, \{K_k\}]] \\ &= \{I_i\} \circ (\{J_j\} \circ \{K_k\}) - \{I_i\} \circ (\{K_k\} \circ \{J_j\}) \\ & - (\{J_j\} \circ \{K_k\}) \circ \{I_i\} + (\{K_k\} \circ \{J_j\}) \circ \{I_i\}, \\ & [\{K_k\}, [\{I_i\}, \{J_j\}]] \\ &= \{K_k\} \circ (\{I_i\} \circ \{J_j\}) - \{K_k\} \circ (\{J_j\} \circ \{I_i\}) \\ & - (\{I_i\} \circ \{J_j\}) \circ \{K_k\} + (\{J_j\} \circ \{I_i\}) \circ \{K_k\}, \\ & [\{J_j\}, [\{K_k\}, \{I_i\}]] \\ &= \{J_j\} \circ (\{K_k\} \circ \{I_i\}) - \{J_j\} \circ (\{I_i\} \circ \{K_k\}) \\ & - (\{K_k\} \circ \{I_i\}) \circ \{J_j\} + (\{I_i\} \circ \{K_k\}) \circ \{J_j\}. \end{aligned}$$

With the help of corollary 3.2 we can get Jacobi identity. \square

The previous discussions can be generalized to more general situations. We assume that every partition $\{I_i\} \in \mathcal{P}_{dis}^2(A)$ assigns to a function $f : \{I_i\} \rightarrow \mathbb{N}$. Now we consider the Lie bracket concerning the pair $(\{I_i\}, f)$. Let $(\{I_i\}, f), (\{J_j\}, g)$ be two pairs, where $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, $\mathcal{R}(\{I_i\}) \cap \mathcal{R}(\{J_j\}) = \emptyset$. Then the insertion of two pairs is defined to be

$$(\{I_i\}, f) \circ_{I_a}^t (\{J_j\}, g) = (\{I_i\} \circ_{I_a}^t \{J_j\}, f \circ_a g), \quad (3.5)$$

where $f \circ_a g$ is defined as follows:

$$f \circ_a g(I_i) = f(I_i) \ (i \neq a), \ f \circ_a g(\iota^{-1}(J_j) \cup J_j) = g(J_j).$$

Above formula shows that $f \circ_a g$ is independent of ι . Now we define the composition of the pairs to be

$$(\{I_i\}, f) \circ_a (\{J_j\}, g) = \sum_{\iota} (\{I_i\}, f) \circ_{I_a}^t (\{J_j\}, g), \quad (3.6)$$

and

$$(\{I_i\}, f) \circ_{pair} (\{J_j\}, g) = \sum_a (-1)^{f(I_a)} (\{I_i\}, f) \circ_a (\{J_j\}, g). \quad (3.7)$$

We hope that the Lie bracket arising from the composition \circ_{pair} will be well defined. hence, we need to prove a conclusion similar to corollary 3.2 is valid. Let $(\{I_i\}, f), (\{J_j\}, g), (\{K_k\}, h)$ be three pairs, it is enough for us to consider the composition $(f \circ_a g) \circ_b h$. Similar to the situation of proposition 3.1, it is necessary for us to discuss the following two possibilities:

- **Case of $(\{I_i\} \circ_{I_a}^t \{J_j\}) \circ_{I_b}^t \{K_k\}$ ($a \neq b$) :** In this situation we have

$$\begin{cases} ((f \circ_a g) \circ_b h)(I_i) = (f \circ_a g)(I_i) = f(I_i), \ i \neq a, b, \\ ((f \circ_a g) \circ_b h)(\iota^{-1}(J_j) \cup J_j) = (f \circ_a g)(\iota^{-1}(J_j) \cup J_j) = g(J_j), \\ ((f \circ_a g) \circ_b h)(\tau^{-1}(K_k) \cup K_k) = h(K_k). \end{cases}$$

In this situation we have

$$((f \circ_a g) \circ_b h) = ((f \circ_b h) \circ_a g).$$

- **Case of $(\{I_i\} \circ_{I_a}^t \{J_j\}) \circ_{\iota^{-1}(J_c) \cup J_c}^{\kappa} \{K_k\}$ ($a \neq b$) :** In this situation we have

$$\begin{cases} ((f \circ_a g) \circ_b h)(I_i) = (f \circ_a g)(I_i) = f(I_i), i \neq a, \\ ((f \circ_a g) \circ_b h)(\iota^{-1}(J_j) \cup J_j) = (f \circ_a g)(\iota^{-1}(J_j) \cup J_j) = g(J_j), j \neq c, \\ ((f \circ_a g) \circ_b h)(\kappa^{-1}(K_k) \cup K_k) = h(K_k). \end{cases}$$

In this situation, it is easy to check that

$$(f \circ_a g) \circ_b h = f \circ_a (g \circ_c h).$$

Similar to previous discussions, we introduce the following notation

$$= \sum_{a,b,a \neq b} (-1)^{f(I_a)+f(I_b)} \sum_{\iota, \tau} \langle (\{I_i\} \circ_{I_a}^t \{J_j\}) \circ_{I_b}^{\tau} \{K_k\}, (f \circ_a g) \circ_b h \rangle_{pair}$$

Now we reach the following conclusion:

Proposition 3.2.

$$\begin{aligned} & \langle (\{I_i\}, f) \circ_{pair} (\{J_j\}, g) \rangle_{pair} \circ_{pair} \langle \{K_k\}, h \rangle_{pair} \\ &= \langle (\{I_i\}, f), (\{J_j\}, g), (\{K_k\}, h) \rangle_{pair} + \langle (\{I_i\}, f) \circ_{pair} ((\{J_j\}, g) \circ_{pair} (\{K_k\}, h)) \rangle_{pair}. \end{aligned} \quad (3.8)$$

$$\langle (\{I_i\}, f), (\{J_j\}, g), (\{K_k\}, h) \rangle_{pair} = \langle (\{I_i\}, f), (\{K_k\}, h), (\{J_j\}, g) \rangle_{pair}. \quad (3.9)$$

Up to now, we can easily see the conclusion similar to corollary 3.2 is valid.

Corollary 3.3.

$$\begin{aligned} & ((\{I_i\}, f) \circ_{pair} (\{J_j\}, g)) \circ_{pair} (\{K_k\}, h) - (\{I_i\}, f) \circ_{pair} ((\{J_j\}, g) \circ_{pair} (\{K_k\}, h)) \\ &= ((\{I_i\}, f) \circ_{pair} (\{K_k\}, h)) \circ_{pair} (\{J_j\}, g) - (\{I_i\}, f) \circ_{pair} ((\{K_k\}, h) \circ_{pair} (\{J_j\}, g)). \end{aligned} \quad (3.10)$$

Summarizing the previous discussions, we know that the following Lie bracket will be well defined.

Definition 3.3.

$$[(\{I_i\}, f), (\{J_j\}, g)]_{pair} = (\{I_i\}, f) \circ_{pair} (\{J_j\}, g) - (\{J_j\}, g) \circ_{pair} (\{I_i\}, f). \quad (3.11)$$

4 Applications to graphs

The construction discussed in section 2 and section 3 is suitable for graphs, for example, Feynman diagrams, Kontsevich's graphs and the ordinary graphs in the sense of graphic theory. For Feynman diagrams, we follow the notations in Jean-Louis Loday and N. M. Nikolov [5], actually, which is one of our motivation for the construction in this article. For simplicity, we do not consider the coloured Feynman diagrams. As a preparation we introduce some notations. Let A be a finite set, $\sigma : A \rightarrow A$ be an involution, $\sigma^2 = \text{id}$. A subset $J \subset A$ is called a σ -**invariant** subset, if $J = \sigma(J)$. Actually, for a subset $J \subset A$, it is easy to check that $J \cap \sigma(J)$ and $J \cup \sigma(J)$ are σ -invariant. Roughly speaking, a graph can be viewed as a finite set endowed with an involution and a decomposition.

4.1 The graphs in the sense of graphic theory

Definition 4.1. • A ordinary graph is a pair $(\sigma, \{I_i\})$, where $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, σ is a map from I to itself without fixed points, $\sigma^2 = \text{id}$, where $I = \mathcal{R}(\{I_i\})$. A graph $(\sigma, \{I_i\})$ is also denoted by $\Gamma_{\sigma, \{I_i\}}$, or, $\Gamma_{\{I_i\}}$ for short.

- If for any I_i and $I_{i'}$, there are some positive integers i_0, i_1, \dots, i_m ($i_0 = i, i_m = i'$), such that $\sigma(I_k) \cap I_{k+1} \neq \emptyset$ ($0 \leq k < m$), we say $\Gamma_{\{I_i\}}$ is a connected graph. Otherwise, we say $\Gamma_{\{I_i\}}$ is disconnected. We call $I_{i_0}, I_{i_1}, \dots, I_{i_m}$ the chain in $\Gamma_{\{I_i\}}$ connecting I_i and $I_{i'}$.
- Let $(\sigma, \{I_i\})$ be an ordinary graph, J be a σ -invariant subset of I . A subgraph of $\Gamma_{\{I_i\}}$ related to J is a pair $(\sigma|_J, \{I_i\}_{(J)})$, denoted by $\Gamma_J \subset \Gamma_{\{I_i\}}$ also. When $J = \mathcal{R}_{\{I_i\}, J} \cap \sigma(\mathcal{R}_{\{I_i\}, J})$, we call Γ_J a induced subgraph.

Remark 4.1. • We call $\{I_i\}$ the set of vertices of $\Gamma_{\{I_i\}}$, denoted by $\mathbf{Vert}(\Gamma_{\{I_i\}})$. I is called the total set of $\Gamma_{\{I_i\}}$. For a subgraph Γ_J , we identify $\mathbf{Vert}(\Gamma_J) = \{I_i \cap J\}$ with $\{I_i\}_{I_i \cap J \neq \emptyset}$.

- By the assumption of the involution σ , we know that there is no $e \in I$ such that $\sigma(e) = e$. Now we define an equivalent relation in I in the following way. We say $e_1 \sim e_2$ ($e_1, e_2 \in I$), if $e_1 = \sigma(e_2)$. It is easy to check that \sim is an equivalent relation, and each equivalent class consists of two pairs $(e, \sigma(e))$ and $(\sigma(e), e)$ ($e \in I$). Each equivalent class assigns an edge of $\Gamma_{\{I_i\}}$, then I/\sim is the set of all edges of $\Gamma_{\{I_i\}}$.
- Let $\tau : \underline{m} \rightarrow \underline{m}$ ($m = |\{I_i\}|$) be a permutation, we identify $(\sigma, \{I_i\})$ with $(\sigma, \{I_{\tau(i)}\})$.

Lemma 4.1. Let $\Gamma_{\sigma, \{I_i\}}$ be a graph, Γ_J and $\Gamma_{J'}$ are two connected subgraphs of $\Gamma_{\sigma, \{I_i\}}$. If $\mathcal{R}_{\{I_i\}, J} \cap \mathcal{R}_{\{I_i\}, J'} \neq \emptyset$, then $\Gamma_{J \cup J'}$ is a connected subgraph of $\Gamma_{\sigma, \{I_i\}}$.

Proof. We need to prove a fact i.e. for any I_i and $I_{i'}$ with $I_i \cap J \neq \emptyset$ and $I_{i'} \cap J' \neq \emptyset$ (or $I_i \subset \mathcal{R}_{\{I_i\}, J}$ and $I_{i'} \subset \mathcal{R}_{\{I_i\}, J'}$) there is a chain in $\Gamma_{J \cup J'}$ connecting $I_i \cap (J \cup J')$ and $I_{i'} \cap (J \cup J')$. Because of $\mathcal{R}_{\{I_i\}, J} \cap \mathcal{R}_{\{I_i\}, J'} \neq \emptyset$, there is I_α such that $I_\alpha \subset \mathcal{R}_{\{I_i\}, J}$ and $I_\alpha \subset \mathcal{R}_{\{I_i\}, J'}$, which means $I_\alpha \cap J \neq \emptyset$ and $I_\alpha \cap J' \neq \emptyset$. Noting both Γ_J and $\Gamma_{J'}$ are connected, thus there is a chain in Γ_J connecting $I_i \cap J$ with $I_\alpha \cap J$, and there is a chain in $\Gamma_{J'}$ connecting $I_{i'} \cap J'$ with $I_\alpha \cap J'$. Above two chains will result in a chain in $\Gamma_{J \cup J'}$ connecting $I_i \cap (J \cup J')$ with $I_{i'} \cap (J \cup J')$. Therefore, we have proved the conclusion. \square

Proposition 4.1. A graph $\Gamma_{\{I_i\}_{i=1}^m}$ is disconnected if and only if $\Gamma_{\{I_i\}_{i=1}^m}$ adapts the following decomposition

$$\Gamma_{\{I_i\}} = \bigcup_j \Gamma_{J_j}, \quad (4.1)$$

where $\{J_j\} \in \mathcal{P}_{dis}^2(A)$ ($2 \leq |\{J_j\}|$) satisfying the following conditions:

- $\mathcal{R}(\{J_j\}) = \mathcal{R}(\{I_i\})$, and $\{J_j\} \sqsubset \{I_i\}$.
- $\sigma(J_j) = J_j$, Γ_{J_j} is connected subgraph for each j .
- $\mathbf{Vert}(\Gamma_{J_j}) \cap \mathbf{Vert}(\Gamma_{J_{j'}}) = \emptyset$, $j \neq j'$.

Each Γ_{J_j} is a connected component of $\Gamma_{\{I_i\}}$. If we ignore the order in the decomposition (4.1), the decomposition (4.1) is unique.

Proof. For I_i and $I_{i'}$, we say $I_i \sim I_{i'}$, if there is a chain connecting I_i and $I_{i'}$. \sim is an equivalent relation obviously. Thus there is a partition $\{K_k\} \in \mathbf{Part}(\underline{m})$ such that

$$\{I_i\}_{i=1}^m = \bigcup_k \{I_i\}_{i \in K_k},$$

where each $\{I_i\}_{i \in K_k}$ is an equivalent class. We take $J_j = \mathcal{R}(\{I_i\}_{i \in K_j})$. We need to prove $\{J_j\}$ constructed in such a way satisfies the conditions in proposition. It is obvious that $\mathcal{R}(\{J_j\}) = \mathcal{R}(\{I_i\})$, and $\{J_j\} \subset \{I_i\}$. We now prove $\sigma(J_j) = J_j$ for each j . It is enough for us to prove for any j and j' ($j \neq j'$), we have $\sigma(J_j) \cap J_{j'} = \emptyset$. Otherwise, there are j and j' such that $\sigma(J_j) \cap J_{j'} \neq \emptyset$. Then, there are $i \in K_j$ and $i' \in K_{j'}$, $\sigma(I_i) \cap I_{i'} \neq \emptyset$, which means $\{I_i\}_{i \in K_j} \cup \{I_i\}_{i \in K_{j'}}$ should be included in some equivalent class. That is a contradiction. The procedure to construct J_j implies each Γ_{J_j} is a connected subgraph of $\Gamma_{\{I_i\}}$. It is easy to check that for each connected subgraph Γ_J there is a j such that $J \subset J_j$. Thus each Γ_{J_j} is a connected component of $\Gamma_{\{I_i\}}$. Furthermore, we know that $\Gamma_{\{I_i\}}$ is disconnected if and only if $|\{J_j\}| \geq 2$. The uniqueness of the decomposition (4.1) is obvious. \square

Remark 4.2. Let $\Gamma_{\sigma, \{I_i\}}$ be a graph, $\{J_j\} \in \mathcal{P}_{dis}^2(A)$. If $\{J_j\} \subset \{I_i\}$ and $\sigma(J_j) = J_j$ for any j , we say $\{J_j\}$ σ -admits to $\{I_i\}$ denoted by $\{J_j\} \subset_{\sigma} \{I_i\}$.

Proposition 4.2. Let $\Gamma_{\sigma, \{I_i\}}$ be a connected graph, $\{J_j\} \in \mathcal{P}_{dis}^2(A)$ satisfy $J \subset I$ and $\sigma(J_j) = J_j$ for any j , where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$. Then, the subgraph Γ_J admits the following decomposition

$$\Gamma_J = \bigcup_l \Gamma_{L_l}, \quad \mathbf{Vert}(\Gamma_{L_l}) \cap \mathbf{Vert}(\Gamma_{L_{l'}}) = \emptyset, \quad l \neq l', \quad (4.2)$$

where $\{L_l\} = \{J_j\}_{ad, \{I_i\}_{(J)}} \subset \{I_i\}_{(J)}$, and each Γ_{L_l} is a connected component of Γ_J .

Proof. According to proposition 4.1, for subgraph Γ_J there is the unique decomposition based on its connected components. Here we need to prove this decomposition is exactly given by $\{L_l\} = \{J_j\}_{ad, \{I_i\}_{(J)}}$.

It is obvious that $\sigma(L_l) = L_l$ for each l , thus each Γ_{L_l} is a subgraph of Γ_J . By proposition 2.6 we know that $\{L_l\} \subset \{I_i\}_{(J)}$. In addition, noting $\mathcal{R}(\{L_l\}) = J$, it is easy to prove $\mathcal{R}_{\{I_i\}_{(J)}, L_l} = L_l$ for each l . Recalling the procedure to construct $\{L_l\}$ in the proof of proposition 2.6, combining with lemma 4.1, we know that Γ_{L_l} is a connected subgraph of Γ_J for each l . It is obvious that as subgraphs of Γ_J , $\mathbf{Vert}(\Gamma_{L_l}) \cap \mathbf{Vert}(\Gamma_{L_{l'}}) = \emptyset$ ($l \neq l'$). Therefore, $\{L_l\}$ satisfies all conditions in proposition 4.2, which means Γ_{L_l} is a connected component of Γ_J for each l . \square

We now turn to the quotient of the ordinary graphs. The discussions below will follow the idea of Connes-Krieger theory, but a different description will be provided based on the setting in this article.

Definition 4.2. Let $\Gamma_{\sigma, \{I_i\}}$ be a connected ordinary graph, Γ_J be a connected subgraph of $\Gamma_{\sigma, \{I_i\}}$ determined by a σ -invariant subset $J \subset I$. We define the quotient of $\Gamma_{\sigma, \{I_i\}}$ by Γ_J to be a pair $(\sigma|_{I \setminus J}, \{I_i\} / \{J_j\})$, denoted by $\Gamma_{\sigma, \{I_i\}} / \Gamma_J$ also, where $I = \mathcal{R}(\{I_i\})$, $\{J_j\} = \{I_i\}_{(J)}$.

Remark 4.3. Let $\{J_j\} \in \mathcal{P}_{dis}^2(A)$ satisfy $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{I_i\})$ and $\sigma(J_j) = J_j$ for any j . Then, it is natural for us to define the quotient of $\Gamma_{\sigma, \{I_i\}}$ by the sequence of the subgraphs $\{J_j\}$ to be $(\sigma|_{I \setminus J}, \{I_i\} / (J_j))$, where $J = \mathcal{R}(\{J_j\})$. Above quotient can also be denoted by $\Gamma_{\sigma, \{I_i\}} / (\Gamma_J)$. Combining proposition 2.7 and proposition 4.1, we have $\Gamma_{\sigma, \{I_i\}} / (\Gamma_J) = \Gamma_{\sigma, \{I_i\}} / (\Gamma_{L_l})$, where

$\{L_i\} = \{J_j\}_{ad, \{I_i\}_{(J)}}$ and each Γ_{L_i} is an connected component of Γ_J . Thus, when we discuss the quotient of the graphs, we can always assume $\{J_j\} \sqsubset_\sigma \{I_i\}_{(J)}$.

Proposition 4.3. *Let $\Gamma_{\sigma, \{I_i\}}$ be a connected ordinary graph, Γ_J be a connected subgraph of $\Gamma_{\sigma, \{I_i\}}$ determined by a σ -invariant subset $J \subset I$. Then $\Gamma_{\sigma, \{I_i\}} / \Gamma_J$ is a connected graph.*

Proof. If $\mathcal{R}_J \setminus J = \emptyset$, the conclusion is obviously valid. Now we assume $\mathcal{R}_J \setminus J \neq \emptyset$. Noting $\text{Vert}(\Gamma_{\sigma, \{I_i\}} / \Gamma_J) = \{I_i\}_{I_i \cap J \neq \emptyset} \cup \{\mathcal{R}_J \setminus J\}$, because $\Gamma_{\sigma, \{I_i\}}$ is a connected graph, we know that $\sigma(\mathcal{R}(\{I_i\}_{I_i \cap J \neq \emptyset})) \cap \mathcal{R}_J \neq \emptyset$. We need to prove $\sigma(\mathcal{R}(\{I_i\}_{I_i \cap J \neq \emptyset})) \cap (\mathcal{R}_J \setminus J) \neq \emptyset$. Noting $\mathcal{R}(\{I_i\}_{I_i \cap J \neq \emptyset}) \cap J = \emptyset$, thus $\sigma(\mathcal{R}(\{I_i\}_{I_i \cap J \neq \emptyset})) \cap J = \emptyset$, which means $\sigma(\mathcal{R}(\{I_i\}_{I_i \cap J \neq \emptyset})) \cap \mathcal{R}_J \subset \mathcal{R}_J \setminus J$. □

Combining proposition 4.2 and proposition 4.3 we have the following corollary.

Corollary 4.1. *Let $\Gamma_{\sigma, \{I_i\}}$ be a connected graph, $\{J_j\} \in \mathcal{P}_{dis}^2(A)$ satisfying $\mathcal{R}(\{J_j\}) \subset \mathcal{R}(\{I_i\})$ and $\sigma(J_j) = J_j$ for any j . Then $\Gamma_{\sigma, \{I_i\}} / (\Gamma_{J_j})$ is a connected graph.*

In the situation of the ordinary graphs there is an analogue of theorem 2.1.

Proposition 4.4. *Let $\Gamma_{\sigma, \{I_i\}}$ be a connected graph, $\{J_j\}, \{K_k\} \in \mathcal{P}_{dis}^2(A)$. $\{I_i\}, \{J_j\}, \{K_k\}$ satisfy the following conditions:*

- $\{J_j\} \sqsubset \{I_i\}, \{K_k\} \sqsubset \{I_i\} / (J_j)$.
- J_j is σ -invariant and Γ_{J_j} is a connected subgraph in $\Gamma_{\sigma, \{I_i\}}$ for each j .
- K_k is $\sigma_{I \setminus J}$ -invariant and Γ_{K_k} is a connected subgraph of $\Gamma_{\sigma, \{I_i\}} / (\Gamma_{J_j})$ for each k , where $I = \mathcal{R}(\{I_i\}), J = \mathcal{R}(\{J_j\})$.

Then there is a partition $\{M_\mu\} \in \mathcal{P}_{dis}^2(A)$ satisfying the following conditions:

- $\sigma(M_\mu) = M_\mu$ and Γ_{M_μ} is connected for each μ .
- $\{M_\mu\} \sqsubset \{I_i\}, \{J_j\} \sqsubset (\cdots, \{I_i\}_{(M_\mu)}, \cdots)$.
- $(\cdots, (\{I_i\} / (J_j))_{(K_k)}, \cdots) = (\cdots, \{I_i\}_{(M_\mu)}, \cdots) / (J_j)$.
- $(\Gamma_{\{I_i\}} / (\Gamma_{J_j})) / (\Gamma_{K_k}) = \Gamma_{\{I_i\}} / (\Gamma_{M_\mu})$.

Proof. The proof of proposition 4.4 is almost same as the proof of proposition 2.7. Here we only need to prove each M_μ is σ -invariant. Recalling proposition 2.7, we know that

$$\begin{aligned} \{M_\mu\} &= \{J_j\}_{\mathcal{R}_{J_j} \cap K = \emptyset} \cup \{N_k\}, \\ N_k &= K_k \cup \left(\bigcup_{\mathcal{R}_{J_j} \cap K_k \neq \emptyset} J_j \right), \quad k = 1, \dots, |\{K_k\}|, \end{aligned}$$

where $K = \mathcal{R}(\{K_k\})$. It is easy to check that $\sigma(M_\mu) = M_\mu$ for each μ . □

We now consider the insertion of the ordinary graphs.

Definition 4.3. *Let $(\sigma, \{I_i\}_{i=1}^m), (\lambda, \{J_j\}_{j=1}^n)$ be two connected ordinary graphs, with structure maps σ and λ respectively, $I = \mathcal{R}(\{I_i\}), J = \mathcal{R}(\{J_j\}), I \cap J = \emptyset$. Then we define the insertion of $\Gamma_{\{J_j\}}$ into $\Gamma_{\{I_i\}}$ at I_a by ι to be a pair $(\delta, \{I_i\} \circ_a^\iota \{J_j\})$ denoted by $\Gamma_{\{I_i\}} \circ_a^\iota \Gamma_{\{J_j\}}$, or, $\Gamma_{\{I_i\}} \circ_a^\iota \{J_j\}$ also, where $1 \leq a \leq m, \iota: I_a \rightarrow \{J_j\}, \delta|_I = \sigma, \delta|_J = \lambda$.*

It is obvious that we have:

Proposition 4.5. *Let $\Gamma_{\sigma, \{I_i\}}, \Gamma_{\delta, \{J_j\}}$ be two connected graphs. Then, $\Gamma_{\{I_i\}} \circ_a^t \Gamma_{\{J_j\}}$ is a connected graph also.*

Based on the discussions in section 3, we can prove there is a well-defined Lie bracket structure on the vector space $\text{Span}_{\mathbb{K}}(\mathcal{G}_c)$, where \mathcal{G}_c denotes the set of all connected graphs, \mathbb{K} is a field of characteristic zero. Here we only consider the composition \circ in definition 3.1 for simplicity. The composition will be defined in a natural way. Let $\Gamma_{\sigma, \{I_i\}}, \Gamma_{\delta, \{J_j\}}$ be two connected graphs. We have:

$$\Gamma_{\sigma, \{I_i\}} \circ \Gamma_{\delta, \{J_j\}} = \sum_{a, t} \Gamma_{\{I_i\}} \circ_a^t \Gamma_{\{J_j\}}. \quad (4.3)$$

The Lie bracket should be defined to be:

$$[\Gamma_{\sigma, \{I_i\}}, \Gamma_{\delta, \{J_j\}}] = \Gamma_{\sigma, \{I_i\}} \circ \Gamma_{\delta, \{J_j\}} - \Gamma_{\delta, \{J_j\}} \circ \Gamma_{\sigma, \{I_i\}}. \quad (4.4)$$

4.2 Feynman diagrams

Definition 4.4. • A Feynman diagram is a pair $(\sigma, \{I_i\})$, where $\{I_i\} \in \mathcal{P}_{dis}^2(A)$, σ is a map from I to itself, $\sigma^2 = \sigma$, where $I = \mathcal{R}(\{I_i\})$. A Feynman diagram $(\sigma, \{I_i\})$ is also denoted by $\Gamma_{\sigma, \{I_i\}}$, or, $\Gamma_{\{I_i\}}$ for short. $I = \mathcal{R}(\{I_i\})$ is called the total set of edges.

- We call $I_{ext} = \{e \in I | \sigma(e) = e\}$ the set of external lines and $I_{int} = (I \setminus I_{ext}) / \sim$ the set of internal lines. We say a Feynman diagram $(\sigma, \{I_i\})$ is connected if $(\sigma|_{I \setminus I_{ext}}, \{I_i\}_{(I \setminus I_{ext})})$ is connected.

Remark 4.4. • It is obvious that $I \setminus I_{ext}$ is σ -invariant, and $\sigma|_{I \setminus I_{ext}}$ has no fixed points, thus $(\sigma|_{I \setminus I_{ext}}, \{I_i\}_{(I \setminus I_{ext})})$ is an ordinary graph denoted by $\Gamma_{\{I_i\}, int}$. $\Gamma_{\{I_i\}, int}$ is a Feynman diagram without external lines. A general Feynman diagram can be regarded as an extension of an ordinary graph. In fact, let $(\sigma, \{I_i\})$ be an ordinary graph, i.e. $I_{ext} = \emptyset$, now we want to add the set of external lines \tilde{I}_{ext} to I by a map $f : \tilde{I}_{ext} \rightarrow \{I_i\}$. The new Feynman diagram is naturally chosen to be the pair $(\omega, \tilde{I}_{ext} \sqcup_f \{I_i\})$, the new structure map ω is defined to be $\omega|_I = \sigma$, $\omega|_{\tilde{I}_{ext}} = id$. A Feynman diagram $(\sigma, \{I_i\})$ with $I_{ext} \neq \emptyset$ can be rewritten as $(\sigma, I_{ext} \sqcup_f \{I_i\}_{(I \setminus I_{ext})})$, which is an extension of $(\sigma|_{I \setminus I_{ext}}, \{I_i\}_{(I \setminus I_{ext})})$, where $f^{-1}(I_i \setminus I_{ext}) = I_i \cap I_{ext}$.

- We call $\{I_i\}$ the set of vertices of $\Gamma_{\{I_i\}}$ denoted by $\mathbf{vert}(\Gamma_{\{I_i\}})$. We identify the vertices of $\Gamma_{\{I_i\}}$ with ones of $\Gamma_{\{I_i\}, int}$, i.e. we have $\mathbf{vert}(\Gamma_{\{I_i\}}) = \mathbf{Vert}(\Gamma_{\{I_i\}, int})$. Actually, for a graph $\Gamma_{\{I_i\}}$, there are two structure maps, σ and projection $p : \mathcal{R}(\{I_i\}) \rightarrow \{I_i\}$, $p(e) = I_i \iff e \in I_i$.

Definition 4.5. Let $(\sigma, \{I_i\})$ be a connected Feynman diagram, $J \subset I$ be a subset, where $I = \mathcal{R}(\{I_i\})$. We call the pair $(\sigma_J, \{I_i\}_{I_i \cap J \neq \emptyset})$ is a sub-diagram of $(\sigma, \{I_i\})$, if J satisfies the following conditions:

- $$\sigma(J) \cap \mathcal{R}_J = J \cap \sigma(\mathcal{R}_J),$$
- If $I_i \cap J \neq \emptyset$, then $I_i \cap [(J \cap \sigma(\mathcal{R}_J)) \setminus I_{ext}] \neq \emptyset$,

where $\mathcal{R}_J = \mathcal{R}(\{I_i\}_{I_i \cap J \neq \emptyset})$ is the total set of $(\sigma_J, \{I_i\}_{I_i \cap J \neq \emptyset})$. The structure map $\sigma_J : \mathcal{R}_J \rightarrow \mathcal{R}_J$ is defined to be:

$$\sigma_J(e) = \begin{cases} \sigma(e), & e \in J \cap \sigma(\mathcal{R}_J), \\ e, & e \in \mathcal{R}_J \setminus (J \cap \sigma(\mathcal{R}_J)). \end{cases}$$

The sub-diagram is also denoted by $\Gamma_J \subset \Gamma_{\{I_i\}}$. When $J \cap \sigma(\mathcal{R}_J) = \mathcal{R}_J \cap \sigma(\mathcal{R}_J)$ we call Γ_J a subgraph.

Remark 4.5. • The first condition in definition 4.5 means that $J \cap \sigma(\mathcal{R}_J)$ is σ -invariant.

- The second condition means that each vertex of Γ_J is the endpoint of at least one internal line in Γ_J .
- For subgraph Γ_J , $\mathcal{R}_J \setminus (J \cap \sigma(\mathcal{R}_J))$ plays the role of external lines of Γ_J . Here, the subset $[(\mathcal{R}_J \cap \sigma(\mathcal{R}_J)) \setminus (J \cap \sigma(\mathcal{R}_J))] \setminus I_{ext}$ is regarded as a subset of external lines of Γ_J . Actually, the external momenta corresponding to this subset will be canceled each other.
- The set of external lines in Γ_J should be

$$(\mathcal{R}_J \setminus (J \cap \sigma(\mathcal{R}_J))) \cup (\mathcal{R}_J \cap I_{ext}).$$

The set of internal lines in Γ_J is

$$((J \cap \sigma(\mathcal{R}_J)) \setminus I_{ext}) / \sim.$$

Let $(\sigma, \{I_i\})$ be a Feynman diagram, we call a σ -invariant subset J is internal, if $J \cap I_{ext} = \emptyset$. We can prove that each sub-diagram of $(\sigma, \{I_i\})$ can be uniquely determined by an internal σ -invariant subset.

Proposition 4.6. *There is an one-one corresponding between the sub-diagrams and internal σ -invariant subsets.*

Proof. Let $(\sigma, \{I_i\})$ be a Feynman diagram, $J \subset I = \mathcal{R}(\{I_i\})$ satisfy the conditions in definition 4.5. Then, $J' = (J \cap \sigma(\mathcal{R}_J)) \setminus I_{ext}$ is an internal σ -invariant subset. Due to the first condition in definition 4.5, we know that $\mathcal{R}_J = \mathcal{R}_{J'}$. Thus $\Gamma_J = \Gamma_{J'}$. Conversely, let $J \subset I$ be an σ -invariant subset, then, J satisfies all conditions in definition 4.5. Thus J determines a sub-diagram of $(\sigma, \{I_i\})$. □

Remark 4.6. • Let $(\sigma, \{I_i\})$ be a Feynman diagram, and $J \subset I = \mathcal{R}(\{I_i\})$ be an internal σ -invariant subset. Then, the set of external lines in Γ_J just be $\mathcal{R}_J \setminus J$, and the set of internal lines in Γ_J is J / \sim .

- By the definition 4.1 and 4.5, we know that there is much difference between the sub-diagrams of Feynman diagrams and subgraphs of the ordinary graphs. Noting $(\sigma|_{I \setminus I_{ext}}, \{I_i\}_{(I \setminus I_{ext})})$ can be regarded as a ordinary graph, the conclusion of proposition 4.6 means that there is an one-one correspondence between the sub-diagrams of Feynman diagram $(\sigma, \{I_i\})$ and the subgraphs of $(\sigma|_{I \setminus I_{ext}}, \{I_i\}_{(I \setminus I_{ext})})$ in the sense of the ordinary graphs. Thus, when we discuss the sub-diagrams of Feynman diagrams, the discussion can be reduced to the situations of the ordinary graphs.

From proposition 4.2 and proposition 4.6 we have,

Proposition 4.7. *Let $\Gamma_{\{I_i\}}$ be a Feynman diagram, Γ_J be a sub-diagram, where J is an internal σ -invariant subset. If Γ_J is a disconnected sub-diagram, then Γ_J admits to the decomposition as follows:*

$$\Gamma_J = \bigcup_j \Gamma_{J_j},$$

where $\{J_j\}$ satisfies the following conditions:

- $\{J_j\} \in \mathbf{Part}(J)$, $|\{J_j\}| \geq 2$,
- each J_j is an internal σ -invariant subset,
- $\{J_j\} \sqsubset \{I_i\}$,
- each Γ_{J_j} is a connected component of Γ_J .

The decomposition of Γ_J is unique.

We now consider the quotient of Feynman diagrams.

Definition 4.6. *Let $\Gamma_{\{I_i\}}$ be a connected Feynman diagram, Γ_J be a connected proper sub-diagram of $\Gamma_{\{I_i\}}$ determined by a internal σ -invariant subset $J \subset I$. We define the quotient of $\Gamma_{\{I_i\}}$ by Γ_J to be a pair $(\sigma|_{I \setminus J}, \{I_i\} / \{J_j\})$, denoted by $\Gamma_{\{I_i\}} / \Gamma_J$ also, where $\{J_j\} = \{I_i\}_{(J)}$.*

Remark 4.7.

- Comparing with definition 4.2, in the situation of $I_{ext} = \emptyset$, the quotient of Feynman diagrams is same as the quotient of the ordinary graphs exactly. Recalling the contents about f -union in section 2 and discussions in remark 4.1, the quotient of general Feynman diagrams can be reduced to the situation of the ordinary graphs. Actually, a Feynman diagram $(\sigma, \{I_i\})$ can be expressed as $(\sigma, I_{ext} \sqcup_f \{I_i\}_{(I \setminus I_{ext})})$, where $f : I_{ext} \rightarrow \{I_i\}_{(I \setminus I_{ext})}$ satisfies $f^{-1}(I_i \cap (I \setminus I_{ext})) = I_i \cap I_{ext}$. Therefore, for an internal σ -invariant subset J , from the viewpoint of partition, the quotient $\Gamma_{\{I_i\}_{(I \setminus I_{ext})}} / \{I_i\}_{(J)}$ results in a projection $p_J : \{I_i\}_{(I \setminus I_{ext})} \rightarrow \{I_i \cap (I \setminus I_{ext})\}_{I_i \cap J = \emptyset} \cup \{(\mathcal{R}_J \cap (I \setminus I_{ext})) \setminus J\}$, such that

$$p_J(I_i \cap (I \setminus I_{ext})) = \begin{cases} (\mathcal{R}_J \cap (I \setminus I_{ext})) \setminus J, & I_i \cap J \neq \emptyset, \\ I_i \cap (I \setminus I_{ext}), & I_i \cap J = \emptyset. \end{cases}$$

Then

$$\Gamma_{I_{ext} \sqcup_f \{I_i\}_{(I \setminus I_{ext})}} / \Gamma_{J_{ind}} = \Gamma_{I_{ext} \sqcup_{p_J \circ f} (\{I_i\}_{(I \setminus I_{ext})} / \{I_i\}_{(J)})}.$$

- If a sub-diagram Γ_J determined by an internal σ -invariant subset J is not connected, then, according to proposition 4.1 or proposition 4.7, J admits to the decomposition $J = \bigcup_{j=1}^p J_j$, such that each Γ_{J_j} is a connected component of Γ_J . The quotient $\Gamma_{\{I_i\}}$ by Γ_J should be

$$\begin{aligned} \Gamma_{\{I_i\}} / \Gamma_J &= (\sigma|_{I \setminus J}, \{I_i\} / (J_j)) \\ &= (\cdots ((\Gamma_{\{I_i\}} / \Gamma_{J_1} / \Gamma_{J_2}) \cdots) / \Gamma_{J_p}). \end{aligned}$$

Above quotient is also denoted by

$$\Gamma_{\{I_i\}} / (\Gamma_{J_j}) = \Gamma_{\{I_i\} / (J_j)}.$$

The previous discussions show that the quotient of Feynman diagrams can be reduced to the situation of the ordinary graphs. Then we have,

Proposition 4.8. *Let $\Gamma_{\sigma, \{I_i\}}$ be a connected Feynman diagram, $J \subset I = \mathcal{R}(\{I_i\})$ be an internal σ -invariant subset. Then, $\Gamma_{\{I_i\}} / \Gamma_J$ is a connected Feynman diagram.*

Furthermore, a conclusion similar to proposition 4.4 or theorem 2.1 is valid.

Definition 4.7. *Let $(\sigma, \{I_i\}_{i=1}^m)$, $(\lambda, \{J_j\}_{j=1}^n)$ be two connected Feynman diagrams satisfying $I \cap J = \emptyset$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$. Then we define the insertion of $\Gamma_{\{J_j\}}$ into $\Gamma_{\{I_i\}}$ at I_a by ι to be a pair $(\delta, \{I_i\} \circ_{I_a}^{\iota} \{J_j\}_{(J \setminus J_{ext})})$ denoted by $\Gamma_{\{I_i\}} \circ_{I_a}^{\iota} \Gamma_{\{J_j\}}$, or, $\Gamma_{\{I_i\} \circ_{I_a}^{\iota} \{J_j\}_{(J \setminus J_{ext})}}$ also, where $1 \leq a \leq m$, $\iota : I_a \rightarrow \{J_j\}$, $\delta|_I = \sigma$, $\delta|_{J \setminus J_{ext}} = \lambda$.*

Remark 4.8. • *In the case of Feynman diagram, by definition 4.4 it is easy to check that $(\{I_i\} \circ_{I_a}^{\iota} \{J_j\}_{(J \setminus J_{ext})})_{(J \setminus J_{ext})} = \{J_j\}_{(J \setminus J_{ext})}$, and $\delta|_{J \setminus J_{ext}} = \lambda|_{J \setminus J_{ext}}$. $J \setminus J_{ext}$ is an internal δ -invariant subset obviously. Then, by proposition 2.3 we have*

$$\Gamma_{(\{I_i\} \circ_{I_a}^{\iota} \{J_j\}_{(J \setminus J_{ext})}) / \{J_j\}_{(J \setminus J_{ext})}} = \Gamma_{\{I_i\}}.$$

If $\Gamma_{\{I_i\}}$ and $\Gamma_{\{J_j\}}$ are connected, so is $\Gamma_{\{I_i\} \circ_{I_a}^{\iota} \{J_j\}_{(J \setminus J_{ext})}}$.

- *Recalling Feynman diagram $(\sigma, \{I_i\})$ can be rewritten as $(\sigma, I_{ext} \sqcup_f \{I_i\}_{(I \setminus I_{ext})})$, where f satisfies $f^{-1}(I_i \setminus I_{ext}) = I_i \cap I_{ext}$, as an application of proposition 2.4, we know that there is a map $F : I_{ext} \rightarrow \{I_i\}_{(I \setminus I_{ext})} \circ_{I_a}^{\iota'} \{J_j\}_{(J \setminus J_{ext})}$ ($\iota' = \iota|_{I_a \setminus I_{ext}}$) such that*

$$(I_{ext} \sqcup_f \{I_i\}_{(I \setminus I_{ext})}) \circ_{I_a}^{\iota} \{J_j\}_{(J \setminus J_{ext})} = I_{ext} \sqcup_F (\{I_i\}_{(I \setminus I_{ext})} \circ_{I_a \setminus I_{ext}}^{\iota'} \{J_j\}_{(J \setminus J_{ext})}).$$

- *In the same way as subsection 4.1, we can define the composition of Feynman diagrams as follows:*

$$\Gamma_{\{I_i\}} \circ \Gamma_{\{J_j\}} = \sum_{a, \iota} \Gamma_{\{I_i\}} \circ_{I_a}^{\iota} \Gamma_{\{J_j\}}.$$

The Lie bracket of Feynman diagrams can be defined to be

$$[\Gamma_{\{I_i\}}, \Gamma_{\{J_j\}}] = \Gamma_{\{I_i\}} \circ \Gamma_{\{J_j\}} - \Gamma_{\{J_j\}} \circ \Gamma_{\{I_i\}}.$$

Lie bracket above mentioned will be well defined obviously.

4.3 Kontsevich graphs

Definition 4.8. *An admissible graph is a triple $(\sigma, \{I_i\}, \{J_j\})$, where $\{I_i\}, \{J_j\} \in \mathcal{P}_{dis}^2(A)$, $I \cap J = \emptyset$, σ is a map from $I \cup J$ to itself without fixed points, $\sigma^2 = \text{id}$, where $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$. $\{I_i\}$ and $\{J_j\}$ satisfy*

$$\sigma(I_i) \cap I_i = \emptyset, \sigma(J_j) \cap J_j = \emptyset, |\sigma(I_i) \cap I_{i'}| \leq 1, |\sigma(I_i) \cap J_j| \leq 1, \quad \forall i, i', j, j', i \neq i'. \quad (4.5)$$

We denote the admissible graph $(\sigma, \{I_i\}, \{J_j\})$ by $\Gamma_{\sigma, \{I_i\}, \{J_j\}}$, or, $\Gamma_{\{I_i\}, \{J_j\}}$ for short. $I \cup J$ is called the total set of $\Gamma_{\{I_i\}, \{J_j\}}$. $\text{Vert}(\Gamma_{\sigma, \{I_i\}, \{J_j\}}) = \{I_i\} \cup \{J_j\}$. The vertices from $\{I_i\}$ are called the vertices of the first type, the vertices from $\{J_j\}$ are called the vertices of the second type.

Remark 4.9. • *The conditions in (4.5) means that:*

- There no loop in $\Gamma_{\{I_i\},\{J_j\}}$.
- For any two vertices there is at most one edge to connect them.
- There is not edge to connect any two vertices of the second type.
- In definition 4.8, we ignore the orientation of the edges. Now we discuss the issues of the orientation in details. Recalling the previous discussions, the edges of a graph are regarded as the equivalent classes under the equivalent relation defined as above, each equivalent class consists of two elements $(e, \sigma(e))$ and $(\sigma(e), e)$ ($e \in I \cup J$). To indicate the orientation of an edge, we can only choose one element in each equivalent class, for example, $(e, \sigma(e))$, to represent an edge. If the first element $e \in I_i$, the second element $\sigma(e) \in I_{i'}$ (or $\sigma(e) \in J_j$), then I_i will be the starting point of the edge $(e, \sigma(e))$, $I_{i'}$ (or J_j) will be the endpoint of this edge. Furthermore, each J_j can not be the starting point of any edge.
- The connectivity of an admissible graph can be defined in the same way as definition 4.1.

We now pay attention to subgraphs.

Lemma 4.2. Let $\Gamma_{\sigma, \{I_i\}, \{J_j\}}$ be an admissible graph. Then $\sigma(J) \cup J$ and $I \setminus \sigma(J)$ are two σ -invariant subsets. Additionally, for a σ -invariant subset $K \subset I \cup J$, we have

$$K = (\sigma(K') \cup K') \cup K'',$$

where $K' = K \cap J$, $K'' = K \cap (I \setminus \sigma(J))$.

Definition 4.9. Let $(\sigma, \{I_i\}, \{J_j\})$ be an admissible graph, and $K \subset I \cup J$ be a σ -invariant subset such that $K \cap J \neq \emptyset$, where $J = \mathcal{R}(\{J_j\})$. A subgraph of $\Gamma_{\sigma, \{I_i\}, \{J_j\}}$ determined by K is a triple $(\sigma|_K, \{I_i\}_{(K)}, \{J_j\}_{(K)})$, also denoted by $\Gamma_{\{I_i\}_{(K)}, \{J_j\}_{(K)}}$, or Γ_K for short.

Now we discuss the quotient and insertion of the admissible graphs.

Definition 4.10. Let $\Gamma_{\sigma, \{I_i\}, \{J_j\}}$ be a connected admissible graph, $K \subset I \cup J$ be a σ -invariant subset satisfying:

- $|(\mathcal{R}_{\{J_j\}, K \cap J} \setminus K) \cap (\mathcal{R}_{\{I_i\}, K \cap I} \setminus K)| \leq 1$,
- $|(\mathcal{R}_{\{J_j\}, K \cap J} \setminus K) \cap I_i| \leq 1$, $I_i \cap K = \emptyset$,
- $|(\mathcal{R}_{\{I_i\}, K \cap I} \setminus K) \cap J_j| \leq 1$, $J_j \cap K = \emptyset$,
- $|(\mathcal{R}_{\{I_i\}, K \cap I} \setminus K) \cap I_i| \leq 1$, $I_i \cap K = \emptyset$.

Then, we define the quotient of $\Gamma_{\sigma, \{I_i\}, \{J_j\}}$ by Γ_K to be a triple

$$(\sigma|_{(I \cup J) \setminus K}, \{I_i\} / \{I_i\}_{(K)}, \{J_j\} / \{J_j\}_{(K)}),$$

denoted by $\Gamma_{\sigma, \{I_i\}, \{J_j\}} / \Gamma_K$ also.

Definition 4.11. Let $\Gamma_{\{I_i\}, \{J_j\}}$ and $\Gamma_{\{L_l\}, \{K_k\}}$ be two connected admissible graphs with structure maps σ and λ ($(I \cup J) \cap (L \cup K) = \emptyset$). Let $1 \leq a \leq |\{I_i\}|$, $1 \leq b \leq |\{J_j\}|$, $\iota : I_a \rightarrow \{L_l\}$, $\kappa : J_b \rightarrow \{K_k\}$, we define the insertion of $\Gamma_{\{L_l\}, \{K_k\}}$ into $\Gamma_{\{I_i\}, \{J_j\}}$ to be the triple

$$(\delta, \{I_i\} \circ'_{I_a} \{L_l\}, \{J_j\} \circ^\kappa_{J_b} \{K_k\}),$$

where δ satisfies $\delta|_{I \cup J} = \sigma$, $\delta|_{L \cup K} = \lambda$, $I = \mathcal{R}(\{I_i\})$, $J = \mathcal{R}(\{J_j\})$, $L = \mathcal{R}(\{L_l\})$, $K = \mathcal{R}(\{K_k\})$. The graph $(\delta, \{I_i\} \circ'_{I_a} \{L_l\}, \{J_j\} \circ^\kappa_{J_b} \{K_k\})$ is also denoted by $\Gamma_{\sigma, \{I_i\}, \{J_j\}} \circ_{I_a, J_b}^{\iota, \kappa} \Gamma_{\lambda, \{L_l\}, \{K_k\}}$.

Similar to the situations in subsection 4.1 and 4.2, it is obvious that the following conclusions are valid.

Proposition 4.9. • *Let admissible graph $\Gamma_{\sigma, \{I_i\}, \{J_j\}}$ and its subgraph Γ_K are connected, where $K \subset (I \cup J)$ is a σ -invariant subset. Then the quotient $\Gamma_{\sigma, \{I_i\}, \{J_j\}} / \Gamma_K$ is connected also.*

- *Let $\Gamma_{\{I_i\}, \{J_j\}}$ and $\Gamma_{\{L_l\}, \{K_k\}}$ be two connected admissible graphs. Then $(\delta, \{I_i\} \circ_{I_a}^{\iota} \{L_l\}, \{J_j\} \circ_{J_b}^{\kappa} \{K_k\})$ is a connected admissible graph.*

Let $(\sigma, \{I_i\}, \{J_j\})$ and $(\lambda, \{L_l\}, \{K_k\})$ be two connected admissible graphs. Then, it is natural for us to define the composition of the admissible graphs to be

$$\begin{aligned} & (\sigma, \{I_i\}, \{J_j\}) \circ (\lambda, \{L_l\}, \{K_k\}) \\ &= \sum_{a, b, \iota, \kappa} (\delta, \{I_i\} \circ_{I_a}^{\iota} \{L_l\}, \{J_j\} \circ_{J_b}^{\kappa} \{K_k\}). \end{aligned} \quad (4.6)$$

Now we define the Lie bracket to be

$$\begin{aligned} & [(\sigma, \{I_i\}, \{J_j\}), (\lambda, \{L_l\}, \{K_k\})] \\ &= (\sigma, \{I_i\}, \{J_j\}) \circ (\lambda, \{L_l\}, \{K_k\}) - (\lambda, \{L_l\}, \{K_k\}) \circ (\sigma, \{I_i\}, \{J_j\}). \end{aligned} \quad (4.7)$$

Observing the definition 4.11 and the formula (4.6), the structure map δ on $(\delta, \{I_i\} \circ_{I_a}^{\iota} \{L_l\}, \{J_j\} \circ_{J_b}^{\kappa} \{K_k\})$ is independent of the choices of a, b, ι, κ , even the order of the insertion, thus, all admissible graphs concerning (4.7) adapt same structure map. Above fact implies that the Lie bracket (4.7) is similar to the cartesian product of Lie algebras, which means that the Lie bracket (4.7) will satisfy the Jacobi identity.

References

- [1] A. Connes, D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem. *Comm. Math. Phys.* 210 (2000), no. 1, 249–273.
- [2] A. Connes, D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. II. The β -function, diffeomorphisms and the renormalization group. *Comm. Math. Phys.* 216 (2001), no. 1, 215–241.
- [3] A. Connes, M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives*. American Mathematical Society, 2007-12-20.
- [4] L. M. Ionescu, *A combinatorial approach to coefficients in deformation quantization*, [arXiv:math/0404389v1].
- [5] Jean-Louis Loday and N. M. Nikolov, Operadic construction of the renormalization group. [arXiv:math-ph.1202.1206v1]
- [6] D. Kreimer, On the Hopf algebra structure of perturbative quantum field theories. *Adv. Theor. Math. Phys.* 2 (1998), no. 2, 303–334.
- [7] M. Kontsevich, *Deformation quantization of Poisson manifolds*, *Letters Math. Phys.* 66 (2003), no.3, 157–216, [arXiv:q-alg/9709040].
- [8] D. E. Radford, *Hopf Algebras*, World Scientific Publishing Co. Pte. Ltd. 2012.

- [9] ZhouMai, *Hopf Algebras Concerning Matrices or Finite Sets and Their Application to Star Product of Scalar fields*, [arXiv:math-ph/2006.11786v3]