

Massive Integrability: From Fishnet Theories to Feynman Graphs and Back

Florian Loebbert* and Julian Miczajka

*Institut für Physik, Humboldt-Universität zu Berlin,
Zum Großen Windkanal 6, 12489 Berlin, Germany*

E-mail: {loebbert,miczajka}@physik.hu-berlin.de

An overview of the massive generalization of Yangian symmetry for Feynman integrals is given. We illustrate the relation to a massive fishnet theory defined as a double-scaling limit of Coulomb-branch $\mathcal{N} = 4$ SYM theory.

*** *The European Physical Society Conference on High Energy Physics (EPS-HEP2021), ****

*** *26-30 July 2021 ****

*** *Online conference, jointly organized by Universität Hamburg and the research center DESY ****

*Speaker

1. Introduction

In the context of the AdS/CFT correspondence, integrability has proven to be a useful tool. Recently, it was found that large classes of Feynman integrals exhibit a conformal Yangian symmetry. Via their interpretation as correlation functions in the so-called fishnet theories, these integrable structures trace back to the integrability of massless planar $\mathcal{N}=4$ super Yang-Mills (SYM) theory [1–4]. In the following, we review the generalization to the massive case [5–7], which corresponds to the first findings of integrable structures in massive quantum field theory in four dimensions.

2. Yangian Symmetry and Integrability in AdS/CFT

The Yangian algebra introduced by Drinfel'd constitutes an infinite dimensional extension of a Lie algebra \mathfrak{g} . It underlies quantum integrable models with a rational R-matrix and plays an important role for the AdS₅/CFT₄ correspondence. In its so-called first realization, the whole Yangian $Y[\mathfrak{g}]$ is generated by two sets of generators. The level-zero generators of the Lie algebra \mathfrak{g} have a trivial tensor product action of the form $J^a = \sum_{k=1}^n J_k^a \in \mathfrak{g}$ and obey the standard Lie algebra commutation relations $[J^a, J^b] = f^{ab}{}_c J^c$. In addition, the Yangian is generated by a set of level-one generators with a non-trivial coproduct:

$$\widehat{J}^a = f^a{}_{bc} \sum_{j < k=1}^n J_j^c J_k^b + \sum_{j=1}^n s_j J_j^a, \quad [J^a, \widehat{J}^b] = f^{ab}{}_c \widehat{J}^c. \quad (1)$$

Moreover, the so-called Serre relations have to be obeyed, cf. Ref. [8] for the massive representation discussed below. In the case of $\mathcal{N} = 4$ SYM theory, the underlying Lie algebra is $\mathfrak{g} = \mathfrak{psu}(2, 2|4)$. In the AdS/CFT context the Yangian can be understood as the closure of an ordinary conformal symmetry and a dual conformal symmetry [9].

3. From $\mathcal{N} = 4$ SYM to Massless Feynman Graphs

A generalization of $\mathcal{N} = 4$ SYM theory is the so-called γ -deformation which is defined by introducing certain phase factors in the Lagrangian. These phases depend on three parameters γ_j for $j = 1, 2, 3$, which correspond to the three Cartan charges of the SU(4) R-symmetry. These parameters together with the coupling constant g allow to take very interesting double-scaling limits of the γ -deformed theory [1]. In the most restrictive limit with $g \rightarrow 0$ and $\gamma_3 \rightarrow i\infty$ one obtains a simple Lagrangian for two complex scalars X and Z (here we use Euclidean signature):


$$\mathcal{L}_F = N_c \operatorname{tr} \left(-\partial_\mu \bar{X} \partial^\mu X - \partial_\mu \bar{Z} \partial^\mu Z + \xi^2 \bar{X} \bar{Z} X Z \right). \quad (2)$$

The new coupling $\xi := g e^{-i\gamma_3/2}$ is kept fix in the limit. Schematically, the route to the so-called *bi-scalar fishnet theory* defined by \mathcal{L}_F takes the form

$$\mathcal{N} = 4 \text{ SYM} \xrightarrow{XY \rightarrow e^{i\gamma_j(\dots)} XY} \gamma\text{-Deformation} \xrightarrow[\xi = g e^{-i\gamma_3/2} \text{ fix}]{g \rightarrow 0, \gamma_3 \rightarrow i\infty} \text{Fishnet Theory} . \quad (3)$$

Notably, the theory defined by (2) is non-unitary due to the chiral four-point interaction, but it has the remarkable feature that correlation functions are in one-to-one correspondence with individual

Feynman graphs of fishnet structure. In particular, these Feynman integrals (alias correlators) inherit a bosonic Yangian symmetry, i.e. the respective integrals are invariant under the level-one differential operators of the Yangian algebra $Y[\mathfrak{so}(1,5)]$ [2, 3]:



$$\widehat{J}^a = 0. \quad (4)$$

Next we discuss how the above integrability properties of massless Feynman integrals can be generalized to the massive situation.

4. Massive Yangian Symmetry of Feynman Graphs

In the massless case, $\mathcal{N} = 4$ SYM theory was the starting point to deduce the integrability of massless Feynman graphs via the sequence (3). Hence, it is suggestive to think about introducing masses into $\mathcal{N} = 4$ SYM in order to follow a similar path. Masses are easily generated by giving a vacuum expectation value (VEV) to one of the scalar fields of $\mathcal{N} = 4$ SYM theory: $\Phi \rightarrow \langle \Phi \rangle + \Phi$. This leads to massive propagators with difference mass, i.e.


$$\frac{1}{x_{jk}^2} \rightarrow \frac{1}{\hat{x}_{jk}^2} := \frac{1}{x_{jk}^2 + (m_j - m_k)^2}. \quad (5)$$

Here we use the notation $x_{jk}^\mu = x_j^\mu - x_k^\mu$ with $\mu = 1, \dots, D$. The mass can be interpreted as a fifth or $(D+1)$ th dimension of the spacetime vector $x^{\hat{\mu}}$ with $\hat{\mu} = 1, \dots, D+1$. For $D = 4$ it plays the role of the radial direction in AdS_5 , i.e. $x^{\hat{\mu}=D+1} = m_j$. There is a well known massive version of dual conformal symmetry in this massive phase, i.e. on the Coulomb branch of $\mathcal{N} = 4$ SYM theory, which is generated by the following differential operators [10]:

$$\begin{aligned} P_j^\mu &= -i\partial_{x_j}^\mu, & L_j^{\mu\nu} &= ix_j^\mu \partial_{x_j}^\nu - ix_j^\nu \partial_{x_j}^\mu, & D_j &= -i(x_{j\mu} \partial_{x_j}^\mu + m_j \partial_{m_j} + \Delta_j), \\ K_j^\mu &= -2ix_j^\mu (x_{j\nu} \partial_{x_j}^\nu + m_j \partial_{m_j} + \Delta_j) + i(x_j^2 + m_j^2) \partial_{x_j}^\mu. \end{aligned} \quad (6)$$

Importantly, however, no massive Yangian symmetry (integrability) is known on the Coulomb branch, i.e. the naive logic to deduce integrability properties of Feynman integrals via the sequence of steps (3) to the massive situation fails. We will come back to this approach in Section 6.

An alternative route is to consider massive Feynman integrals directly, e.g. a diagram of the form



$$= \int \frac{d^D x_0 d^D x_{\bar{0}}}{\hat{x}_{01}^{2a_1} \hat{x}_{02}^{2a_2} x_{0\bar{0}}^{2b_0} \hat{x}_{\bar{0}3}^{2a_3} \hat{x}_{\bar{0}4}^{2a_4}}.$$

Here the region momenta x (black graph) can be related to ordinary momenta (green graph) via $p_j^\mu = x_j^\mu - x_{j+1}^\mu$. Using their defining equation (1), it is straightforward to build Yangian level-one generators composed of densities of the above massive dual conformal (level-zero) generators J^a as given in (6). Here the level-one momentum generator for instance takes the form

$$\widehat{P}^\mu = \frac{i}{2} \sum_{j,k=1}^n \text{sign}(k-j) (P_j^\mu D_k + P_{j\nu} L_k^{\nu\mu}) + \sum_{j=1}^n s_j P_j^\mu. \quad (7)$$

n = generic: ‘experiments’ yield natural conjecture [7]

$$I_n = \frac{\pi^{D/2} \Gamma_{D/2}}{\Gamma_D \prod_{j=1}^n m_j^{a_j}} \sum_{k_{ij}=0}^{\infty} \frac{\prod_{j=1}^n (a_j)_{\sum_{\alpha \in B_{n|j}} k_{\alpha}}}{(\frac{D+1}{2})_{\sum_{\alpha \in B_n} k_{\alpha}}} \prod_{\alpha \in B_n} \frac{u_{\alpha}^{k_{\alpha}}}{k_{\alpha}!}. \quad (10)$$

This n -point conjecture was recently confirmed in Ref. [12].

6. From Feynman Graphs to Massive Fishnet Theory

Having motivated the Yangian symmetry of massless Feynman integrals via the relation to a massless fishnet theory, an obvious question is whether the Yangian symmetry of massive Feynman integrals has a similar origin. In other words: Can we define a massive version of the fishnet theory? The simplest idea would be to take the massless fishnet theory and to introduce mass via spontaneous symmetry breaking. This line of thought was explored in Ref. [6, 13] with the result that masses enter into the propagators of the theory as products, i.e. in the form $p^2 + m_j m_k$. Hence, the resulting Feynman graphs do not correspond to the Yangian-invariant integrals with difference-mass propagators (5). An alternative idea is to first introduce mass in $\mathcal{N} = 4$ SYM theory and to then take a double-scaling limit on the resulting Coulomb branch [6]. In fact, this approach yields the desired difference-mass propagators $p^2 + (m_j - m_k)^2$. However, taking a double-scaling limit on the Coulomb branch of $\mathcal{N} = 4$ SYM theory requires some care. In particular, a critical step is the γ -deformation \mathcal{P}_{γ} , which a priori is only well-defined on R-symmetry singlets. A solution, motivated by the resulting limit theory, is to average over all ways to break up traces in Lagrangian:


$$Q : \text{tr}(\Phi_1 \Phi_2 \dots \Phi_n) \mapsto \frac{1}{n} (\Phi_1 \Phi_2 \dots \Phi_n + \Phi_2 \Phi_3 \dots \Phi_1 + \dots). \quad (11)$$

Then we define the γ -deformed Coulomb-branch Lagrangian as $\mathcal{L}_{\text{Coul}}^{\gamma} = Q^{-1} \mathcal{P}_{\gamma} Q \mathcal{L}_{\text{Coul}}$. Importantly, application of $Q^{-1} \mathcal{P}_{\gamma} Q$ is now a well defined mathematical operation even if the R-symmetry of $\mathcal{L}_{\text{Coul}}$ is broken. Following the massless procedure, the most restrictive double-scaling limit then results in a *massive fishnet theory* defined by the following Lagrangian [6]:

$$\begin{aligned} \mathcal{L}_{\text{MF}} = & N_c \text{tr} \left(-\partial_{\mu} \bar{X} \partial^{\mu} X - \partial_{\mu} \bar{Z} \partial^{\mu} Z + \xi^2 \bar{X} \bar{Z} X Z \right) \\ & - N_c (m_a - m_b)^2 X^a{}_b \bar{X}_a^b - N_c (m_a - m_b)^2 Z^a{}_b \bar{Z}_a^b. \end{aligned} \quad (12)$$

Planar amplitudes in the respective theory are in one-to-one correspondence with Yangian-invariant massive Feynman integrals. As sketched in Section 4, this invariance is proven at one and two loops and conjectural at higher loop orders. In this sense the above massive fishnet theory is integrable.

The relations between the different theories are summarized in the following figure:



This diagram with three integrable corners (black) leads to a natural question: Is planar $\mathcal{N}=4$ SYM theory on the Coulomb branch integrable?

Acknowledgments

We are grateful to Dennis Müller and Hagen Münkler for collaborations on the main topics discussed in this contribution. The work of FL is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)–Projektnummer 363895012. JM is supported by the International Max Planck Research School for Mathematical and Physical Aspects of Gravitation, Cosmology and Quantum Field Theory.

References

- [1] O. Gürdoğan and V. Kazakov, *New Integrable 4D Quantum Field Theories from Strongly Deformed Planar $N = 4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **117** (2016) 201602 [[1512.06704](#)].
- [2] D. Chicherin, V. Kazakov, F. Loebbert, D. Müller and D.-l. Zhong, *Yangian Symmetry for Bi-Scalar Loop Amplitudes*, *JHEP* **05** (2018) 003 [[1704.01967](#)].
- [3] D. Chicherin, V. Kazakov, F. Loebbert, D. Müller and D.-l. Zhong, *Yangian Symmetry for Fishnet Feynman Graphs*, *Phys. Rev. D* **96** (2017) 121901 [[1708.00007](#)].
- [4] F. Loebbert, D. Müller and H. Münkler, *Yangian Bootstrap for Conformal Feynman Integrals*, *Phys. Rev. D* **101** (2020) 066006 [[1912.05561](#)].
- [5] F. Loebbert, J. Miczajka, D. Müller and H. Münkler, *Massive Conformal Symmetry and Integrability for Feynman Integrals*, *Phys. Rev. Lett.* **125** (2020) 091602 [[2005.01735](#)].
- [6] F. Loebbert and J. Miczajka, *Massive Fishnets*, *JHEP* **12** (2020) 197 [[2008.11739](#)].
- [7] F. Loebbert, J. Miczajka, D. Müller and H. Münkler, *Yangian Bootstrap for Massive Feynman Integrals*, *SciPost Phys.* **11** (2021) 010 [[2010.08552](#)].
- [8] J. Miczajka, *The Yangian Bootstrap for Massive Feynman Diagrams*, *PhD thesis* (2021) .
- [9] J.M. Drummond, J.M. Henn and J. Plefka, *Yangian symmetry of scattering amplitudes in $N=4$ super Yang-Mills theory*, *JHEP* **05** (2009) 046 [[0902.2987](#)].
- [10] L.F. Alday, J.M. Henn, J. Plefka and T. Schuster, *Scattering into the fifth dimension of $N=4$ super Yang-Mills*, *JHEP* **01** (2010) 077 [[0908.0684](#)].
- [11] L. Corcoran, F. Loebbert, J. Miczajka and M. Staudacher, *Minkowski Box from Yangian Bootstrap*, *JHEP* **04** (2021) 160 [[2012.07852](#)].
- [12] B. Ananthanarayan, S. Banik, S. Friot and S. Ghosh, *Massive One-loop Conformal Feynman Integrals and Quadratic Transformations of Multiple Hypergeometric Series*, *Phys. Rev. D* **103** (2021) 096008 [[2012.15646](#)].
- [13] G.K. Karananas, V. Kazakov and M. Shaposhnikov, *Spontaneous Conformal Symmetry Breaking in Fishnet CFT*, *Phys. Lett. B* **811** (2020) 135922 [[1908.04302](#)].