

A POINCARÉ-BENDIXSON THEOREM FOR FLOWS WITH ARBITRARILY MANY SINGULAR POINTS

TOMOO YOKOYAMA

ABSTRACT. The Poincaré-Bendixson theorem is one of the most fundamental tools to capture the limit behaviors of orbits of flows. It was generalized and was applied to various phenomena in dynamical systems, differential equations, foliations, group actions, translation lines, and semi-dynamical systems. On the other hand, though the no-slip boundary condition is a fundamental condition in differential equations and appears in various fluid phenomena and since Lakes of Wada attractors naturally occur in discrete and continuous real dynamical systems and complex dynamics, no generalizations of the Poincaré-Bendixson theorem can be applied to any differential equations with no-slip boundary condition on surfaces with boundary and flows with Lakes of Wada attractors. To analyze them, we generalize the Poincaré-Bendixson theorem into one for flows with arbitrarily many singular points on possibly non-compact surfaces by using methods of foliation theory and general topology. Moreover, the generalization implies topological characterizations of recurrence, which are generalizations of the Maier's description of recurrence. In addition, we construct a blow-up operation that makes ω -limit sets not arcwise-connected.

1. INTRODUCTION

The Poincaré-Bendixson theorem is one of the most fundamental tool to capture the limit behaviors of orbits of flows and was applied to various phenomena (e.g. [8, 18, 24, 28, 44, 46–48]). The theorem is stated by Poincaré in [43] for analytic vector fields on the plane and proved by Bendixson in [7] for C^1 vector fields on the plane. In [9], Birkhoff introduced the concepts of ω -limit set and α -limit set of a point. Using these concepts, one can simply describe the limit behaviors of orbits stated in the works of Poincaré and Bendixson in detail. In fact, the Poincaré-Bendixson theorem was generalized for flows on surfaces in various ways [3, 5, 15, 17, 20, 21, 23, 29, 31, 32, 34, 36, 37, 41, 49, 52, 54], and also for foliations [30, 42], translation lines on the sphere [28], geodesics for a meromorphic connection on Riemann surfaces [1, 2], group actions [27], and semidynamical systems [10]. For instance,

Date: September 15, 2022.

2020 Mathematics Subject Classification. 37E35, 37B20, 37B35.

Key words and phrases. Poincaré-Bendixson theorem, ω -limit sets, flows on surfaces, flow boxes.

This work was partially supported by JSPS Kakenhi Grant Number 20K03583.

the following statement holds (cf. [41]): The ω -limit set of an orbit of a flow with finitely many fixed points on a compact surface is either a closed orbit, a semi-attracting limit circuit, or a Q-set, where a Q-set is the closure of non-closed recurrent orbit. A part of this classification is based on the following Maier's result [39] (see Theorem 2.4.4 p.32 [41], Theorem 4.2 [4] for general cases, and Theorem 19 [33] for orientable hyperbolic cases for details): Any point contained in an ω -limit set of some point for a flow on a compact surface whose ω -limit set contains non-closed orbits is positively recurrent. Furthermore, in [32, 53], it is shown that the ω -limit set of a non-recurrent point is the boundary of an open annulus. Moreover, in [36], the Poincaré-Bendixson theorem is generalized for flows with the totally disconnected singular point sets on closed orientable surfaces as follows: The ω -limit set of an orbit of such a flow is either a closed orbit, a union of singular points and of connecting separatrices, or a Q-set. Here a non-singular orbit is a connecting separatrix if each of the ω -limit and the α -limit sets is a singular point. However, any differential equations with no-slip boundary condition on surfaces with boundary can be applied no such generalizations of the Poincaré-Bendixson theorem to, because of the degeneracy of singular points.

On the other hand, the ω -limit sets of orbits of analytic flows on the plane, the sphere, and the projective plane are studied in [31]. However, Buendía and López pointed out a gap in a crucial lemma in the work by constructing a counterexample in the case of the sphere minus two points [19]. Though the lemma is not generally true, they fixed the gap in the case of the sphere, the plane, the projective plane, and the projective plane minus one point. They also showed that ω -limit sets of orbits of analytic flows on open connected subsets of the sphere are essentially the boundaries of simply connected Peano subcontinua [20],

1.1. Statements of main results. To describe the Poincaré-Bendixson theorem for flows with arbitrarily many singular points on surfaces, we recall some concepts and introduce two concepts (quasi-circuit and quasi-Q-set) as follows. A closed connected invariant subset is a **non-trivial quasi-circuit** if it is a boundary component of an open annulus, contains a non-recurrent point, and consists of non-recurrent points and singular points. A non-trivial quasi-circuit γ is a **quasi-semi-attracting limit quasi-circuit** with respect to a small collar \mathbb{A} if there is a point $x \in \mathbb{A}$ with $O^+(x) \subset \mathbb{A}$ such that $\omega(x) = \gamma$. A subset is **locally dense** if its closure has a nonempty interior. A **Q-set** is the closure of a non-closed recurrent orbit. A Q-set is a **transversely Cantor Q-set** if there is a small neighborhood U of a recurrent point of the Q-set \mathcal{M} such that $\mathcal{M} \cap U$ is a product of an open interval and a Cantor set. An ω -limit set of a point is a **quasi-Q-set** if it intersects an essential closed transversal infinitely many times. A non-recurrent orbit is **connecting quasi-separatrix** if each of the ω -limit and the α -limit sets is contained in a boundary component of the singular point

set. A subset is **locally connected** if each point in it has a small open connected neighborhood.

We generalize the Poincaré-Bendixson theorem for flows with the totally disconnected singular point set on a closed orientable surface (see [36, Theorem 3.1] (cf. [38, Theorem 2.1])) as follows.

Theorem A. The following statements hold for a flow with arbitrarily many singular points on a compact surface:

- (a) The ω -limit set of any non-closed orbit is one of the following exclusively:
 - (1) A nowhere dense subset of singular points.
 - (2) A semi-attracting limit cycle.
 - (3) A quasi-semi-attracting limit quasi-circuit that is the image of a circle.
 - (4) A quasi-semi-attracting limit quasi-circuit that is not locally connected.
 - (5) A locally dense Q-set.
 - (6) A transversely Cantor Q-set.
 - (7) A quasi-Q-set that consists of singular points and non-recurrent points.
- (b) Every non-recurrent orbit in the ω -limit set of a point is a connecting quasi-separatrix.
- (c) If the singular point set is totally disconnected, then any non-recurrent orbits in the ω -limit set of a point are connecting separatrices.

The previous theorem can be applied to capture a time-reversal symmetric condition for limit sets. In fact, the characterization of a flow that is either irrational or Denjoy by Athanassopoulos [6] is refined as an application of the previous theorem in [59]. From the construction in [59, Example 3], notice that a subsets of singular points can become Lakes of Wada attractors (or Plykin type attractors) and that quasi-semi-attracting limit quasi-circuits that is not locally connected can contain a Wada-Lakes-like structure. By this construction, cutting closed transversals and collapsing the new boundary components, we can construct flows on a sphere with Lakes of Wada attractors and with an arbitrarily large number of complementary domains, which are flow versions of such attractors of spherical homeomorphisms constructed by Boroński, Činč, and Liu [11] and such an attractor of a transcendental entire function constructed by Martí-Pete, Rempe and Watterman [35]. In particular, such constructions of homeomorphisms are motivated by a generalization of the Poincaré-Bendixson theorem [28]. Moreover, using the previous theorem, for any Hamiltonian flow with arbitrarily many singular points on a compact surface, it is shown that the ω -limit set of any non-closed orbit consists of singular points [58]. In addition, the previous theorem implies a generalization of the Maier's description of recurrence as follows.

Theorem B. Let v be a flow on a compact surface S . The following statements hold for a point $x \in \omega(z)$ for some point $z \in S$:

- (1) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed positively recurrent.
- (2) $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed negatively recurrent.
- (3) $(\omega(x) \cup \alpha(x)) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed recurrent.
- (4) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed Poisson stable.

The previous theorem implies the following topological characterizations of non-closed positive recurrence.

Corollary C. The following are equivalent for a point x for a flow on a compact surface S :

- (1) The point x is non-closed positively recurrent.
- (2) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and there is a point $z \in S$ with $x \in \omega(z)$.
- (3) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and there is a point $z \in S$ with $x \in \alpha(z)$.

Using the end completion of surfaces of finite genus and finitely many boundary components, we can obtain analogous results for such surfaces (see Theorem 4.1, Theorem 4.2, and Corollary 4.3 for details). Furthermore, applying Theorem 4.2, minimal flows on compact surfaces are characterized and the Poincaré recurrence theorem for flows on surfaces is generalized [57].

We also introduce the following blow-up operation, which can modify any limit circuits into quasi-circuits that are not circuits (and more generally modify any non-locally-dense ω -limit sets into ω -limit sets which are not arcwise-connected).

Theorem D. Let v be a flow on a surface S with an ω -limit set ω containing non-singular point x and with a point $y \in S - \omega$ satisfying $\omega(y) = \omega$. Then there is a flow \tilde{v} on S satisfying the following properties:

- (1) The ω -limit set $\omega_{\tilde{v}}(y)$ is not arcwise-connected unless it is locally dense.
- (2) The restriction $v|_{S-\omega}$ of v is topologically equivalent to the restriction $\tilde{v}|_{S-\omega_{\tilde{v}}(y)}$.
- (3) The flow v_x is topologically equivalent to a flow which is topologically semi-conjugate to \tilde{v} , where v_x is the resulting flow of v by replacing x with a singular point (see Lemma 6.1 for details of the definition of v_x).
- (4) The topological conjugacy from \tilde{v} to v_x can be obtained by collapsing a connected closed invariant subset of $\omega_{\tilde{v}}(y)$ into a singleton.

The present paper consists of seven sections. In the next section, as preliminaries, we introduce fundamental concepts. In §3, we generalize the Poincaré-Bendixson theorem to one for a flow with arbitrarily many singular points on a compact surface. In §4, the Poincaré-Bendixson theorem is generalized to one for a flow with arbitrarily many singular points on a surface of finite genus and finitely many boundary components. Moreover, we characterize the recurrence, which are generalizations of the Maier's description of recurrence. In §5, the finiteness of singular points implies that a quasi-Q-set is a Q-set and that a limit quasi-circuit is a limit circuit. These

reductions imply a proof of a generalization of the Poincaré-Bendixson theorem for a flow with finitely many singular points on a compact surface. In §6, we introduce an operation that makes ω -limit sets not arcwise-connected by constructing flow boxes with non-arcwise-connected invariant subsets. In the final section, we state some examples with ω -limit sets which are non-locally-connected subsets of singular points, non-circuits, and non-Q-sets respectively.

2. PRELIMINARIES

2.1. Topological notion. Denote by \overline{A} the closure of a subset A of a topological space, by $\text{int } A$ the interior of A , and by $\partial A := \overline{A} - \text{int } A$ the boundary of A , where $B - C$ is used instead of the set difference $B \setminus C$ when $B \subseteq C$. We define the **coborder** $\partial^+ A$ of A by $\overline{A} - A$ and the **border** $\partial^- A$ by $A - \text{int } A$ of A . Then $\partial A = \partial^- A \sqcup \partial^+ A$, where \sqcup denotes a disjoint union. A boundary component of a subset A is a connected component of the boundary of A . A subset is **locally dense** if its closure has a nonempty interior. A subset of a compact surface S is **essential** if it is not null homotopic in S^* , where S^* is the resulting closed surface from S by collapsing all boundary components into singletons.

2.1.1. Curves and loops. Recall that a curve is a continuous mapping $C : I \rightarrow Y$ where I is a non-degenerate connected subset of a circle \mathbb{S}^1 . A curve is simple if it is injective. We also denote by C the image of a curve C . Denote by $\partial C := C(\partial I)$ the boundary of a curve C , where ∂I is the boundary of $I \subset \mathbb{S}^1$. Put $\text{int } C := C \setminus \partial C$. A simple curve is a simple closed curve if its domain is \mathbb{S}^1 (i.e. $I = \mathbb{S}^1$). A simple closed curve is also called a **loop**. An **arc** is a simple curve whose domain is an interval.

2.2. Notion of dynamical systems. By a surface, we mean a paracompact two dimensional manifold, that does not need to be orientable. A flow is a continuous \mathbb{R} -action on a manifold. From now on, we suppose that flows are on surfaces unless otherwise stated. Let $v : \mathbb{R} \times S \rightarrow S$ be a flow on a surface S . For $t \in \mathbb{R}$, define $v_t : S \rightarrow S$ by $v_t := v(t, \cdot)$. For a point x of S , we denote by $O(x)$ the orbit of x , $O^+(x)$ the positive orbit (i.e. $O^+(x) := \{v_t(x) \mid t > 0\}$), and $O^-(x)$ the negative orbit (i.e. $O^-(x) := \{v_t(x) \mid t < 0\}$). Recall that a point x of S is **singular** if $x = v_t(x)$ for any $t \in \mathbb{R}$ and is **periodic** if there is a positive number $T > 0$ such that $x = v_T(x)$ and $x \neq v_t(x)$ for any $t \in (0, T)$. A point is **closed** if it is singular or periodic. An orbit is singular (resp. periodic, closed) if it contains a singular (resp. periodic, closed) point. Denote by $\text{Sing}(v)$ the set of singular points and by $\text{Per}(v)$ (resp. $\text{Cl}(v)$) the union of periodic (resp. closed) orbits. The subset $\text{Sing}(v)$ is called the singular point set. Recall that the ω -limit (resp. α -limit) set of a point x is $\omega(x) := \bigcap_{n \in \mathbb{R}} \overline{\{v_t(x) \mid t > n\}}$ (resp. $\alpha(x) := \bigcap_{n \in \mathbb{R}} \overline{\{v_t(x) \mid t < -n\}}$). For an orbit O , define $\omega(O) := \omega(x)$ and

$\alpha(O) := \alpha(x)$ for some point $x \in O$. Note that an ω -limit (resp. α -limit) set of an orbit is independent of the choice of point in the orbit.

2.3. Topological equivalence. A flow v on a surface M is **topologically equivalent** to a flow w on a surface N if there is a homeomorphism $h: M \rightarrow N$ such that the images of any orbits of v are orbits of w with preservation of the direction in time. Then the homeomorphism $h: M \rightarrow N$ is called the topological conjugacy.

2.4. Separatrices and orbit arcs. A **separatrix** is a non-singular orbit whose α -limit or ω -limit set is a singular point. A separatrix is **connecting** if each of its ω -limit set and α -limit sets is a singular point. Note that any connecting separatrices are non-recurrent. An arc is an **orbit arc** if it is contained in an orbit. An orbit arc in a subset A is **maximal** in A if it is an orbit arc in A which is maximal with respect to the inclusion order.

2.4.1. Recurrence and invariance. A point x is positively recurrent (or positively Poisson stable) if $x \in \omega(x)$. A point x is negatively recurrent (or negatively Poisson stable) if $x \in \alpha(x)$. A point x is **recurrent** if $x \in \omega(x) \cup \alpha(x)$. A point x of S is **Poisson stable** (or strongly recurrent) if $x \in \omega(x) \cap \alpha(x)$. Denote by $R(v)$ the set of non-closed recurrent points. A point is **wandering** if there are its neighborhood U and a positive number N such that $v_t(U) \cap U = \emptyset$ for any $t > N$. A point is non-wandering if it is not wandering. Note that a recurrent point is non-wandering. An orbit is recurrent (resp. Poisson stable, wandering, non-wandering) if it contains a recurrent (resp. Poisson stable, wandering, non-wandering) point. A non-closed recurrent orbit is also called a non-trivial recurrent orbit.

Definition 1. The closure of a non-closed recurrent orbit is called a **Q-set** (or quasi-minimal set).

A subset is **invariant** (or **saturated**) if it is a union of orbits. The **saturation** of a subset is the union of orbits intersecting it. A nonempty closed invariant subset is **minimal** if it contains no proper nonempty closed invariant subsets. Recall that the **(orbit) class** \hat{O} of an orbit O is the union of orbits each of whose orbit closure corresponds to \overline{O} (i.e. $\hat{O} = \{y \in S \mid \overline{O(y)} = \overline{O}\}$).

2.4.2. Topological properties of orbits. The following properness, local density, and exceptional properties of orbits are analogous concepts of codimension one foliation theory (cf. [13, 26]).

An orbit is **proper** if it is embedded, locally dense if its closure has a nonempty interior, and exceptional if it is neither proper nor locally dense. A point is proper (resp. locally dense) if its orbit is proper (resp. locally dense). Denote by $LD(v)$ (resp. $E(v)$, $P(v)$) the union of locally dense orbits (resp. exceptional orbits, non-closed proper orbits). Note that each exceptional Q-set is a **transversely Cantor set** \mathcal{M} (i.e. there is a small neighborhood U of a recurrent point of the Q-set \mathcal{M} such that $\mathcal{M} \cap U$ is

a product of an open interval and a Cantor set), because a Cantor set is characterized as a compact metrizable perfect totally disconnected space. We have the following observation.

Lemma 2.1. *The following statements hold for a flow v on a paracompact manifold M :*

- (1) *A point of M is non-proper if and only if it is non-closed recurrent.*
- (2) *$M = \text{Cl}(v) \sqcup \text{P}(v) \sqcup \text{R}(v)$.*
- (3) *The union $\text{P}(v)$ is the set of non-recurrent points.*
- (4) *The union $\text{R}(v)$ is the set of non-proper points.*

Proof. By definitions, note that a closed orbit is proper and recurrent. Fix a point $x \in M$. By [56, Corollary 3.4], the orbit $O(x)$ is proper if and only if $O(x) = \hat{O}(x)$. From [14, Theorem VI], the closure of a non-closed recurrent orbit O of a flow on a manifold contains uncountably many non-closed recurrent orbits whose closures are \overline{O} . Therefore the orbit $O(x)$ is non-closed recurrent if and only if $\hat{O}(x)$ consists of uncountably many orbits. Then any non-closed proper orbit is not recurrent.

We claim that the assertion (1) holds. Indeed, suppose that x is non-closed recurrent. Since any non-closed proper orbit is not recurrent, the point x is not proper. Conversely, suppose that x is not proper. Then x is not closed and there is a point $y \in M$ whose orbit does not contain x such that $\overline{O(y)} = \overline{O(x)}$. This implies that $y \in \overline{O(x)} - O(x) \subseteq \alpha(x) \cup \omega(x)$. From the closed invariance of α -limit sets and ω -limit sets, we have $x \in \overline{O(y)} \subset \alpha(x) \cup \omega(x)$. This means that x is recurrent.

Since $\text{Cl}(v)$ is both the set of closed proper points and the set of closed recurrent points, the assertion (1) implies the assertions (2)–(4). \square

For a flow v on a surface S , the previous lemma implies that the union $\text{R}(v) = \text{LD}(v) \sqcup \text{E}(v)$ consists of non-proper orbits and so that $S = \text{Sing}(v) \sqcup \text{Per}(v) \sqcup \text{P}(v) \sqcup \text{LD}(v) \sqcup \text{E}(v)$.

2.4.3. Circuits. An annular subset is homeomorphic to an annulus. An open annular subset \mathbb{A} of a surface is a **collar** of a singular point x if the union $\mathbb{A} \sqcup \{x\}$ is a neighborhood of x . By a **cycle** or a periodic circuit, we mean a periodic orbit. By a **non-trivial circuit**, we mean either a cycle or an image of an oriented circle by a continuous orientation-preserving mapping which is a directed graph but not a singleton and which is the union of separatrices and finitely many singular points. A **trivial circuit** is a singular point.

Definition 2. A **circuit** is either a trivial or non-trivial circuit.

Note that there are non-trivial circuits with infinitely many edges and that any non-trivial non-periodic circuit contains non-recurrent orbits as in Figure 1.




FIGURE 1. A circuit that consists of a degenerate singular point and infinitely many connecting separatrices, and its neighborhood which consists of a singular point, periodic orbits, and non-recurrent orbits.

Definition 3. An open annular subset \mathbb{A} of a surface is a **collar** of a non-trivial circuit γ if γ is a boundary component of \mathbb{A} and there is a neighborhood U of γ such that \mathbb{A} is a connected component of the complement $U - \gamma$.

A circuit γ is a **semi-attracting** (resp. **semi-repelling**) circuit with respect to a small collar \mathbb{A} if $\omega(x) = \gamma$ (resp. $\alpha(x) = \gamma$) and $O^+(x) \subset \mathbb{A}$ (resp. $O^-(x) \subset \mathbb{A}$) for any point $x \in \mathbb{A}$. Then \mathbb{A} is called a semi-attracting (resp. a semi-repelling) collar basin of γ .

Definition 4. A non-trivial circuit γ is a **limit circuit** if it is a semi-attracting or semi-repelling circuit.

2.4.4. *Quasi-circuits.* A **trivial quasi-circuit** is a cycle or a singular point. Recall the non-trivial quasi-circuit as follows.

Definition 5. A closed connected invariant subset is a **non-trivial quasi-circuit** if it is a boundary component of an open annulus, contains a non-recurrent orbit, and consists of non-recurrent orbits and singular points.

Then the open annulus is called a **collar** of the quasi-circuit. A **quasi-circuit** is either a trivial or non-trivial quasi-circuit.

Definition 6. A quasi-circuit γ is a **quasi-semi-attracting** (resp. **quasi-semi-repelling**) quasi-circuit with respect to a small collar \mathbb{A} if there is a point $x \in \mathbb{A}$ with $O^+(x) \subset \mathbb{A}$ (resp. $O^-(x) \subset \mathbb{A}$) such that $\omega(x) = \gamma$ (resp. $\alpha(x) = \gamma$).

Then \mathbb{A} is called a quasi-semi-attracting (resp. a quasi-semi-repelling) collar basin of γ . We introduce the concept of a limit quasi-circuit, which is a generalized concept of a limit circuit.

Definition 7. A non-trivial quasi-circuit γ is a **limit quasi-circuit** if it is a quasi-semi-attracting or quasi-semi-repelling quasi-circuit (i.e. either $\alpha(x) = \gamma$ or $\omega(x) = \gamma$ for some point $x \notin \gamma$).

Note that a collar basin of a circuit γ is a collar of γ . Moreover, a quasi-circuit is not a circuit in general (see an example in § 7.2).

2.4.5. Transversality. Notice that we can define transversality using tangential spaces of surfaces because each flow on a compact surface is topologically equivalent to a C^1 -flow by Gutierrez's smoothing theorem [22].

A C^1 simple curve C is **transverse** at a point $p \in S$ to the flow v_X generated by a vector field X if $T_p S = T_p C \oplus T_p O_{v_X}(p)$, where $T_p C$ is the tangent space of C at p and $T_p O_{v_X}(p)$ is the tangent space of the orbit of p with respect to the flow v_X .

Definition 8. A simple curve C is **transverse** to v at a point $p \in S$ if there are a vector field X on a surface T and a topological conjugacy $h: S \rightarrow T$ such that v is topologically equivalent to the flow v_X generated by X via h and that the image $h \circ C$ is a C^1 simple curve which is transverse at the point $h(p) \in T$ to the flow v_X .

A simple curve C is **transverse** to v if so is it at any point in C . An arc C transverse to v is called a **transverse arc**.

Definition 9. A simple closed curve is a **closed transversal** (cf. [26, Definition 3.4.7 p.41] and [13, Definition 3.3.6 p.86]) if it transverses to v .

Notice that the closed transversal is one of the fundamental tools in foliation theory to analyze transverse relations among leaves of codimension one foliations (cf. [13, 26]).

2.4.6. Flow boxes. Let I, J be intervals which are either $(-1, 1)$, $(-1, 1]$, $[-1, 1)$, or $[-1, 1]$, $\mathbb{D} := I \times J \subset \mathbb{R}^2$ be a square. Write the annulus $\mathbb{A}_{\mathbb{D}} := \mathbb{D} - [-1/2, 1/2]^2$.

Definition 10. A disk B on a surface S is a **trivial flow box** with respect to a flow v on S if there are a continuous vector field $X = \partial/\partial x_1 = (1, 0)$ on the plane \mathbb{R}^2 , a homeomorphism $f: B \rightarrow \mathbb{D} \subset \mathbb{R}^2$ which carries the maximal orbit arc in B to the maximal orbit arc in \mathbb{D} .

Note that any orbit arcs on the square \mathbb{D} are horizontal intervals and satisfy $\dot{x}_1 = 1$ and $\dot{x}_2 = 0$ with respect to the chart (x_1, x_2) as in the left of Figure 2. In other words, a disk B on a surface S is a trivial flow box with respect to a flow v on S if and only if there are a continuous vector field X on a surface T , a topological conjugacy $h: S \rightarrow T$, and a chart $(x_1, x_2): h(B) \rightarrow \mathbb{D} \subset \mathbb{R}^2$ such that v is topologically equivalent to the flow v_X generated by X via h and that X is given by the expression $\partial/\partial x_1$ in the chart (x_1, x_2) .

Definition 11. A disk B on a surface S is a **flow box** with respect to a flow v on S if there are a closed disk $D \subset \text{int } B$, a continuous vector field $X = \partial/\partial x_1$ on the plane \mathbb{R}^2 , a homeomorphism $f: B - D \rightarrow \mathbb{A} \subset \mathbb{R}^2$ which carries the maximal orbit arc in the annulus $B - D$ to the maximal orbit arc in \mathbb{A} .




FIGURE 2. Left, a trivial flow box; middle, the intersection of a flow box and its small neighborhood of the boundary, right; a Cherry flow box.




FIGURE 3. A flow box with a non-arcwise-connected connected invariant subset.

In other words, a disk B on a surface S is a flow box with respect to a flow v on S if and only if there are a continuous vector field X on a surface T , a topological conjugacy $h: S \rightarrow T$, a closed disk $D \subset \text{int } B$, and a chart $(x_1, x_2): h(B - D) \rightarrow \mathbb{A}_{\mathbb{D}} \subset \mathbb{R}^2$ such that v is topologically equivalent to the flow v_X generated by X via h and that X is given by the expression $\partial/\partial x_1 = (1, 0)$ in the chart (x_1, x_2) .

By definition, trivial flow boxes are flow boxes. The orbit arcs on the annulus $\mathbb{A}_{\mathbb{D}}$ in the flow box \mathbb{D} are horizontal intervals as in the middle of Figure 2. Notice that Cherry flow boxes are flow boxes as in the right of Figure 2.

2.4.7. *Quasi-Q-set.* Recall the quasi-Q-set as follows.

Definition 12. An ω -limit (resp. α -limit) set of a point is a **quasi-Q-set** if it intersects an essential closed transversal infinitely many times.

Note that a quasi-Q-set need not be arcwise-connected. In fact, Hastings constructed an attractor of a flow on \mathbb{R}^2 which is homeomorphic to a Warsaw circle but is not an ω -limit set [25, Example 3.3]. By modifying the construction of a Warsaw circle, one can obtain a quasi-Q-set that is not arcwise-connected by replacing a trivial flow box of a point of an exceptional minimal set with a flow box as in Figure 3 (see details § 6.1). Moreover,

quasi-Q-set is not Q-set in general (see an example in § 7.1). Conversely, we will show that a Q-set is a quasi-Q-set (see Lemma 3.10). On the other hand, if v has at most finitely many singular points, then a quasi-Q-set is a Q-set (see Lemma 5.1). We have the following observation.

Lemma 2.2. *Any locally dense quasi-Q-set is a Q-set. In particular, any quasi-Q-set which intersecting a locally dense orbit is a Q-set.*

Proof. Fix a locally dense quasi-Q-set \mathcal{M} . By definition of quasi-Q-set, there is a point x with $\omega(x) = \mathcal{M}$ or $\alpha(x) = \mathcal{M}$. By time reversion if necessary, we may assume that $\omega(x) = \mathcal{M}$. The locally density implies that $O(x) \cap \text{int } \omega(x) = O(x) \cap \text{int } \mathcal{M} \neq \emptyset$ and so that $x \in \text{int } \omega(x)$. This means that x is non-closed recurrent and $\overline{O(x)} = \omega(x) = \mathcal{M}$. Therefore $\mathcal{M} = \overline{O(x)}$ is a Q-set.

If a quasi-Q-set intersects a locally dense orbit, then the quasi-Q-set is locally dense. \square

3. POINCARÉ-BENDIXSON THEOREM FOR SURFACE CONTINUOUS FLOWS WITH ARBITRARILY MANY SINGULAR POINTS

3.1. A generalization of Poincaré-Bendixson theorem. In this subsection, we have the following key lemma to generalize the Poincaré-Bendixson theorem.

Lemma 3.1. *Let v be a flow on a compact surface S . The ω -limit set of any non-closed orbit O is one of the following exclusively:*

- (1) *A nowhere dense subset of singular points.*
- (2) *A semi-attracting limit cycle.*
- (3) *A quasi-semi-attracting limit quasi-circuit.*
- (4) *A locally dense Q-set.*
- (5) *A quasi-Q-set that is not locally dense.*

Moreover, in the case (3), the orbit O is wandering and is not contained in the orbit closures of points outside of it.

To show the previous lemma, we show some technical lemmas. The proof methods use techniques derived from foliation theory (cf. [13, 26]). First, we show the existence of closed transversals near infinite intersections of transverse arcs and orbits.

Lemma 3.2. *Let I be a transverse arc and $x \in I$ such that $|I \cap O(x)| = \infty$. Then there are an orbit arc C in $O(x)$ and a transverse closed arc $J \subseteq I$ such that the union $\mu := J \cup C$ is a loop with $C \cap J = \partial C = \partial J$ and that the return map along C is orientation-preserving between neighborhoods of ∂C in I . Moreover, there is a closed transversal γ intersecting $O(x)$ near μ .*

Proof. By time reversion if necessary, we may assume that $O^+(x) \cap \text{int } I = \infty$. Fix a point $x_0 \in O^+(x) \cap \text{int } I$. Let $f_v : I \rightarrow I$ be the first return map on I induced by v , $x_i := (f_v)^i(x_0)$ the i -th return of x_0 , $C_{a,b} \subset O^+(x)$ the




FIGURE 4. The waterfall construction

orbit arc from a to b , and $I_{a,b} \subset I$ the subinterval between a and b of I . We may assume that $x_0 < x_1$.

Suppose that the restriction of f_v to a neighborhood of x_i for some $i \in \mathbb{Z}_{\geq 0}$ is orientation-preserving. Then put $C := C_{x_i, x_{i+1}}$ and $J := I_{x_i, x_{i+1}}$. By the waterfall construction (cf. [13, Lemma 3.3.7 p.86]) to the loop $\mu := C \cup J$ (see Figure 4), there is a closed transversal γ intersecting $O(x)$ near μ . Thus we may assume that the restriction of f_v to a neighborhood of x_i for any $i \in \mathbb{Z}_{\geq 0}$ is orientation-reversing.

We claim that there is a natural number i such that $x_{i+1} < x_i$. Indeed, otherwise $x_i < x_{i+1}$ for any $i \in \mathbb{Z}_{\geq 0}$. Then each pair of loops $\gamma_i := C_{x_{2i}, x_{2i+1}} \cup I_{x_{2i+1}, x_{2i}}$ has disjoint neighborhoods each of which is a Möbius band. This means that S has infinitely many non-orientable genus, which contradicts the compactness of S .


By renumbering, we may assume that $x_2 < x_1$. From $x_0 < x_1$, the first return map for I_{x_2, x_0} along C_{x_0, x_2} is orientation-preserving such that a pair of $C := C_{x_0, x_2}$ and $J := I_{x_2, x_0}$. As above, the waterfall construction to the loop $\mu := C \cup J$ completes the assertion. \square

We show that the infinite intersection of a transverse closed arc implies the existence of a quasi-circuit under properness.

Lemma 3.3. *If there is a transverse closed arc $J : [-1, 0] \rightarrow S$ with $\{J(0)\} = J([-1, 0]) \cap \omega(J(-1)) \subset \overline{O^+(J(-1))} \cap J([-1, 0])$, then the following properties hold:*

- (1) *The ω -limit set $\omega(J(-1))$ is a quasi-circuit.*
- (2) *The point $J(-1)$ is wandering and $J(-1) \notin \overline{O(p)}$ for any point $p \in S - O(J(-1))$.*

Proof. Let $J : [-1, 0] \rightarrow S$ be a transverse closed arc, $x_0 := J(-1)$ a point, $y := J(0)$ a point, and $I := J([-1, 0])$ a closed interval with $\{y\} = I \cap \omega(x_0) \subset \overline{O^+(x_0)} \cap I$ as in Figure 6. Since $y \in \overline{O^+(x_0)} \cap I \cap \omega(x_0)$, we obtain that $O^+(x_0) \cap I$ is infinite and $\{y\} = I \cap \omega(x_0) \subset \partial I = \{x_0, y\}$. Denote by x_i the i -th return image of x_0 on I . By $x_i \notin \{J(0)\} = J([-1, 0]) \cap \omega(x_0)$ for any $i \in \mathbb{Z}_{\geq 0}$, the point x_i for any $i \in \mathbb{Z}_{> 0}$ is not positively recurrent and

FIGURE 5. A return map on I .

so is x_0 . Since $\{y\} = I \cap \omega(x_0)$, any closed interval contained in $I - \{y\}$ intersects $O^+(x_0)$ at most finitely many times.

We claim that the sequence $(x_i)_{i \in \mathbb{Z}_{\geq N}}$ in I is strictly increasing and converges to y for some $N > 0$. Indeed, we suffice to show that there is a large number $N > 0$ such that $x_i < x_{i+1}$ in I for any natural number $i \geq N$ with respect to a natural total order on the sub-arc I . Otherwise $x_i > x_{i+1}$ holds for infinitely many natural numbers i . Since each closed sub-arc of $\text{int } I$ intersects at most finitely many points of $O^+(x_0)$, there are infinitely many triples $i_{k-} := i_{k_0} - 1 < i_{k_0} < i_{k_0} + 1 := i_{k+}$ of natural numbers with $i_{k+} < i_{k+1-}$ such that either $x_{i_{k-}} < x_{i_{k+}} < x_{i_{k_0}} < x_i$ or $x_{i_{k+}} < x_{i_{k-}} < x_{i_{k_0}} < x_i$ for any $i \geq i_{k+1-}$ as in Figure 6. Denote by $I_{i_{k-}}$ (resp. $I_{i_{k_0+}}$) by the sub-arc of I whose boundary consists of $x_{i_{k-}}$ and $x_{i_{k_0}}$ (resp. $x_{i_{k_0}}$ and $x_{i_{k+}}$), and by $C_{i_{k-}}$ (resp. $C_{i_{k_0+}}$) the curve contained in $O^+(x_0)$ whose boundary consists of $x_{i_{k-}}$ and $x_{i_{k_0}}$ (resp. $x_{i_{k_0}}$ and $x_{i_{k+}}$). Then the unions $T_{i_{k-}} := C_{i_{k-}} \cup I_{i_{k-}}$ and $T_{i_{k_0+}} := C_{i_{k_0+}} \cup I_{i_{k_0+}}$ are simple closed curves whose intersection is a closed arc $I_{i_{k_0+}}$ or $I_{i_{k-}}$ as on the upper of Figure 6. By a deformation like a waterfall construction as on the lower of Figure 6, we obtain two simple closed curves $T'_{i_{k-}}$ and $T'_{i_{k_0+}}$ whose intersection is either $x_{i_{k-}}$ or $x_{i_{k+}}$ and which are close to the original simple closed curves $T_{i_{k-}}$ and $T_{i_{k_0+}}$ respectively. Note that if S is orientable then we can choose $T'_{i_{k-}}$ and $T'_{i_{k_0+}}$ as closed transversals. Since $\max\{x_{i_{k-}}, x_{i_{k_0}}, x_{i_{k+}}\} < x_i$ for any $i \geq i_{k+1-}$, we have $O^+(x_{i_{k+1-}}) \cap (I_{i_{k-}} \cup I_{i_{k_0+}}) = \emptyset$ and so $O^+(x_{i_{k+1-}}) \cap (T'_{i_{k-}} \cup T'_{i_{k_0+}}) = \emptyset$. Hence $(T'_{i_{k-}} \cup T'_{i_{k_0+}}) \cap (T'_{i_{l-}} \cup T'_{i_{l_0+}}) = \emptyset$ for any $k \neq l$. Since these simple closed curves intersect exactly one point, they are essential. Cutting $T'_{i_{l_0+}}$ and collapsing new boundary components into singletons, we obtain the resulting surface whose genus is the genus of S minus one. Since there are infinitely many disjoint bouquets $T'_{i_{k-}} \cup T'_{i_{k_0+}}$, the genus of S is not finite, which contradicts the compactness of S . Thus there is a large number $N > 0$




FIGURE 6. Upper, Two simple closed curves $T_{i_{k-0}} = C_{i_{k-0}} \cup I_{i_{k-0}}$ and $T_{i_{k0+}} = C_{i_{k0+}} \cup I_{i_{k0+}}$; lower, deformed two transverse simple closed curves $T'_{i_{k-0}}$ and $T'_{i_{k0+}}$.




FIGURE 7. A rectangle whose boundary is the union of $I_i \cup C_i \cup I_{i+1} \cup C_{i+1}$.

such that $x_i < x_{i+1}$ in I for any natural number $i \geq N$. This completes the claim.

For any $i \in \mathbb{Z}_{\geq N}$, denote by I_i by the sub-arc of I whose boundary consists of x_i and x_{i+1} , and by C_i the curve contained in $O^+(x_0)$ whose boundary consists of x_i and x_{i+1} . Let D_i be the open subset bounded by the union $I_i \cup C_i \cup I_{i+1} \cup C_{i+1}$. Since there are at most finitely many genus, by renumbering x_i , we may assume that D_i is a rectangle for any $i \in \mathbb{Z}_{\geq N}$ as in Figure 7. Define a union $\mathbb{A}_k := (\bigcup_{i=k}^{\infty} (D_i \cup I_{i+1} \cup C_{i+1})) - \{x_{k+1}\}$ for any $k \in \mathbb{Z}_{\geq N}$, which is an open annulus. By the monotonicity of x_i in

I , each \mathbb{A}_k is a positive invariant open annulus homotopic to \mathbb{A}_N for any $k \in \mathbb{Z}_{\geq N}$. Since $\omega(x_0)$ is connected, by construction, the closure $\overline{D_i}$ of any rectangle D_i does not intersect $\omega(x_0)$ and so $\mathbb{A}_k \cap \omega(x_0) = \emptyset$. Therefore $\omega(x_0) = \bigcap_{n \in \mathbb{R}_{\geq N}} \overline{\{v_t(x_0) \mid t > n\}} = \bigcap_{k \in \mathbb{Z}_{\geq N}} \overline{\{v_t(x_0) \mid t > k\}} = \bigcap_{i \geq N} \overline{\mathbb{A}_i} = \partial \overline{\mathbb{A}_N} - (I_N \cup C_N)$. This means that $\omega(x_0)$ is a boundary component of the annulus \mathbb{A}_N and so a quasi-circuit.

We claim that x_0 is wandering. Indeed, it suffices to show that x_{N+1} is wandering. By definition, we have $x_{N+1} \in \text{int}(I_N \cup I_{N+1})$ and $\partial I_{N+k} = \{x_{N+k}, x_{N+k+1}\}$ for any $k \in \mathbb{Z}_{\geq 0}$. Let f_v be the first return map on I . Since the point x_k for any $k \in \mathbb{Z}_{\geq 0}$ is contained in the domain of f_v , the flow box theorem (cf. [5, Theorem 1.1, p.45]) implies that there is a small closed interval J whose interior contains x_{N+1} such that J and $f_v(J)$ are contained in the domain of f_v . Then $f_v^2(J)$ is contained in \mathbb{A}_{N+2} . Taking J short, we may assume that there is a positive number $T > 0$ such that $x_{N+3} \in v_T(J) \subset \mathbb{A}_{N+2}$. Then there is a small positive number $\varepsilon > 0$ such that $v_T(\bigsqcup_{t \in (-\varepsilon, \varepsilon)} v_t(J)) \subset \mathbb{A}_{N+2}$. The open subset $U := \bigsqcup_{t \in (-\varepsilon, \varepsilon)} v_t(J) \subset \mathbb{A}_{N-1} - \mathbb{A}_{N+2}$ is an open neighborhood of x_{N+1} and $v_T(U) \subset \mathbb{A}_{N+2}$. By the positive invariance of \mathbb{A}_{N+2} , we have $v_t(U) \subset \mathbb{A}_{N+2}$ and so $U \cap v_t(U) = \emptyset$ for any $t > T$. This means that x_{N+1} is wandering.

Since any wandering point w is not contained in the orbit closure of points outside of $O(w)$, we obtain $x_0 \notin \overline{O(z)}$ for any point $z \in S - O(x_0)$. \square

We show that the infinite intersection of an essential closed transversal implies the existence of either a quasi-Q-set, an essential limit cycle, or a quasi-circuit.

Lemma 3.4. *Let x be a point contained in a closed transversal γ such that $O^+(x)$ intersects γ infinitely many times. Then $\omega(x)$ is either a quasi-Q-set, an essential limit cycle, or a quasi-circuit. Moreover, if $\omega(x)$ is a quasi-circuit, then x is wandering and is not contained in the orbit closures of points outside of $O(x)$.*

Proof. Suppose that a positive orbit $O^+(x)$ intersects a closed transversal γ infinitely many times. Then γ is essential such that $\gamma \cap \omega(x) \neq \emptyset$ and so that $\omega(x) \not\subset \text{Sing}(v)$. If $\omega(x)$ contains periodic orbits, then it is an essential limit cycle because the positive orbit $O^+(x)$ intersects the basin of the semi-attracting limit cycle. Thus we may assume that $\omega(x) \cap \text{Per}(v) = \emptyset$. Suppose that $\omega(x)$ is not a quasi-Q-set. Then $\gamma \cap \omega(x)$ is nonempty and finite.

We claim that there is a transverse closed arc $J : [-1, 0] \rightarrow \gamma$ with $J(-1) \in O^+(x)$ and $\{J(0)\} = J([-1, 0]) \cap \omega(J(-1)) \subset \overline{O^+(J(-1))} \cap J([-1, 0])$. Indeed, fix a point $y \in \gamma \cap \omega(x)$. Since $\gamma \cap \omega(x)$ is finite, there are a point $x_0 \in \gamma \cap O^+(x)$ and a transverse closed arc $J : [-1, 0] \rightarrow \gamma$ with $x_0 = J(-1)$ and $y = J(0)$ such that $\{y\} = J([-1, 0]) \cap \omega(x_0) \subset \overline{O^+(x_0)} \cap J([-1, 0])$.

By Lemma 3.3, the ω -limit set $\omega(x)$ is a quasi-circuit such that x is wandering and is not contained in the orbit closures of points outside of $O(x)$. \square

We show that a limit quasi-circuit is not a quasi-Q-set.

Lemma 3.5. *There are no limit quasi-circuits that are also quasi-Q-sets.*

Proof. Let x be a point whose ω -limit set $\omega(x)$ is a limit quasi-circuit. Since a limit quasi-circuit is a boundary component of an annulus and a closed subset, it has no interior and so is not locally dense. Since $\omega(x)$ is a limit quasi-circuit, there is a small open annulus \mathbb{A} of which $\omega(x)$ is a boundary component such that the two boundary components of \mathbb{A} is disjoint. Denote by ∂_1 another boundary component of \mathbb{A} . In other words, we have $\partial\mathbb{A} = \omega(x) \sqcup \partial_1$. Fix any distance function d on S induced by a Riemannian metric. Since the boundary components of \mathbb{A} are compact and disjoint, there is a positive number $d_{\mathbb{A}}$ such that $d_{\mathbb{A}} = \min\{d(y, z) \mid y \in \omega(x), z \in \partial_1\}$.

We claim that $\omega(x)$ is not a quasi-Q-set. Indeed, assume that $\omega(x)$ is a quasi-Q-set. Then there is a closed transversal γ which intersects $\omega(x)$ infinitely many times. From the flow box theorem applying to the compact subset γ , the fact that γ contains no singular point implies that there is a small number $\delta > 0$ with $\delta < d_{\mathbb{A}}$ such that $B_{\delta}(\gamma) := \{y \in S \mid d(y, \gamma) < \delta\}$ is an open annulus and consists of orbit arcs each of which connects the two boundary components as in Figure 8. Moreover, there is a sequence $(C_i)_{i \in \mathbb{Z}_{\geq 0}}$ of orbit arcs which intersect γ exactly once, are contained in $\omega(x)$, and converge to an orbit arc C_{∞} such that $\partial C_i \cap B_{\delta}(\gamma) = \emptyset$. Put $C'_i := C_i \cap B_{\delta}(\gamma) \subset \omega(x)$ and let $x_i \in C'_i \cap \gamma$ be the intersection point. For any $i \in \mathbb{Z}_{\geq 0}$, let $f_i : [0, l_i] \rightarrow \gamma \subset S$ be the closed arc parameterized by arc length from x_i to either x_{i-1} or x_{i+1} such that $f_i((0, r_i)) \subset \mathbb{A}$ and $f_i(r_i) \in \partial_1$ for some real number $r_i \in (0, l_i)$. By definition, we obtain $f_i(l_i) \in \omega(x)$. Since γ is compact, we have $\lim_{i \rightarrow \infty} l_i = 0$. Fix a large integer N such that $l_N < \delta$. Since $l_N < \delta < d_{\mathbb{A}}$, this implies that any closed arc from x_i whose length is less than δ and which intersects \mathbb{A} but does not intersect C'_i except the starting point is contained in the interior \mathbb{A} . In particular, the closed arc $f_N : [0, l_N] \rightarrow S$ is contained in the interior \mathbb{A} except the starting point x_i (i.e. $f_N((0, l_N]) \subset \mathbb{A}$). Therefore $f_N(l_N) \in \mathbb{A}$, which contradicts $f_N(l_N) \in \omega(x) \subset \partial\mathbb{A} = \mathbb{A} - \mathbb{A}$. \square

We have the following equivalence.

Lemma 3.6. *The following statements are equivalent for a point x :*

- (1) $\omega(x) \cap \text{LD}(v) \neq \emptyset$.
- (2) *The ω -limit set $\omega(x)$ is locally dense.*
- (3) *The ω -limit set $\omega(x)$ is a locally dense Q-set which is not transversely Cantor.*

In any case, we have that $x \in \text{LD}(v)$ and $\overline{O(x)} = \omega(x)$.




FIGURE 8. An example of a closed transversal γ , its annular neighborhood $B_\delta(\gamma)$, and a subset of an open annulus \mathbb{A}

Proof. Trivially, the assertion (3) implies the assertions (1) and (2). Suppose that $\omega(x) \cap \text{LD}(v) \neq \emptyset$. [14, Theorem VI] implies that there is a Poisson stable point $y \in \omega(x) \cap \text{LD}(v)$ such that $\emptyset \neq O(x) \cap \text{int}(\overline{O(y)})$ and so that $x \in \overline{O(y)}$. Since $y \in \omega(x)$, this means that $\omega(x) = \omega(y) = \overline{O(y)}$ is a locally dense Q-set. By [55, Lemma 2.3], we have $\omega(y) \cap E(v) = \overline{O(y)} \cap E(v) \subseteq \overline{\text{LD}(v)} \cap E(v) = \emptyset$. The ω -limit set $\omega(y) = \omega(x)$ is not transversely Cantor.

Suppose that $\omega(x)$ is locally dense. Then $\emptyset \neq O(x) \cap \text{int}(\omega(x)) \subseteq O(x) \cap \text{int}(\overline{O(x)})$. Therefore $x \in \text{int}(\omega(x))$ and so $\overline{O(x)} = \omega(x)$. This means that $\omega(x)$ is a locally dense Q-set and that $x \in \omega(x) \cap \text{LD}(v)$. \square

We have the following observation.

Lemma 3.7. *If the ω -limit set $\omega(x)$ of a point x intersects $E(v)$, then $\omega(x)$ is a quasi-Q-set that is not locally dense.*

Proof. Suppose $\omega(x)$ contains an orbit $O \subset E(v)$. We claim that $\omega(x) \cap \text{LD}(v) = \emptyset$. Indeed, assume that $\omega(x) \cap \text{LD}(v) \neq \emptyset$. Lemma 3.6 implies that $x \in \text{LD}(v)$ and that $\overline{O(x)} = \omega(x)$. [55, Lemma 2.3] implies that $O \subseteq \overline{O(x)} \cap E(v) = \overline{\text{LD}(v)} \cap E(v) = \emptyset$, which is a contradiction.

By Lemma 3.6, the ω -limit set $\omega(x)$ is not locally dense. Since O is non-closed recurrent, by Lemma 3.2, taking a small transverse arc if necessary, the waterfall construction implies that there is a closed transversal γ intersecting O infinitely many times. This means that $\omega(x)$ is a quasi-Q-set that is not locally dense. \square

We show the key lemma as follows.

Proof of Lemma 3.1. Let x be a point whose orbit is not closed. We may assume that $\omega(x)$ is not contained in $\text{Sing}(v)$. From Lemma 3.6, we may assume that $\omega(x) \cap \text{LD}(v) = \emptyset$ and that $\omega(x)$ is not locally dense. Lemma 3.7 implies that we may assume that $\omega(x) \cap E(v) = \emptyset$. Then $\omega(x) \cap R(v) = \emptyset$.

If there is a closed transversal that intersects $O^+(x)$ infinitely many times, then Lemma 3.4 implies the assertion. Thus we may assume that $O(x)^+$ has no closed transversal intersecting it infinitely many times. Then $\omega(x)$ is not a quasi-Q-set.

If $\omega(x)$ contains a periodic orbit, then $\omega(x)$ is a semi-attracting limit cycle. Thus we may assume that $\omega(x)$ contains neither periodic orbits nor Q-sets. Since $S = \text{Sing}(v) \sqcup \text{Per}(v) \sqcup P(v) \sqcup R(v)$ and $\omega(x) \not\subset \text{Sing}(v)$, the ω -limit set $\omega(x)$ contains a non-recurrent orbit O such that $\omega(x) \subset \text{Sing}(v) \sqcup P(v)$. Take a non-recurrent point $y \in O \subset \omega(x)$ and a transverse closed arc $J : [-1, 0] \rightarrow S$ with $y = J(0)$ and $x_0 := J(-1) \in O^+(x)$ such that $O^+(x_0)$ intersects $J([-1, 0])$ infinitely many times. Let $f_v : J \rightarrow J$ be the first return map and $x_n := (f_v)^n(x_0)$ n -th return of x_0 . Denote by C_n the orbit arc from x_n to $f_v(x_n)$ and by $I_n \subset J$ the closed arc with $\partial C_n = \partial I_n = \{x_n, f_v(x_n)\}$.

We claim that we may assume that $f_v|_{(-1, 0]}$ is orientation-preserving by shortening the transverse interval $J([-1, 0])$. Indeed, otherwise there are a subsequence $(x_{k_n})_{n \in \mathbb{Z}_{>0}}$ of $(x_n)_{n \in \mathbb{Z}_{>0}}$ in J converging to $J(0)$ and small neighborhoods U_{k_n} of the unions of $I_{k_n} \cup C_{k_n}$ each of which is an open Möbius band such that $U_{k_n} \cap U_{k_m} = \emptyset$ for any $n \neq m \in \mathbb{Z}_{>0}$, which contradicts that S has finitely many non-orientable genus.

We claim that we may assume that $\{J(0)\} = \{y\} = J([-1, 0]) \cap \omega(x) \subset \overline{O^+(J(-1)) \cap J([-1, 0])} \cap P(v)$ by shortening $J([-1, 0])$. Indeed, otherwise there is a point $x_N \in J((-1, 0)) \cap O^+(x)$ with $I_N \cap \omega(x) \neq \emptyset$ such that I_N intersects $O^+(x)$ infinitely many times and that the union of $I_N \cup C_N$ is a loop. By the waterfall construction to the loop $C_N \cup I_N$, there is a closed transversal T_{x_N} near the loop $C_N \cup I_N$ such that T_{x_N} intersects $O^+(x)$ infinitely many times, which contradicts that $O(x)^+$ has no closed transversal intersecting it infinitely many times.

By Lemma 3.3, the ω -limit set $\omega(x)$ is a limit quasi-circuit such that x is wandering and is not contained in the orbit closures of points outside of $O(x)$. Lemma 3.5 implies that the five possible invariant subsets are exclusive. \square

3.2. Classification of quasi-circuits. We have the following dichotomy.

Lemma 3.8. *A quasi-semi-attracting limit quasi-circuit either is the image of a circle or is not locally connected exclusively.*

Proof. Let γ be a quasi-semi-attracting limit quasi-circuit. By definition of non-trivial quasi-circuit, the quasi-circuit γ consists of singular points and non-recurrent points, and there is an open annulus \mathbb{A} such that γ is a boundary component of \mathbb{A} .

We claim that γ can be realized as a boundary component of an open disk. Indeed, taking \mathbb{A} small and using a bump function whose inverse image of the zero is another boundary component μ of \mathbb{A} and collapsing the another boundary component of \mathbb{A} into a singleton, the resulting subset is an open

disk U , and the quasi-circuit γ can be realized as the boundary component of the open disk U . This completes the claim.

Thus we may assume that there is an open disk U such that γ is a boundary component of U . By Riemann mapping theorem, there is a Riemann mapping $\varphi : U \rightarrow D$ from an open disk D from a unit open disk D in a complex plane. By Carathéodory-Torhorst theorem (see [50] and also [40, Theorem 16.6] for detail), the boundary $\partial U = \gamma$ is locally connected if and only if the inverse mapping $\varphi^{-1} : D \rightarrow U$ extends continuously to a map, also denoted by φ^{-1} , from the closed disk \overline{D} onto \overline{U} .

We claim that γ is locally connected if and only if γ is the image of a circle. Indeed, if γ is locally connected, then $\gamma = \varphi^{-1}(\partial D)$ is the image of a circle. Conversely, suppose that γ is the image of a circle. Let $p : \mathbb{S}^1 \rightarrow \gamma$ be the continuous surjection from a circle \mathbb{S}^1 . Since the circle \mathbb{S}^1 is compact and γ is Hausdorff, the continuous surjection p is closed and so is a quotient mapping. Because each quotient space of a locally connected space is locally connected (cf. [12, Proposition 12 (p.112)]), the image $p(\mathbb{S}^1) = \gamma$ is locally connected. \square

3.3. Classification of quasi-Q-sets. In this subsection, we classify quasi-Q-sets.

3.3.1. Inclusion relation between Q-sets and quasi-Q-sets. We describe a property between non-closed recurrent orbits and closed transversals.

Lemma 3.9. *For any non-closed recurrent point $x \in S$, there is a closed transversal γ through $O(x)$ such that the intersection $\gamma \cap O(x)$ is infinite. Moreover, each closed transversal through a non-closed recurrent point y is essential and intersects $O(y)$ infinitely many times.*

Proof. Fix a point $x \in R(v)$ and a transverse arc $I \subset U$ such that x is the interior point of I . Then $|I \cap O(x)| = \infty$. By Lemma 3.2, there are an orbit arc C in $O(x)$ and a transverse closed arc $J \subseteq I$ such that the union $\mu := J \cup C$ is a loop with $C \cap J = \partial C = \partial J$ and that the return map along C is orientation-preserving between neighborhoods of ∂C in I . By the waterfall construction to the loop μ , there is a closed transversal γ intersecting $O(x)$ near μ . Since x is non-closed recurrent, the intersection $\gamma \cap O(x)$ is infinite.

Let ν be a closed transversal through a non-closed recurrent point y . Suppose that ν is not essential. Let S^* be the resulting closed surface from S by collapsing all boundary components into singletons. Then ν is null homotopic in S^* and so the point $y \in P(v)$, which contradicts the recurrence of x . \square

This implies the following corollary.

Corollary 3.10. *A Q-set is a quasi-Q-set.*

$$\begin{array}{ccccc}
S & \xleftarrow{p_{\text{me}}} & S_{\text{me}} & \xrightarrow{p_{\text{col}}} & S_{\text{col}} \\
\cup & & \cup & & \cup \\
S - \text{Sing}(v) & \xlongequal{p_{\text{me}}|} & S_{\text{me}} - \text{Sing}(v_{\text{me}}) & \xlongequal{p_{\text{col}}|} & S_{\text{col}} - \text{Sing}(v_{\text{col}})
\end{array}$$

FIGURE 9. Canonical quotient mappings induced by the metric completion and the collapse

Proof. Let γ be a Q-set. Then there is a non-closed recurrent orbit $O \subset \gamma$. By Lemma 3.9, there is an essential closed transversal intersecting O infinitely many times. \square

3.3.2. Non-recurrent orbits in the ω -limit sets of points. We recall the following Maier's result [39] (cf. Theorem 2.4.4 p.32 [41], Theorem 4.2 [4]).

Lemma 3.11 (Maier). *A point $x \in \omega(z)$ for some point $z \in S$ with $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ is non-closed positively recurrent (i.e. $x \in \omega(x) \cap \text{R}(v)$).*

The existence of recurrence implies following statements.

Lemma 3.12. *The union $\partial^+ \text{LD}(v) \sqcup \partial^+ \text{E}(v)$ consists of singular points and connecting quasi-separatrices.*

Proof. Let x be a non-closed recurrent point. We claim that the difference $\overline{O(x)} - \hat{O}(x)$ consists of singular points and connecting quasi-separatrices. Indeed, [55, Proposition 2.2] implies $\overline{O(x)} - \hat{O}(x) \subseteq \text{Sing}(v) \sqcup \text{P}(v)$. [14, Theorem VI] implies that the orbit class $\hat{O}(x)$ contains infinitely many Poisson stable orbits and so there is a non-closed recurrent point z with $x \in \omega(z) \cap \alpha(z)$. Since any limit cycles are either semi-attracting or semi-repelling, by Lemma 3.11 and its dual statement for α -limit sets, we have $\omega(y) \cup \alpha(y) \subseteq \text{Sing}(v)$ for any point $y \in \overline{O(x)} - \hat{O}(x)$. This means that the union $\partial^+ \text{LD}(v) \sqcup \partial^+ \text{E}(v)$ consists of singular points and connecting quasi-separatrices. \square

Corollary 3.13. *Each non-recurrent orbit contained in a Q-set is a connecting quasi-separatrix.*

3.3.3. Properties of the resulting flows by collapsing connected components of singular points into singletons. Consider a flow v on a surface S . Denote by S_{me} the metric completion of the difference $S - \text{Sing}(v)$ and by v_{me} the resulting \mathbb{R} -action such that the new points are singular points. Let $p_{\text{me}} : S_{\text{me}} \rightarrow S$ be the canonical projection. Then $\text{Sing}(v_{\text{me}}) = p_{\text{me}}^{-1}(\text{Sing}(v))$. Let S_{col} be the resulting space from v_{me} by collapsing any connected components of $\text{Sing}(v_{\text{me}})$ into singletons. By construction, the resulting space S_{col} is a disjoint union of closed surfaces. Let v_{col} the resulting \mathbb{R} -action on S_{col} , and $p_{\text{col}} : S_{\text{me}} \rightarrow S_{\text{col}}$ the canonical projection (see Figure 9). Then

$\text{Sing}(v_{\text{col}}) = p_{\text{col}}(\text{Sing}(v_{\text{me}})) = p_{\text{col}}(p_{\text{me}}^{-1}(\text{Sing}(v)))$. By construction, we obtain $S - \text{Sing}(v) = S_{\text{me}} - \text{Sing}(v_{\text{me}}) = S_{\text{col}} - \text{Sing}(v_{\text{col}})$. We have the following continuity.

Lemma 3.14. *Let $v : \mathbb{R} \times S \rightarrow S$ be a flow on a surface S . The resulting \mathbb{R} -actions v_{me} and v_{col} are flows (i.e. continuous \mathbb{R} -actions).*

To demonstrate the previous lemma, we state the following observation.

Lemma 3.15. *Let $(a_n)_{n \in \mathbb{Z}_{\geq 0}}$ be a sequence of points in S and $(t_n)_{n \in \mathbb{Z}_{\geq 0}}$ be a sequence of points in \mathbb{R} such that the sequence $((-t_n, b_n))_{n \in \mathbb{Z}_{\geq 0}}$ in $\mathbb{R} \times S$ converges a point $(-t_\infty, b_\infty) \in \mathbb{R} \times S$ with $b_\infty \in \text{Sing}(v)$, where $b_n := v(t_n, a_n)$. Then $\lim_{n \rightarrow \infty} a_n = b_\infty = \lim_{n \rightarrow \infty} b_n$.*

Proof. By definition, we have $a_n = v(-t_n, b_n)$. Since the sequence $(t_n)_{n \in \mathbb{Z}_{\geq 0}}$ in \mathbb{R} converges a point $t_\infty \in \mathbb{R}$, there is a number $T_0 > 0$ with $\{t_n \mid n \in \mathbb{Z}_{\geq 0}\} \subset [-T_0, T_0]$. For any positive integer $n \in \mathbb{Z}_{>0}$, there is a neighborhood B_n of b_∞ such that $\sup\{d(b_\infty, v(t, y)) \mid (t, y) \in [-T_0, T_0] \times B_n\} < 1/n$. Taking a subsequence of $((-t_n, b_n))_{n \in \mathbb{Z}_{\geq 0}}$, we may assume that $v(-t_n, b_n) \in B_n$. Then $d(b_\infty, a_n) = d(b_\infty, v(-t_n, b_n)) < 1/n$ for any positive integer $n \in \mathbb{Z}_{>0}$. This means that $b_\infty = \lim_{n \rightarrow \infty} a_n$. \square

Using the previous observation, we show the following continuity of v_{me} and v_{col} .

Proof of Lemma 3.14. By construction, we obtain $S - \text{Sing}(v) = S_{\text{me}} - \text{Sing}(v_{\text{me}}) = S_{\text{col}} - \text{Sing}(v_{\text{col}})$. Therefore $v = v_{\text{me}} = v_{\text{col}}$ on $S - \text{Sing}(v) = S_{\text{me}} - \text{Sing}(v_{\text{me}}) = S_{\text{col}} - \text{Sing}(v_{\text{col}})$. Therefore v_{me} and v_{col} are \mathbb{R} -actions.

We claim that $v_{\text{me}} : \mathbb{R} \times S_{\text{me}} \rightarrow S_{\text{me}}$ is continuous. Indeed, fix any closed subset $A_{\text{me}} \subseteq S_{\text{me}}$. Then the set difference $A_{\text{me}} \setminus \text{Sing}(v_{\text{me}})$ is closed in $S_{\text{me}} - \text{Sing}(v_{\text{me}}) = S - \text{Sing}(v)$, and the intersection $A_{\text{me}} \cap \text{Sing}(v_{\text{me}})$ is closed in S_{me} . From singularity and definition of product topology, the inverse image $v_{\text{me}}^{-1}(A_{\text{me}} \cap \text{Sing}(v_{\text{me}})) = \mathbb{R} \times (A_{\text{me}} \cap \text{Sing}(v_{\text{me}}))$ is closed. By continuity of v , the inverse image $v^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}})) = v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))$ is closed in $S - \text{Sing}(v) = S_{\text{me}} - \text{Sing}(v_{\text{me}})$. Then the closure $\overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))}$ is contained in $\overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))} \sqcup (\mathbb{R} \times \text{Sing}(v_{\text{me}}))$. By Lemma 3.15, since any points in $\overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))}$ are accumulation points of convergence sequences of points in $v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))$, we obtain $\overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))} \subseteq \overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))} \sqcup (\mathbb{R} \times (A_{\text{me}} \cap \text{Sing}(v_{\text{me}}))) = \overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))} \sqcup \overline{v_{\text{me}}^{-1}(A_{\text{me}} \cap \text{Sing}(v_{\text{me}}))} = v_{\text{me}}^{-1}(A_{\text{me}})$. Therefore the union $\overline{v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}}))} \cup \overline{v_{\text{me}}^{-1}(A_{\text{me}} \cap \text{Sing}(v_{\text{me}}))} = v_{\text{me}}^{-1}(A_{\text{me}})$ is closed. This complete the claim.

We claim that $v_{\text{col}} : \mathbb{R} \times S_{\text{col}} \rightarrow S_{\text{col}}$ is continuous. Indeed, fix any closed subset $A_{\text{col}} \subseteq S_{\text{col}}$. The inverse image $p_{\text{col}}^{-1}(A_{\text{col}})$ is closed in S_{me} . Then the set difference $A_{\text{col}} \setminus \text{Sing}(v_{\text{col}}) = A_{\text{me}} \setminus \text{Sing}(v_{\text{me}})$ is closed in $S_{\text{col}} - \text{Sing}(v_{\text{col}}) = S_{\text{me}} - \text{Sing}(v_{\text{me}})$, and the intersection $A_{\text{me}} \cap \text{Sing}(v_{\text{me}})$ is closed in S_{me} . By continuity of v_{me} , the inverse image $v_{\text{me}}^{-1}(A_{\text{me}} \setminus \text{Sing}(v_{\text{me}})) =$

$v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}}))$ is closed in $S_{\text{me}} - \text{Sing}(v_{\text{me}}) = S_{\text{col}} - \text{Sing}(v_{\text{col}})$. By Lemma 3.15, we obtain the following equality:

$$\begin{aligned}
\overline{v_{\text{col}}^{-1}(A_{\text{col}})} &= \overline{v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}})) \sqcup v_{\text{col}}^{-1}(A_{\text{col}} \cap \text{Sing}(v_{\text{col}}))} \\
&= \overline{v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}})) \cup v_{\text{col}}^{-1}(A_{\text{col}} \cap \text{Sing}(v_{\text{col}}))} \\
&= \overline{v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}}))} \cup \overline{v_{\text{col}}^{-1}(A_{\text{col}} \cap \text{Sing}(v_{\text{col}}))} \\
&= \overline{v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}}))} \cup (\mathbb{R} \times (A_{\text{col}} \cap \text{Sing}(v_{\text{col}}))) \\
&= \overline{v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}}))} \sqcup (\mathbb{R} \times (A_{\text{col}} \cap \text{Sing}(v_{\text{col}}))) \\
&= v_{\text{col}}^{-1}(A_{\text{col}} \setminus \text{Sing}(v_{\text{col}})) \sqcup v_{\text{col}}^{-1}(A_{\text{col}} \cap \text{Sing}(v_{\text{col}})) = v_{\text{col}}^{-1}(A_{\text{col}})
\end{aligned}$$

This complete the claim. \square

3.3.4. Properties of connecting quasi-separatrices. We have the following statement.

Proposition 3.16. *Each orbit in the ω -limit set of a point is non-recurrent if and only if it is a connecting quasi-separatrix.*

Proof. By definition, a connecting quasi-separatrix is non-recurrent. Conversely, suppose that O is a non-recurrent orbit in the ω -limit set of a point. Let x_0 be a point with $O \subset \omega(x_0)$. If $\omega(x_0)$ is a Q-set, then Corollary 3.13 implies that O is a connecting quasi-separatrix. Thus we may assume that $\omega(x_0)$ is not a Q-set. Then x_0 is not positively recurrent. Lemma 3.1 implies that $\omega(x_0)$ is either a quasi-semi-attracting limit quasi-circuit or a quasi-Q-set that is not locally dense. Therefore $\omega(x_0)$ consists of singular points and non-recurrent orbits.

We claim that $\omega_{v_{\text{col}}}(x_0)$ consists of singular points and connecting separatrices. Indeed, taking the double of the surface S if necessary, we may assume that S is closed in the proof of the claim. By Gutierrez's smoothing theorem [22], the flow v is topologically equivalent to a C^1 -flow and so is generated by a continuous vector field on S . Since the vector field is lifted to the orientation double covering of S , by the invariance of the claimed property, it suffices to show that the ω -limit set of the lift of O is contained in a boundary component of the singular point set. Therefore we may assume that S is orientable in the proof of the claim. From Theorem 3 [45], since $\text{Sing}(v)$ is closed, each connected component of the open surface $S - \text{Sing}(v)$ is homeomorphic to the resulting surface from a closed surface by removing a closed totally disconnected subset. Let $S_{\text{col}}(O)$ be the connected component of S_{col} containing the orbit $O = p_{\text{col}}(p_{\text{me}}^{-1}(O))$. Then $S_{\text{col}}(O)$ is the orientable closed surface such that the restriction $v_{\text{col}}|_{S_{\text{col}}(O)}$ is a flow whose singular point set is totally disconnected. By construction, the image p_{col} and the inverse image p_{me}^{-1} preserve singular property and non-recurrence. Therefore the lift $p_{\text{col}}(p_{\text{me}}^{-1}(\omega(x_0)))$ is a closed invariant subset that consists of singular points and non-recurrent points. By construction, we have $p_{\text{col}}(p_{\text{me}}^{-1}(\omega(x_0))) = \omega_{v_{\text{col}}}(x_0)$, where $\omega_{v_{\text{col}}}(x_0)$ is the ω -limit set of

x_0 with respect to v_{col} . Since $\omega_{v_{\text{col}}}(x_0)$ consists of singular points and non-recurrent points, by [36, Theorem 3.1] (cf. [38, Theorem 2.1]), the ω -limit set $\omega_{v_{\text{col}}}(x_0)$ consist of singular points and connecting separatrices. This completes the claim.

This means that the orbit O is a connecting quasi-separatrix. \square

Proposition 3.16 implies the following reduction.

Corollary 3.17. *Suppose that the singular point set is totally disconnected. Then each orbit in the ω -limit set of a point is non-recurrent if and only if it is a connecting separatrix.*

We show the following statements.

Lemma 3.18. *For any positively recurrent point $y \in E(v)$ and any non-positive-recurrent point x with $y \in \omega(x)$, the ω -limit set $\omega(x) = \omega(y) = \overline{O(y)}$ is a transversely Cantor Q -set which is not locally dense.*

The following proof is an analogous argument of the proof of [41, Theorem 2.4.1].

Proof of Lemma 3.18. By the positive recurrence of y , we obtain $\omega(y) = \overline{O(y)}$. We claim that $\omega(x) = \overline{O(y)}$. Indeed, assume that there is a point $z \in \omega(x) - \overline{O(y)}$. Then $O(x) \cap O(y) = \emptyset$. Since S is a T_3 -space, there is a closed disk B whose interior contains z such that $B \cap \overline{O(y)} = \emptyset$. Therefore $(O(x) \cup B) \cap O(y) = \emptyset$. The boundary $\gamma := \partial B$ is a simple closed curve. Since the point y is positively recurrent, there are points $\alpha, \beta \in O^+(y)$, a transverse closed arc J whose endpoints are α and β , and there is a sequence $(y_n)_{n \in \mathbb{Z}_{\geq 0}}$ of points in $J \cap O^+(y)$ converging to β such that $y_{n+1} \in O^+(y_n)$ and $J \cap B = \emptyset$. For any $n \in \mathbb{Z}_{\geq 0}$, from $z \in \omega(x)$, $y_{2n+1} \in \omega(x) \cap I$ and $\beta = \lim_{m \rightarrow \infty} y_m$, there are points $x_n, \tilde{x}_n \in O^+(x)$ and an orbit arc $C_n \subset O^+(x)$ as in Figure 10 satisfying the following properties:

- (1) $\tilde{x}_n \in O^+(x_n)$ and $x_{n+1} \in O^+(\tilde{x}_n)$.
- (2) The endpoints of C_n are x_n and \tilde{x}_n .
- (3) $C_n \cap B = \{x_n, \tilde{x}_n\}$, $|C_n \cap J| = 1$, and $C_n \cap J \subset I_{y_{2n}, y_{2n+2}}$.

Here $I_{a,b}$ is the closed sub-arc of I whose endpoints are a and b . Put $C := \gamma \cup \bigsqcup_{n \in \mathbb{Z}_{\geq 0}} C_n$. Since $(O(x) \cup B) \cap O(y) = \emptyset$, we have $O^+(y) \cap C = \emptyset$. Then there are infinitely many connected components D_n of $S - C$ which are bounded by the unions of $C_n \sqcup C_{n+1}$ and two sub-arcs in γ . Since S is compact and so the genus of S is at most finite, taking subsequences of $(x_n)_{n \in \mathbb{Z}_{\geq 0}}$ and $(y_n)_{n \in \mathbb{Z}_{\geq 0}}$, we may assume that any connected components D_n are open disks. By construction, the open disk D_n contains $y_{2n+2} \in O^+(y)$. Since $O^+(y) \cap C = \emptyset$, we have $O^+(y_{2n+2}) \cap \partial D_n = \emptyset$ and so $O^+(y_{2n+2}) \subset D_n$. Therefore $y_{2n+4} \in D_n \cap D_{n+1} = \emptyset$, which is a contradiction.

By [55, Lemma 2.3], we obtain $\omega(x) \cap \text{LD}(v) = \overline{O(y)} \cap \text{LD}(v) \subseteq \overline{E(v)} \cap \text{LD}(v) = \emptyset$. This means that the Q -set $\omega(x)$ is not locally dense. \square




FIGURE 10. A flow box with one singular point with respect to the vector field $\varphi X = (\varphi, 0)$.

Lemma 3.19. *A quasi-Q-set that is not a Q-set consists of singular points and connecting quasi-separatrices.*

Proof. Let \mathcal{M} be a quasi-Q-set that is not a Q-set. By time reversion if necessary, we may assume that \mathcal{M} is the ω -limit set of a non-positive-recurrent point. Then there is a non-positive-recurrent point $x \in S$ with $\omega(x) = \mathcal{M}$. If \mathcal{M} contains a periodic point, then \mathcal{M} is a semi-attracting limit cycle such that it intersects any essential closed transversal at most finitely many times, which contradicts the definition of quasi-Q-set. Thus $\mathcal{M} \cap \text{Per}(v) = \emptyset$. By Lemma 2.2, we have $\mathcal{M} \cap \text{LD}(v) = \emptyset$.

We claim that $E(v) \cap \mathcal{M} = \emptyset$. Indeed, assume that $E(v) \cap \mathcal{M} \neq \emptyset$. By [14, Theorem VI], there is a point $y \in E(v) \cap \mathcal{M}$ with $\overline{O^-(y)} = \alpha(y) = \omega(y)$. Since $\omega(x) = \mathcal{M}$, Lemma 3.18 implies that the quasi-Q-set $\mathcal{M} = \omega(x) = \overline{O^-(y)}$ is a Q-set, which contradicts the hypothesis.

Therefore $\mathcal{M} \subset \text{Sing}(v) \sqcup \text{P}(v)$. Proposition 3.16 implies that \mathcal{M} consists of singular points and connecting quasi-separatrices. \square

We generalize the Poincaré-Bendixson theorem into one for a flow with arbitrarily many singular points on a compact surface and the Maier's description of recurrence as follows.

3.4. Proof of Theorem A. Let v be a flow on a compact surface S and x a non-closed point. Lemma 3.1 implies that $\omega(x)$ is either a nowhere dense subset of singular points, a semi-attracting limit cycle, a quasi-semi-attracting limit quasi-circuit, a locally dense Q-set, or a quasi-Q-set that is not locally dense. Proposition 3.16 implies that any non-recurrent orbit in the ω -limit set of a point is a connecting quasi-separatrix. This means that the assertion (b) holds. Corollary 3.17 implies the assertion (c). By Lemma 3.8, a quasi-semi-attracting limit quasi-circuit either is the image of a circle or is not locally connected exclusively. From Corollary 3.10, a quasi-Q-set that is not locally dense is either a transversely Cantor Q-set

or a quasi-Q-set that is not a Q-set. By Lemma 3.19, a quasi-Q-set that is not a Q-set corresponds to a quasi-Q-set that consists of singular points and connecting quasi-separatrices. This completes the proof of Theorem A.

3.5. Proof of Theorem B. Let v be a flow on a compact surface S and a point $x \in \omega(z)$ for some point $z \in S$. Lemma 3.11 implies that if $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ then x is non-closed positively recurrent. Conversely, if x is non-closed positively recurrent, then $x \in \omega(x) \setminus \text{Cl}(v)$ and so $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$. This completes the assertion (1).

If x is non-closed negatively recurrent, then $x \in \alpha(x) \setminus \text{Cl}(v)$ and so $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$. Conversely, suppose that $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$. Then $x \notin \text{Cl}(v)$ and so $x \in P(v) \sqcup R(v)$. We claim that x is non-closed recurrent. Indeed, assume that x is non-recurrent. Theorem A implies that $O(x)$ is a connecting quasi-separatrix and so that $\alpha(x) \subseteq \text{Sing}(v)$, which contradicts $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$. Thus x is non-closed recurrent. [55, Proposition 2.2] implies that $x \in \overline{O(z)} \setminus (\text{Sing}(v) \sqcup P(v)) \subset R(v)$. By [14, Theorem VI], the orbit class $\hat{O}(x)$ contains infinitely many Poisson stable orbits. Therefore there is a Poisson stable point $y \in \hat{O}(x)$ such that $x \in \alpha(y)$. The dual of Lemma 3.11 implies that x is non-closed negatively recurrent. This completes the assertion (2).

The assertions (3) and (4) are followed from the assertions (1) and (2).

3.6. Proof of Corollary C. By [14, Theorem VI], the orbit class of a non-closed recurrent point contains infinitely many Poisson stable orbits. Therefore Theorem B and its dual statement imply that the conditions (1)–(3) are pairwise equivalent.

4. POINCARÉ-BENDIXSON THEOREM FOR A NON-COMPACT SURFACE AND GENERALIZATION OF MAÏER'S DESCRIPTION OF RECURRENCE

In this section, we generalize the Poincaré-Bendixson theorem to one for a flow with arbitrarily many singular points on a surface of finite genus and finitely many boundary components. Moreover, we also characterize the recurrence, which are generalizations of the Maïer's description of recurrence for such a flow.

4.1. Poincaré-Bendixson theorem for a flow with arbitrarily many singular points on a surface of finite genus and finitely many boundary components. Recall the end completion as follows.

4.1.1. End completion of a topological space. Consider the direct system $\{K_\lambda\}$ of compact subsets of a topological space Y and inclusion maps such that the interiors of K_λ cover Y . There is a corresponding inverse system $\{\pi_0(Y - K_\lambda)\}$, where $\pi_0(Z)$ denotes the set of connected components of a topological space Z . Then the **set of ends** of Y is defined to be the inverse limit of this inverse system. Notice that Y has one end $x_{\mathcal{U}}$ for each sequence $\mathcal{U} := (U_i)_{i \in \mathbb{Z}_{>0}}$ with $U_i \supseteq U_{i+1}$ such that U_i is a connected component of $Y - K_{\lambda_i}$ for some λ_i . Considering the disjoint union Y_{end} of Y

and $\{\pi_0(Y - K_\lambda)\}$ as set, a subset V of the union Y_{end} is an open neighborhood of an end $x_{\mathcal{U}}$ if there is some $i \in \mathbb{Z}_{>0}$ such that $U_i \subseteq V$. Then the resulting topological space Y_{end} is called the **end completion** (or end compactification) of Y . Note that the end completion is not compact in general. Moreover, the surface S_{col} is the end completion of $S - \text{Sing}(v)$. From Theorem 3 [45], any connected surfaces of finite genus and finitely many boundary components are homeomorphic to the resulting surfaces from compact surfaces by removing closed totally disconnected subsets. Therefore the end compactification S_{end} of a connected surface S of finite genus and finitely many boundary components is a compact surface.

4.1.2. Concepts for flows on (possibly non-compact) surfaces. For a flow v on a surface S of finite genus and finitely many boundary components, considering ends to be singular points, we obtain the resulting flow v_{end} on a surface S_{end} which is a union of compact surfaces. A non-recurrent orbit on S is a **virtual quasi-separatrix** if it is a connecting quasi-separatrix on S_{end} with respect to v_{end} . A non-recurrent orbit on S is a **virtual separatrix** if it is a connecting separatrix on S_{end} with respect to v_{end} . An invariant subset on S is a **quasi-semi-attracting limit virtual quasi-circuit** if it is the resulting subset from a quasi-semi-attracting limit quasi-circuit on S_{end} with respect to v_{end} by removing all the ends.

4.1.3. Poincaré-Bendixson theorem for flows on (possibly non-compact) surfaces. By taking end completions, Theorem A implies the following Poincaré-Bendixson theorem for a flow with arbitrarily many singular points on a surface of finite genus and finitely many boundary components.

Theorem 4.1. *The following statements hold for a flow with arbitrarily many singular points on a surface of finite genus and finitely many boundary components:*

- (a) *The ω -limit set of any non-closed orbit is one of the following exclusively:*
 - (1) *A nowhere dense subset of singular points.*
 - (2) *A semi-attracting limit cycle.*
 - (3) *A quasi-semi-attracting limit non-periodic virtual quasi-circuit.*
 - (4) *A locally dense Q -set.*
 - (5) *A transversely Cantor Q -set.*
 - (6) *A quasi- Q -set that consists of singular points and non-recurrent points.*
- (b) *Any non-recurrent orbit in the ω -limit set of a point is a virtual quasi-separatrix.*
- (c) *If the singular point set is totally disconnected, then any non-recurrent orbits in the ω -limit set of a point are virtual separatrices.*

In the previous theorem, notice that the ω -limit set of a point is empty if and only if it is a nowhere dense subset of singular points.

4.2. Topological characterizations of non-closed recurrence for surfaces. For a flow v on a connected surface with finite genus and finitely many boundary components, the end completion S_{end} is a compact connected surface, and the resulting flow v_{end} on it can be obtained by adding exactly new singular points. This implies that Theorem B holds for a flow with arbitrarily many singular points on a surface with finite genus and finitely many boundary components as follows.

Theorem 4.2. *Let v be a flow on a surface S with finite genus and finitely many boundary components. The following statements hold for a point $x \in \omega(z)$ for some point $z \in S$:*

- (1) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed positively recurrent.
- (2) $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed negatively recurrent.
- (3) $(\omega(x) \cup \alpha(x)) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed recurrent.
- (4) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$ if and only if x is non-closed Poisson stable.

Proof. As mentioned above, taking the end completion S_{end} of S , the resulting surface S_{end} is a compact connected surface and the resulting flow v_{end} is obtained by adding singular points. Therefore the conditions that $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and $\alpha(x) \setminus \text{Cl}(v) \neq \emptyset$ are invariant under taking end completions. This means that Theorem B holds for a flow on a connected surface S with finite genus and finitely many boundary components. \square

Corollary C implies the following topological characterizations of non-closed recurrence for a flow with arbitrarily many singular points on a surface of finite genus and finitely many boundary components.

Corollary 4.3. *Let v be a flow on a surface S with finite genus and finitely many boundary components. The following are equivalent for a point $x \in S$:*

- (1) *The point x is non-closed positively recurrent.*
- (2) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and there is a point $z \in S$ with $x \in \omega(z)$.
- (3) $\omega(x) \setminus \text{Cl}(v) \neq \emptyset$ and there is a point $z \in S$ with $x \in \alpha(z)$.

5. REDUCTIONS OF QUASI-Q-SETS AND QUASI-CIRCUITS

Let v be a flow on a compact connected surface S .

5.1. Reductions of quasi-Q-sets into Q-set under finite existence of singular points. We show that a quasi-Q-set is a generalization of a Q-set.

Proposition 5.1. *Suppose that the singular point set is finite. Then a quasi-Q-set corresponds to a Q-set.*

Proof. By Corollary 3.10, a Q-set is a quasi-Q-set. Suppose that the singular point set is finite. Fix a point x whose ω -limit set $\omega(x)$ is a quasi-Q-set. Then $\omega(x)$ intersects an essential closed transversal T infinitely many times. Then x is not a closed orbit. By Theorem A, the finiteness of $\text{Sing}(v)$ implies that any non-recurrent orbits in $\omega(x)$ are connecting separatrices.

If $\omega(x) \cap \text{Per}(v) \neq \emptyset$, then the ω -limit set $\omega(x)$ is a limit cycle. Thus $\omega(x) \cap \text{Per} = \emptyset$.

We claim that $\omega(x) \cap R(v) \neq \emptyset$. Indeed, assume that $\omega(x) \cap R(v) = \emptyset$. Then $\omega(x) \subseteq \text{Sing}(v) \sqcup P(v)$. Then any non-singular orbits of $\omega(x)$ are connecting separatrices. Since $\omega(x)$ intersects an essential closed transversal T infinitely many times, the ω -limit set $\omega(x)$ contains infinitely many connecting separatrices. By the finiteness of $\text{Sing}(v)$, there are singular points α and ω and infinitely many pairwise disjoint connecting separatrices O_i ($i \in \mathbb{Z}_{\geq 0}$) from α to ω . Since S is compact and the genus is finite, taking a subsequence, we may assume that the pair of two orbits O_i and O_j for any $i \neq j \in \mathbb{Z}_{\geq 0}$ is homotopic relative to $\{\alpha, \omega\}$ to each other and that the union $\gamma_i := \{\alpha, \omega\} \sqcup O_i \sqcup O_{i+1}$ for any $i \in \mathbb{Z}_{\geq 0}$ bounds an open disk B_i . Then the disjoint unions $B_i \sqcup O_{i+1} \sqcup B_{i+1}$ are neighborhoods of O_{i+1} . By construction, any orbits intersecting some B_j are contained in B_j . Since $O_1, O_2 \subset \omega(x)$, we have that $O(x) \cap (B_1 \cap B_2) \neq \emptyset$ and $O(x) \cap (B_2 \cap B_3) \neq \emptyset$ and so that $O(x) \subset B_2$. This implies that $O_4 \cap \omega(x) = \emptyset$, which contradicts $O_4 \subset \omega(x)$.

Theorem A implies that $\omega(x)$ is a Q-set. \square

5.2. Reductions of quasi-circuits. We show that a limit quasi-circuit is a generalization of a limit circuit.

Proposition 5.2. *Let v be a flow with finitely many singular points on a compact connected surface S . Then a limit quasi-circuit is a limit circuit (i.e. a continuous image of a circle).*

Proof. Let γ be a limit quasi-circuit and \mathbb{A} a small collar. By Theorem A, the finiteness of $\text{Sing}(v)$ implies that any non-recurrent orbits in γ are connecting separatrices. The invariance of γ implies that each orbit closure in γ is either a singular point, a closed interval, or a loop. Since $\text{Sing}(v)$ is finite, we may assume that \mathbb{A} contains no singular points. By time reversion if necessary, we may assume that $\omega(x) = \gamma$ for a point x . Then the difference $\gamma \setminus \text{Sing}(v)$ is a disjoint union of open intervals.

We claim that γ is locally connected. Indeed, assume that γ is not locally connected. Since each orbit closure in γ is either a point, a closed interval, or a loop, the finiteness of singular points implies that there are non-recurrent points $x_i \in \gamma$ converging a point $x_\infty \in \gamma \setminus (\bigcup_{i \in \mathbb{Z}_{\geq 0}} \alpha(x_i) \cup \omega(x_i))$ such that $O(x_i) \neq O(x_\infty)$ and $O(x_i) \neq O(x_j)$ for any $i \neq j \in \mathbb{Z}_{\geq 0}$. By renumbering, we may assume that there are singular points α and ω in γ such that $x_i \neq x_j$, $\alpha = \alpha(x_i)$, and $\omega = \omega(x_i)$ for any $i \neq j \in \mathbb{Z}_{\geq 0}$. Put $O_i := O(x_i)$. As the proof of the previous lemma, since S is compact and the genus is finite, taking a subsequence, we may assume that the pair of two orbits O_i and O_j for any $i \neq j \in \mathbb{Z}_{\geq 0}$ is homotopic relative to $\{\alpha, \omega\}$ to each other and that the union $\gamma_i := \{\alpha, \omega\} \sqcup O_i \sqcup O_{i+1}$ for any $i \in \mathbb{Z}_{\geq 0}$ bounds an open disk B_i . Then the disjoint unions $B_i \sqcup O_{i+1} \sqcup B_{i+1}$ are neighborhoods of O_{i+1} . By construction, any orbits intersecting some B_j are contained in B_j . Since $O_1, O_2 \subset \omega(x)$, we have that $O(x) \cap (B_1 \cap B_2) \neq \emptyset$ and $O(x) \cap (B_2 \cap B_3) \neq \emptyset$

and so that $O(x) \subset B_2$. This implies that $O_4 \cap \omega(x) = \emptyset$, which contradicts $O_4 \subset \omega(x)$.

We claim that γ is a circuit. Indeed, let ∂ be another boundary component of \mathbb{A} . Cutting ∂ , collapsing the new boundary components into singletons, and taking its universal cover, we may assume that γ is the boundary of an open disk U obtained from \mathbb{A} in a simply connected Riemann surface. By Riemann mapping theorem, there is a Riemann mapping $\varphi : D \rightarrow U$ from a unit open disk in a complex plane. By Carathéodory-Torhorst theorem (see [51] and also Theorem 16.6 [40] for detail), the local connectivity of the boundary $\partial U = \gamma$ implies that φ extends continuously to a map from the closed disk \overline{D} onto \overline{U} . This means that γ is a continuous image of a circle and a boundary component of its collar \mathbb{A} .

Since $\omega(x) = \gamma$, the circuit γ is a limit circuit. \square

5.3. Reduction of the Poincaré-Bendixson theorem for flows with finitely many singular points. The reductions of quasi-Q-sets (Proposition 5.1) and of quasi-circuits (Proposition 5.2) imply a proof of the following generalization of the Poincaré-Bendixson theorem for a flow with finitely many singular points (see for example [41]).

Corollary 5.3. *Let v be a flow with finitely many fixed points on a compact surface S . The ω -limit set of any non-closed orbit is one of the following exclusively:*

- (1) *A singular point.*
- (2) *A semi-attracting limit cycle.*
- (3) *A semi-attracting non-periodic limit circuit.*
- (4) *A locally dense Q-set.*
- (5) *A transversely Cantor Q-set.*

5.3.1. Poincaré-Bendixson theorem for a flow with finitely many singular points on possibly non-compact surfaces. Recall that a non-recurrent orbit on S is a virtual separatrix if it is a connecting separatrix on S_{end} with respect to v_{end} . An invariant subset on S is a **attracting limit virtual circuit** if it is the resulting subset from an attracting limit circuit on S_{end} with respect to v_{end} by removing all the ends.

The previous corollary implies the following generalization of the Poincaré-Bendixson theorem for a flow with finitely many singular points on possibly non-compact surfaces of finite genus and finitely many boundary components.

Corollary 5.4. *The following statements hold for a flow with finitely many singular points on a surface of finite genus and finitely many boundary components:*

- (a) *The ω -limit set of any non-closed orbit is one of the following exclusively:*
 - (0) *The empty set.*
 - (1) *A singular point.*
 - (2) *A semi-attracting limit cycle.*

- (3) *A semi-attracting limit non-periodic virtual circuit.*
 - (4) *A locally dense Q -set.*
 - (5) *A transversely Cantor Q -set.*
- (b) *Any non-recurrent orbit in the ω -limit set of a point is a virtual separator.*

6. CONSTRUCTION OF FLOW BOXES WITH NON-ARCWISE-CONNECTED INVARIANT SUBSETS

In this section, we introduce an operation that makes ω -limit sets not arcwise-connected by constructing flow boxes with non-arcwise-connected invariant subsets. To state the operation, we have the following statement.

Lemma 6.1. *For any flow v on a surface S with a non-singular point x , there is a trivial flow box B containing x such that the resulting flow w by replacing B with a flow box satisfying the following properties:*

- (1) *The restriction $v|_{S-O(x)}$ is topologically equivalent to the restriction $w|_{S-O(x)}$.*
- (2) *For any point $y \in S - O(x)$, we have that $\alpha(y) = \alpha_w(y)$ and $\omega(y) = \omega_w(y)$.*
- (3) *If $O(x)$ is periodic, then $O(x)$ is the disjoint union of one non-recurrent orbit O_0 and one singular point x with $\omega_w(O_0) = x = \alpha_w(O_0)$.*
- (4) *If $O(x)$ is not periodic, then $O(x)$ is the disjoint union of two non-singular orbits O_1, O_2 and one singular point x with $\omega_w(O_1) = x = \alpha_w(O_2)$ such that $\alpha(O_1) = \alpha_w(O_1)$ and $\omega(O_2) = \omega_w(O_2)$.*

We call w in the previous lemma the **resulting flow of v by replacing a non-singular point x with a singular point**, and denote by v_x the resulting flow w . Roughly speaking, the resulting flow by replacing a non-singular point with a singular point is a flow obtained by replacing a trivial flow box with a flow box as in Figure 11.

Proof of Lemma 6.1. By Gutierrez's smoothing theorem [22], we may assume that the flow v is a C^1 -flow generated by a continuous vector field X on S . Fix any open trivial flow box B intersecting $O(x)$. Identifying B with the square $[-1, 1]^2$ such that $\{0\} \times [-1, 1]$ is an orbit arc of $O(x)$ and that $v|_B$ is generated by a vector field $X = (1, 0)$. Take a C^∞ bump function $\varphi: B = [-1, 1]^2 \rightarrow [0, 1]$ with $\varphi^{-1}(0) = \{0\}$ such that φ is one near the boundary $\partial[-1, 1]^2$. Consider the flow box B' on B whose orbits arc are generated by the vector field $\varphi X = (\varphi, 0)$ as in Figure 11. Replacing the trivial flow box B with the flow box B' , the resulting flow is desired. \square

A flow $v: \mathbb{R} \times Z \rightarrow Z$ is **topologically semi-conjugate** to a flow $w: \mathbb{R} \times Y \rightarrow Y$ via $h: Y \rightarrow Z$ if h is a continuous surjection such that $v(t, h(y)) = h(w(t, y))$ for any $(t, y) \in \mathbb{R} \times Y$. In this section, we show the following statement.




FIGURE 11. A flow box with one singular point with respect to the vector field $\varphi X = (\varphi, 0)$.




FIGURE 12. The flow box \mathbb{B} with a non-arcwise-connected invariant subset with respect to the vector field X .

Theorem 6.2. *Let v be a flow on a surface S with an ω -limit set ω containing non-singular point x and with a point $y \in S - \omega$ satisfying $\omega(y) = \omega$. Then there is a trivial flow box B containing x such that the resulting flow w by replacing B with a flow box satisfying the following properties:*

- (1) *The ω -limit set $\omega_w(y)$ is not arcwise-connected unless it is locally dense.*
- (2) *The restriction $v|_{S-\omega}$ to the complement $S - \omega$ is topologically equivalent to the restriction $w|_{S-\omega_w(y)}$.*
- (3) *The flow v_x is topologically equivalent to a flow which is topologically semi-conjugate to w , where v_x is the resulting flow of v by replacing x with a singular point.*
- (4) *The topological conjugacy from w to v_x can be obtained by collapsing a closed invariant subset of $\omega_w(y)$ into a singleton.*

Roughly speaking, the resulting flow w in the previous theorem can be obtained by replacing a trivial flow box with a flow box as in Figure 12. This theorem implies Theorem D.

6.1. Construction of a flow box with a non-arcwise-connected connected invariant subset. Let $\mathbb{D} := [0, 1] \times [-1, 1]$ a closed square and $\partial_{\text{in}} \mathbb{D} := \{0, 1\} \times [-1, 1]$. Define a diffeomorphism $h_0: (0, 1) \rightarrow \mathbb{R}$ by $h_0(x) :=$




FIGURE 13. An ω -limit set which is a non-locally-connected quasi-circuit, which is neither the image of a circle nor a circuit.

$\tan\left(\frac{\pi(x-1)}{2}\right)$. Define a function $f: (0, 1) \rightarrow [-1, 1]$ as follows:

$$f(x) := \cos(h_0(x)) = \cos\left(\tan\left(\frac{\pi(x-1)}{2}\right)\right)$$

Let $G := \{(x, f(x)) \mid x \in (0, 1)\} \subset \mathbb{D}$ be the graph of f and $C := G \sqcup \partial_{\mathfrak{h}} \mathbb{D}$ the union. Consider a closed square $B := [0, 1] \times [-2, 2]$ containing \mathbb{D} . Put $\partial_{\mathfrak{h}} B := \{0, 1\} \times [-2, 2]$. Then the set difference $B - \partial B = (0, 1) \times (-2, 2)$ is an open square $\text{int } B$. We have the following observation.

Lemma 6.3. *The set difference $\text{int } B - G$ consists of two open disks.*

Proof. The closure \overline{G} in \mathbb{R}^2 is the union C . Let \mathcal{F} be a decomposition of \mathbb{R}^2 by $\mathcal{F} := \{\{p\} \mid p \in \mathbb{R}^2 - \partial_{\mathfrak{h}} \mathbb{D}\} \sqcup \{\{0\} \times [-1, 1], \{1\} \times [-1, 1]\}$. By Moore's theorem (cf. p.3 in [16]), the quotient space \mathbb{R}^2/\mathcal{F} of the upper semicontinuous decomposition \mathcal{F} into non-separating continua is homeomorphic to \mathbb{R}^2 , where \mathbb{R}^2/\mathcal{F} is the quotient space $\mathbb{R}^2/\sim_{\mathcal{F}}$ defined by $p \sim_{\mathcal{F}} q$ if there is an element of \mathcal{F} containing p and q . Let $p_{\mathcal{F}}: \mathbb{R}^2 \rightarrow \mathbb{R}^2/\mathcal{F}$ be the quotient map. Then the image $p_{\mathcal{F}}(C) = p_{\mathcal{F}}(G) \sqcup \{p_{\mathcal{F}}((0, 0)), p_{\mathcal{F}}((1, 0))\}$ is a simple curve of a closed interval. The set difference $B - p_{\mathcal{F}}(C)$ consists of two disks that are homeomorphic to $[0, 1] \times (0, 2]$. Therefore the set difference $\text{int } B - p_{\mathcal{F}}(C)$ consists of two open disks that are homeomorphic to $(0, 1) \times (0, 2)$. Since the restriction $p_{\mathcal{F}}|_{\mathbb{R}^2 - C}$ is identical, the set difference $B - C$ are homeomorphic to $B - p_{\mathcal{F}}(C)$ and so consists of two disks that are homeomorphic to $[0, 1] \times (0, 2]$. Moreover, the set difference $\text{int } B - C$ consists of two open disks. \square

Let \mathbb{S}^2 be the one point compactification $\mathbb{R}^2 \sqcup \{\infty\}$ of \mathbb{R}^2 , $J_0 := (\mathbb{R} - [0, 1]) \times \{0\}$ the union of two intervals, and $\gamma := C \sqcup J_0 \sqcup \{\infty\} = G \sqcup \partial_{\mathfrak{h}} \mathbb{D} \sqcup$

$J_0 \sqcup \{\infty\}$. A subset is **annular** if it is homeomorphic to an annulus. A nonempty metrizable space is a **continuum** if it is compact and connected. A continuum Y contained in a surface S is **annular** if there is an open annular neighborhood A of Y such that $A - Y$ consists of two open annuli. We have the following observations.

Lemma 6.4. *The union γ is an annular continuum.*

Proof. Lemma 6.3 implies that the complement $\mathbb{S}^2 - \gamma = \mathbb{R}^2 - C \sqcup J_0$ is the disjoint union of two open disks. Removing two points, we can obtain there is an open annular neighborhood A of γ such that $A - \gamma$ consists of two open annuli. \square

Lemma 6.5. *Let $\mathbb{R}^2/\mathcal{F}_C$ be the resulting space collapsing the continuum $C = G \sqcup \partial_{\mathbb{H}}\mathbb{D}$ into a singleton and $p_{\mathcal{F}_C}: \mathbb{R}^2 \rightarrow \mathbb{R}^2/\mathcal{F}_C$ be the quotient map. Then the resulting space $\mathbb{R}^2/\mathcal{F}_C$ is a plane and the quotient map $p_{\mathcal{F}_C}$ is continuous.*

Proof. Let \mathcal{F} be a decomposition of \mathbb{R}^2 by $\mathcal{F} := \{\{p\} \mid p \in \mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D}\} \sqcup \{\{0\} \times [-1, 1], \{1\} \times [-1, 1]\}$ as in the proof of Lemma 6.3. By Moore's theorem (cf. p.3 in [16]), the quotient space \mathbb{R}^2/\mathcal{F} is homeomorphic to \mathbb{R}^2 . Let $p_{\mathcal{F}}: \mathbb{R}^2 \rightarrow \mathbb{R}^2/\mathcal{F}$ be the quotient map. Then the image $p_{\mathcal{F}}(C) = p_{\mathcal{F}}(G) \sqcup \{p_{\mathcal{F}}((0, 0)), p_{\mathcal{F}}((1, 0))\}$ is a simple curve of a closed interval. Let \mathcal{F}_C be a decomposition of \mathbb{R}^2 by $\mathcal{F}_C := \{\{p\} \mid p \in \mathbb{R}^2 - C\} \sqcup \{C\}$. Then the image $p_{\mathcal{F}}(\mathcal{F}_C)$ is a decomposition on a plane \mathbb{R}^2/\mathcal{F} consisting of the closed interval $p_{\mathcal{F}}(C)$ and points. Define the quotient space $\mathbb{R}^2/\sim_{\mathcal{F}_C}$ by $p \sim_{\mathcal{F}_C} q$ if there is an element of \mathcal{F}_C containing p and q . Let $q: \mathbb{R}^2/\mathcal{F} \rightarrow \mathbb{R}^2/\mathcal{F}_C$ be the quotient map by collapsing the closed interval $p_{\mathcal{F}}(C)$ into a singleton. By Moore's theorem (cf. p.3 in [16]), the quotient space $\mathbb{R}^2/\mathcal{F}_C$ is homeomorphic to \mathbb{R}^2 and the composition $p_{\mathcal{F}_C} := q \circ p_{\mathcal{F}}: \mathbb{R}^2 \rightarrow \mathbb{R}^2/\mathcal{F}_C$ is continuous and is the quotient map. \square

Let $\mathbb{B} := [-1, 2] \times [-5, 5]$ be a closed square containing B , $I_- := [-1, 0) \times \{0\} \subset J_0$ an interval, and $I_+ := (1, 2] \times \{0\} \subset J_0$ an interval. Put $I := C \sqcup I_- \sqcup I_+$. Then we have the following vector field.

Lemma 6.6. *There is a C^∞ vector field X on \mathbb{R}^2 satisfying the following properties:*

- (1) *The square \mathbb{B} is a flow box with respect to X .*
- (2) *The restriction $X|_{\mathbb{R}^2 - \mathbb{B}}$ is $(1, 0)$.*
- (3) *Subsets G , $(-\infty, 0) \times \{0\}$ and $(1, \infty) \times \{0\}$ of \mathbb{R}^2 are orbits of X .*
- (4) *The set $\text{Sing}(X)$ of critical points of X is $\partial_{\mathbb{H}}\mathbb{D} = \{0, 1\} \times [-1, 1]$.*
- (5) *Each of the positive and negative orbits of any points in $\mathbb{R}^2 - (G \sqcup \partial_{\mathbb{H}}\mathbb{D} \sqcup J_0)$ is neither singular nor periodic but is unbounded and closed as subsets.*
- (6) *For any convergence sequence $(y_n)_{n \in \mathbb{Z}_{\geq 0}}$ of non-zero numbers y_n to 0, there is a sequence $(t_n)_{n \in \mathbb{Z}_{\geq 0}}$ such that the sequence $v_X(t_n, (-1, y_n))_{n \in \mathbb{Z}_{\geq 0}}$ converges to a point in G , where v_X is the flow generated by X .*




FIGURE 14. A bump function.

Proof. Define a C^∞ bump function $\varphi: \mathbb{R} \rightarrow [0, 1]$ with $\varphi^{-1}(0) = (-\infty, 1/3]$ and $\varphi^{-1}(1) = [2/3, \infty)$ such that φ is increasing on $[1/3, 2/3]$ as in Figure 14. Put $C_\infty = C'_\infty := \{(x, \cos x) \mid x \in \mathbb{R}\}$ and $D_0 := \mathbb{R} \times (\mathbb{R} - [-3, 3])$. Let $f_0: \mathbb{R} \rightarrow \{3\}$ be a constant map and $g_0: \mathbb{R} \rightarrow \{-3\}$ a constant map.

For any positive integer $n \in \mathbb{Z}_{>0}$, define a C^∞ even function $f_n: \mathbb{R} \rightarrow [-1 + 1/n, 1 + 1/n]$ as follows:

$$f_n(x) := \begin{cases} 1 + \frac{1}{n} & \text{for } x \in \mathbb{R} - [-2\pi n, 2\pi n] \\ \cos x + \frac{1}{n} & \text{for } x \in [-2\pi n + \pi, 2\pi n - \pi] \\ \varphi\left(\frac{|x| - \pi(2n-1)}{\pi}\right) (1 - \cos x) + \cos x + \frac{1}{n} & \text{for } x \in (-2\pi n, -\pi(2n-1)) \sqcup (\pi(2n-1), 2\pi n) \end{cases}$$

Denote by C_n the graph $\{(x, f_n(x)) \mid x \in \mathbb{R}\}$ of f_n for any $n \in \mathbb{Z}_{\geq 0}$. For any point $x \in (-2\pi n, -\pi(2n-1)) \sqcup (\pi(2n-1), 2\pi n)$, we have the following relation:

$$\cos(x) + \frac{1}{n} \leq \varphi\left(\frac{|x| - \pi(2n-1)}{\pi}\right) (1 - \cos x) + \cos x + \frac{1}{n} \leq 1 + \frac{1}{n}$$

Then the family $(C_n)_{n \in \mathbb{Z}_{\geq 0} \sqcup \{\infty\}}$ are pairwise disjoint. Let D_n be the connected component of $\mathbb{R}^2 - \bigsqcup_{n \in \mathbb{Z}_{>0} \sqcup \{\infty\}} C_n$ whose boundary is the union of $C_{n-1} \sqcup C_n$.

Similarly, for any positive integer $n \in \mathbb{Z}_{>0}$, define a C^∞ even function $g_n: \mathbb{R} \rightarrow [-1 - 1/n, 1 - 1/n]$ as follows:

$$g_n(x) := \begin{cases} -1 - \frac{1}{n} & \text{for } x \in \mathbb{R} - [-2\pi n + \pi, 2\pi n - \pi] \\ \cos x - \frac{1}{n} & \text{for } x \in [-2\pi(n-1), 2\pi(n-1)] \\ \varphi\left(\frac{|x| - 2\pi(n-1)}{\pi}\right) (-1 - \cos x) + \cos x - \frac{1}{n} & \text{for } x \in (-2\pi n + \pi, -2\pi(n-1)) \sqcup (2\pi(n-1), 2\pi n - \pi) \end{cases}$$

Denote by C'_n the graph $\{(x, g_n(x)) \mid x \in \mathbb{R}\}$ of g_n for any $n \in \mathbb{Z}_{\geq 0}$. For any point $x \in (-2\pi n + \pi, -2\pi(n-1)) \sqcup (2\pi(n-1), 2\pi n - \pi)$, we have the following relation:

$$-1 - \frac{1}{n} \leq \varphi\left(\frac{|x| - 2\pi(n-1)}{\pi}\right) (-1 - \cos x) + \cos x - \frac{1}{n} \leq \cos(x) - \frac{1}{n}$$

Then the family $(C'_n)_{n \in \mathbb{Z}_{\geq 0} \sqcup \{\infty\}}$ are pairwise disjoint. Let D_{-n} be the connected component of $\mathbb{R}^2 - \bigsqcup_{n \in \mathbb{Z}_{>0} \sqcup \{\infty\}} C'_n$ whose boundary is the union of $C'_{n-1} \sqcup C'_n$. Then $\mathbb{R}^2 = \bigsqcup_{n \in \mathbb{Z}} D_n \sqcup C_\infty \sqcup \bigsqcup_{n \in \mathbb{Z}_{\geq 0}} C_n \sqcup C'_n$.

Define non-singular vector fields X_0 on \mathbb{R}^2 as follows:

$$X_0(x, y) := \begin{cases} (1, 0) & \text{for } (x, y) \in D_0 \\ (1, f'_\infty(x)) = (1, \sin(x)) & \text{for } (x, y) \in C_\infty \\ (1, f'_n(x)) & \text{for } (x, y) \in C_n \\ (1, g'_n(x)) & \text{for } (x, y) \in C'_n \\ \left(1, f'_n(x) + \varphi\left(\frac{y - f_n(x)}{f_{n-1}(x) - f_n(x)}\right)(f'_{n-1}(x) - f'_n(x))\right) & \text{for } (x, y) \in D_n \ (n > 0) \\ \left(1, g'_n(x) + \varphi\left(\frac{g_n(x) - y}{g_n(x) - g_{n-1}(x)}\right)(g'_{n-1}(x) - g'_n(x))\right) & \text{for } (x, y) \in D_n \ (n < 0) \end{cases}$$

By construction, for any point $p = (x, y) \in \bigsqcup_{n \in \mathbb{Z}_{\geq 0} \sqcup \{\infty\}} C_n$, we have a small neighborhood U_p of p such that $X|_{U_p} = (1, f'_n(x))$. Similarly, for any point $p = (x, y) \in \bigsqcup_{n \in \mathbb{Z}_{>0} \sqcup \{\infty\}} C'_n$, we have a small neighborhood U_p of p such that $X_0|_{U_p} = (1, g'_n(x))$. This means that X_0 is a C^∞ vector field on \mathbb{R}^2 . Take a diffeomorphism $h: (0, 1) \times [-4, 4] \rightarrow \mathbb{R} \times [-4, 4]$ by $h(x, y) = (h_0(x), y) = (\tan(\pi(x - 1/2)), y)$. Denote by G the inverse image $h^{-1}(C_\infty)$. The pushforward $X_1 := (h^{-1})_*(X_0|_{\mathbb{R} \times [-4, 4]})$ is a non-singular C^∞ vector field on $B_0 := (0, 1) \times [-4, 4]$. Define C^∞ functions $\phi_1, \phi_2: \mathbb{R} \rightarrow [0, 1]$ with

$$\phi_1^{-1}(0) = \mathbb{R} - (0, 1), \quad \phi_1^{-1}(1) = [1/3, 2/3],$$

$$\phi_2^{-1}(0) = \mathbb{R} - (-4, 4), \quad \text{and } \phi_2^{-1}(1) = [-3, 3]$$

such that ϕ_1 (resp. ϕ_2) is increasing on $[0, 1/3]$ (resp. $[-4, -3]$) and decreasing on $[2/3, 1]$ (resp. $[3, 4]$). Define a C^∞ vector field X_2 on $\mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D}$ as follows:

$$X_2(x, y) := \begin{cases} \phi_1(x)\phi_2(y)X_1(x, y)/|X_1(x, y)| & \text{for } (x, y) \in B_0 \\ 0 & \text{otherwise} \end{cases}$$

Define a C^∞ function $\phi_3: \mathbb{R} \rightarrow [0, 1]$ with

$$\phi_3^{-1}(0) = [0, 1] \text{ and } \phi_3^{-1}(1) = \mathbb{R} - [-1/3, 4/3]$$

such that ϕ_3 is decreasing on $[-1/3, 0]$ and increasing on $[1, 4/3]$. Write $B_{-1} := [-1/3, 0) \times [-1, 1]$ and $B_1 := (1, 4/3] \times [-1, 1]$. Define $B'_{-1} := \{(x, y) \mid x \in [-1/2, 0), y \in [-1 - \phi_3(x), 1 + \phi_3(x)]\}$ and $B'_1 := \{(x, y) \mid x \in (1, 3/2], y \in [-1 - \phi_3(x), 1 + \phi_3(x)]\}$. Then B'_{-1} (resp. B'_1) is a closed neighborhood of B_{-1} (resp. B_1) on $\mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D}$. Therefore there are a C^∞




FIGURE 15. Orbits with respect to X and supports of vector fields X_2 , Y , and Z .

function $f_Y: \mathbb{R}^2 - \partial_{\mathfrak{h}}\mathbb{D} \rightarrow [0, 1]$ and a C^∞ vector field $Y = (0, f_Y(x, y)y)$ on $\mathbb{R}^2 - \partial_{\mathfrak{h}}\mathbb{D}$ such that

$$Y|_{B_{-1} \sqcup B_1} = (0, \phi_3(x)y) \text{ and } Y^{-1}(0) = \mathbb{R}^2 - (\partial_{\mathfrak{h}}\mathbb{D} \sqcup B'_1 \sqcup B'_{-1}).$$

There is a closed neighborhood $U \subseteq h^{-1}(X_0^{-1}((1, 0))) \cup ([-1, 0] \sqcup [1, 2]) \times ([-5, -1] \sqcup [1, 5]) \subseteq \mathbb{B}$ of $\partial B_0 - \partial_{\mathfrak{h}}\mathbb{D}$ on $\mathbb{R}^2 - \partial_{\mathfrak{h}}\mathbb{D}$ with $U \cap (B'_1 \sqcup B'_{-1}) = \emptyset$ and there are a small positive number $\varepsilon \in (0, 1/3)$ and a C^∞ function $\phi_4: \mathbb{R}^2 \rightarrow [0, 1]$ with

$$((0, 1) \times [-7/2, 7/2]) \setminus \text{int } U = \phi_4^{-1}(0) \text{ and}$$

$$\mathbb{R}^2 - ((-2/3, 5/3) \times (-9/2, 9/2)) \subset \phi_4^{-1}(1)$$

such that $\phi_4|_{(-\varepsilon, 0) \times [-1, 1]}(x, y) \leq -\phi_3(x)x$ on any $x \in (-\varepsilon, 0)$ and that $\phi_4|_{(0, \varepsilon) \times [-1, 1]}(x, y) \leq \phi_3(x)(x - 1)$ on any $x \in (0, \varepsilon)$. Define C^∞ vector fields Z and X on $\mathbb{R}^2 - \partial_{\mathfrak{h}}\mathbb{D}$ by $Z(x, y) = (\phi_4(x, y), 0)$ and $X := X_2 + Y + Z$. Then

$$X(x, y) = \begin{cases} X_2(x, y) = \phi_1(x)\phi_2(y)X_1(x, y)/|X_1(x, y)| & \text{for } (x, y) \in B_0 \setminus U \\ X_2(x, y) + Z(x, y) = (\phi_1(x)\phi_2(y) + \phi_4(x, y), 0) & \text{for } (x, y) \in (\mathbb{R}^2 - (B_0 \sqcup B'_{-1} \sqcup B'_1 \sqcup \partial_{\mathfrak{h}}\mathbb{D})) \cup U \\ Y(x, y) + Z(x, y) = (\phi_4(x, y), f_Y(x, y)y) & \text{for } (x, y) \in B'_{-1} \sqcup B'_1 \end{cases}$$

and $X(x, y)|_{B_{-1} \sqcup B_1} = (\phi_4(x, y), \phi_3(x)y)$ as in Figure 15. Adding $\partial_{\mathfrak{h}}\mathbb{D}$ as critical points, we extend X into a vector field on \mathbb{R}^2 , also denoted by X . Let v_X be the \mathbb{R} -action generated by X on \mathbb{R}^2 . We will show that X is desired.

We claim that v_X is continuous on \mathbb{R}^2 . Since $\partial_{\mathbb{H}}\mathbb{D}$ is compact, for any $\varepsilon_0 > 0$, there is a neighborhood V of $\partial_{\mathbb{H}}\mathbb{D}$ such that $\sup_{p \in V} |X(p)| < \varepsilon_0$. This implies the continuity of v_X at any points in the closed subset $\mathbb{R} \times \partial_{\mathbb{H}}\mathbb{D} \subseteq \mathbb{R} \times \text{Sing}(v)$. On the other hand, by the openness of $\mathbb{R} \times (\mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D})$ and the invariance of $\mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D}$, the \mathbb{R} -action v_X is continuous at any points in the complement $\mathbb{R} \times (\mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D})$. This means that v_X is continuous on \mathbb{R}^2 .

We claim that the assertion (4) holds. Indeed, we have that $X(x, y) = X_2(x, y) \neq 0$ on $B_0 \setminus U$, $X(x, y) = X_2(x, y) + Z(x, y) \neq 0$ on $(\mathbb{R}^2 - (B_0 \sqcup B'_{-1} \sqcup B'_1 \sqcup \partial_{\mathbb{H}}\mathbb{D})) \cup U$, and the first component of $X(x, y)$ is $Z(x, y) = \phi_4(x, y) \neq 0$ on $B'_{-1} \sqcup B'_1$. This means that $X(x, y) \neq 0$ on $\mathbb{R}^2 - \partial_{\mathbb{H}}\mathbb{D}$.

We claim that the assertions (1) and (2) hold. Indeed, since $U \subseteq \mathbb{B} = [-1, 2] \times [-5, 5]$, $B_0 = (0, 1) \times [-4, 4]$, $B_{-1} = [-1/3, 0] \times [-1, 1] \subset B'_{-1}$, and $B_1 = (1, 4/3] \times [-1, 1] \subset B'_1$, we have $B_0 \cup (B'_{-1} \sqcup B'_1 \sqcup U) \subset \mathbb{B}$. Then $X(x, y) = Z(x, y) = (\phi_4(x, y), 0) = (1, 0)$ on $\mathbb{R}^2 - ((-2/3, 5/3) \times (-9/2, 9/2)) \subset \phi_4^{-1}(1)$. By $\mathbb{R}^2 - \text{int } \mathbb{B} = \mathbb{R}^2 - ((-1, 2) \times (-5, 5)) \subset \mathbb{R}^2 - ((-2/3, 5/3) \times (-9/2, 9/2))$, the closed square \mathbb{B} is a flow box with respect to X such that the restriction $X|_{\mathbb{R}^2 - \mathbb{B}}$ is $(1, 0)$.

Since \mathbb{B} can be identified with a flow box in a compact surface, by Gutierrez's smoothing theorem [22], we may assume that v is C^∞ and so is X .

We claim that the assertion (3) holds. Indeed, since G is an orbit of X_1 , from $X = X_2 = \phi_1(x)\phi_2(y)X_1(x, y)/|X_1(x, y)|$ on $B_0 \subset \mathbb{B} \setminus U$, the subset G is the orbit of X . By $X_2 = Y = 0$ on $J_0 = (\mathbb{R} - [0, 1]) \times \{0\}$, we obtain that $X(x, y) = Z(x, y) = (\phi_4(x, y), 0)$ and $\phi_4(x, y) > 0$ on J_0 . This means that subsets $(-\infty, 0) \times \{0\}$ and $(1, \infty) \times \{0\}$ are orbits of X respectively.

Finally, we show the assertions (5) and (6). On $[-1, 2] \times ([-5, -4] \sqcup [4, 5])$, the vector field $X = (\phi_4(x, y), 0)$ is non-singular and so the subset $[-1, 2] \times \{y_0\}$ for any $y_0 \in [-5, -4] \sqcup [4, 5]$ is an orbit arc for X .

We claim that the positive orbit in any point in $\{-1\} \times ([-5, 5] - \{0\}) \subset \partial\mathbb{B}$ intersects $\{0\} \times ([-5, 5] - [-1, 1])$. Indeed, fix a point $p_0 = (x_0, y_0) \in \{1\} \times ([-5, 5] - \{0\})$. Suppose that $y_0 > 1$. Then $[-1, 0] \times [y_0, 5]$ is compact and $Z(x, y) = (\phi_4(x, y), 0)$ is non-singular on the domain $[-1, 0] \times [y_0, 5]$. By $\text{supp}(Y) \subset ([-1, 0] \sqcup [1, 2]) \times [-2, 2]$, $Y(x, y) = (0, f_Y(x, y)y)$, and $Y(x, y)y \geq 0$ on $[-1, 0] \times [y_0, 2]$, the positive orbit $O^+((x_1, y_0))$ for any $x_1 \in [-1, 0]$ intersects $\{0\} \times [y_0, 5] \subset \{0\} \times (1, 5]$. In particular, the positive orbit $O^+(p_0)$ intersects $\{0\} \times [y_0, 5] \subset \{0\} \times (1, 5]$. Suppose that $y_0 = 1$. Then $X = (\phi_4(x, 1), 0)$ on $x \in [-1, -1/2]$, $X = (\phi_4(x, 1), f_Y(x, 1))$ on $x \in [-1/2, 0]$, and $f_Y(x, 1) > 0$ on $x \in (-1/2, 0)$. Therefore $O^+(p)$ intersects $[-1, 0] \times (1, 5]$ and so $\{0\} \times (1, 5]$ because of the previous argument. Suppose that $y_0 \in (0, 1]$. Then the positive orbit $O^+(p)$ intersects either $[-1, 0] \times (1, 5]$ or $(-\varepsilon, 0) \times (0, 1)$. If $O^+(p)$ intersects $[-1, 0] \times (1, 5]$, then the previous argument implies that the positive orbit $O^+(p)$ intersects $\{0\} \times [y_0, 5] \subset \{0\} \times (1, 5]$. Thus we may assume that $O^+(p)$ intersects $(-\varepsilon, 0) \times (0, 1)$. By definition, we have that $Y(x, y) = (0, \phi_3(x)y)$ and $\phi_4(x, y) \leq -\phi_3(x)x$ on $(-\varepsilon, 0) \times (0, 1) \subset B_{-1}$. Then $X(x, y) = (\phi_4(x, y), \phi_3(x)y) = \phi_3(x)(-x, y) + (\phi_4(x, y) + \phi_3(x)x, 0)$ and $\phi_4(x, y) + \phi_3(x)x \leq 0$ on $(-\varepsilon, 0) \times (0, 1) \subset B_{-1}$.

Since the orbit of p with respect to the vector field Y' on B_{-1} defined by $Y'(x, y) = (-x, y)$ intersects a point (x', y') in the horizontal boundary $[-1/3, 0] \times \{1\}$, from $\phi_4(x, y) \geq 0$, the orbit $O^+(p)$ with respect to the vector field $X|_{(-\varepsilon, 0) \times (0, 1)} = (\phi_4(x, y), \phi_3(x)y) = \phi_3(x)(-x, y) + (\phi_4(x, y) + \phi_3(x)x, 0)$ intersects a point in $[-1/3, x'] \times \{1\}$. From the previous argument, the orbit $O^+(p)$ intersects $\{0\} \times (1, 5]$. By symmetry, if $y_0 < 0$, then the orbit $O^+(p)$ intersects $\{0\} \times [-5, -1)$.

By symmetry, the negative orbit in any point in $\{2\} \times ([-5, 5] - \{0\}) \subset \partial\mathbb{B}$ intersects $\{1\} \times ([-5, 5] - [-1, 1])$. By construction, the positive (resp. negative) orbit of any point in $B_0 - G$ with respect to X intersects U and so $\{1\} \times ([-5, 5] - [-1, 1])$ (resp. $\{0\} \times ([-5, 5] - [-1, 1])$). Therefore each of the positive and negative orbits of any points in $\mathbb{B} - (G \sqcup \partial_{\text{in}}\mathbb{D} \sqcup I_- \sqcup I_+)$ is neither singular nor periodic but is unbounded. This implies the assertion (5).

Fix a convergence sequence $(y_n)_{n \in \mathbb{Z}_{\geq 0}}$ of non-zero numbers y_n to 0. Denote by z_n the point with $\{z_n\} = O^+(-1, y_n) \cap \{0\} \times (\mathbb{R} - [-1, 1])$. By construction, the sequence $(z_n)_{n \in \mathbb{Z}_{\geq 0}}$ converges to either $(0, 1)$ or $(0, -1)$. Denote by w_n the point with $\{w_n\} = O^+(z_n) \cap \{1/2\} \times \mathbb{R}$. By construction, the sequence $(w_n)_{n \in \mathbb{Z}_{\geq 0}}$ converges to a point in G . This implies the assertion (6). \square

The previous lemma implies the following statement.

Lemma 6.7. *Let v_X be the flow generated by the vector field X as in Lemma 6.6 and $p_{\mathcal{F}_C} : \mathbb{R}^2 \rightarrow \mathbb{R}^2/\mathcal{F}_C$ be the quotient map as in Lemma 6.5. Then the following statements hold:*

(1) *The mapping $v : \mathbb{R} \times \mathbb{R}^2/\mathcal{F}_C \rightarrow \mathbb{R}^2/\mathcal{F}_C$ defined by*

$$v(t, p) := p_{\mathcal{F}_C}(v_X(t, p_{\mathcal{F}_C}^{-1}(p)))$$

is well-defined.

(2) *The mapping v is semi-conjugate to v_X via $p_{\mathcal{F}_C}$.*

(3) *The mapping v is the resulting flow of a unit vector field $(1, 0)$ on \mathbb{R}^2 by replacing a non-singular point with a singular point (as in Lemma 6.1).*

Proof. First, we show the well-definedness of v . Fix $y \in \mathbb{R}^2/\mathcal{F}_C$. Suppose that $y \in p_{\mathcal{F}_C}(C)$. Then $p_{\mathcal{F}_C}^{-1}(y) = G$ and so $v_X(t, p_{\mathcal{F}_C}^{-1}(y)) = v_X(t, G) = G$. Therefore $p_{\mathcal{F}_C}(v_X(t, p_{\mathcal{F}_C}^{-1}(y))) = p_{\mathcal{F}_C}(G) = y$. This means that y is a singular point of v . Suppose that $y \notin p_{\mathcal{F}_C}(C)$. Since $p_{\mathcal{F}_C}^{-1}(x) = y$, we have $v(t, y) = p_{\mathcal{F}_C}(v_X(t, p_{\mathcal{F}_C}^{-1}(y))) = p_{\mathcal{F}_C}(v_X(t, y)) = v_X(t, y)$. This means that v is well-defined.

Since the quotient map $p_{\mathcal{F}_C}$ is continuous and closed, the map $1_{\mathbb{R}} \times p_{\mathcal{F}_C} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}^2/\mathcal{F}_C$ defined by $1_{\mathbb{R}} \times p_{\mathcal{F}_C}(t, x) = (t, p_{\mathcal{F}_C}(x))$ is a quotient map and so is closed. Then $v(t, x) = p_{\mathcal{F}_C} \circ v_X((1_{\mathbb{R}} \times p_{\mathcal{F}_C})^{-1}(t, x))$. For any closed subset $A \subseteq \mathbb{R}^2/\mathcal{F}_C$, the inverse image $v^{-1}(A) = 1_{\mathbb{R}} \times p_{\mathcal{F}_C}(v_X^{-1}(p_{\mathcal{F}_C}^{-1}(A)))$ is closed. This means that the \mathbb{R} -action v is a flow. By

construction, for any $(t, x) \in \mathbb{R} \times \mathbb{R}^2$, we obtain $v(t, p_{\mathcal{F}_C}(x)) = p_{\mathcal{F}_C}(v_X(t, x))$. This implies the semi-conjugacy. \square

We demonstrate Theorem 6.2 as follows.

Proof of Theorem 6.2. Let v be a flow on a surface S with an ω -limit set ω containing non-singular point x and with a point $y \in S - \omega$ satisfying $\omega(y) = \omega$. Take any closed trivial flow box B with $x \in \text{int } B$. Identify x with 0 and the connected component of $O(x) \cap B$ containing x with $[-1, 2] \times \{0\}$. Replacing B with the flow box \mathbb{B} constructed in Lemma 6.6, denote by w the resulting flow. Lemma 6.7 implies that the resulting flow v_x of v by replacing x with a singular point is topologically equivalent to a flow which is topologically semi-conjugate to w by collapsing the closed invariant subset C of $\omega_w(y)$ into a singleton. Then the restriction $v|_{S-\omega}$ is topologically equivalent to the restriction $w|_{S-\omega_w(y)}$.

Suppose that the ω -limit set ω is not locally dense. Then the orbit arc $[-1, 2] \times \{0\}$ is a connected component of $\omega \cap B$ and so the ω -limit set $\omega_w(y)$ is not locally dense. By the assertion (6) in Lemma 6.6, the ω -limit set $\omega_w(y)$ contains C . Therefore the disjoint union $([-1, 0] \times \{0\}) \sqcup C \sqcup ((1, 2] \times \{0\})$ is a connected component of $\omega_w(y) \cap B$. Since C is not arcwise-connected, neither is the ω -limit set $\omega_w(y)$. \square

7. EXAMPLES

We describe some kinds of ω -limit sets that appear in Theorem A.

7.1. Quasi-Q-sets that are not Q-sets. We show that there is a toral flow with a non-locally-dense quasi-Q-set which is not a Q-set as follows.

Lemma 7.1. *There are a toral flow v_φ and a point z whose ω -limit set is a non-locally-dense quasi-Q-set but not a Q-set such that $\overline{\text{Cl}(v_\varphi)} \neq \Omega(v_\varphi)$, where $\Omega(v_\varphi)$ is the non-wandering set of v_φ .*

Proof. Consider a Denjoy diffeomorphism $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ with an exceptional minimal set \mathcal{C} . Let v_f be the suspension of f on the torus $\mathbb{T}^2 := (\mathbb{S}^1 \times \mathbb{R}) / (x, r) \sim (f(x), r + 1)$ and \mathcal{M} the minimal set of v_f .

We will replace the minimal set \mathcal{M} of v_f with a union of singular points and separatrices of the resulting flow v_φ as follows. Fix a bump function $\varphi : \mathbb{T}^2 \rightarrow \mathbb{R}_{\geq 0}$ with $\varphi^{-1}(0) = \hat{\mathcal{C}}$, where $\hat{\mathcal{C}} := \mathcal{M} \cap (\mathbb{S}^1 \times \{1/2\})$ is a lift of \mathcal{M} . Let X be the continuous vector field generating v_f on the mapping torus \mathbb{T}^2 . Since $\hat{\mathcal{C}}$ are covered by finitely many trivial flow boxes, there is an open neighborhood U of $\hat{\mathcal{C}}$ such that the restriction $X|_U$ can be considered as the restriction of a continuous vector field Y on a sphere. By Gutierrez's smoothing theorem [22], we may assume that the vector field Y is C^∞ and so does $X|_U$. Since every closed subset of any paracompact C^∞ manifold is a zero set of some C^∞ function on it, take a C^∞ bump function $\varphi : \mathbb{T}^2 \rightarrow [0, 1]$ with $\varphi^{-1}(0) = \hat{\mathcal{C}}$ and $\varphi|_{\mathbb{T}^2 - U} = 1$. Define a continuous vector field Z by $Z(p) := \varphi(p)X(p)$. Then the restriction $Z|_{\mathbb{T}^2 - U} = X|_{\mathbb{T}^2 - U}$ is non-singular




FIGURE 16. An ω -limit set which is a non-locally-connected quasi-circuit, which is neither the image of a circle nor a circuit.

and the restriction $Z|_U = X|_U$ is C^∞ . Therefore Z is locally Lipschitz continuous and so generates a flow v_φ with $\mathcal{M} = \text{Sing}(v_\varphi) \sqcup \{\text{separatrix of } v_\varphi\}$, $O_{v_\varphi}(p) = O_{v_f}(p)$, and $\omega_v(p) = \omega_{v_\varphi}(p) = \mathcal{M}$ for any point $p \in \mathbb{T}^2 - \mathcal{M}$. Then $\mathbb{T}^2 = \text{Sing}(v_\varphi) \sqcup P(v_\varphi)$ and $\overline{\text{Cl}}(v_\varphi) = \text{Sing}(v_\varphi) = \hat{\mathcal{C}} \neq \mathcal{M} = \Omega(v_\varphi)$, where $P(v_\varphi)$ is the union of non-recurrent orbits of v_φ . \square

7.2. Quasi-circuit that is not circuits. Using the flow box in Lemma 6.6, we can construct a flow with a quasi-circuit that is not a circuit such that it consists of two non-recurrent orbits and two closed intervals contained in the singular point set as in Figure 16. In particular, the quasi-circuit is homeomorphic to the union of G and a curve from a point in G and to a point in G .

Note that Hastings constructed an attractor of a flow on \mathbb{R}^2 which is homeomorphic to a Warsaw circle (i.e. the disjoint union of the graph of the function $f: (0, 1/\pi] \rightarrow [-1, 1]$ by $f(x) = \sin 1/x$, the interval $\{0\} \times [-1, 1]$, and an open arc from $(0, -1)$ to $(1/\pi, 0)$) but is not an ω -limit set in [25, Example 3.3]. By a similar construction of the flow box in Lemma 6.6, we can construct a flow box as in Figure 17 and so a flow with a quasi-circuit that is homeomorphic to a Warsaw circle.

7.3. Non-locally connected subsets of singular points. We construct the following examples of flows.

Lemma 7.2. *There is a toral flow with an ω -limit set consisting of singular points which is a transversely Cantor set.*

Proof. Let be a non-recurrent orbit O whose ω -limit set is an exceptional minimal set \mathcal{M} in a Denjoy flow on a torus \mathbb{T}^2 . Consider the resulting flow of v_x by replacing a non-singular point $x \in \mathcal{M}$ with a singular point. Then $\text{Sing}(v_x) = \{x\}$ and $\overline{O(y)} = \mathcal{M}$ for any non-singular point y . By Gutierrez's smoothing theorem [22], we may assume that the flow v_x is C^∞ . Let X be the C^∞ vector field generating v_x . Since every closed subset of




FIGURE 17. A flow box with a non-arcwise-connected subset.

any paracompact C^∞ manifold is a zero set of some C^∞ function on it by using C^∞ bump functions and partitions of unity, take a C^∞ bump function $\varphi: \mathbb{T}^2 \rightarrow [0, 1]$ with $\varphi^{-1}(0) = \mathcal{M}$. Then the resulting flow v generated by the vector field φX is a C^∞ flow such that the ω -limit set of a non-singular point is $\mathcal{M} = \text{Sing}(v)$. This means that v is desired. \square

Lemma 7.3. *There is a toral C^∞ flow w with an ω -limit set which consists of two non-recurrent orbits and a subset of $\text{Sing}(w)$ which is a transversely Cantor set as a set.*

Proof. Let $X_1 := \varphi X$ be the vector field in the previous example. Then the complement $\mathbb{T}^2 - \mathcal{M}$ is a trivial flow box. Identify the flow box with a square $B := \mathbb{R} \times [-1/2, \pi + 1/2]$ on a chart by the embedding $f: B \rightarrow \mathbb{T}^2 - \mathcal{M}$. Replacing the embedding f of the square B in the torus if necessary, we may assume that the norms of any vector fields on B are no more than the norms of the push-forwards of the vector fields by f on the square $\mathbb{T}^2 - \mathcal{M}$. Define C^∞ bump functions $h, \psi: \mathbb{R} \rightarrow [0, 1]$ with $\psi^{-1}(0) = \mathbb{R} - [-1/2, \pi + 1/2]$ and $\psi^{-1}(1) = [0, 1]$ such that h is an even function which is strictly decreasing on $(0, \infty)$ and $\lim_{x \rightarrow \pm\infty} h(x) = 0$.

Define the following continuous vector field Y_0 on $(-\pi/2, \pi/2) \times [-1/2, \pi + 1/2]$ as follows:

$$Y_0(\theta, y) := \begin{cases} (-\cos \theta, 0) & \text{for } y \in (\pi, \pi + 1/2] \\ (\cos \theta, 0) & \text{for } y \in [-1/2, 0) \\ (\cos \theta \cos y, \sin \theta \sin y) & \text{for } (\theta, y) \in (-\pi/2, \pi/2) \times [0, \pi] \end{cases}$$

Notice that the restriction $Y_0|_{(-\pi/2, \pi/2) \times [0, \pi]}$ is a Taylor-Green vortex. Define the vector field Y_1 on B by $Y_1(x, y) := h(x)\psi(y)Y_0(\tan^{-1}(x), y)$. Then $Y_1(x, 0) = (\psi(y)\cos(\tan^{-1}(x)), 0) = \psi(0)\cos(\tan^{-1}(x))(1, 0)$ on the line $\mathbb{R} \times \{0\}$ and $Y_1(x, \pi) = (-\psi(y)\cos(\tan^{-1}(x)), 0) = \psi(y)\cos(\tan^{-1}(x))(-1, 0)$ on the line $\mathbb{R} \times \{\pi\}$. Since the restrictions $Y_1|_{\mathbb{R} \times ([-1/2, 0) \sqcup (\pi, \pi + 1/2])}$ and $Y_1|_{\mathbb{R} \times [0, \pi]}$ are C^∞ , the vector field Y_1 is locally Lipschitz continuous. Then the push-forward $Y_2 := f_* Y_1$ on $\mathbb{T}^2 - \mathcal{M}$ by f of the vector field Y_1 generates an \mathbb{R} -action v_{Y_2} on B as Figure 18. Since $|Y_1(x, y)| \geq |Y_2(f(x, y))|$ for any




FIGURE 18. A square with a flow.

point $(x, y) \in B$, by $\lim_{x \rightarrow \pm\infty} \max_{y \in [-1/2, \pi+1/2]} |Y_1(x, y)| = 0$, the induced vector field Y_2 can be extended to a continuous vector field Y_3 on \mathbb{T}^2 by $Y_3|_{\mathcal{M}} = 0$.

We claim that the resulting \mathbb{R} -action v_{Y_3} generated by Y_3 is a flow. Indeed, fix any number $T \in \mathbb{R}$, any positive number $\varepsilon > 0$, and any point $p \in \mathcal{M}$. Then there is a neighborhood U of p on \mathbb{T}^2 such that the intersection $V := U \cap f([-T, T] \times [-1/2, \pi+1/2])$ contains at most one connected component and the diameter of $f^{-1}(V)$ is less than ε , where the diameter of a subset A is the superior of distances between two points in A . This implies that the norm of the restriction $Y_1|_{f^{-1}(V)}$ is less than ε and so is one of $Y_3|_V$. This implies the continuity of the resulting \mathbb{R} -action v_{Y_3} .

By Gutierrez's smoothing theorem [22], we may assume that the flow v_{Y_3} is a desired C^∞ flow. \square

REFERENCES

- [1] M. Abate and F. Bianchi. A Poincaré-Bendixson theorem for meromorphic connections on Riemann surfaces. *Math. Z.*, 282(1-2):247–272, 2016.
- [2] M. Abate and F. Tovenà. Poincaré-Bendixson theorems for meromorphic connections and holomorphic homogeneous vector fields. *J. Differential Equations*, 251(9):2612–2684, 2011.
- [3] A. Andronov, E. Leontovich, I. Gordon, and A. Mayer. Qualitative theory of dynamic systems of the second order. *Moscow: Mir*, 1966.
- [4] S. Aranson and E. Zhuzhoma. Maier's theorems and geodesic laminations of surface flows. *Journal of Dynamical and Control Systems*, 2(4):557–582, 1996.
- [5] S. K. Aranson, G. R. Beliskii, and E. Zhuzhoma. *Introduction to the qualitative theory of dynamical systems on surfaces*. American Mathematical Society, 1996.
- [6] K. Athanassopoulos. A characterization of denjoy flows. *Bulletin of the London Mathematical Society*, 24(1):83–86, 1992.
- [7] I. Bendixson. Sur les courbes définies par des équations différentielles. *Acta Math.*, 24(1):1–88, 1901.
- [8] N. Bhatia, A. Lazer, and W. Leighton. Application of the Poincaré-Bendixson theorem. *Annali di Matematica Pura ed Applicata*, 73(1):27–32, 1966.
- [9] G. D. Birkhoff. *Dynamical systems*, volume 9. American Mathematical Soc., 1927.
- [10] E. Bonotto and M. Federson. Limit sets and the Poincaré-Bendixson theorem in impulsive semidynamical systems. *Journal of Differential Equations*, 244(9):2334–2349, 2008.
- [11] J. P. Boroński, J. Činč, and X.-C. Liu. Prime ends dynamics in parametrised families of rotational attractors. *J. Lond. Math. Soc. (2)*, 102(2):557–579, 2020.

- [12] N. Bourbaki. *General Topology: Chapters 1–4*, volume 18. Springer Science & Business Media, 2013.
- [13] A. Candel and L. Conlon. *Foliations. I*, volume 23 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2000.
- [14] T. Cherry. Topological properties of the solutions of ordinary differential equations. *American Journal of Mathematics*, 59(4):957–982, 1937.
- [15] K. Ciesielski. The Poincaré-Bendixson theorems for two-dimensional semiflows. *Topological Methods in Nonlinear Analysis*, 3(1):163–178, 1994.
- [16] R. J. Daverman. *Decompositions of manifolds*. Academic Press, 1986.
- [17] D. P. Demuner, M. Federson, and C. Gutierrez. The Poincaré-Bendixson Theorem on the Klein bottle for continuous vector fields. *Discrete & Continuous Dynamical Systems*, 25(2):495, 2009.
- [18] Z. Du, J. Liu, and Y. Ren. Traveling pulse solutions of a generalized keller-segel system with small cell diffusion via a geometric approach. *Journal of Differential Equations*, 270:1019–1042, 2021.
- [19] J. G. Espín Buendía and V. Jiménez López. Some remarks on the ω -limit sets for plane, sphere and projective plane analytic flows. *Qual. Theory Dyn. Syst.*, 16(2):293–298, 2017.
- [20] J. G. Espín Buendía and V. Jiménez Lopéz. A topological characterization of the ω -limit sets of analytic vector fields on open subsets of the sphere. *Discrete Contin. Dyn. Syst. Ser. B*, 24(3):1143–1173, 2019.
- [21] C. Gardiner. The structure of flows exhibiting nontrivial recurrence on two-dimensional manifolds. *Journal of differential equations*, 57(1):138–158, 1985.
- [22] C. Gutiérrez. Structural stability for flows on the torus with a cross-cap. *Transactions of the American Mathematical Society*, 241:311–320, 1978.
- [23] C. Gutiérrez. Smoothing continuous flows on two-manifolds and recurrences. *Ergodic Theory and dynamical systems*, 6(1):17–44, 1986.
- [24] O. Hájek. *Dynamical systems in the plane*. Academic Press, London-New York, 1968.
- [25] H. M. Hastings. A higher-dimensional Poincaré-Bendixson theorem. *Glasnik Mat. Ser. III*, 14(34)(2):263–268, 1979.
- [26] G. Hector and U. Hirsch. *Introduction to the geometry of foliations. Part A*, volume 1 of *Aspects of Mathematics*. Friedr. Vieweg & Sohn, Braunschweig, 1981. Foliations on compact surfaces, fundamentals for arbitrary codimension, and holonomy.
- [27] J. Hounie. Minimal sets of families of vector fields on compact surfaces. *Journal of Differential Geometry*, 16(4):739–744, 1981.
- [28] A. Koropecki and A. Passeggi. A Poincaré-Bendixson theorem for translation lines and applications to prime ends. *Commentarii Mathematici Helvetici*, 94(1):141–183, 2019.
- [29] G. Levitt. Feuilletages des surfaces. *Ann. Inst. Fourier (Grenoble)*, 32(2):x, 179–217, 1982.
- [30] G. Levitt and H. Rosenberg. Differentiability and topology of labyrinths in the disc and annulus. *Topology*, 26(2):173–186, 1987.
- [31] V. J. López and J. Llibre. A topological characterization of the ω -limit sets for analytic flows on the plane, the sphere and the projective plane. *Advances in Mathematics*, 216(2):677–710, 2007.
- [32] V. J. López and G. S. López. Accumulation points of nonrecurrent orbits of surface flows. *Topology and its Applications*, 137(1-3):187–194, 2004.
- [33] N. Markley and M. Vanderschoot. Recurrent orbits for flows on surfaces. *Journal of Differential Equations*, 244(5):1210–1229, 2008.
- [34] N. G. Markley. The Poincaré-Bendixson theorem for the Klein bottle. *Transactions of the American Mathematical Society*, pages 159–165, 1969.
- [35] D. Martí-Pete, L. Rempe, and J. Waterman. Eremenko’s conjecture, wandering lakes of wada, and maverick points. *arXiv preprint arXiv:2108.10256*, 2021.

- [36] H. Marzougui. Structure des feuilles sur les surfaces ouvertes. *C. R. Acad. Sci. Paris Sér. I Math.*, 323(2):185–188, 1996.
- [37] H. Marzougui. Structure of foliations on 2-manifolds. *Illinois Journal of Mathematics*, 42(3):398–405, 1998.
- [38] H. Marzougui. Flows with infinite singularities on closed two-manifolds. *Journal of dynamical and control systems*, 6(4):461–476, 2000.
- [39] A. Mayer. Trajectories on the closed orientable surfaces. *Rec. Math.[Mat. Sbornik] NS*, 12(54):71–84, 1943.
- [40] J. Milnor. *Dynamics in One Complex Variable.(AM-160):(AM-160)-*, volume 160. Princeton University Press, 2011.
- [41] I. Nikolaev and E. Zhuzhoma. *Flows on 2-dimensional manifolds: an overview*. Number 1705. Springer Science & Business Media, 1999.
- [42] J. Plante. A generalization of the Poincaré-Bendixson theorem for foliations of codimension one. *Topology*, 12(2):177–181, 1973.
- [43] H. Poincaré. Sur les courbes définies par les equations differentielles. *J. Math. Pures Appl.*, 3(8):251–286, 1882.
- [44] A. Pokrovskii, A. Pokrovskiy, and A. Zhezherun. A corollary of the Poincaré–Bendixson theorem and periodic canards. *Journal of Differential Equations*, 247(12):3283–3294, 2009.
- [45] I. Richards. On the classification of noncompact surfaces. *Transactions of the American Mathematical Society*, 106(2):259–269, 1963.
- [46] R. Roussarie. A topological study of planar vector field singularities. *Discrete Contin. Dyn. Syst.*, 40(9):5217–5245, 2020.
- [47] R. Roussarie. Correction to: Some applications of the Poincaré-Bendixson theorem. *Qual. Theory Dyn. Syst.*, 20(3):Paper No. 68, 1, 2021.
- [48] R. Roussarie. Some Applications of the Poincaré-Bendixson Theorem. *Qualitative Theory of Dynamical Systems*, 20(64):17 pp., 2021.
- [49] A. J. Schwartz. A generalization of a Poincaré-Bendixson theorem to closed two-dimensional manifolds. *American Journal of Mathematics*, pages 453–458, 1963.
- [50] M. Torhorst. Über den Rand der einfach zusammenhängenden ebenen Gebiete. *Math. Z.*, 9(1-2):44–65, 1921.
- [51] M. Torhorst. Über den Rand der einfach zusammenhängenden ebenen Gebiete. *Mathematische Zeitschrift*, 9(1):44–65, 1921.
- [52] M. Vanderschoot. Local connectivity of limit sets. *The Rocky Mountain journal of mathematics*, pages 1525–1538, 2003.
- [53] R. Vinograd. On the limit behavior of an unbounded integral curve, moskov. *Gos. Univ. Uc. Zap. Mat.*, 155(5):94–136, 1952.
- [54] K. Yano et al. Asymptotic Cycles on Two Dimensional Manifolds. In *Foliations*, pages 359–377. Mathematical Society of Japan, 1985.
- [55] T. Yokoyama. A topological characterization for non-wandering surface flows. *Proceedings of the American Mathematical Society*, 144(1):315–323, 2016.
- [56] T. Yokoyama. Properness of foliations. *Topology and its Applications*, 254:171–175, 2019.
- [57] T. Yokoyama. Generalization of Poincaré recurrence theorem and characterization of minimal flows on surfaces. *arXiv preprint arXiv:2110.05705*, 2021.
- [58] T. Yokoyama. On the ω -limit set of an orbit of a flow with arbitrarily many singular points on a surface. *arXiv preprint arXiv:2201.01020*, 2022.
- [59] T. Yokoyama. Flows with time-reversal symmetric limit sets on surfaces. *Proceedings of the American Mathematical Society*, to appear.

APPLIED MATHEMATICS AND PHYSICS DIVISION, GIFU UNIVERSITY, YANAGIDO 1-1,
GIFU, 501-1193, JAPAN

Email address: tomoo@gifu-u.ac.jp