

Borderline case of traces and extensions for weighted Sobolev spaces

Manzi Huang, Xiantao Wang, Zhuang Wang*, and Zhihao Xu

Abstract

In this paper, we study the traces and the extensions for weighted Sobolev spaces on upper half spaces when the weights reach to the borderline cases. We first give a full characterization of the existence of trace spaces for these weighted Sobolev spaces, and then study the trace parts and the extension parts between the weighted Sobolev spaces and a new kind of Besov-type spaces (on hyperplanes) which are defined by using integral averages over selected layers of dyadic cubes.

1 Introduction

The study of trace spaces (on the boundary of a domain) for Sobolev spaces on Euclidean domains is associated with the Dirichlet boundary value problem related to elliptic differential equations. Certain types of Dirichlet problems are guaranteed to have solutions when the boundary data arises as the trace of a Sobolev function. The work of Gagliardo identified the classical Besov spaces $B_{p,p}^{1-1/p}(\mathbb{R}^d)$ as the trace spaces for the first order Sobolev spaces $W^{1,p}(\mathbb{R}_+^{d+1})$, where $1 < p < \infty$, $\mathbb{R}_+^{d+1} := \{(x, t) : x \in \mathbb{R}^d, t > 0\}$, and $1 \leq d \in \mathbb{N}$, the set of all natural numbers, see [3, Theorem 1.I]. In the borderline case, that is, $p = 1$, it was also shown that the trace spaces for $W^{1,1}(\mathbb{R}_+^{d+1})$ are $L^1(\mathbb{R}^d)$, see [3, Theorem 1.II]. Here we say that a function space $\mathbb{X}(\mathbb{R}^d)$ is the trace space for the function space $\mathbb{Y}(\mathbb{R}_+^{d+1})$ if every function in $\mathbb{Y}(\mathbb{R}_+^{d+1})$ has a trace in $\mathbb{X}(\mathbb{R}^d)$ and every function in $\mathbb{X}(\mathbb{R}^d)$ is the trace of some function in $\mathbb{Y}(\mathbb{R}_+^{d+1})$. We refer to [3] and monographs [12, 14, 15] for more trace results and properties of classical Besov spaces.

It is natural to seek for the trace spaces for Sobolev spaces associated with weights. The trace spaces for Sobolev spaces with the Muckenhoupt A_p weights (in the following, we briefly denote by A_p the set of all Muckenhoupt A_p weights) (see Definition 2.1 below for the precise definition) were also well studied. For example, early trace results for Sobolev spaces with weights in A_p were given by Nikolskii, Lizorkin and Vašarin, see [9, 11, 19].

2010 **Mathematics Subject classification:** 46E35, 42B35, 30L99

Key words and phrases: Sobolev space, borderline case, trace theorem, Besov-type space, Muckenhoupt A_p weights

* *Corresponding author*

Some recent results about trace spaces of Sobolev spaces with weights belonging to A_p were given by [7, 8, 16–18]. We also refer to [1, 4, 6, 10, 12–15] for more investigations in this line. There is an advantage for the discussions on the trace spaces of Sobolev spaces when the weights are in A_p because under this constraint, the related trace spaces always exist. However, to the best of our knowledge, there are very few results about the trace spaces for Sobolev spaces with weights not belonging to A_p in the literatures.

For $1 \leq p < \infty$, $-1 < \alpha \leq p - 1$ and $\lambda \in \mathbb{R}$, let $\omega_\alpha^\lambda: \mathbb{R}_+^{d+1} \rightarrow (0, \infty)$ denote the weight

$$(1) \quad (x_1, x_2, \dots, x_{d+1}) \mapsto \begin{cases} |x_{d+1}|^\alpha \log^\lambda \frac{4}{|x_{d+1}|}, & x_{d+1} \in (0, 1], \\ \log^\lambda 4, & x_{d+1} \in (1, \infty), \end{cases}$$

and the weighted measure μ_α^λ on $\mathbb{R}_+^{d+1}$ is defined by

$$(2) \quad \mu_\alpha^\lambda(E) = \frac{1}{\log^\lambda 4} \int_E \omega_\alpha^\lambda dm_{d+1}.$$

It is known that, when $-1 < \alpha < p - 1$, for each $p \geq 1$, the weight ω_α^λ belongs to A_p for every $\lambda \in \mathbb{R}$ (cf. Proposition 2.2 below). But when $\alpha = p - 1$, the situation is much different. Again, by Proposition 2.2, we see that if $p > 1$, the weight ω_{p-1}^λ does not belong to A_p for any $\lambda \in \mathbb{R}$, and even for the special case when $p = 1$, the weight ω_0^λ does not belong to A_p for any $\lambda < 0$.




Figure 1: The set Γ .

Let

$$\Pi = \{(p, \lambda) \in \mathbb{R}^2 : p \geq 1\}$$

and

$$\Gamma = \{(p, \lambda) \in \Pi : p > 1, \lambda > p - 1\} \cup \{(p, \lambda) \in \Pi : p = 1, \lambda \geq 0\}$$

(see Figure 1).

The following example shows that there are $(p, \lambda) \in \Pi$ such that the related trace operators $\mathcal{T}$ on $W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ do not exist. The reader is referred to Definition 2.5 below for the precise definition of trace operators.

Example 1.1. Suppose that $(p, \lambda) \in \Pi$ and $\alpha = p - 1$. For any pair $(p, \lambda) \notin \Gamma$, there is a function $u \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$ such that $\mathcal{T}u$ does not exist.

Naturally, one will ask whether for any $(p, \lambda) \in \Gamma$, the related trace operator $\mathcal{T}$ on $W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ exists. Our following result shows that the answer to this question is positive.

Theorem 1.2. *Suppose that $(p, \lambda) \in \Pi$. Then*

- (i) *the trace function $\mathcal{T}u$ belongs to $L^p(\mathbb{R}^d)$ for every $u \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ if and only if $(p, \lambda) \in \Gamma$.*
- (ii) *for every $(p, \lambda) \in \Gamma$, the trace operator $\mathcal{T} : W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda) \rightarrow L^p(\mathbb{R}^d)$ is bounded and linear.*

To investigate the characterization of the trace spaces for weighted Sobolev spaces $W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ with $(p, \lambda) \in \Gamma$ is our main purpose of this paper. In [7], Koskela, Soto and the third author of this paper considered the weights ω_α^λ and the related measures μ_α^λ when $\lambda = 0$ and $\alpha < p - 1$, and they obtained the characterization of the corresponding trace spaces by the aid of a class of Besov spaces. But they did not consider the borderline case $\alpha = p - 1$ because the discussions in [7] do require the constraint $\alpha < p - 1$, and thus, the methods are invalid for the case $\alpha = p - 1$. Hence it needs new ideas to deal with this case.

Under the inspiration of the approach of using integral averages over dyadic cubes in [7] and the idea of choosing a system of tilings in [18], we construct a new kind of Besov-type spaces $\mathcal{B}_p^\gamma(\mathbb{R}^d)$ based on the integral averages over the so-called *selected layers of dyadic cubes* (see Definition 2.4 below for the details). By replacing $L^p(\mathbb{R}^d)$ in Theorem 1.2(ii) with $\mathcal{B}_p^\gamma(\mathbb{R}^d)$, we get the following related trace result.

Theorem 1.3. *Suppose that $(p, \lambda) \in \Gamma$. Then the trace operator $\mathcal{T} : W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda) \rightarrow \mathcal{B}_p^\gamma(\mathbb{R}^d)$ is bounded and linear provided that $0 < \gamma < \lambda - (p - 1)$ if $p > 1$ or $0 < \gamma \leq \lambda$ if $p = 1$.*

By the aid of $\mathcal{B}_p^\lambda(\mathbb{R}^d)$, we also obtain the following extension result.

Theorem 1.4. *Suppose that $(p, \lambda) \in \Gamma$. Then there exists a bounded and linear extension operator $\mathcal{E} : \mathcal{B}_p^\lambda(\mathbb{R}^d) \rightarrow W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ such that $\mathcal{T} \circ \mathcal{E} = \text{id}$ on $\mathcal{B}_p^\lambda(\mathbb{R}^d)$, where “id” denotes “the identity map”.*

The combination of Theorem 1.3 and Theorem 1.4 implies that, when $p = 1$, we get the full characterization of the trace spaces for weighted Sobolev spaces $W^{1,1}(\mathbb{R}_+^{d+1}, \mu_0^\lambda)$, which is formulated in the following corollary.

Corollary 1.5. *Let $\lambda > 0$. Then the Besov-type space $\mathcal{B}_1^\lambda(\mathbb{R}^d)$ is the trace space of the weighted Sobolev space $W^{1,1}(\mathbb{R}_+^{d+1}, \mu_0^\lambda)$.*

Notice that if $\lambda > 0$, the weights ω_0^λ are actually in A_1 (cf. Proposition 2.2). Hence the trace spaces $\mathcal{B}_1^\lambda(\mathbb{R}^d)$ are equivalent to the ones defined in [18]. The trace spaces in [18] are complicated since it needs to work for all weights in A_1 , and then, our concrete trace spaces $\mathcal{B}_1^\lambda(\mathbb{R}^d)$, which can be regarded as an example, would be helpful to understand it.

We remark that we do not consider the case when $\alpha > p - 1$. This is because, when $\alpha > p - 1$, by slightly modifying the proof of Proposition 2.2 and the construction of Example 1.1, we shall know that for any (p, λ) with $p \geq 1$ and $\lambda \in \mathbb{R}$, the weight ω_α^λ does not belong to A_p , and the trace operator $\mathcal{T}$ on $W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$ does not exist either.

The paper is organized as follows. In Section 2, some necessary terminology will be introduced and a proposition will be proved, which classifies the weights w_α^λ into two classes: A_p class and non A_p class. The proofs of Example 1.1, Theorem 1.2 and Theorem 1.3 will be presented in Section 3. Theorem 1.4 will be shown in Section 4.

2 Preliminaries

Throughout this paper, the letter C (sometimes with a subscript) will denote a positive constant that usually depend only on the given parameters of the spaces and may change at different occurrences; if C depends on $a, b, \dots$, then we write $C = C(a, b, \dots)$. The notation $A \lesssim B$ (resp. $A \gtrsim B$) means that there is a constant $C \geq 1$ such that $A \leq C \cdot B$ (resp. $A \geq C \cdot B$). If $A \lesssim B$ and $A \gtrsim B$, then we write $A \approx B$. If (X, μ) is a measure space, for every function $f \in L_{\text{loc}}^1(X, \mu)$ and every measurable subset $A \subset X$, let $\oint_A f d\mu$ stand for the integral average $\frac{1}{\mu(A)} \int_A f d\mu$, i.e.,

$$\oint_A f d\mu = \frac{1}{\mu(A)} \int_A f d\mu.$$

Let us recall the definition of the Muckenhoupt A_p weights with $1 \leq p < \infty$ (see, e.g. [5, Chapter 15] or [2]).

Definition 2.1. We say that a weight $\omega : \mathbb{R}_+^{d+1} \rightarrow [0, \infty)$ belongs to the Muckenhoupt class A_p if ω is locally integrable and there is a constant $C > 0$ such that for any cube $Q \subset \mathbb{R}_+^{d+1}$,

$$\left(\oint_Q \omega dm_{d+1} \right) \left(\oint_Q \omega^{-1/(p-1)} dm_{d+1} \right)^{p-1} \leq C \quad \text{if } p > 1$$

and

$$\left(\oint_Q \omega dm_{d+1} \right) \leq C \operatorname{ess\,inf}_{x \in Q} \omega(x) \quad \text{if } p = 1.$$

Proposition 2.2. (a) Suppose that $(p, \lambda) \in \Pi$, and let $-1 < \alpha < p - 1$. Then the weights ω_α^λ (see (1) for the definition) belong to A_p for all pairs (p, λ) with $p \geq 1$ and $\lambda \in \mathbb{R}$.
(b) Suppose that $(p, \lambda) \in \Pi$, and let $\alpha = p - 1$. Then the weights ω_α^λ belong to A_p if and only if $p = 1$ and $\lambda \geq 0$.

Proof. (a) The proof follows from a direct computation. We omit it here.

(b) We first show that if $p = 1$ and $\lambda \geq 0$, the weights ω_α^λ belong to A_p . If $p = 1$ and $\lambda = 0$, then $\omega_0^\lambda = 1$, and obviously it belongs to A_1 . If $p = 1$ and $\lambda > 0$, then for any $t \geq 0$ and $h > 0$, by letting $Q = (0, h]^d \times (t, t + h]$ and by using the integration by parts, we have

$$(3) \quad \int_Q \omega_0^\lambda dm_{d+1} \leq C_\lambda \log^\lambda \frac{4}{t+h} = C_\lambda \cdot \text{ess inf}_{x \in Q} \omega_0^\lambda(x),$$

which implies that the weight ω_0^λ belongs to A_1 .

For the converse, it suffices to show that for any pair (p, λ) with $p > 1$ and $\lambda \in \mathbb{R}$ or $p = 1$ and $\lambda < 0$, the weight ω_α^λ does not belong to A_p . For $p > 1$, we choose the cube $Q = (0, 2^{-k}]^{d+1} \subset \mathbb{R}_+^{d+1}$. Then

$$\int_Q |x_{d+1}|^\alpha \log^\lambda \frac{4}{|x_{d+1}|} dm_{d+1} = 2^k \int_0^{2^{-k}} t^{p-1} \log^\lambda \frac{4}{t} dt$$

and

$$\int_Q \left(|x_{d+1}|^\alpha \log^\lambda \frac{4}{|x_{d+1}|} \right)^{-1/(p-1)} dm_{d+1} = 2^k \int_{(k+2) \log 2}^\infty t^{-\frac{\lambda}{p-1}} dt.$$

It follows that if $\lambda > p - 1$, then

$$\left(\int_Q |x_{d+1}|^\alpha \log^\lambda \frac{4}{|x_{d+1}|} dm_{d+1} \right) \left(\int_Q \left(|x_{d+1}|^\alpha \log^\lambda \frac{4}{|x_{d+1}|} \right)^{-1/(p-1)} dm_{d+1} \right)^{p-1} \gtrsim (k+2)^{p-1},$$

and if $\lambda \leq p - 1$, then for each $k \geq 0$,

$$\int_{(k+2) \log 2}^\infty t^{-\frac{\lambda}{p-1}} dt = \infty.$$

Hence ω_α^λ does not belong to A_p for any $\lambda \in \mathbb{R}$. For $p = 1$, the estimate (3) will fail for any $\lambda < 0$, since if $t = 0$, we have $\text{ess inf}_{x \in Q} \omega_0^\lambda(x) = 0$. This implies that ω_0^λ does not belong to A_1 for any $\lambda < 0$. Hence the proposition is proved. $\square$

Let us recall that μ_α^λ denotes the weighted measure on $\mathbb{R}_+^{d+1}$ defined as in (2). Then by a direct computation, we know that the measure μ_α^λ is doubling on $\mathbb{R}_+^{d+1}$, i.e., there exists a constant $C \geq 1$ such that for all $x \in \mathbb{R}_+^{d+1}$ and $r > 0$,

$$\mu_\alpha^\lambda(\mathbb{B}(x, 2r) \cap \mathbb{R}_+^{d+1}) \leq C \mu_\alpha^\lambda(\mathbb{B}(x, r) \cap \mathbb{R}_+^{d+1}),$$

where $\mathbb{B}(x, r)$ denotes the open ball with center x and radius r .

Definition 2.3. Suppose that $p \in [1, \infty)$. Then $W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$ is defined as the normed space of all measurable functions $f \in L_{\text{loc}}^1(\mathbb{R}_+^{d+1})$ such that their first-order distributional derivatives, denoted by ∇f , belong to $L_{\text{loc}}^1(\mathbb{R}_+^{d+1})$, and

$$\|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} := \|f\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} + \|\nabla f\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} < +\infty.$$

In order to formulate the dyadic norms of the related Besov-type spaces, let us recall the standard dyadic decompositions of $\mathbb{R}^d$ and $\mathbb{R}_+^{d+1}$, respectively (cf. [7, Section 2]).

Denote by $\mathcal{Q}_d$ the collection of dyadic semi-open cubes in $\mathbb{R}^d$, i.e., the cubes of the form $Q := 2^{-k}((0, 1]^d + m)$, where $k \in \mathbb{Z}$, the set of all integers, $m \in \mathbb{Z}^d$, and $\mathcal{Q}_d^+$ stands for the set of all cubes in $\mathcal{Q}_d$ which are contained in the upper half-space $\mathbb{R}^{d-1} \times (0, \infty)$. Write $\ell(Q)$ for the edge length of $Q \in \mathcal{Q}_d$, i.e., 2^{-k} in the preceding representation, and $\mathcal{Q}_{d,k}$ for the cubes $Q \in \mathcal{Q}_d$ such that $\ell(Q) = 2^{-k}$.

Let $Q \in \mathcal{Q}_{d,2^j}$ for some $j \in \mathbb{N}$. We say that Q' in $\mathcal{Q}_d$ is a *selected neighbor* of Q , denoted by $Q' \prec Q$, if $Q' \in \mathcal{Q}_{d,2^j} \cup \mathcal{Q}_{d,2^{j-1}}$ and $\overline{Q} \cap \overline{Q'} \neq \emptyset$. Here we unify $\mathcal{Q}_{d,2^{-1}}$ as $\mathcal{Q}_{d,1}$, i.e., $\mathcal{Q}_{d,2^{-1}} = \mathcal{Q}_{d,1}$. Note that for every $Q \in \mathcal{Q}_d$, the number of its neighbors is uniformly bounded, and for every $Q \in \cup_{j \in \mathbb{N}} \mathcal{Q}_{d,2^j}$, also, the number of its selected neighbors is uniformly bounded.

Definition 2.4. Let $p \in [1, \infty)$ and $\lambda > 0$. Then the Besov space $\mathcal{B}_p^\lambda(\mathbb{R}^d)$ is defined as the normed space of all measurable functions $f \in L_{\text{loc}}^1(\mathbb{R}^d)$ such that

$$\|f\|_{\mathcal{B}_p^\lambda(\mathbb{R}^d)} := \|f\|_{L^p(\mathbb{R}^d)} + \|f\|_{\dot{\mathcal{B}}_p^\lambda(\mathbb{R}^d)} < +\infty,$$

where

$$\|f\|_{\dot{\mathcal{B}}_p^\lambda(\mathbb{R}^d)}^p := \sum_{j=0}^{\infty} (2^j + 2)^\lambda \sum_{Q \in \mathcal{Q}_{d,2^j}} m_d(Q) \sum_{Q' \prec Q} |f_Q - f_{Q'}|^p.$$

Let us recall the standard $(1, 1)$ -Poincaré inequality satisfied by the functions that are locally $W^{1,1}$ -regular in the upper half-space. If Q is a cube in $\mathbb{R}_+^{d+1}$ such that $\text{dist}(Q, \mathbb{R}^d \times \{0\}) > 0$ and $f \in W^{1,1}(Q)$, then there is a constant $C > 0$ independent of Q and f such that

$$(4) \quad \int_Q |f - f_Q| dm_{d+1} \leq C \ell(Q) \int_Q |\nabla f| dm_{d+1},$$

where “dist” means “distance”.

To formulate the trace and extension operators, we first recall a Whitney decomposition of $\mathbb{R}_+^{d+1}$ related to the dyadic decomposition (cf. [7]). For $Q \in \mathcal{Q}_{d,k}$ and $k \in \mathbb{Z}$, write $\mathcal{W}(Q) := Q \times (2^{-k}, 2^{-k+1}] \in \mathcal{Q}_{d+1,k}^+$. To simplify the notation in the sequel, we further define $\mathcal{Q}_d^1 := \cup_{k \geq 1} \mathcal{Q}_{d,k}$. Then $\{\mathcal{W}(Q) : Q \in \mathcal{Q}_d\}$ is a *Whitney decomposition* of $\mathbb{R}^d \times (0, \infty)$ with respect to the boundary $\mathbb{R}^d \times \{0\}$. For every $Q \in \mathcal{Q}_d^1$, define a smooth function

$$\psi_Q : \mathbb{R}_+^{d+1} \rightarrow [0, 1]$$

such that

- (i) $\text{Lip } \psi_Q \lesssim 1/\ell(Q)$,
- (ii) $\inf_{x \in \mathcal{W}(Q)} \psi_Q(x) > 0$ uniformly in $Q \in \mathcal{Q}_d^1$,
- (iii) $\text{supp } \psi_Q$ is contained in a $\ell(Q)/4$ -neighborhood of $\mathcal{W}(Q)$, and
- (iv) $\sum_{Q \in \mathcal{Q}_d^1} \psi_Q \equiv 1$ in $\bigcup_{Q \in \mathcal{Q}_d^1} \mathcal{W}(Q)$,

where “supp” means “support”.

We say that Q and Q' in $\mathcal{Q}_d$ are *neighbors* and write $Q \sim Q'$ if $\frac{1}{2} \leq \ell(Q)/\ell(Q') \leq 2$ and $\overline{Q} \cap \overline{Q'} \neq \emptyset$. We remark that the sum $\sum_{Q \in \mathcal{Q}_d^1} \psi_Q$ is locally finite – more precisely, it follows from the definition that

$$(5) \quad \text{supp } \psi_Q \cap \text{supp } \psi_{Q'} \neq \emptyset \quad \text{if and only if} \quad Q \sim Q'.$$

Definition 2.5 (Trace). Suppose that $f \in L_{\text{loc}}^1(\mathbb{R}_+^{d+1})$ and $k \in \mathbb{N}$. Define the function $\mathcal{T}_k f : \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$\mathcal{T}_k f := \sum_{Q \in \mathcal{Q}_{d,k}} \left(\int_{\mathcal{N}(Q)} f \, dm_{d+1} \right) \chi_Q,$$

where $\mathcal{N}(Q) = \frac{5}{4}\mathcal{W}(Q) := \{y \in \mathbb{R}_+^{d+1} : \text{dist}(y, \mathcal{W}(Q)) < \frac{1}{4}\ell(Q)\}$, and define the trace function $\mathcal{T}f$ by setting

$$\mathcal{T}f = \lim_{k \rightarrow \infty} \mathcal{T}_k f,$$

if the limit exists m_d -a.e. in $\mathbb{R}^d$.

Before going to the definition of the extension operator, we give one more notation. For any $Q \in \mathcal{Q}_d$, let $\mathcal{S}(Q)$ be the unique cube such that there is $j \in \mathbb{N}$ satisfying $\mathcal{S}(Q) \in \mathcal{Q}_{d,2^j}$ and $Q \subset \mathcal{S}(Q)$ if $Q \in \bigcup_{k=2^j}^{2^{j+1}-1} \mathcal{Q}_{d,k}$.

Definition 2.6 (Extension). Suppose that $f \in L_{\text{loc}}^1(\mathbb{R}^d)$. Then the *selected Whitney extension* $\mathcal{E}f : \mathbb{R}_+^{d+1} \rightarrow \mathbb{R}$ is defined by

$$\mathcal{E}f(x) = \sum_{Q \in \mathcal{Q}_d^1} \left(\int_{\mathcal{S}(Q)} f \, dm_d \right) \psi_Q(x).$$

It is easy to see that the extension operator $\mathcal{E} : L_{\text{loc}}^1(\mathbb{R}^d) \rightarrow C^\infty(\mathbb{R}_+^{d+1})$ is linear.

3 Trace operators and trace spaces

In this section, we shall give the proofs of Example 1.1, Theorem 1.2 and Theorem 1.3.

Proof of Example 1.1

Recall that if $(p, \lambda) \in \Pi$, but $(p, \lambda) \notin \Gamma$, then $\lambda \leq p - 1$ if $p > 1$, and $\lambda < 0$ if $p = 1$. Let φ be a compactly supported smooth function on $\mathbb{R}^d$ with $\varphi(x') = 1$ for $x' \in [-1, 1]^d$ and $\text{supp } \varphi \subset [-2, 2]^d$. Then we define the function u as follows.

For $(x', t) \in \mathbb{R}_+^{d+1}$, let

$$u(x', t) = \varphi(x') \cdot \max\{v(t), 0\},$$

where

$$v(t) = \int_t^1 \frac{1}{t \log(e/t) \left(1 + \log^\beta(\log(e/t))\right)} dt.$$

Here we choose β such that $\beta = 0$ if $p = 1$ and $0 < \beta < 1 < \beta p$ if $p > 1$. Obviously, for any $x' \in [-1, 1]^d$, $\mathcal{T}u(x') = \infty$. This shows that $\mathcal{T}u$ does not exist.

Next, we demonstrate that $u \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$. For any $(x', t) \notin [-2, 2]^d \times (0, 1]$, we have that

$$|u(x', t)| = |\nabla u(x', t)| = 0.$$

For any $(x', t) \in [-2, 2]^d \times (0, 1]$, there is a constant $C > 0$ such that

$$|u(x', t)| \leq C|v(t)| \leq C \int_t^1 \frac{1}{t \log(e/t)} dt = C \log(\log(e/t))$$

and

$$\begin{aligned} |\nabla u(x', t)| &\leq C|v(t)| + \frac{C}{t \log(e/t) \log^\beta(\log(e/t))} \\ &\leq C \log(\log(e/t)) + \frac{C}{t \log(e/t) \log^\beta(\log(e/t))}. \end{aligned}$$

For $p = 1$, $\lambda < 0$ and $\beta = 0$, the above facts guarantee that

$$\|u\|_{L^1(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} \lesssim 4^d \int_0^1 \log^\lambda(4/t) \log(\log(e/t)) dt < \infty$$

and

$$\begin{aligned} \|\nabla u\|_{L^1(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} &\lesssim \|u\|_{L^1(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} + 4^d \int_0^1 \frac{1}{t \log(e/t) \log^{-\lambda}(4/t)} dt \\ &\lesssim \|u\|_{L(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} + \int_0^1 \frac{1}{t \log^{1-\lambda}(4/t)} dt < \infty. \end{aligned}$$

For $p > 1$, $\lambda \leq p - 1$ and $1/p < \beta < 1$, similarly, we obtain that

$$\|u\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p \leq 4^d \int_0^1 t^{p-1} \log^\lambda(4/t) \log^p(\log(e/t)) dt < \infty$$

and

$$\begin{aligned}\|\nabla u\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p &\lesssim \|u\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p + 4^d \int_0^1 \frac{1}{t \log^p(e/t) \log^{-\lambda}(4/t) \log^{\beta p}(\log(e/t))} dt \\ &\lesssim \|u\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p + \int_0^1 \frac{1}{t \log^{p-\lambda}(4/t) \log^{\beta p}(\log(4/t))} dt < \infty.\end{aligned}$$

Hence $u \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$. These imply that the example is true. $\square$

Proof of Theorem 1.2

Proof of (i). From Example 1.1, it suffices to show that if $(p, \lambda) \in \Gamma$, then the trace function $\mathcal{T}u$ belongs to $L^p(\mathbb{R}^d)$ for every $u \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$.

Let $f \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$ with $\alpha = p - 1$ and $(p, \lambda) \in \Gamma$. The first thing is to verify the existence of the limit in Definition 2.5. To reach this goal, it suffices to show that the function

$$f^* := \sum_{k \geq 0} |\mathcal{T}_{k+1}f - \mathcal{T}_k f| + |\mathcal{T}_0 f|$$

belongs to $L^p(\mathbb{R}^d)$ because $f^* \in L^p(\mathbb{R}^d)$ implies that $f^*(x) < \infty$ for m_d -almost every $x \in \mathbb{R}^d$.

For any $P \in \mathcal{Q}_{d,0}$, notice that $m_{d+1}(\mathcal{N}(P)) \approx 1$ and $\omega_\alpha^\lambda \approx 1$ in $\mathcal{N}(P)$, it follows from the Minkowski inequality that

$$\begin{aligned}(6) \quad \left(\int_{\mathbb{R}^d} |f^*|^p dm_d \right)^{1/p} &\lesssim \left(\int_{\mathbb{R}^d} \left(\sum_{k \geq 0} |\mathcal{T}_{k+1}f - \mathcal{T}_k f| \right)^p dm_d \right)^{1/p} + \left(\sum_{P \in \mathcal{Q}_{d,0}} \int_{\mathcal{N}(P)} |f|^p dm_{d+1} \right)^{1/p} \\ &\lesssim \sum_{k \geq 0} \left(\int_{\mathbb{R}^d} |\mathcal{T}_{k+1}f - \mathcal{T}_k f|^p dm_d \right)^{1/p} + \left(\sum_{P \in \mathcal{Q}_{d,0}} \int_{\mathcal{N}(P)} |f|^p d\mu_\alpha^\lambda \right)^{1/p}.\end{aligned}$$

For any $x \in \mathbb{R}^d$, let Q_k^x be the unique cube in $\mathcal{Q}_{d,k}$ containing x . By the definition, we know that the intersection $\mathcal{N}(Q_k^x) \cap \mathcal{N}(Q_{k+1}^x)$ contains a cube $\hat{Q}$ with edge length comparable to 2^{-k} . Since $\omega_\alpha^\lambda(y) \approx 2^{-k\alpha}(2+k)^\lambda$ for all $y \in \mathcal{N}(Q_k^x)$ and $\mu_\alpha^\lambda(\mathcal{N}(Q_k^x)) \approx 2^{-k\alpha}(2+k)^\lambda m_{d+1}(\mathcal{N}(Q_k^x))$, it follows from the Poincaré inequality (4) that

$$\begin{aligned}|\mathcal{T}_{k+1}f(x) - \mathcal{T}_k f(x)| &= \left| \int_{\mathcal{N}(Q_k^x)} f dm_{d+1} - \int_{\mathcal{N}(Q_{k+1}^x)} f dm_{d+1} \right| \\ &\leq \left| \int_{\mathcal{N}(Q_k^x)} f dm_{d+1} - \int_{\hat{Q}} f dm_{d+1} \right| + \left| \int_{\hat{Q}} f dm_{d+1} - \int_{\mathcal{N}(Q_{k+1}^x)} f dm_{d+1} \right| \\ &\lesssim \int_{\mathcal{N}(Q_k^x)} |f - f_{\mathcal{N}(Q_k^x)}| dm_{d+1} + \int_{\mathcal{N}(Q_{k+1}^x)} |f - f_{\mathcal{N}(Q_{k+1}^x)}| dm_{d+1}\end{aligned}$$

$$\begin{aligned}
&\lesssim 2^{-k} \int_{\mathcal{N}(Q_k^x)} |\nabla f| dm_{d+1} + 2^{-k} \int_{\mathcal{N}(Q_{k+1}^x)} |\nabla f| dm_{d+1} \\
&\approx 2^{-k} \int_{\mathcal{N}(Q_k^x)} |\nabla f| d\mu_\alpha^\lambda + 2^{-k} \int_{\mathcal{N}(Q_{k+1}^x)} |\nabla f| d\mu_\alpha^\lambda.
\end{aligned}$$

Applying the Hölder inequality, we arrive at the estimate

$$|\mathcal{T}_{k+1}f(x) - \mathcal{T}_kf(x)| \lesssim 2^{-k} \left(\int_{\mathcal{N}(Q_k^x)} |\nabla f|^p d\mu_\alpha^\lambda \right)^{1/p} + 2^{-k} \left(\int_{\mathcal{N}(Q_{k+1}^x)} |\nabla f|^p d\mu_\alpha^\lambda \right)^{1/p}.$$

Hence

$$\begin{aligned}
\int_{\mathbb{R}^d} |\mathcal{T}_{k+1}f - \mathcal{T}_kf|^p dm_d &= \sum_{Q \in \mathcal{Q}_{d,k}} \int_Q |\mathcal{T}_{k+1}f - \mathcal{T}_kf|^p dm_d(x) \\
&\lesssim \sum_{Q \in \mathcal{Q}_{d,k}} m_d(Q) \left(2^{-kp} \int_{\mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \right. \\
&\quad \left. + \sum_{\substack{Q' \in \mathcal{Q}_{d,k+1} \\ Q' \subset Q}} 2^{-kp} \int_{\mathcal{N}(Q')} |\nabla f|^p d\mu_\alpha^\lambda \right) \\
&\lesssim 2^{-k(d+p)} \sum_{Q \in \mathcal{Q}_{d,k} \cup \mathcal{Q}_{d,k+1}} \int_{\mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \\
(7) \quad &\approx (2+k)^{-\lambda} \sum_{Q \in \mathcal{Q}_{d,k} \cup \mathcal{Q}_{d,k+1}} \int_{\mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda,
\end{aligned}$$

since $\alpha = p - 1$ implies that $\mu_\alpha^\lambda(\mathcal{N}(Q)) \approx 2^{-k(d+p)}(2+k)^\lambda$ for all $Q \in \mathcal{Q}_{d,k} \cup \mathcal{Q}_{d,k+1}$.

Plugging this into (6), we obtain that

$$\|f^*\|_{L^p(\mathbb{R}^d)} \lesssim \sum_{k \geq 0} (2+k)^{-\lambda/p} \left(\sum_{Q \in \mathcal{Q}_{d,k} \cup \mathcal{Q}_{d,k+1}} \int_{\mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \right)^{1/p} + \|f\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}.$$

If $p = 1$ and $\lambda \geq 0$, it is obvious that

$$\|f^*\|_{L^1(\mathbb{R}^d)} \lesssim \sum_{k \geq 0} \sum_{Q \in \mathcal{Q}_{d,k} \cup \mathcal{Q}_{d,k+1}} \int_{\mathcal{N}(Q)} |\nabla f| d\mu_\alpha^\lambda + \|f\|_{L^1(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} \lesssim \|f\|_{W^{1,1}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}.$$

If $p > 1$ and $\lambda > p - 1$, since the sum of $(2+k)^{-\lambda/(p-1)}$ converges, it follows from the Hölder inequality that

$$\|f^*\|_{L^p(\mathbb{R}^d)} \lesssim \|f\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} + \left(\sum_{k \geq 0} \sum_{Q \in \mathcal{Q}_{d,k} \cup \mathcal{Q}_{d,k+1}} \int_{\mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \right)^{1/p} \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}.$$

Thus we obtain that the estimate

$$(8) \quad \|f^*\|_{L^p(\mathbb{R}^d)} \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)}$$

holds for all $1 \leq p < \infty$. Hence $f^* < \infty$ for m_d -almost every $x \in \mathbb{R}^d$, so the trace function $\mathcal{T}f$ exists.

Since $|\mathcal{T}f| \leq |f^*|$ a.e. in $\mathbb{R}^d$, it follows from (8) that

$$(9) \quad \|\mathcal{T}f\|_{L^p(\mathbb{R}^d)} \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)} < \infty.$$

Hence the trace function $\mathcal{T}u$ belongs to $L^p(\mathbb{R}^d)$.

Proof of (ii). For every $(p, \lambda) \in \Gamma$, it follows from the proof of (i) that the trace function $\mathcal{T}u$ belongs to $L^p(\mathbb{R}^d)$ for every $u \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ and that the norm estimate (9) holds for every $f \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$. These guarantee the boundedness of the trace operator $\mathcal{T}$. Since the linearity of the trace operator $\mathcal{T}$ is obvious from the definition, the proof is complete. $\square$

Proof of Theorem 1.3

Let $f \in W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)$ for $(p, \lambda) \in \Gamma$ (i.e., $\alpha = p - 1$). It follows from Theorem 1.2 that the trace function $\mathcal{T}f$ exists and

$$\|\mathcal{T}f\|_{L^p(\mathbb{R}^d)} \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}.$$

The remaining task is to estimate the $\dot{\mathcal{B}}_p^\gamma(\mathbb{R}^d)$ -energy of $\mathcal{T}f =: \tilde{f}$. Let $P \in \mathcal{Q}_{d,2^k}$. Then

$$m_d(P) \sum_{Q \prec P} |\tilde{f}_Q - \tilde{f}_P|^p \leq \sum_{\substack{Q \prec P \\ Q \in \mathcal{Q}_{d,2^k}}} \int_P |\tilde{f} - \tilde{f}_Q|^p dm_d + \sum_{\substack{Q \prec P \\ Q \in \mathcal{Q}_{d,2^{k-1}}}} \int_P |\tilde{f} - \tilde{f}_Q|^p dm_d =: H_P^1 + H_P^2,$$

and so,

$$\begin{aligned} \|\mathcal{T}f\|_{\dot{\mathcal{B}}_p^\gamma(\mathbb{R}^d)}^p &= \|\tilde{f}\|_{\dot{\mathcal{B}}_p^\gamma(\mathbb{R}^d)}^p = \sum_{k=0}^{\infty} (2^k + 2)^\gamma \sum_{P \in \mathcal{Q}_{d,2^k}} m_d(P) \sum_{Q \prec P} |\tilde{f}_Q - \tilde{f}_P|^p \\ &\leq \sum_{k=0}^{\infty} (2^k + 2)^\gamma \sum_{P \in \mathcal{Q}_{d,2^k}} (H_P^1 + H_P^2) \\ &=: H^1 + H^2, \end{aligned}$$

where

$$H^i = \sum_{k=0}^{\infty} (2^k + 2)^\gamma \sum_{P \in \mathcal{Q}_{d,2^k}} H_P^i \quad \text{for } i = 1, 2.$$

Towards the estimate of H^1 , notice that

$$m_d(P) \approx m_d(Q) \approx 2^{2^k p} (2^k + 2)^{-\lambda} \mu_\alpha^\lambda(\mathcal{N}(P)) \approx 2^{2^k p} (2^k + 2)^{-\lambda} \mu_\alpha^\lambda(\mathcal{N}(Q))$$

for any $P, Q \in \mathcal{Q}_{d,2^k}$ with $P \asymp Q$. Hence

$$\begin{aligned} H_P^1 &\leq \sum_{\substack{Q \asymp P \\ Q \in \mathcal{Q}_{d,2^k}}} \left(\int_P |\tilde{f} - f_{\mathcal{N}(P)}|^p dm_d + m_d(P) |f_{\mathcal{N}(P)} - f_{\mathcal{N}(Q)}|^p + \int_Q |\tilde{f} - f_{\mathcal{N}(Q)}|^p dm_d \right) \\ &\lesssim \sum_{\substack{Q \asymp P \\ Q \in \mathcal{Q}_{d,2^k}}} \int_Q |\tilde{f}(x) - \mathcal{T}_{2^k} f(x)|^p dm_d(x) + \sum_{\substack{Q \asymp P \\ Q \in \mathcal{Q}_{d,2^k}}} m_d(P) |f_{\mathcal{N}(P)} - f_{\mathcal{N}(Q)}|^p \\ &\approx \sum_{\substack{Q \asymp P \\ Q \in \mathcal{Q}_{d,2^k}}} \int_Q |\tilde{f}(x) - \mathcal{T}_{2^k} f(x)|^p dm_d(x) + 2^{2^k p} (2^k + 2)^{-\lambda} \sum_{\substack{Q \asymp P \\ Q \in \mathcal{Q}_{d,2^k}}} \mu_\alpha^\lambda(\mathcal{N}(P)) |f_{\mathcal{N}(P)} - f_{\mathcal{N}(Q)}|^p. \end{aligned}$$

Using the Poincaré inequality (4) and the fact that $\#\{Q : Q \asymp P\}$ are uniformly bounded, where “ $\#$ ” means “cardinality”, we get

$$\begin{aligned} \sum_{P \in \mathcal{Q}_{d,2^k}} H_P^1 &\lesssim \int_{\mathbb{R}^d} |\tilde{f}(x) - \mathcal{T}_{2^k} f(x)|^p dm_d(x) + (2^k + 2)^{-\lambda} \int_{\bigcup_{2^{-2^k-1} \leq \ell(Q) \leq 2^{-2^k+1}} \mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \\ &=: I_k + (2^k + 2)^{-\lambda} I'_k. \end{aligned}$$

Since the domains of integration in I'_k 's have bounded overlap, we obtain that

$$\sum_{k \geq 0} I'_k \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p.$$

Next, we are going to estimate the term I_k . Since it follows from the triangle inequality and the Minkowski inequality that

$$(I_k)^{1/p} \leq \left(\int_{\mathbb{R}^d} \left(\sum_{n \geq 2^k} |\mathcal{T}_{n+1} f - \mathcal{T}_n f| \right)^p dm_d \right)^{1/p} \leq \sum_{n \geq 2^k} \left(\int_{\mathbb{R}^d} |\mathcal{T}_{n+1} f - \mathcal{T}_n f|^p dm_d \right)^{1/p},$$

we know from the estimate (7) that

$$\int_{\mathbb{R}^d} |\mathcal{T}_{n+1} f - \mathcal{T}_n f|^p dm_d \lesssim (2+n)^{-\lambda} \sum_{Q \in \mathcal{Q}_{d,n} \cup \mathcal{Q}_{d,n+1}} \int_{\mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda.$$

For the case when $p > 1, \alpha = p - 1$ and $0 < \gamma < \lambda - (p - 1)$, it follows from the Hölder inequality that

$$I_k \lesssim \left(\sum_{n \geq 2^k} (n+2)^{-(\lambda-\gamma)/p} \left((n+2)^{-\gamma} \int_{\bigcup_{2^{-n-2} \leq \ell(Q) \leq 2^{-n+1}} \mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \right)^{1/p} \right)^p$$

$$\lesssim \sum_{n \geq 2^k} (n+2)^{-\gamma} \int_{\cup_{2^{-n-2} \leq \ell(Q) \leq 2^{-n+1}} \mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda,$$

since the fact $\lambda - \gamma > p - 1$ implies that the series $\sum_{n=1}^\infty (n+2)^{-\frac{\lambda-\gamma}{p-1}}$ is convergent.

For the case when $p = 1$ and $0 < \gamma \leq \lambda$, it is obvious that

$$I_k \lesssim \sum_{n \geq 2^k} (n+2)^{-\gamma} \int_{\cup_{2^{-n-2} \leq \ell(Q) \leq 2^{-n+1}} \mathcal{N}(Q)} |\nabla f| d\mu_0^\lambda.$$

Hence the Fubini theorem and the above estimates guarantee that for $p \geq 1$,

$$\begin{aligned} \sum_{k \geq 0} (2^k + 2)^\lambda I_k &\lesssim \sum_{k \geq 0} (2^k + 2)^\gamma \sum_{n \geq 2^k} (n+2)^{-\gamma} \int_{\cup_{2^{-n-2} \leq \ell(Q) \leq 2^{-n+1}} \mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \\ &= \sum_{n \geq 1} (n+2)^{-\gamma} \int_{\cup_{2^{-n-2} \leq \ell(Q) \leq 2^{-n+1}} \mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \sum_{0 \leq 2^k \leq n} (2^k + 2)^\gamma \\ &\lesssim \sum_{n \geq 0} \int_{\cup_{2^{-n-2} \leq \ell(Q) \leq 2^{-n+1}} \mathcal{N}(Q)} |\nabla f|^p d\mu_\alpha^\lambda \\ &\lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p. \end{aligned}$$

Thus the estimate of H^1 is given by

$$H^1 = \sum_{k=0}^\infty (2^k + 2)^\gamma \sum_{P \in \mathcal{Q}_{d,2^k}} H_P^1 \lesssim \sum_{k \geq 0} (2^k + 2)^\gamma I_k + \sum_{k \geq 0} I'_k \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p.$$

For the estimate of H^2 , again, it follows from the Fubini theorem that

$$\begin{aligned} H^2 &= \sum_{k \geq 0} (2^k + 2)^\gamma \sum_{P \in \mathcal{Q}_{d,2^k}} \sum_{\substack{Q \succ P \\ Q \in \mathcal{Q}_{d,2^{k-1}}}} \int_P |\tilde{f} - \tilde{f}_Q|^p dm_d \\ &= \sum_{k \geq -1} (2^{k+1} + 2)^\gamma \sum_{Q \in \mathcal{Q}_{d,2^k}} \sum_{\substack{P \prec Q \\ P \in \mathcal{Q}_{d,2^{k+1}}}} \int_P |\tilde{f} - \tilde{f}_Q|^p dm_d \\ &\leq \sum_{k \geq -1} (2^{k+1} + 2)^\gamma \sum_{Q \in \mathcal{Q}_{d,2^k}} \sum_{\substack{Q' \prec Q \\ Q' \in \mathcal{Q}_{d,2^k}}} \int_{Q'} |\tilde{f} - \tilde{f}_Q|^p dm_d, \end{aligned}$$

where in the last inequality, the fact that for any $Q \in \mathcal{Q}_{d,2^k}$,

$$\bigcup_{\substack{P \prec Q \\ P \in \mathcal{Q}_{d,2^{k+1}}}} P \subset \bigcup_{\substack{Q' \prec Q \\ Q' \in \mathcal{Q}_{d,2^k}}} Q'$$

is applied.

By the reflexivity of the relation $Q' \asymp Q$ when $Q', Q \in \mathcal{Q}_{d,2^k}$, we obtain that

$$\begin{aligned} H^2 &\leq 3^\gamma \sum_{Q \in \mathcal{Q}_{d,2^{-1}}} \sum_{\substack{Q' \asymp Q \\ Q' \in \mathcal{Q}_{d,2^{-1}}}} \int_{Q'} |\tilde{f} - \tilde{f}_Q|^p dm_d + H^1 \\ &\lesssim \sum_{Q \in \mathcal{Q}_{d,1}} \int_Q |\tilde{f}|^p dm_d + H^1 \leq \|\mathcal{T}f\|_{L^p(\mathbb{R}^d)}^p + H^1 \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p. \end{aligned}$$

By combining the estimates of H^1 and H^2 , we obtain that

$$\|\mathcal{T}f\|_{\dot{B}_p^\gamma(\mathbb{R}^d)} \lesssim \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} = \|f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)},$$

which proves the theorem. $\square$

4 Extension operators

The purpose of this section is to prove Theorem 1.4.

Let $f \in \mathcal{B}_p^\lambda(\mathbb{R}^d)$ and $\alpha = p - 1$. For convenience, we rewrite $\mathcal{E}f$ (see Definition 2.6) as follows:

$$\mathcal{E}f(x) = \sum_{k \geq 0} \sum_{j=2^k}^{2^{k+1}-1} \sum_{Q \in \mathcal{Q}_{d,j}} \left(\int_{\mathcal{S}(Q)} f dm_d \right) \psi_Q(x).$$

Here we recall that $\mathcal{S}(Q) \in \mathcal{Q}_{d,2^k}$ is the unique cube with $Q \subset \mathcal{S}(Q)$ for any $Q \in \bigcup_{j=2^k}^{2^{k+1}-1} \mathcal{Q}_{d,j}$.

The first task is to estimate the $L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$ -norm of $\mathcal{E}f$. It follows directly that

$$\begin{aligned} \|\mathcal{E}f\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)}^p &= \int_{\mathbb{R}_+^{d+1}} |\mathcal{E}f|^p d\mu_\alpha^\lambda = \sum_{n \geq 1} \sum_{P \in \mathcal{Q}_{d,n}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda \\ &= \sum_{P \in \mathcal{Q}_{d,1}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda + \sum_{n \geq 2} \sum_{P \in \mathcal{Q}_{d,n}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda \\ &=: I_1 + I_2. \end{aligned}$$

To estimate I_1 , notice that for any $P \in \mathcal{Q}_{d,1}$ and each $x \in \text{supp } \psi_P$, we have $\psi_Q(x) \neq 0$ only for $Q \in \mathcal{Q}_{d,j}$ with $j = 1, 2$ and $Q \sim P$. Since $\mu_\alpha^\lambda(\text{supp } \psi_P) \approx 1$ and $|\psi_Q| \leq 1$ for any $P \in \mathcal{Q}_{d,1}$ and $Q \in \mathcal{Q}_d^1$, we obtain

$$I_1 = \sum_{P \in \mathcal{Q}_{d,1}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda = \sum_{P \in \mathcal{Q}_{d,1}} \int_{\text{supp } \psi_P} \left| \sum_{\substack{Q \in \mathcal{Q}_{d,1} \cup \mathcal{Q}_{d,2} \\ Q \sim P}} f_{\mathcal{S}(Q)} \psi_Q(x) \right|^p d\mu_\alpha^\lambda$$

$$\lesssim \sum_{Q \in \mathcal{Q}_{d,1} \cup \mathcal{Q}_{d,2}} \int_Q |f|^p dm_d \lesssim \|f\|_{L^p(\mathbb{R}^d)}^p,$$

where in the second last inequality, the fact that $\mathcal{S}(Q) = Q$ for any $Q \in \mathcal{Q}_{d,1} \cup \mathcal{Q}_{d,2}$ is used.

Towards the estimate of I_2 , since

$$\bigcup_{n \geq 2} \mathcal{Q}_{d,n} = \bigcup_{k \geq 1} \bigcup_{2^k \leq j \leq 2^{k+1}-1} \mathcal{Q}_{d,j},$$

we have

$$\begin{aligned} I_2 &= \sum_{k \geq 1} \sum_{j=2^k}^{2^{k+1}-1} \sum_{P \in \mathcal{Q}_{d,j}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda \\ &= \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^k}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda + \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^{k+1}-1}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda \\ &\quad + \sum_{k \geq 2} \sum_{j=2^{k+1}}^{2^{k+1}-2} \sum_{P \in \mathcal{Q}_{d,j}} \int_{\text{supp } \psi_P} |\mathcal{E}f|^p d\mu_\alpha^\lambda \\ &=: I_2^A + I_2^B + I_2^C. \end{aligned}$$

For the estimate of I_2^A , recall the relation (5) and notice that for $P \in \mathcal{Q}_{d,2^k}$, if $Q \sim P$, then

$$Q \in \mathcal{Q}_{d,2^{k-1}} \cup \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k+1}}.$$

By the definition of $\mathcal{S}(Q)$, we have

$$\begin{cases} \mathcal{S}(Q) \in \mathcal{Q}_{d,2^k} & \text{if } Q \in \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k+1}}, \\ \mathcal{S}(Q) \in \mathcal{Q}_{d,2^{k-1}} & \text{if } Q \in \mathcal{Q}_{d,2^{k-1}}. \end{cases}$$

Then it follows from the uniformly boundedness of $\#\{Q : Q \sim P\}$ that

$$\begin{aligned} I_2^A &\lesssim \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^k}} \mu_\alpha^\lambda(\mathcal{W}(P)) \sum_{Q \sim P} \int_{\mathcal{S}(Q)} |f|^p dm_d \\ &= \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^k}} \mu_\alpha^\lambda(\mathcal{W}(P)) \sum_{\substack{Q \sim P \\ Q \in \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k+1}}}} \int_{\mathcal{S}(Q)} |f|^p dm_d \\ &\quad + \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^k}} \mu_\alpha^\lambda(\mathcal{W}(P)) \sum_{\substack{Q \sim P \\ Q \in \mathcal{Q}_{d,2^{k-1}}}} \int_{\mathcal{S}(Q)} |f|^p dm_d \end{aligned}$$

$$\begin{aligned}
&\lesssim \sum_{k \geq 1} 2^{-2^k p} (2^k + 2)^\lambda \sum_{Q' \in \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k-1}}} \int_{Q'} |f|^p dm_d \\
&\lesssim \sum_{k \geq 1} 2^{-2^k p} (2^k + 2)^\lambda \int_{\mathbb{R}^d} |f|^p dm_d \lesssim \|f\|_{L^p(\mathbb{R}^d)}^p,
\end{aligned}$$

since $\mu_\alpha^\lambda(\mathcal{W}(P)) \approx 2^{-2^k(d+p)}(2^k + 2)^\lambda$ for $P \in \mathcal{Q}_{d,2^k}$ and $\sum_{k \geq 1} 2^{-2^k p} (2^k + 2)^\lambda$ is convergent.

The similar reasoning as in the estimate of I_2^A ensures that

$$I_2^B \lesssim \sum_{k \geq 1} 2^{-2^{k+1}p} (2^{k+1} + 2)^\lambda \sum_{Q' \in \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k+1}}} \int_{Q'} |f|^p dm_d \lesssim \|f\|_{L^p(\mathbb{R}^d)}^p.$$

The estimate of I_2^C is obtained by using the Fubini theorem as follows:

$$\begin{aligned}
I_2^C &\lesssim \sum_{k \geq 2} \sum_{j=2^{k+1}-2}^{2^{k+1}-2} \sum_{P \in \mathcal{Q}_{d,j}} \mu_\alpha^\lambda(\mathcal{W}(P)) \sum_{Q \sim P} \int_{\mathcal{S}(Q)} |f|^p dm_d \\
&= \sum_{k \geq 2} \sum_{Q' \in \mathcal{Q}_{d,2^k}} 2^{2^k d} \int_{Q'} |f|^p dm_d \left(\sum_{j=2^{k+1}-2}^{2^{k+1}-2} \sum_{\substack{P \in \mathcal{Q}_{d,j}, P \sim Q \\ \mathcal{S}(Q)=Q'}} \mu_\alpha^\lambda(\mathcal{W}(P)) \right) \\
&\approx \sum_{k \geq 2} \sum_{Q' \in \mathcal{Q}_{d,2^k}} 2^{2^k d} \int_{Q'} |f|^p dm_d \left(\sum_{j=2^{k+1}-2}^{2^{k+1}-2} 2^{-j(d+p)} (2+j)^\lambda \left(\frac{2^{-2^k}}{2^{-j}} \right)^d \right) \\
&\lesssim \sum_{k \geq 2} 2^{-2^k p} (2^k + 2)^\lambda \sum_{Q' \in \mathcal{Q}_{d,2^k}} \int_{Q'} |f|^p dm_d \lesssim \|f\|_{L^p(\mathbb{R}^d)}^p.
\end{aligned}$$

By combining the estimates of I_1 , I_2^A , I_2^B and I_2^C , we have

$$(10) \quad \|\mathcal{E}f\|_{L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} \lesssim \|f\|_{L^p(\mathbb{R}^d)}.$$

The next task is to estimate the $L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)$ -norm of $\nabla(\mathcal{E}f)$. To reach this goal, we first divide $\mathbb{R}_+^{d+1}$ into two parts:

$$L^p(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda) = X_1 \cup X_2,$$

where $X_1 = \bigcup_{k \geq 1} \bigcup_{Q \in \mathcal{Q}_{d,k}} \mathcal{W}(P)$ and $X_2 = \mathbb{R}_+^{d+1} \setminus X_1$.

If $x \in X_2$, we have $\psi_Q(x) \neq 0$ only for $Q \in \mathcal{Q}_{d,1}$. So it follows from the Lipschitz continuity of the functions ψ_Q that

$$|\nabla(\mathcal{E}f)(x)| \leq |\text{Lip}(\mathcal{E}f)(x)| \lesssim \sum_{Q \in \mathcal{Q}_{d,1}} |f_Q| \chi_{\text{supp } \psi_Q}(x).$$

Since $\mu_\alpha^\lambda(\text{supp } \psi_Q) \approx 1$ for each $Q \in \mathcal{Q}_{d,1}$, the above estimate yields that

$$(11) \quad \int_{X_2} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda \lesssim \sum_{Q \in \mathcal{Q}_{d,1}} |f_Q|^p \mu_\alpha^\lambda(\text{supp } \psi_Q) \lesssim \|f\|_{L^p(\mathbb{R}^d)}^p.$$

If on the other hand $x \in X_1$, then there exist a $k \geq 0$ and an unique cube $P \in \mathcal{Q}_{d,j}$ such that $2^k \leq j < 2^{k+1}$ and $x \in \mathcal{W}(P)$. So we can have

$$\begin{aligned} \int_{X_1} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda &= \sum_{k \geq 0} \sum_{j=2^k}^{2^{k+1}-1} \sum_{P \in \mathcal{Q}_{d,j}} \int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda \\ &= \sum_{k \geq 0} \sum_{P \in \mathcal{Q}_{d,2^k}} \int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda + \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^{k+1}-1}} \int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda \\ &\quad + \sum_{k \geq 2} \sum_{j=2^k+1}^{2^{k+1}-2} \sum_{P \in \mathcal{Q}_{d,2^{k+1}-1}} \int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda \\ &=: H^A + H^B + H^C. \end{aligned}$$

For the estimate of H^A , let $P \in \mathcal{Q}_{d,2^k}$ with $k \geq 0$. Then for each $x \in \mathcal{W}(P)$,

$$(12) \quad \mathcal{E}f(x) = \sum_{Q \sim P} \left(\int_{\mathcal{S}(Q)} f dm_d \right) \psi_Q(x).$$

Again, it follows from the Lipschitz continuity of the functions ψ_Q that

$$|\nabla(\mathcal{E}f)(x)| \leq |\text{Lip}(\mathcal{E}f(x) - f_P)| \lesssim \sum_{Q \sim P} \frac{1}{2^{-2^k}} |f_{\mathcal{S}(Q)} - f_P|.$$

Since for any $P \in \mathcal{Q}_{d,2^k}$ and $Q \sim P$, the cube $\mathcal{S}(Q)$ is a selected neighbor of P , i.e., $\mathcal{S}(Q) \asymp P$, we obtain

$$\begin{aligned} H^A &\lesssim \sum_{k \geq 0} \sum_{P \in \mathcal{Q}_{d,2^k}} \mu_\alpha^\lambda(\mathcal{W}(P)) 2^{2^k p} \sum_{Q \sim P} |f_{\mathcal{S}(Q)} - f_P|^p \\ &\approx \sum_{k \geq 0} \sum_{P \in \mathcal{Q}_{d,2^k}} (2^k + 2)^\lambda m_d(P) \sum_{Q' \asymp P} |f_{Q'} - f_P|^p \\ &\leq \|f\|_{\mathcal{B}_p^\lambda(\mathbb{R}^d)}^p. \end{aligned}$$

For the estimate of H^B , let $P \in \mathcal{Q}_{d,2^{k+1}-1}$. Then $\mathcal{S}(P) \in \mathcal{Q}_{d,2^k}$ is the unique cube with $P \subset \mathcal{S}(P)$. It follows from (12) and the continuity of the functions ψ_Q that

$$|\nabla(\mathcal{E}f)(x)| \leq |\text{Lip}(\mathcal{E}f(x) - f_{\mathcal{S}(P)})| \lesssim \sum_{Q \sim P} \frac{1}{2^{-2^{k+1}}} |f_{\mathcal{S}(Q)} - f_{\mathcal{S}(P)}|.$$

Since for any $P \in \mathcal{Q}_{d,2^{k+1}-1}$ and $Q \sim P$, the cube $\mathcal{S}(P)$ is a selected neighbor of $\mathcal{S}(Q)$, i.e., $\mathcal{S}(P) \asymp \mathcal{S}(Q)$, it follows from the Fubini theorem that

$$\begin{aligned}
H^B &\lesssim \sum_{k \geq 1} \sum_{P \in \mathcal{Q}_{d,2^{k+1}-1}} \mu_\alpha^\lambda(\mathcal{W}(P)) 2^{2^{k+1}p} \sum_{Q \sim P} |f_{\mathcal{S}(Q)} - f_{\mathcal{S}(P)}| \\
&\lesssim \sum_{k \geq 1} \sum_{j=2^{k+1}-2}^{2^k} \sum_{Q \in \mathcal{Q}_{d,j}} m_d(Q) (2^{k+1} + 2)^\lambda \sum_{P' \asymp \mathcal{S}(Q)} |f_{\mathcal{S}(Q)} - f_{P'}|^p \\
&= \sum_{k \geq 1} (2^{k+1} + 2)^\lambda \sum_{Q' \in \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k+1}}} \sum_{P' \asymp Q'} |f_{P'} - f_{Q'}|^p \left(\sum_{j=2^{k+1}-2}^{2^k} \sum_{\substack{Q \in \mathcal{Q}_{d,j} \\ \mathcal{S}(Q) = Q'}} m_d(Q) \right) \\
&\lesssim \sum_{k \geq 1} (2^{k+1} + 2)^\lambda \sum_{Q' \in \mathcal{Q}_{d,2^k} \cup \mathcal{Q}_{d,2^{k+1}}} m_d(Q') \sum_{P' \asymp Q'} |f_{P'} - f_{Q'}|^p \lesssim \|f\|_{\mathcal{B}_p^\lambda(\mathbb{R}^d)}^p.
\end{aligned}$$

The left is to estimate H^C . For any $2^k + 1 \leq j \leq 2^{k+1} - 2$ with $k \geq 2$, let

$$\mathcal{Q}_{d,j} = Y_{1,j} \cup Y_{2,j},$$

where

$$Y_{1,j} := \{P : P \in \mathcal{Q}_{d,j}, \overline{P} \cap \overline{\mathbb{R}^d \setminus \mathcal{S}(P)} = \emptyset\}$$

and

$$Y_{2,j} := \{P : P \in \mathcal{Q}_{d,j}, \overline{P} \cap \overline{\mathbb{R}^d \setminus \mathcal{S}(P)} \neq \emptyset\}.$$

Let $x \in \mathcal{W}(P)$ for $P \in \mathcal{Q}_{d,j}$. If $P \in Y_{1,j}$, then the definition of $\mathcal{E}f$ implies that $\mathcal{E}f(x) \equiv f_{\mathcal{S}(P)}$, and hence,

$$\int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda = 0.$$

For the case when $P \in Y_{2,j}$, notice that for $Q \sim P$, the cube $\mathcal{S}(Q)$ is a selected neighbor of $\mathcal{S}(P)$, i.e., $\mathcal{S}(Q) \asymp \mathcal{S}(P)$. It follows from (12) that

$$|\nabla(\mathcal{E}f)(x)| \leq |\text{Lip}(\mathcal{E}f(x) - f_{\mathcal{S}(P)})| \lesssim \sum_{Q \sim P} \frac{1}{2^{-j}} |f_{\mathcal{S}(Q)} - f_{\mathcal{S}(P)}| \leq \frac{1}{2^{-j}} \sum_{Q \asymp \mathcal{S}(P)} |f_Q - f_{\mathcal{S}(P)}|.$$

Hence, for any $P \in Y_{2,j}$, we have the estimate

$$\begin{aligned}
\int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda &\lesssim \mu_\alpha^\lambda(\mathcal{W}(P)) \left(\frac{1}{2^{-j}} \sum_{Q \asymp \mathcal{S}(P)} |f_Q - f_{\mathcal{S}(P)}| \right)^p \\
&\lesssim 2^{-jd} (j+2)^\lambda \sum_{Q \asymp \mathcal{S}(P)} |f_Q - f_{\mathcal{S}(P)}|^p.
\end{aligned}$$

It follows from the Fubini theorem that

$$\begin{aligned}
H^C &= \sum_{k \geq 2} \sum_{j=2^{k+1}-2} \sum_{P \in Y_{1,j}} \int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda + \sum_{k \geq 2} \sum_{j=2^{k+1}-2} \sum_{P \in Y_{2,j}} \int_{\mathcal{W}(P)} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda \\
&= \sum_{k \geq 2} \sum_{j=2^{k+1}-2} \sum_{P \in Y_{2,j}} 2^{-jd} (j+2)^\lambda \sum_{Q \sim \mathcal{S}(P)} |f_Q - f_{\mathcal{S}(P)}|^p \\
&\approx \sum_{k \geq 2} (2^k + 2)^\lambda \sum_{P' \in \mathcal{Q}_{d,2^k}} m_d(P') \sum_{Q \sim P'} |f_Q - f_{P'}|^p \left(\sum_{j=2^{k+1}-2} \sum_{\substack{P \in Y_{2,j} \\ P' = \mathcal{S}(P)}} 2^{-jd} / m_d(P') \right).
\end{aligned}$$

Note that for each $P' \in \mathcal{Q}_{d,2^k}$, there are $\left(2^{(j-2^k)d} - (2^{j-2^k} - 2)^d\right)$ many of $P \in Y_{2,j}$ such that $\mathcal{S}(P) = P'$. Hence

$$\begin{aligned}
\sum_{j=2^{k+1}-2} \sum_{\substack{P \in Y_{2,j} \\ P' = \mathcal{S}(P)}} 2^{-jd} / m_d(P') &= \sum_{j=2^{k+1}-2} 2^{(2^k-j)d} \left(2^{(j-2^k)d} - (2^{j-2^k} - 2)^d\right) \\
&\leq \sum_{i=0}^{\infty} 1 - (1 - 2^{-i})^d \\
&\leq \sum_{i=0}^{\infty} d \cdot 2^{-i} < \infty.
\end{aligned}$$

This yields that

$$H^C \lesssim \sum_{k \geq 2} (2^k + 2)^\lambda \sum_{P' \in \mathcal{Q}_{d,2^k}} m_d(P') \sum_{Q \sim P'} |f_Q - f_{P'}|^p \leq \|f\|_{\mathcal{B}_p^\lambda(\mathbb{R}^d)}^p.$$

By combining the estimates of H^A , H^B and H^C , we get that

$$\int_{X_1} |\nabla(\mathcal{E}f)|^p d\mu_\alpha^\lambda = H^A + H^B + H^C \lesssim \|f\|_{\mathcal{B}_p^\lambda(\mathbb{R}^d)}^p.$$

By recalling the estimates (10) and (11), we finally obtain that

$$\|\mathcal{E}f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_{p-1}^\lambda)} = \|\mathcal{E}f\|_{W^{1,p}(\mathbb{R}_+^{d+1}, \mu_\alpha^\lambda)} \lesssim \|f\|_{\mathcal{B}_p^\lambda(\mathbb{R}^d)}.$$

Since all m_d -almost every points in $\mathbb{R}^d$ are the Lebesgue ones of a function $f \in \mathcal{B}_p^\lambda(\mathbb{R}^d)$, it is evident from the definition of the trace operator $\mathcal{T}$ that $\mathcal{T}(\mathcal{E}f) = f$ for m_d -almost every points in $\mathbb{R}^d$, and hence, the theorem is proved. $\square$

Acknowledgments

The first author (Manzi Huang) was partly supported by NNSF of China under the number 11822105. The second author (Xiantao Wang) was partly supported by NNSFs of China under the numbers 12071121 and 11720101003 and the project under the number 2018KZDXM034. The third author (Zhuang Wang) was supported by NNSF of China under the number 12101226.

References

- [1] Aronszajn, N.: *Boundary value of functions with finite Dirichlet integral*, Techn. Report 14, University of Kansas, 1955. [2](#)
- [2] Dyda, B., Ihnatsyeva, L., Lehrbäck, J., Tuominen, H., Vähäkangas, A.: *Muckenhoupt A_p -properties of distance functions and applications to Hardy-Sobolev -type inequalities*, Potential Anal. 50 (2019), no. 1, 83-105. [4](#)
- [3] Gagliardo, E.: *Caratterizzazioni delle tracce sulla frontiera relative ad alcune classi di funzioni in n variabili*, Rend. Sem. Mat. Univ. Padova 27 (1957), 284-305. [1](#)
- [4] Hajłasz, P., Martio, P.: *Trace spaces of Sobolev functions on fractal type sets and characterization of extension domains*, J. Funct. Anal. 143 (1997), no. 1, 221-246. [2](#)
- [5] Heinonen, J., Kilpeläinen, T., Martio, O.: *Nonlinear potential theory of degenerate elliptic equations*, Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1993. vi+363 pp. [4](#)
- [6] Ihnatsyeva, L., Vähäkangas, A.: *Characterization of trace spaces of smooth functions on Ahlfors regular sets*, J. Funct. Anal. 265 (2013), no. 9, 1870-1915. [2](#)
- [7] Koskela, P., Soto, T., Wang, Z.: *Trace spaces of weighted function spaces: dyadic norms and Whitney extensions*, Sci. China Math. 60 (2017), no. 11, 1981-2010. [2](#), [3](#), [6](#)
- [8] Lindquist, J., Shanmugalingam, N.: *Trace spaces and extensions of certain weighted Sobolev spaces on $\mathbb{R}^n$ and Besov functions on Ahlfors regular compact subsets of $\mathbb{R}^n$* , Complex Anal. Synerg. 7 (2021), no. 1, Paper No. 7. [2](#)
- [9] Lizorkin, P. I.: *Boundary properties of functions from “weight” classes* (Russian), Dokl. Akad. Nauk SSSR 132 (1960), 514–517; translated as Soviet Math. Dokl. 1 (1960), 589–593. [1](#)

- [10] Mironescu, P., Russ, E.: *Trace spaces of weighted Sobolev spaces. Old and new*, Nonlinear Anal. 119 (2015), 354-381. [2](#)
- [11] Nikolskii, S. M.: *Properties of certain classes of functions of several variables on differentiable manifolds* (Russian), Mat. Sb. N.S. 33(75), no. 2 (1953), 261-326. [1](#)
- [12] Peetre, J.: *New thoughts on Besov spaces*, Duke University Mathematics Series, No. 1. Mathematics Department, Duke University, Durham, N.C., 1976. [1](#), [2](#)
- [13] Slobodetskii, L. N., Babich, V. M.: *On boundedness of the Dirichlet integrals* (Russian), Dokl. Akad. Nauk SSSR (N.S.) 106 (1956), 604-606. [2](#)
- [14] Triebel, H.: *Theory of function spaces*, Monographs in Mathematics, 78. Birkhäuser Verlag, Basel, 1983. [1](#), [2](#)
- [15] Triebel, H.: *The structure of functions*, Monographs in Mathematics, 97. Birkhäuser Verlag, Basel, 2001. [1](#), [2](#)
- [16] Tyulenev, A. I.: *Description of trace spaces of functions in the Sobolev space with a Muckenhoupt weight*, Proc. Steklov Inst. Math. 284 (2014), no. 1, 280-295. [2](#)
- [17] Tyulenev, A. I.: *Boundary values of functions in a Sobolev space with weight of Muckenhoupt class on some non-Lipschitz domains*, Mat. Sb. 205 (2014), no. 8, 67-94; translation in Sb. Math. 205 (2014), no. 7-8, 1133-1159. [2](#)
- [18] Tyulenev, A. I.: *Traces of weighted Sobolev spaces with Muckenhoupt weight. The case $p = 1$* , Nonlinear Anal. 128 (2015), 248-272. [2](#), [3](#), [4](#)
- [19] Vašarin, A. A.: *The boundary properties of functions having a finite Dirichlet integral with a weight* (Russian), Dokl. Akad. Nauk SSSR (N.S.) 117 (1957), 742-744. [1](#)

Manzi Huang,

MOE-LCSM, School of Mathematics and Statistics, Hunan Normal University, Changsha, Hunan 410081, People's Republic of China, and School of Mathematical Science, Qufu Normal University, Qufu, Shangdong 273165, People's Republic of China

E-mail address: mzhuang@hunnu.edu.cn

Xiantao Wang,

MOE-LCSM, School of Mathematics and Statistics, Hunan Normal University, Changsha, Hunan 410081, People's Republic of China.

E-mail address: xtwang@hunnu.edu.cn

Zhuang Wang,

MOE-LCSM, School of Mathematics and Statistics, Hunan Normal University, Changsha,
Hunan 410081, People's Republic of China.

E-mail address: `zhuang.z.wang@foxmail.com`

Zhihao Xu,

MOE-LCSM, School of Mathematics and Statistics, Hunan Normal University, Changsha,
Hunan 410081, People's Republic of China.

E-mail address: `734669860@qq.com`