

# Matching anti-forcing polynomials of catacondensed hexagonal systems

Shuang Zhao<sup>1,2</sup>

<sup>1</sup>School of Information Engineering, Lanzhou University of Finance and Economics,

Lanzhou, Gansu 730000, P. R. China;

E-mail: zhaosh2018@126.com

## Abstract

In this paper, we derive a recurrence relation of anti-forcing polynomial for catacondensed hexagonal systems.

**Keywords:** Perfect matching; Anti-forcing number; Anti-forcing polynomial; Catacondensed hexagonal system.

## 1. Introduction

Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . A *perfect matching*  $M$  of  $G$  is a set of disjoint edges that covers all vertices of  $G$ . Lei, Yeh and Zhang [4] generalized *anti-forcing number* to single perfect matching, and Klein and Rosenfeld [3] presented the same concept as *(e)-forcing*. An *anti-forcing set*  $S$  of  $M$  is a subset of  $E(G) \setminus M$  such that  $G - S$  has a unique perfect matching  $M$ , where  $G - S$  denotes the subgraph obtained from  $G$  by deleting the edges that belong to  $S$ . The *anti-forcing number* of  $M$  is the minimum cardinality over all anti-forcing sets of  $M$ , denoted by  $af(G, M)$ . The *maximum* (resp. *minimum*) *anti-forcing number* of  $G$  is the maximum (resp. minimum) value of  $af(G, M)$  over all perfect matchings  $M$  of  $G$ , denoted by  $Af(G)$  (resp.  $af(G)$ ). The set of anti-forcing numbers of all perfect matchings of  $G$  is called the *anti-forcing spectrum* of  $G$ .

A *hexagonal system* is a 2-connected finite plane graph such that every interior face is surrounded by a regular hexagon of side length one. Lei et al. [4] showed that the maximum anti-forcing number of a hexagonal system equals the *Fries number*, namely the maximum number of alternating hexagons with respect to a perfect matching, which can measure the stability of a benzenoid hydrocarbon.

Hwang et al. [2] introduced the *anti-forcing polynomial* of a graph  $G$  as

$$Af(G, x) = \sum_{M \in \mathcal{M}(G)} x^{af(G, M)} = \sum_{i=af(G)}^{Af(G)} v(G, i) x^i, \quad (1)$$

where  $\mathcal{M}(G)$  denotes the set of all perfect matchings of  $G$ ,  $v(G, i)$  denotes the number of perfect matchings of  $G$  with anti-forcing number  $i$ . By the definition, we can derive the following lemma, which shows that a disconnected graph can be partitioned into components to consider.

**Lemma 1.1.** *Let  $G$  be a graph with the components  $G_1, G_2, \dots, G_k$ . Then*

$$Af(G, x) = \prod_{i=1}^k Af(G_i, x).$$

From the above lemma, we know that if the anti-forcing spectrum of  $G_i$  is an integer interval  $[a_i, b_i]$  for  $i = 1, 2, \dots, k$ , then the anti-forcing spectrum of  $G$  is  $[\sum_{i=1}^k a_i, \sum_{i=1}^k b_i]$ .

## 2. Some preliminaries

We now introduce some properties of anti-forcing set and anti-forcing number. Let  $G$  be a graph with a perfect matching  $M$ . A cycle of  $G$  is called  *$M$ -alternating* if its edges appear alternately in  $M$  and  $E(G) \setminus M$ .

**Theorem 2.1.** [4] Let  $G$  be a graph with a perfect matching  $M$ . A subset  $S \subseteq E(G) \setminus M$  is an anti-forcing set of  $M$  if and only if each  $M$ -alternating cycle of  $G$  contains at least one edge of  $S$ .

Two  $M$ -alternating cycles is called *compatible* if they either are disjoint or intersect only at edges in  $M$ . A set of  $M$ -alternating cycles of  $G$  is called a *compatible  $M$ -alternating set* if every two members in it are compatible. It is easy to see from the above theorem that the anti-forcing number  $af(G, M)$  is bounded below by  $c'(G, M)$ , the maximum cardinality over all compatible  $M$ -alternating sets of  $G$ . Furthermore, Lei et al. [4] proved that all planar bipartite graphs satisfy the lower bound with equality, especially for hexagonal systems.

**Theorem 2.2.** [4] Let  $G$  be a planar bipartite graph. Then for each perfect matching  $M$  of  $G$ , we have  $af(G, M) = c'(G, M)$ .

For  $S \subseteq E(G) \setminus M$ , an edge  $e$  of  $G - S$  is said to be *forced* by  $S$  if it belongs to all perfect matchings of  $G - S$ , thus belongs to  $M$ . An edge  $g$  of  $G - S$  is said to be *anti-forced* by  $S$  if it belongs to no perfect matchings of  $G - S$ , thus does not belong to  $M$ . Let  $G \odot S$  denote the subgraph obtained from  $G - S$  by deleting the ends of all the edges that forced by  $S$  and deleting all the edges that anti-forced by  $S$ . Obviously we have the following lemma.

**Lemma 2.3.** *Let  $G$  be a graph with a perfect matching  $M$ . A subset  $S \subseteq E(G) \setminus M$  is an anti-forcing set of  $M$  if and only if  $G \ominus S$  is empty.*

**Lemma 2.4.** *Let  $G$  be a graph with a perfect matching  $M$ , and  $\mathcal{C}$  be a compatible  $M$ -alternating set of  $G$ . Given a subset  $S \subseteq E(G) \setminus M$ , which consists of precisely one edge from each cycle in  $\mathcal{C}$ . If  $S$  anti-forces all the other edges in  $E(\mathcal{C}) \cap (E(G) \setminus M)$ , then*

$$af(G, M) = af(G \ominus S, M \cap E(G \ominus S)) + |S|.$$

*Proof.* Let  $S'$  be a minimum anti-forcing set of  $M \cap E(G \ominus S)$  in  $G \ominus S$ . Then  $af(G \ominus S, M \cap E(G \ominus S)) = |S'|$ . Since  $G \ominus (S \cup S') = (G \ominus S) \ominus S'$  is empty,  $S \cup S'$  is an anti-forcing set of  $M$  in  $G$  by Lemma 2.3. Suppose  $S_0$  is another anti-forcing set of  $M$  in  $G$  such that  $|S_0| < |S \cup S'|$ . Then either  $|S_0 \cap E(G \ominus S)| < |S'|$  or  $|S_0 \cap (E(G) \setminus E(G \ominus S))| < |S| = |\mathcal{C}|$ . By assumption, we have  $E(\mathcal{C}) \cap (E(G) \setminus M) \subseteq E(G) \setminus E(G \ominus S)$ . It follows that either  $S_0 \cap E(G \ominus S)$  is not an anti-forcing set of  $M \cap E(G \ominus S)$  in  $G \ominus S$ , or there exists an  $M$ -alternating cycle in  $\mathcal{C}$  containing no edges of  $S_0 \cap (E(G) \setminus E(G \ominus S))$ . This implies that there is an  $M$ -alternating cycle in  $G$  containing no edges of  $S_0$ , a contradiction to Theorem 2.1. Hence  $S \cup S'$  is a minimum anti-forcing set of  $M$  in  $G$ , which implies  $af(G, M) = |S| + |S'|$ .  $\square$

By the above lemma, from a special compatible  $M$ -alternating set we can find a subset that is contained in some minimum anti-forcing set of a perfect matching  $M$ . In particular, by Theorem 2.2 for hexagonal systems we can find a minimum anti-forcing set directly.

### 3. Catacondensed hexagonal system

A hexagonal system  $H$  is called *catacondensed* if no three of its hexagons share a common vertex. If  $H$  contains at least two hexagons, then every hexagon of  $H$  has one, two, or three neighboring hexagons. A hexagon is called *terminal* and *branched* if it has one and three neighboring hexagons respectively. A hexagon with precisely two neighboring hexagons is called a *kink* if it possesses two adjacent vertices with degree two, and is called *linear* otherwise. If  $H$  contains precisely one hexagon, then we call it a *terminal hexagon*. In Fig. 1(a), we illustrate an example of catacondensed hexagonal system, where  $T$  denotes terminal hexagon,  $B$  denotes branched hexagon,  $K$  denotes kink, and  $L$  denotes linear hexagon.

A catacondensed hexagonal system with no branched hexagons is called a *hexagonal chain* (see Fig. 1(b)). A hexagonal chain with no kinks is called *linear*, and furthermore a linear hexagonal chain  $W$  in  $H$  is called *maximal* if it is contained in no other linear ones of  $H$ . Define the *length* of  $W$  as the number of hexagons in  $W$ , and a *vertical edge* of  $W$  as an edge that intersects with a straight line passing through all the centers of hexagons in  $W$ . In Fig. 1(a), we illustrate two maximal linear hexagonal chains with length four and two respectively, where  $e_i$  is a vertical edge of the first one and  $c_j$  is a vertical edge of the second one for  $i = 0, 1, 2, 3, 4$  and  $j = 0, 1, 2$ .




Fig. 1. (a) A catacondensed hexagonal system and (b) a hexagonal chain.

**Lemma 3.1.** [1] Let  $H$  be a catacondensed hexagonal system with a perfect matching  $M$ . Then every maximal linear hexagonal chain of  $H$  has precisely one vertical edge that belongs to  $M$ .

Before discussing anti-forcing polynomial of catacondensed hexagonal systems  $H$ , we give some notations first. A kink of  $H$  is called an *end kink* if  $H$  contains a maximal linear hexagonal chain starting from the kink and ending at a terminal hexagon. In Fig. 1(a) we illustrate an end kink as  $K_E$ , and a kink but not an end one as  $K_N$ . A branched hexagon of  $H$  is called an *end branched hexagon* if  $H$  contains two maximal linear hexagonal chains starting from the branched hexagon and ending at terminal hexagons. In Fig. 1(a) we illustrate an end branched hexagon as  $B_E$ , and a branched hexagon but not an end one as  $B_N$ . Except for linear hexagonal chains, every catacondensed hexagonal system contains a kink or a branched hexagon, thus contains an end kink or an end branched hexagon.

Define a *tail*  $T(t_1, t_2, t_3)$  with respect to  $h$  as a subgraph  $T_1 \cup T_2 \cup T_3$  of  $H$ , where  $h$  is a terminal hexagon of  $H$  if  $H$  is a linear hexagonal chain, and an end kink or an end branched hexagon of  $H$  otherwise. In detail, if  $H$  is a linear hexagonal chain, then define  $T_1 = H$  and  $T_2 = T_3 = h$ , and define  $t_1 + 2$  ( $t_1 \geq -1$ ) as the length of  $T_1$  and  $t_2 = t_3 = -1$  (see Fig. 2(a)). If  $h$  is an end kink, then define  $T_2 = h$ , and  $T_1, T_3$  as maximal linear hexagonal chains starting from  $h$  satisfying that  $T_1$  ends at a terminal hexagon, and define  $t_2 = -1$ ,  $t_1 + 2, t_3 + 2$  ( $t_1, t_3 \geq 0$ ) as the length of  $T_1, T_3$  respectively (see Fig. 2(b)). If  $h$  is an end branched hexagon, then define  $T_1, T_2, T_3$  as maximal linear hexagonal chains starting from  $h$  satisfying that  $T_1, T_2$  end at terminal hexagons, and define  $t_1 + 2, t_2 + 2, t_3 + 2$  ( $t_1, t_2, t_3 \geq 0$ ) as the length of  $T_1, T_2, T_3$  respectively (see Fig. 2(c)).

For a tail  $T(t_1, t_2, t_3)$  with respect to  $h$ , denote  $H_1^T$  the subgraph of  $H$  by deleting all the hexagons in  $T_1 \cup T_2$  except for  $h$ , denote  $H_2^T$  the subgraph of  $H$  by deleting all the hexagons in  $T_1 \cup T_2$ , and denote  $H_3^T$  the subgraph of  $H$  by deleting all the hexagons in  $T(t_1, t_2, t_3)$  (see Figs. 2(a-c)). Note that  $H_1^T$  is a *smaller* catacondensed hexagonal system than  $H$  (namely containing the number of hexagons less than that of  $H$ ) if  $t_1 \geq 0$ .  $H_2^T$  is an empty graph if  $t_3 = -1$  and is a smaller catacondensed hexagonal system than  $H$  otherwise.  $H_3^T$  is an empty graph if  $t_3 = -1$  or  $T_3$  ends at a terminal hexagon, is a




Fig. 2. Illustration of tail  $T(t_1, t_2, t_3)$  and  $H_i^T$  for  $i = 1, 2, 3, 4$ .

smaller catacondensed hexagonal system than  $H$  if  $t_3 \geq 0$  and  $T_3$  ends at a kink, and is a union of two smaller catacondensed hexagonal systems than  $H$  if  $t_3 \geq 0$  and  $T_3$  ends at a branched hexagon. If  $h$  is an end branched hexagon, then denote  $u_1, u_2$  the two edges that adjacent to the last vertical edge of  $T_3$ . Denote  $H_4^T$  the subgraph of  $H_3^T$  by deleting all the hexagons in the maximal linear hexagonal chains starting from the hexagon containing  $u_1$  and the hexagon containing  $u_2$  respectively (see Fig. 2(d)). Note that  $H_4^T$  may be an empty graph, or a union of one, two, three, or four smaller catacondensed hexagonal systems than  $H$ .

**Theorem 3.2.** *The anti-forcing polynomial of a catacondensed hexagonal system  $H$  possessing a tail  $T(t_1, t_2, t_3)$  with respect to  $h$  has the following form:*

(1) if  $t_2 = -1$ , then

$$Af(H, x) = xAf(H_2^T, x) + xAf(H_3^T, x) + \alpha_T; \quad (2)$$

(2) if  $t_2 \geq 0$ , then

$$Af(H, x) = x^2 Af(H_1^T, x) + (3x^2 + 2(t_1 + t_2)x^3 + t_1 t_2 x^4) Af(H_2^T, x) - x^3 Af(H_3^T, x) + \beta_T, \quad (3)$$

where  $\alpha_T = 0$  if  $t_1 = -1$ , and  $\alpha_T = xAf(H_1^T, x) + t_1 x^2 Af(H_2^T, x) - x^2 Af(H_3^T, x)$  if  $t_1 \geq 0$ ; and  $\beta_T = xAf(H_2^T, x) - x^2 Af(H_4^T, x)$  if  $t_3 = 0$ , and  $\beta_T = x^2 Af(H_3^T, x)$  if  $t_3 \geq 1$ .

*Proof.* (1) If  $H$  contains a tail  $T(t_1, -1, t_3)$  ( $t_1, t_3 \geq -1$ , and if  $t_1 = -1$  then  $t_3 = -1$ ), then we in turn denote the hexagon in  $T_1$  by  $C_i$  for  $i = 1, 2, \dots, t_1 + 2$  such that  $C_1 = h$ . Furthermore, denote the vertical edge that  $C_i$  and  $C_{i+1}$  share in common by  $e_i$ , and along the clockwise direction denote the rest edges of  $C_i$  by  $l_i, r_i, e_{i-1}, m_i, n_i$  for  $i = 1, 2, \dots, t_1 + 2$  (see Figs. 2(a,b)). By Lemma 3.1 we can divide  $\mathcal{M}(H)$  in  $t_1 + 3$  subsets:

$$\mathcal{M}_i(H) = \{M \in \mathcal{M}(H) : e_i \in M\}$$

for  $i = 0, 1, \dots, t_1 + 2$ . By Eq. (1), we have

$$Af(H, x) = \sum_{i=0}^{t_1+2} \sum_{M \in \mathcal{M}_i(H)} x^{af(H, M)} := \sum_{i=0}^{t_1+2} Af_i(H, x). \quad (4)$$

Given  $M \in \mathcal{M}_0(H)$ . Then  $r_1 \notin M$ . On the one hand  $r_1$  belongs to  $M$ -alternating hexagon  $C_1$ . And on the other hand  $r_1$  forces edges  $l_1, n_1, e_0$  and anti-forces edges  $e_1, m_1$ . By Lemma 2.4 we know

$$af(H, M) = af(H \ominus r_1, M \cap E(H \ominus r_1)) + 1.$$

In Fig. 3(a), we illustrate the three cases of  $H \ominus r_1$ , where double line denotes  $e_0$ , cross denotes  $r_1$ , and bold lines denote the edges that forced by  $r_1$ . Note that in the last case, the last vertical edge  $v$  of  $T_3$  is a cut edge whose removal from the subgraph  $H - r_1$  makes the two components to be even, which implies that  $v$  is anti-forced by  $r_1$ . Hence we know  $H \ominus r_1 = H_3^T$ . It follows that

$$Af_0(H, x) = \sum_{M \in \mathcal{M}_0(H)} x^{af(H_3^T, M \cap E(H_3^T)) + 1} = \sum_{M \in \mathcal{M}(H_3^T)} x^{af(H_3^T, M)} \cdot x = x Af(H_3^T, x). \quad (5)$$




Fig. 3. (a)  $H \ominus r_1$ , (b)  $H \ominus l_{t_1+2}$ , (c)  $H \ominus \{r_{i+1}, l_i\}$ , and (d)  $H \ominus r_2$ .

Given  $M \in \mathcal{M}_{t_1+2}(H)$ . Then  $l_{t_1+2} \notin M$ . On the one hand  $l_{t_1+2}$  belongs to  $M$ -alternating hexagon  $C_{t_1+2}$ . And on the other hand  $l_{t_1+2}$  forces edges  $e_{t_1+2}, r_{t_1+2}, m_{t_1+2}$  and anti-forces edges  $e_{t_1+1}, n_{t_1+2}$ . By Lemma 2.4 and Fig. 3(b) we know

$$af(H, M) = af(H \ominus l_{t_1+2}, M \cap E(H \ominus l_{t_1+2})) + 1 = af(H_2^T, M \cap E(H_2^T)) + 1.$$

It follows that

$$Af_{t_1+2}(H, x) = \sum_{M \in \mathcal{M}_{t_1+2}(H)} x^{af(H_2^T, M \cap E(H_2^T)) + 1} = \sum_{M \in \mathcal{M}(H_2^T)} x^{af(H_2^T, M)} \cdot x = xAf(H_2^T, x). \quad (6)$$

Substituting Eqs. (5,6) into Eq. (4), we immediately obtain Eq. (2) for  $t_1 = -1$ .

For  $t_1 \geq 1$ , given  $M \in \mathcal{M}_i(H)$  for  $i = 2, 3, \dots, t_1 + 1$ . Then  $r_{i+1}, l_i \notin M$ . On the one hand  $r_{i+1}$  and  $l_i$  belong to compatible  $M$ -alternating hexagons  $C_{i+1}$  and  $C_i$  respectively. And on the other hand  $\{r_{i+1}, l_i\}$  forces edges  $l_{i+1}, n_{i+1}, e_i, r_i, m_i$  and anti-forces edges  $e_{i+1}, m_{i+1}, n_i, e_{i-1}$ . By Lemma 2.4 and Fig. 3(c) we know

$$af(H, M) = af(H \ominus \{r_{i+1}, l_i\}, M \cap E(H \ominus \{r_{i+1}, l_i\})) + 2 = af(H_2^T, M \cap E(H_2^T)) + 2.$$

It follows that

$$Af_i(H, x) = \sum_{M \in \mathcal{M}_i(H)} x^{af(H_2^T, M \cap E(H_2^T)) + 2} = \sum_{M \in \mathcal{M}(H_2^T)} x^{af(H_2^T, M)} \cdot x^2 = x^2 Af(H_2^T, x). \quad (7)$$

For  $t_1 \geq 0$ , given  $M \in \mathcal{M}_1(H)$ . Then  $r_2 \notin M$ . On the one hand  $r_2$  belongs to  $M$ -alternating hexagon  $C_2$ . And on the other hand  $r_2$  forces edges  $l_2, n_2$  and anti-forces edges  $e_2, m_2$ . By Lemma 2.4 and Fig. 3(d) we know

$$af(H, M) = af(H \ominus r_2, M \cap E(H \ominus r_2)) + 1 = af(H_1^T, M \cap E(H_1^T)) + 1.$$

It follows that

$$\begin{aligned} Af_1(H, x) &= \sum_{M \in \mathcal{M}_1(H)} x^{af(H_1^T, M \cap E(H_1^T)) + 1} = \sum_{M \in \mathcal{M}(H_1^T), e_1 \in M} x^{af(H_1^T, M)} \cdot x \\ &:= xAf^*(H_1^T, x) = \sum_{M \in \mathcal{M}(H_1^T)} x^{af(H_1^T, M)} \cdot x - \sum_{M \in \mathcal{M}(H_1^T), e_1 \notin M} x^{af(H_1^T, M)} \cdot x. \end{aligned}$$

In  $H_1^T$ , given  $M \in \mathcal{M}(H_1^T)$  with  $e_1 \notin M$ . Then  $e_0 \in M$ . By a similar argument to the calculation of  $Af_0(H, x)$  we know

$$af(H_1^T, M) = af(H_1^T \ominus r_1, M \cap E(H_1^T \ominus r_1)) + 1 = af(H_3^T, M \cap E(H_3^T)) + 1.$$

It follows that

$$\begin{aligned} Af_1(H, x) &= xAf(H_1^T, x) - \sum_{M \in \mathcal{M}(H_1^T), e_1 \notin M} x^{af(H_3^T, M \cap E(H_3^T)) + 1} \cdot x \\ &= xAf(H_1^T, x) - \sum_{M \in \mathcal{M}(H_3^T)} x^{af(H_3^T, M)} \cdot x^2 = xAf(H_1^T, x) - x^2 Af(H_3^T, x). \quad (8) \end{aligned}$$

Substituting Eqs. (5-8) into Eq. (4), we immediately obtain Eq. (2) for  $t_1 \geq 0$ .

(2) If  $H$  contains a tail  $T(t_1, t_2, t_3)$  ( $t_1, t_2, t_3 \geq 0$ ), then we in turn denote the hexagon in  $T_1$  by  $B_i$  for  $i = 1, 2, \dots, t_1 + 2$  and the hexagon in  $T_2$  by  $D_j$  for  $j = 1, 2, \dots, t_2 + 2$  such that  $B_1 = D_1 = h$ . Furthermore, in turn denote the vertical edge that  $B_i$  and  $B_{i+1}$  (resp.  $D_j$  and  $D_{j+1}$ ) share in common by  $e_i$  (resp.  $e'_j$ ), and along the clockwise direction denote the rest edges of  $B_i$  (resp.  $D_j$ ) by  $l_i, r_i, e_{i-1}, m_i, n_i$  for  $i = 1, 2, \dots, t_1 + 2$  (resp.  $l'_j, r'_j, e'_{j-1}, m'_j, n'_j$  for  $j = 1, 2, \dots, t_2 + 2$ ) (see Fig. 2(c)). By Lemma 3.1 we can divide  $\mathcal{M}(H)$  in  $(t_1 + 2)(t_2 + 2) + 1$  subsets:

$$\mathcal{M}_{i,j}(H) = \{M \in \mathcal{M}(H) : e_i, e'_j \in M\}$$

for  $i = 0, 1, \dots, t_1 + 2$  and  $j = 0, 1, \dots, t_2 + 2$ , and  $i = 0$  if and only if  $j = 0$ . By Eq. (1), we have

$$\begin{aligned} Af(H, x) &= \sum_{M \in \mathcal{M}_{0,0}(H)} x^{af(H,M)} + \sum_{i=1}^{t_1+2} \sum_{j=1}^{t_2+2} \sum_{M \in \mathcal{M}_{i,j}(H)} x^{af(H,M)} \\ &:= Af_{0,0}(H, x) + \sum_{i=1}^{t_1+2} \sum_{j=1}^{t_2+2} Af_{i,j}(H, x). \end{aligned} \quad (9)$$

Given  $M \in \mathcal{M}_{t_1+2, t_2+2}(H)$ . Then  $l_{t_1+2}, l'_{t_2+2} \notin M$ . On the one hand  $l_{t_1+2}$  and  $l'_{t_2+2}$  belong to compatible  $M$ -alternating hexagons  $B_{t_1+2}$  and  $D_{t_2+2}$  respectively. And on the other hand  $\{l_{t_1+2}, l'_{t_2+2}\}$  forces edges  $e_{t_1+2}, r_{t_1+2}, m_{t_1+2}, e'_{t_2+2}, r'_{t_2+2}, m'_{t_2+2}$  and anti-forces edges  $e_{t_1+1}, n_{t_1+2}, e'_{t_2+1}, n'_{t_2+2}$ . By Lemma 2.4 and Fig. 4(a) we know

$$\begin{aligned} af(H, M) &= af(H \ominus \{l_{t_1+2}, l'_{t_2+2}\}, M \cap E(H \ominus \{l_{t_1+2}, l'_{t_2+2}\})) + 2 \\ &= af(H_2^T, M \cap E(H_2^T)) + 2. \end{aligned}$$

It follows that

$$\begin{aligned} Af_{t_1+2, t_2+2}(H, x) &= \sum_{M \in \mathcal{M}_{t_1+2, t_2+2}(H)} x^{af(H_2^T, M \cap E(H_2^T)) + 2} = \sum_{M \in \mathcal{M}(H_2^T)} x^{af(H_2^T, M)} \cdot x^2 \\ &= x^2 Af(H_2^T, x). \end{aligned} \quad (10)$$

Given  $M \in \mathcal{M}_{i, t_2+2}(H)$  for  $i = 2, 3, \dots, t_1 + 1$ . Then  $r_{i+1}, l_i, l'_{t_2+2} \notin M$ . On the one hand  $r_{i+1}, l_i$  and  $l'_{t_2+2}$  belong to compatible  $M$ -alternating hexagons  $B_{i+1}, B_i$  and  $D_{t_2+2}$  respectively. And on the other hand  $\{r_{i+1}, l_i, l'_{t_2+2}\}$  forces edges  $l_{i+1}, n_{i+1}, e_i, r_i, m_i, e'_{t_2+2}, r'_{t_2+2}, m'_{t_2+2}$  and anti-forces edges  $e_{i+1}, m_{i+1}, n_i, e_{i-1}, e'_{t_2+1}, n'_{t_2+2}$ . By Lemma 2.4 and Fig. 4(b) we know

$$\begin{aligned} af(H, M) &= af(H \ominus \{r_{i+1}, l_i, l'_{t_2+2}\}, M \cap E(H \ominus \{r_{i+1}, l_i, l'_{t_2+2}\})) + 3 \\ &= af(H_2^T, M \cap E(H_2^T)) + 3. \end{aligned}$$

It follows that

$$\begin{aligned} Af_{i, t_2+2}(H, x) &= \sum_{M \in \mathcal{M}_{i, t_2+2}(H)} x^{af(H_2^T, M \cap E(H_2^T)) + 3} = \sum_{M \in \mathcal{M}(H_2^T)} x^{af(H_2^T, M)} \cdot x^3 = x^3 Af(H_2^T, x). \end{aligned} \quad (11)$$






Fig. 4. (a)  $H \ominus \{l_{t_1+2}, l'_{t_2+2}\}$ , (b)  $H \ominus \{r_{i+1}, l_i, l'_{t_2+2}\}$ , (c)  $H \ominus \{r_2, l'_{t_2+2}\}$ , and (d)  $H \ominus \{r_2, r'_2\}$ .

Given  $M \in \mathcal{M}_{1,t_2+2}(H)$ . Then  $r_2, l'_{t_2+2} \notin M$ . On the one hand  $r_2$  and  $l'_{t_2+2}$  belong to compatible  $M$ -alternating hexagons  $B_2$  and  $D_{t_2+2}$  respectively. On the other hand  $\{r_2, l'_{t_2+2}\}$  forces edges  $l_2, n_2, e_1, e'_{t_2+2}, r'_{t_2+2}, m'_{t_2+2}$  and anti-forces edges  $e_2, m_2, e'_{t_2+1}, n'_{t_2+2}$ . By Lemma 2.4 and Fig. 4(c) we know

$$af(H, M) = af(H \ominus \{r_2, l'_{t_2+2}\}, M \cap E(H \ominus \{r_2, l'_{t_2+2}\})) + 2 = af(H_2^T, M \cap E(H_2^T)) + 2.$$

It follows that

$$\begin{aligned} Af_{1,t_2+2}(H, x) &= \sum_{M \in \mathcal{M}_{1,t_2+2}(H)} x^{af(H_2^T, M \cap E(H_2^T)) + 2} = \sum_{M \in \mathcal{M}(H_2^T)} x^{af(H_2^T, M)} \cdot x^2 \\ &= x^2 Af(H_2^T, x). \end{aligned} \quad (12)$$

By a similar argument to above, we know for  $i = 2, 3, \dots, t_1 + 1$  and  $j = 2, 3, \dots, t_2 + 1$

$$Af_{t_1+2,j}(H, x) = Af_{1,j}(H, x) = Af_{i,1}(H, x) = x^3 Af(H_2^T, x), \quad (13)$$

$$Af_{t_1+2,1}(H, x) = x^2 Af(H_2^T, x), \quad (14)$$

$$Af_{i,j}(H, x) = x^4 Af(H_2^T, x). \quad (15)$$

Given  $M \in \mathcal{M}_{1,1}(H)$ . Then  $r_2, r'_2 \notin M$ . On the one hand  $r_2$  and  $r'_2$  belong to compatible  $M$ -alternating hexagons  $B_2$  and  $D_2$  respectively. On the other hand  $\{r_2, r'_2\}$  forces edges  $l_2, n_2, l'_2, n'_2$  and anti-forces edges  $e_2, m_2, e'_2, m'_2$ . By Lemma 2.4 and Fig. 3(d) we know

$$af(H, M) = af(H \ominus \{r_2, r'_2\}, M \cap E(H \ominus \{r_2, r'_2\})) + 2 = af(H_1^T, M \cap E(H_1^T)) + 2.$$

By a similar argument to the calculation of Eq. (8) we have

$$\begin{aligned} Af_{1,1}(H, x) &= \sum_{M \in \mathcal{M}_{1,1}(H)} x^{af(H_1^T, M \cap E(H_1^T)) + 2} = \sum_{M \in \mathcal{M}(H_1^T), e_1, e'_1 \in M} x^{af(H_1^T, M)} \cdot x^2 \\ &= x^2 Af^*(H_1^T, x) = x^2 Af(H_1^T, x) - x^3 Af(H_3^T, x). \end{aligned} \quad (16)$$

It remains to consider  $Af_{0,0}(H, x)$ . Denote  $s$  the edge adjacent to  $n_1$  and  $m_1$ ,  $t$  the edge adjacent to  $e_0$  and  $m_1$ . Obviously there is a bijection  $g$  between  $\mathcal{M}_{0,0}(H)$  (see Fig. 5(a))

and  $\{M \in \mathcal{M}(H_2^T) : m_1 \in M\}$  (see Fig. 5(b)) through replacing the  $M$ -alternating path  $n_1 m_2 n_2 \cdots m_{t_1+2} n_{t_1+2} e_{t_1+2} l_{t_1+2} r_{t_1+2} \cdots l_2 r_2 l_1 m'_2 n'_2 \cdots m'_{t_2+2} n'_{t_2+2} e'_{t_2+2} l'_{t_2+2} r'_{t_2+2} \cdots l'_2 r'_2 e_0$  by  $g(M)$ -matched edge  $m_1$ . Given  $M \in \mathcal{M}_{0,0}(H)$  and a minimum anti-forcing set  $S$  of  $g(M)$  in  $H_2^T$ . Then  $m_1 \notin S$ . We claim that  $S \cup \{m_1\}$  is an anti-forcing set of  $M$ . Obviously every  $M$ -alternating cycle in  $T_1 \cup T_2$  must contain  $m_1$ , and every  $M$ -alternating cycle in  $H_2^T$  must contain some edge in  $S$ . If there is no other  $M$ -alternating cycles in  $H$ , then the claim holds. Otherwise, denote a such one by  $X$ . Then  $X$  must contain  $M$ -alternating paths  $n_1 s$  and  $e_0 t$ . Since  $X$  can be transformed into a  $g(M)$ -alternating cycle in  $H_2^T$  through replacing the  $M$ -alternating path in  $T_1 \cup T_2$  starting from  $n_1$  and ending at  $e_0$  by  $g(M)$ -matched edge  $m_1$ , we have that  $X$  must contain some edge in  $S$ . Then the claim holds in this case, which implies  $af(H, M) \leq af(H_2^T, g(M)) + 1$ .




Fig. 5. Illustration of calculation for  $Af_{0,0}(H, x)$ .

Given a maximum compatible  $g(M)$ -alternating set  $\mathcal{C}$  of  $H_2^T$ . We claim that  $H$  contains a compatible  $M$ -alternating set with cardinality  $|\mathcal{C}| + 1$ . Obviously every  $g(M)$ -alternating cycle in  $\mathcal{C}$  that does not contain  $m_1$  is also an  $M$ -alternating cycle of  $H$ , and the  $M$ -alternating cycle  $h$  is compatible with the above every such one in  $H$ . If  $\mathcal{C}$  does not contain  $m_1$ , then  $\mathcal{C} \cup \{h\}$  is a compatible  $M$ -alternating set of  $H$  and the claim holds. Otherwise, such one is unique since it contains  $s$ , hence we denote it by  $Y$ . Since  $M$ -alternating cycle  $Z$  in  $H$  through replacing  $m_1$  by  $n_1 m_2 n_2 e_2 l_2 r_2 l_1 m'_2 n'_2 e'_2 l'_2 r'_2 e_0$  is compatible with every member in  $\mathcal{C} \setminus \{Y\} \cup \{h\}$ , we have that  $\mathcal{C} \setminus \{Y\} \cup \{h, Z\}$  is a compatible  $M$ -alternating set in  $H$ . Then the claim holds in this case, which implies  $af(H, M) \geq af(H_2^T, g(M)) + 1$ . It follows that  $af(H, M) = af(H_2^T, g(M)) + 1$  and

$$Af_{0,0}(H, x) = \sum_{M \in \mathcal{M}_{0,0}(H)} x^{af(H_2^T, g(M)) + 1} = \sum_{M \in \mathcal{M}(H_2^T), m_1 \in M} x^{af(H_2^T, M)} \cdot x := x Af^*(H_2^T, x).$$

If  $t_3 = 0$ , then by a similar argument to the calculation of Eq. (8) and Fig. 5(c), we have

$$\begin{aligned} Af_{0,0}(H, x) &= \sum_{M \in \mathcal{M}(H_2^T)} x^{af(H_2^T, M)} \cdot x - \sum_{M \in \mathcal{M}(H_2^T), m_1 \notin M} x^{af(H_2^T, M)} \cdot x \\ &= x Af(H_2^T, x) - x^2 Af(H_4^T, x). \end{aligned} \quad (17)$$

If  $t_3 \geq 1$ , then by a similar argument to the above and Fig. 5(d), we have

$$Af_{0,0}(H, x) = x^2 Af(H_3^T, x). \quad (18)$$

Substituting Eqs. (10-18) into Eq. (9), we immediately obtain Eq. (3).  $\square$

Deng and Zhang [1] confirmed the continuity of anti-forcing spectrum for catacondensed hexagonal systems by analysing properties of graphs directly, and here we show the result by degrees of anti-forcing polynomial.

**Corollary 3.3.** [1] *The anti-forcing spectrum of a catacondensed hexagonal system  $H$  is continuous, namely it is an integer interval.*

*Proof.* We proceed it by induction on the number of hexagons  $m$  of  $H$ . If  $m = 1$ , then  $H = T(-1, -1, -1)$  and has anti-forcing polynomial  $2x$ . Thus the anti-forcing spectrum is  $\{1\}$ . Suppose that the result holds for the cases of less than  $m (\geq 2)$ . Now we consider the case of  $m$ . Pick a tail  $T(t_1, t_2, t_3)$  of  $H$ , where  $t_1 \geq 0$ . By assumption, the anti-forcing spectrum of  $H_k^T$  is continuous, say integer interval  $[i_k, j_k]$  for  $k = 1, 2, 3, 4$ .

Obviously there is a bijection  $g$  between  $\mathcal{M}(H_3^T)$  and  $\{M \in \mathcal{M}(H_2^T) : m_1 \in M\}$ , where  $g(M)$  is obtained from  $M$  by adding some  $g(M)$ -matched edges in  $T_3$  such that  $m_1$  is the vertical edge of  $T_3$  contained in  $g(M)$ . Since every compatible  $M$ -alternating set of  $H_3^T$  is also a compatible  $g(M)$ -alternating set of  $H_2^T$ , and the union of  $\{s\}$  and an anti-forcing set of  $M$  is an anti-forcing set of  $g(M)$ , we have that  $af(H_3^T, M) \leq af(H_2^T, g(M)) \leq af(H_3^T, M) + 1$ . Then the set of degrees of  $Af^*(H_2^T, x)$  is contained in  $[i_3, j_3 + 1]$ , and the set  $[i_2, j_2] \cup [i_3, j_3]$  of degrees of  $Af(H_2^T, x) + Af(H_3^T, x)$  is continuous.

**Case 1.**  $t_2 = -1$ . Denote  $\hat{H}$  the subgraph of  $H$  by deleting  $C_{t_1+2}$ . Hence  $\hat{H}$  is a catacondensed hexagonal system with  $m - 1$  hexagons and suppose its anti-forcing spectrum is an integer interval  $[a, b]$ .

**Subcase 1.1.**  $t_1 \geq 1$ . Pick the tail  $T(t_1 - 1, -1, t_3)$  of  $\hat{H}$  obtained from  $T(t_1, -1, t_3)$  by deleting  $C_{t_1+2}$ . By the calculation of Eq. (2) we have

$$\begin{aligned} Af(H, x) &= (x + t_1 x^2) Af(H_2^T, x) + x Af(H_3^T, x) + x Af^*(H_1^T, x), \\ Af(\hat{H}, x) &= (x + (t_1 - 1)x^2) Af(H_2^T, x) + x Af(H_3^T, x) + x Af^*(H_1^T, x) \\ &= Af(H, x) - x^2 Af(H_2^T, x). \end{aligned}$$

Obviously the degrees of  $x Af(H_2^T, x)$  and degrees of  $x^2 Af(H_2^T, x)$  form  $[i_2 + 1, j_2 + 1]$  and  $[i_2 + 2, j_2 + 2]$ , respectively. Since the existence of  $x Af(H_2^T, x)$  in  $Af(\hat{H}, x)$ , we know  $a \leq i_2 + 1 \leq j_2 + 1 \leq b$ . It follows that the anti-forcing spectrum of  $H$  is  $[a, b + 1]$  if  $j_2 + 1 = b$ , and  $[a, b]$  if  $j_2 + 1 < b$ .

**Subcase 1.2.**  $t_1 = 0$ . Then  $\hat{H} = H_1^T$  and by a similar argument to the calculation of Eq. (2) we have

$$\begin{aligned} Af(H, x) &= x Af(H_2^T, x) + x Af(H_3^T, x) + x Af^*(H_1^T, x), \\ Af(\hat{H}, x) &= x Af(H_3^T, x) + Af^*(H_1^T, x). \end{aligned}$$

By assumption, we know that the union of  $[i_3 + 1, j_3 + 1]$  and the set of degrees of  $Af^*(H_1^T, x)$  form  $[a, b]$ . We claim that either  $xAf(H_2^T, x)$  or  $xAf^*(H_1^T, x)$  has degree  $j_3 + 2$ . If the claim holds, then the anti-forcing spectrum of  $H$  is continuous since the union of set of degrees of  $xAf(H_2^T, x) + xAf(H_3^T, x)$  and  $\{j_3 + 2\}$  is an integer interval  $[i_2 + 1, j_2 + 1] \cup [i_3 + 1, j_3 + 2]$ .

Given a perfect matching  $M$  such that  $af(H_3^T, M) = j_3$ . If  $af(H_2^T, g(M)) = af(H_3^T, M) + 1$ , then  $xAf(H_2^T, x)$  has degree  $j_3 + 2$ . If  $af(H_2^T, g(M)) = af(H_3^T, M)$ , then  $af(H_3^T, M) + 2 = af(H, d(M))$ , where  $d(M) \in \mathcal{M}_1(H)$  is obtained from  $g(M)$  by adding  $l_2, n_2, e_1, r_1$ . This implies that  $xAf^*(H_1^T, x)$  has degree  $j_3 + 2$ . Then the claim holds.

**Case 2.**  $t_2 \geq 0$ .

**Subcase 2.1.**  $t_1 \geq 1$ . Denote  $\tilde{H}$  the subgraph of  $H$  by deleting  $B_{t_1+2}$ . Hence  $\tilde{H}$  is a catacondensed hexagonal system with  $m - 1$  hexagons. Pick the tail  $T(t_1 - 1, t_2, t_3)$  of  $\tilde{H}$  obtained from  $T(t_1, t_2, t_3)$  by deleting  $B_{t_1+2}$ . By the calculation of Eq. (3) we have

$$\begin{aligned} Af(H, x) &= (3x^2 + 2(t_1 + t_2)x^3 + t_1t_2x^4)Af(H_2^T, x) + x^2Af^*(H_1^T, x) + xAf^*(H_2^T, x), \\ Af(\tilde{H}, x) &= (3x^2 + 2(t_1 + t_2 - 1)x^3 + (t_1 - 1)t_2x^4)Af(H_2^T, x) + x^2Af^*(H_1^T, x) \\ &\quad + xAf^*(H_2^T, x) \\ &= Af(H, x) - (2x^3 + t_2x^4)Af(H_2^T, x). \end{aligned}$$

By a similar argument to Subcase 1.1, if  $t_2 = 0$ , then the anti-forcing spectrum of  $H$  is continuous since the existence of  $3x^2Af(H_2^T, x)$  in  $Af(\tilde{H}, x)$ ; and if  $t_2 \geq 1$ , then the anti-forcing spectrum of  $H$  is continuous since the existence of  $2(t_1 + t_2 - 1)x^3Af(H_2^T, x)$  in  $Af(\tilde{H}, x)$ .

**Subcase 2.2.**  $t_1 = 0$  and  $t_2 \geq 1$ . Denote  $\tilde{H}$  the subgraph of  $H$  by deleting  $D_{t_2+2}$ . By a similar to Subcase 2.1, we know that the anti-forcing spectrum of  $H$  is continuous.

**Subcase 2.3.**  $t_1 = t_2 = 0$ . Denote  $\tilde{H}$  the subgraph of  $H$  by deleting  $D_2$ . Hence  $\tilde{H}$  is a catacondensed hexagonal system with  $m - 1$  hexagons and suppose its anti-forcing spectrum is  $[p, q]$ . Pick the tail  $T(0, -1, t_3)$  of  $\tilde{H}$  obtained from  $T(0, 0, t_3)$  by deleting  $D_2$ . By the calculation of Eqs. (2,3) we have

$$\begin{aligned} Af(H, x) &= 3x^2Af(H_2^T, x) + x^2Af^*(H_1^T, x) + xAf^*(H_2^T, x), \\ Af(\tilde{H}, x) &= xAf(H_2^T, x) + xAf^*(H_1^T, x) + xAf(H_3^T, x). \end{aligned}$$

By assumption, we know that the union of the set of degrees of  $xAf(H_2^T, x) + xAf^*(H_1^T, x)$  and  $[i_3 + 1, j_3 + 1]$  form  $[p, q]$ . Note that the set of degrees of  $xAf^*(H_2^T, x)$  is contained in  $[i_3 + 1, j_3 + 2]$ . We claim that either  $xAf^*(H_2^T, x)$  or  $3x^2Af(H_2^T, x)$  has degree  $w + 2$  for every  $w \in [i_3, j_3]$ . If the claim holds, then the anti-forcing spectrum of  $H$  is continuous since the union of set of degrees of  $3x^2Af(H_2^T, x) + x^2Af^*(H_1^T, x)$  and  $[i_3 + 2, j_3 + 2]$  is an integer interval  $[p + 1, q + 1]$ .

Given a perfect matching  $M$  such that  $af(H_3^T, M) = w$ . If  $af(H_2^T, g(M)) = af(H_3^T, M) + 1$ , then  $xAf^*(H_2^T, x)$  has degree  $w + 2$ . If  $af(H_2^T, g(M)) = af(H_3^T, M)$ , then  $3x^2Af(H_2^T, x)$  has degree  $w + 2$ . Then the claim holds.  $\square$

We now give some anti-forcing polynomials of particular catacondensed hexagonal systems. Given a hexagonal chain with  $n(\geq 1)$  maximal linear hexagonal chains and in which containing  $r_1 + 2, r_2 + 2, \dots, r_n + 2$  ( $r_i \geq 0$  for  $i = 1, 2, \dots, n$ ) hexagons in turn (see Fig. 1(b)). For convenience, we denote it by  $Hl(r_1, r_2, \dots, r_n)$ , and make a convention that  $Hl(r_1, r_2, \dots, r_i, -1) = Hl(r_1, r_2, \dots, r_i)$  for  $i \geq 0$ ,  $Hl(r_1, \dots, r_0)$  is a hexagon, and  $Hl(r_1, \dots, r_{-1} - 1) = Hl(r_1, \dots, r_0 - 1)$  is an empty graph. Pick the tail  $T(r_n, -1, t_3)$ , where  $t_3 = -1$  if  $n = 1$  and  $t_3 = r_{n-1}$  if  $n \geq 2$ . Then  $H_1^T = Hl(r_1, r_2, \dots, r_{n-1})$ ,  $H_2^T = Hl(r_1, r_2, \dots, r_{n-1} - 1)$ , and  $H_3^T = Hl(r_1, r_2, \dots, r_{n-2} - 1)$ . Then we have the following conclusion.

**Corollary 3.4.** [2] *The anti-forcing polynomial of a hexagonal chain  $Hl(r_1, r_2, \dots, r_n)$  with at least two hexagons is*

$$\begin{aligned} Af(Hl(r_1, \dots, r_n), x) = & (x + r_n x^2) Af(Hl(r_1, \dots, r_{n-1} - 1), x) \\ & + (x - x^2) Af(Hl(r_1, \dots, r_{n-2} - 1), x) + x Af(Hl(r_1, \dots, r_{n-1}), x), \end{aligned}$$

where  $Af(Hl(r_1, r_2, \dots, r_i, -1), x) = Af(Hl(r_1, r_2, \dots, r_i), x)$  for  $i \geq 0$ ,  $Af(Hl(r_1, \dots, r_0), x) = 2x$ , and  $Af(Hl(r_1, \dots, r_{-1} - 1), x) = Af(Hl(r_1, \dots, r_0 - 1), x) = 1$ .

Denote the anti-forcing polynomials of graphs illustrated in Figs. 6(a-c) by  $P_n, Q_n, R_n$  ( $n \geq 1$ ), where the third graph consists of  $n$  rows of maximal linear hexagonal chains of length two, the second graph is obtained from the third one by adding a hexagon above the first row or below the last row, and the first graph is obtained from the third one by adding a hexagon above the first row and adding a hexagon below the last row.




Fig. 6. Examples of catacondensed hexagonal system.

**Example 3.5.**  $P_n, Q_n, R_n$  have the following recurrence relations:

$$P_n = x^2 P_{n-1} + (x + 3x^2) Q_{n-1} - 2x^4 Q_{n-2} - 2x^3 Q_{n-3} \quad \text{for } n \geq 3, \quad (19)$$

$$Q_n = x^2 Q_{n-1} + (x + 3x^2) R_{n-1} - 2x^4 R_{n-2} - 2x^3 R_{n-3} \quad \text{for } n \geq 3, \quad (20)$$

$$Q_n = x P_{n-1} + x Q_{n-1} + (2x^2 - 2x^3) Q_{n-2} \quad \text{for } n \geq 2, \quad (21)$$

$$R_n = x Q_{n-1} + x R_{n-1} + (2x^2 - 2x^3) R_{n-2} \quad \text{for } n \geq 2, \quad (22)$$

where  $P_0 = x^2 + 2x$ ,  $P_1 = x^4 + 7x^3 + x^2$ ,  $P_2 = x^6 + 6x^5 + 11x^4 + 6x^3$ ,  $Q_0 = 2x$ ,  $Q_1 = x^3 + 3x^2 + x$ ,  $Q_2 = x^5 + 4x^4 + 8x^3 + x^2$ ,  $R_0 = 1$ ,  $R_1 = x^2 + 2x$ ,  $R_2 = x^4 + 2x^3 + 5x^2$ .

Furthermore, for  $n \geq 0$  the explicit expression of  $R_n$  is

$$\begin{aligned}
R_n = & \sum_{k=0}^n \sum_{a=0}^{\lfloor \frac{n-k}{2} \rfloor} \sum_{c=\lceil \frac{n-k-2a}{2} \rceil}^{\lfloor \frac{n-k-a}{2} \rfloor} \binom{k+a}{a} \binom{a}{n-k-a-2c} \binom{k+c}{c} 2^c x^{k+2a+2c} \\
& + \sum_{k=0}^{n-1} \sum_{a=0}^{\lfloor \frac{n-k-1}{2} \rfloor} \sum_{c=\lceil \frac{n-k-2a-1}{2} \rceil}^{\lfloor \frac{n-k-a-1}{2} \rfloor} \binom{k+a}{a} \binom{a}{n-k-a-2c-1} \binom{k+c}{c} 2^c x^{k+2a+2c+1} \\
& - \sum_{k=0}^{n-2} \sum_{a=0}^{\lfloor \frac{n-k-2}{2} \rfloor} \sum_{c=\lceil \frac{n-k-2a-2}{2} \rceil}^{\lfloor \frac{n-k-a-2}{2} \rfloor} \binom{k+a}{a} \binom{a}{n-k-a-2c-2} \binom{k+c}{c} 2^c x^{k+2a+2c+3} \\
& - \sum_{k=0}^{n-3} \sum_{a=0}^{\lfloor \frac{n-k-3}{2} \rfloor} \sum_{c=\lceil \frac{n-k-2a-3}{2} \rceil}^{\lfloor \frac{n-k-a-3}{2} \rfloor} \binom{k+a}{a} \binom{a}{n-k-a-2c-3} \binom{k+c}{c} 2^c x^{k+2a+2c+3}, \quad (23)
\end{aligned}$$

which can be applied to derive explicit expressions of  $P_n$  and  $Q_n$  by Eqs. (21,22).

*Proof.* The recurrence relations Eqs. (19,20) (resp. Eqs. (21,22)) can be obtained from Eq. (3) (resp. Eq. (2)) directly. Furthermore from Eq. (22) for  $n \geq 2$  we can derive

$$xQ_{n-1} = R_n - xR_{n-1} + (2x^3 - 2x^2)R_{n-2}.$$

Substituting it into Eq. (20), for  $n \geq 4$  we have the following recurrence relation:

$$R_n - (x + x^2)R_{n-1} - 3x^2R_{n-2} + 2x^4R_{n-3} + 2x^4R_{n-4} = 0.$$

Define the generating function of  $R_n$  as  $W = \sum_{n=0}^{\infty} R_n t^n$ . By the above equation we have

$$\sum_{n=4}^{\infty} R_n t^n - (x + x^2) \sum_{n=4}^{\infty} R_{n-1} t^n - 3x^2 \sum_{n=4}^{\infty} R_{n-2} t^n + 2x^4 \sum_{n=4}^{\infty} R_{n-3} t^n + 2x^4 \sum_{n=4}^{\infty} R_{n-4} t^n = 0.$$

Substituting  $R_0 = 1$ ,  $R_1 = x^2 + 2x$ ,  $R_2 = x^4 + 2x^3 + 5x^2$ , and  $R_3 = x^6 + 3x^5 + 8x^4 + 10x^3$  into it, we have

$$\begin{aligned}
& W - 1 - (x^2 + 2x)t - (x^4 + 2x^3 + 5x^2)t^2 - (x^6 + 3x^5 + 8x^4 + 10x^3)t^3 \\
& - (x + x^2)t[W - 1 - (x^2 + 2x)t - (x^4 + 2x^3 + 5x^2)t^2] \\
& - 3x^2t^2[W - 1 - (x^2 + 2x)t] + 2x^4t^3[W - 1] + 2x^4t^4W = 0.
\end{aligned}$$

Simplifying the above equation gives

$$W \left[ 1 - \frac{xt}{(1 - x^2t - x^2t^2)(1 - 2x^2t^2)} \right] = \frac{1 + xt - x^3t^2 - x^3t^3}{(1 - x^2t - x^2t^2)(1 - 2x^2t^2)}.$$

This yields

$$\begin{aligned} \sum_{n=0}^{\infty} R_n t^n &= \sum_{k=0}^{\infty} \frac{x^k t^k (1 + xt - x^3 t^2 - x^3 t^3)}{(1 - x^2 t - x^2 t^2)^{k+1} (1 - 2x^2 t^2)^{k+1}} \\ &= \sum_{k=0}^{\infty} x^k t^k (1 + xt - x^3 t^2 - x^3 t^3) \sum_{a=0}^{\infty} \binom{k+a}{a} \sum_{b=0}^a \binom{a}{b} x^{2a} t^{a+b} \sum_{c=0}^{\infty} \binom{k+c}{c} 2^c x^{2c} t^{2c}. \end{aligned}$$

Extracting the coefficient of  $t^n$  from both sides of the above equation gives Eq. (23).  $\square$

## References

- [1] K. Deng, H. Zhang, Anti-forcing spectrum of any cata-condensed hexagonal system is continuous, *Front. Math. China* **12** (2017) 19–33.
- [2] H.-K. Hwang, H. Lei, Y.-N. Yeh, H. Zhang, Distribution of forcing and anti-forcing numbers of random perfect matchings on hexagonal chains and crowns, preprint, 2015. <http://140.109.74.92/hk/?p=873>.
- [3] D.J. Klein, V. Rosenfeld, Forcing, freedom, & uniqueness in graph theory & chemistry, *Croat. Chem. Acta* **87** (2014) 49–59.
- [4] H. Lei, Y.-N. Yeh, H. Zhang, Anti-forcing numbers of perfect matchings of graphs, *Discrete Appl. Math.* **202** (2016) 95–105.