

Scaling Behavior of the Hirsch Index for Failure Avalanches, Percolation Clusters and Paper Citations

Asim Ghosh,^{1,*} Bikas K. Chakrabarti,^{2,3,†} Dachepalli R. S. Ram,^{4,‡} Manipushpak Mitra,^{3,§} Raju Maiti,^{3,¶} Soumyajyoti Biswas,^{4,**} and Suchismita Banerjee^{3,††}

¹*Department of Physics, Raghunathpur College, Raghunathpur, Purulia - 723133, West Bengal, India.*

²*Saha Institute of Nuclear Physics, Kolkata 700064, India.*

³*Economic Research Unit, Indian Statistical Institute, Kolkata 700108, India.*

⁴*Department of Physics, SRM University-AP, Andhra Pradesh - 522240, India*

A popular measure for citation inequalities of individual scientists has been the Hirsch index (h). If for any scientist the number n_c of citations is plotted against the serial number n_p of the paper having those many citations (when the papers are ordered from highest cited to lowest) then h corresponds to the nearest lower integer value of n_p below the fixed point of the non-linear citation function (or given by $n_c = h = n_p$ if both n_p and n_c are dense set of integers near the h value). The same index can be estimated (from $h = s = n_s$) for the avalanche or cluster of size (s) distributions (n_s) in elastic fiber bundle or percolation models. Another such inequality index, called the Kolkata index (k) says that $(1 - k)$ fraction of papers attract k fraction of citations ($k = 0.80$ corresponds to the 80-20 law of Pareto). We find, for stress (σ), lattice occupation probability (p) or Kolkata index (k) near the bundle failure threshold (σ_c) or percolation threshold (p_c) or critical value of Kolkata index k_c , good fit to Widom-Stauffer like scaling $h/[\sqrt{N}/\log N] = f(\sqrt{N}[\sigma_c - \sigma]^\alpha)$, $h/[\sqrt{N}/\log N] = f(\sqrt{N}[p_c - p]^\alpha)$ or $h/[\sqrt{N_c}/\log N_c] = f(\sqrt{N_c}[k_c - k]^\alpha)$ respectively, with asymptotically defined scaling function f , for systems of size N (total number of fibers or lattice sites) or N_c (total number of citations), and α denoting the appropriate scaling exponent. We also show that if the number (N_m) of members of parliaments or national assemblies of different countries (with population N) is identified as their respective h -index, then the data fit the scaling relation $N_m \sim \sqrt{N}/\log N$, resolving a major recent controversy.

I. INTRODUCTION

Monotonically and non linearly decaying inequality functions are ubiquitous. When the number (n_p) of papers by any author (or, for that matter, an institution) are arranged according to the number (n_c) of the citations they received, the citation inequality function becomes a monotonically decaying one (see e.g., [1]). The same is true for avalanches in materials failure or in earthquakes (see e.g., [2]), cluster size distributions in percolation problems (see e.g., [3]), etc., where the number (n_s) of avalanches or clusters (giving the size inequality function) decrease monotonically and nonlinearly with the size (s) of the avalanche or cluster. Large avalanches, strong quakes, or big size clusters come or occur in small numbers, while the smaller or weaker the avalanches or quakes, or clusters, the larger are their abundance in occurrence. How does one statistically measure these inequalities in occurrence frequencies of citation numbers or avalanche or cluster sizes? Obviously, the corresponding distribution functions for inequalities in citations or sizes would contain the entire statistics. However, they are not convenient to handle. One can consider the citation number of the best cited paper (as sometimes done for some unique awards, etc.), or study the statistical (self-similar) structure of the biggest avalanche or the largest (percolating) cluster (as in statistical physics [3]). Hirsch proposed [1] an index to measure these inequalities, by locating the fixed point of the nonlinear inequality function. The Hirsch index (h) corresponds to the citation number or occurrence frequency which is commensurate in magnitude with the number of publications or avalanche cluster sizes.

* Email: asimghosh066@gmail.com

† Email: bikask.chakrabarti@saha.ac.in

‡ Email: dachepalli_ravi@srmmap.edu.in

§ email: manipushpak.mitra@gmail.com; corresponding author

¶ Email: rajumaiti@gmail.com

** Email: soumyajyoti.b@srmmap.edu.in

†† Email: suchib.1993@gmail.com

Systems near their critical points, self-organized or tuned, have mostly been studied in the self-similar limit of their divergent correlations (see e.g., [2, 3]). The corresponding critical exponent values arising out of such self-similarities have helped to classify vastly different physical systems based on their symmetries, dimensionality and such broad qualifiers. One powerful tool has been the scaling relations among the critical exponents that helped to build an interconnected and precise relation between experimental observables in such systems near criticality. In this work, however, we focus on quantifying the response of near-critical systems through the corresponding inequality statistics (for example, in n_c versus n_p of citations or in n_s versus s for avalanches or clusters). Specifically, we measure the widely used Hirsch index (h), which in effect, gives a measure of a size that is commensurate with its relative abundance. It turns out to be possibly even more robust than the critical behavior (characterized by a set of exponents). We demonstrate this by choosing a wide variety of systems: citation statistics, percolation cluster statistics and avalanche statistics in Fiber Bundle Models (FBMs) and even in the statistics of Parliament sizes in different countries of the world. They differ in their dimensionality (two dimensions for the percolation model studied here, mean-field for the fracture model, possibly small-world networks for citation and parliament statistics). They further differ in their measurement variables, and their size distribution statistics, making them widely different in terms of their prominently apparent features. However, we show here that in spite of their obvious differences, the scaling behavior of the Hirsch index shows remarkable universality.




FIG. 1. A schematic drawing of the citation function of a typical scientist. h -index (an integer) is given by lower value of the paper serial number below the fixed point value (intersection point of the 45 degree line from the origin), when the papers are ordered from highest cited to the lowest cited one. When the citation number and the serial number of the paper (having those citations) are both sequential integers near the fixed point, the citation number equals the number of papers and both become h . Similar will be the case where citations are replaced by the failure of the materials avalanche sizes (or cluster sizes) and the paper numbers are replaced by the number of such avalanches (or of the clusters in pre- or post-percolating systems).

How does the h index scale with a total number of publications (N_p) by the author (institution) or the total number (N_c) of citations received by the author (institution)? Young [4] suggested analytically that h -index value should scale with the total number of citations N_c ($= \sum n_c$) as $\sqrt{N_c}$ asymptotically. In ref. [5], a brief analysis of the Google Scholar data indicated independently that $h \sim \sqrt{N_p}$ for large values of N_p which corresponds to the highest value of n_p , the total number of publications by the author. Note that if both these relationships are valid then, statistically speaking, for a prolific author, the total number of citations would be linearly proportional to the total number

of papers published and the proportionality constant would perhaps be determined by the size of the existing author network in the subject (suggesting an effective Dunbar number [7, 8], for the community of authors).

In a recent Monte Carlo study [9] on the avalanche sizes and their numbers in the Fiber Bundle Models (FBM) of materials failure (see e.g., [2, 10]) due to increasing stress on such bundles, the numerical analysis of the data for the nonlinearly decaying numbers of avalanches with their sizes (or released elastic energies) suggested $h \sim \sqrt{N}/\log N$.

Our numerical study here of avalanche size distributions in FBMs (for stress or load per fiber σ less than its global failure stress σ_c), of cluster size distributions for lattice occupation concentration p near the percolation point p_c (for p both below and above p_c) in the percolating system and the previous analysis [5] of citation distributions of scientists with Kolkata index k (see e.g., [6] for a review, giving the fraction k of citations/wealth attracted/possessed by $1 - k$ fraction of publications /people) near (both above and below) the threshold point $k_c (= 0.86)$, all show excellent fit to a Widom-Stauffer like scaling relation between the Hirsch index (h) and system size N (or individual's total citation size N_c), following Widom Scaling for the free energy away from the critical point and the subsequent Stauffer Scaling [3] for the number of a particular sized cluster, identified here as the equivalent Hirsch index, at and away from the percolation point),

$$\frac{h}{[\sqrt{N}/\log N]} = f(\sqrt{N}[\sigma_c - \sigma]^\alpha) \quad (1a)$$

$$\frac{h}{[\sqrt{N}/\log N]} = f(\sqrt{N}|p_c - p|^\alpha) \quad (1b)$$

$$\frac{h}{[\sqrt{N}/\log N]} = f(\sqrt{N_c}|k_c - k|^\alpha), \quad (1c)$$

with asymptotically well-defined finite size scaling function f ($f(x) = \text{constant}$ at $x = 0$ and f remaining continuous and finite as x approaches infinity) for systems of size N (total number of fibers or lattice sites/bonds) or size N_c (total number of citations of all the publications by an individual scientist), and α denoting the appropriate scaling exponent.

Traditionally the Hirsch index h for different authors have been fitted [4] to the scaling form $\sqrt{N_c}$. For cases where the Kolkata index values of the authors are not known, we simply fitted, following the scaling relation (1) to the form appropriate for the critical point

$$h \sim \sqrt{N_c}/\log N_c \quad (2a)$$

$$\sim \sqrt{N_p}/\log N_p. \quad (2b)$$

We also show that if the number (N_m) of members of the parliaments or national assemblies of different countries (with the corresponding population denoted by N) are identified as those countries' respective h indices, then the data fit well to the scaling relation $N_m \sim \sqrt{N}/\log N$. This helps to comprehend the discrepancies (c.f. [11]) with the \sqrt{N} relationship, reported in a very recent analysis [12], resolving a recent major controversy.

II. DATA ANALYSIS & NUMERICAL STUDIES

A. Failure avalanches in fiber bundle model (FBM)

The fiber bundle model (FBM) is a generic model for failure of disordered solids. An ensemble of N fibers are set between two rigid parallel plates and a load is applied on the bottom plate. Each fiber is linear elastic, with the same elastic constant and has a failure threshold selected randomly from a distribution. This failure threshold is the source of disorder and non-linearity in the otherwise linear model.




FIG. 2. Finite size scaling analysis of the fiber bundle model h -index (from avalanche size distribution). The scaling fit to Widom-Stauffer relations (1) are obtained by making all the data points of different sizes ($N = 1000, 5000, 10000$ and 100000) and stress level (different σ values) collapse together. It turns out that the scaling form (1b) with exponent $\alpha = 1$ gives good data collapse. The inset shows that the data collapse clearly gets worsened by dropping the $\log N$ term in the scaling relation (1b).

When a small load (W) is applied, the weakest fiber breaks, and the load carried by that fiber is shared equally by all the remaining fibers, which can trigger further failures. Through gradual increase of load, therefore, the model goes through intermittent stable states, which are subsequently perturbed by increasing the load slowly. In going from one stable state to another, the number of fibers that break is the avalanche size (s). The size distribution of this follows a power-law statistics $P(s) \sim s^{-5/2}$ for $s \rightarrow \infty$ [13]. The avalanche dynamics continues until the load per fiber value $\sigma = W/N$ reaches a critical limit σ_c , when the entire system breaks down.

The avalanches are arranged in the ascending order to estimate h index values for stress level σ below σ_c . It was shown ([9]) that the terminal value of h ($= h_f$) at the critical point ($\sigma = \sigma_c$) follows a scaling relation $h_f \sim \sqrt{N}/\log(N)$. Here we look for the scaling of h with the critical interval $(\sigma_c - \sigma)$. It turns out that the scaling relation (1a) fits very well with our numerical the data (see Fig. 2). We also attempted the scaling fit without the $\log(N)$ term in the above mentioned relation (see inset of Fig. 2) and it clearly worsens the data collapse.

The Fig.3 shows the scaling behavior for the average h -index (in the range $4 \leq h \leq 20$) at the breaking point ($\sigma = \sigma_c$) of the bundles with total number N (in the range $200 \leq N \leq 20000$) of fibers in the equal-load-sharing FBM (with uniform distribution of fiber breaking thresholds) considered here (cf. [9]). The Fig. shows the best fit of h to $\sqrt{N}/\log N$.




FIG. 3. Scaling behavior for the average h -index (in the range $4 \leq h \leq 20$) at the breaking point ($\sigma = \sigma_c$) of the bundles with total number N (in the range $200 \leq N \leq 20000$) of fibers in the equal-load-sharing FBM (with uniform distribution of fiber breaking thresholds) considered here (cf. [9]). The Fig. shows the best fit of h to $\sqrt{N}/\log N$. The inset shows statistical deviations $\Sigma = \sqrt{\sum_i (h_d(i) - h_\gamma(i))^2 / \sum_i 1}$ to the scaling forms $h \sim N^\gamma$ and $h \sim N^\gamma/\log(N)$ with respect to γ , where i represents the individual (random) realization of the FBM with the corresponding h ($= h_d$) value obtained from the simulation result and the scaling fit (h_γ) from the above mentioned scaling relations. The inset shows that the statistical deviation becomes minimum for the scaling with log correction term with the appropriate value of γ near $1/2$.




FIG. 4. Finite size scaling analysis of the 2D site percolation h -index (from the cluster size distribution). The scaling fit to Widom-Stauffer relations (1) are obtained by making all the data points of different sizes ($N = 4000 \times 4000, 8000 \times 8000$ and 16000×16000) and concentrations (different p values, $p_c = 0.5927$) collapse together. It turns out that the scaling form (1b) with exponent $\alpha = 3$ gives good data collapse. The inset shows that the data collapse seems to get worsened by dropping the $\log N$ term in the scaling relation (1b).

B. Percolation clusters in 2D lattice

Here we consider two-dimensional (2D) site percolation in a square lattice with site occupancy probability p ($0 < p < 1$). For a given p , we measure cluster distribution, and hence the h -index is computed (here h -index measures h number of clusters each having cluster sizes greater than equal to h). In our simulation, we took four different system sizes ($N = 4000 \times 4000, 8000 \times 8000$ and 16000×16000) and h -index was estimated for different p values.

Here we carry out the finite size scaling analysis [14] (see also [15]) of the h index for different system sizes. The critical exponent α is determined by fit to scaling relation (1b) for all the data points of different sizes ($N = 4000 \times 4000, 8000 \times 8000$ and 16000×16000). We obtain good data collapse with $\alpha = 3$ and different scaling function f for $p > p_c$ and $p < p_c$ while they converge to the same value at $p = p_c$ (see Fig 4). The inset of Fig. 4 shows the scaling fit to relation (1b) without the $\log(N)$ term and it is observed that right hand side scaling fit is not that well.

The cluster size distribution in the percolation problem has also been studied for estimating the h -index scaling with the total number N of lattice sites at the percolation threshold of site percolation on square lattice. We of course find here the best fit scaling form to be $h \sim \sqrt{N}/\log N$ (see Fig. 5).




FIG. 5. Scaling behavior for the h -index (in the range $185 \leq h \leq 4421$) for cluster size distributions at the percolation point (with size N on square lattice; $10^6 < N < 10^9$). The Fig. shows the best fit of h to $\sqrt{N}/\log N$. The inset shows the statistical deviations $\Sigma = \sqrt{\Sigma_i (h_d(i) - h_\gamma(i))^2 / \Sigma_i 1}$ to the scaling forms $h \sim N^\gamma$ and $h \sim N^\gamma / \log(N)$ with respect to γ , where i represents the individual (random) realization of the percolating system with the corresponding h ($= h_d$) value obtained from the simulation result and the scaling fit (h_γ) from the above mentioned scaling relations. The inset shows that the statistical deviation becomes minimum for the scaling with log correction term, though with a value of γ somewhat above $1/2$.

C. Paper citations

We first analyze the data for h -index and its scaling with the total number of publications N_p and of citations N_c for the one hundred scientists (in mathematics, physics, chemistry, medicine, biology, economics, sociology; including those of twenty Nobel Laureates in those subjects), given in ref. [5]. Next, we collected (in May-June, 2021) the same kind of data¹ for one thousand scientists (mostly physicists) in all the above-mentioned subjects from Google Scholar.

Figs. 6 and 7 show the scaling behavior for the h -index (in the range $20 \leq h \leq 222$) of the 100 scientists (data taken from [5]), with the total number of citations N_c (in the range $1819 \leq N_c \leq 323473$). Our analysis shows the best fit to $h \sim \sqrt{N_c}/\log N_c$.

¹ The data will be available on request to corresponding author




FIG. 6. Finite size (N_c) scaling analysis of the citation distributions of 100 individual scientists (table 1 of Ref. [5]). The scaling fit to Widom-Stauffer relation (1c) is obtained by fitting all the data for their h index and the N_c values of total citations of the publications by individual scientists and their corresponding Kolkata index values k near the critical value $k_c \simeq 0.87$ [5]. It turns out that the scaling form (1c) with exponent $\alpha = 0.7$ gives good data collapse. The inset shows that the data collapse seems to get worsened by dropping the $\log N$ term in the scaling relation (1c), though at the critical point $k = k_c$, one gets $h = a\sqrt{N_c}$, with $a = 0.45 \pm 0.10$, which includes the analytical prediction [4] that $a = 0.54$ in the large N_c limit. The continuous lines (in both the main figure and the inset) represent the scaling functions for $k < k_c$ (green) and $k > k_c$ (violet), and are drawn for a guide to the eye.




FIG. 7. Scaling behavior for the h -index (in the range $20 \leq h \leq 222$) of the 100 scientists (data taken from [5]), with the total number of citations N_c (in the range $1819 \leq N_c \leq 323473$). Our analysis shows the best fit to $h \sim \sqrt{N_c}/\log N_c$. The inset shows the statistical deviations $\Sigma = \sqrt{\sum_i (h_d(i) - h_\gamma(i))^2 / \sum_i 1}$ to the scaling forms $h \sim N_c^\gamma$ and $h \sim N_c^\gamma/\log(N_c)$ with respect to γ , where i represents the individual scientist with the corresponding h ($= h_d$) obtained from Google Scholar and the scaling fit (h_γ) obtained from the above mentioned scaling relations. The inset shows that the statistical deviation becomes minimum for the scaling with log correction term with the value of γ a little higher than $1/2$.

The Figs 8(a,b) scaling behavior for the h -index (in the range $17 \leq h \leq 221$) of the 1000 scientists with the total number of citations N_c (in the range $996 \leq N_c \leq 348680$) in (a) and with the total number of papers N_p (in the range $100 \leq N_p \leq 2987$) in (b). The data are taken from Google Scholar in June 2021. The Figs. show the best fits to $h \sim \sqrt{N_c}/\log N_c$ in (a) and to $h \sim \sqrt{N_p}/\log N_p$ in (b).




FIG. 8. Scaling behavior for the h -index (in the range $17 \leq h \leq 221$) of the 1000 scientists with the total number of citations N_c (in the range $996 \leq N_c \leq 348680$) in (a) and with the total number of papers N_p (in the range $100 \leq N_p \leq 2987$) in (b). The data are taken from Google Scholar in June 2021. The Figs. show the best fits to $h \sim \sqrt{N_c}/\log N_c$ in (a) and to $h \sim \sqrt{N_p}/\log N_p$ in (b). The insets show the statistical deviations $\Sigma = \sqrt{\sum_i (h_d(i) - h_\gamma(i))^2 / \sum_i 1}$ to the scaling forms $h \sim N_p^\gamma$ (or $h \sim N_c^\gamma$) and $h \sim N_p^\gamma/\log(N_p)$ (or $h \sim N_c^\gamma/\log(N_c)$) with respect to γ , where i represents the individual scientist with the corresponding h ($= h_d$) value obtained from Google Scholar and the scaling fit (h_γ) from the above mentioned scaling relations. The inset shows that though the statistical deviation does not become minimum for the scaling with log correction term (presumably due to inclusion of some tail-end points), the best fit for γ assumes the desired value $1/2$.

D. Number of the representatives in the National Assemblies

Finally, we note that if the number N_m of the representatives in the National Assemblies or in the Parliaments of different countries of the world are identified as the h -index for respective country, having population N , then we find (see Fig. 9) that the data analyzed in ref [11]² shows the best fit of N_m to $\sqrt{N}/\log N$. This also resolves the discrepancy noted in the analysis if the same data ([12] and references therein).




FIG. 9. Scaling behavior for the number N_m (in the range $15 \leq N_m \leq 3040$) of the representatives in National assemblies or Parliaments of different countries of the world (1972 data from the original study [11]; see also [12]) with total population N (in the range $19 \times 10^4 \leq N \leq 70.5 \times 10^7$) of the respective countries. When N_m is identified as the h -index for the countries, we get $N_m \sim \sqrt{N}/\log N_m$ which is fitted here with the data. The inset shows the statistical deviations $\Sigma = \sqrt{\sum_i (N_{md}(i) - N_{m\gamma}(i))^2 / \sum_i 1}$ to the scaling forms $N_m \sim N^\gamma$ and $N_m \sim N^\gamma / \log(N)$ with respect to γ , where i represents the individual country with the corresponding members of parliaments $N_m (= N_{md})$ and the scaling fit ($N_{m\gamma}$) from the above mentioned scaling relations. This resolves the discrepancy noted in the analysis of the same data in ref. [12]. Here again, as seen from the inset, though the statistical deviation does not become minimum for the scaling with log correction term (presumably due to inclusion of some tail-end points), the best fit for γ assumes the desired value $1/2$.

III. SUMMARY AND CONCLUSION

We have studied here the scaling behaviors of Hirsch index h for inequalities [1] applied to the unequal distributions of responses and statistics in different physical (fracture and percolation) and social systems (citations and parliament sizes). We have shown that Hirsch index follow remarkable off-critical Widom-Stauffer scaling in these systems that are widely different in terms of their dimensionality and symmetry.

We have studied avalanche or cluster sizes in physical systems like the fiber bundle models (see e.g., [2][10]) and percolating systems (see e.g., [3, 14]). Indeed, surprising successes of such Hirsch-like social inequality measures were already seen (see e.g., [9, 15]) in predicting the global failures in fiber

² The data extracted from this paper can be found in <https://sciencehistory.epfl.ch/physics-and-sociology/>

bundles and the self-organized critical points of sand-pile systems. We show here, from the Monte Carlo simulation results (see sections IIA and IIB), that the Widom-Stauffer-like scaling relation (1a,b) of the Hirsch index h (defined here for the size s distributions n_s , through $h = s = n_s$) fit remarkably well with the system size N (see Figs. 2 and 4). We also show (see the insets of Figs. 2 and 4) that the scaling collapse breaks down without the $\log N$ term in (1) (visibly clear for the fiber bundle results in Fig. 2).

As is well known, the Hirsch index (h) was introduced originally to measure the inequalities in success (through citations in the subsequent literature) of the contributions (papers) of individual scientists. The citation function (see Fig. 1), is an well documented nonlinear and monotonically decaying function (cf. Zipf law [16]). h corresponds to the fixed point of this citation function. Analytical study [4], as well as numerical data analysis (see e.g., [5]), suggested $h \sim \sqrt{N_c} \sim \sqrt{N_p}$, for any author having N_p total papers and N_c total citations. Since for different scientists, we do not have any knowledge of the critical intervals (like $|\sigma_c - \sigma|$ or $|p_c - p|$ in relations (1)) here the detailed data for h are fitted to the relations (2) (see section IIC, Figs. 7, 8(a,b)) assuming the critical interval to be zero for the chosen authors. We show a better fit is $h \sim \sqrt{N_c}/\log N_c \sim \sqrt{N_p}/\log N_p$ (see relations (2a,b)). It may be noted that analysis of larger data set (see e.g., [17]) indicated significantly lower value ($\simeq 0.42$) of γ . As may be seen from our analysis (see Figs. 7 and 8), this is probably due to the choices of data corresponding to very large deviations from the respective critical points to fit power law scaling form and the missing $\log N$ term (in relations (2)) in the scaling form. It suggests, in absence of this information, extending the data sets for scientists who may not be that competitive may lead to wrong conclusions. In fact, while all the data for h -index of FBM avalanches and percolation clusters fit so well with the Widom-Stauffer scaling (2) with $\gamma = 1/2$, when the data for their critical points are fitted to $h \sim N^\gamma$ (see insets of Figs. 3 and 5), one gets best fits for γ around 0.40 and 0.48 respectively.

We also show that when the number N_m of the members of national assemblies or of parliaments for different countries of the world are identified effectively as their Hirsch index h , then N_m indeed would scale (see Fig. 9 in section IID) with the total population N as $\sqrt{N}/\log N$ for the respective countries. This observation should help comprehending the discrepancies with the proposed $N_m \sim \sqrt{N}$ relationship, reported in a very recent analysis [12].

For an additional check for the $\log N$ term in the scaling behaviour of the Hirsch index, we fitted the data for h in Fiber bundle models (section IIA, for stress values σ near the failure point σ_c), for percolation systems (section IIB, for site occupation concentrations p both above and below the percolation threshold p_c), paper citation data (section IIC, for authors with Kolkata index values k both above and below the critical value k_c) and parliament membership data (section IID) to the scaling forms $h = a\sqrt{N}$ and $h = \bar{a}\sqrt{N}/\log(N)$, where N denotes the appropriate system size (total number N of fibers in the FBM, total number N of lattice sites, total number of citations N_c of the author, or the total population N of the country). The estimated best fit values of the pre-factors a or \bar{a} are given in Table I. The range of error bars in these estimates of a and \bar{a} are such that, more than 95% of the data points fall within the indicated ranges. If we assume the value of the pre-factor to be the same across the systems considered (Young's asymptotic result [4] suggested the value of the pre-factor a for citation to be about 0.54), the observed fitting error ranges for other systems (see Table I) then clearly suggest the fit with $\log N$ term to be more appropriate. In fact, the average value of the pre-factor ($\bar{a} \simeq 3.0$ in Table I) fits reasonably across the systems considered and also agrees with the values of $f(0)$ in Figs. 2 (for FBM), 4 (for percolating systems) and 6 (for paper citations by an individual scientist). The comparative agreements of the values of a (with large percentile errors) and \bar{a} ($= f(0)$ in Figs. 2, 4 and 6 having smaller percentile errors) in table I clearly indicates that the fitting of h -index to the scaling relation (1) is much better. These errors due to the fittings with scaling relations (2) therefore become much more prominent for parliament member number data, where the critical intervals or the country's distance from the respective critical point are completely unknown.

TABLE I. Values of the pre-factors a and \bar{a} for h -index scaling fits.

	FBM avalanche sizes	percolation cluster sizes	citation sizes	parliament sizes
fit to $h = a\sqrt{N}$ $a =$	0.15 ± 0.10	0.15 ± 0.10	0.45 ± 0.15	0.10 ± 0.08
fit to $h = \bar{a}\sqrt{N}/\log(N)$ $\bar{a} =$	2.00 ± 1.00	2.50 ± 1.00	3.50 ± 0.50	2.00 ± 1.25

In conclusion, we have explored the best fit of the Hirsch index values h for system sizes N (or N_c) with a Widom-Stauffer like finite size scaling form Eq. (1). This is essentially based on the Monte Carlo simulation data collapse in the fiber bundle model and percolating systems (for σ or p away from σ_c or p_c respectively; see Figs. 2 and 4) and data analysis for citations of 100 scientists (k away from k_c ; see Fig. 6) analyzed in ref. [5]. Data for parliament member numbers (identified as the corresponding country's h index; see Fig 9) are fitted to the relation Eq. (2), as the equivalent critical interval ($[\sigma_c - \sigma]$, $[p_c - p]$ or $[k_c - k]$) are unknown. We find the scaling fit (to relations Eq. (1) and (2)) deteriorate considerably if the $\log N$ term is dropped. We give in the table I the estimated error in the pre-factors a and \bar{a} , and assuming the pre-factor to have the same value across the systems considered here, we again find the finite size scaling relation for the Hirsch index h with $\log N$ term ($h = \bar{a}\sqrt{N}/\log N$, $\bar{a} \equiv f(0)$ of the Widom-Stauffer relation Eq. (1) for finite system size N) to be more appropriate for the scaling behaviour of Hirsch index near the respective critical points.

ACKNOWLEDGMENTS

We are grateful to Parongama Sen for useful comments and suggestions. BKC is thankful to the Indian National Science Academy for their Senior Scientist Research Grant.

-
- [1] Jorge E Hirsch. An index to quantify an individual's scientific research output. *Proceedings of the National Academy of Sciences*, 102(46):16569–16572, 2005.
 - [2] Soumyajyoti Biswas, Purusattam Ray, and Bikas K Chakrabarti. *Statistical physics of fracture, breakdown, and earthquake: effects of disorder and heterogeneity*. John Wiley & Sons, 2015.
 - [3] Dietrich Stauffer and Ammon Aharony. *Introduction to percolation theory*. CRC press, 2018.
 - [4] Alexander Yong. Critique of Hirsch's citation index: A combinatorial Fermi problem. *Notices of the American Mathematical Society*, 61(9):1040–1050, 2014.
 - [5] Asim Ghosh and Bikas K Chakrabarti. Limiting value of the Kolkata Index for social inequality and a possible social constant. *Physica A: Statistical Mechanics and its Applications*, 573:125944, 2021.
 - [6] Suchismita Banerjee, Bikas K. Chakrabarti, Manipushpak Mitra and Suresh Mutuswami, Inequality Measures: The Kolkata Index in Comparison With Other Measures, *Frontiers in Physics* 8(2020) 562182. doi: 10.3389/fphy.2020.562182
 - [7] Robin IM Dunbar. Neocortex size as a constraint on group size in primates. *Journal of Human Evolution*, 22(6):469–493, 1992.
 - [8] Kunal Bhattacharya, Asim Ghosh, Daniel Monsivais, Robin IM Dunbar, and Kimmo Kaski. Sex differences in social focus across the life cycle in humans. *Royal Society Open Science*, 3(4):160097, 2016.
 - [9] Soumyajyoti Biswas and Bikas K Chakrabarti. Social inequality analysis of Fiber Bundle Model statistics and prediction of materials failure. *Physical Review E*, 104(4):044308, 2021.
 - [10] Srutarshi Pradhan, Alex Hansen, and Bikas K Chakrabarti. Failure processes in elastic Fiber Bundles. *Reviews of Modern Physics*, 82(1):499, 2010.
 - [11] Rein Taagepera. The size of national assemblies. *Social Science Research*, 1(4):385–401, 1972.
 - [12] Giorgio Margaritondo. Size of national assemblies: The classic derivation of the cube-root law is conceptually flawed. *Frontiers in Physics*, 8:606, 2021.
 - [13] Per C. Hemmer and Alex Hansen. The distribution of simultaneous fiber failures in Fiber Bundles. *Journal of Applied Mechanics*, 59:909, 1992.

- [14] Dietrich Stauffer. Scaling theory of percolation clusters. *Physics reports*, 54(1):1–74, 1979.
- [15] Subhrangshu Sekhar Manna, Soumyajyoti Biswas, and Bikas K Chakrabarti. Near universal values of social inequality indices in self-organized critical models. *Physica A: Statistical Mechanics and its Applications*, 596:127121, 2022.
- [16] George Kingsley Zipf. *Selected Studies of the Principle of Relative Frequency in Language*. Harvard University Press, Cambridge, MA, 1932.
- [17] Filippo Radicchi and Claudio Castellano. Analysis of bibliometric indicators for individual scholars in a large data set. *Scientometrics*, 97(3):627–637, 2013.