

A Deformation Quantization for Non-Flat Spacetimes and Applications to QFT

Albert Much*¹

¹Institut für Theoretische Physik, Universität Leipzig, D-04103 Leipzig

January 31, 2024

Abstract

We provide a deformation quantization, in the sense of Rieffel, for *all* globally hyperbolic spacetimes with a Poisson structure. The Poisson structures have to satisfy Fedosov type requirements in order for the deformed product to be associative. We apply the novel deformation to quantum field theories and their respective states and we prove that the deformed state (i.e. a state in non-commutative spacetime) has a singularity structure resembling Minkowski, i.e. is *Hadamard*, if the undeformed state is Hadamard. This proves that the Hadamard condition, and hence the quantum field theoretical implementation of the equivalence principle is a general concept that holds in spacetimes with quantum features (i.e. a non-commutative spacetime).

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Mathematical Preliminaries | 3 |
| 2.1 | Geometrical Setup | 3 |
| 2.2 | Embedding Formalism | 3 |
| 2.3 | Moving in the Embedded Spacetime | 5 |
| 2.4 | Poisson Structure | 10 |
| 3 | A Generalized Rieffel Deformation | 11 |
| 3.1 | Oscillatory Integrals | 11 |
| 3.2 | The Deformed Product | 12 |
| 3.3 | Deformed Product for Two Different Points | 18 |
| 4 | Quantum Field Theory in Globally Hyperbolic Spacetimes | 21 |
| 4.1 | The Quantized Scalar Field | 21 |
| 4.2 | States and GNS Theorem | 21 |
| 4.3 | The Microlocal Spectrum Condition and Hadamard | 22 |
| 5 | Deformation of QFT in Globally Hyperbolic Spacetimes | 24 |
| 5.1 | Representation of the Deformation | 24 |
| 5.2 | The Microlocal Spectrum Condition under Deformation Quantization | 25 |
| 6 | Discussion and Outlook | 26 |

*much@itp.uni-leipzig.de

1 Introduction

Quantum field theory (**QFT**) in curved spacetimes (**CST**) (see monographs [Ful89, Wal94, Few08, BD82, BGP08] and [KW91]) is regarded as a zero-order approximation of a complete theory of quantum gravity. The approximation pertains to the treatment of gravity (or spacetime) as a classical entity rather than a quantum one, as in general relativity.

Even in this approximation, the framework has already provided various insights and achievements, such as thermal radiations by black holes ([Haw75, Wal75, FH90, KW91]), the micro-local spectrum condition ([Rad92, Rad96, BFK96]), Pauli's spin-statistic theorem in curved spacetime ([Ver01]), quantum inequalities ([FP78, For91, Few00, FH05, FS08, SV08, Few12]), perturbation theory ([BF00, HW01, HW02, Rej16]), descriptions of the early universe and extreme astrophysical environments [Hac15] (and references therein).

Moreover, it has led to questioning the foundations of physics, such as the very notion (or existence) of particles (Fulling-Davies-Unruh effect, [Ful73, Dav75, Unr76, Dav84]).

The states generalizing the Minkowski vacuum in QFT in CST are the so-called Hadamard states. These states exhibit the Hadamard singularity structure and exist for a large class of quantum field theories in globally hyperbolic spacetimes, [FNW81, SV01], see also [GW13]. An essential reason to work with them is the lack of a vacuum (lack of an isometry group¹) in curved spacetimes. Motivated by the equivalence principle, one demands states resembling the same singularity structure as the Minkowski vacuum at short distances.

Due to the foundational importance of QFT in CST and the zero-order approximation argument, it is clear that any fundamental and complete theory of quantum gravity should at least contain QFT in CST in some limit. In addition, this theory should become important where the domain of validity of QFT in CST breaks down. This breakdown occurs at distances (or curvatures) near the Planck scales, [Wal94, Chapter 1].

It is exactly around those distances that the hypothesis of a quantum (or noncommutative) spacetime (geometry) comes into play as the solution to the geometrical measurement problem, see Refs.[Ahl94, DFR95]. The problem arises due to limitations of localization of spacetime points, resulting from combining the quantum-mechanical uncertainty principle with the formation of black holes in general relativity. An n -dimensional quantum spacetime is defined as an algebra \mathcal{V} , generated by elements \hat{x} , that obey

$$[\hat{x}^\mu, \hat{x}^\nu] = 2i\Theta^{\mu\nu}, \quad (1.1)$$

where Θ is a real, constant, skew symmetric $n \times n$ matrix. By representing the generating elements \hat{x} on a Hilbert space, one derives an uncertainty principle for the coordinates of spacetime. In [DFR95], the authors defined a quantum field theory in noncommutative Minkowski space, by defining the representation space of the QFT to be the tensor product of $\mathcal{V} \otimes \mathcal{H}$, where \mathcal{H} is the Hilbert space of the quantum field under consideration. Next, the authors in Ref. [GL07] constructed a unitary operator that maps the tensor product space $\mathcal{V} \otimes \mathcal{H}$ to the Hilbert space \mathcal{H} , which is a strict deformation of QFT called warped convolutions [BS08, BLS11].

This prescription is connected to the Rieffel product [Rie93] (also known as the Moyal-Weyl-Groenewold), which is a noncommutative associative product of functions provided by

$$f \star_\theta g = (2\pi)^{-n} \iint \alpha_{\Theta x}(f) \alpha_y(g) e^{ix \cdot y} d^n x d^n y, \quad (1.2)$$

where α denotes the action of the translation group \mathbb{R}^n , i.e. $\alpha_a(\phi)(x) = \phi(x + a)$. The oscillatory integral is well defined for smooth functions of compact support $f, g \in C_0^\infty(\mathbb{R}^n)$. To introduce perturbations stemming from the *non-commutative spacetime*, one replaces the pointwise product by the Rieffel product. This procedure is equal to defining the n -point functions for a QFT in non-commutative Minkowski space, as done in [Oec00, Oec01], using the fields in [GL07].

The Rieffel deformation, yet fruitful, is generally inapplicable to curved spacetimes due to the essential requirement of smooth actions of \mathbb{R}^n , i.e., mutually commutative generators. In rare cases, one can use commuting Lie vector fields [MM11] or a spacetime with commuting Killing vector fields [DLMM11].

Recently, this limitation was circumvented for the De Sitter spacetime by using the embedding formalism. The idea was to embed an n -dimensional curved spacetime manifold into a higher

¹In flat spacetime, this is the Poincaré group, which leaves the vacuum invariant.

dimensional Minkowski spacetime, thus supplying us with an n -dimensional group of translations with which we define the Rieffel product.

In what follows, we first use [MS11, Theorem 1.1], which provides an isometric embedding for all globally hyperbolic spacetimes into a higher dimensional Minkowski spacetime \mathbb{L}^N . Then, we use those embeddings together with recent techniques from optimization problems on smooth manifolds, see [AMS07, Bou23], to define the equivalent of translations (including the exponential map), namely retractions on those manifolds. The retractions form a group, and using this group structure, we define a generalized Rieffel product for curved manifolds. This product is parameterized by a deformation parameter, and it reduces to the well-established Rieffel product in the flat case. Furthermore, at first order in the deformation parameter, the generalized deformed product aligns with the Poisson bracket. We prove that under a certain assumption, namely a Fedosov type requirement on the Poisson structure π , i.e. $\nabla_U \pi = 0$, that the product is associative up to second order in the deformation parameter.

Since our main motivation to study these deformations lies in the application to QFT in curved spacetimes, and in particular to the two-point function, we generalize the deformed product to two points.

In the final section, we demonstrate that if a state satisfies the microlocal spectrum condition—implying, according to the Radzikowski theorem, that it is Hadamard (possessing the same singularity structure as in Minkowski spacetime), then the deformed state also satisfies this condition. Consequently, we conclude that the deformed state is Hadamard. Moreover, beyond the aforementioned results, we establish that the Hadamard property is independent of the chosen isometric embedding and/or retraction.

2 Mathematical Preliminaries

We start with basic definitions needed for the next results.

2.1 Geometrical Setup

DEFINITION 2.1 (Globally hyperbolic spacetime). *We denote a spacetime by (\mathcal{M}, g) where \mathcal{M} is a smooth, four-dimensional manifold and g is a Lorentzian metric on \mathcal{M} with signature $(-1, +1, +1, +1)$. Besides, we assume time-orientability of the manifold. This means that there exists a C^∞ -vector-field u on \mathcal{M} that is everywhere time-like, i.e. $g(u, u) < 0$. A smooth curve $\gamma : I \rightarrow \mathcal{M}$, I being a connected subset of \mathbb{R} , is called causal if $g(\dot{\gamma}, \dot{\gamma}) \leq 0$ where $\dot{\gamma}$ denotes the tangent vector of γ . A causal curve is called future directed if $g(\dot{\gamma}, u) < 0$ and past directed if $g(\dot{\gamma}, u) > 0$ all along γ and for a global time-like vector-field u . For any point $x \in \mathcal{M}$, $J^\pm(x)$ denotes the set of all points in \mathcal{M} which can be connected to x by a future(+)/past (-)-directed causal curve. A time-orientable spacetime is called globally hyperbolic if for each pair of points $x, y \in \mathcal{M}$ the set $J^-(x) \cap J^+(y)$ is compact whenever it is non-empty.*

The definition is equal to the existence of a smooth foliation of \mathcal{M} in Cauchy surfaces, where a smooth hypersurface of \mathcal{M} is called a *Cauchy surface* if it is intersected exactly once by each inextendible causal curve. A functional point of view of these spacetimes is the admittance of a choice of time.

2.2 Embedding Formalism

In Ref. [MS11] the authors proved the existence of an isometric embedding of any globally hyperbolic manifold in to a higher dimensional Minkowski space \mathbb{L}^N . First, we define a smooth embedded submanifold of a linear space \mathcal{E} of dimension N , [Bou23, Defintion 3.10].²

DEFINITION 2.2. (Embedded submanifold) *Let \mathcal{E} be a linear space of dimension N . A non-empty subset \mathcal{M} of \mathcal{E} is a (smooth) embedded submanifold of \mathcal{E} of dimension $d = N - k$ if for some $k \geq 1$ and, for each $X \in \mathcal{M}$, there exists a neighbourhood U of X in \mathcal{E} and a smooth function $F : U \rightarrow \mathbb{R}^k$ such that*

²For example the Minkowski space \mathbb{L}^N and the Euclidean space \mathbb{R}^n fall into this category.

1. if X is in U , then $F(X) = 0$ if and only if $X \in \mathcal{M}$; and
2. $\text{rank } DF(X) = k$.

Such a function F is called a **local defining function** for \mathcal{M} at X .

Example 2.3. For example, in the case of embedding the unit sphere \mathbb{S}^2 into \mathbb{R}^3 , the differentiable map $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ is given by the spherical equation

$$F(X) = X_1^2 + X_2^2 + X_3^2 - 1,$$

or for the N -dimensional de Sitter $d\mathbb{S}^N$ a possible map $F : \mathbb{R}^{N+1} \mapsto \mathbb{R}$ is specified by

$$F(X) = -X_0^2 + X_i^2 + X_N^2 - H^{-2},$$

with index $i = 1, \dots, N-1$ and where H is the Hubble parameter.

For more embeddings of spacetimes that are familiar in general relativity into a higher dimensional Minkowski spacetimes, see [Ros65].

The Nash embedding theorem states that every Riemannian manifold (compact or not) can be isometrically embedded in \mathbb{R}^N for some large enough N . The authors in Ref. [MS11] extended this result to Lorentzian manifolds that are stably causal with a steep temporal function, i.e. a smooth function τ such that $g(\nabla\tau, \nabla\tau) \leq -1$. In particular, one can prove that *any* globally hyperbolic spacetime (\mathcal{M}, g) admits a steep temporal function τ and thus a Cauchy orthogonal decomposition, i.e.

$$\mathcal{M} = \mathbb{R} \times \Sigma, \quad g = -\beta d\tau^2 + g_\tau, \quad (2.1)$$

where $\beta > 0$ is a function on \mathcal{M} , g_τ is a Riemannian metric on $\Sigma_\tau := \{\tau\} \times \Sigma$ smoothly varying with τ , with upper bounded function β . And it is exactly in those spacetimes in which quantum field theories are rigorously defined. The theorem that establishes the embedding is the following, [MS11, Theorem 1.1].

THEOREM 2.4. *Let (\mathcal{M}, g) be a Lorentzian manifold. The following assertions are equivalent:*

1. (\mathcal{M}, g) admits a isometric embedding in \mathbb{L}^N for some $N \in \mathbb{N}$.
2. (\mathcal{M}, g) is a stably causal spacetime with a steep temporal function, i.e., a smooth function τ such that $g(\nabla\tau, \nabla\tau) \leq -1$.

This theorem is used in the following to define movements resembling translations in non-flat manifolds. In particular, we choose a local coordinate system for the globally hyperbolic manifold (\mathcal{M}, g) and one for the embedding space. Hence, let the local coordinate basis expanded in terms of the metric g be specified by

$$g = g_{\mu\nu} dx^\mu dx^\nu,$$

where $\mu, \nu = 0, 1, 2, 3$. A smooth local isometric embedding in to a higher dimensional Minkowski space \mathbb{L}^N is given by a smooth map (the local defining function) $F : \mathcal{M} \rightarrow \mathbb{L}^N$, with local coordinates $X^A = (X^\mu, X^a)$ with $a = 4, \dots, N$, such that

$$\sum_{A=0}^N \frac{\partial X^A}{\partial x^\mu} \frac{\partial X_A}{\partial x^\nu} = g_{\mu\nu}. \quad (2.2)$$

The existence of an embedding is equivalent to existence of solutions for this system of coupled differential equations presented by the parametric representation, see [SKM⁺03, Chapter 37], as

$$X^A = X^A(x^\mu).$$

Note that we have $(N-4)$ constraints on the coordinates X^a , (see [Goe74]). Due to the full rank of the sub-matrix of the Jacobian DF , w.r.t. the constrained coordinates X^a , it follows by [Lee13,

Proposition B.25] and the implicit function theorem that *locally* the embedding coordinates X^a can be written as functions of the variables X^μ , i.e.

$$X^a = X^a(X^\mu). \quad (2.3)$$

Furthermore, due to the invertibility of the Jacobi-matrix $\partial X^\mu / \partial x^\nu$ (which follows from Equation (2.2)³) we can use the inverse function theorem to prove that the functions $X^\mu(x^\nu)$ can be inverted, i.e.

$$x^\mu = x^\mu(X^\nu). \quad (2.4)$$

DEFINITION 2.5. For a function f on a globally hyperbolic spacetime (\mathcal{M}, g) , we denote $f(X)$ as its extension to the embedding spacetime (\mathbb{R}^N, η) using the parametrization given in Equation (2.4).

DEFINITION 2.6. Let $\iota_{\mathcal{M},g}$ denote the isometric embedding of a globally hyperbolic manifold (\mathcal{M}, g) in \mathbb{L}^N .

Remark 2.7. While we consider the embedding of the entire manifold (\mathcal{M}, g) , the upcoming deformation can be as well applied to submanifolds of the spacetime (\mathcal{M}, g) . The most obvious is to embed the Riemannian manifold, denoted by Σ in Equation (2.1) using the Nash embedding theorem. Other examples include the case of product manifolds, $(\mathcal{M} = \mathcal{M}_1 \times \mathcal{M}_2, g = g_1 \oplus g_2)$ where (\mathcal{M}_1, g_1) is globally hyperbolic and (\mathcal{M}_2, g_2) is a geodesically complete Riemannian manifold, hence by [BEE96, Theorem 3.68] securing that (\mathcal{M}, g) is globally hyperbolic. In this case one could either embed the manifold (\mathcal{M}_1, g_1) by Theorem 2.4 or the Riemannian manifold (\mathcal{M}_2, g_2) .

2.3 Moving in the Embedded Spacetime

For the following considerations we need the concept of tangent spaces w.r.t points on the embedded manifolds. Thus, we first define the tangent space $T_x\mathcal{M}$ at the point $x \in \mathcal{M}$, [Bou23, Definition 3.14.].

DEFINITION 2.8 (Tangent space). Let \mathcal{M} be a subset of \mathcal{E} . For all $x \in \mathcal{M}$, define

$$T_x\mathcal{M} = \{c'(0) | c : I \rightarrow \mathcal{M} \text{ is smooth and } c(0) = x\}, \quad (2.5)$$

where I is any open interval containing $t = 0$. That is, v is in $T_x\mathcal{M}$ if and only if there exists a smooth curve on \mathcal{M} passing through x with velocity v .

To keep this presentation self-contained, we also introduce the concept of smoothness w.r.t. maps on the embedded submanifolds, [Bou23, Definition 3.30.].

DEFINITION 2.9 (Smooth maps). Let \mathcal{M} and \mathcal{M}' be embedded submanifolds of \mathcal{E} and \mathcal{E}' (respectively). A map $f : \mathcal{M} \rightarrow \mathcal{M}'$ is smooth at $x \in \mathcal{M}$ if there exists a function $\bar{f} : U \rightarrow \mathcal{E}'$ which is smooth on a neighborhood U of x in \mathcal{E} and such that f and \bar{f} coincide on $\mathcal{M} \cap U$. We call \bar{f} a (local) smooth extension of f around x . The map f is smooth if it is smooth at all $x \in \mathcal{M}$.

A more intuitive notion of smoothness goes as follows. A real-valued function $f : \mathcal{M} \rightarrow \mathbb{R}$ is smooth if and only if there exists a smooth extension $\bar{f} : U \rightarrow \mathbb{R}$, where U is a neighbourhood of \mathcal{M} in \mathcal{E} and the extension coincides with f on \mathcal{M} . Next, we define the differential of a function, [Bou23, Definition 3.34.].

DEFINITION 2.10 (Differential). The differential of $f : \mathcal{M} \rightarrow \mathcal{M}'$ at the point $x \in \mathcal{M}$ is the linear map $Df(x) : T_x\mathcal{M} \rightarrow T_{f(x)}\mathcal{M}'$ defined by

$$Df(x)[v] = \left. \frac{d}{dt} f(c(t)) \right|_{t=0} = (f \circ c)'(0) = Df(c(0))[c'(0)],$$

where c is a smooth curve passing through x at $t = 0$ with velocity v .

³This is seen by rewriting Equation (2.2) using the chain rule as $(\eta_{\mu\lambda} + \frac{\partial X^A}{\partial X^\mu} \frac{\partial X_A}{\partial X^\lambda}) \frac{\partial X^\mu}{\partial x^\nu} \frac{\partial X^\nu}{\partial x^\rho} = g_{\lambda\rho}$ and using twice the determinant product rule $\det(A \cdot B) = \det(A) \det(B)$.

DEFINITION 2.11 (Tangent space). We call $T_x\mathcal{M}$ the tangent space to \mathcal{M} at x , and the vectors in $T_x\mathcal{M}$ are called tangent vectors to \mathcal{M} at x . The dimension of $T_x\mathcal{M}$ coincides with the dimension of \mathcal{M} .

For the case of embedded manifolds, we have the following result, [Bou23, Theorem 3.15].

THEOREM 2.12. Let \mathcal{M} be an embedded submanifold of \mathcal{E} . Then, the tangent space $T_X\mathcal{M}$ is given as the kernel of the differential of any local defining function F at X ,

$$T_X\mathcal{M} = \ker DF(X). \quad (2.6)$$

We endow the embedded manifolds with a metric by using the fact that each tangent space $T_X\mathcal{M}$ is a linear subspace of (\mathbb{R}^N, η) or (\mathbb{R}^N, δ) . Hence, we can define the metric product (not necessarily positive definite)

$$(\cdot, \cdot)_X : T_X\mathcal{M} \times T_X\mathcal{M} \rightarrow \mathbb{R}$$

by the restriction of the metric product in \mathbb{L}^N or \mathbb{R}^N to $T_X\mathcal{M}$. Next, we define the vector field ∇f on a manifold, for the (Pseudo-)Riemannian case see [Bou23, Definition 3.58].

DEFINITION 2.13 (Gradient). Let $f : \mathcal{M} \rightarrow \mathbb{R}$ be a smooth function on (Pseudo-)Riemannian manifold \mathcal{M} . The (Pseudo-)Riemannian gradient of the function f is the vector field ∇f on \mathcal{M} defined by the following identities:

$$\forall (x, v) \in T\mathcal{M}, \quad Df(x)[v] = (v, \nabla f)_x,$$

where $Df(x)$ is as in Definition 2.10 and $(\cdot, \cdot)_x$ refers to the (Pseudo-)Riemannian metric restricted to $T_x\mathcal{M}$.

Next, we define the concept of the projector, see [Bou23, Definition 3.60]. The idea of a projector is to define gradients or tangent vectors, by natural definitions in the embedding manifold and projecting those quantities back to the embedded manifold. An example in the case of the $N-1$ dimensional (unit) sphere embedded into a one dimensional higher Euclidean space is $(\text{Proj}_X)_A^B = \delta_A^B - X_A X^B$, where $\text{Proj}_X : \mathbb{R}^N \rightarrow T_X\mathbb{S}^{N-1}$,

$$\begin{aligned} (X, \text{Proj}_X(V))_X &= X^A(\delta_A^B - X_A X^B)V_B \\ &= X^A V_A - \underbrace{X^2}_{=1} X^B V_B = 0. \end{aligned}$$

DEFINITION 2.14 (Orthogonal Projector). Let \mathcal{M} be an embedded submanifold of a (Pseudo-)Euclidean space \mathcal{E} equipped with a (Pseudo-)Euclidean metric (\cdot, \cdot) . The orthogonal projector to $T_X\mathcal{M}$ is the linear map $\text{Proj}_X : \mathcal{E} \rightarrow \mathcal{E}$ characterized by the following properties:

1. Range: $\text{im}(\text{Proj}_X) = T_X\mathcal{M}$;
2. Projector: $\text{Proj}_X \circ \text{Proj}_X = \text{Proj}_X$
3. Orthogonal $(u - \text{Proj}_X(u), v) = 0$ for all $v \in T_X\mathcal{M}$ and $u \in \mathcal{E}$.

LEMMA 2.15. The orthogonal projector, which is a smooth map [RS22, Lemma 3.1.2], can be represented by a matrix $P \in \mathbb{R}^{N \times N}$

$$\text{Proj}_Z(V^A) = P_C^A(Z) V^C.$$

which is uniquely determined by the conditions (see [Bou23, Proposition 3.63], [RS22, Chapter 3, Equation 3.1.2, Equation 3.1.3])

$$P(Z) = P(Z)^2 = P(Z)^T$$

for $Z \in M$ and $V \in T_Z M$

$$P(Z)V = V.$$

Using the orthogonal projector we have the following (extended) result, see [Bou23, Proposition 3.61., Equation (3.36)].

PROPOSITION 2.16. *Let (\mathcal{M}, g) be a (Pseudo-)Riemannian submanifold of $(\mathcal{E}, (\cdot, \cdot))$ and let $f : \mathcal{M} \rightarrow \mathbb{R}$ be a smooth function. The (Pseudo-)Riemannian gradient of f is defined by*

$$\nabla_A f(X) = \text{Proj}_X(\partial_A \bar{f}(X))$$

where \bar{f} is any smooth extension of f to a neighborhood of \mathcal{M} in \mathcal{E} and ∂ denotes the (Pseudo-)Euclidean gradient w.r.t. the coordinate X . Moreover, $\forall (X, V) \in T\mathcal{M}$

$$(V, \nabla f)_X = (V, \partial \bar{f}). \quad (2.7)$$

Thus, the scalar product with a tangent vector induces the following relation and useful definition

$$V^A \nabla_A f = V^B \partial_B \bar{f} =: V^B \frac{\partial f}{\partial X^B} = V^B \partial_B f. \quad (2.8)$$

In order to make these definitions more applicable, we give the following examples.

Example 2.17. *Consider the flat space $\mathbb{R}^{N-1} = \{X \in \mathbb{R}^N : X^N = 0\}$ (trivially) embedded into \mathbb{R}^N , where the embedding function is $F(X) = X^N$. The orthogonal projector in this case is the identity map. The differential of F is given by $DF(X)[V] = V^A \nabla_A F$. The tangent space $T_X \mathbb{R}^{N-1}$ is the set of all vectors V where $DF(X)[V] = 0$. Since the gradient of F vanishes if $A \neq N$ and is 1 if $A = N$ the kernel of $DF(X)$ consists of all $V \in \mathbb{R}^N$ such that $V^N = 0$, i.e.*

$$T_X \mathbb{R}^{N-1} = \ker DF(X) = \mathbb{R}^{N-1}.$$

This of course is a well-known fact that serves as the simplest example.

Example 2.18. *The unit sphere $\mathbb{S}^{N-1} = \{X \in \mathbb{R}^N : X_A X^A = 1, A = 1, \dots, N\}$ is the zero level set $M = F^{-1}(0)$ of $F(X) = X_A X^A - 1$, which is smooth from \mathbb{R}^N to \mathbb{R} . Since $DF(X)[V] = X_A V^A$, it is clear that $\text{rank } DF(X)[V] = 1$ for all $X \in \mathbb{S}^{N-1}$. The tangent spaces $T_X \mathbb{S}^{N-1}$ for $X \in \mathbb{S}^{N-1}$ are determined by the set of vectors V where $DF(X)[V] = 0$, i.e.*

$$T_X \mathbb{S}^{N-1} = \ker DF(X) = \{V \in \mathbb{R}^N : X_A V^A = 0\}.$$

Example 2.19. *The de Sitter space (where for the moment we set the Hubble parameter equal to one) $d\mathbb{S}^N = \{X \in \mathbb{R}^{N+1} : -X_0^2 + X_i X^i + X_N X^N = 1, i = 1, \dots, N-1\}$ is the zero level set of the $F(X) = -X_0^2 + X_i X^i + X_N X^N - 1$, which is smooth from \mathbb{R}^{N+1} to \mathbb{R} . The orthogonal projector in this case is*

$$(\text{Proj}_X)_A^B = \eta_A^B - X_A X^B$$

The rank of the differential of the local defining function is $\text{rank } DF(X)[V] = 1$ for all $X \in d\mathbb{S}^{N-1}$ and the tangent spaces $T_X \mathbb{S}^N$ for X are determined by the set of vectors V where $DF(X)[V] = 0$, i.e.

$$T_X d\mathbb{S}^N = \ker DF(X) = \{V \in \mathbb{R}^{N+1} : X_A V^A = -X_0 V^0 + X_i V^i + X_N V^N = 0\}.$$

DEFINITION 2.20. *The tangent bundle of a manifold \mathcal{M} is the disjoint union of the tangent spaces of \mathcal{M} :*

$$T\mathcal{M} = \{(x, v) : x \in \mathcal{M} \text{ and } v \in T_x \mathcal{M}\}$$

To consider terms involving second derivatives (Hessian) we introduce in the for submanifolds of \mathcal{E} the concept of a connection, see [Bou23, Equation 5.4] and [Bou23, Theorem 5.2]

DEFINITION 2.21. (Connection) *If a (Pseudo-)Riemannian manifold \mathcal{M} is a submanifold of the linear vector space \mathcal{E} , we define the connection as*

$$\nabla_U V = \text{Proj}_X(\bar{\nabla}_U \bar{V}(X)), \quad (2.9)$$

where V is a smooth vector field on \mathcal{M} , $U \in T_X \mathcal{M}$ is a tangent vector at the point X , \bar{V} is any smooth extension of V and $\bar{\nabla}$ is the connection on \mathcal{E} .

This connection is the (Pseudo-)Riemannian connection on \mathcal{M} , see [Bou23, Theorem 5.9]. Next, we define Hessian, see [Bou23, Definition 5.14].

DEFINITION 2.22 (Hessian). *Let $u \in T_x\mathcal{M}$ be a tangent vector and \mathcal{M} be (Pseudo-)Riemannian manifold and with its respective Riemannian connection ∇_u . The Riemannian Hessian of a smooth function f at $x \in \mathcal{M}$ is the linear map $\text{Hess}f(x) : T_x\mathcal{M} \rightarrow T_x\mathcal{M}$ defined as follows:*

$$\text{Hess}f(x)[u] = \nabla_u \nabla f(x). \quad (2.10)$$

The following result allows us to express the Hessian more explicitly, [Bou23, Corollary 5.16].

Corollary 2.23. *Let \mathcal{M} be a (Pseudo-)Riemannian submanifold of a (Pseudo-)Euclidean space. Consider a smooth function $f : \mathcal{M} \rightarrow \mathbb{R}$. Let \bar{G} be a smooth extension of ∇f — that is, \bar{G} is any smooth vector field defined on a neighborhood of \mathcal{M} in the embedding space such that $\bar{G}(X) = \nabla f(X)$ for all $X \in \mathcal{M}$. Then,*

$$\text{Hess}f(X)[U] = \text{Proj}_X(\bar{\nabla}_U \bar{G}(X)). \quad (2.11)$$

Using the former Definition and Corollary, we can write the Hessian as

$$\begin{aligned} \nabla_U \nabla_B f(X) &= \text{Proj}_X(U^A \partial_A \nabla_B f(X)) \\ &= P_B^C(U^A \partial_A \nabla_C f(X)). \end{aligned} \quad (2.12)$$

In order to define the Rieffel deformation we want to move away from a spacetime point $x \in \mathcal{M}$ along a tangent vector, while remaining on the manifold \mathcal{M} . For that reason, we introduce the concept of retraction at the point $x \in \mathcal{M}$, see [Bou23, Definition 3.47.] and [AMS07, Definition 4.1.1]

DEFINITION 2.24. *A retraction on a manifold \mathcal{M} is a smooth map R from the tangent bundle $T\mathcal{M}$ onto the manifold \mathcal{M} with the following properties*

$$R : T\mathcal{M} \rightarrow \mathcal{M} : (x, v) \mapsto R_x(v)$$

We can as well define retractions without the referral of a curve as follows, see [Bou23, Section 3.6].

DEFINITION 2.25. *Let $R_x : T_x\mathcal{M} \rightarrow \mathcal{M}$ denote the restriction of a smooth map $R : T\mathcal{M} \rightarrow \mathcal{M}$ to the tangent space at x . By the chain rule, each curve $c(t) = R_x(tv)$ satisfies $c(0) = R_x(0)$ and $c'(0) = DR_x(0)[v]$. Thus, R is a retraction exactly if, for all $(x, v) \in T\mathcal{M}$ we have*

1. $R_x(0) = x$, and
2. $DR_x(0) : T_x\mathcal{M} \rightarrow T_x\mathcal{M}$ is the identity map: $DR_x(0)[v] = v$.

If \mathcal{M} is an embedded manifold of a vector space \mathcal{E} , the retraction can be understood as follows. Since $T_x\mathcal{M}$ is a linear subspace of $T_x\mathcal{E}$ which is identified with \mathcal{E} , we can consider the sum $X + \xi_X$, where $X \in \mathcal{M}$ viewed as a point in \mathcal{E} and $\xi_X \in T_x\mathcal{E}$. The sum is well defined as an element in \mathcal{E} (with a slight abuse of notation). Projecting that element back to the embedded manifold \mathcal{M} gives the sum as an element in \mathcal{M} . This procedure defines a retraction. If the manifold \mathcal{M} is a Riemannian or pseudo-Riemannian manifold it admits a (second-order) retraction that is defined by the exponential mapping, see [Bou23, Definition 5.42. and Proposition 10.18]. Yet, this is computationally challenging, and hence we give in the following an algorithm given in [AMS07, Section 4.1.1.] to calculate a retraction that is more natural in the aspect of translations. This is done to keep close to the original definition of Rieffel, i.e. for the case where the manifold is Minkowski, we should get the definition of the deformed integral via translations.

Let \mathcal{M} be an embedded manifold of vector space \mathcal{E} and let \mathcal{N} be an abstract manifold such that $\dim(\mathcal{M}) + \dim(\mathcal{N}) = \dim(\mathcal{E})$. Assume that there is a diffeomorphism.

$$\phi : \mathcal{M} \times \mathcal{N} \rightarrow \mathcal{E}_* : (f, g) \mapsto \phi(f, g),$$

where \mathcal{E}_* is an open subset of \mathcal{E} , with a neutral element $I \in \mathcal{N}$ satisfying

$$\phi(f, I) = f, \quad \forall f \in \mathcal{M},$$

where I denotes the neutral element in \mathcal{N} . The following proposition gives a path to define a retraction, [AMS07, Proposition 4.1.2.]

PROPOSITION 2.26. *Under the above assumptions on ϕ , the mapping*

$$R_X(\xi) := \pi_1(\phi^{-1}(X + \xi)),$$

where $\pi_1 : \mathcal{M} \times \mathcal{N} \rightarrow \mathcal{M} : (f, g) \mapsto f$ is the projection onto the first component, defines a retraction on \mathcal{M} .

Example 2.27. *The retraction on a linear manifold \mathcal{E} is a translation*

$$R_X(\xi) = x + \xi,$$

where $x \in \mathcal{E}$ and $\xi \in T_x\mathcal{E}$.

Example 2.28. *Let $\mathcal{M} = \mathbb{S}^{n-1}$ be the $(n-1)$ -dimensional sphere, then a retraction is specified by*

$$R_X(\xi) = \frac{X + \xi}{\|X + \xi\|} = \frac{X + \xi}{\sqrt{1 + \|\xi\|^2}}$$

where $X \in \mathbb{S}^{n-1}$ and $\xi \in T_X\mathbb{S}^{n-1}$.

Example 2.29. *In the case of de Sitter spacetime (with Hubble parameter $H = 1$) $\mathcal{M} = d\mathbb{S}^{n-1}$ a retraction is provided by*

$$R_X(\xi) = \frac{X + \xi}{\sqrt{\eta_{AB}(X^A + \xi^A)(X^B + \xi^B)}} = \frac{X + \xi}{\sqrt{1 + \eta_{AB}\xi^A\xi^B}}$$

with $A, B = 0, 1, \dots, n$ and where $X \in d\mathbb{S}^{n-1}$ and $\xi \in T_X d\mathbb{S}^{n-1}$ as long as $\eta_{AB}\xi^A\xi^B \neq -1$.

DEFINITION 2.30. *A second-order retraction R on a (Pseudo-)Riemannian manifold \mathcal{M} is a retraction such that, for all $x \in \mathcal{M}$ and all $v \in T_x\mathcal{M}$, the curve $c(t) = R_x(tv)$ has zero acceleration at $t = 0$, that is, $c''(0) = 0$.*

Using the gradients, Hessians, projectors and the second-order retraction we have a Taylor expansion on curves given by, [Bou23, Proposition 5.44.]

$$f(R_x(v)) = f(x) + (v, \nabla f(x))_x + \frac{1}{2}(v, \nabla_v \nabla f(x))_x + \mathcal{O}(v^3). \quad (2.13)$$

DEFINITION 2.31. *Let (\mathcal{M}, g) be a globally hyperbolic spacetime, and let $T\mathcal{M}$ denote the tangent bundle of the manifold. Then, we denote by $\mathcal{G}_{\mathcal{M}}$ the set of transformations that are generated by second-order retractions $R : T\mathcal{M} \rightarrow \mathcal{M}$.*

DEFINITION 2.32. *The set $\mathcal{G}_{\mathcal{M}}$ forms a group by the natural identification of the group decomposition denoted by the symbol \circ*

$$R_{R_X(V)}(U) = R_X(V \circ U) \quad (2.14)$$

for $X \in \mathcal{M}$ and $V \in T_X\mathcal{M}, U \in T_{R_X(V)}\mathcal{M}$. The associativity of the composition rule is a direct consequence of the associativity inherent in the composition of maps and the identity element follows by a property of the retraction $R_X(0) = X$, see Definition 2.25.

Remark 2.33. *The definition of the group $\mathcal{G}_{\mathcal{M}}$ is sensible since one could build retractions from the exponential map, which under certain conditions can be considered as a map $\exp : \mathfrak{g} \rightarrow G$ from a Lie algebra \mathfrak{g} (thought of as the tangent space to the identity element of the group G) to a Lie group G . See, for example, [Mü01] for the proof of the group properties w.r.t. the exponential map.*

2.4 Poisson Structure

A Poisson manifold is defined by $(\mathcal{M}, \{\cdot, \cdot\})$, where \mathcal{M} is smooth manifold and $\{\cdot, \cdot\}$ is a Poisson bracket that can be equivalently represented by a Poisson bivector, see [Vai94, Chapter 1.1] or [Wal07, Proposition 4.1.20].

DEFINITION 2.34. *A Poisson bivector on a smooth manifold \mathcal{M} is a smooth bivector field $\pi \in \Gamma^\infty(\Lambda^2(T\mathcal{M}))$, i.e. π is a smooth skew-symmetric tensor, where Λ^2 denotes the exterior power of the tangent bundle and Γ^∞ denotes the space of smooth sections, satisfying the non-linear partial differential equation [Vai94, Equation 1.5] (Jacobi identity)*

$$\pi^{AB}\partial_A\pi^{CD} + \pi^{AC}\partial_A\pi^{DB} + \pi^{AD}\partial_A\pi^{BC} = 0.$$

Choosing local coordinates (U, X) , any Poisson bivector is given by

$$\pi|_U = \frac{1}{2} \sum_{A,B} \pi^{AB} \frac{\partial}{\partial X^A} \wedge \frac{\partial}{\partial X^B}.$$

The connection between the Poisson bracket and the Poisson bivector is provided by

$$\{f, g\} = \sum_{A,B} \pi^{AB} \partial_A f \partial_B g.$$

Remark 2.35. *With a slight abuse of notation we drop the bar symbol denoting the smooth extensions, where there is no risk of confusion.*

In the following result we make a connection between the product of a Poisson bivector and the orthogonal projector.

LEMMA 2.36. *For a smooth bivector field $\pi \in \Gamma^\infty(\Lambda^2(T_Z\mathcal{M}))$ and the orthogonal projector the following relations hold,*

$$\begin{aligned} \pi^{AB}(Z)P_B^C(Z) &= \pi^{AC}(Z), & \pi^{BD}(Z)P_B^C(Z) &= \pi^{CD}(Z), \\ \pi^{AD}(Z)\pi^{BC}(Z)\partial_C P_D^E(Z) &= \pi^{AC}(Z)\pi^{BD}(Z)\partial_C P_D^E(Z). \end{aligned}$$

Proof. We use the orthogonality condition given in Definition 2.14 and write an element $V \in T_Z\mathcal{M}$ as

$$V^A = \pi^{AC}W_C.$$

Then, the orthogonality condition for any element $U \in \mathbb{R}^N$ reads

$$(U_A - P_A^B U_B)\pi^{AC}W_C = 0$$

which reads

$$(\pi^{AC} - P_B^A \pi^{BC})U_A W_C = 0$$

since this holds for all $U \in \mathbb{R}^N$ and $W \in T_Z\mathcal{M}$, the proof of the first statement is concluded. By the skew-symmetry of the Poisson bivector and the first relation, the second statement is proved,

$$\pi^{BC}P_B^A = -\pi^{CB}P_B^A = -\pi^{CA} = \pi^{AC}.$$

For the second part, we use the first part of the lemma and the Jacobi identity, i.e. we write the equation as

$$\begin{aligned} \pi^{AD}\pi^{BC}\partial_C P_D^E &= \pi^{AC}\pi^{BD}\partial_C P_D^E \\ \pi^{BC}\partial_C(\pi^{AD}P_D^E) - \pi^{BC}P_D^E\partial_C\pi^{AD} &= \pi^{AC}\partial_C(\pi^{BD}P_D^E) - \pi^{AC}P_D^E\partial_C\pi^{BD} \\ \pi^{BC}\partial_C\pi^{AE} - \pi^{BC}P_D^E\partial_C\pi^{AD} &= \pi^{AC}\partial_C\pi^{BE} - \pi^{AC}P_D^E\partial_C\pi^{BD} \end{aligned}$$

Rearranging the terms we have,

$$\pi^{BC} \partial_C \pi^{AE} + \pi^{AC} \partial_C \pi^{EB} = P_D^E (\pi^{BC} \partial_C \pi^{AD} + \pi^{AC} \partial_C \pi^{DB}).$$

Next, we use the Jacobi identity to which the equation reduces to

$$-\pi^{EC} \partial_C \pi^{BA} = -P_D^E \pi^{DC} \partial_C \pi^{BA} = -\pi^{EC} \partial_C \pi^{BA}.$$

Reading the equalities from the bottom to the top we conclude the proof. \square

LEMMA 2.37. *The covariant derivative of the Poisson bivector field $\pi \in \Gamma^\infty(\Lambda^2(T_Z\mathcal{M}))$ is given by*

$$\nabla_U \pi^{DC} = U^A \partial_A \pi^{DC} - U^A \pi^{EC} \partial_A P_E^D - U^A \pi^{DE} \partial_A P_E^C.$$

Proof. Note that with the use of the orthogonal projection w.r.t. the embedding point Z , we can write the covariant derivative of a vector field $V \in T_Z\mathcal{M}$, using Definition 2.21, as

$$\begin{aligned} \nabla_U V^B &= U^A \nabla_A V^B = U^A P_C^B \partial_A \bar{V}^C \\ &= U^A \partial_A (P_C^B \bar{V}^C) - U^A V^C \partial_A P_C^B \\ &= U^A \partial_A \bar{V}^B - U^A V^C \partial_A P_C^B. \end{aligned}$$

Applying this using standard methods (see for example [Bou23, Section 4.7] to the two-tensor π gives us

$$\nabla_U \pi^{DC} = U^A \partial_A \pi^{DC} - U^A \pi^{EC} \partial_A P_E^D - U^A \pi^{DE} \partial_A P_E^C$$

and using the notation in Equation (2.8) we conclude the proof. \square

3 A Generalized Rieffel Deformation

Using the embedding formalism, we define the Rieffel product for a globally hyperbolic spacetime (\mathcal{M}, g) with a certain Poisson structure π , as a smooth action of the group $\mathcal{G}_\mathcal{M}$. See [Rie93] for the case $\mathcal{G}_{\mathbb{R}^4} = \mathbb{R}^4$ and [BW07, HNW07, DLS20] for constructions that use the exponential map to define a deformation quantization. The smooth action α of $\mathcal{G}_\mathcal{M}$ acts as a retraction in the embedding coordinates on functions extended to the embedding space.

3.1 Oscillatory Integrals

In order to define the Rieffel product a few definitions and results w.r.t. oscillatory integrals are briefly given. First, we define an oscillatory integral as in Ref. [Hör89, Ch. 7.8].

DEFINITION 3.1. *Let $X \subset \mathbb{R}^M$ be open and let Γ be an open cone on $X \times (\mathbb{R}^N \setminus \{0\})$ for some N , i.e., for all $(x, y) \in \Gamma$ also $(x, \lambda y) \in \Gamma$ for all $\lambda > 0$. Then an integral of the form (see Ref. [Hör89, Eq. 7.8.1])*

$$\int e^{-ix \cdot y} b(x, y) d^N y \quad (3.1)$$

*is called an **oscillatory integral**.*

Next, we define a symbol, see Ref. [Hör89, Def. 7.8.1].

DEFINITION 3.2. *Let $X \subset \mathbb{R}^M$ be open and let m, ρ, δ be real numbers with $0 < \rho \leq 1$ and $0 \leq \delta < 1$. Then we denote by $S_{\rho, \delta}^m(X \times \mathbb{R}^N)$ the set of all $b \in C^\infty(X \times \mathbb{R}^N)$ such that for every compact set $K \subset X$ and all α, β there exists some constant $C_{\alpha, \beta, K}$ such that the estimate*

$$|\partial_x^\beta \partial_y^\alpha b(x, y)| \leq C_{\alpha, \beta, K} (1 + |y|)^{m - \rho|\alpha| + \delta|\beta|} \quad (3.2)$$

is valid for all $(x, y) \in K \times \mathbb{R}^N$. The elements $S_{\rho, \delta}^m$ are called symbols of order m and type ρ, δ .

Remark 3.3. *It is proven in Refs. [Hör89, Thm. 7.8.2], (see also Refs. [LW16]), that if $b \in S_{\rho,\delta}^m$ and $m < -N$ the oscillatory integral (3.1) converges absolutely. In the case $m \geq -N$, the oscillatory integral has to be defined in a distributional manner. Using [Hör71, Proposition 1.1.11, Lemma 1.2.1, Proposition 1.2.2] one can prove the convergence of the oscillatory integral whenever $b(x, y)$ is a symbol even for the case $m \geq -N$ by using a mollifying function.*

DEFINITION 3.4. *We define the domain $\mathcal{D}_{\rho,\delta}^m(\mathcal{M})$*

$$\mathcal{D}_{\rho,\delta}^m(\mathcal{M}) := \{f, g \in C^\infty(\mathcal{M}) \mid \alpha_{\Theta X}(f)\alpha_Y(g) \in S_{\rho,\delta}^m, \text{ with } X, Y \in T\mathcal{M}\}. \quad (3.3)$$

as the collection of a pair of two smooth functions (f, g) that are symbols w.r.t. the retractions. Analogously, we define the domain $\mathcal{D}_{0,\rho,\delta}^m(\mathcal{M})$

$$\mathcal{D}_{0,\rho,\delta}^m(\mathcal{M}) := \{f, g \in C_0^\infty(\mathcal{M}) \mid \alpha_{\Theta X}(f)\alpha_Y(g) \in S_{\rho,\delta}^m, \text{ with } X, Y \in T\mathcal{M}\}. \quad (3.4)$$

as the collection of a pair of two smooth functions with compact support (f, g) that are symbols w.r.t. the retractions.

3.2 The Deformed Product

DEFINITION 3.5. *Let the smooth action α of the group $\mathcal{G}_{\mathcal{M}}$ denote second-order retractions w.r.t. the manifold $(\mathcal{M}, g) \subset (\mathbb{R}^N, \eta)$ (or (\mathbb{R}^N, δ)) and let θ be a real, constant deformation parameter and $\pi \in \Gamma^\infty(\Lambda^2(T_Z\mathcal{M}))$ a Poisson bivector that define the matrix $\Theta := \theta\pi$. Then, the (formal) generalized Rieffel product of two functions $(f, g) \in \mathcal{D}_{\rho,\delta}^m(\mathcal{M})$ is defined as*

$$\begin{aligned} (f \star_\theta g)(z) &\equiv \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \alpha_{\Theta X}(f(Z)) \alpha_Y(g(Z)) e^{-i(X,Y)_z} d^N X d^N Y \\ &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_Z(\Theta X)) g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y, \end{aligned}$$

where Z is the embedding point corresponding to z , $X, Y \in T_Z\mathcal{M}$ and the integrations are w.r.t. the non-vanishing components (i.e. maximally N) and the scalar product \cdot is w.r.t. the flat Minkowski (or Euclidean) metric η_{AB} (δ_{AB}) at the point Z . Moreover, the cut-off function $\chi \in C_0^\infty(\mathbb{R}^N \times \mathbb{R}^N)$ is chosen such that condition $\chi(0, 0) = 1$ is fulfilled.

By identifying $\alpha_{\Theta X}(f(Z))\alpha_Y(g(Z))$ with the function $b(X, Y)$ in Definition 3.1, it is clear that the Rieffel product is given as an oscillatory integral.

The generalized Rieffel product satisfies the following properties, where the first two are standard for a star product, see [Wal07, Defintion 6.1.1] and [Kon03].

PROPOSITION 3.6. *For functions $(f, g) \in \mathcal{D}_{\rho,\delta}^m(\mathcal{M})$ the generalized Rieffel product given in Definition 3.5 is well-defined and satisfies the following properties*

- *Unital,*

$$1 \star_\theta f = f \star_\theta 1 = f$$

- *The commutative limit,*

$$\lim_{\theta \rightarrow 0} (f \star_\theta g)(z) = (f \cdot g)(z),$$

- *The flat limit, i.e. in case that the manifold is the four-dimensional flat Minkowski (or Euclidean) space, with retractions provided by the translation group and constant Poisson bivector the generalized Rieffel product turns to the standard Rieffel product.*

Proof. To see that the Rieffel product is well-defined, for functions $(f, g) \in \mathcal{D}_{\rho,\delta}^m(\mathcal{M})$, we refer the reader to results mentioned in Remark 3.3. Unitality is seen by a straight-forward calculation,

$$(f \star_\theta 1)(z) \equiv \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \alpha_{\Theta X}(f(Z)) \alpha_Y(1) e^{-i(X,Y)_z} d^N X d^N Y$$

$$\begin{aligned}
&= \lim_{\epsilon \rightarrow 0} \iint \chi_1(\epsilon X) f(R_Z(\Theta X)) \delta^N(X) d^N X \\
&= f(Z),
\end{aligned}$$

where we used the property $R_Z(0) = Z$. The proof for $(1 \star_\theta f)(z)$ is analogous.

The commutative limit $\lim_{\theta \rightarrow 0}$ is proven as follows. We distinguish between two cases. The first case where the (translated) functions $\alpha_{\Theta X}(f)\alpha_Y(g)$ are symbols with $m < -4$ is trivial since for such symbols no mollifier is needed. In this case we interchange the limit $\lim_{\theta \rightarrow 0}$ with the integral by using the dominated Lebesgue convergence theorem, since we know that the function inside the integral is bounded (by the very definition of a symbol) by a measurable function. The explicit term is then given by,

$$\begin{aligned}
\lim_{\theta \rightarrow 0} (f \star_\theta g)(z) &= \lim_{\theta \rightarrow 0} \iint \alpha_{\Theta X}(f(Z)) \alpha_Y(g(Z)) e^{-iX \cdot Y} d^N X d^N Y \\
&= f(Z) \iint \alpha_Y(g(Z)) \delta(Y) d^N Y \\
&= f(z) \cdot g(z)
\end{aligned}$$

where we used the continuity of f and the property $R_Z(0) = Z$ of the retraction. In the case where the function $\alpha_{\Theta X}(f)\alpha_Y(g)$ is a symbol with $m \geq -4$ the deformed product is defined in a distributional manner, which we denote by

$$T(\varphi_\theta) := \lim_{\epsilon \rightarrow 0} \iint e^{-iX \cdot Y} \varphi_\theta^\epsilon(X, Y) d^N X d^N Y.$$

By continuity of the distribution, it is clear that $T(\varphi_\theta)$ converges to $T(\varphi_0)$ in the limit $\theta \rightarrow 0$, since $\varphi_\theta \rightarrow \varphi$ in $C_0^\infty(\mathbb{R}^4 \times \mathbb{R}^4)$. The commutative limit is as well given in [Rie93, Corollary 2.8] for functions belonging to certain Fréchet spaces.

We can (trivially) embed $(\mathcal{M}, g) = (\mathbb{R}^4, \eta)$ into a one dimension higher flat space (\mathbb{R}^5, η) by the embedding function $F = Z^5$, where the tangent space is $T_Z \mathbb{R}^4 = \mathbb{R}^4$, see Example 2.17. Since the 5-th component vanishes for both tangent vectors, we only integrate over four components. Taking the retractions to be translations we have the deformed integral

$$\begin{aligned}
(f \star_\theta g)(z) &\equiv \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \alpha_{\Theta X}(f(Z)) \alpha_Y(g(Z)) e^{-iX \cdot Y} d^4 X d^4 Y \\
&= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon x, \epsilon y) f(z + \Theta x) g(z + y) e^{-ix \cdot y} d^4 x d^4 y
\end{aligned}$$

□

Further properties that are standard for a star product, namely associativity and Poisson-compatibility, are satisfied and are subject of the following proposition.

PROPOSITION 3.7. *The generalized Rieffel product is Poisson compatible,*

$$\frac{i}{2}(f \star_\theta g - g \star_\theta f) = \theta \{f, g\}_\pi$$

and hence associative up to first order in θ , i.e.,

$$((f \star_\theta g) \star_\theta h)(z) = (f \star_\theta (g \star_\theta h))(z) + \mathcal{O}(\theta^2). \quad (3.5)$$

Proof. We prove the proposition by using the Taylor expansion of the retractions in order of the deformation parameter θ , i.e.

$$f(R_Z(\Theta X)) = f(Z) + (\Theta X)^A \nabla_A f(Z) + \mathcal{O}(\theta^2),$$

plugging this into the deformation formula we have up to first order in θ ,

$$(f \star_\theta g)(z) = \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_Z(\Theta X)) g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y,$$

$$\begin{aligned}
&= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) (f(Z) + (\Theta X)^A \nabla_A f(Z)) g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y, \\
&= f(z)g(z) + \Theta^{AB} \nabla_A f(Z) \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) X_B g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y, \\
&= f(z)g(z) - i\Theta^{AB} \nabla_A f \nabla_B g
\end{aligned}$$

where in the last lines we used the form of the retractions $g(R_Z(Y)) = g(Z) + Y^A \nabla_A g(Z) + \mathcal{O}(Y^2)$. To prove associativity we first consider

$$\begin{aligned}
((f \star_\theta g) \star_\theta h)(z) &= (F \star_\theta h)(z) = F(z)h(z) - i\Theta^{AB} \nabla_A F \nabla_B h \\
&= (f(z)g(z) - i\Theta^{AB} \nabla_A f \nabla_B g)h(z) - i\Theta^{AB} (\nabla_A f(z)g(z) + f(z)\nabla_A g(z)) \nabla_B h
\end{aligned}$$

Next we consider,

$$\begin{aligned}
(f \star_\theta (g \star_\theta h))(z) &= (f \star_\theta H)(z) = f(z)H(z) - i\Theta^{AB} \nabla_A f \nabla_B H \\
&= f(z)(g(z)h(z) - i\Theta^{AB} \nabla_A g \nabla_B h) - i\Theta^{AB} \nabla_A f (\nabla_B g(z)h(z) + g(z)\nabla_B h(z))
\end{aligned}$$

Using this expression we have for the commutator

$$\begin{aligned}
\frac{i}{2}(f \star_\theta g - g \star_\theta f) &= \Theta^{AB} \nabla_A f \nabla_B g \\
&= \theta \pi^{AB} \nabla_A f \nabla_B g \\
&= \theta \pi^{AB} \partial_A f \partial_B g,
\end{aligned}$$

where in the last line we used Lemma 2.36. \square

Next, we turn to the issue of associativity of the generalized product.

THEOREM 3.8. *Let the Poisson bivector $\pi \in \Gamma^\infty(\Lambda^2(T_Z \mathcal{M}))$, have a vanishing covariant derivative,*

$$\nabla_U \pi^{CD}(Z) = 0$$

for all $U \in T_Z \mathcal{M}$ and $Z \in \mathcal{M}$. Then, the generalized Rieffel product is associative up to second order in θ , and it is explicitly given by

$$(f \star_\theta g)(z) = fg - i\Theta^{AB} \partial_A f \partial_B g - \frac{1}{2}\Theta^{AC}\Theta^{BD} \partial_A \nabla_B f \partial_C \nabla_D g,$$

for functions $(f, g) \in \mathcal{D}_{\rho, \delta}^m(\mathcal{M})$.

Proof. We prove the proposition by using the Taylor expansion of the retractions in order of the deformation parameter θ ,

$$f(R_Z(\Theta X)) = f(Z) + (\Theta X)^A \nabla_A f(Z) + \frac{1}{2}(\Theta X)^A \nabla_{\Theta X} \nabla_A f(Z) + \mathcal{O}(\theta^3)$$

We use the following notation

$$\nabla_A f(Z) = \text{Proj}_Z(\partial_A \bar{f}), \quad \nabla_{\Theta X} \nabla_A f(Z) = \text{Proj}_Z((\Theta X)^B \partial_B \nabla_A f) \quad (3.6)$$

plugging this into the deformation formula we have up to second order in θ ,

$$\begin{aligned}
(f \star_\theta g)(z) &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_Z(\Theta X)) g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y, \\
&= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \left(f(Z) + (\Theta X)^A \nabla_A f(Z) + \frac{1}{2}(\Theta X)^A \text{Proj}_Z((\Theta X)^B \partial_B \nabla_A f) \right) \\
&\quad \times g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y,
\end{aligned}$$

$$\begin{aligned}
&= f(z)g(z) + \Theta^{AB} \nabla_A f(Z) \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) X_B g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y \\
&+ \frac{1}{2} \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) ((\Theta X)^A \text{Proj}_Z((\Theta X)^B \partial_B \nabla_A f(Z))) g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y,
\end{aligned}$$

We write the orthogonal projector as a matrix (see Lemma 2.15) and use the orthogonality of this projection rendering the former integral into

$$\begin{aligned}
(f \star_\theta g)(z) &= f(z)g(z) - i\Theta^{AB} \nabla_A f(Z) \nabla_B g(Z) \\
&+ \frac{1}{2} \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) ((\Theta X)^A \text{Proj}_Z((\Theta X)^B \partial_B \nabla_A f(Z))) g(R_Z(Y)) e^{-iX \cdot Y} d^N X d^N Y \\
&= f(z)g(z) - i\Theta^{AB} \nabla_A f(Z) \nabla_B g(Z) \\
&+ \frac{1}{4} \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) ((\Theta X)^A (\Theta X)^B \partial_B \nabla_A f(Z)) Y^D Y^F \partial_F \nabla_D g(Z) e^{-iX \cdot Y} d^N X d^N Y \\
&= f(z)g(z) - i\Theta^{AB} \nabla_A f(Z) \nabla_B g(Z) \\
&+ \frac{1}{4} \partial_B \nabla_A f(Z) \partial_F \nabla_D g(Z) \\
&\times \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) ((\Theta X)^A (\Theta X)^B Y^D Y^F e^{-iX \cdot Y} d^N X d^N Y \\
&= f(z)g(z) - i\Theta^{AB} \nabla_A f(Z) \nabla_B g(Z) \\
&- \frac{1}{4} \partial_B \nabla_A f(Z) \partial_C \nabla_D g(Z) (\Theta^{AC} \Theta^{BD} + \Theta^{AD} \Theta^{BC}) \\
&= f(z)g(z) - i\Theta^{AB} \nabla_A f(Z) \nabla_B g(Z) - \frac{1}{4} (\Theta^{AC} \Theta^{BD} + \Theta^{AD} \Theta^{BC}) \partial_A \nabla_B f(Z) \partial_C \nabla_D g(Z).
\end{aligned}$$

Using Lemma 2.36 and the following relation

$$\partial_C \nabla_D g(Z) = \partial_C \partial_D \bar{g}(Z) + (\partial_C P_D^E) \partial_E \bar{g}(Z)$$

we see that

$$\Theta^{AD} \Theta^{BC} \partial_C \nabla_D g(Z) = \Theta^{AC} \Theta^{BD} \partial_C \nabla_D g(Z).$$

Therefore, the deformed product of two functions is given up to second order in the deformation parameter as,

$$\begin{aligned}
(f \star_\theta g)(z) &= fg - i\Theta^{AB} \partial_A f \partial_B g - \frac{1}{4} (\Theta^{AC} \Theta^{BD} + \Theta^{AD} \Theta^{BC}) \partial_A \nabla_B f \partial_C \nabla_D g \\
&= fg - i\Theta^{AB} \partial_A f \partial_B g - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f \partial_C \nabla_D g.
\end{aligned}$$

Next, we prove associativity where we first consider

$$\begin{aligned}
((f \star_\theta g) \star_\theta h) &= (F \star_\theta h) \\
&= Fh - i\Theta^{AB} \partial_A F \partial_B h - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B F \partial_C \nabla_D h \\
&= (fg - i\Theta^{AB} \partial_A f \partial_B g - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f \partial_C \nabla_D g)h \\
&- i\Theta^{AB} \partial_A (fg - i\Theta^{CD} \partial_C f \partial_D g) \partial_B h \\
&- \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B (fg) \partial_C \nabla_D h
\end{aligned}$$

$$\begin{aligned}
&= (fg - i\Theta^{AB} \partial_A f \partial_B g - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f \partial_C \nabla_D g) h \\
&\quad - i\Theta^{AB} (\partial_A f g + f \partial_A g - i\partial_A \Theta^{CD} \partial_C f \partial_D g - i\Theta^{CD} \partial_A \partial_C f \partial_D g - i\Theta^{CD} \partial_C f \partial_A \partial_D g) \partial_B h \\
&\quad - \frac{1}{2} \Theta^{AC} \Theta^{BD} (\partial_A \nabla_B f g + \partial_A f \partial_B g + \partial_B f \partial_A g + f \partial_A \nabla_B g) \partial_C \nabla_D h
\end{aligned}$$

Next, we consider

$$\begin{aligned}
(f \star (g \star_\theta h)) &= (f \star_\theta G) \\
&= fG - i\Theta^{AB} \partial_A f \partial_B G - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f \partial_C \nabla_D G \\
&= f(gh - i\Theta^{AB} \partial_A g \partial_B h - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B g \partial_C \nabla_D h) \\
&\quad - i\Theta^{AB} \partial_A f \partial_B (gh - i\Theta^{CD} \partial_C g \partial_D h) \\
&\quad - \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f \partial_C \nabla_D (gh) \\
&= f(gh - i\Theta^{AB} \partial_A g \partial_B h - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B g \partial_C \nabla_D h) \\
&\quad - i\Theta^{AB} \partial_A f (\partial_B g h + g \partial_B h - i\partial_B \Theta^{CD} \partial_C g \partial_D h - i\Theta^{CD} \partial_B \partial_C g \partial_D h - i\Theta^{CD} \partial_C g \partial_B \partial_D h) \\
&\quad - \frac{1}{2} \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f (\partial_C \nabla_D g h + \partial_C g \partial_D h + \partial_D g \partial_C h + g \partial_C \nabla_D h)
\end{aligned}$$

This reduces to

$$\begin{aligned}
&(-\Theta^{AD} \partial_A \Theta^{CB} \partial_C f \partial_B g \partial_D h - \Theta^{AB} \Theta^{CD} \partial_A \partial_C f \partial_D g \partial_B h) - \Theta^{AC} \Theta^{BD} \partial_A f \partial_B g \partial_C \nabla_D h \\
&= (-\Theta^{CA} \partial_A \Theta^{BD} \partial_C f \partial_B g \partial_D h - \Theta^{AB} \Theta^{CD} \partial_A f \partial_C g \partial_B \partial_D h) - \Theta^{AC} \Theta^{BD} \partial_A \nabla_B f (\partial_C g \partial_D h)
\end{aligned}$$

which further reduces to

$$\Theta^{AB} \partial_A \Theta^{DC} - \Theta^{CA} \Theta^{BE} \partial_A P_E^D + \Theta^{BE} \Theta^{DA} \partial_A P_E^C = 0,$$

or using Lemma 2.36 can be written as

$$\Theta^{AB} \partial_A \Theta^{DC} - \Theta^{CE} \Theta^{BA} \partial_A P_E^D + \Theta^{BA} \Theta^{DE} \partial_A P_E^C = 0.$$

Using Lemma 2.37, this equality is equivalent to

$$\Theta^{AB} \nabla_A \Theta^{DC} = 0,$$

hence concluding the proof of associativity to second order. \square

A few remarks regarding the vanishing covariant derivative of the Poisson tensor, in order to assure associativity, are in order. A triple (\mathcal{M}, π, g) with the additional condition $\nabla \pi = 0$, is called a (Pseudo-)Riemannian Poisson manifold, see [Bou03, Definition 1.1.] and references therein. For a definition of Poisson-Riemannian manifold, where the covariant constant condition on the Poisson tensor is called Poisson compatibility, see [BM17, Equation 3.1].

In case, where the Poisson tensor has maximal rank $2n$ equal to the dimension of the manifold, the inverse of π^{AB} is given by the matrix ω_{AB} , a symplectic structure, which defines a symplectic manifold. In this case, Fedosov, [Fed94] has proven the existence of a deformation quantization if there is a symplectic connection, which is a torsion-free connection that preserves the symplectic tensor, defined by condition, [Fed94, Definition 2.3].

$$\nabla_C \omega_{AB} = 0.$$

A collection of a symplectic manifold and a symplectic connection is referred to as a Fedosov manifold, see [GRS97]. We proved associativity of the generalized deformed product, up to second order in the deformation parameter, if the covariant derivative of the Poisson structure vanishes. In the symplectic case, the vanishing Poisson structure induces a symplectic connection, which can be seen from the following identity,

$$\nabla(\pi\omega) = 0, \quad (3.7)$$

the vanishing covariant derivative of the Poisson tensor and the Leibniz rule.

Example 3.9. Let for the two-sphere S^2 the Poisson bivector at the embedding point Z and the orthogonal projection be assigned by,

$$\pi^{AB}(Z) = \varepsilon^{ABC} Z_C, \quad P_A^C(Z) = \delta_A^C - Z_A Z^C. \quad (3.8)$$

Then, $\nabla_U \pi^{AB} = 0$. This is explicitly proven in the following. The terms in Lemma 2.37 read

$$\begin{aligned} & -\frac{1}{2}(\Theta^{CA}\Theta^{BE} + \Theta^{CE}\Theta^{BA})\partial_A P_E^D = \frac{1}{2}(\Theta^{CA}\Theta^{BE} + \Theta^{CE}\Theta^{BA})\delta_{AE} Z^D \\ & = \frac{\theta^2}{2}(\varepsilon^{CAR}\varepsilon^{BES} + \varepsilon^{CES}\varepsilon^{BAR}) Z_R Z_S \delta_{AE} Z^D \\ & = (\varepsilon^{CAR}\varepsilon^{BES}) Z_R Z_S \delta_{AE} Z^D \\ & = \theta^2 (\delta^{CB}\delta^{RS} - \delta^{CS}\delta^{RB}) Z_R Z_S Z^D \\ & = -2\theta^2 (\delta^{CB} \underbrace{Z^2}_{=1} - Z^B Z^C) Z^D \\ & = \theta^2 (\delta^{CB} - Z^B Z^C) Z^D \end{aligned}$$

and

$$\begin{aligned} & +\frac{1}{2}(\Theta^{DA}\Theta^{BE} + \Theta^{DE}\Theta^{BA})\partial_E P_A^C = -\frac{1}{2}(\Theta^{DA}\Theta^{BE} + \Theta^{DE}\Theta^{BA})\delta_{AE} Z^C \\ & = -\frac{\theta^2}{2}(\varepsilon^{DAR}\varepsilon^{BES} + \varepsilon^{DES}\varepsilon^{BAR}) Z_R Z_S \delta_{AE} Z^C \\ & = -\theta^2 (\varepsilon^{DAR}\varepsilon^{BES}) Z_R Z_S \delta_{AE} Z^C \\ & = -\theta^2 (\delta^{DB}\delta^{RS} - \delta^{DS}\delta^{RB}) Z_R Z_S Z^C \\ & = -\theta^2 (\delta^{DB}\delta^{RS} - \delta^{DS}\delta^{RB}) Z_R Z_S Z^C \\ & = -\theta^2 (\delta^{DB} \underbrace{Z^2}_{=1} - Z^B Z^D) Z^C \\ & = -\theta^2 (\delta^{DB} - Z^B Z^D) Z^C \end{aligned}$$

and

$$\begin{aligned} \Theta^{AB}\partial_A\Theta^{DC} & = \theta^2 \varepsilon^{ABF}\varepsilon^{DCE} Z_F \delta_{AE} \\ & = \theta^2 (\delta^{BD}\delta^{FC} - \delta^{BC}\delta^{DF}) Z_F \\ & = \theta^2 (\delta^{BD} Z^C - \delta^{BC} Z^D). \end{aligned}$$

The sum of all those terms is zero.

PROPOSITION 3.10. *The commutation relations between the embedding coordinates are, up to second order in the deformation parameter θ , explicitly presented by*

$$[X^A, X^B]_\theta := X^A \star_\theta X^B - X^B \star_\theta X^A = -2i \Theta^{AB}$$

Proof. Using Theorem 3.8 we have for the deformed product of the coordinates X

$$\begin{aligned} X^A \star_\theta X^B &= X^A X^B - i \Theta^{CD} \partial_C X^A \partial_D X^B - \frac{1}{2} \Theta^{EC} \Theta^{FD} \partial_E \nabla_F X^A \partial_C \nabla_D X^B \\ &= X^A X^B - i \Theta^{CD} \delta_C^A \delta_D^B - \frac{1}{2} \Theta^{EC} \Theta^{FD} (\partial_E P_F^R \delta_R^A) \partial_C P_D^S \delta_S^B \\ &= X^A X^B - i \Theta^{AB} - \frac{1}{2} \Theta^{EC} \Theta^{FD} (\partial_E P_F^A) \partial_C P_D^B \end{aligned}$$

Next, we consider the deformed commutator

$$[X^A, X^B]_\theta = -2i \Theta^{AB} - \frac{1}{2} \Theta^{EC} \Theta^{FD} ((\partial_E P_F^A) \partial_C P_D^B - (\partial_E P_F^B) \partial_C P_D^A),$$

by interchanging the indices E and C and as well F and D , the second order term cancels. \square

Remark 3.11. *Following the definition of the generalized deformed product and taking into account the well-know fact that $\{X^A, X^B\} = \pi^{AB}$, the noncommutativity of the coordinates, given by*

$$[X^A, X^B]_\theta = -2i \theta \pi^{AB}$$

emerges in a canonical fashion. For example in the flat case, the canonical Poisson bivector is presented by the canonical, constant, skew-symmetric tensor and the noncommutativity has the Moyal-Weyl structure,

$$[x^\mu, x^\nu] = -2i \theta \pi^{\mu\nu},$$

where the embedding is determined by the trivial one. In the case of the two-sphere the canonical induced Poisson structure gives us the well-studied fuzzy sphere, i.e.

$$[X^A, X^B] = -2i \theta \varepsilon^{AB}_C X^C.$$

3.3 Deformed Product for Two Different Points

In the previous section we defined a generalization of the Rieffel product to the case of non-flat manifolds and established their associativity under a certain assumption on the Poisson bivector. Next, we broaden this framework to encompass the deformed product of functions defined at two distinct points. The main motivation for this extension is the application of those methods to quantum field theory and in particular to the two-point function.

DEFINITION 3.12. *Let the smooth action α of the group \mathcal{G}_M denote second-order retractions w.r.t. the manifold $(\mathcal{M}, g) \subset (\mathbb{R}^N, \eta)$ (or (\mathbb{R}^N, δ)) and let θ be a real constant deformation parameter and $\pi \in \Gamma^\infty(\Lambda^2(T_{Z_1}\mathcal{M}))$ a Poisson bivector that define the matrix $\Theta := \theta \pi$. Then, the (formal) generalized Rieffel product of two functions $(f, g) \in \mathcal{D}_{\rho, \delta}^m(\mathcal{M})$ at two different points is defined as*

$$\begin{aligned} f(z_1) \star_\theta g(z_2) &\equiv \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \alpha_{\Theta \delta(Z_1, Z_2)X}(f(Z_1)) \alpha_Y(g(Z_2)) e^{-i(X, Y)_{z_2}} d^N X d^N Y \\ &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_{Z_1}(\Theta \delta(Z_1, Z_2)X)) g(R_{Z_2}(Y)) e^{-iX \cdot Y} d^N X d^N Y, \end{aligned} \quad (3.9)$$

where Z_1 and Z_2 are the embedding point corresponding to z_1 and z_2 , $X, Y \in T_{Z_2}\mathcal{M}$ and the integration is w.r.t. the non-vanishing components (i.e. maximally N) and the scalar product \cdot is w.r.t. the flat Minkowski (or Euclidean) metric η_{AB} (δ_{AB}). Moreover, $\delta_B^{A'}(Z_1, Z_2)$ is the operator of geodesic transport from $T_{Z_2}\mathcal{M}$ to $T_{Z_1}\mathcal{M}$,⁴ the cut-off function $\chi \in C_0^\infty(\mathbb{R}^N \times \mathbb{R}^N)$ is chosen such that condition $\chi(0, 0) = 1$ is fulfilled.

⁴We use here the same notation as in [Mor03]. This bitensor is as well notated by $g_\rho^{\mu'}$, see for example [Hac10, Chapter III].

Note that when both points are equal, i.e. $z_1 = z_2$, the product assumes the form previously defined for a single point. The following propositions deal with the uniqueness of this definition. In particular, we prove that various sensible definitions all agree with the product given in the former Definition 3.12.

PROPOSITION 3.13. *Let θ be a real, constant deformation parameter and $\pi \in \Gamma^\infty(\Lambda^2(T_{Z_1}\mathcal{M}))$ a Poisson bivector that define the matrix $\Theta := \theta\pi$. Then, the definition of the deformed product for two different points given in Definition 3.12 is equivalent to the following product,*

$$f(z_1) \star_\theta g(z_2) = \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon U, \epsilon V) f(R_{Z_1}(\Theta U)) g(R_{Z_2}(\delta(Z_2, Z_1)V)) e^{-iU \cdot V} d^N U d^N V,$$

where Z_1 and Z_2 are the embedding point corresponding to z_1 and z_2 , $U, V \in T_{Z_1}\mathcal{M}$ and the integration is w.r.t. the non-vanishing components, where $\delta_B^{A'}(Z_2, Z_1)$ is the operator of the geodesic transport from $T_{Z_1}\mathcal{M}$ to $T_{Z_2}\mathcal{M}$.

Proof. To see the equivalence to Definition 3.12 we perform the following variable substitutions,

$$\delta(Z_2, Z_1)V = Y, \quad U = \delta(Z_1, Z_2)X$$

where $X, Y \in T_{Z_2}\mathcal{M}$. Using the relation of the determinant of the parallel propagator, namely ([PPV11, Equation (7.7)]),

$$\det \delta_B^{A'} = \frac{\det(-\eta)}{\det(-\eta')} = 1$$

and using the inverses of the parallel-transports (see [PPV11, Equation 5.7]) the scalar product

$$\begin{aligned} (U, V)_{z_1} &= \delta_A^{B'}(Z_1, Z_2) X_{B'} \delta_{C'}^A(Z_1, Z_2) Y^{C'} \\ &= \delta_{C'}^A(Z_2, Z_1) \delta_A^{B'}(Z_1, Z_2) X_{B'} Y^{C'} \\ &= (X, Y)_{z_2} \end{aligned}$$

where in the last lines we used relation $\delta_{C'}^A(Z_1, Z_2) = \delta_{C'}^A(Z_2, Z_1)$ (see [PPV11, Equation (5.8)]) and the inverse relation. □

PROPOSITION 3.14. *Let θ be a real, constant deformation parameter and $\pi \in \Gamma^\infty(\Lambda^2(T_{Z_2}\mathcal{M}))$ a Poisson bivector that define the matrix $\Theta^{A'B'}(Z_2) = \theta\pi^{A'B'} = \delta_A^{A'}(Z_2, Z_1)\delta_B^{B'}(Z_2, Z_1)\Theta^{AB}(Z_1)$. Then, the definition of the deformed product for two different points given in Definition 3.12 is equivalent to the following product,*

$$\begin{aligned} f(z_1) \star_\theta g(z_2) &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_{Z_1}(\delta(Z_1, Z_2)(\Theta(Z_2)X))) g(R_{Z_2}(Y)) e^{-iX \cdot Y} d^N X d^N Y, \\ &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_{Z_1}(\delta(Z_1, Z_2)\Theta(Z_2)\delta(Z_2, Z_1)U)) g(R_{Z_2}(\delta(Z_2, Z_1)V)) \\ &\quad \times e^{-iU \cdot V} d^N U d^N V, \end{aligned}$$

where Z_1 and Z_2 are the embedding point corresponding to z_1 and z_2 , $X, Y \in T_{Z_2}\mathcal{M}$, $U, V \in T_{Z_1}\mathcal{M}$, and the integration is w.r.t. the non-vanishing components.

Proof. The first equality is easily proven, since

$$\delta(Z_1, Z_2)\delta(Z_1, Z_2)\Theta(Z_2) = \Theta(Z_1)$$

and the second equality is proven analogously to the former Proposition. □

LEMMA 3.15. *For a smooth bivector field $\pi \in \Gamma^\infty(\Lambda^2(T_{Z_1}\mathcal{M}))$ and the operator of the geodesic transport $\delta(Z_1, Z_2)$ from $T_{Z_2}\mathcal{M}$ to $T_{Z_1}\mathcal{M}$, we define the following object*

$$\pi^{AC'}(Z_1, Z_2) := \pi^{AB}(Z_1) \delta_B^{C'}(Z_1, Z_2).$$

Then, for the tensor $\Theta(Z_1, Z_2)$ and the orthogonal projector $P(Z_2)$ the following relation holds,

$$\pi^{AA'}(Z_1, Z_2) = \pi^{AD'}(Z_1, Z_2) P_{D'}^{A'}(Z_2)$$

Proof. The proof is analogous to the proof of Lemma 2.36. \square

PROPOSITION 3.16. *The deformed product for two different points, according to Definition 3.12, is given up to second order in the deformation parameter as follows,*

$$\begin{aligned} f(z_1) \star_\theta g(z_2) &= f(z_1)g(z_2) - i\Theta^{AB'}(Z_1, Z_2) \partial_A f(Z_1) \partial_{B'} g(Z_2) \\ &\quad - \Theta^{ABC'D'} \partial_A \nabla_B f(Z_1) \partial_{C'} \nabla_{D'} g(Z_2) + \mathcal{O}(\Theta^3), \end{aligned}$$

where we defined

$$\begin{aligned} \Theta^{AB'}(Z_1, Z_2) &:= \Theta^{AB}(Z_1) \delta_B^{B'}(Z_1, Z_2), \\ \Theta^{ABC'D'}(Z_1, Z_2) &:= \frac{1}{4} (\Theta^{AC} \Theta^{BD} + \Theta^{AD} \Theta^{BC}) \delta_D^{D'}(Z_1, Z_2) \delta_C^{C'}(Z_1, Z_2), \end{aligned}$$

for the functions $(f, g) \in \mathcal{D}_{\rho, \delta}^m(\mathcal{M})$.

Proof. We prove the proposition by using the Taylor expansion of the retractions in order of the deformation parameter θ ,

$$f(R_{Z_1}(\Theta X)) = f(Z_1) + (\Theta \delta(Z_1, Z_2)) X^A \nabla_A f(Z_1) + \frac{1}{2} (\Theta \delta(Z_1, Z_2)) X^A \nabla_{\Theta \delta(Z_1, Z_2) X} \nabla_A f(Z_1)$$

We use the following notation

$$\nabla_A f(Z_1) = \text{Proj}_Z(\partial_A \bar{f}(Z_1)), \quad \nabla_Z \nabla_A f(Z_1) = \text{Proj}_{Z_1}(Z^B \partial_B \nabla_A f(Z_1)) \quad (3.10)$$

plugging this into the deformation formula we have up to second order in θ ,

$$\begin{aligned} f(z_1) \star_\theta g(z_2) &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) f(R_{Z_1}(\Theta \delta(Z_1, Z_2) X)) g(R_{Z_2}(Y)) e^{-iX \cdot Y} d^N X d^N Y \\ &= \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \left(f(Z_1) + (\Theta \delta(Z_1, Z_2) X)^A \nabla_A f(Z_1) \right. \\ &\quad \left. + \frac{1}{2} (\Theta \delta(Z_1, Z_2) X)^A \text{Proj}_{Z_1}((\Theta \delta(Z_1, Z_2) X)^B \partial_B \partial_A f(Z_1)) \right) g(R_{Z_2}(Y)) e^{-iX \cdot Y} d^N X d^N Y \\ &= f(z_1)g(z_2) + \Theta^{AB} \delta_B^{C'}(Z_1, Z_2) \nabla_A f(Z_1) \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) X_{C'} g(R_{Z_2}(Y)) e^{-iX \cdot Y} d^N X d^N Y \\ &\quad + \frac{1}{2} \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) ((\Theta \delta(Z_1, Z_2) X)^A \text{Proj}_{Z_1}((\Theta \delta(Z_1, Z_2) X)^B \partial_B \nabla_A f(Z_1))) \\ &\quad \times g(R_{Z_2}(Y)) e^{-iX \cdot Y} d^N X d^N Y \\ &= f(z_1)g(z_2) - i\Theta^{AB} \delta_B^{C'}(Z_1, Z_2) \nabla_A f(Z_1) \nabla_{C'} g(Z_2) \\ &\quad + \frac{1}{4} \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) ((\Theta \delta(Z_1, Z_2) X)^A \text{Proj}_{Z_1}((\Theta \delta(Z_1, Z_2) X)^B \partial_B \nabla_A f(Z_1))) \\ &\quad \times Y^{A'} Y^{B'} P_{A'}^{D'}(Z_2) \partial_{B'} \nabla_{D'} g(Z_2) e^{-iX \cdot Y} d^N X d^N Y \\ &= f(z_1)g(z_2) - i\Theta^{AB} \delta_B^{C'}(Z_1, Z_2) \nabla_A f(Z_1) \nabla_{C'} g(Z_2) \\ &\quad + \frac{1}{4} \Theta^{AC} \delta_C^{C'}(Z_1, Z_2) \Theta^{BD} \delta_D^{D'}(Z_1, Z_2) \partial_B \nabla_A f(Z_1) P_{A'}^{D'}(Z_2) \partial_{B'} \nabla_{D'} g(Z_2) \\ &\quad \times \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) X_{C'} X_{D'} Y^{A'} Y^{B'} e^{-iX \cdot Y} d^N X d^N Y \\ &= f(z_1)g(z_2) - i\Theta^{AB} \delta_B^{C'}(Z_1, Z_2) \nabla_A f(Z_1) \nabla_{C'} g(Z_2) \\ &\quad - \frac{1}{4} \Theta^{AC} \delta_C^{C'}(Z_1, Z_2) \Theta^{BD} \delta_D^{D'}(Z_1, Z_2) \partial_B \nabla_A f(Z_1) P_{A'}^{E'}(Z_2) \partial_{B'} \nabla_{E'} g(Z_2) \\ &\quad \times \left(\delta_{D'}^{A'} \delta_{C'}^{B'} + \delta_{D'}^{B'} \delta_{C'}^{A'} \right) \\ &= f(z_1)g(z_2) - i\Theta^{AB} \delta_B^{C'}(Z_1, Z_2) \nabla_A f(Z_1) \nabla_{C'} g(Z_2) \\ &\quad - \frac{1}{4} \Theta^{AC} \delta_C^{C'}(Z_1, Z_2) \Theta^{BD} \delta_D^{A'}(Z_1, Z_2) \partial_B \nabla_A f(Z_1) P_{A'}^{E'}(Z_2) \partial_{C'} \nabla_{E'} g(Z_2) \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{4}\Theta^{AC}\delta_C^{A'}(Z_1, Z_2)\Theta^{BD}\delta_D^{C'}(Z_1, Z_2)\partial_B\nabla_{A'}f(Z_1)P_{A'}^{E'}(Z_2)\partial_{C'}\nabla_{E'}g(Z_2) \\
& = f(z_1)g(z_2) - i\Theta^{AB}\delta_B^{C'}(Z_1, Z_2)\nabla_{A'}f(Z_1)\nabla_{C'}g(Z_2) \\
& -\frac{1}{4}(\Theta^{AC}\Theta^{BD} + \Theta^{AD}\Theta^{BC})\delta_D^{A'}(Z_1, Z_2)\delta_C^{C'}(Z_1, Z_2)\partial_B\nabla_{A'}f(Z_1)P_{A'}^{E'}(Z_2)\partial_{C'}\nabla_{E'}g(Z_2),
\end{aligned}$$

Next, we use Lemma 2.36 and Lemma 3.15 and simplify the terms to

$$\begin{aligned}
f(z_1) \star_{\theta} g(z_2) & = f(z_1)g(z_2) - i\Theta^{AC'}(Z_1, Z_2)\partial_{A'}f(Z_1)\partial_{C'}g(Z_2) \\
& -\frac{1}{4}\left(\Theta^{AC'}(Z_1, Z_2)\Theta^{BD'}(Z_1, Z_2) + \Theta^{AD'}(Z_1, Z_2)\Theta^{BC'}(Z_1, Z_2)\right)\partial_{A'}\nabla_{B'}f(Z_1)(Z_2)\partial_{C'}\nabla_{D'}g(Z_2)
\end{aligned}$$

□

Although the deformed product is well-defined for a single point, its extension to two points requires an additional tool of geodesic parallel transport. In the context of quantum field theory and associated expectation values w.r.t. the two-point function, the inversion is the case. Specifically, the two-point function is singular for a single point, prompting the requirement for regularization. After regularization, one takes the coinciding point limit via the geodesic parallel transport.

4 Quantum Field Theory in Globally Hyperbolic Spacetimes

4.1 The Quantized Scalar Field

In this section we give a quick overview and fix the notation of the free quantized scalar field ϕ in a globally hyperbolic spacetime (\mathcal{M}, g) . For a more complete discussion, see [Wal94]. Due to the global hyperbolicity of (\mathcal{M}, g) there exists unique advanced and retarded fundamental solutions $G^{\text{adv/ret}}$ (“Green’s operators”) for the Klein-Gordon operator $\square_g - m^2$ defined on smooth scalar test-functions on \mathcal{M} , where \square_g is the Laplace-Beltrami operator w.r.t. the manifold (\mathcal{M}, g) . Next, we define a $*$ -algebra $\mathcal{A} = \mathcal{A}(\mathcal{M}, g)$ generated by the elementary algebraic objects, namely the field operators $\phi(F)$, $F \in C_0^\infty(\mathcal{M})$, and a unit element $\mathbf{1}$. It is called the CCR algebra of the quantum field $\phi(F)$ over the manifold \mathcal{M} and it fulfills (see [KM15, Definition 8])

$$\begin{aligned}
(i) \quad & F \mapsto \phi(F) \text{ is } \mathbb{R}\text{-linear} & (ii) \quad & \phi((\nabla^a \nabla_a - m^2)F) = 0 \\
(iii) \quad & \phi(F)^* = \phi(F) & (iv) \quad & [\phi(F), \phi(H)] = iE(F, H) \cdot \mathbf{1}, \quad F, H \in C_0^\infty(\mathcal{M})
\end{aligned}$$

where $E(F, F')$ is the causal Green’s function (or causal propagator) defined by the following expression

$$E(F, H) = \int_{\mathcal{M}} (F(x)(G^{\text{adv}}H)(x) - F(x)(G^{\text{ret}}H)(x)) \, d\text{vol}_g(x), \quad F, H \in C_0^\infty(\mathcal{M}),$$

where $d\text{vol}_g$ is the volume form w.r.t. the metric g .

An important consequence of the global hyperbolicity of the manifold (\mathcal{M}, g) is given by the causality condition (see [KM15, Proposition 4]).

PROPOSITION 4.1. *Referring to $\mathcal{A}(\mathcal{M}, g)$, $\phi(F)$ and $\phi(H)$ commute if the supports of F and H are causally separated, i.e.*

$$E(F, H) = 0,$$

if the support of F does not intersect $J_M^+(suppH) \cup J_M^-(suppH)$.

4.2 States and GNS Theorem

Up to this point, the field operators and their generating algebra are abstract algebraic objects. To represent those algebraic objects on a Hilbert space we use the GNS (Gelfand-Naimark-Segal) construction. This theorem supplies us with such a representation in the following manner, see for example [KM15, Chapter 1, Theorem 1] and references therein. For a given state ω over the (unital) $*$ -algebra \mathcal{A} , i.e. a positive \mathbb{C} -linear map $\omega : \mathcal{A} \rightarrow \mathbb{C}$ which is *positive* ($\omega(a^*a) \geq 0$, for all $a \in \mathcal{A}$) and *normalized*, one obtains a quadruple $(\mathcal{H}_\omega, D_\omega, \pi_\omega, \Psi_\omega)$. This quadruple consists of a

complex Hilbert space \mathcal{H}_ω , a dense subspace $\mathcal{D}_\omega \subset \mathcal{H}_\omega$, a $*$ -representation $\pi_\omega : \mathcal{A}(\mathcal{M}, g) \rightarrow \mathcal{L}(\mathcal{D}_\omega)$ of \mathcal{A} on \mathcal{H}_ω with domain \mathcal{D}_ω and a cyclic and separating vector Ψ_ω . The field operators are given by densely defined symmetric operators,

$$\phi_\omega(F) = \pi_\omega(\phi(f)) : \mathcal{D}_\omega \rightarrow \mathcal{H}_\omega.$$

The expectation value of n functions or the n -point function of elements of the algebra \mathcal{A} is determined by

$$\omega_n(F_1, \dots, F_n) = \langle \Psi_\omega | \pi_\omega(\phi(F_1)) \cdots \pi_\omega(\phi(F_n)) \Psi_\omega \rangle.$$

Next, we list some basic properties of the two point function in the following proposition ([KM15, Proposition 7]).

PROPOSITION 4.2. *Consider a state $\omega : \mathcal{A}(\mathcal{M}) \rightarrow \mathbb{C}$ and define $P := \square_M + m^2 + \xi R$ (for arbitrarily fixed values of $m^2, \xi \in \mathbb{R}$). The two-point function, ω_2 satisfies the following relations*

$$\omega_2(F, F) \geq 0, \quad \omega_2(F', F) = \omega_2(\bar{F}, F'), \quad \text{Im } \omega_2(F, F') = \frac{1}{2}E(F, F'), \quad (4.1)$$

$$\omega_2(PF, F') = 0 = \omega_2(F, PF'), \quad \omega_2(F, F') - \omega_2(F', F) = iE(F, F') \quad (4.2)$$

for $F, F' \in C_0^\infty(\mathcal{M})$.

Furthermore, we define the Hadamard parametrix, where for more details see [KW91],[KM15, Section 3.2]. Let $\sigma(x, x')$ be the half of the squared geodesic distance (the Synge function) between x and x' and let in a convex neighborhood C of a four dimensional spacetime the parametrix H_ϵ (that is a limit of the Hadamard parametrix) be given as,

$$H_\epsilon(x, y) = \frac{u(x, y)}{(2\pi)^2 \sigma_\epsilon(x, y)} + v(x, y) \log \left(\frac{\sigma_\epsilon(x, y)}{\lambda^2} \right) \quad (4.3)$$

where u, v are C^∞ functions on $C \times C$, with $u(x, x) = 1$ and where $x, y \in C$, T is any local time coordinate increasing towards the future, $\lambda > 0$ a length scale and

$$\sigma_\epsilon(x, y) \stackrel{\text{def}}{=} \sigma(x, y) + 2i\epsilon(T(x) - T(y)) + \epsilon^2, \quad (4.4)$$

finally, the cut in the complex domain of the log function is assumed along the negative axis.

DEFINITION 4.3. *A (not necessarily quasifree) state ω on $\mathcal{A}(\mathcal{M})$ and its two point function ω_2 are **Hadamard** if $\omega_2 \in \mathcal{D}'(M \times M)$ and every point of \mathcal{M} admits an open normal neighborhood C where*

$$\omega_2(x, y) - H_{0^+}(x, y) = w(x, y) \quad \text{for some } w \in C^\infty(C \times C). \quad (4.5)$$

Here 0^+ indicates the standard weak distributional limit as $\epsilon \rightarrow 0^+$.

4.3 The Microlocal Spectrum Condition and Hadamard

This section is a recollection of definitions and theorems about microlocal analysis, see [Few08, Chapter 6] (and references therein) for more details. The principle idea of microlocal analysis is to use the decay properties of the Fourier-transformation, to gain information about the underlying singular structure. This has to do with the fundamental connection between smooth functions and their rapidly decaying Fourier transformations. In the context of microlocal analysis, the wavefront set plays a fundamental role and we define it next, [Few08, Definition 6.1].

DEFINITION 4.4. *A) If $u \in \mathcal{D}'(\mathbb{R}^n)$, a pair $(x, k) \in \mathbb{R}^n \times (\mathbb{R}^n \setminus \{0\})$ is a regular direction for u if there exist*

1. $\phi \in C_0^\infty(\mathbb{R}^n)$ with $\phi(x) \neq 0$
2. a conic neighborhood V of k

3. constants C_N , $N \in \mathbb{N}$ so that

$$\left| \hat{\phi}u(k) \right| < \frac{C_N}{1 + |k|^N}, \quad \forall k \in V$$

i.e. $\hat{\phi}u$ decays rapidly as $k \rightarrow \infty$ in V .

B) The wavefront set of u is defined to be

$$WF(u) = \{(x, k) \in \mathbb{R}^n \times (\mathbb{R}^n \setminus \{0\}) : (x, k) \text{ is not a regular direction for } u\}.$$

Some useful properties of the wavefront set are given in the following.

1. If $f \in C^\infty$ it has an empty wavefront set $WF(f) = \emptyset$.
2. $WF(\alpha u + \beta v) \subset WF(u) \cup WF(v)$ for $u, v \in \mathcal{D}'(\mathbb{R}^n)$, $\alpha, \beta \in \mathbb{C}$.
3. If P is any differential operator with smooth coefficients, then

$$WF(Pu) \subset WF(u)$$

for any $u \in \mathcal{D}'(\mathbb{R}^n)$.

Before turning to quantum field theory and applying the tools of microlocal analysis, we define \mathcal{N} to be the bundle of nonzero null covectors on \mathcal{M} :

$$\mathcal{N} = \{(x, \xi) \in T^*M : \xi \text{ a non-zero null at } p\}.$$

The first application is the realization that the wavefront set of the Minkowski vacuum ω_2^{vac} is a subset of the product $\mathcal{N}^+ \times \mathcal{N}^-$, i.e.

$$WF(\omega_2^{vac}) \subset \mathcal{N}^+ \times \mathcal{N}^-,$$

where

$$\mathcal{N}^\pm = \{(p, \xi) \in \mathcal{N} : \xi \text{ is future(+)/past(-)directed}\}.$$

Since for QFT in globally hyperbolic spacetimes the singularity structure of the state has to be analog to the one given in the Minkowski case (due to the equivalence principle), we elevate this rule to a general context and give the following definition, [Few08, Definition 6.2] (see as well [Rad92, Rad96, BFK96]),

DEFINITION 4.5. A state ω obeys the **Microlocal Spectrum Condition** (μSC) if

$$WF(\omega_2) \subset \mathcal{N}^+ \times \mathcal{N}^-.$$

Intuitively this condition states that the singular behaviour of the two-point function is of positive frequency in the first slot and negative in the second. This corresponds to the well-known frequency splitting that is given in the Minkowski case. We further state another important result in the context of states in QFT.

THEOREM 4.6. If ω and ω' obey μSC then

$$\omega_2 - \omega'_2 \in C^\infty(\mathcal{M} \times \mathcal{M}), \quad (4.6)$$

i.e. the μSC determines an equivalence of class of states under equality of the two-point functions modulo C^∞ .

In particular, if we start with a field algebra (defined in the former section) and pick two states ω and ω' to represent the algebra (using the GNS-construction) it might happen that these two states are not unitary equivalent. Hence, they might lead to different (or nonequivalent) representations of the field algebra. Yet, the significant physical content of a QFT can be the two-point function.

Hence, if the two different two point functions of the different states ω and ω' fulfill the μSC then their difference is smooth, which equates their corresponding wavefront set. This fact is seen by the properties of the wavefront set, i.e.

$$WF(\omega - \omega') = 0 \quad WF(\omega) = WF(\omega').$$

In addition to the advantages of the wavefront set we state a theorem by Radzikowski ([Rad92, Rad96]) about the connection to the Hadamard condition.

THEOREM 4.7. *The μSC is equivalent to the **Hadamard condition**.*

Hence, given a state that is Hadamard, which is proven to exist for all massive Klein-Gordon theories, the theorem tells us that all such states fulfill the physically reasonable microlocal spectrum condition.

5 Deformation of QFT in Globally Hyperbolic Spacetimes

5.1 Representation of the Deformation

As before we start with a globally hyperbolic spacetime (\mathcal{M}, g) , a $*$ -algebra $\mathcal{A} = \mathcal{A}(\mathcal{M}, g)$ generated by the field operators that satisfy the Klein Gordon equation and the choice of a state $\omega : \mathcal{A}(\mathcal{M}, g) \rightarrow \mathbb{C}$, i.e. a normalized positive linear functional. This state then determines as described in Subsection 4.2 the quadruple $(\mathcal{H}_\omega, D_\omega, \pi_\omega, \Psi_\omega)$. The deformation, w.r.t. the noncommutative Rieffel product, enters the framework not via the states or the representation but via changing the point-wise product of the representations of the algebra. Summarizing, the *deformed* 2-point function is given in the following definition.

DEFINITION 5.1. *For a $*$ -algebra $\mathcal{A} = \mathcal{A}(\mathcal{M}, g)$ defined on a globally hyperbolic spacetime (\mathcal{M}, g) generated by Klein-Gordon fields, the (formal) deformed 2-point function is defined as the expectation value of the generalized Rieffel product (Definition 3.12) of the test functions of the representations of the fields, i.e.*

$$\begin{aligned} \omega_2^\Theta(\phi(F_1)\phi(F_2)) &:= \langle \Psi_\omega | \pi_\omega(\phi(F_1)) \star_\Theta \pi_\omega(\phi(F_2)) \Psi_\omega \rangle \\ &= \int F_1(x_1) \star_\Theta F_2(x_2) \langle \Psi_\omega | \pi_\omega(\phi(x_1)) \pi_\omega(\phi(x_2)) \Psi_\omega \rangle d\text{vol}_g(x_1) d\text{vol}_g(x_2) \\ &= \iint \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) \alpha_{\Theta \delta(X_1, X_2)X}(F_1(X_1)) \alpha_Y(F_2(X_2)) e^{-i(X, Y)_{X_2}} d^N X d^N Y \\ &\quad \times \omega_2(\phi(X_1)\phi(X_2)) d^N X_1 d^N X_2. \end{aligned}$$

for test-functions $F_1, F_2 \in C_0^\infty(\mathcal{M})$.

THEOREM 5.2. *The deformed smeared two-point function given in Definition 5.1 is for functions $(F_1, F_2) \in \mathcal{D}_{0, \rho, \delta}^m(\mathcal{M})$ well-defined.*

Proof. To prove that the oscillatory integral is well-defined we write the (smeared) deformed two-point function as follows.

$$\omega_2^\Theta(\phi(F_1)\phi(F_2)) = \lim_{\epsilon \rightarrow 0} \iint \chi(\epsilon X, \epsilon Y) b_\Theta(X, Y, F_1, F_2) e^{-iX \cdot Y} d^N X d^N Y \quad (5.1)$$

with $b_\Theta(X, Y, F_1, F_2)$ formally given by

$$b_\Theta(X, Y, F_1, F_2) = \iint \alpha_{\Theta \delta(X_1, X_2)X}(F_1(X_1)) \alpha_Y(F_2(X_2)) \langle \Psi_\omega | \phi(X_1)\phi(X_2) \Psi_\omega \rangle d^N X_1 d^N X_2. \quad (5.2)$$

Following Ref. [Hör71, Proposition 1.1.11, Lemma 1.2.1, Proposition 1.2.2] the oscillatory integral is well defined if we prove that the function $b_\Theta(X, Y, F_1, F_2)$ belongs to the symbol space and thus we have for $|b_\Theta(X, Y, F_1, F_2)|$ the following equality

$$\left| \iint F_1(R_{X_1}(\Theta \delta(X_1, X_2)X)) F_2(R_{X_2}(Y)) \langle \Psi_\omega | \phi(X_1)\phi(X_2) \Psi_\omega \rangle d^N X_1 d^N X_2 \right|$$

$$\begin{aligned}
&= \left| \iint F_1(R_{X_1}(\Theta \delta(X_1, X_2)X)) F_2(R_{X_2}(Y)) \langle \phi(X_1)\Psi_\omega | \phi(X_2)\Psi_\omega \rangle d^N X_1 d^N X_2 \right| \\
&\leq C \|\alpha_{\Theta \delta X}(F_1)\alpha_Y(F_2)\|_\infty,
\end{aligned}$$

where we used the Cauchy-Schwarz inequality, the conclusion in [FM21, Proof Proposition III.3] and the supremum-norm $\|G\|_\infty$ of the product of the product of the translated and retracted test-functions $G_{X,Y} := \alpha_{\Theta \delta X}(F_1)\alpha_Y(F_2)$ determined by

$$\|G_{X,Y}\|_\infty = \sup_{X_1, X_2 \in \mathcal{M}} |G_{X,Y}(X_1, X_2)|,$$

where we write $X_1, X_2 \in \mathcal{M}$ for $X_1, X_2 \in \mathbb{R}^N : F(X_{1,2}) = 0$, where F is the local defining function. Since we choose $G_{X,Y} \in \mathcal{D}_{0,\rho,\delta}^m(\mathcal{M})$ the absolute value of the function $b_\Theta(X, Y, F_1, F_2)$ and derivatives thereof are bounded such that b is a symbol and therefore the Integral 5.1 is well-defined. \square

5.2 The Microlocal Spectrum Condition under Deformation Quantization

In this section we prove that the micro-local spectrum condition holds for deformed states, if the undeformed state is Hadamard. This will be proven up to the second order in the deformation parameter (see the expansion given in Proposition 3.16).

LEMMA 5.3. *The deformed two point function defined by the star product in Proposition 3.16 is, up to second order in Θ , explicitly presented by*

$$\begin{aligned}
\omega_2^\Theta(X_1, X_2) &= P^\Theta \omega_2(X_1, X_2) \\
&= \omega_2(X_1, X_2) - i \partial_A \partial_{B'} (\Theta^{AB'}(X_1, X_2) \omega_2(X_1, X_2)) \\
&\quad - \nabla_A \partial_B \nabla_{C'} \partial_{D'} (\Theta^{ABC'D'}(X_1, X_2) \omega_2(X_1, X_2)),
\end{aligned}$$

where P^Θ is a fourth-order differential operator with smooth coefficients, that depend on the Poisson bivector, the orthogonal projection and the parallel transport.

Proof. Using Definition 5.1, Proposition 3.16 and assuming that the functions $(F_1, F_2) \in \mathcal{D}_{\rho,\delta}^m(\mathcal{M})$ we write the smeared deformed two-point function as

$$\begin{aligned}
\omega_2^\Theta(\phi(F_1)\phi(F_2)) &:= \langle \Psi_\omega | \pi_\omega(\phi(F_1)) \star_\Theta \pi_\omega(\phi(F_2)) \Psi_\omega \rangle \\
&= \int F_1(X_1) \star_\Theta F_2(X_2) \omega_2(\phi(X_1)\phi(X_2)) d^N X_1 d^N X_2 \\
&=: \int F_1(X_1) F_2(X_2) \omega_2^\Theta(\phi(X_1)\phi(X_2)) d^N X_1 d^N X_2 \\
&= \int F_1(X_1) F_2(X_2) \left(\omega_2(X_1, X_2) - i \partial_A \partial_{B'} (\Theta^{AB'}(X_1, X_2) \omega_2(X_1, X_2)) \right. \\
&\quad \left. - \nabla_A \partial_B \nabla_{C'} \partial_{D'} (\Theta^{ABC'D'}(X_1, X_2) \omega_2(X_1, X_2)) \right) + \mathcal{O}(\theta^3) \\
&= \int F_1(X_1) F_2(X_2) P^\Theta \omega_2(X_1, X_2) + \mathcal{O}(\theta^3)
\end{aligned}$$

where in the last lines we inserted the explicit form of the deformed product given in Proposition 3.16 and we integrated by parts. The smoothness of the differential operator P follows from the smoothness of the Poisson bivector (see Definition 2.34), the orthogonal projection (see Lemma 2.15) and the parallel transport. \square

THEOREM 5.4. *Let the state ω_2 obey the microlocal spectrum condition. Then, the deformed state ω_2^Θ obeys the microlocal spectrum condition up to second order in Θ , i.e.*

$$WF(\omega_2^\Theta) \subset \mathcal{N}^+ \times \mathcal{N}^-,$$

where the deformed two-point function ω_2^Θ is given explicitly in Lemma 5.3.

Proof. By using the smoothness of the differential operator P^Θ (see Lemma 5.3), the following chain follows from Property 3 of the wavefront set and the fact that ω_2 fulfills the microlocal spectrum condition

$$WF(\omega_2^\Theta) = WF(P^\Theta \omega_2) \subset WF(\omega_2) \subset \mathcal{N}^+ \times \mathcal{N}^-.$$

□

Corollary 5.5. *The deformed two-point function ω_2^Θ is Hadamard.*

Proof. The fact that the deformed two-point function is Hadamard follows directly from Theorem 5.4 and Theorem 4.7. □

THEOREM 5.6. *Let the deformed states ω_2^θ and $\omega_2^{\theta'}$ be defined using either two different isomorphic embeddings or/and retractions. Then, the deformed two-point functions ω_2^Θ and $\omega_2^{\Theta'}$ have the same wavefront set, i.e.*

$$WF(\omega_2^\Theta) = WF(\omega_2^{\Theta'}).$$

Proof. Due to the smoothness of the embedding or retractions, the deformed two-point functions ω_2^Θ and $\omega_2^{\Theta'}$ obey the μ SC and according to Theorem 4.6, their difference is smooth. Hence, by Property 1 we have

$$WF(\omega_2^\Theta - \omega_2^{\Theta'}) = 0$$

and Property 2 equates the two wavefront sets. □

6 Discussion and Outlook

In this work, we provided a specific extension of the Rieffel deformation for all globally hyperbolic spacetimes with a certain Poisson structure. This novel deformation technique is a symbiosis between the Kontsevich deformation [Kon03] (since we define a star product for Poisson manifolds), the Rieffel product and the Fedosov quantization (due to the condition $\nabla\pi = 0$). It is tailored such that it can be directly applied to quantum field theories and in particular to the central physical object, namely the two-point function.

In addition to the introduction of a quantum-type feature (non-commutativity) to spacetime itself (see Proposition 3.10), we proved that if a state fulfills the microlocal spectral condition, i.e. has a certain sensible singularity structure, then its deformed counterpart will fulfill the μ SC as well. Thus, by the Radzikowski theorem we conclude that the deformed quasi-free states are Hadamard.

This result sheds light into the question of whether the behavior at short distances (Hadamard criterion) has a physical foundation due to the break down of the spacetime continuum at scales around the Planck length. In particular, this means that the introduction of a quantum nature (or non-commutativity) to spacetime does not violate the equivalence principle, from a quantum field theoretical perspective.

These basic deformation principles can be applied to the various achievements and insights of quantum field theory in curved spacetimes to evaluate its physical validity as being a first-order approximation to a theory of quantum gravity. In addition to introducing a quantum nature to spacetime, the deformation quantization can be considered a first-order approximation since in the limit $\Theta \rightarrow 0$ we obtain the undeformed, i.e. regular, version of QFT in CST.

The first place to look for the physical consequences imposed by the deformation quantization are quantum energy inequalities. These are restrictions obtained from QFT towards the domain of violation of classical energy conditions. These inequalities are sensitive to the spacetime geometry, e.g. restrict certain exotic spacetimes and effects (such as wormholes, time-machines or warp drives), and take into account the quantum behaviour implied by

quantum field theory. Hence, implementing a quantum behavior on spacetime itself may change these inequalities in a non-trivial manner and therewith induce new physical and measurable effects.

Although we defined a deformed product for the case of globally hyperbolic spacetimes, by using the existence of an isometric embedding, no such result exists for non-globally hyperbolic manifolds. These cases are in particular interesting when investigating black-holes. Although no isometric embedding will exist in general (e.g. in the presence of closed time-like curves), one still can define deformations for QFTs in non-globally hyperbolic spacetimes by using the very idea of how to study these theories [Kay92]. In principle, one requires the existence of a globally hyperbolic neighborhood N for every point in a given spacetime \mathcal{M} . Hence, the observable algebra of the spacetime \mathcal{M} restricted to N is equal to the algebra one would obtain if one regards N as a globally hyperbolic spacetime in its own right. Regarding deformation, we can therefore deform the product in the globally hyperbolic neighborhood N by using the embedding formalism and thus define a deformed state.

Acknowledgements

The author would like to thank R. Verch for various discussions on this topic. Moreover, we extend our gratitude to H. Grosse, in particular on discussions related to the quasi-free deformed states. We would further like to extend our thanks to M. Fröb whose encouragement, sharp remarks and questions shaped many ideas in this paper. O. Müller is gratefully acknowledged for various discussions regarding the embedding formalism. Furthermore, the author expresses his gratitude to R. Ballal, D. Vidal-Cruzprieto and P. Dorau for various discussions that cleared the ideas set in this paper. Last but not least, the author thanks Nicolas Boumal for promptly answering questions in regards to optimization techniques.

References

- [Ahl94] Dharam Vir Ahluwalia. Quantum measurements, gravitation, and locality. *Phys. Lett. B*, 339:301–303, 1994.
- [AMS07] P.-A. Absil, R. Mahony, and R. Sepulchre. *Optimization Algorithms on Matrix Manifolds*. Princeton University Press, USA, 2007.
- [BD82] N. D. Birrell and P. C. W. Davies. *Quantum Fields in Curved Space*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1982.
- [BEE96] John. K. Beem, Paul. E. Ehrlich, and Kevin. L. Easley. *Global Lorentzian Geometry*, volume 202 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, second edition, 1996.
- [BF00] Romeo Brunetti and Klaus Fredenhagen. Microlocal analysis and interacting quantum field theories: Renormalization on physical backgrounds. *Communications in Mathematical Physics*, 208(3):623–661, Jan 2000.
- [BFK96] R. Brunetti, K. Fredenhagen, and M. Köhler. The microlocal spectrum condition and wick polynomials of free fields on curved spacetimes. *Communications in Mathematical Physics*, 180(3):633–652, Oct 1996.
- [BGP08] Christian Baer, Nicolas Ginoux, and Frank Pfäeffle. Wave Equations on Lorentzian Manifolds and Quantization. *arXiv e-prints*, page arXiv:0806.1036, Jun 2008.
- [BLS11] Detlev Buchholz, Gandalf Lechner, and Stephen J. Summers. Warped Convolutions, Rieffel Deformations and the Construction of Quantum Field Theories. *Commun. Math. Phys.*, 304:95–123, 2011.
- [BM17] Edwin J. Beggs and Shahn Majid. Poisson–riemannian geometry. *Journal of Geometry and Physics*, 114:450–491, 2017.

- [Bou03] Mohamed Boucetta. Riemann–poisson manifolds and kähler–riemann foliations. *Comptes Rendus Mathématique*, 336(5):423–428, 2003.
- [Bou23] Nicolas Boumal. *An introduction to optimization on smooth manifolds*. Cambridge University Press, 2023.
- [BS08] Detlev Buchholz and Stephen J. Summers. Warped Convolutions: A Novel Tool in the Construction of Quantum Field Theories. In Erhard Seiler and Klaus Sibold, editors, *Quantum Field Theory and Beyond*, pages 107 – 121. World Scientific, Singapore, 2008.
- [BW07] Dorothea Bahns and Stefan Waldmann. Locally noncommutative space-times. *Reviews in Mathematical Physics*, 19(03):273–305, April 2007.
- [Dav75] P. C. W. Davies. Scalar production in Schwarzschild and Rindler metrics. *Journal of Physics A Mathematical General*, 8(4):609–616, April 1975.
- [Dav84] P. C. W. Davies. Particles do not Exist. In S. M. Christensen, editor, *Quantum Theory of Gravity*, page 66. 1984.
- [DFR95] Sergio Doplicher, Klaus Fredenhagen, and John E. Roberts. The Quantum structure of space-time at the Planck scale and quantum fields. *Commun. Math. Phys.*, 172:187–220, 1995.
- [DLMM11] Claudio Dappiaggi, Gandalf Lechner, and Eric Morfa-Morales. Deformations of Quantum Field Theories on Spacetimes with Killing Vector Fields. *Commun. Math. Phys.*, 305:99–130, 07 2011.
- [DLS20] Jan Dereziński, Adam Latosiński, and Daniel Siemssen. Pseudodifferential weyl calculus on (pseudo-)riemannian manifolds. *Annales Henri Poincaré*, 21(5):1595–1635, 2020.
- [Fed94] Boris V. Fedosov. A simple geometrical construction of deformation quantization. *Journal of Differential Geometry*, 40(2):213 – 238, 1994.
- [Few00] Christopher J Fewster. A general worldline quantum inequality. *Classical and Quantum Gravity*, 17(9):1897–1911, Apr 2000.
- [Few08] Christopher J. Fewster. Lectures on quantum field theory in curved spacetime. Preprint, 2008.
- [Few12] Christopher J. Fewster. Lectures on quantum energy inequalities, 2012.
- [FH90] Klaus Fredenhagen and Rudolf Haag. On the Derivation of Hawking Radiation Associated With the Formation of a Black Hole. *Commun. Math. Phys.*, 127:273, 1990.
- [FH05] Christopher J. Fewster and Stefan Hollands. Quantum energy inequalities in two-dimensional conformal field theory. *Reviews in Mathematical Physics*, 17(05):577–612, Jun 2005.
- [FM21] M. B. Fröb and A. Much. Strict deformations of quantum field theory in de sitter spacetime. *Journal of Mathematical Physics*, 62(6):062302, Jun 2021.
- [FNW81] S. A. Fulling, F. J. Narcowich, and Robert M. Wald. Singularity Structure of the Two Point Function in Quantum Field Theory in Curved Space-time. II. *Annals Phys.*, 136:243–272, 1981.
- [For91] L. H. Ford. Constraints on negative-energy fluxes. *Phys. Rev. D*, 43:3972–3978, Jun 1991.
- [FP78] L. H. Ford and Roger Penrose. Quantum coherence effects and the second law of thermodynamics. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 364(1717):227–236, 1978.
- [FS08] Christopher J. Fewster and Calvin J. Smith. Absolute quantum energy inequalities in curved spacetime. *Annales Henri Poincaré*, 9(3):425–455, May 2008.

- [Ful73] Stephen A. Fulling. Nonuniqueness of canonical field quantization in riemannian space-time. *Phys. Rev. D*, 7:2850–2862, May 1973.
- [Ful89] Stephen A. Fulling. Aspects of Quantum Field Theory in Curved Space-time. *London Math. Soc. Student Texts*, 17:1–315, 1989.
- [GL07] Harald Grosse and Gandalf Lechner. Wedge-Local Quantum Fields and Noncommutative Minkowski Space. *JHEP*, 0711:012, 2007.
- [Goe74] G. H. Goedecke. On global embedding. *Journal of Mathematical Physics*, 15(6):789–791, 1974.
- [GRS97] Israel Gelfand, Vladimir Retakh, and M. Shubin. Fedosov manifolds, 1997.
- [GW13] C. Gérard and M. Wrochna. Construction of hadamard states by pseudo-differential calculus. *Communications in Mathematical Physics*, 325(2):713–755, Nov 2013.
- [Hac10] Thomas-Paul Hack. On the backreaction of scalar and spinor quantum fields in curved spacetimes - from the basic foundations to cosmological applications, 2010.
- [Hac15] T.P. Hack. *Cosmological Applications of Algebraic Quantum Field Theory in Curved Spacetimes*. SpringerBriefs in Mathematical Physics. Springer International Publishing, 2015.
- [Haw75] S. W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43(3):199 – 220, 1975.
- [HNW07] Jakob G. Heller, Nikolai Neumaier, and Stefan Waldmann. A c^* -algebraic model for locally noncommutative spacetimes. *Letters in Mathematical Physics*, 80(3):257–272, June 2007.
- [Hör71] Lars Hörmander. Fourier integral operators. I. *Acta Mathematica*, 127(none):79 – 183, 1971.
- [Hör89] L. Hörmander. *The Analysis of Linear Partial Differential Operators I: Distribution Theory and Fourier Analysis*. 2. Edition, Springer, 1989.
- [HW01] Stefan Hollands and Robert M. Wald. Local wick polynomials and time ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics*, 223(2):289–326, Oct 2001.
- [HW02] Stefan Hollands and Robert M. Wald. Existence of local covariant time ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics*, 231(2):309–345, Dec 2002.
- [Kay92] Bernard S. Kay. The Principle of locality and quantum field theory on (nonglobally hyperbolic) curved space-times. *Rev. Math. Phys.*, 4:167–195, 1992.
- [KM15] Igor Khavkine and Valter Moretti. Algebraic qft in curved spacetime and quasifree hadamard states: An introduction. *Mathematical Physics Studies*, page 191–251, 2015.
- [Kon03] Maxim Kontsevich. Deformation quantization of poisson manifolds. *Letters in Mathematical Physics*, 66(3):157–216, December 2003.
- [KW91] Bernard S. Kay and Robert M. Wald. Theorems on the Uniqueness and Thermal Properties of Stationary, Nonsingular, Quasifree States on Space-Times with a Bifurcate Killing Horizon. *Phys. Rept.*, 207:49–136, 1991.
- [Lee13] J.M. Lee. *Introduction to Smooth Manifolds*. Graduate Texts in Mathematics. Springer New York, 2013.
- [LW16] Gandalf Lechner and Stefan Waldmann. Strict deformation quantization of locally convex algebras and modules. *J. Geom. Phys.*, 99:111–144, 2016.

- [MM11] Eric Morfa-Morales. Deformations of quantum field theories on de Sitter spacetime. *J. Math. Phys.*, 52:102304, 2011.
- [Mor03] Valter Moretti. Comments on the stress-energy tensor operator in curved spacetime. *Communications in Mathematical Physics*, 232(2):189–221, January 2003.
- [MS11] Olaf Mueller and Miguel Sanchez. Lorentzian manifolds isometrically embeddable in L^N . *Trans. Am. Math. Soc.*, 363:5367–5379, 2011.
- [Mü01] Wolfgang Mück. Ideas on the semi-classical path integral over embedded manifolds. *Fortschritte der Physik*, 49(4-6):607–615, 2001.
- [Oec00] Robert Oeckl. Untwisting noncommutative and the equivalence of quantum field theories. *Nuclear Physics B*, 581(1-2):559–574, aug 2000.
- [Oec01] Robert Oeckl. Braided quantum field theory. *Communications in Mathematical Physics*, 217(2):451–473, mar 2001.
- [PPV11] Eric Poisson, Adam Pound, and Ian Vega. The motion of point particles in curved spacetime. *Living Reviews in Relativity*, 14(1), September 2011.
- [Rad92] M. J. Radzikowski. The Hadamard condition and Kay’s conjecture in (axiomatic) quantum field theory on curved space-time. *PhD thesis, Princeton University*, October 1992.
- [Rad96] M. J. Radzikowski. Micro-local approach to the Hadamard condition in quantum field theory on curved space-time. *Commun. Math. Phys.*, 179:529–553, 1996.
- [Rej16] Kasia Rejzner. *Perturbative Algebraic Quantum Field Theory*, volume 10.1007/978-3-319-25901-7. Springer International Publishing, 2016.
- [Rie93] M.A. Rieffel. Deformation quantization for actions of \mathbb{R}^d . *Memoirs A.M.S.*, 506, 1993.
- [Ros65] Joe Rosen. Embedding of various relativistic riemannian spaces in pseudo-euclidean spaces. *Rev. Mod. Phys.*, 37:204–214, Jan 1965.
- [RS22] J.W. Robbin and D.A. Salamon. *Introduction to Differential Geometry*. Springer Studium Mathematik (Master). Springer Berlin Heidelberg, 2022.
- [SKM⁺03] Hans Stephani, Dietrich Kramer, Malcolm MacCallum, Cornelius Hoenselaers, and Eduard Herlt. *Exact solutions of Einstein’s field equations*. Cambridge monographs on mathematical physics. Cambridge University Press, 2nd ed edition, 2003.
- [SV01] Hanno Sahlmann and Rainer Verch. Microlocal spectrum condition and hadamard form for vector-valued quantum fields in curved spacetime. *Reviews in Mathematical Physics*, 13(10):1203–1246, Oct 2001.
- [SV08] Jan Schlemmer and Rainer Verch. Local thermal equilibrium states and quantum energy inequalities. *Annales Henri Poincaré*, 9(5):945–978, Aug 2008.
- [Unr76] W. G. Unruh. Notes on black-hole evaporation. *Phys. Rev. D*, 14(4):870–892, August 1976.
- [Vai94] Izu Vaisman. *Lectures on the Geometry of Poisson Manifolds*. Progress in Mathematics No. 118. Birkhäuser, 1 edition, 1994.
- [Ver01] Rainer Verch. A spin statistics theorem for quantum fields on curved space-time manifolds in a generally covariant framework. *Commun. Math. Phys.*, 223:261–288, 2001.
- [Wal75] Robert M. Wald. On particle creation by black holes. *Communications in Mathematical Physics*, 45:9–34, 1975.
- [Wal94] Robert M. Wald. *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*. Chicago Lectures in Physics. University of Chicago Press, 1994.

-
- [Wal07] S. Waldmann. *Poisson-Geometrie und Deformationsquantisierung: Eine Einführung*. Masterclass. Springer Berlin Heidelberg, 2007.