

RING EPIMORPHISMS FROM PATH ALGEBRAS TO MATRIX ALGEBRAS

Jakub Kopřiva

November 25, 2021

Abstract

In this text, we are concerned with ring epimorphisms, and more specifically universal localisations, from path algebras to matrix algebras. We are mainly focused on constructing ring epimorphisms and universal localisations by extending them from smaller path algebras to larger path algebras. At first, we discuss some simple results, and then we present generalizations thereof in several directions.

Key words: Ring epimorphism, universal localisation, path algebra, matrix algebra, free associative algebra

MSC2010: 16S10, 16S50, 16S85 (Primary) 16G20 (Secondary)

Contents

1	Ring epimorphisms and universal localisations	2
1.1	Change of rings	2
1.2	Ring epimorphisms	3
1.3	Universal localisations	5
2	Ring epimorphisms from path algebras to matrix rings	7
2.1	Preliminaries	8
2.2	Simple examples	10
2.3	Extending ring epimorphisms	12
2.4	Extending universal localisations	16

Introduction

This document is structured into two sections: in section 1, we review, and, in section 2, we set out to construct ring epimorphisms from path algebras to matrix algebras. Our starting point is formed by Proposition 19, which yields a ring epimorphism to matrix algebra $M_n(k)$ arising from a brick, a module with trivial endomorphisms, and Theorem 20, which shows that a ring epimorphism to $M_n(k\langle x_1, \dots, x_n \rangle)$ can be obtained, informally speaking,

by adding an additional edge to a brick. If, moreover, the brick has no self-extensions, we obtain universal localisations.

Our goal in this text is to apply the ideas underpinning the basic results in broader contexts. Among other results, we generalize Proposition 19 and Theorem 20 in Proposition 23 and Theorem 24, respectively, for ring epimorphisms that are not necessarily universal localisations. Moreover, in Proposition 32, we present a generalization of Theorem 20 in terms of universal localisations using discussion on extending universal localisations in general.

1 Ring epimorphisms and universal localisations

In this section, we give an overview of ring epimorphisms and related results; we also derive some useful corollaries of these results.

We begin by recalling the functors between categories of modules arising from a homomorphism between rings (all rings in this text are assumed to be unital); then, we define ring epimorphisms. The concept of ring epimorphism is examined from both ring-theoretic and category-theoretic point of view. Finally, we focus on universal localisations, specific type of ring epimorphisms that can be viewed as a vast, but natural, generalization of the concept of localisation in a multiplicative set in commutative algebra.

Among other sources, this section draws on nice reviews of the subject in subsection 2.1 *Ring epimorphisms* in [MV15] and subsection 2.1 *Localisations of rings* in [HMS⁺18].

1.1 Change of rings

In this subsection, we briefly recapitulate some notions regarding change of rings induced by a homomorphism between them.

Given a ring homomorphism $f : A \rightarrow B$, there are several functors between $\text{Mod-}A$ and $\text{Mod-}B$:

- (i) *restriction of scalars* functor $\varphi^* : \text{Mod-}B \rightarrow \text{Mod-}A$ such that:

$$f^*(N_B) = N_A$$

- (ii) *induced module* functor $f_! : \text{Mod-}A \rightarrow \text{Mod-}B$ such that:

$$f_!(M_A) = M_A \otimes_A B_B$$

- (iii) *coinduced module* functor $f_* : \text{Mod-}A \rightarrow \text{Mod-}B$ such that:

$$f_*(M_A) = \text{Hom}_A({}_B B_A, M_A)$$

These functors form two pairs of adjoint functors:

$$f_! : \text{Mod-}A \dashv \text{Mod-}B : f^* \text{ and } f^* : \text{Mod-}B \dashv \text{Mod-}A : f_*$$

In other words, for any $M \in \mathbf{Mod}\text{--}A$ and $N \in \mathbf{Mod}\text{--}B$, we have the following functorial isomorphisms:

$$\mathrm{Hom}_B(M_A \otimes_A {}_A B_B, N_B) \cong \mathrm{Hom}_A(M_A, N_A),$$

$$\mathrm{Hom}_A(N_A, M_A) \cong \mathrm{Hom}_B(N_B, \mathrm{Hom}_A({}_B B_A, M_A)).$$

For details, we refer the reader to section 10.4 *Tensor Product of Modules* in [DF04] for instance.

1.2 Ring epimorphisms

In this section, we try to give a rather representative overview of results on ring epimorphisms and to illustrate the variety approaches with which this concept have been treated in the literature thus far.

Definition 1 (Ring epimorphism). *We say that a ring homomorphism $\varphi : R \rightarrow S$ is a ring epimorphism if for any two ring homomorphisms $\varrho_1, \varrho_2 : S \rightarrow T$ with $\varrho_1 \varphi = \varrho_2 \varphi$, also $\varrho_1 = \varrho_2$.*

Definition 2 (Dominion). *Let $A \subseteq B$ be rings. The dominion of A in B is the maximal subset D of B such that homomorphisms from B that agree on A must agree on D .*

Remark. It is easy to see that dominion D needs to be a subring of B . For more details, see section 1 *Preliminaries and corrections* in [Isb69] or section *Maximal epic subrings and dominions in simple artinian rings* in chapter 7 in [Sch85].

Theorem 3 (Ring-theoretic characterization of ring epimorphisms, Theorem 1.1 in [Isb69] attributed to Silver and Mazet and [Ste12]). *For a ring homomorphism $f : A \rightarrow B$, the following statements are equivalent:*

- (i) *f is a ring epimorphism;*
- (ii) *The dominion of $\varphi(A)$ in B is equal to B .*
- (iii) *$f \otimes_A B = B \otimes_A f : B \rightarrow B \otimes_A B$ is an isomorphism of $B\text{--}B$ –bimodules;*
- (iv) *$\mathrm{Coker}(f) \otimes_A B = 0$.*

Moreover, the dominion of $\varphi(A)$ is a set of all $XPY \in B$ where X is a $1 \times n$ matrix over B , Y is $n \times 1$ matrix over B , and P is $n \times n$ matrix over $\varphi(A)$ such that XP is a $1 \times n$ matrix over $\varphi(A)$ and PY a $n \times 1$ matrix over $\varphi(A)$.

Theorem 4 (Categorical characterization of ring epimorphisms, [Ste12]). *For a ring homomorphism $f : A \rightarrow B$, the following statements are equivalent:*

- (i) *f is a ring epimorphism;*
- (ii) *The restriction of scalars functor $f^* : \mathbf{Mod}\text{--}B \rightarrow \mathbf{Mod}\text{--}A$ (respectively the restriction of scalars from $B\text{--}\mathbf{Mod}$ to $A\text{--}\mathbf{Mod}$) is fully faithful.*

Theorem 5 (Categorical view of ring epimorphisms, Theorem 1.2 in [GDLP87], [GL91], [Iya03]). *There is a bijection between:*

- (i) *ring epimorphisms $A \rightarrow B$ up to equivalence (given by isomorphisms of targets thereof);*
- (ii) *bireflective subcategories \mathcal{X}_B of $\mathbf{Mod}\text{-}A$ (respectively, $A\text{-}\mathbf{Mod}$), i.e., strict full subcategories of closed under products, coproducts, kernels and cokernels.*

If, furthermore, A is a finitely generated algebra over k , this bijection can be restricted between:

- (i) *k -algebra epimorphisms $A \rightarrow B$ up to equivalence, where B is a finitely generated algebra over k ;*
- (ii) *bireflective subcategories \mathcal{X}_B of $\mathbf{mod}\text{-}A$ (respectively, $A\text{-}\mathbf{mod}$), i.e., strict full functorially finite subcategories closed under kernels and cokernels.*

Now, we establish some useful consequences of the results above.

Corollary 6 (Ring epimorphisms and Morita equivalence). *Provided that $f : A \rightarrow B$ is a ring epimorphism and A' is Morita equivalent to A , then there is a ring epimorphism $f' : A' \rightarrow B'$ with B' Morita equivalent to B .*

Proof. Let $F : \mathbf{Mod}\text{-}A \rightarrow \mathbf{Mod}\text{-}A'$ be an equivalence of categories, and let \mathcal{X}_B be the bireflective subcategory corresponding to the ring epimorphism $f : A \rightarrow B$ by Theorem 5.

The functor F restricts to an equivalence between \mathcal{X}_B and $(\mathcal{X}_B)'$ its essential image under F . As F is an equivalence of categories, $(\mathcal{X}_B)'$ is strictly full subcategory of $\mathbf{Mod}\text{-}A'$, closed under products, coproducts, kernels and cokernels, hence bireflective, since \mathcal{X}_B is such.

From Theorem 5, we obtain that there is a ring epimorphism $f' : A' \rightarrow B'$ such that the essential image of the restriction functor $(f')^*$ is $(\mathcal{X}_B)'$. We have that $\mathbf{Mod}\text{-}B$ is equivalent to \mathcal{X}_B , \mathcal{X}_B to $(\mathcal{X}_B)'$, and $(\mathcal{X}_B)'$ to $\mathbf{Mod}\text{-}B'$. This yields that $\mathbf{Mod}\text{-}B$ and $\mathbf{Mod}\text{-}B'$ are equivalent. \square

Lemma 7. *Let A be a ring, $M, N, L \in \mathbf{Mod}\text{-}A$, and*

$$\begin{array}{ccc} M & & \\ \downarrow \varphi & \searrow \psi & \\ L & \xrightarrow{\varrho} & N \end{array}$$

be a commutative square in \mathbf{Set} with φ, ψ A -homomorphisms, φ surjective, then ϱ is an A -homomorphism.

Proof. Choose $\ell_1, \ell_2 \in L$ and $a \in A$. Since φ is surjective, there are m_1, m_2 such that $\varphi(m_i) = \ell_i, i = 1, 2$. Now, using commutativity of the diagram, we compute $\varrho(\ell_1 + \ell_2) = \varrho(\varphi(m_1 + m_2)) = \psi(m_1 + m_2) = \psi(m_1) + \psi(m_2) = \varrho(\ell_1) + \varrho(\ell_2)$. For $\varrho(\ell_1 \cdot a) = \varrho(\ell_1) \cdot a$, we proceed similarly. \square

Theorem 8. *A ring homomorphism $f : A \rightarrow B$ is a ring epimorphism if and only if, for all finitely generated $M, N \in \mathbf{Mod}\text{-}B$, $\text{Hom}_A(M, N) = \text{Hom}_B(M, N)$ and, for any $L \in \mathbf{Mod}\text{-}B$ and an A -homomorphism $\psi : B \rightarrow L$, the $\text{Im } \psi$ is contained in a finitely generated B -submodule of L .*

Proof. (\Rightarrow) The first part follows from Theorem 3. The second part from the fact that homomorphisms in $\text{Hom}_A(B, L) = \text{Hom}_S(B, L)$ correspond to cyclic B -submodules of L via their image for any $L \in \mathbf{Mod}\text{-}B$.

(\Leftarrow) Let $M, N \in \mathbf{Mod}\text{-}B$ and $\tau : M \rightarrow N$ be an A -homomorphism between them. Choose $\varrho : B^{(I)} \twoheadrightarrow M$ surjective B -homomorphism. For each $i \in I$, there is a N_i a finitely generated B -submodule such that $\text{Im } \tau \varrho \mu_i$ lies inside it. Since, by our assumption, $\text{Hom}_A(B, N_i) = \text{Hom}_B(B, N_i)$, we have that $\tau \varrho \mu_i$ is an B -homomorphism. By universal property of $B^{(I)}$, there is a unique B -homomorphism $\psi : B^{(I)} \rightarrow N$ such that, for all $i \in I$, $\tau \varrho \mu_i = \psi \mu_i$. Due to its uniqueness, we get that $\tau \varrho = \psi$, we and apply Lemma 7 to obtain that τ is an B -homomorphism. \square

Corollary 9. *A ring homomorphism $f : A \rightarrow B$ that turns B into a finitely generated A -module is a ring epimorphism if and only if, for all finitely generated $M, N \in \mathbf{Mod}\text{-}B$, $\text{Hom}_A(M, N) = \text{Hom}_B(M, N)$.*

Proof. For any $L \in \mathbf{Mod}\text{-}B$ and an A -homomorphism $\psi : B \rightarrow L$, the $\text{Im } \psi$ is a finitely generated A -module since B is such. A B -submodule of L generated by these finitely many generators clearly contains $\text{Im } \psi$. \square

1.3 Universal localisations

In this subsection, we gather some notions and results regarding universal localisations that will be of use later in this text.

Theorem 10 (Universal localisation, [Sch85]). *Let Σ be a set of morphisms between finitely generated projective right A -modules. Then there are: a ring A_Σ and a homomorphism of rings $f_\Sigma : A \rightarrow A_\Sigma$, called the universal localisation of A at Σ , such that*

- (i) f_Σ is Σ -inverting, i.e. if $\alpha : P \rightarrow Q$ belongs to Σ , then $\alpha \otimes_A A_\Sigma : P \otimes_A A_\Sigma \rightarrow Q \otimes_A A_\Sigma$ an isomorphism of right A_Σ -modules, and
- (ii) f_Σ is universal Σ -inverting, i.e. for any Σ -inverting ring homomorphism $\psi : A \rightarrow B$, there is a unique ring homomorphism $\bar{\psi} : A_\Sigma \rightarrow B$ such that $\psi = \bar{\psi} f_\Sigma$.

Moreover, the homomorphism f_Σ is a ring epimorphism and $\text{Tor}_1^A(A_\Sigma, A_\Sigma) = 0$. The restrictions of A_Σ -modules are modules $M \in \mathbf{Mod}\text{-}A$ such that, for each $\alpha \in \Sigma$, the map $\text{Hom}_A(\alpha, M)$ is invertible.

Theorem 11 (Theorems 4.7 and 4.8 in [Sch85]). *Let $f : A \rightarrow B$ be a ring epimorphism; then the following conditions are equivalent:*

- (i) $\text{Ext}_A^1 = \text{Ext}_B^1$ on $\text{Mod-}B$;
- (ii) $\text{Tor}_1^A(B, B) = 0$;
- (iii) $\text{Tor}_1^A = \text{Tor}_1^B$ on $\text{Mod-}B$;
- (iv) $\text{Ext}_R^1 = \text{Ext}_S^1$ on $B\text{-Mod}$

Moreover, a universal localisation satisfies all these conditions.

Remark. Provided that $f : A \rightarrow B$ a ring epimorphism turns B into a finitely generated A -module and A is Noetherian, using a finite free presentation of B in the proof in [Sch85] (Theorem 4.8 therein) yields that $\text{Ext}_A^1 = \text{Ext}_B^1$ on finitely generated B -modules implies $\text{Tor}_1^A(B, B) = 0$.

Definition 12 ([GL91]). *A ring epimorphism $f : A \rightarrow B$ is called homological ring epimorphism if $\text{Tor}_n^A(B, B) = 0$ for all $n \in \mathbb{N}$.*

Theorem 13 ([KŠ10]). *For hereditary rings, universal localisations and homological epimorphisms coincide.*

Similarly as in the previous section, we now derive some useful conclusions from the results above:

Corollary 14 (Universal localisations and Morita equivalence). *Provided that $f : A \rightarrow B$ is a universal localisation and A' is Morita equivalent to A ; then there is a universal localisation $f' : A' \rightarrow B'$ with B' Morita equivalent to B .*

Proof. Let $B = A_\Sigma$ for some Σ , and let $F : \text{Mod-}A \rightarrow \text{Mod-}A'$ be an equivalence of categories. As F sends finitely generated projectives in $\text{Mod-}A$ to finitely generated projectives in $\text{Mod-}A'$, $F(\Sigma)$ is also a set of maps between finitely generated projectives. Let $f' : A' \rightarrow B'$ be the ring epimorphism by Corollary 9 such that B' is Morita equivalent to A_Σ . Since F restricted between \mathcal{X}_{A_Σ} and $\mathcal{X}_{B'}$ is an equivalence, then f' is $F(\Sigma)$ -inverting. The fact that f' is universal $F(\Sigma)$ -inverting is then proved by going back to $\text{Mod-}A$ by equivalence F and using that f is universal Σ -inverting. Consequently, $B' = A'_{F(\Sigma)}$. \square

Lemma 15. *Let $A \subseteq B$ be rings, and Σ be a set of maps between finitely generated projectives over A . Denote $\Sigma' = \{\alpha \otimes_A B \mid \alpha \in \Sigma\}$; this is a set of maps between finitely generated projectives over B . If a B -module N can be viewed as a restriction of a $B_{\Sigma'}$ -module, then N_A is a restriction of a A_Σ -module.*

Proof. By Theorem 10, N can be viewed as a restriction of a $B_{\Sigma'}$ -module if $\text{Hom}_B(\alpha', N)$ is an isomorphism for all $\alpha' \in \Sigma'$. Write α' as $\alpha \otimes_A B : P_1 \otimes_A B \rightarrow P_2 \otimes_A B$ for P_i finitely generated projectives over A .

The adjunction of induced module and restriction of scalars functors yields functorial isomorphisms:

$$\text{Hom}_B(P_i \otimes_A B, N) \cong \text{Hom}_A(P_i, N_A)$$

Provided that $\text{Hom}_B(\alpha', N)$ is an isomorphism, the map $\text{Hom}_A(\alpha, N_A)$ needs to be an isomorphism as well.

Thus, if $\text{Hom}_B(\alpha', N)$ is an isomorphism for all $\alpha' \in \Sigma'$ (and N is a restriction of a $B_{\Sigma'}$ -module), then $\text{Hom}_A(\alpha, N_A)$ is an isomorphism for all $\alpha \in \Sigma$ (and N_A is a restriction of a A_Σ -module). \square

Lemma 16. *Let Σ be a set of morphisms between finitely generated projective right A -modules and $I \subseteq A$ be an ideal of A . Then $A_\Sigma / \langle I \rangle_{A_\Sigma}$ and $(A/I)_{\Sigma \otimes_A A/I}$ are isomorphic as rings where $\Sigma \otimes A/I = \{\alpha \otimes_A A/I \mid \alpha \in \Sigma\}$*

Proof. We prove this by showing that both $A'_1 = A_\Sigma / \langle I \rangle_{A_\Sigma}$ and $A'_2 = (A/I)_{\Sigma \otimes_A A/I}$ are pushouts of the following diagram: $A/I \leftarrow A \rightarrow A_\Sigma$; hence they are isomorphic as rings.

We begin our discussion with A'_1 . Naturally, we have a projection $\pi_{\langle I \rangle_{A_\Sigma}} : A_\Sigma \rightarrow A'_1$. If we denote $f_\Sigma : A \rightarrow A_\Sigma$ the universal localisation, we have a map $\pi_{\langle I \rangle_{A_\Sigma}} \circ f_\Sigma : A \rightarrow A'_1$. Clearly, this map maps elements of I to zero. By the homomorphism theorem, there exists a unique map $g : A/I \rightarrow A'_1$ such that $g \circ \pi_I = \pi_{\langle I \rangle_{A_\Sigma}} \circ f_\Sigma$. Suppose we have two maps $h_1 : A/I \rightarrow B$ and $h_2 : A_\Sigma \rightarrow B$ such that $h_1 \circ \pi_I = h_2 \circ f_\Sigma$. Since $h_2 \circ f_\Sigma$ maps all elements of I to zero, h_2 needs to map all elements of $\langle I \rangle_{A_\Sigma}$ to zero. This gives us, by the homomorphism theorem, that there is a unique homomorphism h' such that $h_2 = h' \circ \pi_{\langle I \rangle_{A_\Sigma}}$. Now, we have two ring homomorphisms h_1 and $h' \circ g$ from A/I to B ; they agree on A , since $h_1 \circ \pi_I = h_2 \circ f_\Sigma = (h' \circ \pi_{\langle I \rangle_{A_\Sigma}}) \circ f_\Sigma = (h' \circ g) \circ \pi_I$. As π_I is a ring epimorphism, h_1 and $h' \circ g$ need to be equal. The uniqueness of h' is warranted by the universal property of the quotient $A_\Sigma / \langle I \rangle_{A_\Sigma}$.

Second, let us have A'_2 . We have a universal localisation $f_{\Sigma \otimes_A A/I} : A/I \rightarrow A'_2$; we proceed in an analogous manner. Given that $f_{\Sigma \otimes_A A/I}$ is Σ -inverting, by Theorem 8 stating the universal property of universal localisation, there exists a unique ring homomorphism $g : A_\Sigma \rightarrow A'_2$ such that $g \circ f_\Sigma = f_{\Sigma \otimes_A A/I} \circ \pi_I$. Assume we have two maps $h_1 : A/I \rightarrow B$ and $h_2 : A_\Sigma \rightarrow B$ such that $h_1 \circ \pi_I = h_2 \circ f_\Sigma$. This gives us that h_1 is $\Sigma \otimes_A A/I$ -inverting yielding a unique ring homomorphism $h' : A'_2 \rightarrow B$ arising from its universal property as a universal localisation such that $h_1 = h' \circ f_{\Sigma \otimes_A A/I}$. Again, we have two maps h_2 and $h' \circ g$ from A_Σ to B . However, as there two homomorphism agree when precomposed with f_Σ , the fact that f_Σ is a ring epimorphisms gives us that, indeed, $h' \circ g = h_2$. \square

Remark. Provided that A is an R -algebra, $A_\Sigma / \langle I \rangle_{A_\Sigma}$ and $(A/I)_{\Sigma \otimes_A A/I}$ are isomorphic as R -algebras.

2 Ring epimorphisms from path algebras to matrix rings

In this section, we discuss ring epimorphisms, and more specifically universal localisations, from a finite-dimensional path algebra of quiver Q over k , an

algebraically closed field, to matrices over algebras over k , particularly of type $k\langle x_1, \dots, x_n \rangle$.

This section is structured as follows. At first, we recall some preliminary results. Subsequently, we look at an example of a ring epimorphism (universal localisation) in question arising from bricks (exceptional modules) over path algebra kQ and their simple extensions. Inspired by these examples of constructing ring epimorphisms and universal localisations from smaller to larger path algebras, we provide generalizations of them in several directions.

The quiver Q is a finite acyclic quiver $Q = (Q_0, Q_1, t, h)$. For brevity, we denote the corresponding path algebra $A = kQ$ in the first two subsections. We note that A is a finite dimensional hereditary algebra and that $\text{Rep}_k(Q)$ and $\text{Mod-}A$ (or respectively, $\text{rep}_k(Q)$ and $\text{mod-}A$) are naturally equivalent. For foundations of the theory of representations of quivers and finite-dimensional algebras, we refer the reader to [ASS06].

2.1 Preliminaries

In this section, we frequently work with matrix rings $M_n(R)$ over a given ring R . For this end, we establish some notation. We set:

$$e_{ij} = \begin{cases} 1 & \text{at position } ij; \\ 0 & \text{elsewhere.} \end{cases}$$

Recall that $e_{ij}e_{kl} = \delta_{jk} \cdot e_{il}$ and that $e_{ii}Ae_{jj} = a_{ij} \cdot e_{ij}$ for any $A \in M_n(R)$.

Definition 17 ([Coh06]). *Given a ring R , we say that:*

- (i) *R is projective-free if every finitely generated projective module is free and of unique rank ;*
- (ii) *R is a free ideal ring (fir) if all one-sided ideals of R are free as R -modules;*
- (iii) *R is a semifir if all finitely generated one-sided ideals of R are free as R -modules.*

Remark. All semifirs are projective-free by Corollary 2.3.4 in [Coh06].

Remark. For any n , the free associative algebra $k\langle x_1, \dots, x_n \rangle$ is a fir. See Corollary 2.5.2 in [Coh06] and ensuing discussion therein.

Using these notions and results, we formulate, and prove some auxiliary results for this section.

Lemma 18. *Let S be a projective-free ring, and $\varphi : R \rightarrow M_n(S)$ be a ring homomorphism. Given a full set of orthogonal idempotents $e_1, \dots, e_m \in R$, there exists an automorphism ψ of S such that, for each $i = 1, \dots, m$, the image $\psi\varphi(e_i)$ is equal to:*

$$\psi\varphi(e_i) = \sum_j e_{jj}.$$

Proof. If $e_1, \dots, e_m \in R$ form a full set of orthogonal idempotents, then their images, $\varphi(e_1), \dots, \varphi(e_m) \in M_n(S)$, also form such a set. Every $\varphi(e_i)$ yields a direct summand of S^n ; since S is projective-free, each such summand is a free S -module of a unique rank r_i . We also have A_i an $n \times r_i$ matrix over S and B_i an $r_i \times n$ matrix over S such that $A_i B_i = \varphi(e_i)$ and $B_i A_i = I_{r_i}$, the $r_i \times r_i$ identity matrix. The matrix A_i gives the injection from S^{r_i} to S^n ; whereas the matrix B_i represents the projection from S^n to S^{r_i} .

As the idempotents $\varphi(e_1), \dots, \varphi(e_m) \in M_n(S)$ are orthogonal, we obtain that $B_j A_i = 0$ for $i \neq j$. This yields that:

$$\begin{pmatrix} B_1 \\ \dots \\ B_m \end{pmatrix} (A_1 \dots A_m) = I_n.$$

The identity matrix I_n should be thought of a block diagonal matrix with identity matrices I_{r_1}, \dots, I_{r_m} on the diagonal. As S^n is projective, the image of $B^T = (B_1 \dots B_m)^T$ splits; we obtain that $S^n \cong P \oplus S^n$. Since S is projective-free, the module P needs to be free; however, due to unique rank of S^n the module S^n is zero. Thus, the matrices B and $A = (A_1 \dots A_m)$ give mutually an S -automorphisms of S^n .

Therefore, we may set ψ to be the inner automorphism of $M_n(S)$ given by conjugation by A^{-1} . We calculate $\psi\varphi(e_i)$ as follows:

$$\psi\varphi(e_i) = A^{-1}\varphi(e_i)A = \begin{pmatrix} B_1 \\ \dots \\ B_m \end{pmatrix} A_i B_i (A_1 \dots A_m) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & I_{r_i} & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Where the matrix on the right is to be understood as a block matrix. □

Remark. Suppose $f : A \rightarrow M_m(k\langle x_1, \dots, x_n \rangle)$ is a ring homomorphism. Then, by the lemma above, there exists a dimension vector α such that for any finite-dimensional module N over $M_m(k\langle x_1, \dots, x_n \rangle)$, the dimension vector of N_A is a multiple of α . The values of α are given by unique ranks of free $k\langle x_1, \dots, x_n \rangle$ -modules given by images orthogonal idempotents $f(e_v)$ for $v \in Q_0$.

Hence, we can write $f : A \rightarrow M_\alpha(k\langle x_1, \dots, x_n \rangle)$ for a ring homomorphism from A to matrices over $k\langle x_1, \dots, x_n \rangle$ such that the A -modules that lie in the image of f^* have a multiple of α as their dimension vector.

One could also establish this fact by means of algebraic geometry. We note that, for any finite-dimensional module N over the algebra $M_m(k\langle x_1, \dots, x_n \rangle)$, we have that $\dim_k N = \ell m$ for some $\ell \geq 0$. Also, up to isomorphism, any such N is parameterized by a choice of $X_1, \dots, X_n \in M_\ell(k)$ that $g_{X_1, \dots, X_n} : M_m(k\langle x_1, \dots, x_n \rangle) \rightarrow M_{\ell m}(k)$ maps $g_N(U)_{ij} = U_{ij}(X_1, \dots, X_n)$ for all $U \in M_m(k\langle x_1, \dots, x_n \rangle)$. The restriction f^* is represented by $g_{X_1, \dots, X_n} \circ f$ in this setting.

This means that each representation of $M_m(k\langle x_1, \dots, x_n \rangle)$ can be thought of as a point in $\mathbb{A}_k^{n \cdot \ell^2}$ representing the choice of n matrices $\ell \times \ell$ over k . Now, for

each $v \in Q$, the set of points of $(Y_1, \dots, Y_n) \in \mathbb{A}_k^{n \cdot \ell^2}$ such that $g_{Y_1, \dots, Y_n}[f(e_v)]$ has maximal rank α_v is dense open. Naturally, the set of points $\mathbb{A}_k^{n \cdot \ell^2}$ at which each of $f(e_v)$ are simultaneously mapped to a matrix with maximum possible rank α_v is dense open. This means that $\sum_{v \in Q_0} \alpha_v = \ell m$, as $N = \bigoplus_{v \in Q_0} g_N f(e_v) N$ as a vector space and that there no points in $\mathbb{A}_k^{n \cdot \ell^2}$ with any $f(e_v)$ mapped to a matrix of rank smaller than α_v as they would need to be of lower dimension than ℓm .

2.2 Simple examples

In this subsection, we present some simple results on ring epimorphisms and universal localisations from path algebras to matrix algebras over $k\langle x_1, \dots, x_n \rangle$. Later in this text, we will strive to find generalizations of these results.

Proposition 19. *Let $B \in \mathbf{mod}\text{-}A$ be a module, then the following are equivalent:*

- (i) *B is a brick, i.e. $\text{End}_A(B) \cong k$, (and $\text{Ext}_A^1(B, B)$ is zero),*
- (ii) *The ring homomorphism $f_B : A \rightarrow \text{End}_k(B)$ is a ring epimorphism (universal localisation).*

Proof. (i) \Rightarrow (ii): Given $B \in \mathbf{mod}\text{-}A$ a brick over A , we have a natural ring homomorphism $f_B : A \rightarrow \text{End}_k(B)$ giving B_k as a vector space the structure of an A -module. Denote $\Lambda = \text{End}_k(B)$. In accordance with Theorem 4, we prove that $(f_B)^*$ is fully faithful. By Corollary 9, as f_B makes Λ a finitely generated module over A , it suffices to show that $(f_B)^*$ is fully faithful on finite-dimensional modules.

Clearly, the essential image of $(f_B)^*$ on finitely generated Λ -modules are only direct sums of finitely many copies of B up to isomorphism. Let I, J be finite sets, then we have:

$$\text{Hom}_A(B^I, B^J) = \prod_{I, J} \text{Hom}_A(B, B) = \prod_{I, J} \text{Hom}_\Lambda(B, B) = \text{Hom}_\Lambda(B^I, B^J).$$

Therefore, $f_B : A \rightarrow \Lambda$ is a ring epimorphism. If it is also given that $\text{Ext}_A^1(B, B) = 0$, then by Remark after Theorem 11 (as A is right Noetherian and Λ is a finitely generated module over A) we obtain that $\text{Tor}_1^A(\Lambda, \Lambda) = 0$ (all higher Tor groups vanish since A is hereditary). This makes f_B a homological ring epimorphism, and, therefore, a universal localisation by Theorem 13.

(ii) \Rightarrow (i): Clearly, B is a brick over Λ , and, thus, over A since f_B is a ring epimorphism. Provided that f_B is, furthermore, a universal localisation, then $\text{Ext}_A^1(B, B) = \text{Ext}_\Lambda^1(B, B) = 0$ by Theorem 11 and since all non-trivial Ext groups vanish for Λ . \square

Theorem 20. *Let $B \in \mathbf{mod}\text{-}A$ be an exceptional module of dimension vector α ; Q' be a quiver such that $Q_0 = Q'_0$ and $Q_1 \subseteq Q'_1$, and $A' = kQ'$.*

Set $n = \sum_{e \in Q'_1 - Q_1} \alpha_{s(e)} \alpha_{t(e)}$. There exists a universal localisation $f'_B : A' \rightarrow M_\alpha(k\langle x_1, \dots, x_n \rangle)$ that restricts to the universal localisation $f_B : A \rightarrow \text{End}_k(B)$.

Proof. Let denote f_B the universal localisation of A in Σ , and set $\Sigma' = \{\sigma \otimes_A A' : P \otimes_A A' \rightarrow Q \otimes_A A' \mid \sigma : P \rightarrow Q \in \Sigma\}$. Modules over the universal localisation $A'_{\Sigma'}$ among modules over A are those for which $\text{Hom}_{A'}(\sigma \otimes_A A', M)$ are invertible for all $\sigma \otimes_A A' \in \Sigma'$.

Using Lemma 15, we obtain that for $\sigma \in \Sigma$, $\text{Hom}_{A'}(\sigma \otimes_A A', M)$ is invertible if and only if $\text{Hom}_A(\sigma, M_A)$ is invertible. Naturally, M_A is the image of M under the restriction of scalars functor induced by the inclusion $A \subseteq A'$.

From Proposition 19, we have a universal localisation $f_B : A \rightarrow M_\alpha(k)$. By change of basis, we may assume that $f_B(e_v)$ maps to a matrix with ones at positions $(n_v, n_v), (n_v + 1, n_v + 1), \dots, (n_v + \alpha_v, n_v + \alpha_v)$ and zeros elsewhere. We now define a k -algebra homomorphism from $f'_B : A' \rightarrow M_\alpha(k\langle x_1, \dots, x_n \rangle)$ as extension of f_B . It suffices to define the image of $e \in Q'_1 - Q_1$; it is defined as follows:

$$[f'_B(e)]_{ij} = \begin{cases} x_{ij,e} & n_{s(e)} \leq i \leq n_{s(e)} + \alpha_{s(e)}, n_{s(e)} \leq j \leq n_{s(e)} + \alpha_{s(e)} \\ 0 & \text{otherwise} \end{cases}$$

We observe that $f'_B(e_{s(e)}) \cdot f'_B(e) \cdot f'_B(e_{t(e)})$ for all $e \in Q'_1 - Q_1$. Due to the universal property of path algebras (Theorem 1.8 in chapter II in [ASS06]), f' is a k -algebra homomorphism. Since f_B is a ring epimorphisms, the dominion D of $f'_B(A')$ in $M_\alpha(k\langle x_1, \dots, x_n \rangle)$ contains $M_\alpha(k)$. To prove that D equals $M_\alpha(k\langle x_1, \dots, x_n \rangle)$, it suffices to show that $x_\ell \cdot I_\alpha \in D$ for all $1 \leq \ell \leq n$. For any ℓ , x_ℓ corresponds to $x_{i_\ell j_\ell, e_\ell}$ for some $1 \leq i_\ell, j_\ell \leq \alpha$ and $e_\ell \in Q'_1 - Q_1$. We obtain

$$\begin{pmatrix} x_\ell & & \\ & \ddots & \\ & & x_\ell \end{pmatrix} = \sum_{i=1}^n e_{i, i_\ell} \cdot f'_B(e_\ell) \cdot e_{j_\ell, i},$$

which lies in D as it is a subring. Matrices e_{ij} belong to D and so does $f'_B(e_\ell)$.

Finally, we show that all $A'_{\Sigma'}$ -modules are actually $M_\alpha(k\langle x_1, \dots, x_n \rangle)$ modules via f'_B . Any module M that is an $A'_{\Sigma'}$ -module is $M_\alpha(k)$ -module via the universal localisation f_B . Therefore as a representation, M is isomorphic to $[(k^{(I)})^{\alpha_v}, \varphi_e]_{v \in Q_0, e \in Q'_1}$ where $\varphi_e \in \text{Hom}((k^{(I)})^{\alpha_{s(e)}}, (k^{(I)})^{\alpha_{t(e)}})$. We can think of it as a $n_{s(e)} \times n_{s(e)}$ matrix with entries in $\text{End}_k(k^{(I)})$. Naturally, $[(k^{(I)})^{\alpha_v}, \varphi_e]_{v \in Q_0, e \in Q'_1}$ is a $M_\alpha(k\langle x_1, \dots, x_n \rangle)$ -module as x_ℓ acts as the $(i_\ell - n_{t(e_\ell)}, j_\ell - n_{s(e_\ell)})$ entry of φ_{e_ℓ} . \square

Remark. The construction of a universal localisation $A' \rightarrow M_n(k)$ based on a universal localisations $A \rightarrow M_n k$ arising from a brick B with no self-extension performed in the theorem above can be carried out in a similar manner even if B has self-extensions. However, the maps are only ring epimorphisms, not universal localisations in such a case.

2.3 Extending ring epimorphisms

In this subsection, we aim to generalize Theorem 20 by constructing ring epimorphisms, without requiring that they be universal localisations, from path algebras to matrices over $k\langle x_1, \dots, x_n \rangle$ using a well-suited representations thereof.

Lemma 21. *Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between additive categories, then F is full if and only if for every $A \in \mathcal{A}$ the group homomorphism $F_{A,A} : \text{Hom}_{\mathcal{A}}(A, A) \rightarrow \text{Hom}_{\mathcal{B}}(F(A), F(A))$ is surjective.*

Proof. Suppose that $F_{A,A}$ is surjective for every $A \in \mathcal{A}$, and let us have $f' \in \text{Hom}_{\mathcal{B}}(F(A'), F(B'))$ for some $A', B' \in \mathcal{A}$.

Consider the biproduct $A' \oplus B'$ in \mathcal{A} with morphisms $\iota_{A'} : A' \rightarrow A' \oplus B'$, $\pi_{A'} : A' \oplus B' \rightarrow A'$ and $\iota_{B'}, \pi_{B'}$ similarly.

From f' , we obtain $F(\iota_{B'}) \circ f' \circ F(\pi_{A'}) : F(A') \oplus F(B') \rightarrow F(A') \oplus F(B')$ in \mathcal{B} . Since F is additive, $F(A') \oplus F(B') \cong F(A' \oplus B')$. Our assumptions gives us that there is $f : A' \oplus B' \rightarrow A' \oplus B'$ such that $F(f) = F(\iota_{B'}) \circ f' \circ F(\pi_{A'})$. It is easy to see that $F(\pi_{B'}) \circ F(f) \circ F(\iota_{A'}) = f'$. Therefore, F is full. \square

Theorem 22. *Let $f : A \rightarrow M_n(B)$ be a homomorphism of k -algebras. Then f is a ring epimorphism if and only if the ideal I_f of $B * k\langle (v_{ij})_{i,j=1}^n \rangle$ generated by $(v_{ij})_{i,j=1}^n \cdot f(a) - f(a) \cdot (v_{ij})_{i,j=1}^n$ for $a \in A$ understood element-wise contains $v_{ii} - v_{jj}$, $v_{i'j'}$, and $bv_{ii} - v_{ii}b$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$ and $b \in B$.*

Proof. (\Leftarrow) Suppose that I_f of $B * k\langle (v_{ij})_{i,j=1}^n \rangle$ contains $v_{ii} - v_{jj}$, $v_{i'j'}$, and $bv_{ii} - v_{ii}b$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$ and $b \in B$. We will show that every A -endomorphism of a right $M_n(B)$ -module is already an $M_n(B)$ -endomorphism. Combining this with Theorem 4 and Lemma 21 above will give us that f is a ring epimorphism.

Any $M_n(B)$ -module can be obtained as follows: given a vector space W over k and a homomorphism $\mu : B \rightarrow \text{End}_k(W)$, we define a right $M_n(B)$ -module structure on W^n such that $b \in B$ act diagonally and e_{ij} via itself. The structure of an A -module on W^n is induced by restriction of scalars per $f : A \rightarrow M_n(B)$.

Suppose $(w_{ij})_{i,j=1}^n$ is an A -endomorphism of W^n ; this means that for all $a \in A$ we have $(w_{ij})_{i,j=1}^n \mu(f(a)) - \mu(f(a))(w_{ij})_{i,j=1}^n = 0$. There is a unique homomorphism from $B * k\langle (v_{ij})_{i,j=1}^n \rangle$ to $\text{End}_k(V)$ defined by μ on B and by $v_{ij} \mapsto w_{ij}$ on $k\langle (v_{ij})_{i,j=1}^n \rangle$. Moreover, since $(w_{ij})_{i,j=1}^n$ is an A -endomorphism of W^n , this homomorphism factors through $B * k\langle (v_{ij})_{i,j=1}^n \rangle / I_f$. As $v_{ii} - v_{jj}, v_{i'j'} \in I_f$ and $bv_{ii} - v_{ii}b \in I_f$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$ and $b \in B$, $(w_{ij})_{i,j=1}^n$ clearly is also an $M_n(B)$ -endomorphism.

(\Rightarrow) Assume now that $f : A \rightarrow M_n(B)$ is an epimorphism. Denote $V = B * k\langle (v_{ij})_{i,j=1}^n \rangle / I_f$ as vector space over k . Consider a homomorphism $B \rightarrow \text{End}_k(V)$, $b \mapsto \mu(b)$, where $\mu(b)$ is the linear map of on V such that $v \mapsto v \cdot b$; this gives a right $M_n(B)$ -module structure on V^n . By definition, $(\mu(v_{ij}))_{i,j=1}^n$ is a well-defined endomorphism of V^n as a right A -module.

Because f is a ring epimorphism, $(\mu(v_{ij}))_{i,j=1}^n$ is also an endomorphism of V^n as a right $M_n(B)$ -module. This means that $\mu(v_{ii}) - \mu(v_{jj}) = 0$, $\mu(v_{i'j'}) = 0$,

and $\mu(v_{ii})\mu(b) - \mu(b)\mu(v_{ii}) = 0$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$ and $b \in B$. Applying these maps to $1 \in V$ yields the claim. \square

The following proposition may be seen as a generalization of Proposition 19 above, and it will be used in a subsequent proof of a generalization of Theorem 20 in a similar way too.

Proposition 23. *Suppose that A is finite-dimensional algebra over k and $M \in \text{mod} - A$ of dimension n . The A -module structure on M is given by a homomorphism $f : A \rightarrow M_n(k)$; this homomorphism gives us I_f an ideal of $k * k$. $k\langle (v_{ij})_{i,j=1}^n \rangle \cong k\langle (v_{ij})_{i,j=1}^n \rangle$. If $\sum_{i,j=1}^n \lambda_{ij} m_{ij} = 0$ for all $(m_{ij})_{i,j=1}^n$ A -endomorphisms of $M_A \cong k_A^n$, via f , and $\lambda_{ij} \in k$, then $\sum_{i,j=1}^n \lambda_{ij} v_{ij} \in I_f$.*

Proof. Let $k^{(L)}$ for a set I be a vector space over k . We define the A -module structure on $(k^{(L)})^n$ using $f : A \rightarrow M_n(k)$. Actually, as an A -module, $(k^{(L)})^n \cong (k_A^n)^{(L)}$ with $k_A^n \cong M$. Since M is a finitely presented A -module, we have that:

$$\text{Hom}_A\left(\bigoplus_{\ell \in L} M_\ell, \bigoplus_{\ell \in L} M_\ell\right) \cong \prod_{\ell_1 \in L} \bigoplus_{\ell_2 \in L} \text{Hom}_A(M_{\ell_1}, M_{\ell_2})$$

Suppose we have an A -endomorphism of $(k^{(L)})^n$ given by $(M_{ij})_{i,j=1}^n$ for $M_{ij} : k^{(L)} \rightarrow k^{(L)}$. Every $M_{ij} = ((m_{ij}^{\ell_1 \ell_2})_{\ell_2 \in L})_{\ell_1 \in L}$ such that for any $\ell_1 \in L$ only finitely many $m_{ij}^{\ell_1 \ell_2}$ are non-zero and for every $\ell_1, \ell_2 \in L$, $(m_{ij}^{\ell_1 \ell_2})_{i,j=1}^n$ form an endomorphism of M as an A -module. If $\sum_{i,j=1}^n \lambda_{ij} m_{ij} = 0$ holds for all $(m_{ij})_{i,j=1}^n$ A -endomorphisms of M , clearly, $\sum_{i,j=1}^n \lambda_{ij} M_{ij} = 0$ holds as well. The claim follows by applying this observation to $k\langle (v_{ij})_{i,j=1}^n \rangle / I_f$ as an A -module and its A -endomorphism $(\mu(v_{ij}))_{i,j=1}^n$, where $(\mu(v_{ij}))_{i,j=1}^n$ is the multiplication by v_{ij} from the right. \square

Theorem 24. *Let us have a finite acyclic quiver Q' and a subquiver Q with $Q_0 = Q'_0$ and $Q_1 \cup \{e'\} = Q'_1$. Suppose there is a representation $M' = (M_v, \varphi_\alpha)_{v \in Q'_0, \alpha \in Q'_1}$ of Q' over k of dimension n such that e' is brick and $\varphi_{e'}$ is not of full rank. Consider the representation $M = (M_v, \varphi_\alpha)_{v \in Q_0, \alpha \in Q_1}$ of Q over k . Denote $M'_{s(e')}$ and $M'_{t(e')}$ such that $\text{Ker } \varphi_{e'} \oplus M'_{s(e')} = M_{s(e')}$ and $\text{Im } \varphi_{e'} \oplus M'_{t(e')} = M_{t(e')}$, respectively. We write $\varphi_{e'} : \text{Ker } \varphi_{e'} \oplus M'_{s(e')} \rightarrow \text{Im } \varphi_{e'} \oplus M'_{t(e')} = M_{t(e')}$ as follows:*

$$\varphi_{e'} = \begin{pmatrix} 0 & \varphi \\ 0 & 0 \end{pmatrix}$$

Provided that $\text{Ker } \varphi_{e'}$ and $\text{Im } \varphi_{e'}$ are invariant under all endomorphisms of M , there exists a ring epimorphism $f : kQ' \rightarrow M_n(k\langle X \rangle)$ such that on kQ this coincides with $kQ \rightarrow M_n(k)$ giving the action of kQ on M with:

$$s(e') \mapsto \begin{pmatrix} I_{s(e')} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \text{ and } t(e') \mapsto \begin{pmatrix} 0 & 0 & 0 \\ 0 & I_{t(e')} & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

where $I_{s(e')}$ is $\dim_k M_{s(e')} \times \dim_k M_{s(e')}$ identity matrix and $I_{t(e')}$ similarly, and:

$$e' \mapsto \begin{pmatrix} 0 & 0 & 0 \\ f_{e'} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The matrix $f_{e'}$ is given as follows:

(i)

$$f_{e'} = \begin{pmatrix} 0 & \varphi \\ X_{21} & 0 \end{pmatrix}$$

(ii) if moreover $M'_{s(e')}$ is invariant under all endomorphisms of M , then:

$$f_{e'} = \begin{pmatrix} X_{11} & \varphi \\ X_{21} & 0 \end{pmatrix}$$

(iii) if moreover $M'_{t(e')}$ is invariant under all endomorphisms of M , then:

$$f_{e'} = \begin{pmatrix} 0 & \varphi \\ X_{21} & X_{22} \end{pmatrix}$$

(iv) if moreover $M'_{s(e')}$ and $M'_{t(e')}$ are invariant under all endomorphisms of M , then:

$$f_{e'} = \begin{pmatrix} X_{11} & \varphi \\ X_{21} & X_{22} \end{pmatrix}$$

Where X_{ij} are matrices of indeterminates of appropriate dimensions and $X = X_{11} \cup X_{21} \cup X_{22}$.

Proof. We only prove the first assertion as the other ones follow in an analogous way.

Denote $f' : kQ' \rightarrow M_n(k)$ defining the action of kQ' on M' . By Proposition 23 above, the relation from $f'(e')$ has, without loss of generality, this form:

$$\begin{pmatrix} 0 & \varphi \\ 0 & 0 \end{pmatrix} \begin{pmatrix} s_{11} & s_{12} \\ s_{21} & s_{22} \end{pmatrix} - \begin{pmatrix} t_{11} & t_{12} \\ t_{21} & t_{22} \end{pmatrix} \begin{pmatrix} 0 & \varphi \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} \varphi s_{21} & \varphi s_{22} - t_{11} \varphi \\ 0 & -t_{21} \varphi \end{pmatrix}$$

Where s_{ij} and t_{ij} correspond to components of k -endomorphisms of $M_{s(e')} = \text{Ker } \varphi_{e'} \oplus M'_{s(e')}$ and $M_{t(e')} = \text{Im } \varphi_{e'} \oplus M'_{t(e')}$. Note that M_v for all $v \in Q'_0$ and thus all maps between them are zero.

Naturally, $I_{f'} \subseteq k\langle (v_{ij})_{i,j=1}^n \rangle$ can be seen as being contained in $I_f \subseteq k\langle X \rangle *_k k\langle (v_{ij})_{i,j=1}^n \rangle$. If $\text{Ker } \varphi_{e'}$ and $\text{Im } \varphi_{e'}$ are invariant under all endomorphisms of M , by Proposition 23, we obtain that $s_{12}, t_{12} \in I_f$ as they lie in the ideal $I_{kQ \rightarrow M_n(k)}$. SO the relation from $f(e')$ amounts to the following:

$$\begin{pmatrix} 0 & \varphi \\ X & 0 \end{pmatrix} \begin{pmatrix} s_{11} & 0 \\ s_{21} & s_{22} \end{pmatrix} - \begin{pmatrix} t_{11} & 0 \\ t_{21} & t_{22} \end{pmatrix} \begin{pmatrix} 0 & \varphi \\ X & 0 \end{pmatrix} =$$

$$= \begin{pmatrix} \varphi s_{21} & \varphi s_{22} - t_{11}\varphi \\ X_{21}s_{11} - t_{22}X_{21} & -t_{21}\varphi \end{pmatrix}$$

Since $I_{f'}$ contains $v_{ii} - v_{jj}, v_{i'j'}$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$ and I_f contains the generators of $I_{f'}$, we have that $v_{ii} - v_{jj}, v_{i'j'} \in I_f$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$. Finally, using the element $X_{21}s_{11} - t_{22}X_{21} \in I_f$, we derive that $xv_{ii} - v_{ii}x \in I_f$ for every $x \in X$ and $i = 1, \dots, n$. By Theorem 22, f is an epimorphism. \square

Remark. The theorem above is the aforementioned generalization of Theorem 20. Theorem 20, obviously without the reference to universal localisations, is a special case thereof for the brick B , whose all k -linear subspaces are invariant under all its endomorphisms as they are trivial, corresponding to M and φ_e being zero.

There is also a way how to construct ring epimorphisms from a path algebra to matrices over noncommutative polynomials by adding a vertex rather than an edge in a similar manner to Theorem 20, which can be easily proved using Theorem 22. Actually, this proposition can be also generalized in terms of constructing universal localisations as we show in the next subsection.

Proposition 25. *Let Q be a finite acyclic quiver and $M = (M_v, \varphi_\alpha)_{v \in Q_0, \varphi \in Q_1}$ a representation of Q over k of dimension n that is brick and such that there exists $w \in Q_0$ with $\dim_k M_w = m > 1$. We define a quiver Q' with $Q'_0 = Q_0 \cup \{w'\}$, $Q'_1 = Q_1 \cup \{e'\}$, $s_{Q'}(e') = w', t_{Q'}(e') = w$ and $s_{Q'}(e) = s_Q(e), t_{Q'}(e) = t_Q(e)$ for all $e \in Q_1$. Denote $f : kQ \rightarrow M_n(k)$ defining the action of kQ on M . There is a ring epimorphism $g : kQ' \rightarrow M_{n+1}(k\langle x_1, \dots, x_{m-1} \rangle)$ defined as follows:*

$$g(a) = \begin{pmatrix} f(a) & 0 \\ 0 & 0 \end{pmatrix} \text{ for } a \in kQ, g(w') = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, g(e') = \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}$$

where $X = (1 \ x_1 \dots x_{m-1} \ 0 \dots 0)^T$.

Proof. Consider the ideal I_g of $k\langle x_1, \dots, x_{m-1} \rangle * k\langle (v_{ij})_{i,j=1}^{n+1} \rangle$. Since M is brick, we already have that $v_{ii} - v_{jj}, v_{i'j'} \in I_g$ for all $i, j, i' \neq j' \in \{1, \dots, n\}$. The relation from $g(w')$ gives us that $v_{ij} = 0$ for all $i \neq j$ and $i = n+1$ or $j = n+1$.

Therefore, the relation arising from $g(e')$ reads as follows:

$$\begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix} \begin{pmatrix} v_{11} \cdot I_n & 0 \\ 0 & v_{(n+1)(n+1)} \end{pmatrix} - \begin{pmatrix} v_{11} \cdot I_n & 0 \\ 0 & v_{(n+1)(n+1)} \end{pmatrix} \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}$$

This yields that $Xv_{(n+1)(n+1)} = v_{11} \cdot I_n X$, which can be written as

$$(v_{(n+1)(n+1)} v_{(n+1)(n+1)} x_1 \dots v_{(n+1)(n+1)} x_{m-1} 0 \dots 0)^T - (v_{11} v_{11} x_1 \dots v_{11} x_{m-1} 0 \dots 0)^T \in I_g.$$

We obtain that $v_{(n+1)(n+1)} - v_{11}$, and we employ this to show that v_{ii} commute with x_j for all $i = 1, \dots, n+1$ and $j = 1, \dots, m-1$. The map g is an epimorphism by Theorem 22. \square

2.4 Extending universal localisations

In this subsection, we explore possible generalizations of Theorem 20 by extending universal localisations from smaller to larger path algebras. For this end, we make use of the matrix reduction functor, and we show that, due to their similar universal property, matrix algebras over $k\langle x_1, \dots, x_n \rangle$ arise naturally when dealing with certain universal localisations of path algebras.

Definition 26 (Matrix reduction functor, section 0.2 in [Coh06]). *There is a left adjoint to the $n \times n$ matrix functor $M_n(-) : k\text{-Alg} \rightarrow k\text{-Alg}$; it is denoted $\sqrt[n]{-} : k\text{-Alg} \rightarrow k\text{-Alg}$ and referred to as the n -matrix reduction functor.*

The adjunction gives us a functorial isomorphism $\text{Hom}_{k\text{-Alg}}(A, M_n(\sqrt[n]{B})) \cong \text{Hom}_{k\text{-Alg}}(\sqrt[n]{A}, B)$ for all $A, B \in k\text{-Alg}$. It follows that there is a map $\mu_{A,n} : A \rightarrow M_n(\sqrt[n]{A})$ such that for any $f : A \rightarrow M_n(B)$ there is a unique map $f' : \sqrt[n]{A} \rightarrow B$ with $f = M_n(f')\mu_{A,n}$. This situation can be summarized by the following commutative diagram:

$$\begin{array}{ccc} \text{Hom}_{k\text{-Alg}}(A, M_n(\sqrt[n]{A})) & \xrightarrow{\cong} & \text{Hom}_{k\text{-Alg}}(\sqrt[n]{A}, \sqrt[n]{A}) \\ \text{Hom}_{k\text{-Alg}}(A, M_n(f')) \downarrow & & \downarrow \text{Hom}_{k\text{-Alg}}(\sqrt[n]{A}, f') \\ \text{Hom}_{k\text{-Alg}}(A, M_n(B)) & \xrightarrow{\cong} & \text{Hom}_{k\text{-Alg}}(\sqrt[n]{A}, B) \end{array}$$

Actually, it is possible to describe $\sqrt[n]{A}$ and $\mu_{A,n}$ rather simply. The ring $M_n(\sqrt[n]{A})$ is isomorphic to $M_n(K) *_k A$; $\sqrt[n]{A}$ is isomorphic to the centralizer of e_{ij} in $M_n(K) *_k A$, which comprises of elements of form $\sum_v e_{iv} a e_{vj}$ for $a \in A$ (compare Theorem 0.2.3 in [Coh06]); $\mu_{A,n}(a) = \sum_{ij} e_{ij} \cdot (\sum_v e_{iv} a e_{vj})$ for all $a \in A$ and, given $f : A \rightarrow M_n(B)$, the $f' : \sqrt[n]{A} \rightarrow B$ such that $f = M_n(f')\mu_{A,n}$ is defined simply as $f'(\sum_v e_{iv} a e_{vj}) = (f(a))_{ij}$.

Lemma 27. *Suppose that $\ell_\Sigma : A \rightarrow A_\Sigma$ is the universal localisation in a set of maps between finitely generated projectives over A . Let $f : A \rightarrow B$ be a map such that there is a map $g : B \rightarrow A_\Sigma$ such that $\ell_\Sigma = gf$, then there exists a surjective map $h : B_{\Sigma \otimes_A B} \rightarrow A_\Sigma$ such that $g = h\ell_{\Sigma \otimes_A B}$, where $\Sigma \otimes_A B = \{\sigma \otimes_A B \mid \sigma \in \Sigma\}$ and $\ell_{\Sigma \otimes_A B} : B \rightarrow B_{\Sigma \otimes_A B}$ is the universal localisation of B in $\Sigma \otimes_A B$.*

Proof. Due to the universal property of the universal localisation ℓ_Σ , we have a unique map $i : A_\Sigma \rightarrow B_{\Sigma \otimes_A B}$ with the property that $\ell_{\Sigma \otimes_A B} f = i\ell_\Sigma$ as every $\sigma \otimes_A B_{\Sigma \otimes_A B}$, $\sigma \in \Sigma$, is invertible in $B_{\Sigma \otimes_A B}$. Similarly, there exists a unique map $j : B_{\Sigma \otimes_A B} \rightarrow A_\Sigma$ extending the map g since every $(\sigma \otimes_A B) \otimes A_\Sigma = \sigma \otimes_A A_\Sigma$, $\sigma \in \Sigma$, is invertible. We can calculate that $j i \ell_\Sigma = j \ell_{\Sigma \otimes_A B} f = g f = \ell_\Sigma$. By universal property of ℓ_Σ , only the identity map on A_Σ satisfies that $\ell_\Sigma = m \ell_\Sigma$ for $m : A_\Sigma \rightarrow A_\Sigma$; therefore, $j i = \text{id}_{A_\Sigma}$. This implies that j is surjective. \square

Corollary 28. *Suppose that $\ell_\Sigma : A \rightarrow A_\Sigma$ is a universal localisation in Σ such that $A_\Sigma \cong M_n(B)$ for some B and n , then A_Σ is isomorphic to a factor of $M_n(\sqrt[n]{A})_{\Sigma'}$ where $\Sigma' = \{\sigma \otimes_A M_n(\sqrt[n]{A}) \mid \sigma \in \Sigma\}$.*

Proposition 29. *Let $f : A \rightarrow M_n(B)$ be map of k -algebras, then the following are equivalent:*

- (i) *The map f is an epimorphism.*
- (ii) *The dominion of $f(A)$ in $M_n(B)$ contains all e_{ij} for $1 \leq i, j \leq n$ and the corresponding map from $\sqrt[n]{A}$ to B is an epimorphism.*
- (iii) *The map $\sqrt[n]{f} : \sqrt[n]{A} \rightarrow \sqrt[n]{M_n(B)}$ is an epimorphism.*

Proof. (i) \Leftrightarrow (ii) : Provided that $f : A \rightarrow M_n(B)$ is an epimorphism, then the dominion of $f(A)$ in $M_n(B)$ contains all of $M_n(B)$. The fact that $f : A \rightarrow M_n(B)$ is an epimorphism implies that $\text{Hom}_{k\text{-Alg}}(f, M_n(C))$ is injective for any C . However, the adjunction implies that so is $\text{Hom}_{k\text{-Alg}}(f', C)$ where f' corresponds to f under the natural isomorphism $\text{Hom}_{k\text{-Alg}}(A, M_n(B)) \cong \text{Hom}_{k\text{-Alg}}(\sqrt[n]{A}, B)$. Therefore, $f' : \sqrt[n]{A} \rightarrow B$ is an epimorphism.

Suppose that the dominion of $f(A)$ in $M_n(B)$ contains all e_{ij} for $1 \leq i, j \leq n$ and the corresponding map $f' : \sqrt[n]{A} \rightarrow B$ is an epimorphism. Let us have C and two maps $g_1, g_2 : M_n(B) \rightarrow C$ such that $g_1 f = g_2 f$. We observe that images of e_{ij} under g_1 and g_2 are equal as they lie in the dominion of $f(A)$ in $M_n(B)$ and they induce a matrix ring structure on C represented by an isomorphism $h : C \rightarrow M_n(C')$ with C' being the centralizer of all $g_1(e_{ij}) = g_2(e_{ij})$. Moreover, hg_1 and hg_2 restrict to maps g'_1, g'_2 between B and C' because B as centralizer of e_{ij} maps to the centralizer of $g_1(e_{ij})$ or $g_2(e_{ij})$ under g_1 or g_2 , respectively. In other words, $M_n(g'_1) = hg_1$ and similarly for g_2 . This allows us to employ the adjunction: $M_n(g'_1)f = M_n(g'_2)f$ means that $g'_1 f' = g'_2 f'$, but, as f' is an epimorphism, g'_1 need to equal g'_2 . It follows easily that $g_1 = g_2$.

(i) \Leftrightarrow (iii) : Let us have a homomorphism $g : M_n(B) \rightarrow C'$. There is a C a k -algebra such that $C' \cong M_n(C)$. Hence, $f : A \rightarrow M_n(B)$ is an epimorphism is equivalent to:

$$\text{Hom}_{k\text{-Alg}}(f, M_n(C)) : \text{Hom}_{k\text{-Alg}}(A, M_n(C)) \rightarrow \text{Hom}_{k\text{-Alg}}(M_n(B), M_n(C))$$

being surjective for every k -algebra C , which, by the adjunction, is equivalent to $\text{Hom}_{k\text{-Alg}}(\sqrt[n]{f}, C) : \text{Hom}_{k\text{-Alg}}(\sqrt[n]{A}, C) \rightarrow \text{Hom}_{k\text{-Alg}}(\sqrt[n]{M_n(B)}, C)$ being surjective for every k -algebra C . The last statement is equivalent to $\sqrt[n]{f} : \sqrt[n]{A} \rightarrow \sqrt[n]{M_n(B)}$ being an epimorphism. \square

Theorem 30. *Let $i : A \rightarrow A'$ be a homomorphism, and let $\ell_\Sigma : A \rightarrow A_\Sigma$ is a universal localisation such that $A_\Sigma \cong M_n(B)$. We claim that A'_Σ , with $\Sigma' = \{\sigma \otimes_A A' \mid \sigma \in \Sigma\}$ is a factor of the universal localisation of $M_n(\sqrt[n]{A'})$ in induced maps arising only from relations in representation of A_Σ as a factor of the universal localisation $M_n(\sqrt[n]{A})$ in induced maps.*

Proof. At first, we denote I the ideal of $M_n(\sqrt[n]{A})$ such that $M_n(\sqrt[n]{A})/I \cong A_\Sigma$. We claim that there exists the following commutative diagram:

$$\begin{array}{ccc}
A & \xrightarrow{f} & A' \\
\mu_{A,n} \downarrow & & \downarrow \mu_{A',n} \\
M_n(\sqrt[n]{A}) & \xrightarrow{i_1=M_n(\sqrt[n]{f})} & M_n(\sqrt[n]{A'}) \\
\ell_{\Sigma_n} \downarrow & & \downarrow \ell_{\Sigma'_n} \\
M_n(\sqrt[n]{A})_{\Sigma_n} & \xrightarrow{i_2=(i_1)_{\Sigma_n}} & M_n(\sqrt[n]{A'})_{\Sigma'_n} \\
\pi_i \downarrow & & \downarrow \pi_{I'} \\
M_n(\sqrt[n]{A})_{\Sigma_n}/I & \xrightarrow{i_3} & M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'
\end{array}$$

Where $\Sigma_n = \{\sigma \otimes_A M_n(\sqrt[n]{A}) \mid \sigma \in \Sigma\}$, $\Sigma'_n = \{\sigma \otimes_A M_n(\sqrt[n]{A'}) \mid \sigma \in \Sigma\}$ and I' is the ideal generated by images of elements of I under i_2 . Each horizontal map below i is obtained using the universal property of the vertical map on the left applied to the composition of the horizontal map above and the vertical map on the right. Moreover, all horizontal maps between the matrix algebras respect the matrix structure. In other words, they are M_n of some map between the centralizers of e_{ij} .

We will prove two statements:

- (i) The map $\pi_{I'} \circ \ell_{\Sigma'_n} \circ \mu_{A',n} : A' \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$ from the commutative diagram map is an epimorphism.
- (ii) The universal localisation $\ell_{\Sigma'} : A' \rightarrow A'_{\Sigma'}$ factors through said map $A' \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$.

Together, these two statements imply that said map $A' \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$ is the universal localisation of A' in Σ' .

The fact that the map $A \rightarrow M_n(\sqrt[n]{A})_{\Sigma}/I$ from the commutative diagram map is an epimorphism yields that e_{ij} for $1 \leq i, j \leq n$ lie in the dominion of A in $M_n(\sqrt[n]{A})_{\Sigma}/I$. Combining it with the fact that homomorphic image of a dominion of a subring lies in the dominion of the homomorphic image of the subring gives us that e_{ij} for $1 \leq i, j \leq n$ lie in the dominion of $i_3 \circ \pi_i \circ \ell_{\Sigma_n} \circ \mu_{A,n}$ in $M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$, specifically also in the dominion of image of A' . We recall that taking universal localisation or a factor by an ideal effectively commutes with forming a matrix algebra; this gives that the said map $A' \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$ corresponds, by abuse of notation, to a map $\sqrt[n]{A'} \rightarrow \sqrt[n]{A'}_{\Sigma'_n}/I'$, which is an epimorphism. We have now established that the said map $A' \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$ is an epimorphism.

Finally, we ought to show that the universal localisation $\ell_{\Sigma'} : A' \rightarrow A'_{\Sigma'}$ factors through said map $A' \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$. Consider the following com-

mutative diagram:

$$\begin{array}{ccccc}
A & \longrightarrow & M_n(\sqrt[n]{A})_{\Sigma_n} & \longrightarrow & M_n(\sqrt[n]{A})_{\Sigma_n}/I \\
\downarrow & & \downarrow & & \downarrow \text{dotted} \\
A' & \longrightarrow & M_n(\sqrt[n]{A'})_{\Sigma'_n} & \longrightarrow & A'_{\Sigma'_n}
\end{array}$$

The dotted map results from using the universal property of ℓ_{Σ} to the map $A \rightarrow M_n(\sqrt[n]{A})_{\Sigma_n} \rightarrow M_n(\sqrt[n]{A'})_{\Sigma'_n} \rightarrow A'_{\Sigma'_n}$, which is a Σ -inverting map. Commutativity of the square on the right is due to the universal property of $A \rightarrow M_n(\sqrt[n]{A})$ and to $M_n(\sqrt[n]{A}) \rightarrow M_n(\sqrt[n]{A})_{\Sigma_n}$ being an epimorphism; since $A'_{\Sigma'_n}$ can be given structure of $n \times n$ matrices over some algebra, the image of A fully determines the image of $M_n(\sqrt[n]{A})$, and also $M_n(\sqrt[n]{A})_{\Sigma_n}$, indeed. From the commutative diagram, we read that images of elements of I in $M_n(\sqrt[n]{A'})_{\Sigma'_n}$ need to map to zero, so the morphism $M_n(\sqrt[n]{A'})_{\Sigma'_n} \rightarrow A'_{\Sigma'_n}$ factors through $M_n(\sqrt[n]{A'})_{\Sigma'_n}/I'$ as I' is generated by such elements. \square

Let us have a path algebra kQ with the vertices of Q_0 labelled $1, \dots, n$ and a dimension vector $\alpha \in \mathbb{N}_0^{Q_0}$. We will construct a map $q_{Q,\alpha}$ from kQ to $M_{\alpha}(k\langle X_{Q,\alpha} \rangle)$ which is the algebra of $\sum_{i=1}^n \alpha_i \times \sum_{i=1}^n \alpha_i$ matrices over with:

$$X_{Q,\alpha} = \bigcup_{e \in Q_1} \{x_{ij}^{(e)} \mid 1 \leq i \leq \alpha_{s(e)}, 1 \leq j \leq \alpha_{s(e)}\}$$

In order to simplify working with $\sum_{i=1}^n \alpha_i \times \sum_{i=1}^n \alpha_i$ matrices over $k\langle X_{Q,\alpha} \rangle$, we think of elements of $M_{\alpha}(k\langle X_{Q,\alpha} \rangle)$ as of $n \times n$ block matrices such that the block at position ij for $1 \leq i, j \leq n$ is an $\alpha_i \times \alpha_j$ matrix.

For each $m \in Q_0$, we set $q_{Q,\alpha}(e_m)_{ij} = I_{\alpha_m}$ for $i = j = m$ and $q_{Q,\alpha}(e_m)_{ij} = 0$, otherwise, when viewed as an $n \times n$ block matrix. For each $e \in Q_1$, we set $q_{Q,\alpha}(e)_{ij} = X_{Q,\alpha}^{(e)}$ for $i = s(e), j = s(e)$ and $q_{Q,\alpha}(e)_{ij} = 0$, otherwise; $X_{Q,\alpha}^{(e)}$ is a $\alpha_{t(e)} \times \alpha_{s(e)}$ matrix with $x_{ij}^{(e)}$ at position ij .

Thus, by universal property of the path algebra kQ , this assignment can be extended to a map from $q_{Q,\alpha} : kQ \rightarrow M_{\alpha}(k\langle X_{Q,\alpha} \rangle)$. This map factors through $\mu_{A,n}$ via $q'(Q, \alpha)$, and $q'_{Q,\alpha}$ is surjective as the entries of matrices of image of $q_{Q,\alpha}$ generate $k\langle X_{Q,\alpha} \rangle$ as a k -algebra by discussion below Definition 26.

Proposition 31. *Suppose $f : kQ \rightarrow R'$ is a map and S is a module over R such that $e_i(kQ) \otimes_{kQ} R' \cong S^{\alpha_i}$ for all $1 \leq i \leq n$, then there exists R such that $R' \cong M_n(R)$. The map f viewed now as going from kQ to $M_n(R')$ uniquely factors via $M_n(f')$ through $q_{Q,\alpha}$.*

Proof. Since $kQ \cong e_1(kQ) \oplus \dots \oplus e_n(kQ)$ as a right module over itself and $kQ \otimes_{kQ} R' \cong R'$, it follows that $R' \cong S^{\alpha_1} \oplus \dots \oplus S^{\alpha_n}$. Hence, there is an isomorphism $R' \rightarrow M_{\alpha}(\text{End}_{R'}(S))$. Set $R = \text{End}_{R'}(S)$, and let us work with f as with a map from $kQ \rightarrow M_n(R)$. Without loss of generality, we can assume that $f(e_m)_{ij} = I_{\alpha_m}$ for $i = j = m$ and $f(e_m)_{ij} = 0$, otherwise. For an arrow

$e \in Q_1$ from $s(e)$ to $t(e)$, this means that $f(e)_{ij} \neq 0$ if and only if $i = t(e)$ and $j = s(e)$ and $f(e)_{(t(e))(s(e))}$ is an $\alpha_{t(e)} \times \alpha_{s(e)}$ matrix. We define $f'k\langle X_{Q,\alpha} \rangle \rightarrow R$ such that $x_{ij}^{(e)} \mapsto (f(e)_{(t(e))(s(e))})_{ij}$; we obtain $f = M_n(f') \circ q_{Q,\alpha}$. Uniqueness thereof follows from the same universal property of $k\langle X_{Q,\alpha} \rangle$. \square

Using the theorem above, we can formulate another generalization of Theorem 20, which in extending universal localisation from a smaller path algebra to a larger one.

Proposition 32. *Let Q be a finite acyclic quiver and $M \in \mathbf{mod} - kQ$ be an exceptional module of dimension vector α , then the map $f : kQ \rightarrow M_\alpha(k)$ is a universal localisation in Σ set of finitely generated maps between projectives. Let Q' be a finite acyclic quiver such that $Q_0 = Q'_0$ and arrows in Q arise from paths in Q' ; kQ is a naturally subalgebra of kQ' . Then, the universal localisation of kQ' in the set maps induced from Σ has the following form:*

$$kQ'_{\Sigma'} \cong M_\alpha(k\langle X_{Q',\alpha} \rangle)/I$$

where I is generated by $q_{Q',\alpha}(a) - \iota f(a)$ for all $a \in kQ$ viewed as a subalgebra of kQ' with ι being M_α of the map $k \rightarrow k\langle X_{Q',\alpha} \rangle$.

Proof. This is an easy consequence of Lemma 15, Theorem 30, Proposition 31, and the fact that $q_{Q',\alpha}$ factors through $\mu_{kQ',\alpha}$ via a surjective map. \square

As mentioned above, Proposition 25 can be generalized in the context of universal localisations; this idea is formalized in the following proposition.

Proposition 33. *Suppose that we have finite acyclic quivers Q_1 and Q_2 and universal localisations $f_{\Sigma_1} : kQ_1 \rightarrow M_{\alpha_1}(k\langle x_1, \dots, x_{n_1} \rangle)$ and $f_{\Sigma_2} : kQ_2 \rightarrow M_{\alpha_2}(k\langle x_1, \dots, x_{n_2} \rangle)$ for Σ_1 and Σ_2 sets of maps between finitely generated projectives over kQ_1 and kQ_2 , respectively.*

Let Q be a quiver such that $Q_0 = (Q_1)_0 \cup (Q_2)_0$, Q_1 such that for all $e \in Q'_1$ either is $s(e) \in (Q_1)_0, t(e) \in (Q_2)_0$ or $s(e) \in (Q_2)_0, t(e) \in (Q_1)_0$. Denote Σ the set of maps between finitely generated projectives that correspond to maps in Σ_1 and Σ_2 .

Then kQ_Σ is Morita equivalent to representations of a quiver Q' with two vertices v_1 and v_2 with n_1 loops for v_1 , n_2 loops for v_2 , $\alpha_1(s(e)) \cdot \alpha_2(t(e))$ arrows from v_1 to v_2 for each $e \in Q_1$ with $s(e) \in (Q_1)_0$ and $t(e) \in (Q_2)_0$, and $\alpha_2(s(e)) \cdot \alpha_1(t(e))$ arrows from v_2 to v_1 for each $e \in Q_1$ with $s(e) \in (Q_2)_0$ and $t(e) \in (Q_1)_0$.

Proof. Denote I_1 and I_2 the ideal generated by $v \in (Q_2)_0$ and $v \in (Q_1)_0$, respectively. We have that $kQ/I_j \cong kQ_j$ and that:

$$kQ_\Sigma / \langle I_j \rangle_{kQ_\Sigma} \cong (kQ_j)_{\Sigma_j} \cong M_{\alpha_j}(k\langle x_1, \dots, x_{n_j} \rangle)$$

for $j = 1, 2$. Given a representation $M = (M_v, \varphi_e)_{v \in Q_0, e \in Q_1}$ of Q over k that is a restriction of module over kQ_Σ , we find that $(M_v, \varphi_e)_{v \in (Q_2)_0, e \in (Q_2)_1}$ is a

restriction of a module over $(kQ_1)_{\Sigma_1}$ and, therefore, corresponds to a representation $(M_1, \lambda_1^M, \dots, \lambda_{n_1}^M)$ of $k\langle x_1, \dots, x_{n_1} \rangle$. Analogously, $(M_v, \varphi_e)_{v \in (Q_2)_0, e \in (Q_2)_1}$ corresponds to $(M_2, \mu_1^M, \dots, \mu_{n_2}^M)$. Also, $M_v = (M_1)^{\alpha_1(v)}$ and $M_v = (M_2)^{\alpha_2(v)}$ for $v \in (Q_1)_0$ and $v \in (Q_2)_0$, respectively.

For $e \in Q'_1$ with source in $(Q_1)_0$ and target in $(Q_2)_0$, we have that $\varphi_e : M_1^{\alpha_1(s(e))} \rightarrow M_2^{\alpha_2(t(e))}$, which means it is determined by a $\alpha_2(t(e)) \times \alpha_1(s(e))$ matrix of maps from $M_1 \rightarrow M_2$, similarly for $e' \in Q'_1$ with source in $(Q_2)_0$. Therefore, M corresponds to a representation $F(M)$ of Q' given by $F(M)_{v_1} = M_1$, $F(M)_{v_2} = M_2$, the loops are given by $\lambda_1, \dots, \lambda_{n_1}$ and μ_1, \dots, μ_{n_2} and other arrows are given by maps from M_1 to M_2 or vice versa that form components of respective $e \in Q'_1$ as discussed above.

By considering the components pertaining to Q_1 and Q_2 , we obtain that any possible homomorphism f between M and N has the following form $f = ((f_1 I_{\alpha_1(v)})_{v \in (Q_1)_0}, (f_2 I_{\alpha_2(v)})_{v \in (Q_2)_0})$ such that f_1 commutes with all λ and f_2 commutes with all μ . We also have that $f_2 I_{\alpha_2(t(e))} \cdot \varphi_e^M = \varphi_e^N \cdot f_1 I_{\alpha_1(s(e))}$ for every $e' \in Q'_1$ with source in $(Q_1)_0$ and target in $(Q_2)_0$. Similarly, it holds for $e' \in Q'_1$ from a vertex in $(Q_2)_0$ and to a vertex in $(Q_1)_0$. This gives us that f commutes with M_1 to M_2 or vice versa representing arrows in Q' . We set $F(f)_{v_1} = f_1$ and $F(f)_{v_2} = f_2$.

We have constructed a functor F from $\mathbf{Mod} - kQ_{\Sigma}$ to $\mathbf{Mod} - kQ'$. It is easy to show that the functor F is well-defined, fully faithful and essentially surjective. \square

Acknowledgments

This text was prepared as in the course of the author's work on a doctoral thesis at Faculty of Mathematics and Physics, Charles University advised by Jan Šťovíček, to whom the author wishes to thank for his help and support.

The author was partially supported from grant *GAČR 17-23112S* of the Czech Science Foundation.

References

- [ASS06] Ibrahim Assem, Andrzej Skowronski, and Daniel Simson. *Elements of the Representation Theory of Associative Algebras: Volume 1: Techniques of Representation Theory*, volume 65. Cambridge University Press, 2006.
- [Coh06] Paul Moritz Cohn. *Free ideal rings and localization in general rings*, volume 3. Cambridge University Press, 2006.
- [DF04] David Steven Dummit and Richard M Foote. *Abstract algebra*, volume 3. Wiley Hoboken, 2004.

- [GDLP87] Peter Gabriel and Jose Antonio De La Peña. Quotients of representation-finite algebras. *Communications in Algebra*, 15(1-2):279–307, 1987.
- [GL91] Werner Geigle and Helmut Lenzing. Perpendicular categories with applications to representations and sheaves. *Journal of Algebra*, 144(2):273–343, 1991.
- [HMS⁺18] Lidia Angeleri Hügel, Frederik Marks, Jan Stovicek, Ryo Takahashi, and Jorge Vitória. Flat ring epimorphisms and universal localisations of commutative rings. *Preprint arXiv:1807.01982*, 2018.
- [Isb69] John R Isbell. Epimorphisms and dominions. iv. *Journal of the London Mathematical Society*, 2(1):265–273, 1969.
- [Iya03] Osamu Iyama. Rejective subcategories of artin algebras and orders. *arXiv preprint math/0311281*, 2003.
- [KŠ10] Henning Krause and Jan Št’ovíček. The telescope conjecture for hereditary rings via ext-orthogonal pairs. *Advances in Mathematics*, 225(5):2341–2364, 2010.
- [MV15] Frederik Marks and Jorge Vitoria. From ring epimorphisms to universal localisations. *Forum Mathematicum*, 27(2):1139–1161, 2015.
- [Sch85] Aidan Harry Schofield. *Representations of rings over skew fields*, volume 92. Cambridge University Press, 1985.
- [Ste12] Bo Stenström. *Rings of Quotients: An Introduction to Methods of Ring Theory*, volume 217. Springer Science & Business Media, 2012.

CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF ALGEBRA, SOKOLOVSKA 83, 186 75 PRAHA, CZECH REPUBLIC

Email address: jakub.kopriva@outlook.com