

## On the origin of the holographic universe

Haidar Sheikahmadi

*School of Astronomy, Institute for Research in Fundamental Sciences (IPM), P. O. Box  
19395-5531, Tehran, Iran,*

*Center for Space Research, North-West University, Potchefstroom, South Africa,  
Canadian Quantum Research Center 204-3002 32 Avenue Vernon, British Columbia V1T 2L7,  
Canada*

*h.sh.ahmadi@gmail.com ; h.sheikahmadi@ipm.ir*

Received (Day Month Year)

Revised (Day Month Year)

In this work, we reexamine the holographic dark energy concept proposed already for cosmological applications. By considering, more precisely, the bounds on the entropy arising from lattice field theory on one side and Bekenstein-Hawking entropy of black holes on another side, it is shown that the so-called holographic dark energy cannot be mimicked as easily as claimed in the literature. In addition, the limits on the electron ( $g - 2$ ) experiments are taken into account again. It is shown that the corrections to the electron magnetic momentum are of the order of  $\mathcal{O}(10^{-23})$ .

**keywords:** Holography concept; Lattice Field Theory; Bekenstein-Hawking entropy; Electron gyromagnetic anomaly

**PACS:** 04.60.-m, 04.60.Nc, 04.62.+v, 04.70.Dy, 12.20.Fv

### 1. Introduction

Determining at what scales both in length and energy, and consequently in mass, the correct interpretation of the concept of quantum gravity (at least still as a gedanken experiment) has been one of the most serious studies in the last half-century <sup>1</sup>. A model that is all-encompassing and can bring all the forces and interactions of nature under one umbrella, at least, is not yet built <sup>2</sup>. In fact, the main problem goes back to the structure of the frame in which the remnant of the particles and the interactions are placed. Unlike flat space time, gravity, or in other words bend geometry, makes it possible for an active and consequently dynamic spacetime, that can interact with the rest of the contents of this framework. From the point of view of quantum mechanics, any uncertainty in the position of the particle implies an uncertainty in the momentum. In the presence of gravity and due to the gravity-energy interaction, we expect an additional uncertainty in the position of the particle. By means of a relatively straightforward and model-independent calculation, in the presence of gravity, it can be shown that the maximum distance between two events can be considered as small as the Plank length, i.e.,  $\sqrt{g}L \leq L_{Pl}$ ,

where we denote the gravitational potential by  $\bar{g}$  and a wavelength of the photon by  $\simeq L$  where  $L$  stands for the radius of a tesseract, for instance see [3](#). Now let us look at the minimum length problem by considering the vision arising from loop quantum gravity, LQG. In a non-perturbative version of the LQG, we do not need to perturb the geometry itself, and therefore the background will not directly play a role in subsequent evolutions, but we can instead look for operators that carry information about the structure of the geometry. In LQG, such quantum operators can be defined, e.g., the surface operator, and are established in such a way that at lower energies, for example in effective field theory, EFT, limits they return to the classical geometry [4,5](#). The significance of such an interpretation can be understood in Bekenstein's work on Kerr black hole [6](#). This interpretation is important, in that we need a black hole entropy for our analysis, that is proportional to the surface of the black hole, to constrain the infrared, IR, and ultraviolet, UV, cut-offs connection to eliminate the strong gravitational effects aiming at achieving a feasible EFT limit. As discussed in [7](#), henceforth CKN, this IR-UV relationship can supply a third possibility to overcome the cosmological constant problem. To interpret the consequences of such a connection, one can consider both the theoretical and experimental works in this regard, especially electron  $(g - 2)$  criterion [8,9,10](#). Basically, the  $(g - 2)$  experiment allows us to calculate the deviations that occur due to the presence of gravity. For a technical review on both the UV-IR cutoffs correlation and  $(g - 2)$  test, and to avoid prolonging the discussion, let us refer the reader to [7,10,11,12](#) and references therein. Now we turn our attention to the so called holographic dark energy model obviously as an emergent of CKN model. The concept of holography principle that raised from string theory, first was introduced in [13,14](#) and [15](#) aiming at solving shortcomings that appeared to construct a feasible solution for quantum gravity. In such a proposal, it can be supposed that the description of a special volume of space on a lower-dimensional boundary to this region is achievable [16](#). There are some outstanding examples of the applications of holography concept in high energy physics, of which the anti De Sitter/conformal field theory, AdS/CFT, correspondence [17,18](#) and entropy of black holes [6,19](#) are well-known to physics community.

Here also it deserves to note that, the importance of the relation between mass and UV cutoff from different perspectives, by taking into account for instance supersymmetry and supergravity, are addresses satisfactorily, where we refer the reader to [20,21,22,23,24,25](#) and references therein. The main motivation for doing this research is to re-examine the concept of holography, which has recently found a special place in the discussion of dark energy investigations. As will be seen, the proportion defined by the inverse of the area defined for the dark energy density will deviate. This work is organised as follows: Sec. [1](#) was a brief introduction aiming to address the problems that the ill-defined holographic dark energy model is faced. In Sec. [2](#) the claimed problems are worked out then in order to constrain the cut-offs bounds the  $(g - 2)$  criterion are utilized. Ultimately Sec. [3](#) is devoted to the discussion and conclusions of the work.

## 2. Basic Properties of Holographic Universe

With the help of the quantum field theory, for a hypothetical box with the size of  $L$  and the UV cutoff  $\Lambda$  the entropy will be an extensive quantity and reads  $L^3\Lambda^3$  [7](#). This is the result that will be obtained with the help of the normalization in the lattice field theory as well. On the other side, from the quantum states and their re-normalization near the black hole horizon, from its standard point of view, the entropy of a black hole through the relationship with the surface is closely related to length of the black hole [6,19](#). Now, based on the discussions appeared in the introduction, if we want to have an effective version of quantum field theory, say EFT, there must be a relationship between the entropy of the black hole and the state of the lattice field theory [7,14](#). This actually guarantees that we will be at lower orders of magnitude that the Planck energy is, because as mentioned we do not have an achievable version of quantum gravity and we do need to work with effective theories. Besides these, such a relationship is necessary to guarantee the Bekenstein upper bounds aiming at avoiding the breakdown of black hole formation within the EFT context based on [7,11,12,14,26,27](#), there should be a bound for the entropy of black hole as

$$L^3\Lambda^3 \lesssim (S_{BH}) \equiv \pi M_{Pl}^2 L^2, \quad (1)$$

where the holographic Bekenstein-Hawking, HBH, entropy denoted by  $S_{BH} \sim M_{Pl}^2 L^2$  and Planck mass by  $M_{Pl}$  [19](#). In  $L^3\Lambda^3$  that is the lattice field theory, LFT, entropy,  $L$  stands for IR cutoff and  $\Lambda$  for UV one, **and in large scale contexts stands for cosmological constant** [26,27,28,29](#). To obtain a definition for energy density in large scale limits, one can multiply both sides of Eq.(1) by  $\Lambda$  then readily it expresses

$$L^3\Lambda^4 \lesssim \pi M_{Pl}^2 L^2 \Lambda, \quad (2)$$

then by means of EFT techniques the energy density that corresponds to zero point quantum fluctuations, that appears as  $\Lambda^4$ , using (2) can be rewritten as

$$\rho_\Lambda = \Lambda^4 \lesssim \pi M_{Pl}^2 L^{-1} \Lambda = 3c^2 M_{Pl}^2 L^{-1} \Lambda, \quad (3)$$

where even if IR cutoff behaves like  $\Lambda^{-2}$ , according to [7,12](#), the holography concept can not be mimicked anymore, at least for a cosmological purpose. The numerical constant  $3c^2$  is introduced for convenience and it is not dimensionless as appeared usually in the literature. Here also we should emphasis that, in some works this mater were declared that the so called holographic parameter  $c^2$  can not be considered as a constant in general, see [30,31,32](#). Comparing Eq.(3) to its analogues that appeared in [7](#) and [12](#), it is not allowed to easily consider this equation as a holographic dark energy equation, that should mimic such a formula  $\Lambda^4 = \rho_\Lambda \approx \pi M_{Pl}^2 L^{-2}$ . In this regard, as discussed in CKN and CK, [7,12](#), and obviously from Eq.(1), the IR and UV cutoffs can not be treated as independent

parameters. Therefore, one cannot introduce an arbitrary scale for these cutoffs aiming at getting rid of this shortcoming, appeared due to inequality (2) here. In this regard, to reconstruct holographic dark energy model, one maybe would like to rescale the length as  $L \propto \Lambda^{-1}$ , but obviously it is forbidden based on aforementioned discussions, inequality (2) and the following explanations. These results besides Eq.(3) and (4) are important because they set a boundary on which the EFT rules can be employed properly. In other words, they determine how and where the effects of quantum gravity can be relinquished.

Now let's turn our attention to the so called electron ( $g - 2$ ) experiment. Following CKN and CK, [7,12](#), and by utilizing Eq.(1) we can obtain the constraint between IR and UV cutoffs as follows,

$$L\Lambda^3 \lesssim \pi M_{Pl}^2. \quad (4)$$

This relation indicates a correlation between the energy and the size of the box, that is the Universe, and such a coupling remains for even perturbative EFT calculations. The analogues of this equation appeared in CK paper as

$$L\Lambda^2 \lesssim M_{Pl}, \quad (5)$$

and obviously Eqs.(4) and (5) behave completely different, as it is shown, in determining the amount of bonds on the energy and the size of the box. In fact, the problem goes back to ignoring  $L\Lambda$ , by CKN and CK, on the right hand side of Eq.(2), which allows getting a square root of both sides of this equation, and therefore allowed them to claim a holographic like dark energy model for the Universe. As discussed, the relationship between the UV and IR cutoffs and how they are renormalized play an important role in investigating the thermodynamics of a black hole and especially its entropy [33,34](#). In fact, many works have been done to investigate the vacuum polarization around a black hole, all of which were to obtain a finite value for the expectation value of the energy density and its relationship with the temperature and finally the entropy of the black hole [35,36,37](#). On the other hand, in the asymptotic limits and far from the horizon of the black hole, investigations are also important for the role of the IR cutoff and its renormalization, which may even include interesting phenomena such as the cosmological constant [38](#).

In the next stage, we want to examine this model in the concordance the well known electron gyromagnetic anomaly terrestrial experiment.

### 2.1. Constraints from $g-2$ experiment

Here, following CKN and CK, we want to calculate the corrections to the usual calculations by virtue of the ( $g - 2$ ) experiment. In order to reexamine such calculations one can consider the 1-loop results for electron gyromagnetic anomaly,  $a = (g - 2)/2$ , for a lepton like electron of mass  $m$ , i.e.

$$a(L, \Lambda) = \left( \frac{\alpha}{2\pi} \right) \left( 1 - \frac{\pi^2}{mL} - \frac{m^2}{3\Lambda^2} + \dots \right), \quad (6)$$

see [7,12](#). As discussed by CK, to eliminate the effects of strong gravitational effects and consequently the sensitivity of the model, to such gravitational effects, one possible way is minimizing IR-UV corrections. In doing so by utilizing Eqs.(4) and (6) it leads to the following results for  $\Lambda$  and  $L$ ,

$$\Lambda \sim m_{eff}^{2/3} \left( \frac{M_{Pl}^2}{m} \right)^{\frac{1}{6}} GeV, \quad (7)$$

and

$$L \sim \pi m^{-1} \left( \frac{M_{Pl}^2}{\tilde{m}_{eff}} \right)^{\frac{1}{2}} GeV^{-1}, \quad (8)$$

where  $m_{eff} \equiv \kappa m$  and  $\tilde{m}_{eff} \equiv \lambda m$  are introduced for dimensional reasons and  $\kappa$  and  $\lambda$  are constants of the order of unity. Comparing the results in Eqs.(7) and (8) and their analogues in CKN or CK, one immediately observe that they are not as same as each other. By introducing dimension of  $[\kappa] \propto \mathcal{O}(MeV^{1/4})$  and  $[\lambda] \propto \mathcal{O}(MeV)$ , and considering  $m$  as electron mass, the amount of  $\Lambda$  and  $L$  immediately read

$$\Lambda \simeq 10 GeV, \quad L \simeq 200 cm. \quad (9)$$

In CK, they obtained respectively  $\Lambda \simeq 200 GeV$  and  $L \simeq 6 cm$ . Then from Eqs.(6),(7) and (8) for the minimum deviation from the Schwinger 1-loop result one gets

$$a - \frac{\alpha}{2\pi} \sim \frac{\alpha}{2\pi} \left( \pi \sqrt{\frac{\tilde{m}_{eff}}{M_{Pl}^2}} + \frac{m^{2/3}}{3\kappa^{4/3}} \sqrt[3]{\frac{m}{M_{Pl}^2}} \right) \sim \mathcal{O}(10^{-23}), \quad (10)$$

that is different comparing to the results of [7,12](#).

### 3. Concluding remarks

Our main aim of doing this study was to check the validity of the proposed holographic dark energy model that appeared in the literature and especially in CKN paper. It has been shown that, by considering precisely the bounds on the entropies arising from LFT and black holes the so-called holographic dark energy model can not be claimed anymore. It has been also figured out that, it is not arbitrary, to re-scale the UV cutoff nor like  $\propto L^{-2}$  neither  $\propto L^{-1}$  as discussed in CKN, CK or references therein. The consequences of this correction obviously affect the so-called entropy-corrected holographic dark energy [28](#), agegraphic dark energy [39](#), new agegraphic dark energy [40](#) and other emergent prototypes. This new constraint which has appeared in (4) is examined to check the order of the theoretical corrections that appeared in the electron ( $g-2$ ) experiment and it was of the order of  $\mathcal{O}(10^{-23})$ . Ultimately, in this report, a correct version of the UV-IR bound resulted in a new version for holographic-like dark energy models obtained.

*Examining the corrections that will appear for the aforementioned emergent models through this study will be the subject of future work.*

## Acknowledgments

H.S. is very grateful to the anonymous referees for their valuable comments resulted in improvement of the paper. H.S. would like also to thank H. Firouzjahi for very constructive and interesting discussions on the vacuum polarization around black holes and its renormalization. He is grateful to T. Harko for the discussions on the original draft of the work.

## References

1. C. J. Isham, “Prima facie questions in quantum gravity,” *Lect. Notes Phys.* **434**, 1-21 (1994) doi:10.1007/3-540-58339-4\_13 [arXiv:gr-qc/9310031 [gr-qc]].
2. L. J. Garay, “Quantum gravity and minimum length,” *Int. J. Mod. Phys. A* **10**, 145-166 (1995) doi:10.1142/S0217751X95000085 [arXiv:gr-qc/9403008 [gr-qc]].
3. J. A. Wheeler, in *Relativity, Groups and Topology*, eds. B. S. DeWitt and C. DeWitt (Gordon and Breach, London, 1964). Geometrodynamics and the issue of the final state.
4. C. Rovelli and L. Smolin, “Loop Space Representation of Quantum General Relativity,” *Nucl. Phys. B* **331**, 80-152 (1990) doi:10.1016/0550-3213(90)90019-A
5. A. Ashtekar, C. Rovelli and L. Smolin, “Weaving a classical geometry with quantum threads,” *Phys. Rev. Lett.* **69**, 237-240 (1992) doi:10.1103/PhysRevLett.69.237 [arXiv:hep-th/9203079 [hep-th]].
6. J. D. Bekenstein, “The quantum mass spectrum of the Kerr black hole,” *Lett. Nuovo Cim.* **11**, 467 (1974) doi:10.1007/BF02762768
7. A. G. Cohen, D. B. Kaplan and A. E. Nelson, “Effective field theory, black holes, and the cosmological constant,” *Phys. Rev. Lett.* **82**, 4971-4974 (1999) doi:10.1103/PhysRevLett.82.4971 [arXiv:hep-th/9803132 [hep-th]].
8. F. H. Combley, “(g-2) factors for muon and electron and the consequences for QED”, *Reports on Progress in Physics.* **42** (12), 1889 (1979).
9. L. Morel, Z. Yao, P. Cladé and S. Guellati-Khélifa, “Determination of the fine-structure constant with an accuracy of 81 parts per trillion,” *Nature* **588**, no.7836, 61-65 (2020) doi:10.1038/s41586-020-2964-7
10. T. Aoyama, N. Asmussen, M. Benayoun, J. Bijnens, T. Blum, M. Bruno, I. Caprini, C. M. Carloni Calame, M. Cè and G. Colangelo, *et al.* “The anomalous magnetic moment of the muon in the Standard Model,” *Phys. Rept.* **887**, 1-166 (2020) doi:10.1016/j.physrep.2020.07.006 [arXiv:2006.04822 [hep-ph]].
11. T. Banks and P. Draper, “Remarks on the Cohen-Kaplan-Nelson bound,” *Phys. Rev. D* **101**, no.12, 126010 (2020) doi:10.1103/PhysRevD.101.126010 [arXiv:1911.05778 [hep-th]].
12. A. G. Cohen and D. B. Kaplan, “Gravitational contributions to the electron  $g$ -factor,” [arXiv:2103.04509 [hep-ph]].
13. G. 't Hooft, “Dimensional reduction in quantum gravity,” *Conf. Proc. C* **930308**, 284-296 (1993) [arXiv:gr-qc/9310026 [gr-qc]].
14. G. 't Hooft, “The Holographic principle: Opening lecture,” *Subnucl. Ser.* **37**, 72-100 (2001) doi:10.1142/9789812811585\_0005 [arXiv:hep-th/0003004 [hep-th]].
15. L. Susskind, “The World as a hologram,” *J. Math. Phys.* **36**, 6377-6396 (1995) doi:10.1063/1.531249 [arXiv:hep-th/9409089 [hep-th]].
16. R. Bousso, “The Holographic principle,” *Rev. Mod. Phys.* **74**, 825-874 (2002) doi:10.1103/RevModPhys.74.825 [arXiv:hep-th/0203101 [hep-th]].

17. J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231-252 (1998) doi:10.1023/A:1026654312961 [arXiv:hep-th/9711200 [hep-th]].
18. H. Lin, O. Lunin and J. M. Maldacena, “Bubbling AdS space and 1/2 BPS geometries,” *JHEP* **10**, 025 (2004) doi:10.1088/1126-6708/2004/10/025 [arXiv:hep-th/0409174 [hep-th]].
19. J. D. Bekenstein, “Black holes and entropy,” *Phys. Rev. D* **7**, 2333-2346 (1973) doi:10.1103/PhysRevD.7.2333
20. S. M. Carroll, “Quintessence and the rest of the world,” *Phys. Rev. Lett.* **81**, 3067-3070 (1998) doi:10.1103/PhysRevLett.81.3067 [arXiv:astro-ph/9806099 [astro-ph]].
21. B. Bellazzini, F. Riva, J. Serra and F. Sgarlata, “Massive Higher Spins: Effective Theory and Consistency,” *JHEP* **10**, 189 (2019) doi:10.1007/JHEP10(2019)189 [arXiv:1903.08664 [hep-th]].
22. R. A. El-Nabulsi, “Phase transitions in the early universe with negatively induced supergravity cosmological constant,” *Chin. Phys. Lett.* **23**, 1124-1127 (2006) doi:10.1088/0256-307X/23/5/017
23. M. Kawasaki and T. Takesako, “Hubble Induced Mass in Radiation Dominated Universe,” *Phys. Lett. B* **711**, 173-177 (2012) doi:10.1016/j.physletb.2012.03.069 [arXiv:1112.5823 [hep-ph]].
24. R. A. El-Nabulsi, “Effective cosmological constant from supergravity arguments and non-minimal coupling,” *Phys. Lett. B* **619**, 26-29 (2005) doi:10.1016/j.physletb.2005.06.002
25. R. Kallosh, A. D. Linde, S. Prokushkin and M. Shmakova, “Supergravity, dark energy and the fate of the universe,” *Phys. Rev. D* **66**, 123503 (2002) doi:10.1103/PhysRevD.66.123503 [arXiv:hep-th/0208156 [hep-th]].
26. B. Guberina, R. Horvat and H. Nikolic, “Nonsaturated Holographic Dark Energy,” *JCAP* **01**, 012 (2007) doi:10.1088/1475-7516/2007/01/012 [arXiv:astro-ph/0611299 [astro-ph]].
27. G. Lopes Cardoso, B. de Wit and T. Mohaupt, “Corrections to macroscopic supersymmetric black hole entropy,” *Phys. Lett. B* **451**, 309-316 (1999) doi:10.1016/S0370-2693(99)00227-0 [arXiv:hep-th/9812082 [hep-th]].
28. H. Wei, “Entropy-Corrected Holographic Dark Energy,” *Commun. Theor. Phys.* **52**, 743-749 (2009) doi:10.1088/0253-6102/52/4/35 [arXiv:0902.0129 [gr-qc]].
29. H. Sheikahmadi, A. Aghamohammadi and K. Saaidi, “The effect of de Sitter like background on increasing the zero point budget of dark energy,” *Adv. High Energy Phys.* **2016**, 2594189 (2016) doi:10.1155/2016/2594189 [arXiv:1407.0125 [gr-qc]].
30. N. Radicella and D. Pavon, “On the  $c^2$  term in the holographic formula for dark energy,” *JCAP* **10**, 005 (2010) doi:10.1088/1475-7516/2010/10/005 [arXiv:1007.4129 [gr-qc]].
31. R. A. El-Nabulsi, “Maxwell brane cosmology with higher-order string curvature corrections, a nonminimally coupled scalar field, dark matter-dark energy interaction and a varying speed of light,” *Int. J. Mod. Phys. D* **18**, 289-318 (2009) doi:10.1142/S0218271809014431
32. A. Sheykhi, S. Ghaffari and N. Roshanshah, “A note on holographic dark energy with varying  $c^2$  term,” *Int. J. Theor. Phys.* **56**, no.6, 1845-1860 (2017) doi:10.1007/s10773-017-3329-3 [arXiv:1612.03040 [physics.gen-ph]].
33. P. C. W. Davies, “Quantum Field Theory in Curved Space-Time,” *Nature* **263**, 377-380 (1976) doi:10.1038/263377a0
34. T. S. Bunch and P. C. W. Davies, “Quantum Field Theory in de Sitter Space: Renormalization by Point Splitting,” *Proc. Roy. Soc. Lond. A* **360**, 117-134 (1978)

- doi:10.1098/rspa.1978.0060
35. S. M. Christensen, "Vacuum Expectation Value of the Stress Tensor in an Arbitrary Curved Background: The Covariant Point Separation Method," *Phys. Rev. D* **14**, 2490-2501 (1976) doi:10.1103/PhysRevD.14.2490
  36. P. Candelas, "Vacuum Polarization in Schwarzschild Space-Time," *Phys. Rev. D* **21**, 2185-2202 (1980) doi:10.1103/PhysRevD.21.2185
  37. P. Candelas and K. W. Howard, "VACUUM ( $\Phi^2$ ) IN SCHWARZSCHILD SPACE-TIME," *Phys. Rev. D* **29**, 1618-1625 (1984) doi:10.1103/PhysRevD.29.1618
  38. H. Firouzjahi, "Cosmological constant and vacuum zero point energy in black hole backgrounds," *Phys. Rev. D* **106**, no.4, 045015 (2022) doi:10.1103/PhysRevD.106.045015 [arXiv:2205.06561 [gr-qc]].
  39. R. G. Cai, "A Dark Energy Model Characterized by the Age of the Universe," *Phys. Lett. B* **657**, 228-231 (2007) doi:10.1016/j.physletb.2007.09.061 [arXiv:0707.4049 [hep-th]].
  40. H. Wei and R. G. Cai, "Interacting Agegraphic Dark Energy," *Eur. Phys. J. C* **59**, 99-105 (2009) doi:10.1140/epjc/s10052-008-0799-8 [arXiv:0707.4052 [hep-th]].