

Compact groups with probabilistically central monothetic subgroups

João Azevedo and Pavel Shumyatsky

ABSTRACT. If K is a closed subgroup of a compact group G , the probability that randomly chosen pair of elements from K and G commute is denoted by $Pr(K, G)$. Say that a subgroup $K \leq G$ is ϵ -central in G if $Pr(\langle g \rangle, G) \geq \epsilon$ for any g in K . Here $\langle g \rangle$ denotes the monothetic subgroup generated by $g \in G$. Our main result is that if K is ϵ -central in G , then there is an ϵ -bounded number e and a normal subgroup $T \leq G$ such that the index $[G : T]$ and the order of the commutator subgroup $[K^e, T]$ both are finite and ϵ -bounded. In particular, if G is a compact group for which there is $\epsilon > 0$ such that $Pr(\langle g \rangle, G) \geq \epsilon$ for any $g \in G$, then there is an ϵ -bounded number e and a normal subgroup T such that the index $[G : T]$ and the order of $[G^e, T]$ both are finite and ϵ -bounded.

1. Introduction

In this paper, all compact groups are Hausdorff topological spaces. By a subgroup of a topological group we mean a closed subgroup unless explicitly stated otherwise. If S is a subset of a topological group G , then we denote by $\langle S \rangle$ the subgroup (topologically) generated by S . A subgroup of G is monothetic if it is generated by a single element.

The Borel σ -algebra \mathcal{M} of a compact group G is the one generated by all closed subsets of G . We say that a measure on (G, \mathcal{M}) is a (left) Haar measure provided μ is both inner and outer regular, $\mu(K) < \infty$ and $\mu(xE) = \mu(E)$ for all compact subsets K and measurable subsets E of G (see [14, Chapter 4] or [22, Chapter II]). Recall that there is a unique Haar measure μ on (G, \mathcal{M}) such that $\mu(G) = 1$.

2020 *Mathematics Subject Classification.* 20F24, 20P05, 22C05.

Key words and phrases. Commuting probability, Haar measure, compact groups, monothetic subgroups.

Supported by CNPq and FAPDF.

Let G be a compact group and let K be a subgroup of G . Consider the set $C = \{(x, y) \in K \times G \mid xy = yx\}$. This is closed in $K \times G$ since it is the preimage of 1 under the continuous map $f : K \times G \rightarrow G$ given by $f(x, y) = [x, y]$. Denoting the normalized Haar measures of K and G by ν and μ , respectively, the probability that a random element from K commutes with a random element from G is defined as $Pr(K, G) = (\nu \times \mu)(C)$. This is a well-studied concept (see in particular [5, 6, 10, 12, 15, 18, 19, 23, 28]).

Recently Detomi and the second author proved in [3] that if G is finite and $Pr(K, G) \geq \epsilon$ for some $\epsilon > 0$, then there is a normal subgroup T of G and a subgroup B of K such that the indices $[G : T]$ and $[K : B]$ and the order of the subgroup $[T, B]$ are ϵ -bounded. Throughout the article we use the expression “ (a, b, \dots) -bounded” to mean that a quantity is bounded from above by a number depending only on the parameters a, b, \dots . If B and T are subgroups of a group G , we denote by $[T, B]$ the subgroup generated by all commutators $[t, b]$ with $t \in T$ and $b \in B$. In the case where $K = G$, this is a well-known theorem due to P.M. Neumann [25]. Conversely, if K is a subgroup of a finite group G , and if $T \leq G$ and $B \leq K$, then $Pr(K, G)$ is bounded away from zero in terms of the indices $[G : T]$ and $[K : B]$ and the order of $[T, B]$. The present work grew out of a desire to understand the impact of similar probabilistic considerations on the structure of a compact group. Our first goal is to extend the main result of [3] to compact groups.

PROPOSITION 1.1. *Let $\epsilon > 0$ and let G be a compact group having a subgroup K such that $Pr(K, G) \geq \epsilon$. Then there is a normal subgroup $T \leq G$ and a subgroup $B \leq K$ such that the indices $[G : T]$ and $[K : B]$ and the order of $[T, B]$ are ϵ -bounded.*

Let G be a compact group and let K be a subgroup of G . If $Pr(\langle g \rangle, G) \geq \epsilon$ for every $g \in K$ we say that K is ϵ -central in G , and in the case where $K = G$ we say that G is ϵ -central. If e is a positive integer, we denote by G^e the subgroup generated by all e th powers of elements of G . Recall that a group G has finite exponent e if $G^e = 1$ and e is the least positive number with this property. It is easy to see that if G has exponent e , then $Pr(\langle g \rangle, G) \geq \frac{1}{e}$ for all $g \in G$. More generally, if $G^e \leq Z(G)$, where $Z(G)$ denotes the center of G , then $Pr(\langle g \rangle, G) \geq \frac{1}{e}$ for all $g \in G$. We shall prove the following theorem.

THEOREM 1.2. *Let $\epsilon > 0$ and assume that the subgroup K is ϵ -central in G . Then there is an ϵ -bounded number e and a finite-index normal subgroup $T \leq G$ such that the index $[G : T]$ and the order of $[K^e, T]$ are ϵ -bounded.*

The theorem implies that if G is an ϵ -central compact group, then there is an ϵ -bounded number e and a normal subgroup T such that the index $[G : T]$ and the order of $[G^e, T]$ are ϵ -bounded. Moreover, the exponent of the commutator subgroup $[T, T]$ is ϵ -bounded. Indeed, passing to the quotient over $[G^e, T]$ we can assume that all e th powers of elements of G centralize T . In particular, $T/Z(T)$ has exponent e , and a theorem of Zelmanov [34] ensures that $T/Z(T)$ is locally finite. A result of Mann [20] can then be used to deduce that $[T, T]$ has finite e -bounded exponent.

As usual, we denote the conjugacy class of an element $x \in G$ by x^G . It is easy to see that if K is a subgroup of G such that $|x^G| \leq n$ for every $x \in K$, then K is $\frac{1}{n}$ -central in G . More generally, let l, n be positive integers and suppose that K is a subgroup of a compact group G such that any conjugacy class containing an l th power x^l of an element $x \in K$ is of size at most n . It is not difficult to see that K is $\frac{1}{ln}$ -central in G . It turns out that this admits a converse: if K is ϵ -central in G , then there exist ϵ -bounded integers l and n such that every conjugacy class containing an l th power of an element of K has cardinality at most n . Indeed, let $x \in K$. Since $\text{Pr}(\langle x \rangle, G) \geq \epsilon$, in view of Proposition 1.1 there is a normal subgroup T of G and a subgroup B of $\langle x \rangle$ such that the indices $[G : T]$ and $[\langle x \rangle : B]$ and the order of $[T, B]$ are ϵ -bounded. Hence, as required, there are ϵ -bounded numbers l and n such that $[G : C_G(x^l)] \leq n$ for all $x \in K$. Therefore we have proved that

For every $0 < \epsilon \leq 1$ there are positive integers l and n depending only on ϵ with the property that if K is an ϵ -central subgroup of the compact group G , then $[G : C_G(g^l)] \leq n$ for all $g \in K$.

Taking this into consideration, Theorem 1.2 will follow from the next proposition.

PROPOSITION 1.3. *Let G be a compact group and let l, n be positive integers. Suppose that there is a subgroup K of G such that $[G : C_G(g^l)] \leq n$ for every $g \in K$. Then there exist a positive integer e , depending only on l and n , and a normal subgroup T of G , such that the index $[G : T]$ and the order of $[K^e, T]$ are (l, n) -bounded.*

Recall that a group is said to be a BFC-group if its conjugacy classes are finite and have bounded size. A famous theorem of B. H. Neumann says that in a BFC-group the commutator subgroup G' is finite [24]. It follows that if $|x^G| \leq n$ for every $x \in G$, then the order of G' is bounded by a number depending only on n . A first explicit bound for the order of G' was found by J. Wiegold [31], and the best known was

obtained in [9] (see also [26] and [30]). Proposition 1.3 is an extension of the Neumann theorem (in the particular case where $K = G$ and $l = 1$ the proof shows that we can take $T = G$ and $e = 1$). Recently, some other generalisations of Neumann's theorem have been obtained (see in particular [1, 2, 4]).

We end this introduction with the remark that Theorem 1.1 admits a converse:

For any positive integers s, e, m there is $0 < \epsilon \leq 1$ depending only on s, e, m with the property that if K is a subgroup of a compact group G and if G has a normal subgroup T of index at most s such that $[K^e, T]$ has order at most m , then K is ϵ -central in G .

To see this simply note that if K is as above, then $[G : C_G(g^e)] \leq ms$ for every $g \in K$.

2. Preliminaries

In this section we record some results needed in the proofs of the main theorems. The following lemma is [28, Lemma 3.1].

LEMMA 2.1. *Let G be a compact group and let K be a subgroup of G . Then either $\mu(K) = 0$ or $\mu(K) > 0$ and K is open on G . Furthermore, in the latter case, $\mu(K) = [G : K]^{-1}$.*

LEMMA 2.2. *Let G be a compact group, and let K and H be subgroups of G with $K \leq H$. Assume further that $\mu(K) \geq \epsilon \mu(H) > 0$ for some positive ϵ . Then $[H : K] \leq \epsilon^{-1}$.*

PROOF. Since $\mu(K), \mu(H) > 0$, the previous lemma implies that both subgroups are of finite index and $\mu(K) = [G : K]^{-1}$ and $\mu(H) = [G : H]^{-1}$. Hence the result. \square

For every $x \in G$, the centralizer $C_G(x)$ equals $f_x^{-1}(1)$, where f_x is the continuous function $f_x(y) = [x, y]$, so this subgroup is closed and measurable.

LEMMA 2.3. *Let H and K be subgroups of a compact group G , with $H \leq K$. Then*

$$Pr(K, G) \leq Pr(H, G) \leq Pr(H, K).$$

In particular, $Pr(G, G) \leq Pr(K, G) \leq Pr(K, K)$.

PROOF. Let μ, ν and λ be the normalized Haar measures of G, K and H , respectively. Given $x \in G$, the map $\alpha : \{hC_H(x) \mid h \in H\} \rightarrow$

$\{kC_K(x) \mid k \in K\}$ taking $hC_H(x)$ to $hC_K(x)$ is injective. We deduce that $[H : C_H(x)] \leq [K : C_K(x)]$ and $\nu(C_K(x)) \leq \lambda(C_H(x))$. We have

$$Pr(H, G) = \int_G \lambda(C_H(x)) d\mu(x) \geq \int_G \nu(C_K(x)) d\mu(x) = Pr(K, G).$$

The other inequality is proved in an analogous way. \square

Suppose that G is a compact group and let N be a normal subgroup of G . The normalized Haar measure on G/N coincides with the one induced by the normalized Haar measure on G . If A is a measurable subset of G , we denote by χ_A the characteristic function of A . We say that $x \in G$ is an FC-element if the conjugacy class of x in G is finite.

In the case of finite groups the next lemma was established in [3].

LEMMA 2.4. *Let G be a compact group and let N be a normal subgroup of G . For any subgroup K of G , we have*

$$Pr(K, G) \leq Pr(KN/N, G/N) Pr(K \cap N, N).$$

PROOF. If X is a compact group, we denote by μ_X the normalized Haar measure of X . We have

$$Pr(K, G) = \int_K \mu_G(C_G(x)) d\mu_K(x).$$

If $x \in G$ is an FC-element, then

$$\mu_G(C_G(x)N) = [C_G(x)N : C_G(x)] \mu_G(C_G(x)) = [N : C_N(x)] \mu_G(C_G(x)),$$

so $\mu_G(C_G(x)N) \mu_N(C_N(x)) = \mu_G(C_G(x))$. Let $FC(K)$ be the abstract subgroup of K consisting of elements having finite conjugacy class in G . Then

$$\begin{aligned} \int_K \mu_G(C_G(x)) d\mu_K(x) &= \int_{FC(K)} \mu_G(C_G(x)) d\mu_K(x) \\ &= \int_{FC(K)} \mu_G(C_G(x)N) \mu_N(C_N(x)) d\mu_K(x) \\ &\leq \int_K \mu_G(C_G(x)N) \mu_N(C_N(x)) d\mu_K(x). \end{aligned}$$

We now apply the extended Weil formula [27, p.88] to the last integral and obtain

$$\begin{aligned}
Pr(K, G) &\leq \\
&\int_{\frac{K}{K \cap N}} \left(\int_{K \cap N} \mu_G(C_G(xk)N) \mu_N(C_N(xk)) d\mu_{K \cap N}(k) \right) d\mu_{\frac{K}{K \cap N}}(x(K \cap N)) \\
&\leq \int_{\frac{K}{K \cap N}} \left(\int_{K \cap N} \mu_{\frac{G}{N}}(C_{\frac{G}{N}}(xN)) \mu_N(C_N(xk)) d\mu_{K \cap N}(k) \right) d\mu_{\frac{K}{K \cap N}}(x(K \cap N)) \\
&= \int_{\frac{K}{K \cap N}} \mu_{\frac{G}{N}}(C_{\frac{G}{N}}(xN)) \left(\int_{K \cap N} \mu_N(C_N(xk)) d\mu_{K \cap N}(k) \right) d\mu_{\frac{K}{K \cap N}}(x(K \cap N)). \quad (1)
\end{aligned}$$

If x is any element of K , define the set

$$\begin{aligned}
A_x &= \{(k, n) \in (K \cap N) \times N \mid [xk, n] = 1\} \\
&= \{(k, n) \in (K \cap N) \times N \mid xk \in C_G(n) \cap x(K \cap N)\}.
\end{aligned}$$

If $C_G(n) \cap x(K \cap N)$ is nonempty, then it equals $tC_{K \cap N}(k)$ for some $t \in x(K \cap N)$. Thus,

$$\begin{aligned}
A_x &= \{(k, n) \in (K \cap N) \times N \mid xk \in tC_{K \cap N}(k)\} \\
&= \{(k, n) \in (K \cap N) \times N \mid k \in x^{-1}tC_{K \cap N}(k)\}.
\end{aligned}$$

We use the Lebesgue-Fubini Theorem to give an estimate for the expression in parenthesis in (1):

$$\begin{aligned}
\int_{K \cap N} \mu_N(C_N(xk)) d\mu_{K \cap N}(k) &= \int_{(K \cap N) \times N} \chi_{A_x}(k, n) d(\mu_{K \cap N} \times \mu_N)(k, n) \\
&\leq \int_N \mu_{K \cap N}(x^{-1}tC_{K \cap N}(n)) d\mu_N(n) \\
&= \int_N \mu_{K \cap N}(C_{K \cap N}(n)) d\mu_N(n) \\
&= Pr(K \cap N, N).
\end{aligned}$$

Replacing this back in (1) we have

$$\begin{aligned} Pr(K, G) &\leq \int_{\frac{K}{K \cap N}} \mu_{\frac{G}{N}}(C_{\frac{G}{N}}(xN)) \left(\int_{K \cap N} \mu_N(C_N(xn)) d\mu_{K \cap N}(n) \right) d\mu_{\frac{K}{K \cap N}}(x(K \cap N)) \\ &\leq Pr(K \cap N, N) \int_{\frac{K}{K \cap N}} \mu_{\frac{G}{N}}(C_{\frac{G}{N}}(xN)) d\mu_{\frac{K}{K \cap N}}(x(K \cap N)). \end{aligned}$$

Finally, since $K/K \cap N$ and KN/N are isomorphic, we can apply Corollary 2.5 in [28] with respect to the last integral above to conclude that

$$\begin{aligned} \int_{\frac{K}{K \cap N}} \mu_{\frac{G}{N}}(C_{\frac{G}{N}}(xN)) d\mu_{\frac{K}{K \cap N}}(x(K \cap N)) &= \int_{\frac{KN}{N}} \mu_{\frac{G}{N}}(C_{\frac{G}{N}}(xN)) d\mu_{\frac{KN}{N}}(xN) \\ &= Pr(KN/N, G/N). \end{aligned}$$

The lemma follows. \square

If A and B are normal subgroups of a group G such that $[A : C_A(B)] \leq m$ and $[B : C_B(A)] \leq m$, then $[A, B]$ has m -bounded order. This well-known result is due to Baer, cf. [29, 14.5.2]. We need a variation of it, which is Lemma 2.1 in [3].

LEMMA 2.5. *Let $m \geq 1$ and let G be a group containing a normal subgroup A and a subgroup B such that $[A : C_A(y)] \leq m$ and $[B : C_B(x)] \leq m$ for all $x \in A, y \in B$. Assume further that $\langle B^G \rangle$ is abelian. Then $[A, B]$ has finite m -bounded order.*

The next theorem holds in any group and plays a key role in the proof of Theorem 1.1. It is taken from [1].

THEOREM 2.6. *Let m be a positive integer, G a group having a subgroup K such that $|x^G| \leq m$ for each $x \in K$, and let $H = \langle K^G \rangle$. Then the order of the commutator subgroup $[H, H]$ is finite and m -bounded.*

The next lemma is essentially Lemma 2.1 in [5].

LEMMA 2.7. *Let G be a compact group with normalized Haar measure μ and let $r \geq 1$. Suppose that X is a closed symmetric subset of G containing the identity. If $\mu(X) > \frac{1}{r+1}$, then $\langle X \rangle = X^{3r}$.*

PROOF. Suppose $x_i \in X^{3i+1} \setminus X^{3i}$ for $i = 0, \dots, r$. Then for each i , as long as $X^{3i+1} \setminus X^{3i}$ is nonempty, we have

$$x_i X \subseteq X^{3i+2} \setminus X^{3i-1}.$$

So, assume that the sets $X^{3i+1} \setminus X^{3i}$ are nonempty for $i = 0, \dots, r$. Then x_0X, \dots, x_rX are disjoint subsets of G , each of measure $\mu(X)$, and

$$\mu \left(\bigcup_{i=0}^r x_iX \right) = (r+1)\mu(X) > 1.$$

Therefore $X^{3i+1} = X^{3i}$ for some $i \leq r$. In particular, $X^{3r} = \langle X \rangle$. \square

In a compact group G the set $\{x \in G \mid |x^G| \leq n\}$ is closed and thus measurable.

LEMMA 2.8. *Let G be a compact group and n a positive integer. The set $X = \{x \in G \mid |x^G| \leq n\}$ is a closed subset of G .*

PROOF. It is sufficient to show that if $a \in G \setminus X$, then a is contained in an open subset U which has empty intersection with X . Since $a \notin X$, we can choose $n+1$ elements x_1, \dots, x_{n+1} in such a way that the conjugates a^{x_i} are distinct for $i = 1, \dots, n+1$. Set

$$U = \{u \in G; [u, x_i x_j^{-1}] \neq 1 \text{ for } 1 \leq i, j \leq n+1\}.$$

Observe that $a \in U$ and every element in U has at least $n+1$ conjugates, whence $U \cap X = \emptyset$. Further, since the commutator map is continuous, U is open. The proof is complete. \square

REMARK 2.9. If S is any subset of G , we denote by x^S the set of S -conjugates of x in G . The previous argument also proves that the set $\{x \in G \mid |x^S| \leq n\}$ is closed in G for any (not necessarily closed) subset S of G .

3. Proof of Proposition 1.1

Now we are able to prove Proposition 1.1, which we restate here for the reader's convenience.

Let $\epsilon > 0$ and let G be a compact group having a subgroup K such that $\Pr(K, G) \geq \epsilon$. Then there is a normal subgroup $T \leq G$ and a subgroup $B \leq K$ such that the indices $[G : T]$ and $[K : B]$ and the order of $[T, B]$ are ϵ -bounded.

PROOF. Let μ and ν be the normalized Haar measures of G and K , respectively. Set

$$X = \{x \in K \mid |x^G| \leq 2/\epsilon\}.$$

Note that X is measurable, by Lemma 2.8: it is the intersection of K and the closed set $\{x \in G \mid |x^G| \leq 2/\epsilon\}$. We have

$$K \setminus X = \{x \in K \mid |x^G| > 2/\epsilon\}.$$

Since $\mu(C_G(x)) < \epsilon/2$ for all $x \in K \setminus X$, it follows that

$$\begin{aligned}
\epsilon &\leq Pr(K, G) = \int_K \mu(C_G(x)) d\nu(x) \\
&= \int_X \mu(C_G(x)) d\nu(x) + \int_{K \setminus X} \mu(C_G(x)) d\nu(x) \\
&\leq \int_X d\nu(x) + \int_{K \setminus X} \frac{\epsilon}{2} d\nu(x) \\
&= \nu(X) + \frac{\epsilon}{2}(1 - \nu(X)) \leq \nu(X) + \frac{\epsilon}{2}.
\end{aligned}$$

This implies that $\epsilon/2 \leq \nu(X)$. Let B be the subgroup generated by X . Then, by Lemma 2.7, every element of B is a product of at most $6/\epsilon$ elements of X . Clearly, $\nu(B) \geq \nu(X) \geq \epsilon/2$, so the index of B in K is at most $2/\epsilon$, by Lemma 2.2. Furthermore, $|b^G| \leq (2/\epsilon)^{6/\epsilon}$ for every $b \in B$.

Let $L = \langle B^G \rangle$. Theorem 2.6 tells us that the commutator subgroup $[L, L]$ has finite ϵ -bounded order. Let us use the bar notation for images of subgroups of G in $G/[L, L]$. By Lemma 2.4, $Pr(\overline{K}, \overline{G}) \geq Pr(K, G) \geq \epsilon$. Moreover, $[\overline{K} : \overline{B}] \leq [K : B] \leq \epsilon/2$ and $|\overline{b}^{\overline{G}}| \leq |b^G| \leq (2/\epsilon)^{6/\epsilon}$ for any $b \in B$. Thus we can pass to the quotient over $[L, L]$ and assume that L is abelian.

Now we set

$$Y = \{y \in G \mid |y^K| \leq 2/\epsilon\}.$$

Observe that Y is closed, by Remark 2.9. Arguing as before, since $\nu(C_K(y)) < \epsilon/2$ for all $y \in G \setminus Y$, we have

$$\begin{aligned}
\epsilon &\leq Pr(K, G) = \int_G \nu(C_K(y)) d\mu(y) \\
&= \int_Y \nu(C_K(y)) d\mu(y) + \int_{G \setminus Y} \nu(C_K(y)) d\mu(y) \\
&\leq \int_Y d\mu(y) + \int_{G \setminus Y} \epsilon/2 d\mu(y) \\
&= \mu(Y) + \epsilon/2(1 - \mu(Y)) \leq \mu(Y) + \epsilon/2.
\end{aligned}$$

Therefore, $\mu(Y) \geq \epsilon/2$. Let E be the subgroup generated by Y . Lemma 2.7 ensures that every element of E is a product of at most $6/\epsilon$ elements of Y . Also, we have $\mu(E) \geq \mu(Y) \geq \epsilon/2$, so the index of E in G is at most $2/\epsilon$, by Lemma 2.2. Since $|y^K| \leq 2/\epsilon$ for every $y \in Y$, it follows

that $|g^K| \leq (2/\epsilon)^{6/\epsilon}$ for every $g \in E$. Let T be the maximal normal subgroup of G contained in E . Then the index $[G : T]$ is ϵ -bounded. Moreover, $|b^G| \leq (2/\epsilon)^{6/\epsilon}$ for every $b \in B$ and $|g^K| \leq (2/\epsilon)^{6/\epsilon}$ for every $g \in T$. As L is abelian, we can apply Lemma 2.5 and deduce that $[T, B]$ has finite ϵ -bounded order. The proposition follows. \square

4. About ϵ -central subgroups

Let G be a topological group generated by a symmetric set X . If it is possible to write $g \in G$ as a product of finitely many elements from X , we denote by $w(g)$ the shortest length of such an expression. If g cannot be written as a product of finitely many elements of X , we simply say that $w(g)$ is infinite. The next result is Lemma 2.1 in [4].

LEMMA 4.1. *Let G be a group generated by a symmetric set X and let D be a subgroup of index m in G . Then every coset Db contains an element such that $w(g) \leq m - 1$.*

We remark that Lemma 4.1 holds for topological groups and their closed (open) subgroups of finite index. Indeed, for an integer $r \geq 0$ let D_r be the union of the cosets of D containing some element g with $w(g) \leq r$. Then $D_r \subseteq D_{r+1}$ and $D_r X \subseteq D_{r+1}$ for all r . Let R be the minimal integer such that $D_{R+1} = D_R$. Then D_R is a closed set containing the group generated by X , so $D_R = G$. Since D has m cosets and $D_0 = D$, $R < m$.

We now proceed to the proof of Proposition 1.3, which we restate here for the reader's convenience.

Let G be a compact group and let l, n be positive integers. Suppose that there is a subgroup K of G such that $[G : C_G(g^l)] \leq n$ for every $g \in K$. Then there exists a positive integer e , depending only on l and n , and a normal subgroup T of G , such that the index $[G : T]$ and the order of $[K^e, T]$ both are finite and (l, n) -bounded.

Let G be a compact group satisfying the hypothesis of Proposition 1.3. Let X be the union of the conjugacy classes of G containing an l th power of an element of K and let H be the subgroup generated by X . Define m as the maximum of the indices of $C_H(x)$ in H , where $x \in X$. Obviously, $m \leq n$.

LEMMA 4.2. *For any $x \in X$ the order of the subgroup $[H, x]$ is m -bounded.*

PROOF. Since the index of $C_H(x)$ in H is at most m , Lemma 4.1 guarantees that there are elements y_1, \dots, y_m in H such that each y_i is a product of at most $m - 1$ elements of X and the subgroup $[H, x]$

is generated by the commutators $[y_i, x]$, for $i = 1, \dots, m$. For any such i write $y_i = y_{i1} \dots y_{i(m-1)}$, where y_{ij} belongs to X . Using the standard commutator identities, we can rewrite $[y_i, x]$ as a product of conjugates in H of the commutators $[y_{ij}, x]$. Let $\{h_1, \dots, h_s\}$ be the set of conjugates in H of all elements from the set $\{x, y_{ij} \mid 1 \leq i, j \leq m-1\}$. Note that the number s here is m -bounded. This follows from the fact that $C_H(x)$ has index at most m in H for every $x \in X$. Let D be the subgroup of H generated by h_1, \dots, h_s . Since $[H, x]$ is contained in the commutator subgroup D' , it suffices to show that D' has finite m -bounded order. Observe that the center $Z(D)$ has index at most m^s in D , since the index of $C_H(h_i)$ is at most m for every h_i . Thus, by Schur's theorem [29, 10.1.4], we conclude that D' has finite m -bounded order. \square

We now argue by induction on m . If $[H : C_H(g^{l^2})] \leq m-1$ for all $g \in K$, then by induction the result holds. We therefore assume that there is $d \in K$ such that $[H : C_H(d^{l^2})] = m$. Of course, necessarily, $[H : C_H(d^l)] = m$. Set $a = d^l$ and choose b_1, \dots, b_m in H such that $a^H = \{a^{b_i} \mid i = 1, \dots, m\}$ and $w(b_i) \leq m-1$ (the existence of the elements b_i is guaranteed by Lemma 4.1). Since $C_H(a) = C_H(a^l)$, it follows that $(a^l)^H = \{(a^l)^{b_i} \mid i = 1, \dots, m\}$. Set $U = C_G(\langle b_1, \dots, b_m \rangle)$. Note that the index of U in G is n -bounded. Indeed, since $w(b_i) \leq m-1$ we can write $b_i = b_{i1} \dots b_{i(m-1)}$, where $b_{ij} \in X$ and $i = 1, \dots, m$. By the hypothesis, the index of $C_G(b_{ij})$ in G is at most n for any such element b_{ij} . Thus, $[G : U] \leq n^{(m-1)m}$.

LEMMA 4.3. *Suppose that $u \in U$ and $ua \in X$. Then $[H, u] \leq [H, a]$.*

PROOF. For each $i = 1, \dots, m$ we have $(ua)^{b_i} = ua^{b_i}$, since u belongs to U . By hypothesis, $ua \in X$. Hence, taking into account the assumption on the cardinality of the conjugacy class of ua in H , we deduce that $(ua)^H$ consists exactly on the elements $(ua)^{b_i}$, for $i = 1, \dots, m$. Therefore, given an arbitrary element $h \in H$, there exists $b \in \{b_1, \dots, b_m\}$ such that $(ua)^h = (ua)^b$ and so $u^h a^h = ua^b$. It follows that $[u, h] = a^b a^{-h} \in [H, a]$, and the lemma holds. \square

Let R be the normal closure in G of the subgroup $[H, a]$, that is, $R = [H, a^{b_1}] \dots [H, a^{b_n}]$, where a^{b_i} are all the conjugates of a in G (if $|a^G| \leq n-1$, then not all the a^{b_1}, \dots, a^{b_n} are pairwise distinct). By Lemma 4.2, each of the subgroups $[H, a^{b_i}]$ has n -bounded order. Thus, the order of R is n -bounded as well.

Let $Y_1 = Xa^{-l} \cap U$ and $Y_2 = Xa^{-1} \cap U$. Note that for any $y \in Y_1$, the product ya^l belongs to X . So, by Lemma 4.3 applied with a^l in place of a and y in place of u , the subgroup $[H, y]$ is contained in $[H, a^l]$,

which is contained in R . Similarly, for any $y \in Y_2$, we have $[H, y] \leq R$. Set $Y = Y_1 \cup Y_2$. Thus, $[H, Y] \leq R$.

Let U_0 be the maximal normal subgroup of G contained in U . Observe that the index of U_0 in G is n -bounded. Observe further that for any $u \in U_0$ the commutators $[u, a^{-l}]$ and $[u, a^{-1}]$ lie in Y . Since $[U_0, a^{-1}] = [U_0, a]$, we deduce that

$$[H, [U_0, a]] \leq R.$$

Let $K_0 = K \cap U_0$. We remark that $(ua)^l(a^l)^{-1} \in Y$ whenever $u \in K_0$. We pass to the quotient $\overline{G} = G/R$ and use the bar notation to denote images in \overline{G} . We know that \overline{Y} is central in \overline{H} . We also deduce that $[\overline{U}_0, \overline{a}] \leq Z(\overline{H})$.

Since $(ua)^l(a^l)^{-1} \in Y$ whenever $u \in K_0$ and since $\overline{Y} \leq Z(\overline{H})$, it follows that in the quotient $\overline{G}/Z(\overline{H})$ the element \overline{a} commutes with \overline{U}_0 and $(\overline{ua})^l(\overline{a})^{-l} = 1$ for every $\overline{u} \in \overline{K}_0$. This implies that \overline{K}_0 has exponent dividing l modulo $Z(\overline{H})$. It follows that \overline{K}_0^l is abelian and every element of $\overline{K}_0^{l^2}$ is again an l th power of an element in \overline{K}_0 . We therefore deduce that

$$Pr(\overline{K}_0^{l^2}, \overline{G}) \geq \frac{1}{n}.$$

By Proposition 1.1 there is a normal subgroup \overline{T} in \overline{G} and a subgroup \overline{V} in $\overline{K}_0^{l^2}$ such that the indices $[\overline{G} : \overline{T}]$ and $[\overline{K}_0^{l^2} : \overline{V}]$ and the order of $[\overline{T}, \overline{V}]$ are (l, n) -bounded. Let T be the inverse image of \overline{T} in G and V the inverse image of \overline{V} in $K_0^{l^2}$. Bearing in mind that the order of R is n -bounded, we conclude that the indices $[G : T]$ and $[K_0^{l^2} : V]$ are (n, l) -bounded, as also is the order of $[T, V]$. As the index of V in $K_0^{l^2}$ is bounded, there is a positive (n, l) -bounded integer e such that $K^e \leq V$. This completes the proof of the proposition. \square

COROLLARY 4.4. *Let G be a compact group and let G_0 be the connected component of identity in G . Then, the following are equivalent*

- (i) *The probability $Pr(G_0, G)$ is positive;*
- (ii) *The centralizer $C_G(G_0)$ is open in G ;*
- (iii) *The connected component G_0 is ϵ -central in G for some $\epsilon > 0$.*

PROOF. Suppose first that $Pr(G_0, G) > 0$. Then there are subgroups of finite index T of G and B of G_0 such that $[T, B]$ is finite. Since G_0 is divisible [21], the only finite index subgroup of G_0 is G_0 itself. Therefore, the set of commutators $\{[x, y] \mid x \in G_0, y \in T\}$ is connected and finite, hence is trivial and $T \leq C_G(G_0)$.

Now, assume that $C_G(G_0)$ is open in G and let μ and μ_0 be the normalized Haar measures of G and G_0 , respectively. For any $x \in G_0$,

the inclusion $C_G(G_0) \leq C_G(x)$ holds, thus $\mu(C_G(G_0)) \leq \mu(C_G(x))$. We have

$$Pr(\langle x \rangle, G) = \int_{G_0} \mu(C_G(x)) d\mu_0(x) \geq \mu(C_G(G_0)) > 0.$$

We conclude that G_0 is ϵ -central, where $\epsilon = [G : C_G(G_0)]^{-1}$, and so (ii) implies (iii).

Now we assume the validity of (iii) and prove that (i) holds. By Theorem 1.2, there are a finite index subgroup T of G and a natural number e such that $[G_0^e, T]$ is finite. Since G_0 is divisible, $G_0 = G_0^e$ and so T centralizes G_0 and $T \leq C_G(x)$ for any $x \in G_0$. Writing μ and μ_0 for the normalized Haar measures of G and G_0 , respectively, we have

$$Pr(G_0, G) = \int_{G_0} \mu(C_G(x)) d\mu_0(x) \geq \mu(T) > 0.$$

□

5. Corollaries for Finite Groups

In this section we collect some easy corollaries of Theorem 1.2 for finite groups. Roughly, we show that many well-known results on the exponent of a finite group admit a probabilistic interpretation in the spirit of Theorem 1.2.

The restricted Burnside problem was whether the order of an r -generator finite group of exponent e is bounded in terms of r and e alone. This was famously solved in the affirmative by Zelmanov [32, 33]. Theorem 1.2 enables us to obtain the following extension of Zelmanov's theorem.

THEOREM 5.1. *Let G be a finite ϵ -central r -generator group. Then G has a normal subgroup N such that both the index $[G : N]$ and the order of the commutator subgroup $[N, N]$ are (r, ϵ) -bounded.*

PROOF. By Theorem 1.2 there is an ϵ -bounded number e and a normal subgroup $T \leq G$ such that the index $[G : T]$ and the order of $[G^e, T]$ are ϵ -bounded. The solution of the Restricted Burnside problem implies that the index $[G : G^e]$ is (e, r) -bounded. Therefore the subgroup $N = G^e \cap T$ has the required properties. □

An important part of the eventual solution of the restricted Burnside problem was developed by Hall and Higman in their paper [13]. They proved that if p is a prime and G is a finite group with Sylow p -subgroups of exponent p^s , then G has a normal series of s -bounded

length all of whose factors are p -groups, or p' -groups, or direct products of nonabelian simple groups of order divisible by p .

THEOREM 5.2. *Let p be a prime and G a finite group with ϵ -central Sylow p -subgroups. Then G has a normal series of ϵ -bounded length all of whose factors are p -groups, or p' -groups, or direct products of nonabelian simple groups of order divisible by p .*

PROOF. Let P be a Sylow p -subgroup of G . By Theorem 1.2 there is an ϵ -bounded number e and a normal subgroup $T \leq G$ such that the index $[G : T]$ and the order of $[P^e, T]$ are ϵ -bounded. Set $P_0 = P^e$. It is sufficient to show that T has a normal series with the required properties. Note that $[P_0, T]$ is normal in T . Let Z be the inverse image in T of the center of $T/[P_0, T]$. Consider the normal series

$$1 \leq [P_0, T] \leq Z \leq T.$$

Here $[P_0, T]$ has ϵ -bounded order, $Z/[P_0, T]$ is abelian, and T/Z has Sylow p -subgroups of exponent dividing e . According to the Hall-Higman theory T/Z has a normal series of ϵ -bounded length with the required properties. Thus, the result follows. \square

Given a group-word w , we write $w(G)$ for the corresponding verbal subgroup of a group G , that is, the subgroup generated by the values of w in G . The word is said to be a law in G if $w(G) = 1$. In view of Theorem 1.2 it is easy to see that if $w(G)$ is ϵ -central in G , then a law of (ϵ, w) -bounded length holds in the group G . This simple observation provides a tool for obtaining extensions of results about finite groups satisfying certain laws. We will illustrate this with a theorem bounding the nonsoluble length of a finite group. The concept of nonsoluble length $\lambda(G)$ of a finite group G was introduced in [17]. This is the minimum number of nonsoluble factors in a normal series of G in which every factor either is soluble or is a direct product of non-abelian simple groups. (In particular, the group is soluble if and only if its nonsoluble length is 0.)

It was shown in [7] that if a word w is a law in the Sylow 2-subgroup of a finite group G , then $\lambda(G)$ is bounded in terms of the length of the word w only. This can be extended as follows.

THEOREM 5.3. *Let w be a group-word and P a Sylow 2-subgroup of a finite group G such that $w(P)$ is ϵ -central in P . Then $\lambda(G)$ is (ϵ, w) -bounded.*

PROOF. As explained above, this is straightforward combining the result in [7] and Theorem 1.2. \square

For a group of automorphisms A of a group G we write $C_G(A)$ for the centralizer of A in G . It is well-known that if a finite group G admits a coprime group of automorphisms A , then $C_{G/N}(A) = NC_G(A)/N$ for any A -invariant normal subgroup N of G (see for example [8, Theorem 6.2.2 (iv)]). Here the group A is a coprime group of automorphisms if $(|G|, |A|) = 1$. The symbol $A^\#$ stands for the set of nontrivial elements of the group A . The main result of [16] states that if a finite group G admits an elementary abelian coprime group of automorphisms A of order p^2 such that $C_G(\phi)$ has exponent dividing ϵ for each $\phi \in A^\#$, then the exponent of G is (ϵ, p) -bounded. We can now extend this in the following way.

THEOREM 5.4. *Let $\epsilon > 0$, and let G be a finite group admitting an elementary abelian coprime group of automorphisms A of order p^2 such that $C_G(\phi)$ is ϵ -central in G for each $\phi \in A^\#$. Then there is a (p, ϵ) -bounded number e and an A -invariant normal subgroup T such that the index $[G : T]$ and the order of $[G^e, T]$ are (p, ϵ) -bounded.*

PROOF. Let A_1, \dots, A_{p+1} be the subgroups of order p in A and set $G_i = C_G(A_i)$ for $i = 1, \dots, p+1$. According to Theorem 1.2 there is an ϵ -bounded number d and, for $i = 1, \dots, p+1$, A -invariant normal subgroups $T_i \leq G$ such that the index $[G : T_i]$ and the order of $[G_i^d, T_i]$ are ϵ -bounded. Set $T = \bigcap T_i$ and observe that T is A -invariant and the index of T in G is (p, ϵ) -bounded. Let $N_i = [G_i^d, T]$ and $N_0 = \prod N_i$. Note that the subgroup N_0 is normal in T and has (p, ϵ) -bounded order. Let $N = \langle N_0^G \rangle$ be the normal closure of N_0 in G . Since the index of T in G is (p, ϵ) -bounded, it follows that the order of N is (p, ϵ) -bounded as well. We also observe that N is A -invariant since the subgroups N_i are. Let C be the centralizer of T modulo N , that is, $C = \{x \in G \mid [T, x] \leq N\}$. Clearly, the subgroup C is A -invariant. Moreover $G_i^d \leq C$ for $i = 1, \dots, p+1$. Hence, $C_{G/C}(A_i)$ has exponent dividing d for each $i = 1, \dots, p+1$. Now the main result of [16] says that the exponent of G/C is (d, p) -bounded. Therefore there exists a (p, ϵ) -bounded number e such that $G^e \leq C$, that is, $[G^e, T] \leq N$. This completes the proof. \square

COROLLARY 5.5. *Under the hypotheses of Theorem 5.4 there exists a number $\epsilon_0 > 0$ depending only on ϵ and p such that G is ϵ_0 -central.*

PROOF. Assume the hypotheses of Theorem 5.4. The theorem tells us that there is a (p, ϵ) -bounded number e and a normal subgroup T such that the index $[G : T]$ and the order of $[G^e, T]$ are (p, ϵ) -bounded. As explained in the introduction, Theorem 1.2 admits a converse. Hence the result. \square

In the spirit of the work [11] we record the following theorem.

THEOREM 5.6. *Let $\epsilon > 0$, and let G be a finite group admitting an elementary abelian coprime group of automorphisms A of order p^3 such that the commutator subgroup of $C_G(\phi)$ is ϵ -central in G for each $\phi \in A^\#$. Then there is a (p, ϵ) -bounded number e and an A -invariant normal subgroup T such that the index $[G : T]$ and the order of $[[G, G]^e, T]$ are (p, ϵ) -bounded.*

PROOF. Let A_1, \dots, A_s be the subgroups of order p of A and let D_i denote the commutator subgroup of $C_G(A_i)$ for $i = 1, \dots, s$. According to Theorem 1.2 there is an ϵ -bounded number d and, for $i = 1, \dots, s$, A -invariant normal subgroups $T_i \leq G$ such that the index $[G : T_i]$ and the order of $[D_i^d, T_i]$ are ϵ -bounded. Set $T = \bigcap T_i$ and observe that T is A -invariant and the index of T in G is (p, ϵ) -bounded. Let $N_i = [D_i^d, T]$ and $N_0 = \prod N_i$. Note that N_0 is normal in T and has (p, ϵ) -bounded order. Let $N = \langle N_0^G \rangle$ be the normal closure of N_0 in G . Since the index of T in G is (p, ϵ) -bounded, it follows that the order of N is (p, ϵ) -bounded. We also observe that N is A -invariant since the subgroups N_i are. Let C be the centralizer of T modulo N , that is, $C = \{x \in G \mid [T, x] \leq N\}$. Clearly, the subgroup C is A -invariant. Moreover $D_i^d \leq C$ for $i = 1, \dots, s$. Hence, $C_{G/C}(A_i)$ has commutator subgroup of exponent dividing d for each $i = 1, \dots, s$. Now the main result of [11] says that the exponent of the commutator subgroup of G/C is (d, p) -bounded. Therefore there exists a (p, ϵ) -bounded number e such that $[G, G]^e \leq C$, that is, $[[G, G]^e, T] \leq N$. This completes the proof. \square

References

- [1] Acciarri, C., Shumyatsky, P.: A stronger form of Neumann's BFC-theorem. *Israel J. Math.* 242, 269-278 (2021)
- [2] Detomi, E., Morigi, M., Shumyatsky, P.: BFC-theorems for higher commutator subgroups. *Quarterly J. Math.* 70, no. 3, 849 - 858 (2019)
- [3] Detomi, E., Shumyatsky, P.: On the commuting probability for subgroups of a finite group. *Proc. Roy. Soc. Edinburgh Sect. A*, (2022), DOI:10.1017/prm.2021.68
- [4] Dierings, G., Shumyatsky, P.: Groups with boundedly finite conjugacy classes of commutators. *Quarterly J. Math.* 69, no. 3, 1047 - 1051 (2018)
- [5] Eberhard, S.: Commuting probabilities of finite groups. *Bull. London Math. Soc.* 47, 796 - 808 (2015)
- [6] Erfanian, A., Rezaei, R., Lescot, P.: On the relative commutativity degree of a subgroup of a finite group. *Comm. Algebra* 35, 4183-4197 (2007)
- [7] Fumagalli, F., Leinen, F., Puglisi, O.: An upper bound for the nonsolvable length of a finite group in terms of its shortest law. *arXiv:2101.09119*
- [8] Gorenstein, D.: *Finite Groups*. Chelsea Publishing Company, New York (1980)

- [9] Guralnick, R.M., Maroti, A.: Average dimension of fixed point spaces with applications. *J. Algebra* 226, 298 - 308 (2011)
- [10] Guralnick, R.M., Robinson, G.: On the commuting probability in finite groups. *J. Algebra* 300, 509–528 (2006)
- [11] Guralnick, R.M., Shumyatsky, P.: Derived subgroups of fixed points. *Israel J. Math.* 126, 345 - 362 (2001)
- [12] Gustafson, W.H.: What is the probability that two group elements commute? *Amer. Math. Monthly* 80, 1031–1304 (1973)
- [13] Hall, P., Higman, G.: The p -length of a p -soluble group and reduction theorems for Burnside's problem. *Proc. London Math. Soc.* (3) 6, 1 - 42 (1956)
- [14] Hewitt, E., Ross, K.: *Abstract Harmonic Analysis*. Vol. I Springer, Berlin (1963)
- [15] Hofmann, K.H., Russo, F.G.: The probability that x and y commute in a compact group. *Math. Proc. Cambridge Phil. Soc.* 153, 557 - 571 (2012)
- [16] Khukhro, E.I., Shumyatsky, P.: Bounding the exponent of a finite group with automorphisms. *J. Algebra* 212, 363 - 374 (1999)
- [17] Khukhro, E.I., Shumyatsky, P.: Nonsoluble and non- p -soluble length of finite groups. *Israel J. Math.* 207, 507 - 525 (2015)
- [18] Lescot, P.: Sur certains groupes finis. *Rev. Math. Spéciales*, Avril 1987 276-277
- [19] Lescot, P.: Degré de commutativité et structure d'un groupe fini (1). *Rev. Math. Spéciales*, Avril 1988 276–279
- [20] Mann, A.: The exponent of central factors and commutator groups. *J. Group Theory* 10, 435 - 436 (2007)
- [21] Mycielski, J.: Some properties of connected compact groups. *Colloq. Math.* 5, 162 - 166 (1958)
- [22] Nachbin, L.: *The Haar Integral*, Van Nostrand (1965)
- [23] Nath, R.K., Yadav, M.K.: Some results on relative commutativity degree. *Rend. Circ. Mat. Palermo.* 64 (2), 229 - 239 (2015)
- [24] Neumann, B.H.: Groups covered by permutable subsets. *J. London Math. Soc.* (3) 29, 236 - 248 (1954)
- [25] Neumann, P.M.: Two combinatorial problems in group theory. *Bull. Lond. Math. Soc.* 21, 456 - 458 (1989)
- [26] Neumann, P.M., Vaughan-Lee, M.R.: An essay on BFC groups. *Proc. Lond. Math. Soc.* 35, 213 - 237 (1977).
- [27] Reiter, H.: *Classical Harmonic Analysis and Locally Compact Groups*, Oxford, (2000)
- [28] Rezaei, R., Erfanian, A.: On the commutativity degree of compact groups. *Arch. Math.* 43, 345-356 (2009)
- [29] Robinson, D.J.S.: *A course in the theory of groups*, Second edition. Graduate Texts in Mathematics, 80. Springer-Verlag, New York, 1996
- [30] Segal, D., Shalev, A.: On groups with bounded conjugacy classes. *Quart. J. Math. Oxford* 50, 505 - 516 (1999)
- [31] Wiegold, J.: Groups with boundedly finite classes of conjugate elements. *Proc. Roy. Soc. London Ser. A* 238, 389 - 401 (1957)
- [32] Zelmanov, E.I.: Solution of the restricted Burnside problem for groups of odd exponent. *Math. USSR-Izv.* 36, 41 - 60 (1991)
- [33] Zelmanov, E.I.: Solution of the restricted Burnside problem for 2-groups. *Math. USSR-Sb.* 72, 543 - 565 (1992)

- [34] Zelmanov, E.I.: On periodic compact groups. *Isr. J. Math.* 77(1-2), 83 - 95 (1992)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA, BRASILIA-DF,
70910-900 BRAZIL

Email address: J.P.P.Azevedo@mat.unb.br

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA, BRASILIA-DF,
70910-900 BRAZIL

Email address: pavel@unb.br