

Observational constraints of the modified cosmology through Barrow entropy

Mahnaz Asghari ^{*1,2} and Ahmad Sheykhi ^{†1,2}

¹*Physics Department, College of Sciences, Shiraz University, Shiraz 71454, Iran*

²*Biruni Observatory, College of Sciences, Shiraz University, Shiraz 71454, Iran*

Abstract

Taking into account a fractal structure for the black hole horizon, Barrow argued that the area law of entropy is modified due to quantum-gravitational effects (Barrow in *Phys Lett B* 808:135643, <https://doi.org/10.1016/j.physletb.2020.135643>, 2020). Accordingly, the corrected entropy takes the form $S \sim A^{1+\Delta/2}$, where $0 \leq \Delta \leq 1$, indicates the amount of the quantum-gravitational deformation effects. In this paper, based on Barrow entropy, we first derive the modified gravitational field equations through the Clausius relation. We then consider the Friedmann-Lemaître-Robertson-Walker (FLRW) metric as the background metric and derive the modified Friedmann equations inspired by Barrow entropy. In order to explore observational constraints on the modified Barrow cosmology, we employ two different combinations of available datasets, mainly "Planck + Pantheon + BAO" and "Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORSD" datasets. According to numerical results, we observe that the "Planck + Pantheon + BAO" dataset predicts higher values of H_0 in Barrow cosmology with a phantom dark energy compared to Λ CDM, so tensions between low redshift determinations of the Hubble constant and cosmic microwave background (CMB) results are slightly reduced. On the other hand, in case of dataset "Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORSD" there is a slight amelioration in σ_8 tension in Barrow cosmology with a quintessential dark energy compared to the standard model of cosmology. Additionally, for a more reliable comparison, we also constrain the w CDM model with the same datasets, where our results exhibit a satisfying compatibility between Barrow cosmology and w CDM.

1 Introduction

The profound connection between thermodynamics and gravitational field equations has received considerable attention since the discovery of the thermodynamic properties of black holes [1–3]. It has been confirmed that the field equations of general relativity are nothing but an equation of state for the spacetime [4]. In other words, when the spacetime is regarded as a thermodynamic system, the law of thermodynamics on the large scale can be interpreted as the law of gravity. The thermodynamics-gravity conjecture has now been well explored in the literatures [5–13]. The investigation has been generalized to the cosmological background, where it has been shown that the Friedmann equations describing the evolution of the universe can be rewritten in the form of the first law of thermodynamics and vice versa [14–21]. Furthermore, one can apply this remarkable connection in the context of braneworld scenarios [22–25].

In the cosmological approach, it is possible to extract the Friedmann equations of the Friedmann-Lemaître-Robertson-Walker (FLRW) universe by applying the first law of thermodynamics $dE = TdS + WdV$ at the apparent horizon [20]. It was argued that, in any gravity theory, one can consider the entropy expression associated with the apparent horizon in the form of the black hole entropy in the same gravity theory. The only change needed is to replace the black hole horizon radius r_+ in the entropy expression by the apparent horizon radius $\tilde{r}_A$ in the entropy expression associated with the apparent horizon. While it is more convenient to apply the Bekenstein-Hawking area law relation defined as $S_{BH} = A/(4G)$ for the black hole entropy [3, 26] (with the black hole horizon area $A = 4\pi r_+^2$), it should be noted that there are several types of corrections to the area law entropy. Two possible corrections that occur due to quantum effects are known as logarithmic corrections arising from loop quantum gravity [27–34], and also power-law corrections established in the entanglement between quan-

*e-mail: mahnaz.asghari@shirazu.ac.ir

†e-mail: asheykhi@shirazu.ac.ir (corresponding author)

tum fields inside and outside the horizon [35–37]. The other appropriate correction concerning the area law of entropy comes from the fact that the Boltzmann-Gibbs (BG) additive entropy should be generalized to non-additive entropy in divergent partition function systems such as gravitational systems [38–44]. Therefore, it has been argued that the entropy of these systems should be in the form of non-extensive Tsallis entropy $S \sim A^\beta$, where β is the non-extensive or Tsallis parameter [45]. Studies on the non-additive Tsallis entropy and its applications in cosmology have been carried out in [46–55].

Moreover, it is worth turning our attention to Barrow correction to area law entropy due to quantum-gravitational effects [56]. Around 1 year ago, J. D. Barrow considered an intricate and fractal geometry for the black hole horizon, which leads to an increase in surface area. The modified entropy relation based on Barrow corrections takes the form [56]

$$S = \left(\frac{A}{A_0} \right)^{1+\Delta/2}, \quad (1)$$

where A is the black hole horizon area, A_0 is the Planck area, and the exponent Δ , which quantifies the quantum-gravitational deformation, is in the range of $0 \leq \Delta \leq 1$ [56–58]. The area law entropy expression is restored by choosing $\Delta = 0$, which accordingly corresponds to the simplest horizon structure, while $\Delta = 1$ expresses the most intricate surface structure. There are some investigations on Barrow entropy in the cosmological and gravitational setups [59–75].

In order to rewrite the modified field equations from Barrow entropy at apparent horizon of FLRW universe, one should replace the entropy in Clausius relation $\delta Q = T \delta S$ by the entropy expression (1) and consider A as the apparent horizon area given by $A = 4\pi \tilde{r}_A^2$. Our approach in the present study is similar to that in [76], which considers the non-additive Tsallis entropy correction to area law relation. However, it should be mentioned that the physical motivation and principles are completely different between the two investigations. In particular, the Tsallis non-additive entropy correction is motivated by generalizing standard thermodynamics to a non-extensive one, while the Barrow correction to entropy originates from intricate, fractal structure on the horizon induced by quantum-gravitational effects.

It is worth noting that applying the Bekenstein-Hawking entropy in the Clausius relation leads to the Einstein field equations in the standard Λ CDM model [7]. It is known that the standard model of cosmology is in excellent agreement with the majority of observational constraints; however, it suffers from some observational tensions which inspire investigations beyond the standard model. Specifi-

cally, low redshift measurements of the Hubble constant [77–80], and local determinations of the growth of structure [81] are inconsistent with the Planck cosmic microwave background (CMB) observations [82]. Accordingly, in this paper we study whether the discrepancies between local and global measurements can be resolved in Barrow cosmology. Throughout the paper we set $k_B = c = \hbar = 1$ for simplicity.

This paper is structured as follows. In Sect. 2 we derive modified field equations describing the evolutions of the universe when the horizon entropy is given by Eq. (1). Numerical solutions based on Barrow entropy corrections to the field equations are presented in Sect. 3. Section 4 is dedicated to constraining Barrow cosmology with observational data. We summarize our conclusions in Sect. 5.

2 Modified gravitational field equations from Barrow corrections

In this part, we will derive the corresponding field equations in the cosmological setup when the entropy associated with the apparent horizon is in the form of Barrow entropy. In this respect, we consider a spatially flat FLRW universe with the background metric given by

$$ds^2 = a^2(\tau)(-d\tau^2 + d\mathbf{x}^2), \quad (2)$$

where τ is the conformal time. Also, the perturbed line element in linear perturbation theory in the conformal Newtonian gauge reads

$$ds^2 = a^2(\tau) \left(-(1+2\Psi)d\tau^2 + (1-2\Phi)d\mathbf{x}^2 \right), \quad (3)$$

with gravitational potentials Ψ and Φ . Similarly in the synchronous gauge we have

$$ds^2 = a^2(\tau) \left(-d\tau^2 + (\delta_{ij} + h_{ij})dx^i dx^j \right), \quad (4)$$

where $h_{ij} = \text{diag}(-2\eta, -2\eta, h+4\eta)$, with scalar perturbations h and η .

Furthermore, we consider the universe consists of radiation (R), matter (M) (dark matter [DM] and baryons [B]), and dark energy (DE), which are assumed to be perfect fluids, with the energy-momentum tensor defined as

$$T_{\mu\nu(i)} = (\rho_i + p_i)u_{\mu(i)}u_{\nu(i)} + g_{\mu\nu}p_i, \quad (5)$$

where $\rho_i = \bar{\rho}_i + \delta\rho_i$ is the energy density, $p_i = \bar{p}_i + \delta p_i$ is the pressure and $u_{\mu(i)}$ is the four-velocity of the i^{th} component in the universe (and a bar indicates the background level).

Employing the Clausius relation, $\delta Q = T\delta S$, which is satisfied on a local causal horizon $\mathcal{H}$, we derive the gravitational field equations. Considering the universe as a thermodynamic system, we apply the Clausius relation on the apparent horizon of the universe with radius $\tilde{r}_A$, defined as [83]

$$\tilde{r}_A = \left(H^2 + \frac{K}{a^2}\right)^{-1/2}, \quad (6)$$

where H is the Hubble parameter and $K = -1, 0, 1$ is the curvature constant corresponding to an open, flat and closed universe, respectively. Also, the associated temperature with the apparent horizon is given by

$$T = \frac{\kappa}{2\pi}, \quad (7)$$

where κ is the surface gravity at the apparent horizon. As mentioned before, in the interest of deriving field equations in Barrow cosmology, we should apply the Barrow entropy defined in Eq. (1) in the Clausius relation. Thus, δS takes the form

$$\delta S = \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} A^{\Delta/2} \delta A. \quad (8)$$

According to Refs. [4, 7], δQ can be written as

$$\delta Q = -\kappa \int_{\mathcal{H}} \lambda T_{\mu\nu} k^\mu k^\nu d\lambda dA, \quad (9)$$

and δA is given by

$$\delta A = \int_{\mathcal{H}} \theta d\lambda dA, \quad (10)$$

with expansion θ defined as

$$\theta = -\lambda R_{\mu\nu} k^\mu k^\nu. \quad (11)$$

Thus, considering Eqs. (7), (8) and (9), the Clausius relation takes the form

$$\begin{aligned} & \kappa \int_{\mathcal{H}} (-\lambda) T_{\mu\nu} k^\mu k^\nu d\lambda dA \\ &= \frac{\kappa}{2\pi} \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \int_{\mathcal{H}} (-\lambda) R_{\mu\nu} k^\mu k^\nu A^{\Delta/2} d\lambda dA, \end{aligned} \quad (12)$$

$$\begin{aligned} & \rightarrow \int_{\mathcal{H}} (-\lambda) \left(-2\pi T_{\mu\nu} + \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} R_{\mu\nu} A^{\Delta/2} \right) \\ & \quad \times k^\mu k^\nu d\lambda dA = 0. \end{aligned} \quad (13)$$

Then, for all null vectors k^μ we find

$$-2\pi T_{\mu\nu} + \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} R_{\mu\nu} A^{\Delta/2} = f g_{\mu\nu}, \quad (14)$$

where f is a scalar. Thus, according to energy-momentum conservation ($\nabla^\mu T_{\mu\nu} = 0$), we must have

$$\nabla^\mu \left(\left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} R_{\mu\nu} A^{\Delta/2} - f g_{\mu\nu} \right) = 0, \quad (15)$$

$$\rightarrow \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \left(\frac{1}{2} (\partial_\nu R) A^{\Delta/2} + R_{\mu\nu} \partial^\mu A^{\Delta/2} \right) = \partial_\nu f. \quad (16)$$

Considering Eq. (16), the left-hand side is not the gradient of a scalar, which means that this corresponds to non-equilibrium behaviour of thermodynamics, and the Clausius relation is not satisfied. Thus, the Clausius relation should be replaced by the entropy balance relation [7]

$$\delta S = \frac{\delta Q}{T} + d_i S, \quad (17)$$

where $d_i S$ is the entropy produced inside the system caused by irreversible transformations of the system [84]. Then, in order to resolve the contradiction with energy-momentum conservation, we assume $d_i S$ as the following form

$$d_i S = \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \int_{\mathcal{H}} (-\lambda) \nabla_\mu \nabla_\nu A^{\Delta/2} k^\mu k^\nu d\lambda dA. \quad (18)$$

Substituting $d_i S$ in relation (17) results in

$$\begin{aligned} & \int_{\mathcal{H}} (-\lambda) \left(\left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} R_{\mu\nu} A^{\Delta/2} - 2\pi T_{\mu\nu} \right. \\ & \quad \left. - \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \nabla_\mu \nabla_\nu A^{\Delta/2} \right) k^\mu k^\nu d\lambda dA = 0. \end{aligned} \quad (19)$$

Again, for all null vectors k^μ we should have

$$\begin{aligned} & \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} R_{\mu\nu} A^{\Delta/2} - 2\pi T_{\mu\nu} \\ & - \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \nabla_\mu \nabla_\nu A^{\Delta/2} = f g_{\mu\nu}. \end{aligned} \quad (20)$$

Then, considering energy-momentum conservation, and after doing some calculations, we obtain

$$\left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \left(\frac{1}{2} (\partial_\nu R) A^{\Delta/2} - \partial_\nu \square A^{\Delta/2} \right) = \partial_\nu f. \quad (21)$$

Here, we can choose the scalar $\mathcal{L}$ as $\mathcal{L} = R A^{\Delta/2}$, which reads

$$\begin{aligned} & \frac{\partial \mathcal{L}}{\partial R} = A^{\Delta/2}, \\ & \partial_\nu R = \frac{\partial R}{\partial x^\nu} = \frac{\partial R}{\partial \mathcal{L}} \frac{\partial \mathcal{L}}{\partial x^\nu} = A^{-\Delta/2} \partial_\nu \mathcal{L}, \\ & \rightarrow (\partial_\nu R) A^{\Delta/2} = \partial_\nu \mathcal{L}. \end{aligned} \quad (22)$$

Therefor, the scalar f becomes

$$\begin{aligned} f &= \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \left(\frac{1}{2} \mathcal{L} - \square A^{\Delta/2}\right) \\ &= \left(1 + \frac{\Delta}{2}\right) A_0^{-1-\Delta/2} \left(\frac{1}{2} R A^{\Delta/2} - \square A^{\Delta/2}\right). \end{aligned} \quad (23)$$

Thus, the modified field equations in Barrow cosmology can be written as

$$\begin{aligned} R_{\mu\nu} A^{\Delta/2} - \nabla_\mu \nabla_\nu A^{\Delta/2} - \frac{1}{2} R A^{\Delta/2} g_{\mu\nu} + \square A^{\Delta/2} g_{\mu\nu} \\ = \frac{4\pi}{2+\Delta} A_0^{1+\Delta/2} T_{\mu\nu}, \end{aligned} \quad (24)$$

in which for a flat spacetime, we have

$$A^{\Delta/2} = (4\pi)^{\Delta/2} \left(\frac{a'}{a^2}\right)^{-\Delta} = (4\pi)^{\Delta/2} H^{-\Delta}, \quad (25)$$

where a prime indicates a deviation with respect to the conformal time. In this way, we derive the modified Einstein field equations based on Barrow corrections to area law entropy caused by quantum-gravitational effects on the apparent horizon of the FLRW universe. Considering $\Delta = 0$, the standard field equations in Einstein gravity will be recovered. The (00) and (ii) components of gravitational field equations (24) at background level take the form

$$\left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ \left(\frac{a'}{a}\right)^2 - \Delta \left(\frac{a''}{a} - 2\left(\frac{a'}{a}\right)^2\right) \right\} = \frac{8\pi G}{3} a^2 \sum_i \bar{\rho}_i, \quad (26)$$

$$\begin{aligned} \left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ -2\frac{a''}{a} + \left(\frac{a'}{a}\right)^2 + \Delta \left[-\frac{a''}{a} + \frac{a'''}{a'} - \left(\frac{a''}{a'}\right)^2 \right. \right. \\ \left. \left. - \Delta \left(\frac{a''}{a'} - 2\frac{a'}{a}\right)^2 \right] \right\} = 8\pi G a^2 \sum_i \bar{p}_i, \end{aligned} \quad (27)$$

where we have defined A_0 as

$$A_0 = (4\pi)^{(\Delta-2)/(2+\Delta)} \left((2+\Delta)8\pi G\right)^{2/(2+\Delta)}.$$

Also, the background level equations in term of the Hubble parameter are given by

$$H^{2-\Delta} - \Delta H^{-\Delta} H' \frac{1}{a} = \frac{8\pi G}{3} \sum_i \bar{\rho}_i, \quad (28)$$

$$\begin{aligned} H^{-\Delta} \left\{ (\Delta-2) H' \frac{1}{a} - 3H^2 \right. \\ \left. + \Delta \left(\frac{H''}{H} - (1+\Delta) \left(\frac{H'}{H}\right)^2 \right) \frac{1}{a^2} \right\} = 8\pi G \sum_i \bar{p}_i. \end{aligned} \quad (29)$$

Then, from (28) and (29), the first modified Friedmann equation can be derived as

$$H^{2-\Delta} = \frac{1}{1+2\Delta} \frac{8\pi G}{3} \sum_i \bar{\rho}_i. \quad (30)$$

It is also convenient to rewrite the Friedmann equation in term of the total density parameter defined as $\Omega_{\text{tot}} = \bar{\rho}_{\text{tot}}/\rho_{\text{cr}}$, where $\rho_{\text{cr}} = 3H^2/(8\pi G)$ and $\bar{\rho}_{\text{tot}} = \sum_i \bar{\rho}_i$. Hence, Eq. (30) can be written as

$$\Omega_{\text{tot}} = (1+2\Delta) H^{-\Delta}. \quad (31)$$

It should be noted that in the limit $\Delta \rightarrow 0$, the Friedmann equation takes the standard form in general relativity.

Defining the total equation of state parameter w_{tot} as $w_{\text{tot}} = \bar{p}_{\text{tot}}/\bar{\rho}_{\text{tot}}$, the accelerated expansion of the universe in Barrow cosmology is satisfied when $w_{\text{tot}} < -(1+\Delta)/3$. Taking into account the fact that $0 \leq \Delta \leq 1$, Barrow corrections predict a more negative equation of state parameter in an accelerating universe (for example, for $\Delta = 1$ we obtain $w_{\text{tot}} < -2/3$), while in the limit $\Delta \rightarrow 0$, the accelerated expansion condition in standard cosmology will be restored.

Taking into account the modified field equations (24) to linear order of perturbations, in the conformal Newtonian gauge (con) we have

$$\begin{aligned} \left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ 3\frac{a'}{a} \Phi' + k^2 \Phi + 3\left(\frac{a'}{a}\right)^2 \Psi \right. \\ \left. - \frac{3}{2} \Delta \left(\frac{a''}{a'} - 2\frac{a'}{a}\right) \left(\Phi' + 2\frac{a'}{a} \Psi\right) \right\} = -4\pi G a^2 \sum_i \delta \rho_{i(\text{con})}, \end{aligned} \quad (32)$$

$$\begin{aligned} \left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ k^2 \Phi' + \frac{a'}{a} k^2 \Psi - \frac{1}{2} \Delta \left(\frac{a''}{a'} - 2\frac{a'}{a}\right) k^2 \Psi \right\} \\ = 4\pi G a^2 \sum_i (\bar{\rho}_i + \bar{p}_i) \theta_{i(\text{con})}, \end{aligned} \quad (33)$$

$$\Phi = \Psi, \quad (34)$$

$$\begin{aligned} \left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ 2\frac{a''}{a} \Psi - \left(\frac{a'}{a}\right)^2 \Psi + \frac{a'}{a} \Psi' + 2\frac{a'}{a} \Phi' + \Phi'' \right. \\ \left. + \frac{k^2}{3} (\Phi - \Psi) + \Delta \left\{ \Psi \left[\Delta \left(\frac{a''}{a'} - 2\frac{a'}{a}\right)^2 - \frac{a'''}{a'} + \frac{a''}{a} \right. \right. \right. \\ \left. \left. + \left(\frac{a''}{a'}\right)^2 \right] + \Psi' \left(\frac{a'}{a} - \frac{1}{2} \frac{a''}{a'}\right) + \Phi' \left(2\frac{a'}{a} - \frac{a''}{a'}\right) \right\} \right\} \\ = 4\pi G a^2 \sum_i \delta p_{i(\text{con})}, \end{aligned} \quad (35)$$

while in the synchronous gauge (syn) we can write

$$\left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ \frac{a'}{a} h' - 2k^2 \eta - \frac{1}{2} \Delta h' \left(\frac{a''}{a'} - 2 \frac{a'}{a} \right) \right\} = 8\pi G a^2 \sum_i \delta \rho_{i(\text{syn})}, \quad (36)$$

$$\left(\frac{a'}{a^2}\right)^{-\Delta} k^2 \eta' = 4\pi G a^2 \sum_i (\bar{\rho}_i + \bar{p}_i) \theta_{i(\text{syn})}, \quad (37)$$

$$\left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ \frac{1}{2} h'' + 3\eta'' + \frac{a'}{a} h' + 6 \frac{a'}{a} \eta' - k^2 \eta - \frac{1}{2} \Delta \left(\frac{a''}{a'} - 2 \frac{a'}{a} \right) \left(h' + 6\eta' \right) \right\} = 0, \quad (38)$$

$$\left(\frac{a'}{a^2}\right)^{-\Delta} \left\{ -2 \frac{a'}{a} h' - h'' + 2k^2 \eta + \Delta \left(\frac{a''}{a'} - 2 \frac{a'}{a} \right) h' \right\} = 24\pi G a^2 \sum_i \delta p_{i(\text{syn})}. \quad (39)$$

Choosing $\Delta = 0$ would result in standard field equations at the perturbation level in Einstein gravity.

Furthermore, considering energy-momentum conservation, continuity and Euler equations would not be affected by Barrow corrections, and so conservation equations are identical to those in general relativity. In the rest of the paper, we explore Barrow cosmology in the synchronous gauge; hence, conservation equations for matter and dark energy components take the form

$$\delta'_{\text{M(syn)}} = -\theta_{\text{M(syn)}} - \frac{1}{2} h', \quad (40)$$

$$\theta'_{\text{M(syn)}} = -\frac{a'}{a} \theta_{\text{M(syn)}}, \quad (41)$$

$$\delta'_{\text{DE(syn)}} = -3 \frac{a'}{a} (c_{s,\text{DE}}^2 - w_{\text{DE}}) \delta_{\text{DE(syn)}} - \frac{1}{2} h' (1 + w_{\text{DE}}) - (1 + w_{\text{DE}}) \left(1 + 9 \left(\frac{a'}{a} \right)^2 (c_{s,\text{DE}}^2 - c_{a,\text{DE}}^2) \frac{1}{k^2} \right) \theta_{\text{DE(syn)}}, \quad (42)$$

$$\theta'_{\text{DE(syn)}} = \frac{a'}{a} (3c_{s,\text{DE}}^2 - 1) \theta_{\text{DE(syn)}} + \frac{k^2 c_{s,\text{DE}}^2}{1 + w_{\text{DE}}} \delta_{\text{DE(syn)}}. \quad (43)$$

In the next section, we study Barrow cosmology using a modified version of the Boltzmann code CLASS¹ [85], in

¹Cosmic linear anisotropy solving system

which we have included the Barrow parameter Δ that quantifies deviations from standard cosmology. Moreover, in order to constrain cosmological parameters from current observations, we employ the Markov chain Monte Carlo (MCMC) method by making use of the MONTE PYTHON code [86, 87].

3 Numerical analysis

In order to study the Barrow cosmology model numerically, we modify the CLASS code according to the Barrow cosmology field equations described in Sect. 2. In this direction, we consider Planck 2018 results [82] for the cosmological parameters, which read $\Omega_{\text{B},0} h^2 = 0.02242$, $\Omega_{\text{DM},0} h^2 = 0.11933$, $H_0 = 67.66 \frac{\text{km}}{\text{sMpc}}$, $A_s = 2.105 \times 10^{-9}$, and $\tau_{\text{reio}} = 0.0561$, and also for the dark energy fluid we choose $w_{\text{DE}} = -0.98$ (to avoid divergences in dark energy perturbations) and $c_{s,\text{DE}}^2 = 1$.

Figure 1 displays the CMB temperature anisotropy and matter power spectra diagrams in Barrow cosmology compared to Λ CDM (it should be noted that the case $\Delta = 0$ corresponds to the w CDM model, where it consists of cold dark matter and a dark energy fluid with a constant equation of state).

Matter power spectra diagrams show an enhancement in structure growth in the Barrow cosmology model, which is inconsistent with low redshift measurements of galaxy clustering. The evolution of matter density contrast illustrated in Fig. 2 would also reflect the increase in the growth of structures in Barrow cosmology.

On the other hand, considering the Friedmann equation (30), it is possible to explore the expansion history of the universe in Barrow cosmology as demonstrated in Fig. 3. According to this figure, it can be understood that the current value of the Hubble parameter in Barrow cosmology is more compatible with its local determinations in comparison with the Λ CDM model.

Moreover, in Fig. 4 we show the evolution of dark energy density, which illustrates an enhancement in $\bar{\rho}_{\text{DE}}$ of Barrow cosmology in comparison with the Λ CDM model, and consequently confirms the increase in the current value of the Hubble parameter in the Barrow model.

4 Comparison with observational data

In this section, we put constraints on the parameters of Barrow cosmology by applying an MCMC approach through the MONTE PYTHON code [86, 87]. The set of cosmological parameters used in MCMC analysis consists of $\{100\Omega_{\text{B},0} h^2, \Omega_{\text{DM},0} h^2, 100\theta_s, \ln(10^{10} A_s), n_s, \tau_{\text{reio}}, w_{\text{DE}},$




Figure 1: The CMB power spectra (upper left) and their relative ratios with respect to Λ CDM (upper right) for different values of Δ . Lower panels show the analogous diagrams for the matter power spectrum




Figure 2: Matter density contrast diagrams in terms of conformal time in Barrow cosmology compared to Λ CDM




Figure 3: Hubble parameter in terms of conformal time in Barrow cosmology compared to Λ CDM

Δ }, where $\Omega_{B,0}h^2$ and $\Omega_{DM,0}h^2$ represent the baryon and cold dark matter densities relative to the critical density, respectively, θ_s is the ratio of the sound horizon to the angular diameter distance at decoupling, A_s stands for the amplitude of the primordial scalar perturbation spectrum, n_s is the scalar spectral index, τ_{reio} is the optical depth to reionization, w_{DE} is the dark energy equation of state parameter, and Δ indicates the Barrow parameter. Furthermore, we have four derived parameters including the reionization redshift (z_{reio}), the matter density parameter ($\Omega_{M,0}$), the Hubble constant (H_0), and the root-mean-square mass fluctuations on scales of $8 h^{-1} \text{Mpc}$ (σ_8). Considering preliminary nu-

merical studies, we choose the prior range $[0, 0.015]$ for the Barrow parameter, and additionally, we set no prior range on w_{DE} .

The following likelihoods are utilized in the MCMC method: The Planck likelihood with Planck 2018 data (containing high- l TT, TE, EE, low- l EE, low- l TT, and lensing) [82], the Planck-SZ likelihood for the Sunyaev-Zeldovich (SZ) effect measured by Planck [88, 89], the CFHTLenS likelihood with the weak lensing data [90, 91], the Pantheon likelihood with the supernovae data [92], the BAO likelihood with the baryon acoustic oscillations data [93, 94], and the BAORS likelihood for BAO and redshift-space distortions.




Figure 4: The evolution of dark energy density in terms of conformal time in Barrow cosmology compared to Λ CDM

tion (RSD) measurements [95, 96].

In order to constrain the cosmological model under consideration, we use two different dataset combinations: "Planck + Pantheon + BAO" [hereafter dataset (I)] and "Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORS" [hereafter dataset (II)]. Tables 1 and 2 represent the observational constraints from two different datasets (I) and (II), respectively, where we have considered Λ CDM and also wCDM as our base models.

Marginalized 1σ and 2σ confidence level contour plots from datasets (I) and (II) for selected cosmological parameters of Barrow cosmology are also depicted in Fig. 5. Furthermore, in order to compare Barrow cosmology with Tsallis modified gravity (TMG), the results based on the TMG model according to Ref. [76] are also displayed.

Considering dataset (I), we can see an enhancement in the Hubble constant in Barrow cosmology compared to standard model, which indicates a slight amelioration in H_0 tension. Moreover, dark energy has a phantom behaviour according to the best fit and mean value of w_{DE} , which is influential in alleviating the H_0 tension. However, a quintessential character of dark energy is also allowed within the 1σ confidence level, where $-1.048 < w_{DE} < -0.9735$. Also, to be more precise, we compare Barrow cosmology with the wCDM model, which also represents a phantom behaviour of dark energy. Then, there is a minor increase in Hubble constant of the wCDM model in comparison with Λ CDM, which confirms that Barrow cosmology is in acceptable agreement with wCDM. Furthermore, according to the obtained constraints on Barrow parameter Δ and dark energy equation of state, one can conclude that Barrow cosmology is in a reasonable consistency with the Λ CDM model. For the last point, it should be emphasized that best fit and mean value of Δ are compatible with Λ CDM, and the minor increase in the Hubble constant caused by the phantom nature of dark energy would only slightly alleviate the H_0

tension and not solve it completely.

On the other hand, dataset (II) results show a suppression in the growth of structure in Barrow cosmology with respect to the Λ CDM model. Actually, one anticipates higher values of σ_8 in Barrow cosmology according to numerical results described in Sect. 3. However, the quintessential behaviour of dark energy can reduce the structure growth and consequently provide more compatible results with local galaxy surveys. Likewise, the wCDM model has a quintessential character of a dark energy equation of state, which yields a lower σ_8 growth rate than the standard cosmological model. Thus, MCMC analysis implies that Barrow cosmology is in good agreement with wCDM. Additionally, observational constraints on w_{DE} and Δ confirm that the Barrow model is also compatible with Λ CDM.

It should be noted that, considering datasets (I) and (II), simultaneous alleviation of existing discrepancies between local observations and CMB measurements is not possible in Barrow cosmology due to the correlation between σ_8 and H_0 .

Moreover, it is worthwhile to compare Barrow cosmology with the TMG model based on the numerical results from dataset (II). Considering the fact that Barrow cosmology and the TMG model are established on absolutely different physical principles, the growth of structure is slightly reduced in both the Barrow and Tsallis scenarios, related to the quintessential behaviour of dark energy, which is also consistent with the wCDM model. The structure growth suppression is more important in the TMG model than in Barrow cosmology, according to the behaviour of β and Δ . Specifically, the derived best fit and mean value of Δ correspond to higher values of σ_8 , in contrast to the quintessential character of dark energy which results in a decrease in the growth of structure.

Also, the Akaike information criterion (AIC) defined as [97, 98]

$$AIC = -2 \ln \mathcal{L}_{\max} + 2K, \quad (44)$$

with $\mathcal{L}_{\max}$ the maximum likelihood function and K the number of free parameters, is helpful for evaluating which model is better supported by observational data. Considering Λ CDM and wCDM as reference models, we obtain the following results

$$\begin{aligned} \text{dataset(I)} : AIC_{(\Lambda\text{CDM})} &= 3817.12, \quad AIC_{(w\text{CDM})} = 3818.08, \\ AIC_{(\text{Barrow})} &= 3820.20, \\ \rightarrow \begin{cases} AIC_{(\text{Barrow})} - AIC_{(\Lambda\text{CDM})} &= 3.08, \\ AIC_{(\text{Barrow})} - AIC_{(w\text{CDM})} &= 2.12, \end{cases} \end{aligned}$$

Table 1: Best-fit values and 68% and 95% confidence level intervals for cosmological parameters of Λ CDM, wCDM, and Barrow cosmology, considering dataset (I)

parameter	Planck + Pantheon + BAO					
	Λ CDM		wCDM		Barrow cosmology	
	best fit	68% & 95% limits	best fit	68% & 95% limits	best fit	68% & 95% limits
$100\Omega_{B,0}h^2$	2.246	$2.245^{+0.013+0.027}_{-0.014-0.028}$	2.238	$2.239^{+0.014+0.028}_{-0.014-0.028}$	2.235	$2.239^{+0.015+0.030}_{-0.015-0.028}$
$\Omega_{DM,0}h^2$	0.1191	$0.1189^{+0.00096+0.0018}_{-0.00086-0.0020}$	0.1198	$0.1194^{+0.0011+0.0020}_{-0.00099-0.0021}$	0.1203	$0.1197^{+0.0012+0.0021}_{-0.00095-0.0023}$
$100\theta_s$	1.042	$1.042^{+0.00029+0.00057}_{-0.00030-0.00059}$	1.042	$1.042^{+0.00029+0.00057}_{-0.00031-0.00056}$	1.042	$1.042^{+0.00029+0.00055}_{-0.00029-0.00057}$
$\ln(10^{10}A_s)$	3.048	$3.047^{+0.013+0.028}_{-0.014-0.028}$	3.040	$3.045^{+0.014+0.030}_{-0.014-0.027}$	3.040	$3.046^{+0.012+0.029}_{-0.015-0.027}$
n_s	0.9675	$0.9673^{+0.0036+0.0076}_{-0.0038-0.0074}$	0.9647	$0.9660^{+0.0040+0.0081}_{-0.0041-0.0078}$	0.9685	$0.9665^{+0.0042+0.0079}_{-0.0040-0.0081}$
τ_{reio}	0.05699	$0.05662^{+0.0063+0.014}_{-0.0073-0.013}$	0.05223	$0.05536^{+0.0061+0.014}_{-0.0076-0.014}$	0.05135	$0.05483^{+0.0061+0.014}_{-0.0080-0.014}$
w_{DE}	—	—	-1.018	$-1.034^{+0.035+0.065}_{-0.032-0.067}$	-1.026	$-1.009^{+0.036+0.070}_{-0.039-0.072}$
Δ	—	—	—	—	0.0008549	$0.0007799^{+0.00021+0.0011}_{-0.00078-0.00078}$
z_{reio}	7.921	$7.870^{+0.64+1.4}_{-0.72-1.3}$	7.463	$7.757^{+0.64+1.4}_{-0.73-1.4}$	7.376	$7.702^{+0.62+1.4}_{-0.78-1.4}$
$\Omega_{M,0}$	0.3038	$0.3022^{+0.0057+0.011}_{-0.0050-0.011}$	0.3015	$0.2963^{+0.0088+0.016}_{-0.0076-0.016}$	0.2922	$0.2953^{+0.0077+0.016}_{-0.0084-0.016}$
$H_0 \left[\frac{\text{km}}{\text{sMpc}} \right]$	68.28	$68.41^{+0.37+0.90}_{-0.46-0.82}$	68.67	$69.20^{+0.88+1.9}_{-0.97-1.7}$	69.86	$69.39^{+0.96+1.8}_{-0.90-1.9}$
σ_8	0.8225	$0.8217^{+0.0056+0.012}_{-0.0061-0.011}$	0.8264	$0.8319^{+0.011+0.024}_{-0.012-0.022}$	0.8408	$0.8343^{+0.012+0.024}_{-0.012-0.024}$

dataset(II) : $\text{AIC}_{(\Lambda\text{CDM})} = 3847.12$, $\text{AIC}_{(\text{wCDM})} = 3849.38$, the deviation from standard cosmology. Furthermore, Einstein field equations can be derived from the first law of thermodynamics at the apparent horizon of the FLRW universe, and vice versa, inspired by the remarkable analogy between thermodynamics and gravity. In this direction, it is possible to associate an entropy to the apparent horizon as the same expression of black hole entropy, just by replacing the black hole horizon radius by the apparent horizon radius. Accordingly, we derived modified field equations in Barrow cosmology by applying Barrow entropy in a Clausius relation. Then, by performing MCMC analysis, we put constraints on the cosmological parameters, specifically the Barrow parameter Δ which quantifies deviations from Λ CDM, based on two different combinations of datasets: "Planck + Pantheon + BAO" [dataset (I)] and "Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORSD" [dataset (II)].

$$\begin{aligned} & \text{AIC}_{(\text{Barrow})} = 3850.72 , \\ & \rightarrow \begin{cases} \text{AIC}_{(\text{Barrow})} - \text{AIC}_{(\Lambda\text{CDM})} = 3.6 , \\ \text{AIC}_{(\text{Barrow})} - \text{AIC}_{(\text{wCDM})} = 1.34 . \end{cases} \end{aligned}$$

Therefor, we can understand that the Λ CDM model is better fitted to both datasets (I) and (II) compared to Barrow cosmology; however, one can not rule out the Barrow cosmology model. On the other hand, according to dataset (I), wCDM is preferred to Barrow cosmology, while the Barrow model is still valid. Moreover, dataset (II) indicates that the Barrow model is supported by observational data as well as wCDM.

5 Conclusions

It is argued by Barrow [56] that a black hole horizon might have an intricate, fractal structure caused by quantum gravitational corrections to the area law of entropy. In this respect, the Barrow entropy relation (1) is associated with the black hole horizon, with the new exponent Δ that measures

Numerical results from dataset (I) indicate that the H_0 tension can be slightly relieved in Barrow cosmology with a phantom behaviour of dark energy compared to the Λ CDM model; however, the tension is not solved completely.

Table 2: Best-fit values and 68% and 95% confidence level intervals for cosmological parameters of Λ CDM, wCDM, Barrow cosmology (with Barrow parameter Δ), and also the TMG model (with Tsallis parameter β) from Ref. [76], considering dataset (II)

Planck + Planck-SZ + CFHTLenS + Pantheon + BAO + BAORSD								
	Λ CDM		wCDM		Barrow cosmology		TMG model	
parameter	best fit	68% & 95% limits	best fit	68% & 95% limits	best fit	68% & 95% limits	best fit	68% & 95% limits
$100\Omega_{B,0}h^2$	2.261	$2.263^{+0.012+0.026}_{-0.013-0.025}$	2.260	$2.265^{+0.012+0.026}_{-0.013-0.025}$	2.270	$2.264^{+0.013+0.027}_{-0.013-0.027}$	2.272	$2.268^{+0.014+0.027}_{-0.015-0.028}$
$\Omega_{DM,0}h^2$	0.1163	$0.1164^{+0.00078+0.0015}_{-0.00079-0.0015}$	0.1161	$0.1162^{+0.00083+0.0016}_{-0.00080-0.0017}$	0.1161	$0.1164^{+0.00092+0.0016}_{-0.00076-0.0017}$	0.1166	$0.1160^{+0.00097+0.0016}_{-0.00077-0.0017}$
$100\theta_s$	1.042	$1.042^{+0.00029+0.00055}_{-0.00026-0.00053}$	1.042	$1.042^{+0.00030+0.00059}_{-0.00029-0.00056}$	1.042	$1.042^{+0.00029+0.00059}_{-0.00029-0.00056}$	1.042	$1.042^{+0.00026+0.00054}_{-0.00028-0.00052}$
$\ln(10^{10}A_s)$	3.034	$3.024^{+0.010+0.023}_{-0.014-0.021}$	3.029	$3.028^{+0.010+0.025}_{-0.015-0.023}$	3.020	$3.028^{+0.012+0.025}_{-0.014-0.022}$	3.026	$3.028^{+0.011+0.024}_{-0.014-0.025}$
n_s	0.9712	$0.9719^{+0.0036+0.0072}_{-0.0039-0.0074}$	0.9735	$0.9723^{+0.0036+0.0071}_{-0.0035-0.0074}$	0.9706	$0.9728^{+0.0038+0.0078}_{-0.0041-0.0077}$	0.9706	$0.9720^{+0.0038+0.0074}_{-0.0040-0.0076}$
τ_{reio}	0.05358	$0.04963^{+0.0041+0.010}_{-0.0074-0.0096}$	0.05220	$0.05090^{+0.0052+0.012}_{-0.0075-0.011}$	0.04839	$0.05074^{+0.0054+0.012}_{-0.0077-0.011}$	0.05102	$0.05158^{+0.0059+0.011}_{-0.0072-0.012}$
$w_{\text{DE}} (w_f)$	—	—	-0.9731	$-0.9733^{+0.031+0.054}_{-0.025-0.055}$	-0.9762	$-0.9569^{+0.033+0.065}_{-0.032-0.066}$	-0.9677	$-0.9944^{+0.041+0.089}_{-0.046-0.084}$
$\Delta (\beta)$	—	—	—	—	0.0002247	$0.0005340^{+0.00013+0.00092}_{-0.00053-0.00053}$	0.9999	$0.9997^{+0.00047+0.00098}_{-0.00048-0.00090}$
z_{reio}	7.502	$7.084^{+0.50+1.0}_{-0.69-1.0}$	7.366	$7.211^{+0.53+1.2}_{-0.76-1.2}$	6.951	$7.196^{+0.55+1.1}_{-0.78-1.2}$	7.231	$7.275^{+0.59+1.2}_{-0.72-1.2}$
$\Omega_{M,0}$	0.2871	$0.2876^{+0.0043+0.0086}_{-0.0044-0.0086}$	0.2934	$0.2941^{+0.0072+0.016}_{-0.0083-0.016}$	0.2897	$0.2938^{+0.0078+0.017}_{-0.0084-0.017}$	0.2994	$0.2945^{+0.0076+0.016}_{-0.0074-0.015}$
$H_0 \left[\frac{\text{km}}{\text{sMpc}}\right]$	69.56	$69.54^{+0.37+0.73}_{-0.36-0.71}$	68.76	$68.74^{+0.87+1.8}_{-0.87-1.8}$	69.23	$68.82^{+0.94+1.9}_{-0.91-1.9}$	68.20	$68.62^{+0.79+1.7}_{-0.89-1.8}$
σ_8	0.8079	$0.8044^{+0.0045+0.0096}_{-0.0051-0.0091}$	0.7980	$0.7974^{+0.0093+0.018}_{-0.0090-0.017}$	0.7958	$0.7978^{+0.0096+0.019}_{-0.0092-0.019}$	0.7927	$0.7956^{+0.0097+0.019}_{-0.0092-0.019}$




Figure 5: The one-dimensional posterior distribution and two-dimensional posterior contours with 68% and 95% confidence limits for the selected cosmological parameters of Barrow cosmology from dataset (I) (blue) and dataset (II) (orange) (left panel), and also for the TMG model from dataset (II) according to Ref. [76] (right panel)

Moreover, w CDM describes a phantom dark energy equation of state which is consistent with Barrow cosmological model.

On the other hand, numerical results based on dataset (II) indicate that the quintessential nature of dark energy can slightly alleviate the σ_8 tension in Barrow cosmology compared to the Λ CDM model, while there is a satisfactory agreement between the Barrow model and w CDM.

In general, our MCMC investigation based on both datasets shows compatibility between Barrow cosmology and the reference models (Λ CDM and w CDM). Further, the correlation between σ_8 and H_0 shows that a full reconciliation between local and global observations is not possible.

For the last point, it is interesting to compare the obtained constraints on Barrow cosmology with the results based on Tsallis cosmology reported in [76]. Considering both Barrow and Tsallis scenarios, there is a slight alleviation in σ_8 tension with a quintessential behaviour of dark energy according to dataset (II), which is more significant in the TMG model. Also, the quintessential character of dark energy in Barrow and Tsallis cosmologies is in reasonable agreement with the w CDM model. However, considering physical principles and the motivation of correction, Barrow cosmology and the TMG model describe two different corrections to the area law of entropy.

Acknowledgements We thank the Shiraz University Research Council. We are also grateful to an anonymous referee for valuable comments which helped us improve the paper significantly.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: Observational data we have used in this paper are publicly available in Refs. [82, 88–96].

References

- [1] J. M. Bardeen, B. Carter, and S. W. Hawking. The four laws of black hole mechanics. *Comm. Math. Phys.*, 31(2):161–170, 1973.
- [2] Jacob D. Bekenstein. Black holes and entropy. *Phys. Rev. D*, 7:2333–2346, Apr 1973.
- [3] S. W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43:199–220, Aug 1975.
- [4] Ted Jacobson. Thermodynamics of spacetime: The Einstein equation of state. *Phys. Rev. Lett.*, 75:1260–1263, Aug 1995.
- [5] T Padmanabhan. Classical and quantum thermodynamics of horizons in spherically symmetric spacetimes. *Classical and Quantum Gravity*, 19(21):5387–5408, oct 2002.
- [6] T. Padmanabhan. Gravity and the thermodynamics of horizons. *Physics Reports*, 406(2):49 – 125, 2005.
- [7] Christopher Eling, Raf Guedens, and Ted Jacobson. Nonequilibrium thermodynamics of spacetime. *Phys. Rev. Lett.*, 96:121301, Mar 2006.
- [8] M. Akbar and Rong-Gen Cai. Friedmann equations of FRW universe in scalar-tensor gravity, $f(R)$ gravity and first law of thermodynamics. *Physics Letters B*, 635(1):7 – 10, 2006.
- [9] Aseem Paranjape, Sudipta Sarkar, and T. Padmanabhan. Thermodynamic route to field equations in Lanczos-Lovelock gravity. *Phys. Rev. D*, 74:104015, Nov 2006.
- [10] T. Padmanabhan and Aseem Paranjape. Entropy of null surfaces and dynamics of spacetime. *Phys. Rev. D*, 75:064004, Mar 2007.
- [11] M. Akbar and Rong-Gen Cai. Thermodynamic behavior of field equations for $f(R)$ gravity. *Physics Letters B*, 648(2):243 – 248, 2007.
- [12] Dawood Kothawala, Sudipta Sarkar, and T. Padmanabhan. Einstein's equations as a thermodynamic identity: The cases of stationary axisymmetric horizons and evolving spherically symmetric horizons. *Physics Letters B*, 652(5):338 – 342, 2007.
- [13] T Padmanabhan. Thermodynamical aspects of gravity: new insights. *Reports on Progress in Physics*, 73(4):046901, mar 2010.
- [14] Erik Verlinde. On the holographic principle in a radiation dominated universe. arXiv:hep-th/0008140, Aug 2000.
- [15] Bin Wang, Elcio Abdalla, and Ru-Keng Su. Relating friedmann equation to Cardy formula in universes with cosmological constant. *Physics Letters B*, 503(3):394 – 398, 2001.
- [16] Andrei V Frolov and Lev Kofman. Inflation and de Sitter thermodynamics. *Journal of Cosmology and Astroparticle Physics*, 2003(05):009–009, may 2003.
- [17] Ulf H. Danielsson. Transplanckian energy production and slow roll inflation. *Phys. Rev. D*, 71:023516, Jan 2005.
- [18] Rong-Gen Cai and Sang Pyo Kim. First law of thermodynamics and friedmann equations of Friedmann–Robertson–Walker universe. *Journal of High Energy Physics*, 2005(02):050–050, feb 2005.
- [19] Raphael Bousso. Cosmology and the S matrix. *Phys. Rev. D*, 71:064024, Mar 2005.
- [20] M. Akbar and Rong-Gen Cai. Thermodynamic behavior of the Friedmann equation at the apparent horizon of the FRW universe. *Phys. Rev. D*, 75:084003, Apr 2007.
- [21] Ahmad Sheykhi. Thermodynamics of apparent horizon and modified Friedmann equations. *The European Physical Journal C*, 69:265, Sep 2010.
- [22] Gianluca Calcagni. de Sitter thermodynamics and the braneworld. *Journal of High Energy Physics*, 2005(09):060–060, sep 2005.
- [23] Ahmad Sheykhi, Bin Wang, and Rong-Gen Cai. Deep connection between thermodynamics and gravity in Gauss-Bonnet braneworlds. *Phys. Rev. D*, 76:023515, Jul 2007.
- [24] Ahmad Sheykhi, Bin Wang, and Rong-Gen Cai. Thermodynamical properties of apparent horizon in warped DGP braneworld. *Nuclear Physics B*, 779(1):1 – 12, 2007.

- [25] Ahmad Sheykhi and Bin Wang. Generalized second law of thermodynamics in Gauss-Bonnet braneworld. *Physics Letters B*, 678(5):434 – 437, 2009.
- [26] S. W. Hawking. Black hole explosions? *Nature*, 248(5443):30–31, March 1974.
- [27] Carlo Rovelli. Black hole entropy from loop quantum gravity. *Phys. Rev. Lett.*, 77:3288–3291, Oct 1996.
- [28] Robert B. Mann and Sergey N. Solodukhin. Quantum scalar field on a three-dimensional (BTZ) black hole instanton: Heat kernel, effective action, and thermodynamics. *Phys. Rev. D*, 55:3622–3632, Mar 1997.
- [29] A. Ashtekar, J. Baez, A. Corichi, and K. Krasnov. Quantum geometry and black hole entropy. *Phys. Rev. Lett.*, 80:904–907, Feb 1998.
- [30] Romesh K. Kaul and Parthasarathi Majumdar. Logarithmic correction to the Bekenstein-Hawking entropy. *Phys. Rev. Lett.*, 84:5255–5257, Jun 2000.
- [31] Saurya Das, Parthasarathi Majumdar, and Rajat K Bhaduri. General logarithmic corrections to black-hole entropy. *Classical and Quantum Gravity*, 19(9):2355–2367, apr 2002.
- [32] Rabin Banerjee and Bibhas Ranjan Majhi. Quantum tunneling and back reaction. *Physics Letters B*, 662(1):62 – 65, 2008.
- [33] Rabin Banerjee and Bibhas Ranjan Majhi. Quantum tunneling beyond semiclassical approximation. *Journal of High Energy Physics*, 2008(06):095–095, jun 2008.
- [34] Jingyi Zhang. Black hole quantum tunnelling and black hole entropy correction. *Physics Letters B*, 668(5):353 – 356, 2008.
- [35] Saurya Das, S. Shankaranarayanan, and Sourav Sur. Power-law corrections to entanglement entropy of horizons. *Phys. Rev. D*, 77:064013, Mar 2008.
- [36] Saurya Das, S. Shankaranarayanan, and Sourav Sur. Black hole entropy from entanglement: A review. arXiv:0806.0402, 2008.
- [37] Ninfa Radicella and Diego Pavón. The generalized second law in universes with quantum corrected entropy relations. *Physics Letters B*, 691(3):121 – 126, 2010.
- [38] Josiah Willard Gibbs. *Elementary Principles in Statistical Mechanics: Developed with Especial Reference to the Rational Foundation of Thermodynamics*. Cambridge Library Collection - Mathematics. Cambridge University Press, 2010.
- [39] Constantino Tsallis. Possible generalization of Boltzmann-Gibbs statistics. *Journal of Statistical Physics*, 52:479, 1988.
- [40] M. L. Lyra and C. Tsallis. Nonextensivity and multifractality in low-dimensional dissipative systems. *Phys. Rev. Lett.*, 80:53–56, Jan 1998.
- [41] Constantino Tsallis, Renio S. Mendes, and A.R. Plastino. The role of constraints within generalized nonextensive statistics. *Physica A: Statistical Mechanics and its Applications*, 261(3):534 – 554, 1998.
- [42] Constantino Tsallis. The nonadditive entropy s_q and its applications in physics and elsewhere: Some remarks. *Entropy*, 13(10):1765–1804, Sep 2011.
- [43] Constantino Tsallis. From nonlinear statistical mechanics to nonlinear quantum mechanics - concepts and applications. arXiv:1202.3178, 2012.
- [44] Rafael da C. Nunes, Edésio M. Barboza Jr., Everton M. C. Abreu, and Jorge Ananias Neto. Dark energy models through nonextensive Tsallis’ statistics. arXiv:1403.5706, 2014.
- [45] Constantino Tsallis and Leonardo J. L. Cirto. Black hole thermodynamical entropy. *The European Physical Journal C*, 73(7):2487, Jul 2013.
- [46] Ahmad Sheykhi. Modified friedmann equations from Tsallis entropy. *Physics Letters B*, 785:118 – 126, 2018.
- [47] A. Sayahian Jahromi, S. A. Moosavi, H. Moradpour, J. P. Morais Graça, I. P. Lobo, I. G. Salako, and A. Jawad. Generalized entropy formalism and a new holographic dark energy model. *Physics Letters B*, 780:21–24, May 2018.
- [48] M. Abdollahi Zadeh, A. Sheykhi, H. Moradpour, and Kazuharu Bamba. Note on tsallis holographic dark energy. *The European Physical Journal C*, 78(11):940, Nov 2018.
- [49] Emmanuel N. Saridakis, Kazuharu Bamba, R. Myrzakulov, and Fotios K. Anagnostopoulos. Holographic dark energy through tsallis entropy. *Journal of Cosmology and Astroparticle Physics*, 2018(12):012–012, Dec 2018.
- [50] Ehsan Sadri. Observational constraints on interacting tsallis holographic dark energy model. *The European Physical Journal C*, 79(9):762, Sep 2019.
- [51] Ahmad Sheykhi. New explanation for accelerated expansion and flat galactic rotation curves. *The European Physical Journal C*, 80:25, 2020.
- [52] Abdulla Al Mamon, Amir Hadi Ziaie, and Kazuharu Bamba. A generalized interacting tsallis holographic dark energy model and its thermodynamic implications. *The European Physical Journal C*, 80(10):974, Oct 2020.
- [53] Anish Ghoshal and Gaetano Lambiase. Constraints on Tsallis cosmology from big bang nucleosynthesis and dark matter freeze-out. arXiv:2104.11296, 2021.
- [54] W. J. C. da Silva and R. Silva. Cosmological perturbations in the tsallis holographic dark energy scenarios. *The European Physical Journal Plus*, 136(5):543, May 2021.
- [55] Shin’ichi Nojiri, S. D. Odintsov, and Tanmoy Paul. Different faces of generalized holographic dark energy. arXiv:2105.08438, 2021.
- [56] John D. Barrow. The area of a rough black hole. *Physics Letters B*, 808:135643, 2020.
- [57] S.W. Hawking. Spacetime foam. *Nuclear Physics B*, 144(2):349–362, 1978.
- [58] Gerard ’t Hooft. On the quantum structure of a black hole. *Nuclear Physics B*, 256:727–745, 1985.

- [59] Everton M. C. Abreu, Jorge Ananias Neto, and Edésio M. Barboza. Barrow's black hole entropy and the equipartition theorem. *EPL (Europhysics Letters)*, 130(4):40005, Jun 2020.
- [60] Emmanuel N. Saridakis. Modified cosmology through spacetime thermodynamics and Barrow horizon entropy. *Journal of Cosmology and Astroparticle Physics*, 2020(07):031–031, Jul 2020.
- [61] Fotios K. Anagnostopoulos, Spyros Basilakos, and Emmanuel N. Saridakis. Observational constraints on Barrow holographic dark energy. *The European Physical Journal C*, 80:826, Sep 2020.
- [62] Everton M.C. Abreu and Jorge Ananias Neto. Barrow black hole corrected-entropy model and Tsallis nonextensivity. *Physics Letters B*, 810:135805, 2020.
- [63] Biswajit Das and Biswajit Pandey. A study of holographic dark energy models using configuration entropy. arXiv:2011.07337, 2020.
- [64] Emmanuel N. Saridakis. Barrow holographic dark energy. *Phys. Rev. D*, 102:123525, Dec 2020.
- [65] S. Srivastava and U. K. Sharma. Barrow holographic dark energy with hubble horizon as IR cutoff. *International Journal of Geometric Methods in Modern Physics*, 18:2150014, 2021.
- [66] U. K. Sharma, G. Varshney, and V. C. Dubey. Barrow age-graphic dark energy. *International Journal of Modern Physics D*, 30:2150021, Jan 2021.
- [67] A.A. Mamon, A. Paliathanasis, and S. Saha. Dynamics of an interacting Barrow holographic dark energy model and its thermodynamic implications. *Eur. Phys. J. Plus*, 136:134, Jan 2021.
- [68] John D. Barrow, Spyros Basilakos, and Emmanuel N. Saridakis. Big bang nucleosynthesis constraints on Barrow entropy. *Physics Letters B*, 815:136134, 2021.
- [69] A. Pradhan, A. Dixit, and V. K. Bhardwaj. Barrow HDE model for statefinder diagnostic in FLRW universe. *International Journal of Modern Physics A*, 36:2150030, Feb 2021.
- [70] Archana Dixit, Vinod Kumar Bharadwaj, and Anirudh Pradhan. Barrow HDE model for statefinder diagnostic in non-flat FRW universe. arXiv:2103.08339, Mar 2021.
- [71] Priyanka Adhikary, Sudipta Das, Spyros Basilakos, and Emmanuel N. Saridakis. Barrow holographic dark energy in non-flat universe. arXiv:2104.13118, Apr 2021.
- [72] Vinod Kumar Bhardwaj, Archana Dixit, and Anirudh Pradhan. Statefinder hierarchy model for the barrow holographic dark energy. *New Astronomy*, 88:101623, 2021.
- [73] Ahmad Sheykhi. Barrow entropy corrections to friedmann equations. *Phys. Rev. D*, 103:123503, Jun 2021.
- [74] Emmanuel N. Saridakis and Spyros Basilakos. The generalized second law of thermodynamics with barrow entropy. *The European Physical Journal C*, 81:644, 2021.
- [75] Genly Leon, Juan Magaña, A. Hernández-Almada, Miguel A. García-Aspeitia, Tomás Verdugo, and V. Motta. Barrow entropy cosmology: an observational approach with a hint of stability analysis. arXiv:2108.10998, 2021.
- [76] Mahnaz Asghari and Ahmad Sheykhi. Observational constraints on Tsallis modified gravity. *Monthly Notices of the Royal Astronomical Society*, 09 2021. stab2671.
- [77] Adam G. Riess et al. A 2.4% DETERMINATION OF THE LOCAL VALUE OF THE HUBBLE CONSTANT. *The Astrophysical Journal*, 826(1):56, Jul 2016.
- [78] Adam G. Riess et al. New parallaxes of galactic cepheids from spatially scanning the Hubble space telescope: Implications for the Hubble constant. *The Astrophysical Journal*, 855(2):136, mar 2018.
- [79] Adam G. Riess, Stefano Casertano, Wenlong Yuan, Lucas M. Macri, and Dan Scolnic. Large magellanic cloud cepheid standards provide a 1% foundation for the determination of the Hubble constant and stronger evidence for physics beyond Λ CDM. *The Astrophysical Journal*, 876(1):85, May 2019.
- [80] Adam G. Riess, Stefano Casertano, Wenlong Yuan, J. Bradley Bowers, Lucas Macri, Joel C. Zinn, and Dan Scolnic. Cosmic distances calibrated to 1% precision with Gaia EDR3 parallaxes and Hubble space telescope photometry of 75 milky way cepheids confirm tension with Λ CDM. *The Astrophysical Journal*, 908(1):L6, Feb 2021.
- [81] S. W. Allen, R. W. Schmidt, A. C. Fabian, and H. Ebeling. Cosmological constraints from the local X-ray luminosity function of the most X-ray-luminous galaxy clusters. *Monthly Notices of the Royal Astronomical Society*, 342(1):287, 2003.
- [82] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641:A6, Sep 2020.
- [83] Dongsu Bak and Soo-Jong Rey. Cosmic holography. *Classical and Quantum Gravity*, 17(15):L83–L89, jul 2000.
- [84] S.R. De Groot and P. Mazur. *Non-equilibrium Thermodynamics*. Dover books on physics and chemistry. North-Holland Publishing Company, 1962.
- [85] Diego Blas, Julien Lesgourgues, and Thomas Tram. The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes. *Journal of Cosmology and Astroparticle Physics*, 2011(07):034, 2011.
- [86] Benjamin Audren, Julien Lesgourgues, Karim Benabed, and Simon Prunet. Conservative Constraints on Early Cosmology: an illustration of the MONTE PYTHON cosmological parameter inference code. *JCAP*, 1302:001, 2013.
- [87] Thejs Brinckmann and Julien Lesgourgues. MontePython 3: boosted MCMC sampler and other features. arXiv:1804.07261, 2018.
- [88] P. A. R. Ade et al. Planck 2015 results. XXIV. Cosmology from Sunyaev-Zeldovich cluster counts. *Astron. Astrophys.*, 594:A24, September 2016.
- [89] P. A. R. Ade et al. Planck 2013 results. XX. Cosmology from Sunyaev-Zeldovich cluster counts. *Astron. Astrophys.*, 571:A20, November 2014.

- [90] Martin Kilbinger et al. CFHTLenS: Combined probe cosmological model comparison using 2D weak gravitational lensing. *Monthly Notices of the Royal Astronomical Society*, 430(3):2200, 2013.
- [91] Catherine Heymans et al. CFHTLenS tomographic weak lensing cosmological parameter constraints: Mitigating the impact of intrinsic galaxy alignments. *Monthly Notices of the Royal Astronomical Society*, 432(3):2433, 2013.
- [92] D. M. Scolnic et al. The complete light-curve sample of spectroscopically confirmed SNe Ia from pan-STARRS1 and cosmological constraints from the combined pantheon sample. *The Astrophysical Journal*, 859(2):101, May 2018.
- [93] Florian Beutler, Chris Blake, Matthew Colless, D. Heath Jones, Lister Staveley-Smith, Lachlan Campbell, Quentin Parker, Will Saunders, and Fred Watson. The 6DF Galaxy Survey: Baryon Acoustic Oscillations and the Local Hubble Constant. *Monthly Notices of the Royal Astronomical Society*, 416(4):3017–3032, 2011.
- [94] Lauren Anderson et al. The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations in the Data Releases 10 and 11 Galaxy Samples. *Monthly Notices of the Royal Astronomical Society*, 441(1):24–62, 2014.
- [95] Shadab Alam et al. The clustering of galaxies in the completed SDSS-III baryon oscillation spectroscopic survey: cosmological analysis of the DR12 galaxy sample. *Monthly Notices of the Royal Astronomical Society*, 470(3):2617–2652, 2017.
- [96] Manuel A. Buen-Abad, Martin Schmaltz, Julien Lesgourgues, and Thejs Brinckmann. Interacting Dark Sector and Precision Cosmology. *JCAP*, 1801(01):008, 2018.
- [97] Hirotugu Akaike. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19(6):716–723, December 1974.
- [98] K.P. Burnham and D.R. Anderson. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. Springer Verlag, 2002.