

Construction of meta-GGA functionals through restoration of exact constraint adherence to regularized SCAN functionals.

James W. Furness,^{1,*} Aaron D. Kaplan,² Jinliang Ning,¹ John P. Perdew,^{2,3} and Jianwei Sun^{1,†}

¹*Department of Physics and Engineering Physics, Tulane University, New Orleans, LA 70118*

²*Department of Physics, Temple University, Philadelphia, PA 19122*

³*Department of Chemistry, Temple University, Philadelphia, PA 19122*

(Dated: October 5, 2021)

The SCAN meta-GGA exchange-correlation functional [Phys. Rev. Lett. **115**, 036402 (2015)] is constructed as a chemical environment-determined interpolation between two separate energy densities: one describes single orbital electron densities accurately, and another describes slowly-varying densities accurately. To conserve constraints known for the exact exchange-correlation functional, the derivatives of this interpolation vanish in the slowly-varying limit. While theoretically convenient, this choice introduces numerical challenges that degrade the functional’s efficiency. We have recently reported a modification to the SCAN functional, termed r^2 SCAN [J. Phys. Chem. Lett. **11**, 8208 (2020)] that introduces two regularizations into SCAN which improve its numerical performance at the expense of not recovering the fourth order term of the slowly-varying density gradient expansion for exchange. Here we show the derivation of a progression of functionals (r SCAN, r^{++} SCAN, r^2 SCAN, and r^4 SCAN) with increasing adherence to exact conditions while maintaining a smooth interpolation. The greater smoothness of r^2 SCAN seems to lead to better general accuracy than the additional exact constraint of SCAN or r^4 SCAN does.

I. INTRODUCTION

The importance of efficient computational modeling in chemistry and materials science cannot be understated, and for many applications, Kohn–Sham density functional theory presents the most appealing compromise between accuracy and efficiency. The favorable position of this compromise has been enabled by the steady progression of ever more accurate density functionals produced over the last 60 years. These functionals are commonly characterized by the Perdew–Schmidt hierarchy [1], a progression of increasing non-locality where successive levels can be expected to give greater accuracy at the cost of increased computational complexity.

The meta-generalized gradient approximations (meta-GGAs) stand as an appealing level at which the highest accuracy can be expected from semi-local ingredients, including the electron density, its gradient, and the kinetic energy density. While hybrid functionals incorporating admixtures of non-local exact exchange have become most prominent for molecular applications, the prohibitive cost scaling of exact-exchange with number of electrons has limited their utility for extended systems.

A meta-GGA is commonly designed by either enforcing constraints on the exchange-correlation (XC) functional, or by fitting to reference data sets. Those that take the latter route, called empirical functionals, can be inaccurate for systems outside their respective fitting set, or can suffer from difficulties due to over-fitting. General purpose functionals that are accurate for diverse systems have tended to be of the former, so-called non-

empirical, variety in which transferable accuracy is promoted by adherence to physical constraints that are necessarily true for all systems of electrons. The first generation of meta-GGAs were non-empirical, and predate most GGAs. Becke and Roussel [2, 3] derived generalized Taylor series of the exact-exchange hole by enforcing sum rule and non-negativity constraints on a hole model. Perdew [4] derived a Laplacian-level meta-GGA for the exchange energy by enforcing the same set of constraints.

At the meta-GGA level, the strongly-constrained and appropriately-normed (SCAN) functional has incorporated all 17 known constraints on the exact XC energy appropriate to the semi-local level [5]. (These constraints are listed together in the Supplementary Material of Ref. [5], and the references for them are presented in the main text of Ref. [5].) From this foundation, further works have proposed modifications to the SCAN energy densities to improve its accuracy in certain domains. revSCAN is a simple modification to the slowly-varying limit of SCAN’s correlation energy that modifies the fourth-order term in its density-gradient expansion [6]. The TASK functional is a complete revision of SCAN, retaining only its fulfillment of exact constraints for the exchange energy [7]. TASK is designed to accurately predict band gaps, and has recently been itself extended for accuracy in 2D systems, in a modification termed “mTASK” [8]. Note TASK and mTASK use a correlation density functional at the local density approximation.

The SCAN functional has proved broadly transferable and has shown good accuracy for many systems normally challenging for DFT methods [9–21], though its numerical difficulties have hindered some applications such as pseudo-potential generation [22, 23]. To address this, Bartók and Yates proposed a regularized SCAN, termed “ r SCAN”, that aims to control numerical challenges while remaining as close to the original SCAN

* jfurness@tulane.edu

† jsun@tulane.edu

functional as possible. While the regularizations are effective in improving numerical performance, removing the grid sensitivity of SCAN that is problematic in some electronic structure codes, they break five of the exact conditions SCAN was designed to obey. Recent work by Mejía-Rodríguez and Trickey shows that some transferability is lost in rSCAN, with atomization energies particularly degraded [24, 25]. We have recently proposed a restored-regularized-SCAN, called r²SCAN, which maintains the regularizations of rSCAN while restoring exact constraint adherence [26]. Compared to SCAN, the resulting r²SCAN functional has shown pronounced improvements in numerical efficiency, alongside small systematic improvements in accuracy [21, 26–30]. On the extensive GMTKN55 test set [31] for main-group chemistry, the overall error measure WTMAD-2 was [29] 8.6 kcal/mol for SCAN+D4 and 7.5 kcal/mol for r²SCAN+D4, where D4 is a dispersion correction. Note that SCAN and r²SCAN are not fitted to any bonded system, but are genuinely predictive for bonded systems. Applying SCAN+D4 to the unrestricted Hartree-Fock density [32] instead of its own self-consistent density leads to a remarkably small WTMAD-2 of 5.079 kcal/mol, better than nearly all the (necessarily empirical) hybrid functionals tested thus far. This “density correction” [33] to SCAN also leads to a nearly-perfect many-body expansion and molecular dynamics for water [34].

The present publication completes Ref [26], providing the necessary detail of how each exact constraint was restored in r²SCAN. We show how these restorations affect the numerical performance of the functional and how smoothness can be maintained for all constraints except the fourth order term of the slowly-varying density gradient expansion for exchange, which is less easy to enforce following the present interpolation-based model. This work also provides context for the exact constraints enforced by SCAN, and demonstrates how a meta-GGA can be constructed to enforce those constraints.

This work builds upon Ref. [26] by expanding it to a progression of functionals (rSCAN, r++SCAN, r²SCAN, and r⁴SCAN) that ultimately restore all the exact constraints obeyed by SCAN to a regularized form. Thus our presentation expands upon and completes the letter version of Ref. [26], by filling in the details in the constructions of the last three functionals, and by presenting numerical results for r++SCAN and r⁴SCAN. We also present individual errors of these functionals on their appropriate norms (used to determine their parameters) in Table I, and on the lattice constants of solids in Table IV. Figures 3, 5, 6, and 7 will be familiar to readers of Ref [26], but are expanded to incorporate results for the new meta-GGAs. We also include a more detailed analysis of the construction of r²SCAN than was presented in Ref. [26]: in addition to a derivation of the r²SCAN gradient expansion in supplemental material A, Fig. 4 shows how the damping factor d_{p_2} of r²SCAN was determined. Variations of Figs. 1 and 2 were presented in Ref. [26]; they are included here for completeness. The

Tables in Appendices C–F report results for the same test sets considered in Ref. [26], but including the novel meta-GGAs.

II. CONSTRAINT RESTORATION

A. Coordinate scaling and uniform density limit

The SCAN functional is comprised of independent exchange and correlation functionals each constructed as an interpolation and extrapolation of two semi-local energy densities: one for single-orbital densities $\epsilon_{x/c}^0$, and one for slowly-varying densities $\epsilon_{x/c}^1$, where “x/c” is either exchange or correlation, respectively. Here, we will use ϵ to refer to the energy density, and $\varepsilon = \epsilon/n$ to refer to the energy per electron. The single-orbital and slowly-varying energy densities are joined by way of an interpolation function,

$$\epsilon_{x/c}^{\text{SCAN}}(\mathbf{r}) = \epsilon_{x/c}^1(\mathbf{r}) + f_{x/c}(\alpha(\mathbf{r})) \left[\epsilon_{x/c}^0(\mathbf{r}) - \epsilon_{x/c}^1(\mathbf{r}) \right], \quad (1)$$

which is controlled by the iso-orbital indicator variable,

$$\alpha = \frac{\tau - \tau_W}{\tau_U}. \quad (2)$$

α is built from three kinetic energy densities: the positive-definite conventional $\tau = 1/2 \sum_i^{\text{occ.}} |\nabla \phi_i|^2$ defined with the occupied Kohn-Sham orbitals $\{\phi_i\}$, von Weizsäcker $\tau_W = |\nabla n|^2/(8n)$ that is the single-orbital limit of τ as a function of the electron density $n = \sum_i^{\text{occ.}} |\phi_i|^2$, and $\tau_U = \frac{3}{10} k_F^2 n d_s(\zeta)$ the uniform electron gas limit of τ . Here, the Fermi wavevector $k_F = [3\pi^2 n]^{1/3}$, and $d_s(\zeta) = [(1 + \zeta)^{5/3} + (1 - \zeta)^{5/3}]/2$ is a function of the spin polarization $\zeta = (n_\uparrow - n_\downarrow)/(n_\uparrow + n_\downarrow)$. We refer to Refs. [5, 23, 35–39] for a detailed discussion of the properties of α and related quantities.

The first change made in rSCAN [22] is to regularize the iso-orbital indicator α to prevent divergence of the XC potential in the asymptotic regions of single orbital systems [23]. In this region, the derivative $\partial\alpha/\partial\tau$ diverges faster than the decay of $\epsilon_x^{\text{LDA}} = -3k_F n/(4\pi)$ (the local density approximation for exchange), resulting in a diverging exchange-correlation potential when $\partial\epsilon_{xc}/\partial\alpha \neq 0$ [23], as is the case for SCAN. This diverging potential is problematic for pseudo-potential generation [22, 23] and is avoided in rSCAN using a regularized α' ,

$$\tilde{\tau}_U = \left(\frac{3}{10} (3\pi^2)^{2/3} n^{5/3} + \tau_r \right) d_s(\zeta), \quad (3)$$

$$\tilde{\alpha} = \frac{\tau - \tau_W}{\tilde{\tau}_U}, \quad (4)$$

$$\alpha' = \frac{\tilde{\alpha}^3}{\tilde{\alpha}^2 + \alpha_r}. \quad (5)$$

The regularizing constants are $\tau_r = 10^{-4}$ and $\alpha_r = 10^{-3}$. Whilst successful in preventing the rSCAN potential from

diverging, the α' regularization breaks two exact constraints: the 1) uniform density limit, and 2) the uniform coordinate scaling of the exchange energy [40].

The exact uniform density limit is recovered in SCAN by recognizing,

$$\lim_{|\nabla n| \rightarrow 0} \tau = \lim_{|\nabla n| \rightarrow 0} \tau_U, \quad (6)$$

and,

$$\lim_{|\nabla n| \rightarrow 0} \tau_W = 0, \quad (7)$$

hence,

$$\lim_{|\nabla n| \rightarrow 0} \alpha = 1. \quad (8)$$

Then by construction,

$$\lim_{|\nabla n| \rightarrow 0} f_{x/c}^{\text{SCAN}}(\alpha) = 0, \quad (9)$$

and Eq. 1 exclusively selects ϵ_x^1 and ϵ_c^1 in the uniform density limit. These energy densities satisfy the uniform density limit by design.

The uniform density limit is broken by the regularization parameters in α' as,

$$\lim_{|\nabla n| \rightarrow 0} \tilde{\tau}_U \neq \lim_{|\nabla n| \rightarrow 0} \tau, \quad (10)$$

$$\lim_{|\nabla n| \rightarrow 0} \tilde{\alpha} = \frac{\lim_{|\nabla n| \rightarrow 0} \tau_U}{\lim_{|\nabla n| \rightarrow 0} \tilde{\tau}_U} \neq 1, \quad (11)$$

$$\lim_{|\nabla n| \rightarrow 0} \alpha' \neq 1, \quad (12)$$

hence,

$$\lim_{|\nabla n| \rightarrow 0} f_{x/c}^{\text{rSCAN}}(\alpha') \neq 0. \quad (13)$$

The final inequality results in a slight scaling of $\epsilon_{x/c}^1$ and a small inclusion of $\epsilon_{x/c}^0$, which does not recover the uniform density limit. Hence the constraint is broken. The uniform density limit is important for metallic elements. For a uniform electron gas of density parameter $r_s = 4$ (roughly characteristic of the valence electron density in solid sodium) $\alpha' \approx 0.719$, for example.

The regularized uniform electron gas kinetic energy density, $\tilde{\tau}_U$, also causes the exchange energy density to scale incorrectly under the uniform coordinate scaling transformations. To see this, we define a uniform coordinate scaling of the density n and Kohn-Sham orbital ϕ_i as

$$n_\lambda(\mathbf{r}) = \lambda^3 n(\lambda\mathbf{r}), \quad (14)$$

$$\phi_{i,\lambda}(\mathbf{r}) = \lambda^{3/2} \phi_i(\lambda\mathbf{r}), \quad (15)$$

with $\lambda \geq 0$, such that the standard meta-GGA variables scale as,

$$\tau_\lambda(\mathbf{r}) = \lambda^5 \frac{1}{2} \sum_i^{\text{occ.}} \left| \frac{\partial \phi_i(\lambda\mathbf{r})}{\partial(\lambda\mathbf{r})} \right|^2 = \lambda^5 \tau(\lambda\mathbf{r}) \quad (16)$$

$$\tau_{W,\lambda}(\mathbf{r}) = \lambda^5 \tau_W(\lambda\mathbf{r}) \quad (17)$$

$$\tau_{U,\lambda}(x, y, z) = \lambda^5 \tau_U(\lambda\mathbf{r}). \quad (18)$$

Thus, while $\alpha(\mathbf{r}) \rightarrow \alpha(\lambda\mathbf{r})$ under uniform coordinate scaling, $\tilde{\alpha}$ does not have this correct behavior except in the limit $\lambda \rightarrow \infty$, because the regularization τ_r in $\tilde{\tau}_U$ doesn't vary with the coordinate scaling parameter λ . This clearly violates the uniform coordinate scaling behavior of the exchange energy [40]

$$E_x[n_\lambda] = \lambda E_x[n], \quad (19)$$

as $\tilde{\alpha}$ does not scale correctly, and the exchange and correlation models here are highly nonlinear in $\tilde{\alpha}$. The exact correlation energy evaluated on a uniformly scaled density tends to distinct limits [41, 42]

$$\lim_{\lambda \rightarrow \infty} E_c[n_\lambda] = \text{constant} \leq 0 \quad (20)$$

$$\lim_{\lambda \rightarrow 0} E_c[n_\lambda] = \lambda D_c[n], \quad (21)$$

where the constant and functional D_c are unknown. It can be seen that SCAN, and the functionals presented here satisfy both limits, but rSCAN satisfies only the $\lambda \rightarrow \infty$ limit.

It should be noted that under nonuniform coordinate scaling, rSCAN does not violate known exact constraints [41, 43, 44] because of the robustness of the underlying SCAN model. It does, however, tend to distinct limits from SCAN, likely impacting performance for real systems. To see this, we define a non-uniform coordinate scaling of the density n and Kohn-Sham orbital ϕ_i in one dimension as,

$$n_\lambda^x(x, y, z) = \lambda n(\lambda x, y, z), \quad (22)$$

$$\phi_{i,\lambda}^x(x, y, z) = \lambda^{1/2} \phi_i(\lambda x, y, z), \quad (23)$$

again with $\lambda \geq 0$. Under this coordinate transformation, the exact exchange energy satisfies [41, 43]

$$\lim_{\lambda \rightarrow 0} \frac{1}{N} E_x[n_\lambda^x] > -\infty \quad (24)$$

$$\lim_{\lambda \rightarrow \infty} \frac{1}{N} E_x[n_\lambda^x] > -\infty, \quad (25)$$

with N the number of electrons. Identical inequalities hold for the exact correlation energy [43, 44]. It should be emphasized that these constraints imply that the exact exchange and correlation energies per electron tend to finite constants under either limit of non-uniform coordinate scaling. The constant limit for Eq. 25 is a non-zero negative constant [44], the exchange energy per electron for a two-dimensional system.

To recover these constraints on an approximate exchange energy functional, the exchange enhancement factor $F_x = \epsilon_x / \epsilon_x^{\text{LDA}}$ must satisfy [45, 46]

$$\lim_{p \rightarrow \infty} F_x \propto p^{-1/4}. \quad (26)$$

$p = [|\nabla n| / (2k_F n)]^2$ is the square of a dimensionless gradient of the density on the appropriate length scale for the exchange energy. We will discuss p further in the

ensuing section on gradient expansions, but it suffices here to note that p scales as $\lambda^{-2/3}$ as $\lambda \rightarrow 0$, and as $\lambda^{4/3}$ as $\lambda \rightarrow \infty$, as shown in supplemental material B. Therefore, p is always divergent under the extreme limits of non-uniform coordinate scaling. In SCAN, rSCAN, and the functionals developed here, the set of coordinate scaling constraints for exchange are imposed through a function $g_x(p)$

$$F_x(n, |\nabla n|, \tau) = \{h_{1x} + f_x(\alpha) [h_{0x} - h_{1x}]\} g_x(p) \quad (27)$$

$$g_x(p) = 1 - \exp[-a_1 p^{-1/4}]. \quad (28)$$

Referring to Eq. (1), it can be seen that $h_{jx} = \epsilon_x^j / [\epsilon_x^{\text{LDA}} g_x]$, with $j = 0, 1$. In the limit $p \rightarrow \infty$,

$$\lim_{p \rightarrow \infty} g_x(p) = a_1 p^{-1/4} + \mathcal{O}(p^{-1/2}) \quad (29)$$

$$\lim_{p \rightarrow \infty} h_{jx} = \text{constant} > 0, \quad j = 0, 1. \quad (30)$$

Thus $F_x \sim p^{-1/4}$. In SCAN, rSCAN, and this work, $h_{x0} = 1.174$ identically, and $h_{x1}(p \rightarrow \infty) = 1.065$. The LDA, most GGAs, and most meta-GGAs do not recover the right asymptotic behavior for exchange.

Recovering the analogous set of non-uniform coordinate scaling constraints for correlation is more straightforward, and requires that, for $j = 0, 1$,

$$\lim_{p \rightarrow \infty} \epsilon_c^j = \text{constant} \leq 0 \quad (31)$$

$$\lim_{r_s \rightarrow 0} \epsilon_c^j = \text{constant} \leq 0, \quad (32)$$

where $r_s = [3/(4\pi n)]^{1/3}$ is the Wigner-Seitz radius. In SCAN, rSCAN, and the functionals developed here, both constants are chosen to be zero. Many non-empirical GGAs and meta-GGAs for correlation satisfy the non-uniform coordinate scaling constraints. LDA, which has a logarithmic divergence as $r_s \rightarrow 0$, does not.

We can now consider the iso-orbital indicators used in SCAN, rSCAN, and r²SCAN. Under the non-uniform coordinate scaling defined in Eqs. 22 and 23, the standard meta-GGA variables scale as

$$\tau_\lambda^x(x, y, z) = \frac{\lambda}{2} \sum_i^{\text{occ.}} \left| \hat{\mathbf{x}} \lambda \frac{\partial \phi_i(\lambda x, y, z)}{\partial(\lambda x)} + \nabla_\perp \phi_i(\lambda x, y, z) \right|^2 \quad (33)$$

$$\tau_{W,\lambda}^x(x, y, z) = \lambda \frac{\left| \hat{\mathbf{x}} \lambda \frac{\partial n(\lambda x, y, z)}{\partial(\lambda x)} + \nabla_\perp n(\lambda x, y, z) \right|^2}{8n(\lambda x, y, z)} \quad (34)$$

$$\tau_{U,\lambda}^x(x, y, z) = \lambda^{5/3} \tau_U(\lambda x, y, z), \quad (35)$$

with

$$\nabla_\perp = \hat{\mathbf{y}} \frac{\partial}{\partial y} + \hat{\mathbf{z}} \frac{\partial}{\partial z}.$$

From these equations, we see that, when $\lambda \rightarrow 0$, τ and τ_W scale as λ , but τ_U scales as $\lambda^{5/3}$. Thus, α scales with a leading order of $\lambda^{-2/3}$ in this limit. When $\lambda \rightarrow \infty$, α

can either scale as $\lambda^{4/3}$ or $\lambda^{-2/3}$. Examples and analysis of both scaling limits are given in supplemental material B.

Due to the τ_r constant in the denominator of $\tilde{\alpha}$ (Eq. 4), the leading order behavior of the regularized $\tilde{\alpha}$ under non-uniform scaling, with $\lambda \rightarrow 0$, is λ . Then, whereas α tends to infinity in the $\lambda \rightarrow 0$ limit, $\tilde{\alpha}$ tends to zero. $\tilde{\alpha}$ has the correct leading-order behavior in the limit $\lambda \rightarrow \infty$, which can be a single-orbital limit where exact constraints, including the finite exchange and correlation energies under the nonuniform coordinate scaling, were built for SCAN.

In Ref. [47], we proposed an alternative regularization of α to restore these constraints,

$$\bar{\alpha} = \frac{\tau - \tau_W}{\tau_U + \eta \tau_W} = \frac{\alpha}{1 + \eta \frac{5}{3} p}, \quad (36)$$

where η is a regularization parameter to be determined later. Clearly, $\bar{\alpha}$ has the correct behavior, $\bar{\alpha}(\mathbf{r}) \rightarrow \bar{\alpha}(\lambda \mathbf{r})$ under uniform coordinate scaling. This regularization eliminates the asymptotic region ($|\mathbf{r}| \rightarrow \infty$) divergence and, by Eq. (7), does not change the uniform density limit,

$$\lim_{|\nabla n| \rightarrow 0} \bar{\alpha} = 1. \quad (37)$$

The nonuniform coordinate scaling of $\bar{\alpha}$ is also maintained as $\lambda^{-2/3}$ to leading order in the $\lambda \rightarrow 0$ limit. But, for $\lambda \rightarrow \infty$, the leading-order term of $\bar{\alpha}$ is independent of λ , for non-homogeneous densities. This is demonstrated in supplemental material B.

Figure 1 shows a comparison of α , α' , and $\bar{\alpha}$ for the krypton atom. The divergence of the conventional α is apparent in the asymptotic region, while α' and $\bar{\alpha}$ decay to 0. Close to the nucleus, the α_r regularization constant causes α' to behave differently to α and $\bar{\alpha}$, but otherwise all three indicators behave similarly.

Substituting $\bar{\alpha}$ for α' in rSCAN is sufficient to restore the uniform density limit and coordinate scaling behaviors, and we refer to rSCAN with this replacement as “r++SCAN” throughout.

B. Gradient expansions

The interpolative design of SCAN allows construction of the single-orbital (ϵ^0) and slowly-varying (ϵ^1) energy densities that consider only the exact constraints relevant to their respective limits. In SCAN, the interpolation function is a piece-wise combination of two exponential terms,

$$f_{x/c}(\alpha) = \begin{cases} \exp\left[\frac{-c_{1x/c}\alpha}{1-\alpha}\right] & \alpha \leq 1 \\ -d_{x/c} \exp\left[\frac{c_{2x/c}}{1-\alpha}\right] & \alpha > 1, \end{cases} \quad (38)$$

where $\{d_{x/c}, c_{1x/c}, c_{2x/c}\}$ are separate parameters for exchange and correlation determined by fitting to appropriate norms [5]. This function was chosen such that:




FIG. 1. Comparison between the conventional α (e.g. from SCAN), α' (from rSCAN), and the new $\tilde{\alpha}$ as a function of distance from the Kr nucleus (in Bohr radii) computed from accurate spherical Hartree–Fock Slater type orbitals [47, 48]. Regularization parameters are $\tau_r = 10^{-4}$ and $\alpha_r = 10^{-3}$ in α' , and $\eta = 10^{-3}$ in $\tilde{\alpha}$.

1) $f(\alpha = 0) = 1$ exclusively selects ϵ^0 for single-orbital densities, 2) $f(\alpha = 1) = 0$ exclusively selects ϵ^1 in slowly-varying densities, and 3)

$$\left. \frac{df(\alpha)}{d\alpha} \right|_{\alpha \rightarrow 1} = \left. \frac{d^2f(\alpha)}{d\alpha^2} \right|_{\alpha \rightarrow 1} = 0, \quad (39)$$

which prevents ϵ^0 contributing to the slowly-varying density gradient expansions to the 4th order in $|\nabla n|$. Note $\left. \frac{d^m f(\alpha)}{d\alpha^m} \right|_{\alpha \rightarrow 1} = 0$ with m to be any integer in Eq. 38 by design. While theoretically convenient, Eq. 38 introduces a twist into the function around $\alpha = 1$, see Figure 2. This twist destroys the overall smoothness of the functional, introduces oscillations into the XC potential [22, 49], and harms its performance on numerical integration grids.

The rSCAN functional uses a smooth polynomial interpolation function in place of the SCAN piece-wise exponential for the range $0 \leq \alpha' < 2.5$,

$$f(\alpha') = \begin{cases} \sum_{i=0}^7 c_i \alpha'^i & 0 \leq \alpha' \leq 2.5 \\ -d_{x/c} \exp\left[\frac{c_{2x/c}}{1-\alpha'}\right] & \alpha' > 2.5 \end{cases}, \quad (40)$$

where $\{c_i\}$ are polynomial coefficients determined to smoothly join $f^{\text{SCAN}}(\alpha')$ at $\alpha' = 0$ and 2.5, see Ref. 22. A comparison of the two interpolation functions is shown in Figure 2. While this replacement smooths the XC energy density and potential, it breaks the third constraint on the interpolation function, and hence ϵ^0 makes spurious contributions to the slowly-varying gradient density expansion of ϵ^1 . Here, we restore the correct expansions by directly subtracting the extra terms that result from breaking the third condition (Eq. (39)) around $p \rightarrow 0$ and $\alpha \rightarrow 1$ where the expansion is relevant.

The exact gradient expansion for exchange around the slowly-varying density limit to the 2nd order (GE2X) and




FIG. 2. The exchange (solid) and correlation (dashed) interpolation functions for the SCAN (blue, Eq. 38) and rSCAN (orange, Eq. 40) functionals.

to the 4th order (GE4X) was derived in terms of the exchange enhancement factor F_x in Refs. [50, 51] as,

$$\lim_{|\nabla n| \rightarrow 0} F_x^{\text{GE}} = 1 + \mu p + \frac{146}{2025} \tilde{q}^2 - \frac{73}{405} p \tilde{q} + \mathcal{O}[|\nabla n|^6], \quad (41)$$

where $\mu = 10/81$ and $\tilde{q} = (9/20)(\tilde{\alpha} - 1) + (\eta 3/4 + 2/3)p$ recovers the reduced density Laplacian at $|\nabla n| \rightarrow 0$. The gradient expansion for correlation was derived to the 2nd order (GE2C) in Ref. [5, 52–54] as,

$$\epsilon_c = \epsilon_c^{\text{LSDA}} + \beta(r_s) \phi(\zeta)^3 t(r_s, \zeta, p)^2 + \mathcal{O}[|\nabla n|^4], \quad (42)$$

where $\phi(\zeta) = [(1 + \zeta)^{2/3} + (1 - \zeta)^{2/3}]/2$ and $t(r_s, \zeta, p) = (3\pi^2/16)^{1/3} \sqrt{p/r_s}/\phi(\zeta)$. We will restore each expansion to r++SCAN in turn.

1. Exchange

The SCAN exchange enhancement factor for a spin-unpolarized system is,

$$F_x^{\text{SCAN}}(p, \alpha) \quad (43)$$

$$= \{h_{1x}(p, \alpha) + f_x(\alpha) [h_{0x} - h_{1x}(p, \alpha)]\} g_x(p),$$

$$g_x(p) = 1 - \exp[-a_1 p^{-1/4}], \quad (44)$$

$$h_{0x} = 1 + \kappa_0 = 1.174, \quad (45)$$

$$h_{1x}(p, \alpha) = 1 + \kappa_1 - \frac{\kappa_1}{1 + \frac{x(p, \alpha)}{\kappa_1}}, \quad (46)$$

$$x(p, \alpha) = \mu p \left[1 + \left(\frac{b_4 p}{\mu} \right) \exp\left(\frac{-|b_4| p}{\mu} \right) \right] + \{b_1 p + b_2(1 - \alpha) \exp[-b_3(1 - \alpha)^2]\}^2, \quad (47)$$

where $\mu = 10/81$ and $\{b_1, b_2, b_3, b_4\}$ are chosen such that SCAN yields GE2X and GE4X, noting that the expansion of $g_x(p)$ around $p = 0$ has only zero-order contributions (see supplemental material A 3).

In rSCAN, the interpolation function derivatives are non-zero at $\bar{\alpha} = 1$, so h_{0x} also contributes to the gradient expansion. This changes the functional's gradient expansion, spoiling the correct gradient expansion of the $x(p, \alpha)$ inherited from SCAN. Thus both rSCAN and r++SCAN fail to recover GE2X and GE4X.

We restore GE2X to give the r²SCAN exchange functional by redesigning $x(p, \alpha)$ as,

$$x(p) = (C_\eta C_{2x} \exp[-p^2/d_{p2}^4] + \mu) p, \quad (48)$$

where C_η and C_{2x} are constants set to cancel spurious contributions from h_{0x} to the 2nd order in ∇n . The restoring constants are multiplied by the damping function $\exp[-p^2/d_{p2}^4]$ to prevent them dominating as p becomes large. The damping parameter d_{p2} derives from scaling the reduced density gradient as $s \rightarrow s/d_{p2}$ with d_{p2} fit to recover the appropriate norms in Section IV.

To find C_η and C_{2x} we take the Taylor expansion of the rSCAN interpolation function (Eq. 40) around $\bar{\alpha} = 1$, noting that $1 - \bar{\alpha}$ is $\mathcal{O}[|\nabla n|^2]$,

$$\begin{aligned} \lim_{|\nabla n| \rightarrow 0} f^{\text{rSCAN}}(\bar{\alpha}) & \quad (49) \\ &= -(1 - \bar{\alpha})\Delta f_2 + \frac{(1 - \bar{\alpha})^2}{2}\Delta f_4 + \mathcal{O}[|\nabla n|^6], \end{aligned}$$

where,

$$\Delta f_2 = \sum_{i=1}^7 i c_i, \quad (50)$$

$$\Delta f_4 = \sum_{i=2}^7 i(i-1)c_i, \quad (51)$$

are determined by the first and second derivatives of the interpolation function with respect to $\bar{\alpha}$ respectively.

The $(1 - \bar{\alpha})$ term of Eq. 49 indicates a fixed slope for the $\bar{\alpha}$ dependence of the enhancement factor across the slowly-varying limit that is found to be numerically problematic, analyzed further in Section III. This can be avoided to second order by expressing $(1 - \bar{\alpha})$ in terms of p through an integration by parts on the exchange energy density [55],

$$\lim_{|\nabla n| \rightarrow 0} (1 - \bar{\alpha}) = \left(\frac{20}{27} + \eta \frac{5}{3} \right) p + \mathcal{O}[|\nabla n|^4]. \quad (52)$$

This substitution, derived and discussed in supplemental material A 2, is used in r²SCAN and identifies,

$$C_\eta = (20/27 + \eta 5/3). \quad (53)$$

To second order the slowly-varying gradient expansion of r²SCAN is then,

$$\lim_{|\nabla n| \rightarrow 0} F_x^{\text{r}^2\text{SCAN}} = \lim_{|\nabla n| \rightarrow 0} h_{1x} - C_\eta p \Delta f_2 \left(h_{0x} - \lim_{|\nabla n| \rightarrow 0} h_{1x} \right). \quad (54)$$

Finding,

$$\lim_{|\nabla n| \rightarrow 0} h_{1x} = 1 + (\mu + C_\eta C_{2x}) p + \mathcal{O}[|\nabla n|^4], \quad (55)$$

and collecting terms gives,

$$\begin{aligned} \lim_{|\nabla n| \rightarrow 0} F_x^{\text{r}^2\text{SCAN}} & \quad (56) \\ &= 1 + \mu p + C_\eta [C_{2x} - \Delta f_2 h_{0x} + \Delta f_2] p + \mathcal{O}[|\nabla n|^4], \end{aligned}$$

equating this to GE2X (second order and below terms of Eq. 41) and solving for C_{2x} gives,

$$C_{2x} = -\Delta f_2 (1 - h_{0x}) \approx -0.162742, \quad (57)$$

as shown in supplemental material A1.

GE4X can be restored to give the ‘‘r⁴SCAN’’ functional by including a further correcting term in the exchange enhancement factor outside the interpolation. This introduces three more constants, derived in supplemental material A 3, for all terms in Eq. 41:

$$\begin{aligned} F_x^{\text{r}^4\text{SCAN}}(p, \bar{\alpha}) &= \{h_{1x}(p) + f_x(\bar{\alpha}) [h_{0x} - h_{1x}(p)] \\ & \quad + \Delta F_4(p, \bar{\alpha})\} g_x(p) \quad (58) \\ \Delta F_4(p, \bar{\alpha}) &= \{C_{2x} [(1 - \bar{\alpha}) - C_\eta p] + C_{\bar{\alpha}\bar{\alpha}} (1 - \bar{\alpha})^2 \\ & \quad + C_{p\bar{\alpha}} p (1 - \bar{\alpha}) + C_{pp} p^2\} \Delta F_4^{\text{damp}}(p, \bar{\alpha}) \quad (59) \end{aligned}$$

$$\Delta F_4^{\text{damp}}(p, \bar{\alpha}) = \frac{2\bar{\alpha}^2}{1 + \bar{\alpha}^4} \exp \left[-\frac{(1 - \bar{\alpha})^2}{d_{\bar{\alpha}4}^2} - \frac{p^2}{d_{p4}^4} \right] \quad (60)$$

$$C_{\bar{\alpha}\bar{\alpha}} = \frac{73}{5000} - \frac{\Delta f_4}{2} [h_{0x} - 1] \approx -0.0593531 \quad (61)$$

$$\begin{aligned} C_{p\bar{\alpha}} &= \frac{511}{13500} - \frac{73}{1500} \eta - \Delta f_2 [C_\eta C_{2x} + \mu] \\ &\approx 0.0402684 \quad (62) \end{aligned}$$

$$\begin{aligned} C_{pp} &= \frac{146}{2025} \left\{ \eta \frac{3}{4} + \frac{2}{3} \right\}^2 - \frac{73}{405} \left\{ \eta \frac{3}{4} + \frac{2}{3} \right\} \\ & \quad + \frac{(C_\eta C_{2x} + \mu)^2}{k_1} \approx -0.0880769. \quad (63) \end{aligned}$$

Further damping functions, ΔF_4^{damp} , are included to prevent the correction terms dominating as $(1 - \bar{\alpha})$ and p become large, introducing $d_{\bar{\alpha}4}$ and d_{p4} as additional parameters. These are again set to recover the appropriate norms in Section IV. For the fourth order expansion, the integration by parts substitution of Eq. 52 cannot be applied, and hence $(1 - \bar{\alpha})$ cannot be removed.

2. Correlation

The SCAN model of the correlation energy per electron $\varepsilon_c = \varepsilon_c/n$ is

$$\varepsilon_c^{\text{SCAN}} = \varepsilon_c^1 + f_c^{\text{SCAN}}(\alpha) [\varepsilon_c^0 - \varepsilon_c^1], \quad (64)$$

with the $\alpha = 0$ correlation energy per electron given by

$$\varepsilon_c^0 = (\varepsilon_c^{\text{LDA0}} + H_c^0) g_c(\zeta), \quad (65)$$

$$\varepsilon_c^{\text{LDA0}} = -\frac{b_{1c}}{1 + b_{2c}\sqrt{r_s} + b_{3c}r_s}, \quad (66)$$

$$H_c^0 = b_{1c} \ln\{1 + w_0[1 - g_\infty(p)]\}, \quad (67)$$

$$w_0 = \exp[-\varepsilon_c^{\text{LDA0}}/b_{1c}] - 1, \quad (68)$$

$$g_\infty(p) = (1 + 4\chi_\infty p)^{-1/4}, \quad (69)$$

$$g_c(\zeta) = \{1 - 2.3631[d_x(\zeta) - 1]\}(1 - \zeta^{12}), \quad (70)$$

with $d_x(\zeta) = [(1 + \zeta)^{4/3} + (1 - \zeta)^{4/3}]/2$.

Similarly, the $\alpha = 1$ limit is given by

$$\varepsilon_c^1 = \varepsilon_c^{\text{LSDA}} + H_1, \quad (71)$$

$$H_1 = \gamma\phi^3 \ln\{1 + w_1[1 - g(y)]\}, \quad (72)$$

$$w_1 = \exp\left[-\frac{\varepsilon_c^{\text{LSDA}}}{\gamma\phi^3}\right] - 1, \quad (73)$$

$$g(y) = (1 + 4y)^{-1/4}, \quad (74)$$

$$y = \frac{\beta(r_s)}{\gamma w_1} t^2, \quad (75)$$

$$\beta(r_s) = \beta_{\text{MB}} \frac{1 + Ar_s}{1 + Br_s}, \quad (76)$$

where $b_{1c} = 0.0285764$, $b_{2c} = 0.0889$, $b_{3c} = 0.125541$, $\chi_\infty = 0.128026$, $\gamma = 0.0310907$, $\beta_{\text{MB}} = 0.066725$, $A = 0.1$, and $B = 0.1778$. $\varepsilon_c^{\text{LSDA}}$ is the local spin-density approximation for correlation from Ref. [56].

As r++SCAN takes the same correlation model, the violation of Eq. 39 by the rSCAN interpolation function breaks GE2C. The GE2C correction terms are restored to r++SCAN by replacing $g(y)$ in ε_c^1 with,

$$g(y, \Delta y) = [1 + 4(y - \Delta y)]^{-1/4}, \quad (77)$$

$$\Delta y = \frac{C_{2c}}{27\gamma d_s(\zeta)\phi^3 w_1} \left\{ 20r_s \left[g_c(\zeta) \frac{\partial \varepsilon_c^{\text{LDA0}}}{\partial r_s} - \frac{\partial \varepsilon_c^{\text{LSDA}}}{\partial r_s} \right] - 45\eta [\varepsilon_c^{\text{LDA0}} g_c(\zeta) - \varepsilon_c^{\text{LSDA}}] \right\} p \exp[-p^2/d_{p2}^4] \quad (78)$$

where the damping function $\exp[-p^2/d_{p2}^4]$ is the same as in Eq. 48. Similarly to exchange, we restore the second order slowly-varying density gradient expansion when,

$$C_{2c} = \Delta f_2 \approx -0.711402. \quad (79)$$

The derivation of this expression is shown in supplemental material A 4.

Making these replacements to r++SCAN gives the ‘‘r²SCAN’’ functional which only breaks GE4X. Including the full correction of Eq. 58 gives the ‘‘r⁴SCAN’’ functional, which obeys all the exact constraints SCAN does. For convenience, a collected definition of the working equations for all new functionals is given in supplemental material C.

3. Summary of Changes

Here, we summarize the changes made from SCAN for each of the functionals.

1. rSCAN replaces α with α' , which contains two regularization parameters, $\tau_r = 10^{-4}$ and $\alpha_r = 10^{-3}$. It also replaces the SCAN interpolation function with a polynomial between $0 \leq \alpha' \leq 2.5$.
2. r++SCAN evolves from rSCAN, by replacing α' with $\bar{\alpha}$ that uses only a single regularization parameter, $\eta = 0.001$.
3. r²SCAN inherits all the changes of r++SCAN. Additionally, for exchange, it replaces $x(p, \alpha)$ in h_{1x} with Eq. 48. For correlation, it replaces $g(y)$ with Eqs. 77 and 78 in ε_c^1 .
4. r⁴SCAN inherits all the changes from r²SCAN. Additionally, for exchange, it replaces F_x with Eq. 58, which introduces ΔF_4 of Eq. 60.

III. NUMERICAL CHALLENGES

The corrections to restore the slowly-varying density gradient expansions for exchange and correlation contain terms linear in $(1 - \bar{\alpha})$. These terms are necessary to restore the 2nd or 4th order gradient expansion, for example, of Eq. 41 for exchange, if the integration in parts is not used as we did for r²SCAN in Eq. 52. These corrections inevitably twist the slope of the interpolation function $f(\bar{\alpha})$ to that of F_x^{GE} with respect to $\bar{\alpha}$ around $\bar{\alpha} \rightarrow 1$ as $|\nabla n| \rightarrow 0$, illustrated in Fig. 3. This introduces oscillations into the derivatives of the enhancement factor with respect to $\bar{\alpha}$, and hence into the overall exchange-correlation potential. Such oscillations are undesirable and reintroduce the numerical problems rSCAN regularizes away. As the gradient expansion constraint requires

$$\left. \frac{\partial F_x}{\partial \bar{\alpha}} \right|_{\bar{\alpha}=1, p=0} \propto \left. \frac{df(\bar{\alpha})}{d\bar{\alpha}} \right|_{\bar{\alpha}=1}, \quad (80)$$

this oscillation in derivatives must be present, at least in the interpolation scheme discussed here, and cannot be removed by damping. Figure 3, compares uncorrected exchange enhancement, $d_{\bar{\alpha}4} \rightarrow 0$, the exchange enhancement with no damping on the correction terms, $d_{\bar{\alpha}4} \rightarrow \infty$, and the $d_{\bar{\alpha}4} = 0.178$ determined in Section IV, showing this effect.

These numerical problems are not present in r²SCAN, as the corrections do not depend upon $(1 - \bar{\alpha})$. Thus, the corresponding oscillation in derivative is avoided, allowing r²SCAN to recover GE2X and GE2C whilst maintaining a smooth potential. As the integration by parts is not possible to fourth order, r⁴SCAN necessarily suffers an oscillatory XC potential in order to recover GE4X.




FIG. 3. a) Exchange enhancement factor for r^4 SCAN as a function of $\bar{\alpha}$ at $p = 0$. The uncorrected enhancement with $d_{\bar{\alpha}4} \rightarrow 0$ (red) is contrasted against the un-damped corrections with $d_{\bar{\alpha}4} \rightarrow \infty$ (black), the proposed r^4 SCAN damping of $d_{\bar{\alpha}4} = 0.178$ (dashed, dark red). b) Derivative of the exchange enhancement with respect to $\bar{\alpha}$ at $p = 0$ for the same conditions.

IV. DETERMINING PARAMETERS

The regularization of the $\bar{\alpha}$ indicator in r^{++} SCAN, r^2 SCAN, and r^4 SCAN is controlled by the parameter η , with larger values increasing regularization strength. We find performance is largely insensitive to η within the range of $0 \leq \eta \leq 0.001$ and take the upper value of $\eta = 0.001$.

We introduce a single damping parameter, d_{p2} , in r^2 SCAN through Eqs. 48 and 78, and set it using the appropriate norms philosophy of the SCAN functional. The parameter was chosen to minimize the sum of the mean absolute percentage errors in XC energy for four rare gas atoms: Ne, Ar, Kr, and Xe (evaluated for spherical Hartree–Fock orbitals [48] relative to [57–59] reference energies), and four jellium surface formation energies with $r_s = 2, 3, 4,$ and 6 bohr (relative to reference energies from Ref. [60]). As the parameters are not fit to any bound systems, we regard the resulting functionals as non-empirical.

Objective error as a function of damping parameter, d_{p2} , is shown in Figure 4. Setting the damping parameter too high degrades accuracy for the rare gas atoms (while mildly improving the jellium surface formation energies) as the gradient expansion terms dominate too far




FIG. 4. The mean absolute percentage error for (left axis, blue) the exchange-correlation energies of Ne, Ar, Kr, Xe [57–59], and (right axis, red) the exchange-correlation jellium surface energy for $r_s = \{2, 3, 4, 6\}$ [60] as a function of second order gradient expansion damping parameter d_{p2} for r^2 SCAN. The optimal value is chosen as the largest for which the rare gas error is $< 0.1\%$ and the jellium error is $< 5\%$.

from $|\nabla n| \rightarrow 0$. Conversely, setting d_{p2} too small degrades accuracy for the jellium surfaces, as the second order gradient expansion is not sufficiently corrected. As a sharper damping function causes sharper features in XC potential, we take the largest value for d_{p2} which meets the accuracy threshold defined by SCAN: a mean absolute percentage error (MAPE) of 0.1% for rare gas XC energies and 5% for jellium surfaces. The optimizing value is found as $d_{p2} = 0.361$.

Two additional parameters are introduced in r^4 SCAN that control damping of the fourth-order gradient expansion terms. These were determined similarly as those which minimize a normalized sum of the rare gas and jellium surface mean absolute percentage errors. The minimizing parameters were found as $d_{\bar{\alpha}4} = 0.178$ and $d_{p4} = 0.802$, as shown in Figure 5.

V. RESULTS

A. Enhancement Factors and Derivatives

An important principle in functional design is to take an ‘‘Occam’s razor’’ approach and determine a functional that is free of twists and kinks. In this way the functional avoids over-fitting to data and ensures smooth functional derivatives that are easy to render on numerical grids.

Figure 6 compares the XC enhancement factors of SCAN, r^2 SCAN, and r^4 SCAN for the Xenon atom. The SCAN enhancement factor shows sharp plateau like regions from the twists in its interpolation function around $\alpha = 1$. The smooth polynomial interpolation function removes these plateaus from r^2 SCAN, though some twists are re-introduced in r^4 SCAN by the $(1 - \bar{\alpha})$ terms in the




FIG. 5. Mean absolute percentage error in rare gas XC energy (upper) and jellium surface exchange-correlation energy (lower) as a function of damping parameters $d_{\bar{\alpha}4}$ and d_{p4} . Optimal parameters are identified by the white circled cross at $d_{p4} = 0.802$, $d_{\bar{\alpha}4} = 0.178$.

GE4X restoration.

The effect of twists in F_{xc} can be seen in the semi-local and non-local XC potential components of the XC potential, shown in Figure 6. The SCAN functional shows sharp oscillations around $\alpha = 1$ points and sharp drops in its non-local component. In contrast, the r^2 SCAN functional is a smooth function of its ingredients, and hence has smooth semi-local and non-local components to its XC potential. While the potential components of r^2 SCAN and r^4 SCAN coincide for much of space they differ significantly around $\bar{\alpha} = 1$ points. Here the $(1 - \bar{\alpha})$ correction terms in r^4 SCAN required for GE4X cause sharp oscillations that return the oscillatory behavior we aim to remove. As these terms cannot be removed by partial integration to the 4th order in ∇n , we therefore conclude that GE4X is incompatible with functional smoothness under the present SCAN interpolation-based model: one must either twist the interpolation function to enforce Eq. 39, or include correcting terms that reintroduces oscillatory factors.




FIG. 6. The XC enhancement factor (top), the multiplicative component of the XC potential (middle), and the non-local component, i.e., the derivative of the XC energy density with respect to the orbital dependent kinetic energy density, τ (bottom) in the Xe atom. Shown for the SCAN, r^2 SCAN, and r^4 SCAN functionals, calculated from reference Hartree–Fock Slater orbitals [47, 48]. Points where $\alpha = 1$ are shown by black vertical lines.

B. Appropriate Norms

An appropriate norm is defined in Ref. [5] as “systems for which semilocal functionals can be exact or extremely accurate”. Here, like SCAN, we take these as the exchange and correlation energies of four rare gas atoms (Ne, Ar, Kr, and Xe), the exchange and correlation surface energies of four jellium slabs ($\bar{r}_s = 2, 3, 4$, and 6), and the interaction energy of Ar_2 at repulsive interatomic distances ($R_{\text{Ar}-\text{Ar}} = 1.6, 1.8$, and 2.0 Å). Table I compares the accuracy of SCAN, rSCAN, r++SCAN, r^2 SCAN, and r^4 SCAN for the appropriate norms.

Care must be taken when computing the jellium surface energy for rSCAN as the uniform bulk density energy is changed by the α' regularization. To evaluate the exchange-correlation contribution to the jellium surface formation energy, we compute [61]

$$\sigma_{xc}(\bar{r}_s) = \int_{-\infty}^{\infty} [\varepsilon_{xc}(n, \nabla n, \tau) - \varepsilon_{xc}(\bar{n}, 0, \bar{\tau}_U)] n(x) dx, \quad (81)$$

where $\bar{r}_s = (4\pi\bar{n}/3)^{-1/3}$ is the density parameter of the corresponding bulk jellium, and $\bar{\tau}_U = 3(3\pi^2)^{2/3}\bar{n}^{5/3}/10$ is its kinetic energy density. Here, we have assumed that the surface lies along the x direction. This ensures that the uniform density limit of a given functional is used, regardless of whether that uniform limit is the LSDA. Building upon our previous example of the valence density in solid sodium: when $r_s = 4$, the rSCAN exchange uniform density limit is

$$\epsilon_x^{\text{rSCAN}}(\bar{n}, 0, \bar{\tau}_U) \Big|_{\bar{r}_s=4} \approx 1.051 \epsilon_x^{\text{LDA}}(\bar{r}_s),$$

making a substantial error over LDA exchange. For $\bar{r}_s = 6$, this error is increased to roughly 14%.

The importance of recovering the second order gradient expansion is seen in the relative accuracy of the functionals for the rare gas atoms. The two functionals which do not recover the gradient expansions (rSCAN and r++SCAN) have MAPEs of $\approx 0.25\%$ whereas the functionals that do have MAPEs of $\approx 0.1\%$. Restoring the fourth order gradient expansion for exchange improves accuracy further, though r²SCAN already has similar accuracy to SCAN for all the appropriate norms suggesting GEX4 is less important for these systems. Outside of these differences, all functionals performed similarly for the appropriate norm systems.

C. Atomization Energies

The work of Ref. [24] shows the performance of rSCAN is relatively poor for atomization energy prediction, as measured by its increased error for the G3 test set of molecules [63]. Table II compares the errors for this test set for all the functionals derived above. Consistent with Ref. [24] we find a large error from rSCAN, with this error only slightly corrected in r++SCAN. As in Ref. 26, restoration of GE2X and GE2C in r²SCAN restores the good accuracy of SCAN, showing the importance of these constraints for atomization energies. The good accuracy of r²SCAN suggests that recovering GE4X is not essential for accurately predicting atomization energies.

The improved numerical performance of the functionals is illustrated by examining the convergence of atomization energy predictions as a function of numerical grid density, as shown in Figure 7. The original SCAN functional shows wild variation with changing grid density, with a range of over 6 kcal/mol in mean absolute error! While there is some indication that a convergence




FIG. 7. Mean absolute percentage error of atomization energies (kcal/mol) for the G3 set of 226 molecules [63] as a function of increasing numerical integration grid density expressed relative to the smallest grid. The grids chosen were defined by the default TURBOMOLE grid levels. The mean number of grid points per atom over the G3 set is about 1,326 for the smallest (gridsize=1) and 53,590 for the largest (gridsize=7) grid shown here.

is approached for the most dense grids, the results from more computationally efficient grids are problematic and clearly untrustworthy.

All four regularized functionals show very fast grid convergence, with all sparse grids giving close agreement to the dense grids. Given the sharp oscillations in Figure 6 and analysis of Section III, it is somewhat unexpected that r⁴SCAN shows good convergence with grid density. We attribute the improved performance over SCAN to the reduction in plateau like behavior of F_{xc} but caution that grid convergence behavior will likely be more system dependent for r⁴SCAN than r++SCAN and r²SCAN as a result of the oscillations.

It is similarly unexpected to find that the accuracy of r²SCAN is degraded by the inclusion of the GE4X in r⁴SCAN. This supports our previous conclusion that GE4X is not important for the properties tested here and elsewhere[29, 30]. We take the mild degradation of G3 accuracy from extending to r⁴SCAN as further evidence that this method of including GE4X into interpolation based meta-GGA functionals is problematic. The inclusion of two additional fitting parameters beyond those in r²SCAN may also contribute.

D. Further Testing

Beyond atomization energies, we have tested the accuracy of the progression of SCAN-like functionals for the interaction energies of closed shell complexes (S22)[64] and reaction barrier heights (BH76)[65], summarized in Table III. Additionally, Table IV summarizes accuracy for the LC20 set of lattice constants for solids [66], obtained by fitting the stabilized jellium equation of state

TABLE I. Accuracy for appropriate norms. Rare gas and jellium surface exchange-correlation energies given in Hartrees (E_h), Ar₂ interaction energies in kcal/mol. Benchmark data for rare gas atom exchange-correlation, jellium surface exchange-correlation, and Ar₂ interaction energy are from Refs. [57–59], [60], and [62] respectively. Error summaries are given as mean absolute percentage errors (MAPE). Full calculation details in main text.

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN	Benchmarks	
Ne	E_x	-12.164 (0.48%)	-12.183 (0.64%)	-12.176 (0.58%)	-12.144 (0.32%)	-12.146 (0.34%)	-12.105 E_h
	E_c	-0.345 (-11.81%)	-0.346 (-11.53%)	-0.347 (-11.36%)	-0.347 (-11.24%)	-0.347 (-11.24%)	-0.391 E_h
	E_{xc}	-12.508 (0.10%)	-12.529 (0.26%)	-12.522 (0.21%)	-12.491 (-0.04%)	-12.493 (-0.03%)	-12.496 E_h
Ar	E_x	-30.264 (0.29%)	-30.295 (0.40%)	-30.281 (0.35%)	-30.182 (0.02%)	-30.196 (0.07%)	-30.175 E_h
	E_c	-0.690 (-4.81%)	-0.695 (-4.24%)	-0.696 (-4.06%)	-0.697 (-3.90%)	-0.697 (-3.90%)	-0.725 E_h
	E_{xc}	-30.955 (0.17%)	-30.990 (0.29%)	-30.977 (0.25%)	-30.879 (-0.07%)	-30.893 (-0.02%)	-30.901 E_h
Kr	E_x	-94.071 (0.25%)	-94.215 (0.41%)	-94.186 (0.37%)	-93.820 (-0.02%)	-93.940 (0.11%)	-93.834 E_h
	E_c	-1.756 (-5.10%)	-1.765 (-4.59%)	-1.768 (-4.47%)	-1.770 (-4.34%)	-1.770 (-4.34%)	-1.850 E_h
	E_{xc}	-95.827 (0.15%)	-95.980 (0.31%)	-95.953 (0.28%)	-95.590 (-0.10%)	-95.710 (0.03%)	-95.684 E_h
Xe	E_x	-179.315 (0.14%)	-179.614 (0.31%)	-179.567 (0.28%)	-178.827 (-0.13%)	-179.136 (0.04%)	-179.064 E_h
	E_c	-2.899 (-3.43%)	-2.910 (-3.07%)	-2.914 (-2.96%)	-2.918 (-2.82%)	-2.918 (-2.82%)	-3.002 E_h
	E_{xc}	-182.214 (0.08%)	-182.524 (0.25%)	-182.480 (0.23%)	-181.745 (-0.18%)	-182.053 (-0.01%)	-182.066 E_h
MAPE[E_x]	0.29%	0.44%	0.40%	0.12%	0.14%		
MAPE[E_c]	6.29%	5.86%	5.71%	5.58%	5.58%		
MAPE[E_{xc}]	0.13%	0.28%	0.24%	0.10%	0.02%		
r_s 2	E_x	2631.022 (-0.27%)	2198.716 (16.21%)	2259.204 (13.90%)	2318.763 (11.63%)	2419.739 (7.78%)	2624.000 E_h
	E_c	811.437 (-5.66%)	989.604 (-28.85%)	971.939 (-26.55%)	963.796 (-25.49%)	963.796 (-25.49%)	768.000 E_h
	E_{xc}	3442.459 (-1.49%)	3188.320 (6.00%)	3231.143 (4.74%)	3282.559 (3.23%)	3383.535 (0.25%)	3392.000 E_h
r_s 3	E_x	489.080 (7.02%)	352.217 (33.04%)	394.134 (25.07%)	412.474 (21.58%)	424.755 (19.25%)	526.000 E_h
	E_c	299.340 (-23.69%)	361.776 (-49.49%)	342.788 (-41.65%)	339.987 (-40.49%)	339.987 (-40.49%)	242.000 E_h
	E_{xc}	788.419 (-2.66%)	713.992 (7.03%)	736.922 (4.05%)	752.461 (2.02%)	764.741 (0.42%)	768.000 E_h
r_s 4	E_x	126.894 (19.18%)	82.262 (47.60%)	92.568 (41.04%)	100.800 (35.80%)	102.789 (34.53%)	157.000 E_h
	E_c	146.586 (-40.95%)	152.932 (-47.05%)	162.499 (-56.25%)	161.117 (-54.92%)	161.117 (-54.92%)	104.000 E_h
	E_{xc}	273.480 (-4.78%)	235.194 (9.89%)	255.068 (2.27%)	261.917 (-0.35%)	263.906 (-1.11%)	261.000 E_h
r_s 6	E_x	6.264 (71.53%)	30.717 (-39.62%)	-1.577 (107.17%)	1.210 (94.50%)	1.604 (92.71%)	22.000 E_h
	E_c	52.651 (-69.84%)	20.481 (33.93%)	56.057 (-80.83%)	55.508 (-79.06%)	55.508 (-79.06%)	31.000 E_h
	E_{xc}	58.915 (-11.16%)	51.199 (3.40%)	54.480 (-2.79%)	56.719 (-7.02%)	57.112 (-7.76%)	53.000 E_h
MAPE[E_x]	24.50%	34.12%	46.79%	40.88%	38.57%		
MAPE[E_c]	35.03%	39.83%	51.32%	49.99%	49.99%		
MAPE[E_{xc}]	5.02%	6.58%	3.46%	3.15%	2.39%		
Ar ₂ 1.6Å	360.936 (-1.19%)	361.031 (-1.17%)	360.200 (-1.39%)	362.516 (-0.76%)	360.616 (-1.28%)	365.292 kcal/mol	
Ar ₂ 1.8Å	195.723 (-1.28%)	197.023 (-0.62%)	196.277 (-1.00%)	198.109 (-0.07%)	196.431 (-0.92%)	198.255 kcal/mol	
Ar ₂ 2.0Å	102.465 (-0.73%)	103.577 (0.35%)	102.889 (-0.32%)	104.242 (1.00%)	103.151 (-0.06%)	103.215 kcal/mol	
MAPE	1.07%	0.71%	0.90%	0.61%	0.75%		

TABLE II. Summary of atomization energy errors (in kcal/mol) for the G3 test set [63] using the most dense numerical integration grid (TURBOMOLE level 7).

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
ME	-5.036	-14.010	-12.912	-5.042	-6.939
MAE	6.121	14.258	13.239	5.866	7.716

[67] to single point energy calculations at a range of lattice volumes. All five functionals gave comparable good accuracy across all three test sets, showing that SCAN’s good performance is not significantly changed by the regularizations or exact constraint restoration for these properties.

TABLE III. Mean error (ME) and Mean absolute error (MAE) of SCAN[5], rSCAN[22], r++SCAN, r²SCAN [26], and r⁴SCAN for the S22 set of 22 interaction energies between closed shell complexes[64], and the BH76 set of 76 chemical barrier heights[65]. Full data is presented in supplemental material Tables VI and VII. All calculations use the most dense integration grid (TURBOMOLE level 7).

		SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
S22	ME	-0.524	-1.153	-0.554	-0.937	-0.874
	MAE	0.786	1.273	0.846	1.057	1.015
BH76	ME	-7.653	-7.365	-7.488	-7.125	-7.463
	MAE	7.724	7.434	7.556	7.182	7.527

VI. CONCLUSIONS

To summarize, we have shown how exact constraints obeyed by SCAN are broken by the regularizations in

TABLE IV. Error in lattice constant prediction for the LC20 lattice constant test set [66]. Errors are in Å relative to zero-point anharmonic expansion corrected experimental data from Ref. [68]. Lattice constants were obtained by fitting the stabilized jellium equation of state [67] to single point energy calculations at a range of lattice volumes.

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
Ag	0.012	0.028	0.026	0.034	0.017
Al	-0.012	-0.027	-0.031	-0.032	-0.014
Ba	0.049	0.100	0.046	0.076	0.062
C	-0.004	-0.001	-0.001	0.005	0.002
Ca	-0.009	0.017	-0.011	0.018	0.016
Cu	-0.030	-0.023	-0.025	-0.020	-0.027
GaAs	0.020	0.031	0.026	0.029	0.017
Ge	0.029	0.043	0.038	0.039	0.028
Li	0.011	0.021	0.006	0.016	0.010
LiCl	0.016	0.021	0.017	0.034	0.032
LiF	0.004	0.008	0.008	0.021	0.020
MgO	0.018	0.019	0.018	0.027	0.024
NaCl	0.010	0.025	0.015	0.036	0.036
NaF	0.006	0.015	0.012	0.028	0.028
Na	-0.007	0.024	-0.012	0.004	0.031
Pd	0.016	0.026	0.026	0.032	0.019
Rh	-0.005	0.006	0.005	0.008	-0.005
Si	0.005	0.012	0.009	0.018	0.016
SiC	0.002	-0.001	-0.002	0.006	0.007
Sr	0.039	0.064	0.019	0.056	0.095
ME	0.009	0.020	0.009	0.022	0.021
MAE	0.015	0.025	0.017	0.027	0.025

rSCAN. Through this analysis we have shown how the exact constraint adherence can be restored and how this can be achieved without sacrificing the good numerical performance of rSCAN. This results in three new functionals with increasing exact constraint adherence: r++SCAN, r²SCAN, and r⁴SCAN. Additional parameters introduced to the new functionals are set without reference to any real bonded systems, thus we can still regard the resulting functionals as non-empirical and expect them to be applicable to a wide range of systems. Figure 7 suggests that restoring GEX4 in r⁴SCAN gives similar accuracy to SCAN with some improvement in grid efficiency. We therefore expect the new r²SCAN functional to remain the preferred choice for situations where the accuracy of SCAN is desired but its use is prohibited by poor numerical performance [21, 69].

Further improvement over r²SCAN might be achieved by a smoother and fuller incorporation of the fourth-

order density-gradient terms for the exchange energy in a SCAN-like functional. Work on this is underway.

Figure 7 shows that rSCAN and r++SCAN, which lose the correct second-order gradient expansions for densities that vary slowly over space, also lose accuracy for atomization energies of molecules, and that the restoration of this limit in SCAN, r²SCAN, and r⁴SCAN also restores accurate atomization energies. This result is in line with arguments made in Ref. [70].

Experience with SCAN and r²SCAN (and with atomic densities [71]) suggests that smoothness at fixed electron number could be elevated to the status of an 18th exact constraint that a meta-GGA can satisfy, or at least to the status of a construction principle: By Occam’s Razor, the simplest assumption, consistent with what is known, should be preferred.

ACKNOWLEDGMENTS

J.F., J.N., and J.S. acknowledge the support of the U.S. DOE, Office of Science, Basic Energy Sciences Grant No. DE-SC0019350. J.S. also acknowledges the support of the US National Science Foundation under Grant No. DMR-2042618. A.D.K. acknowledges the support of the U.S. Department of Energy, Office of Science, Basic Energy Sciences, through Grant No. DE-SC0012575 to the Energy Frontier Research Center: Center for Complex Materials from First Principles, and also support from Temple University. JPP acknowledges the support of the US National Science Foundation under Grant No. DMR-1939528. We thank Albert Bartók-Partay and Daniel Mejía-Rodríguez for their invaluable discussions around the ideas presented here. J.P.P. and J.S. thank Natalie Holzwarth for pointing out that the SCAN exchange-correlation potential for an atom diverges in the tail of the density, making pseudo-potential construction difficult.

MATERIALS AVAILABILITY

r²SCAN and r⁴SCAN subroutines are made freely available at <https://gitlab.com/dhamil/r2scan-subroutines>. The data that support the findings of this study are available from the corresponding author upon reasonable request.

-
- [1] J. P. Perdew and K. Schmidt, AIP Conf. Proc. **577**, 1 (2001).
[2] A. D. Becke, Int. J. Quantum Chem. **23**, 1915 (1983), <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qua.560230605>.
[3] A. D. Becke and M. R. Roussel, Phys. Rev. A **39**, 3761 (1989).
[4] J. P. Perdew, Phys. Rev. Lett. **55**, 1665 (1985), *ibid.*, **55**, 2370 (1985).
[5] J. Sun, A. Ruzsinszky, and J. P. Perdew, Phys. Rev. Lett. **115**, 036402 (2015).
[6] P. D. Mezei, G. I. Csonka, and M. Kallay, J. Chem. Theory Comput. 10.1021/acs.jctc.8b00072 (2018).

- [7] T. Aschebrock and S. Kümmel, *Phys. Rev. Res.* **1**, 033082 (2019).
- [8] B. Neupane, H. Tang, N. K. Nepal, S. Adhikari, and A. Ruzsinszky, *arXiv*, 1 (2021), [arXiv:2104.04053](https://arxiv.org/abs/2104.04053).
- [9] D. A. Kitchaev, H. Peng, Y. Liu, J. Sun, J. P. Perdew, and G. Ceder, *Phys. Rev. B* **93**, 045132 (2016).
- [10] J. Sun, R. C. Remsing, Y. Zhang, Z. Sun, A. Ruzsinszky, H. Peng, Z. Yang, A. Paul, U. Waghmare, X. Wu, M. L. Klein, and J. P. Perdew, *Nat. Chem.* **8**, 831 (2016).
- [11] H. Peng and J. P. Perdew, *Phys. Rev. B* **96**, 100101(R) (2017).
- [12] Y. Zhang, J. Sun, J. P. Perdew, and X. Wu, *Phys. Rev. B* **96**, 035143 (2017).
- [13] M. Chen, H.-Y. Ko, R. C. Remsing, M. F. Calegari Andrade, B. Santra, Z. Sun, A. Selloni, R. Car, M. L. Klein, J. P. Perdew, and X. Wu, *Proc. Natl. Acad. Sci. U. S. A.* **114**, 10846 (2017), [arXiv:1709.10493](https://arxiv.org/abs/1709.10493).
- [14] J. W. Furness, Y. Zhang, C. Lane, I. G. Buda, B. Barbiellini, R. S. Markiewicz, A. Bansil, and J. Sun, *Commun. Phys.* **1**, 1 (2018).
- [15] C. Lane, J. W. Furness, I. G. Buda, Y. Zhang, R. S. Markiewicz, B. Barbiellini, J. Sun, and A. Bansil, *Phys. Rev. B* **98**, 125140 (2018).
- [16] G. Sai Gautam and E. A. Carter, *Phys. Rev. Mater.* **2**, 095401 (2018).
- [17] Y. Zhang, J. W. Furness, B. Xiao, and J. Sun, *J. Chem. Phys.* **150**, 014105 (2019).
- [18] Y. Zhang, C. Lane, J. W. Furness, B. Barbiellini, J. P. Perdew, R. S. Markiewicz, A. Bansil, and J. Sun, *Proc. Natl. Acad. Sci. U.S.A.* **117**, 68 (2020), <https://www.pnas.org/content/117/1/68.full.pdf>.
- [19] A. Pulkkinen, B. Barbiellini, J. Nokelainen, V. Sokolovskiy, D. Baigutlin, O. Miroshkina, M. Zagrebin, V. Buchelnikov, C. Lane, R. S. Markiewicz, A. Bansil, J. Sun, K. Pussi, and E. Lähderanta, *Phys. Rev. B* **101**, 075115 (2020).
- [20] R. Zhang, B. Singh, C. Lane, J. Kidd, Y. Zhang, B. Barbiellini, R. S. Markiewicz, A. Bansil, and J. Sun, *arXiv:2003.11052 [cond-mat.str-el]* (2020).
- [21] J. Ning, J. W. Furness, and J. Sun, *arXiv:2107.11850 [cond-mat.mtrl-sci]* (2021).
- [22] A. P. Bartók and J. R. Yates, *J. Chem. Phys.* **150**, 161101 (2019).
- [23] J. W. Furness and J. Sun, *Phys. Rev. B* **99**, 041119 (2019).
- [24] D. Mejía-Rodríguez and S. B. Trickey, *J. Chem. Phys.* **151**, 207101 (2019).
- [25] A. P. Bartók and J. R. Yates, *J. Chem. Phys.* **151**, 207102 (2019).
- [26] J. W. Furness, A. D. Kaplan, J. Ning, J. P. Perdew, and J. Sun, *J. Phys. Chem. Lett.* **11**, 8208 (2020).
- [27] D. Mejía-Rodríguez and S. B. Trickey, *Phys. Rev. B* **102**, 121109(R) (2020).
- [28] D. Mejía-Rodríguez and S. B. Trickey, *J. Phys. Chem. A* **124**, 9889 (2020).
- [29] S. Ehlert, U. Huniar, J. Ning, J. W. Furness, J. Sun, A. D. Kaplan, J. P. Perdew, and J. G. Brandenburg, *J. Chem. Phys.* **154**, 061101 (2021).
- [30] S. Grimme, A. Hansen, S. Ehlert, and J.-M. Mewes, *J. Chem. Phys.* **154**, 064103 (2021).
- [31] L. Goerigk, A. Hansen, C. Bauer, S. Ehrlich, A. Najibi, and S. Grimme, *Phys. Chem. Chem. Phys.* **19**, 32184 (2017).
- [32] G. Santra and J. M. L. Martin, *J. Chem. Theory Comput.* **17**, 1368–1379 (2021).
- [33] S. Song, S. Vuckovic, E. Sim, and K. Burke, *The Journal of Physical Chemistry Letters* **12**, 800 (2021), pMID: 33411542, <https://doi.org/10.1021/acs.jpcclett.0c03545>.
- [34] S. Dasgupta, E. Lambros, J. Perdew, and F. Paesani, *chemRxiv* 10.33774/chemrxiv-2021-hstgf-v2 (2021).
- [35] A. D. Becke and K. E. Edgecombe, *J. Chem. Phys.* **92**, 5397 (1990).
- [36] A. Ruzsinszky, J. Sun, B. Xiao, and G. I. Csonka, *J. Chem. Theory Comput.* **8**, 2078 (2012).
- [37] J. Sun, B. Xiao, and A. Ruzsinszky, *J. Chem. Phys.* **137**, 051101 (2012), <https://doi.org/10.1063/1.4742312>.
- [38] J. Sun, B. Xiao, Y. Fang, R. Haunschield, P. Hao, A. Ruzsinszky, G. I. Csonka, G. E. Scuseria, and J. P. Perdew, *Phys. Rev. Lett.* **111**, 106401 (2013).
- [39] J. W. Furness, N. Sengupta, J. Ning, A. Ruzsinszky, and J. Sun, *J. Chem. Phys.* **152**, 244112 (2020), <https://doi.org/10.1063/5.0008014>.
- [40] M. Levy and J. P. Perdew, *Phys. Rev. A* **32**, 2010 (1985).
- [41] M. Levy, *Phys. Rev. A* **43**, 4637 (1991).
- [42] M. Levy and J. P. Perdew, *Phys. Rev. B* **48**, 11638 (1993).
- [43] A. Görling and M. Levy, *Phys. Rev. A* **45**, 1509 (1992).
- [44] L. Pöhlmann and J. P. Perdew, *J. Phys. Condens. Matter* **12**, 1239 (2000).
- [45] L. Chiodo, L. A. Constantin, E. Fabiano, and F. Della Sala, *Phys. Rev. Lett.* **108**, 126402 (2012).
- [46] J. P. Perdew, A. Ruzsinszky, J. Sun, and K. Burke, *J. Chem. Phys.* **140**, 18A533 (2014), <https://doi.org/10.1063/1.4870763>.
- [47] J. W. Furness and S. Lehtola, *Hartree-Fock Orbitals for Spherical Atoms - a python toolbox* (2021).
- [48] E. Clementi and C. Roetti, *At. Data Nucl. Data Tables* **14**, 177 (1974).
- [49] Z.-h. Yang, H. Peng, J. Sun, and J. P. Perdew, *Phys. Rev. B* **93**, 205205 (2016).
- [50] D. A. Kirzhnits, *J. Exp. Theor. Phys.* **5**, 64 (1957).
- [51] P. Svendsen and U. von Barth, *Phys. Rev. B* **54**, 17402 (1996).
- [52] S.-K. Ma and K. A. Brueckner, *Phys. Rev.* **165**, 18 (1968).
- [53] Y. Wang and J. P. Perdew, *Phys. Rev. B* **43**, 8911 (1991).
- [54] J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 3865 (1996).
- [55] J. Sun, J. P. Perdew, and A. Ruzsinszky, *Proc. Natl. Acad. Sci.* **112**, 685 (2015).
- [56] J. P. Perdew and Y. Wang, *Phys. Rev. B* **45**, 13244 (1992).
- [57] A. D. Becke, *Phys. Rev. A* **38**, 3098 (1988).
- [58] S. J. Chakravorty, S. R. Gwaltney, and E. R. Davidson, *Phys. Rev. A* **47**, 3649 (1993).
- [59] S. P. McCarthy and A. J. Thakkar, *J. Chem. Phys.* **134**, 044102 (2011).
- [60] B. Wood, N. D. M. Hine, W. M. C. Foulkes, and P. García-González, *Phys. Rev. B* **76**, 035403 (2007).
- [61] N. D. Lang and W. Kohn, *Phys. Rev. B* **1**, 4555 (1970).
- [62] K. Patkowski, G. Murdachaew, C. M. Fou, and K. Szalewicz, *Mol. Phys.* **103**, 2031 (2005).
- [63] L. A. Curtiss, K. Raghavachari, P. C. Redfern, and J. A. Pople, *J. Chem. Phys.* **112**, 7374 (2000).
- [64] P. Jurecka, J. Šponer, J. Černý, and P. Hobza, *Phys. Chem. Chem. Phys.* **8**, 1985 (2006).

- [65] Y. Zhao, N. González-Garda, and D. G. Truhlar, *J. Phys. Chem. A* **109**, 2012 (2005).
- [66] J. Sun, M. Marsman, G. I. Csonka, A. Ruzsinszky, P. Hao, Y. S. Kim, G. Kresse, and J. P. Perdew, *Phys. Rev. B* **84**, 035117 (2011).
- [67] V. N. Staroverov, G. E. Scuseria, J. Tao, and J. P. Perdew, *Phys. Rev. B* **69**, 075102 (2004).
- [68] P. Hao, Y. Fang, J. Sun, G. I. Csonka, P. H. T. Philipsen, and J. P. Perdew, *Phys. Rev. B* **85**, 014111 (2012).
- [69] R. Kingsbury, A. Gupta, C. J. Bartel, J. M. Munro, S. Dwaraknath, M. Horton, and K. A. Persson, [chemrXiv:10.33774/chemrxiv-2021-gwm9m-v2](https://arxiv.org/abs/10.33774/chemrxiv-2021-gwm9m-v2) (2021).
- [70] A. D. Kaplan, B. Santra, P. Bhattarai, K. Wagle, S. T. u. R. Chowdhury, P. Bhetwal, J. Yu, H. Tang, K. Burke, M. Levy, and J. P. Perdew, *J. Chem. Phys.* **153**, 074114 (2020), <https://doi.org/10.1063/5.0017805>.
- [71] M. G. Medvedev, I. S. Bushmarinov, J. Sun, J. P. Perdew, and K. A. Lyssenko, *Science* **355**, 49 (2017), <https://science.sciencemag.org/content/355/6320/49.full.pdf>.
- [72] M. Brack, B. K. Jennings, and Y. H. Chu, *Phys. Lett. B* **65**, 1 (1976).
- [73] G. L. Oliver and J. P. Perdew, *Phys. Rev. A* **20**, 397 (1979).
- [74] J. Tao, J. P. Perdew, V. N. Staroverov, and G. E. Scuseria, *Phys. Rev. Lett.* **91**, 146401 (2003).
- [75] C.-O. Almbladh and U. von Barth, *Phys. Rev. B* **31**, 3231 (1985).
- [76] A. D. Kaplan, K. Wagle, and J. P. Perdew, *Phys. Rev. B* **98**, 085147 (2018).
- [77] T. H. Dunning, *J. Chem. Phys.* **90**, 1007 (1989).
- [78] D. E. Woon and T. H. Dunning, *J. Chem. Phys.* **103**, 4572 (1995).

SUPPLEMENTAL MATERIAL: CONSTRUCTION OF META-GGA FUNCTIONALS THROUGH RESTORATION OF EXACT CONSTRAINT ADHERENCE TO REGULARIZED SCAN FUNCTIONALS.

Appendix A: The slowly-varying limit of r²SCAN and r⁴SCAN

This Appendix sketches the derivation of r²SCAN and r⁴SCAN. We will presume that both functionals have the structure of Eq. 1, where the interpolation functions $f_{x/c}(\mathbf{r})$, are taken from rSCAN. As discussed in the main text, this functional is termed r++SCAN.

From the starting point of r++SCAN, we will derive the corrections needed to restore the second order gradient expansions for exchange and correlation (r²SCAN), and those for the fourth-order gradient expansion for exchange (r⁴SCAN). Thus r⁴SCAN can be viewed as a correction to r²SCAN exchange, and we begin with r²SCAN.

1. The gradient expansion of $\bar{\alpha}$

The gradient expansion of $\tau[n]$, the one-body, spin-unpolarized kinetic energy density, was derived in Ref. [72]

$$\tau[n] = \tau_U(n) + \frac{1}{6}\nabla^2 n + \frac{1}{72}\frac{|\nabla n|^2}{n} + \mathcal{O}[|\nabla n|^4] \quad (\text{A1})$$

(in Hartree atomic units). Here, $\mathcal{O}[|\nabla n|^4]$ indicates that the next term in the series of higher order is of the form $|\nabla n|^4$, $\nabla^2 n |\nabla n|^2$, $(\nabla^2 n)^2$, etc. The gradient expansion is more useful in terms of dimensionless (length-scale invariant) variables

$$p(n) = \left[\frac{|\nabla n|}{2k_F n} \right]^2 = \frac{3}{40} \frac{|\nabla n|^2}{\tau_U(n)n} \quad (\text{A2})$$

$$q(n) = \frac{\nabla^2 n}{4k_F^2 n} = \frac{3}{40} \frac{\nabla^2 n}{\tau_U(n)}, \quad (\text{A3})$$

where we have used

$$\tau_U(n) = \frac{3}{10} k_F^2 n \quad (\text{A4})$$

$$k_F = (3\pi^2 n)^{1/3}. \quad (\text{A5})$$

Then the gradient expansion of τ_σ can be cast as

$$\tau[n] = \tau_U(n) \left[1 + \frac{20}{9}q(n) + \frac{5}{27}p(n) \right] + \mathcal{O}[|\nabla n|^4]. \quad (\text{A6})$$

The integrated kinetic energy scales with the spin-densities in the same manner as the exchange energy [73]

$$T[n_\uparrow, n_\downarrow] = \frac{1}{2}\{T[2n_\uparrow] + T[2n_\downarrow]\}. \quad (\text{A7})$$

This implies a local spin-scaling relation

$$\tau[n_\uparrow, n_\downarrow] = \frac{1}{2}\{\tau[2n_\uparrow] + \tau[2n_\downarrow]\}. \quad (\text{A8})$$

We will seek a gradient expansion in terms of n and ζ , where

$$\zeta = \frac{n_\uparrow - n_\downarrow}{n_\uparrow + n_\downarrow}, \quad (\text{A9})$$

rather than the individual spin-densities. After simplification, one finds that

$$\tau(n_\uparrow, n_\downarrow) = \tau_U(n)d_s(\zeta) \left[1 + \frac{20}{9d_s(\zeta)}q + \frac{5}{27d_s(\zeta)}p + \frac{5}{27} \frac{\xi^2}{d_s(\zeta)(1-\zeta^2)} \right] + \mathcal{O}[|\nabla n|^4], \quad (\text{A10})$$

where

$$d_s(\zeta) = [(1+\zeta)^{5/3} + (1-\zeta)^{5/3}]/2 \quad (\text{A11})$$

describes the spin-scaling of the uniform electron gas kinetic energy density, and

$$\xi = \frac{|\nabla \zeta|}{2k_F}, \quad (\text{A12})$$

which also appeared in TPSS [74].

The spin resolved $\bar{\alpha}$ tends to

$$\bar{\alpha}(n_\uparrow, n_\downarrow) = \frac{\tau(n_\uparrow, n_\downarrow) - \tau_W}{\tau_U(n)d_s(\zeta) + \eta\tau_W} \quad (\text{A13})$$

$$= \left[\frac{\tau}{\tau_U(n)d_s(\zeta)} - \frac{5}{3d_s(\zeta)}p \right] \left[1 + \frac{5\eta}{3d_s(\zeta)}p \right]^{-1}. \quad (\text{A14})$$

After performing a Taylor expansion in p , and inserting Eq. A10 for $\tau(n_\uparrow, n_\downarrow)$, we find

$$\bar{\alpha}(n_\uparrow, n_\downarrow) = 1 + \frac{20}{9d_s(\zeta)}q - \frac{5(8+9\eta)}{27d_s(\zeta)}p + \frac{5}{27d_s(\zeta)(1-\zeta^2)}\xi^2 + \mathcal{O}[|\nabla n|^4]. \quad (\text{A15})$$

In the special case where $\zeta = 0$ (needed for the exchange energy),

$$\bar{\alpha}(n) = 1 + \frac{20}{9}q - \frac{5(8+9\eta)}{27}p + \mathcal{O}[|\nabla n|^4]. \quad (\text{A16})$$

2. Exchange, second order gradient expansion

For any spin-unpolarized exchange energy density $\epsilon_x(n)$, the spin-scaled exchange energy density is [73]

$$\epsilon_x(n_\uparrow, n_\downarrow) = \frac{1}{2}[\epsilon_x(2n_\uparrow) + \epsilon_x(2n_\downarrow)]. \quad (\text{A17})$$

Therefore we need only consider the spin-unpolarized exchange energy density, and apply the spin-scaling relationship as needed.

We start with an explicit expression for the $r^2\text{SCAN}$ exchange enhancement factor

$$F_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) = \{h_x^1(p) + f_x(\bar{\alpha})[h_x^0 - h_x^1(p)]\}g_x(p), \quad (\text{A18})$$

with $h_x^0 = 1.174$. The function

$$g_x(p) = 1 - \exp(-a_x/p^{1/4}) = 1 + \mathcal{O}[|\nabla n|^\infty] \quad (\text{A19})$$

in the slowly-varying limit (its derivatives of all order vanish in the limit $p \rightarrow 0$). In $r^2\text{SCAN}$, we take

$$h_x^1(p) = 1 + k_1 - k_1 \{1 + [D_x \exp(-p^2/d_{p2}^4) + \mu_{\text{AK}}]\}p/k_1\}^{-1}, \quad (\text{A20})$$

which has the following Taylor series:

$$h_x^1(p) = 1 + (D_x + \mu_{\text{AK}})p - \frac{(D_x + \mu_{\text{AK}})^2}{k_1}p^2 + \mathcal{O}[|\nabla n|^6]. \quad (\text{A21})$$

The $r^2\text{SCAN}$ interpolation function is taken from $r\text{SCAN}$, and has the structure

$$f_x(\bar{\alpha}) = \begin{cases} \sum_{i=0}^7 c_{x,i} \bar{\alpha}^i, & \bar{\alpha} \leq 2.5 \\ -c_{\text{dx}}^{\text{SCAN}} \exp\left[\frac{c_{2x}^{\text{SCAN}}}{1-\bar{\alpha}}\right], & \bar{\alpha} > 2.5 \end{cases}, \quad (\text{A22})$$

with Taylor series

$$f_x(\bar{\alpha}) = (\bar{\alpha} - 1) \sum_{i=1}^7 i c_{x,i} + \frac{1}{2}(\bar{\alpha} - 1)^2 \sum_{i=2}^7 i(i-1) c_{x,i} + \mathcal{O}[(\bar{\alpha} - 1)^3] \quad (\text{A23})$$

in the slowly-varying limit. It's important here to note that $\bar{\alpha}$ has a gradient expansion to much higher order than $\mathcal{O}[|\nabla n|^2]$, as we derived in Eqs. A15 and A16. Therefore, the term of lowest order in $(1 - \bar{\alpha})$ is $\mathcal{O}[|\nabla n|^2]$, and the term of lowest order in $(1 - \bar{\alpha})^2$ is $\mathcal{O}[|\nabla n|^4]$.

Inserting the Taylor series of Eqs. A19, A21, and A23 into $F_x^{\text{r}^2\text{SCAN}}$, we find, to $\mathcal{O}[|\nabla n|^4]$,

$$\begin{aligned} F_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) &= 1 + (D_x + \mu_{\text{AK}})p \\ &+ \left[(h_x^0 - 1) \sum_{i=1}^7 i c_{x,i} \right] (\bar{\alpha} - 1) - \frac{(D_x + \mu_{\text{AK}})^2}{k_1} p^2 \\ &- \left[(D_x + \mu_{\text{AK}}) \sum_{i=1}^7 i c_{x,i} \right] (\bar{\alpha} - 1)p \\ &+ \left[\frac{h_x^0 - 1}{2} \sum_{i=2}^7 i(i-1) c_{x,i} \right] (\bar{\alpha} - 1)^2 + \mathcal{O}[|\nabla n|^6], \end{aligned} \quad (\text{A24})$$

with the terms written in (generally) increasing order. For $r^2\text{SCAN}$, we demand that the exchange enhancement factor recover the exchange gradient expansion [50, 51]

$$F_x^{\text{GE}}(p, q) = 1 + \mu_{\text{AK}}p + \frac{146}{2025}q^2 - \frac{73}{405}pq + \mathcal{O}[|\nabla n|^6] \quad (\text{A25})$$

only to second order, i.e. $1 + \mu_{\text{AK}}p$. Therefore, we can ignore all terms of order higher than $\mathcal{O}[|\nabla n|^2]$ that are written explicitly in Eq. A24, but also those that are included implicitly in $(\bar{\alpha} - 1)$,

$$F_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) = 1 + \left[D_x - \frac{5(8+9\eta)}{27}(h_x^0 - 1) \sum_{i=1}^7 i c_{x,i} + \mu_{\text{AK}} \right] p + \frac{20(h_x^0 - 1)}{9} \left(\sum_{i=1}^7 i c_{x,i} \right) q + \mathcal{O}[|\nabla n|^4], \quad (\text{A26})$$

where we have evaluated $(\bar{\alpha} - 1)$ using Eq. A16.

To eliminate the term linear in q , we perform an integration by parts. The ‘‘gauge variance’’ of the exchange energy density implies that two exchange enhancement factors can yield the same integrated exchange energy, but different exchange energy densities,

$$\int_{\Omega} F_x \epsilon_x^{\text{LSDA}} d^3r = \int_{\Omega} [\tilde{F}_x \epsilon_x^{\text{LSDA}} + n^{-4/3} \nabla \cdot \mathbf{G}_x] d^3r = \int_{\Omega} \tilde{F}_x d^3r - \frac{3}{4\pi} (3\pi^2)^{1/3} \int_{\text{bdy}\Omega} \mathbf{G} \cdot d\mathbf{S}. \quad (\text{A27})$$

The second equality is a straightforward application of the divergence theorem. Provided that the gauge function \mathbf{G}_x vanishes sufficiently rapidly over the bounding surface $\text{bdy}\Omega$ of the integration volume Ω , the surface integral vanishes. Note also that the LSDA exchange energy density is

$$\epsilon_x^{\text{LSDA}} = -\frac{3}{4\pi} (3\pi^2)^{1/3} n^{4/3}. \quad (\text{A28})$$

Consider then

$$q \epsilon_x^{\text{LSDA}} = \left\{ \frac{p}{3} + n^{-4/3} \nabla \cdot \left[\frac{\nabla n}{4(3\pi^2)^{2/3} n^{1/3}} \right] \right\} \epsilon_x^{\text{LSDA}}, \quad (\text{A29})$$

thus

$$F_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) = 1 + \left[D_x - \frac{5(4+9\eta)}{27} (h_x^0 - 1) \sum_{i=1}^7 i c_{x,i} + \mu_{\text{AK}} \right] p + n^{-4/3} \nabla \cdot \left\{ \left[\frac{5(h_x^0 - 1)}{9(3\pi^2)^{2/3}} \left(\sum_{i=1}^7 i c_{x,i} \right) \right] \frac{\nabla n}{n^{1/3}} \right\} + \mathcal{O}[|\nabla n|^4]. \quad (\text{A30})$$

The rightmost term in curly braces is the gauge function \mathbf{G}_x . Except in certain situations, like the density tail of an atom (generally, outside the Kohn-Sham turning surface if one exists), where the gradient expansion does not apply, $n^{-1/3} \nabla n$ vanishes sufficiently rapidly at infinity. Moreover, we generalize $F_x^{\text{r}^2\text{SCAN}}$ by cutting off the divergent gradient expansion terms and keeping only the integrated-by-parts expression

$$\tilde{F}_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) = 1 + \left[D_x - \frac{5(4+9\eta)}{27} (h_x^0 - 1) \sum_{i=1}^7 i c_{x,i} + \mu_{\text{AK}} \right] p + \mathcal{O}[|\nabla n|^4], \quad (\text{A31})$$

allowing for validity even outside the Kohn-Sham turning surface.

Thus, we recover the correct second-order gradient expansion for exchange by demanding

$$D_x = \frac{5(4+9\eta)}{27} (h_x^0 - 1) \sum_{i=1}^7 i c_{x,i} \quad (\text{A32})$$

which is the product of C_η and C_{2x} defined in Eqs. 53 and 57, respectively,

$$C_\eta = \frac{5(4+9\eta)}{27}$$

$$C_{2x} = (h_x^0 - 1) \sum_{i=1}^7 i c_{x,i}.$$

3. Exchange, fourth-order gradient expansion

r^4SCAN prescribes an explicit correction to the r^2SCAN enhancement factor,

$$F_x^{\text{r}^4\text{SCAN}} = F_x^{\text{r}^2\text{SCAN}} + \Delta F_4(p, \bar{\alpha}) g_x(p) \quad (\text{A33})$$

where $g_x(p)$ is unchanged from r^2SCAN , and

$$\Delta F_4(p, \bar{\alpha}) = \left\{ -C_{2x}[(\bar{\alpha} - 1) + C_\eta p] + C_{\bar{\alpha}\bar{\alpha}}(1 - \bar{\alpha})^2 + C_{p\bar{\alpha}} p(1 - \bar{\alpha}) + C_{pp} p^2 \right\} \frac{2\bar{\alpha}^2}{1 + \bar{\alpha}^4} \times \exp \left[-\frac{(1 - \bar{\alpha})^2}{d_{\bar{\alpha}^4}^2} - \frac{p^2}{d_{p^4}^4} \right]. \quad (\text{A34})$$

Despite the complexity of ΔF_4 , the damping function used to modulate these corrections, it has a simple Taylor

series

$$\Delta F_4(p, \bar{\alpha}) = -C_{2x}[(\bar{\alpha} - 1) + C_\eta p] + C_{\bar{\alpha}\bar{\alpha}}(1 - \bar{\alpha})^2 + C_{p\bar{\alpha}} p(1 - \bar{\alpha}) + C_{pp} p^2 + \mathcal{O}[|\nabla n|^6]. \quad (\text{A35})$$

We will now take $D_x = C_\eta C_{2x}$ in the r^2SCAN exchange enhancement factor. Returning to Eq. A24 for the r^2SCAN exchange enhancement factor, and adding in the r^4SCAN corrections,

$$F_x^{\text{r}^4\text{SCAN}}(p, \bar{\alpha}) = 1 + \mu_{\text{AK}} p + \left[C_{pp} - \frac{(C_\eta C_{2x} + \mu_{\text{AK}})^2}{k_1} \right] p^2 + \left[C_{p\bar{\alpha}} + (C_\eta C_{2x} + \mu_{\text{AK}}) \sum_{i=1}^7 i c_{x,i} \right] (1 - \bar{\alpha}) p + \left[C_{\bar{\alpha}\bar{\alpha}} + \frac{h_x^0 - 1}{2} \sum_{i=2}^7 i(i-1) c_{x,i} \right] (\bar{\alpha} - 1)^2 + \mathcal{O}[|\nabla n|^6]. \quad (\text{A36})$$

Now, again using Eq. A16 for the gradient expansion of $\bar{\alpha}$ to second-order,

$$(1 - \bar{\alpha}) p = -\frac{20}{9} p q + \frac{5(8+9\eta)}{27} p^2 + \mathcal{O}[|\nabla n|^6] \quad (\text{A37})$$

$$(1 - \bar{\alpha})^2 = \frac{400}{81} q^2 + \left[\frac{5(8+9\eta)}{27} \right]^2 p^2 - \frac{200(8+9\eta)}{243} p q + \mathcal{O}[|\nabla n|^6]. \quad (\text{A38})$$

The second-order gradient expansion of $\bar{\alpha}$ is valid here, because any higher order terms in $1 - \bar{\alpha}$ will yield, to lowest order, sixth-order terms in these products. Using Eqs. A37 and A38, one can show that

$$\frac{73}{5000} (\bar{\alpha} - 1)^2 + \left[\frac{511}{13500} - \frac{73}{1500} \eta \right] (1 - \bar{\alpha}) p + \left[\frac{146}{2025} \left(\frac{2}{3} + \frac{3\eta}{4} \right)^2 - \frac{73}{405} \left(\frac{2}{3} + \frac{3\eta}{4} \right) \right] p^2 = \frac{146}{2025} q^2 - \frac{73}{405} p q + \mathcal{O}[|\nabla n|^6], \quad (\text{A39})$$

the same fourth-order terms as in Eq. A25.

Then to recover the fourth-order gradient expansion for exchange in r^4SCAN , we equate A36 and A39, and

find

$$C_{pp} = \frac{(C_\eta C_{2x} + \mu_{AK})^2}{k_1} + \frac{146}{2025} \left(\frac{2}{3} + \frac{3\eta}{4} \right)^2 - \frac{73}{405} \left(\frac{2}{3} + \frac{3\eta}{4} \right) \quad (\text{A40})$$

$$C_{p\bar{\alpha}} = \frac{511}{13500} - \frac{73}{1500} \eta - (C_\eta C_{2x} + \mu_{AK}) \sum_{i=1}^7 i c_{x,i} \quad (\text{A41})$$

$$C_{\bar{\alpha}\bar{\alpha}} = \frac{73}{5000} - \frac{h_x^0 - 1}{2} \sum_{i=2}^7 i(i-1) c_{x,i}, \quad (\text{A42})$$

as presented in Eqs. 61–63.

4. Correlation, second-order gradient expansion

The gradient expansion for the correlation energy per electron is known only to second order [5, 52–54]

$$\varepsilon_c(r_s, \zeta, t) = \varepsilon_c^{\text{LSDA}}(r_s, \zeta) + \beta(r_s) \phi^3(\zeta) t^2. \quad (\text{A43})$$

The density-dependent function $\beta(r_s)$ is known only for small values of r_s , and we take the parameterization used in Ref. [5],

$$\beta(r_s) = \beta_{\text{MB}} \frac{1 + 0.1 r_s}{1 + 0.1778 r_s}, \quad (\text{A44})$$

constructed to cancel with the second-order gradient expansion term for exchange in the limit $r_s \rightarrow \infty$. Two other quantities enter this expansion: the spin-scaling function

$$\phi(\zeta) = [(1 + \zeta)^{2/3} + (1 - \zeta)^{2/3}]/2, \quad (\text{A45})$$

and dimensionless density gradient on the length scale of the Thomas-Fermi wavevector

$$t^2 = \left(\frac{3\pi^2}{16} \right)^{2/3} \frac{p}{\phi^2(\zeta) r_s}. \quad (\text{A46})$$

In both r²SCAN and r⁴SCAN, we propose that the correlation energy per electron is

$$\varepsilon_c^{\text{r}^2\text{SCAN}}(r_s, \zeta, p, \bar{\alpha}) = \varepsilon_c^1(r_s, \zeta, p) + f_c(\bar{\alpha}) [\varepsilon_c^0(r_s, \zeta, p) - \varepsilon_c^1(r_s, \zeta, p)] \quad (\text{A47})$$

with $f_c(\bar{\alpha})$ taken from rSCAN. It has a Taylor series about $\bar{\alpha} = 1$ that is identical in structure (but not value) to the Taylor series for $f_x(\bar{\alpha})$. The individual energies per electron are

$$\varepsilon_c^0(r_s, \zeta, p) = [\varepsilon_c^{\text{LDA0}}(r_s, \zeta) + H_0(r_s, \zeta, p)] g_c(\zeta) \quad (\text{A48})$$

$$\varepsilon_c^1(r_s, \zeta, p) = \varepsilon_c^{\text{LSDA1}}(r_s, \zeta) + H_1(r_s, \zeta, p), \quad (\text{A49})$$

with ε_c^0 unchanged from SCAN (see also Eqs. C23–C28). In r²SCAN, we posit that

$$H_1(r_s, \zeta, p) = \gamma \phi^3(\zeta) \ln \{1 + w_1 [1 - g(y, \Delta y)]\} \quad (\text{A50})$$

$$y = \frac{\beta(r_s)}{\gamma w_1} t^2 \quad (\text{A51})$$

$$g(y, \Delta y) = [1 + 4(y - \Delta y)]^{-1/4} \quad (\text{A52})$$

$$\Delta y = D_c p \exp[-p^2/d_{p2}^4], \quad (\text{A53})$$

with d_{p2} unchanged from the exchange component of r²SCAN. It can readily be seen that these have the following Taylor series:

$$\varepsilon_c^0(r_s, \zeta, p) = \varepsilon_c^{\text{LDA0}}(r_s, \zeta) g_c(\zeta) + \chi_\infty g_c(\zeta) p + \mathcal{O}[|\nabla n|^4] \quad (\text{A54})$$

$$\varepsilon_c^1(r_s, \zeta, p) = \varepsilon_c^{\text{LSDA1}}(r_s, \zeta) + \beta(r_s) \phi^3(\zeta) t^2 - \gamma \phi^3(\zeta) w_1 D_c p + \mathcal{O}[|\nabla n|^4]. \quad (\text{A55})$$

Then the full gradient expansion of the r²SCAN correlation energy per electron is, after simplification,

$$\varepsilon_c^{\text{r}^2\text{SCAN}}(r_s, \zeta, p, \bar{\alpha}) = \varepsilon_c^{\text{LSDA1}} + \beta(r_s) \phi^3(\zeta) t^2 - \gamma \phi^3(\zeta) w_1 D_c p + \left(\sum_{i=1}^7 i c_{c,i} \right) (\bar{\alpha} - 1) + \mathcal{O}[|\nabla n|^4], \quad (\text{A56})$$

where $\varepsilon_c^{\text{LSDA0}} = \varepsilon_c^{\text{LDA0}} g_c(\zeta)$. Let

$$\Delta f_{c2} \equiv \sum_{i=1}^7 i c_{c,i}. \quad (\text{A57})$$

We can now use Eq. A15 for the gradient expansion of $\bar{\alpha}$ at arbitrary spin polarization to find that

$$\begin{aligned}
\varepsilon_c^{\text{r}^2\text{SCAN}}(r_s, \zeta, p, \bar{\alpha}) &= \varepsilon_c^{\text{LSDA1}}(r_s, \zeta) + \beta(r_s)\phi^3(\zeta)t^2 + \frac{20\Delta f_{c2}}{9d_s(\zeta)} [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)]q \\
&- \left\{ \gamma\phi^3(\zeta)w_1D_c + \frac{5(8+9\eta)\Delta f_{c2}}{27d_s(\zeta)} [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)] \right\} p \\
&+ \frac{5\Delta f_{c2}}{27d_s(\zeta)(1-\zeta^2)} [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)]\xi^2 + \mathcal{O}[|\nabla n|^4].
\end{aligned} \tag{A58}$$

We can eliminate the term linear in q using a similar gauge variance principle for the correlation energy:

$$\int_{\Omega} \varepsilon_c n d^3r = \int_{\Omega} [\tilde{\varepsilon}_c + n^{-1}\nabla \cdot \mathbf{G}_c] n d^3r = \int_{\Omega} \tilde{\varepsilon}_c n d^3r + \int_{\text{bdy}\Omega} \mathbf{G}_c \cdot d\mathbf{S}. \tag{A59}$$

Again, provided that the gauge function \mathbf{G}_c vanishes sufficiently rapidly at the bounding surface $\text{bdy}\Omega$, we may replace the r^2SCAN correlation energy per electron with the integrated-by-parts expression $\tilde{\varepsilon}_c$. To do this, consider that for a general function $f(r_s, \zeta)$,

$$\nabla f(r_s, \zeta) = -\frac{r_s}{3n} \frac{\partial f}{\partial r_s} \nabla n + \frac{\partial f}{\partial \zeta} \nabla \zeta, \tag{A60}$$

where we have used $r_s = [4\pi n/3]^{-3}$. Then

$$\begin{aligned}
\frac{\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)}{d_s(\zeta)} q n &= \left\{ \frac{2}{3} \frac{\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)}{d_s(\zeta)} + \frac{r_s}{3d_s(\zeta)} \left[\frac{\partial \varepsilon_c^{\text{LSDA0}}}{\partial r_s} - \frac{\partial \varepsilon_c^{\text{LSDA1}}}{\partial r_s} \right] \right\} p n \\
&- \frac{\nabla n \cdot \nabla \zeta}{4(3\pi^2)n^{5/3}} \frac{\partial}{\partial \zeta} \left[\frac{\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)}{d_s(\zeta)} \right] n + n^{-1}\nabla \cdot \left[\frac{\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)}{d_s(\zeta)} \frac{\nabla n}{4(3\pi^2n)^{2/3}} \right].
\end{aligned} \tag{A61}$$

The rightmost term in square brackets is the gauge function \mathbf{G}_c . Then the integrated-by-parts r^2SCAN correlation energy per electron is

$$\begin{aligned}
\tilde{\varepsilon}_c^{\text{r}^2\text{SCAN}}(r_s, \zeta, p, \bar{\alpha}) &= \varepsilon_c^{\text{LSDA1}}(r_s, \zeta) + \beta(r_s)\phi^3(\zeta)t^2 \\
&- \left\{ \gamma\phi^3(\zeta)w_1D_c + \frac{45\Delta f_{c2}\eta}{27d_s(\zeta)} [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)] - \frac{20\Delta f_{c2}r_s}{27d_s(\zeta)} \left[\frac{\partial \varepsilon_c^{\text{LSDA0}}}{\partial r_s} - \frac{\partial \varepsilon_c^{\text{LSDA1}}}{\partial r_s} \right] \right\} p \\
&- \frac{5\nabla n \cdot \nabla \zeta}{9(3\pi^2)n^{5/3}} \frac{\partial}{\partial \zeta} \left[\frac{\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)}{d_s(\zeta)} \right] + \frac{5\Delta f_{c2}}{27d_s(\zeta)(1-\zeta^2)} [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)]\xi^2 + \mathcal{O}[|\nabla n|^4].
\end{aligned} \tag{A62}$$

In r^2SCAN , we make the simplification that $\nabla \zeta \approx 0$. Thus $\xi \approx 0$, and

$$\begin{aligned}
\tilde{\varepsilon}_c^{\text{r}^2\text{SCAN}}(r_s, \zeta, p, \bar{\alpha}) &= \varepsilon_c^{\text{LSDA1}}(r_s, \zeta) + \beta(r_s)\phi^3(\zeta)t^2 \\
&- \left\{ \gamma\phi^3(\zeta)w_1D_c + \frac{45\Delta f_{c2}\eta}{27d_s(\zeta)} [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)] - \frac{20\Delta f_{c2}r_s}{27d_s(\zeta)} \left[\frac{\partial \varepsilon_c^{\text{LSDA0}}}{\partial r_s} - \frac{\partial \varepsilon_c^{\text{LSDA1}}}{\partial r_s} \right] \right\} p + \mathcal{O}[|\nabla n|^4].
\end{aligned} \tag{A63}$$

To recover the second order gradient expansion for correlation, we take

$$D_c = \frac{\Delta f_{c2}}{27\gamma\phi^3(\zeta)d_s(\zeta)w_1(r_s, \zeta)} \left\{ 20r_s \left[\frac{\partial \varepsilon_c^{\text{LSDA0}}}{\partial r_s} - \frac{\partial \varepsilon_c^{\text{LSDA1}}}{\partial r_s} \right] - 45\eta [\varepsilon_c^{\text{LSDA0}}(r_s, \zeta) - \varepsilon_c^{\text{LSDA1}}(r_s, \zeta)] \right\}, \tag{A64}$$

which is the factor appearing Eq. 78.

Appendix B: Further discussion of non-uniform coordinate scaling

As described in Sec. II A, a density n and Kohn-Sham orbital ϕ_i that are non-uniformly scaled along one dimension,

here the x coordinate, have the form

$$n_{\lambda}^z(x, y, z) = \lambda n(u, y, z) \tag{B1}$$

$$[\phi_i]_{\lambda}^z(x, y, z) = \lambda^{1/2} \phi_i(u, y, z), \tag{B2}$$

where $u = \lambda x$. Using Eqs. 33–35, one can show that the iso-orbital indicator α scales as

$$\alpha_\lambda^x = \frac{5}{12(3\pi^2)^{2/3}[n(u, y, z)]^{8/3}} \left\{ \lambda^{4/3} f_{\alpha 1}(u, y, z) + \lambda^{-2/3} f_{\alpha 2}(u, y, z) \right\}, \quad (\text{B3})$$

where

$$f_{\alpha 1}(u, y, z) = 4n(u, y, z) \left(\sum_i^{\text{occ.}} \left| \frac{\partial \phi_i(u, y, z)}{\partial u} \right|^2 \right) - \left(\frac{\partial n(u, y, z)}{\partial u} \right)^2 \quad (\text{B4})$$

$$f_{\alpha 2}(u, y, z) = \left(\sum_i^{\text{occ.}} |\nabla_\perp \phi_i(u, y, z)|^2 \right) - |\nabla_\perp n(u, y, z)|^2. \quad (\text{B5})$$

Similarly, the dimensionless gradient p can be expressed as

$$\frac{5}{3} p_\lambda^x = \frac{5}{12(3\pi^2)^{2/3}[n(u, y, z)]^{8/3}} \left\{ \lambda^{4/3} f_{p 1}(u, y, z) + \lambda^{-2/3} f_{p 2}(u, y, z) \right\}, \quad (\text{B6})$$

where

$$f_{p 1}(u, y, z) = \left[\frac{\partial n(u, y, z)}{\partial u} \right]^2 \quad (\text{B7})$$

$$f_{p 2}(u, y, z) = |\nabla_\perp n(u, y, z)|^2. \quad (\text{B8})$$

Recall that the iso-orbital indicator used in r++SCAN, r²SCAN, and r⁴SCAN is

$$\bar{\alpha} = \frac{\alpha}{1 + 5\eta p/3}. \quad (\text{B9})$$

Now, in the limit $\lambda \rightarrow 0$, $\lambda^{-2/3} \gg 1 \gg \lambda^{4/3}$, and the leading order of the scaled α and scaled p will be $\lambda^{-2/3}$. Consequently,

$$\lim_{\lambda \rightarrow 0} \bar{\alpha} = \frac{f_{\alpha 2}(u, y, z)}{\eta f_2(u, y, z)}, \quad (\text{B10})$$

which is independent of the scaling parameter (here we are assuming that a change of coordinates from $x \rightarrow u = \lambda x$ is used to evaluate all necessary integrals as well).

As we will show, the $\lambda \rightarrow \infty$ limit can result in two different scaling behaviors for α . It appears that systems that are finite in at least one dimension have $\alpha \sim \lambda^{-2/3}$ as $\lambda \rightarrow \infty$, which yields an iso-orbital ($\alpha = 0$) character in the hard limit. Completely extended systems (like the uniform electron gas) will have the other scaling, $\alpha \sim \lambda^{4/3}$ as $\lambda \rightarrow \infty$.

To understand why this might be, consider a generic isolated atom. This is a prototype for finite systems, so we expect the ensuing analysis to hold qualitatively for

related systems like molecules. As is well-known, far from the nucleus, the density is dominated by the character of the highest-occupied (HO) Kohn-Sham orbital, and thus decays exponentially like [75]

$$\phi_{\text{HO}}(r) \sim e^{-\kappa r} \quad (\text{B11})$$

$$n(r) \sim e^{-2\kappa r}, \quad (\text{B12})$$

where $\kappa = \sqrt{-2I}$ and I is the ionization potential. Under the non-uniform coordinate scaling described here, $r \rightarrow r_u = \sqrt{u^2 + y^2 + z^2}$. Thus as $\lambda \rightarrow \infty$, $r \rightarrow \infty$ as well and the density and HO Kohn-Sham orbital should tend to their respective asymptotic behaviors. Then

$$\sum_i^{\text{occ.}} \left| \frac{\partial \phi_i(u, y, z)}{\partial u} \right|^2 \sim \left| \frac{\partial \phi_{\text{HO}}(u, y, z)}{\partial u} \right|^2$$

$$\sum_i^{\text{occ.}} \left| \frac{\partial \phi_i(u, y, z)}{\partial u} \right|^2 \sim \kappa^2 e^{-2\kappa r_u} \frac{u^2}{r_u^2} \quad (\text{B13})$$

$$\left(\frac{\partial n(u, y, z)}{\partial u} \right)^2 \sim 4\kappa^2 e^{-4\kappa r_u} \frac{u^2}{r_u^2}, \quad (\text{B14})$$

and the $f_{\alpha 1}(u, y, z)$ and $f_{\alpha 2}(u, y, z)$ functions vanish identically. Thus we anticipate that finite systems have $\alpha \sim \lambda^{-2/3}$ for large λ , as the other scaling behavior cannot yield an iso-orbital asymptotic behavior. Note that, in this case,

$$\lim_{\lambda \rightarrow \infty} \bar{\alpha} \rightarrow \lambda^{-2} \frac{f_{\alpha 2}(u, y, z)}{\eta f_{p 1}(u, y, z)}, \quad (\text{B15})$$

which tends to zero like α .

Additionally, systems that are finite only in one dimension can exhibit similar scaling behavior for α . Consider the quasi-two dimensional electron gas in the infinite barrier model [44]. This system is a two-dimensional uniform electron gas in the yz plane with a finite, small thickness along the x axis. Moreover, the kinetic energy density of this system can be shown to be [76]

$$\tau(x) = \tau_W(x) + \frac{1}{2[r_s^{2D}]^2} n(x), \quad (\text{B16})$$

where r_s^{2D} is taken to be a constant Wigner-Seitz density parameter for the two-dimensional gas. Under non-uniform coordinate scaling,

$$\alpha_\lambda^x = \lambda^{-2/3} \frac{10}{6(3\pi^2)^{2/3}[r_s^{2D}]^2[n(u)]^{2/3}}. \quad (\text{B17})$$

Finally, if the scaled $\alpha \sim \lambda^{4/3}$ as $\lambda \rightarrow \infty$,

$$\lim_{\lambda \rightarrow \infty} \bar{\alpha} \rightarrow \frac{f_{\alpha 1}(u, y, z)}{\eta f_{p 1}(u, y, z)}. \quad (\text{B18})$$

This is again independent of the scale parameter. In a three-dimensional uniform electron gas, it is straightforward to show (using cylindrical coordinates) that

$$\alpha_\lambda^x = \frac{1}{3}(\lambda^{4/3} + 2\lambda^{-2/3}). \quad (\text{B19})$$

It should be emphasized that this example is only illustrative, and serves to demonstrate that such a $\lambda^{4/3}$ scaling

is possible in the large λ limit. We cannot assert that all extended systems will exhibit similar scaling behavior for α .

Appendix C: Working Equations

The full equations required for implementing $r^4\text{SCAN}$ and $r^4\text{SCAN}$ are given below. Note that by construction $\bar{\alpha} \geq 0$ (also α and α'). In pseudo-potential codes (e.g. VASP) or through rounding errors in very small density regions $\bar{\alpha}$ can become negative however, which can cause numerical problems for interpolation functions that do not consider this possibility. An additional condition was included to Eqs. C7 and C29 to consistently handle negative $\bar{\alpha}$ regions. These provisions were essential to reliably converge calculations in VASP, as neither extending the polynomial interpolation nor setting a constant $f(\bar{\alpha} < 0) = 1$ were sufficient.

A list of constants needed for both functionals follows the lists of equations.

1. Exchange $r^2\text{SCAN}$

$$E_x^{\text{r}^2\text{SCAN}}[n_\uparrow, n_\downarrow] = \frac{1}{2} \{E_x^{\text{r}^2\text{SCAN}}[2n_\uparrow] + E_x^{\text{r}^2\text{SCAN}}[2n_\downarrow]\} \quad (\text{C1})$$

$$E_x^{\text{r}^2\text{SCAN}}[n] = \int \varepsilon_x^{\text{r}^2\text{SCAN}} n d^3r \quad (\text{C2})$$

$$\varepsilon_x^{\text{r}^2\text{SCAN}} = \varepsilon_x^{\text{LDA}}(r_s) F_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) \quad (\text{C3})$$

$$\varepsilon_x^{\text{LDA}}(r_s) = -\frac{\frac{3}{4\pi} \left(\frac{9\pi}{4}\right)^{1/3}}{r_s} \quad (\text{C4})$$

$$F_x^{\text{r}^2\text{SCAN}}(p, \bar{\alpha}) = \{h_{1x}(p) + f_x(\bar{\alpha}) [h_{0x} - h_{1x}(p)]\} g_x(p) \quad (\text{C5})$$

$$\bar{\alpha}(p, \alpha) = \frac{\alpha}{1 + \eta \frac{5}{3} p} = \frac{\tau - \tau_W}{\tau_U + \eta \tau_W} \quad (\text{C6})$$

$$f_x(\bar{\alpha}) = \begin{cases} \exp\left[-\frac{c_{1x}^{\text{SCAN}} \bar{\alpha}}{1 - \bar{\alpha}}\right] & \bar{\alpha} < 0 \\ \sum_{i=0}^7 c_{xi} \bar{\alpha}^i & 0 \leq \bar{\alpha} \leq 2.5 \\ -c_{dx}^{\text{SCAN}} \exp\left[\frac{c_{2x}^{\text{SCAN}}}{1 - \bar{\alpha}}\right] & \bar{\alpha} > 2.5 \end{cases} \quad (\text{C7})$$

$$h_{0x} = 1 + k_0 \quad (\text{C8})$$

$$h_{1x}(p) = 1 + k_1 - \frac{k_1}{1 + \frac{x(p)}{k_1}} \quad (\text{C9})$$

$$x(p) = (C_\eta C_2 \exp[-p^2/d_{p2}^4] + \mu) p \quad (\text{C10})$$

$$C_\eta = \left[\frac{20}{27} + \eta \frac{5}{3} \right] \quad (\text{C11})$$

$$C_2 = -\sum_{i=1}^7 i c_{xi} [1 - h_{0x}] \approx -0.162742 \quad (\text{C12})$$

$$g_x(p) = 1 - \exp\left[\frac{-a_1}{p^{1/4}}\right] \quad (\text{C13})$$

2. Exchange r⁴SCAN

$$F_x^{r^4\text{SCAN}}(p, \bar{\alpha}) = \{h_{1x}(p) + f_x(\bar{\alpha})[h_{0x} - h_{1x}(p)] + \Delta F_4(p, \bar{\alpha})\} g_x(p) \quad (\text{C14})$$

$$\Delta f_{x2} = -(1 - \bar{\alpha}) \sum_{i=1}^7 i c_{xi} \quad (\text{C15})$$

$$\Delta f_{x4} = \frac{(1 - \bar{\alpha})^2}{2} \sum_{i=2}^7 i(i-1) c_{xi} \quad (\text{C16})$$

$$\Delta F_4(p, \bar{\alpha}) = \{C_2 [(1 - \bar{\alpha}) - C_\eta p] + C_{\bar{\alpha}\bar{\alpha}}(1 - \bar{\alpha})^2 + C_{p\bar{\alpha}}p(1 - \bar{\alpha}) + C_{pp}p^2\} \frac{2\bar{\alpha}^2}{1 + \bar{\alpha}^4} \exp \left[-\frac{(1 - \bar{\alpha})^2}{d_{\bar{\alpha}^4}} - \frac{p^2}{d_{p^4}} \right] \quad (\text{C17})$$

$$C_{\bar{\alpha}\bar{\alpha}} = \frac{73}{5000} - \frac{1}{2} \sum_{i=2}^7 i(i-1) c_i [h_{0x} - 1] \approx -0.0593531 \quad (\text{C18})$$

$$C_{p\bar{\alpha}} = \frac{511}{13500} - \frac{73}{1500} \eta - \sum_{i=1}^7 i c_{xi} [C_\eta C_2 + \mu] \approx 0.0402684 \quad (\text{C19})$$

$$C_{pp} = \frac{146}{2025} \left\{ \eta \frac{3}{4} + \frac{2}{3} \right\}^2 - \frac{73}{405} \left\{ \eta \frac{3}{4} + \frac{2}{3} \right\} + \frac{(C_\eta C_2 + \mu)^2}{k_1} \approx -0.0880769 \quad (\text{C20})$$

3. Correlation (both r²SCAN and r⁴SCAN)

$$E_c^{\text{r}^2\text{SCAN}}[n_\uparrow, n_\downarrow] = \int \varepsilon_c^{\text{r}^2\text{SCAN}} n d^3r \quad (\text{C21})$$

$$\varepsilon_c^{\text{r}^2\text{SCAN}} = \varepsilon_c^1 + f_c(\bar{\alpha})(\varepsilon_c^0 - \varepsilon_c^1) \quad (\text{C22})$$

$$\varepsilon_c^0 = (\varepsilon_c^{\text{LDA0}} + H_0) g_c(\zeta) \quad (\text{C23})$$

$$H_0 = b_{1c} \ln \{1 + w_0[1 - g_\infty(\zeta = 0, s)]\} \quad (\text{C24})$$

$$\varepsilon_c^{\text{LDA0}} = -\frac{b_{1c}}{1 + b_{2c}\sqrt{r_s} + b_{3c}r_s} \quad (\text{C25})$$

$$g_\infty(\zeta = 0, s) = \frac{1}{(1 + 4\chi_\infty s^2)^{1/4}} \quad (\text{C26})$$

$$w_0 = \exp\left[-\frac{\varepsilon_c^{\text{LDA0}}}{b_{1c}}\right] - 1 \quad (\text{C27})$$

$$g_c(\zeta) = \{1 - 2.363[d_x(\zeta) - 1]\}(1 - \zeta^{12}) \quad (\text{C28})$$

$$f_c(\bar{\alpha}) = \begin{cases} \exp\left[-\frac{c_{1c}^{\text{SCAN}} \bar{\alpha}}{1 - \bar{\alpha}}\right] & \bar{\alpha} < 0 \\ \sum_{i=0}^7 c_{ci} \bar{\alpha}^i & 0 \leq \bar{\alpha} \leq 2.5 \\ -c_{dc} \exp\left[\frac{c_{2c}}{1 - \bar{\alpha}}\right] & \bar{\alpha} > 2.5 \end{cases} \quad (\text{C29})$$

$$\varepsilon_c^1 = \varepsilon_c^{\text{LSDA}} + H_c^1 \quad (\text{C30})$$

$$\Delta f_{c2} = \sum_{i=1}^7 i c_{ci} \quad (\text{C31})$$

$$C_2 = -C_\eta \Delta f_{c2} \quad (\text{C32})$$

$$d_s(\zeta) = \frac{(1 + \zeta)^{5/3} + (1 - \zeta)^{5/3}}{2} \quad (\text{C33})$$

$$H_c^1 = \gamma \phi^3 \ln [1 + w_1(1 - g(y, \Delta y))] \quad (\text{C34})$$

$$w_1 = \exp\left[-\frac{\varepsilon_c^{\text{LSDA}}}{\gamma \phi^3}\right] - 1 \quad (\text{C35})$$

$$g(y, \Delta y) = \frac{1}{(1 + 4(y - \Delta y))^{1/4}} \quad (\text{C36})$$

$$y = \frac{\beta(r_s)}{\gamma w_1} t^2 \quad (\text{C37})$$

$$\beta(r_s) = \beta_{\text{MB}} \frac{1 + 0.1r_s}{1 + 0.1778r_s} \quad (\text{C38})$$

$$\Delta y = \frac{\Delta f_{c2}}{27\gamma d_s(\zeta) \phi^3 w_1} \left\{ 20r_s \left[g_c(\zeta) \frac{\partial \varepsilon_c^{\text{LDA0}}}{\partial r_s} - \frac{\partial \varepsilon_c^{\text{LSDA}}}{\partial r_s} \right] - 45\eta [\varepsilon_c^{\text{LDA0}} g_c(\zeta) - \varepsilon_c^{\text{LSDA}}] \right\} p \exp[-p^2/d_{p2}^4] \quad (\text{C39})$$

4. Constants needed for r²SCAN and r⁴SCAN

Constants needed for both exchange and correlation in r²SCAN and r⁴SCAN:

$$d_{p2} = 0.361 \quad (\text{C40})$$

$$\eta = 0.001 \quad (\text{C41})$$

Constants needed for exchange:

$$\mathbf{c}_x = (1, -0.667, -0.4445555, -0.663086601049, 1.451297044490, \\ -0.887998041597, 0.234528941479, -0.023185843322) \quad (\text{C42})$$

$$c_{1x}^{\text{SCAN}} = 0.667 \quad (\text{C43})$$

$$c_{2x}^{\text{SCAN}} = 0.8 \quad (\text{C44})$$

$$c_{dx}^{\text{SCAN}} = 1.24 \quad (\text{C45})$$

$$k_0 = 0.174 \quad (\text{C46})$$

$$k_1 = 0.065 \quad (\text{C47})$$

$$\mu = 10/81 \quad (\text{C48})$$

$$a_1 = 4.9479 \quad (\text{C49})$$

$$(\text{C50})$$

Constants for r⁴SCAN exchange,

$$d_{p^4} = 0.802 \quad (\text{C51})$$

$$d_{\bar{\alpha}^4} = 0.178 \quad (\text{C52})$$

The \mathbf{c}_x are taken from Ref. [22]; all other constants in the preceding list are taken from Ref. [5].

Constants needed for correlation:

$$\mathbf{c}_c = (1, -0.64, -0.4352, -1.535685604549, 3.061560252175, \\ -1.915710236206, 0.516884468372, -0.051848879792) \quad (\text{C53})$$

$$c_{1c}^{\text{SCAN}} = 0.64 \quad (\text{C54})$$

$$c_2^{\text{SCAN}} = 1.5 \quad (\text{C55})$$

$$c_{dc}^{\text{SCAN}} = 0.7 \quad (\text{C56})$$

$$b_{1c} = 0.0285764 \quad (\text{C57})$$

$$b_{2c} = 0.0889 \quad (\text{C58})$$

$$b_{3c} = 0.125541 \quad (\text{C59})$$

$$\beta_{\text{MB}} \approx 0.066725 \quad (\text{C60})$$

$$\chi_\infty = \left(\frac{3\pi^2}{16}\right)^{2/3} \frac{\beta_{\text{MB}}}{1.778\{0.9 - 3[3/(16\pi)]^{2/3}\}} \approx 0.128025 \quad (\text{C61})$$

$$\gamma = (1 - \ln 2)/\pi^2 \approx 0.031090690869655 \quad (\text{C62})$$

The \mathbf{c}_c are taken from Ref. [22]; all other constants in the preceding list are taken from Ref. [5].

Appendix D: Reference Atomic Calculations With Standard Basis Set

	r++SCAN	r ⁴ SCAN
E	-128.5891915	-128.5891915
E _x	-12.14196798	-12.14196798

TABLE V. Reference self-consistent total energy (E) **using only the exchange part of the functional**, and corresponding exchange energies (E_x) in Hartree for the neon atom. Calculated using the cc-pVTZ basis set [77, 78] and the “reference” level Turbomole grid.

	r++SCAN	r ⁴ SCAN
E	-128.6015519	-128.5718446
E _{xc}	-12.15409076	-12.12073578

TABLE VI. Reference self-consistent total atomic energy (E) and exchange-correlation energies (E_{xc}) in Hartree for the neon atom. Calculated using the cc-pVTZ basis set [77, 78] and the “reference” level Turbomole grid.

Appendix E: Individual Xenon Potential Components




FIG. 8. The XC enhancement factor (top), the multiplicative component of the XC potential (middle), and the derivative of the XC energy density with respect to the orbital dependent kinetic energy density, τ , (bottom). Shown for the SCAN, r++SCAN, r²SCAN, and r⁴SCAN functionals, calculated from reference Hartree–Fock Slater orbitals [47, 48].

Appendix F: G3 Atomization Energies

TABLE VII: Deviation of SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
LiH	2.186	2.549	2.778	2.524	2.449	2.448	2.518

Continued on next page

TABLE VII: Deviation of SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
BeH	-10.520	-10.879	-11.005	-10.701	-10.472	-10.447	-10.508
CH	1.438	1.265	0.127	1.596	2.317	2.277	1.914
CH ₂ (³ B ₁)	-6.715	-6.243	-7.581	-6.325	-5.790	-5.911	-6.400
CH ₂ (¹ A ₁)	4.207	4.836	2.635	4.362	5.415	5.626	5.269
CH ₃	-5.871	-5.316	-7.022	-5.373	-4.543	-4.490	-4.867
Methane (CH ₄)	0.207	-0.173	-1.744	-0.707	0.398	0.746	0.570
NH	-2.770	-3.772	-1.302	-1.004	-1.294	-1.994	-1.230
NH ₂	-4.177	-6.533	-4.185	-3.224	-2.816	-3.294	-2.677
Ammonia (NH ₃)	3.112	-1.255	1.896	2.334	3.021	2.795	3.564
OH	-0.045	-1.446	-3.032	-4.284	-3.519	-2.676	-2.609
Water (H ₂ O)	6.827	3.340	1.985	0.484	1.593	2.886	2.971
Hydrogen fluoride (HF)	0.847	4.405	3.877	2.541	3.018	3.642	3.051
SiH ₂ (¹ A ₁)	3.181	3.272	2.264	2.696	2.471	2.316	2.179
SiH ₂ (³ B ₁)	-5.900	-8.309	-6.847	-7.640	-7.529	-7.592	-7.717
SiH ₃	-3.738	-4.588	-4.814	-4.743	-4.794	-4.948	-5.102
Silane (SiH ₄)	-0.270	0.160	-1.469	-0.594	-0.809	-1.062	-1.248
PH ₂	-1.783	-5.033	-1.050	-4.662	-4.630	-4.332	-4.024
PH ₃	2.702	-1.543	3.158	-1.232	-1.054	-0.669	-0.227
Hydrogen sulfide (H ₂ S)	3.024	-1.871	2.961	-1.635	-0.892	-0.288	-0.089
Hydrogen chloride (HCl)	-1.589	2.333	0.073	-0.551	-0.208	0.417	0.270
Li ₂	5.462	6.040	6.290	5.925	5.823	5.839	5.963
LiF	-0.790	6.241	3.731	2.530	3.046	3.713	3.052
Acetylene (C ₂ H ₂)	1.980	3.157	-1.206	1.534	3.594	4.140	3.719
Ethylene (H ₂ C=CH ₂)	-0.949	-0.497	-4.482	-1.982	0.149	0.781	0.411
Ethane (H ₃ C-CH ₃)	-2.103	-2.564	-5.813	-3.641	-1.396	-0.705	-1.086
CN	2.446	-1.058	0.804	1.916	3.371	3.249	3.741
Hydrogen cyanide (HCN)	3.210	0.741	0.811	2.758	4.491	4.558	5.123
CO	7.760	2.989	1.109	0.178	2.614	4.290	4.095
HCO	-1.365	-4.601	-7.396	-7.510	-5.389	-3.973	-4.253
Formaldehyde (H ₂ C=O)	2.472	-0.627	-4.063	-4.200	-1.969	-0.340	-0.491
Methanol (CH ₃ -OH)	2.170	-1.072	-4.520	-4.598	-2.383	-0.781	-0.914
N ₂	7.481	1.813	5.957	6.966	8.419	8.119	9.743
Hydrazine (H ₂ N-NH ₂)	2.081	-5.216	0.374	1.449	2.908	2.434	3.919
NO	3.446	-2.180	-1.047	-1.996	-0.187	0.682	1.385
O ₂	-5.548	-2.950	-8.604	-10.389	-9.051	-7.782	-7.366
Hydrogen peroxide (HO-OH)	8.488	2.977	-0.946	-3.916	-1.679	0.915	1.153
F ₂	3.598	1.020	2.414	0.247	1.739	2.998	1.675
Carbon dioxide (CO ₂)	3.055	-6.629	-8.421	-11.459	-7.974	-5.193	-5.300
Na ₂	3.816	2.868	2.944	2.635	2.673	2.742	2.715
Si ₂	2.134	3.216	1.044	1.917	1.490	1.153	0.910
P ₂	11.197	2.793	12.012	2.790	3.048	3.876	4.847
S ₂	-1.598	-8.305	-2.055	-8.723	-8.039	-7.419	-7.062
Cl ₂	-2.926	2.723	-1.299	-0.482	-0.361	0.330	0.242
NaCl	-0.624	3.843	1.340	-1.178	-0.257	0.736	0.409
Silicon monoxide (SiO)	11.761	6.310	5.488	3.433	4.601	5.804	5.689
CS	6.627	1.874	4.535	0.883	2.809	3.779	3.781
SO	-1.665	-5.249	-4.847	-8.817	-7.946	-6.717	-6.318
ClO	-5.098	-2.972	-6.678	-7.048	-6.655	-5.098	-5.017
Chlorine monofluoride (FCl)	-2.146	1.092	-0.103	-0.486	-0.330	0.607	0.001
Si ₂ H ₆	-3.243	-1.993	-5.529	-3.635	-4.149	-4.672	-5.078
Methyl chloride (CH ₃ Cl)	-4.788	-0.833	-4.970	-4.039	-2.638	-1.778	-2.122
Methanethiol (H ₃ CSH)	0.962	-3.920	-0.991	-4.221	-2.401	-1.510	-1.517
Hypochlorous acid (HOCl)	2.196	1.714	-1.983	-2.735	-1.745	-0.095	-0.010
Sulfur dioxide (SO ₂)	11.258	-3.155	1.681	-7.031	-3.987	-0.835	-0.505
BF ₃	-13.344	3.819	-0.202	-3.545	-2.246	-0.998	-2.619
BCl ₃	-18.533	-4.845	-10.795	-14.932	-13.937	-11.983	-12.188
AlF ₃	-5.436	13.042	6.542	3.670	5.393	7.153	5.023

Continued on next page

TABLE VII: Deviation of SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
AlCl ₃	-13.365	-0.538	-7.735	-12.913	-10.628	-7.882	-8.833
Carbon tetrafluoride (CF ₄)	-16.947	-4.795	-7.932	-12.136	-8.825	-6.169	-8.845
Carbon tetrachloride (CCl ₄)	-15.691	-2.150	-12.259	-11.792	-9.380	-7.114	-7.830
Carbon oxide sulfide (COS)	-0.318	-10.165	-7.227	-13.138	-9.944	-7.652	-7.564
Carbon bisulfide (CS ₂)	-1.993	-12.356	-4.488	-13.602	-10.788	-9.010	-8.666
Carbonic difluoride (COF ₂)	-4.111	-1.675	-5.331	-8.262	-4.952	-2.230	-3.699
Silicon tetrafluoride (SiF ₄)	-2.075	20.933	12.645	8.344	10.686	12.757	9.832
Silicon tetrachloride (SiCl ₄)	-12.793	5.713	-6.801	-11.262	-8.975	-5.948	-7.288
Dinitrogen monoxide (N ₂ O)	1.907	-11.626	-5.341	-6.886	-4.318	-3.535	-1.796
Nitrogen chloride oxide (ClNO)	-4.239	-7.362	-8.386	-10.665	-8.118	-6.275	-5.736
Nitrogen trifluoride (NF ₃)	-13.166	-11.202	-7.881	-11.179	-9.032	-7.592	-8.806
PF ₃	-3.746	9.737	9.021	1.565	3.328	5.379	3.662
O ₃	8.819	1.429	-4.593	-9.460	-5.877	-1.907	-1.540
F ₂ O	-2.176	-3.190	-3.783	-8.446	-6.101	-3.631	-4.739
Chlorine trifluoride (ClF ₃)	-21.279	-9.913	-12.221	-15.243	-14.226	-11.734	-13.726
Tetrafluoro Ethene (F ₂ C=CF ₂)	-26.837	-13.667	-18.805	-21.367	-17.450	-14.484	-17.353
Tetrachloro Ethene (C ₂ Cl ₄)	-22.264	-7.569	-20.099	-18.276	-14.972	-12.304	-13.199
Acetonitrile, trifluoro- (CF ₃ CN)	-14.441	-7.706	-10.425	-11.188	-6.724	-4.573	-6.107
Propyne (C ₃ H ₄)	-1.947	-1.162	-7.009	-3.257	-0.066	0.829	0.218
Allene (C ₃ H ₄)	-5.655	-5.048	-10.969	-7.274	-4.114	-3.157	-3.693
Cyclopropene (C ₃ H ₄)	-1.682	-2.244	-7.391	-3.737	-0.512	0.418	-0.168
Propylene (C ₃ H ₆)	-3.717	-3.562	-9.116	-5.572	-2.309	-1.329	-1.887
Cyclopropane (C ₃ H ₆)	-4.806	-6.203	-10.831	-7.431	-4.119	-3.136	-3.715
Propane (C ₃ H ₈)	-4.377	-4.827	-9.806	-6.533	-3.143	-2.108	-2.683
Trans-1,3-butadiene (C ₄ H ₆)	-6.072	-5.055	-12.979	-8.058	-3.778	-2.499	-3.232
Dimethylacetylene (C ₄ H ₆)	-4.579	-4.215	-11.520	-6.764	-2.429	-1.187	-1.994
Methylenecyclopropane (C ₄ H ₆)	-9.776	-10.506	-17.299	-12.681	-8.321	-7.016	-7.777
Bicyclo[1.1.0]butane (C ₄ H ₆)	-6.294	-8.637	-14.492	-10.004	-5.592	-4.284	-5.068
Cyclobutene (C ₄ H ₆)	-5.894	-5.805	-12.924	-8.198	-3.758	-2.441	-3.221
Cyclobutane (C ₄ H ₈)	-8.111	-8.449	-14.936	-10.455	-5.925	-4.546	-5.345
Isobutene (C ₄ H ₈)	-6.141	-6.059	-13.169	-8.595	-4.202	-2.873	-3.619
Trans-butane(C ₄ H ₁₀)	-6.084	-7.066	-13.471	-9.148	-4.621	-3.230	-4.006
Isobutane (C ₄ H ₁₀)	-5.313	-6.576	-13.083	-8.769	-4.244	-2.856	-3.624
Spiropentane (C ₅ H ₈)	-10.924	-13.199	-20.757	-15.224	-9.674	-8.010	-8.983
Benzene (C ₆ H ₆)	-15.017	-12.995	-25.134	-17.775	-11.333	-9.378	-10.433
Difluoromethane (CH ₂ F ₂)	-8.938	-2.334	-5.282	-6.285	-4.207	-2.740	-4.218
Trifluoromethane(CHF ₃)	-13.277	-3.304	-6.503	-9.068	-6.385	-4.307	-6.429
CH ₂ Cl ₂	-8.892	-1.098	-7.560	-6.777	-5.080	-3.688	-4.201
CHCl ₃	-12.121	-1.528	-9.932	-9.342	-7.287	-5.413	-6.055
Methylamine (H ₃ C-NH ₂)	0.146	-4.125	-2.783	-1.053	0.790	0.878	1.427
Acetonitrile (CH ₃ -CN)	-0.621	-3.511	-4.898	-1.901	0.943	1.336	1.719
Nitromethane (CH ₃ -NO ₂)	-1.237	-13.405	-14.234	-15.988	-11.686	-8.940	-8.265
Methyl nitrite (CH ₃ -O-N=O)	-0.797	-10.986	-13.084	-14.071	-9.969	-7.281	-6.659
Methyl silane (CH ₃ SiH ₃)	-0.794	-0.749	-3.827	-1.938	-1.058	-0.956	-1.329
Formic acid (HCOOH)	3.336	-4.913	-8.475	-10.566	-7.098	-4.202	-4.308
Methyl formate (HCOOCH ₃)	-1.889	-9.602	-15.235	-15.937	-11.404	-8.210	-8.531
Acetamide (CH ₃ CONH ₂)	-2.840	-11.631	-11.952	-11.514	-7.305	-5.527	-5.086
Aziridine (C ₂ H ₄ NH)	-4.201	-8.136	-8.863	-5.701	-2.832	-2.429	-2.077
Cyanogen (NCCN)	0.979	-4.074	-3.793	0.056	3.558	3.670	4.761
Dimethylamine ((CH ₃) ₂ NH)	-2.847	-7.190	-7.529	-4.657	-1.668	-1.242	-0.903
Trans ethylamine (CH ₃ CH ₂ NH ₂)	-3.064	-7.742	-7.872	-5.088	-2.110	-1.670	-1.314
Ketene (CH ₂ CO)	-3.098	-7.970	-11.730	-11.340	-8.040	-6.158	-6.457
Oxirane (C ₂ H ₄ O)	-2.841	-6.122	-11.133	-10.172	-6.793	-4.861	-5.194
Acetaldehyde (CH ₃ CHO)	-0.740	-4.341	-9.163	-8.325	-4.977	-2.999	-3.328
Glyoxal (HCOCOH)	1.423	-4.682	-11.485	-11.814	-7.378	-4.102	-4.379
Ethanol (CH ₃ CH ₂ OH)	0.083	-3.551	-8.409	-7.498	-4.123	-2.176	-2.506
Dimethylether (CH ₃ OCH ₃)	-2.158	-5.078	-10.448	-9.185	-5.872	-3.941	-4.279

Continued on next page

TABLE VII: Deviation of SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Thiirane (C ₂ H ₄ S)	-3.180	-7.754	-6.951	-8.929	-6.041	-4.826	-5.048
Dimethyl sulfoxide ((CH ₃) ₂ SO)	0.834	-9.436	-7.573	-12.239	-8.226	-5.714	-5.754
Ethanethiol (C ₂ H ₅ SH)	-0.864	-5.927	-4.626	-6.788	-3.826	-2.580	-2.784
Dimethyl sulfide (CH ₃ SCH ₃)	-1.565	-6.376	-5.184	-7.159	-4.269	-3.076	-3.276
Vinyl fluoride (CH ₂ =CHF)	-8.324	-4.326	-8.855	-7.468	-4.903	-3.702	-4.729
Ethyl chloride (C ₂ H ₅ Cl)	-7.346	-3.430	-9.191	-7.307	-4.729	-3.489	-4.054
Vinyl chloride (CH ₂ =CHCl)	-10.376	-5.774	-12.217	-9.983	-7.539	-6.344	-6.896
Acrylonitrile (CH ₂ =CHCN)	-0.798	-2.889	-6.695	-2.279	1.600	2.292	2.491
Acetone (CH ₃ COCH ₃)	-3.776	-7.361	-13.726	-11.860	-7.383	-5.050	-5.563
Acetic acid (CH ₃ COOH)	0.568	-7.639	-12.837	-13.850	-9.272	-6.035	-6.328
Acetyl fluoride (CH ₃ COF)	-6.596	-6.380	-11.830	-12.278	-8.352	-5.850	-6.876
CH ₃ COCl (acetyl chloride)	-7.092	-7.391	-14.071	-14.207	-10.353	-7.736	-8.329
CH ₃ CH ₂ CH ₂ Cl (propyl chloride)	-9.588	-5.784	-13.263	-10.223	-6.519	-4.937	-5.700
Isopropanol (CH ₃) ₂ CHOH)	-1.932	-5.921	-12.449	-10.441	-5.956	-3.651	-4.163
Methyl ethyl ether (C ₂ H ₅ OCH ₃)	-4.759	-7.910	-14.820	-12.496	-8.034	-5.753	-6.285
Trimethylamine ((CH ₃) ₃ N)	-5.909	-10.460	-12.649	-8.570	-4.487	-3.721	-3.584
Furan (C ₄ H ₄ O)	-8.459	-11.623	-20.469	-17.107	-11.696	-9.109	-9.780
C ₄ H ₄ S (thiophene)	-7.630	-12.231	-15.052	-14.851	-9.797	-7.860	-8.393
Pyrrrole (C ₄ H ₅ N)	-10.190	-14.989	-18.491	-13.809	-8.686	-7.556	-7.517
Pyridine (C ₅ H ₅ N)	-14.825	-16.824	-24.095	-17.381	-11.314	-9.932	-10.058
H ₂	2.072	2.088	2.088	2.089	2.088	2.088	2.088
HS	-0.164	-3.056	-0.181	-2.786	-2.409	-2.111	-2.052
CCH	-1.141	-0.784	-3.802	-1.578	0.092	0.360	-0.195
C ₂ H ₃ (² A')	-7.440	-7.003	-10.392	-7.823	-5.980	-5.577	-6.114
CH ₃ CO (² A')	-4.740	-8.392	-12.771	-11.763	-8.559	-6.824	-7.308
H ₂ COH (² A)	-2.919	-5.761	-9.061	-8.986	-6.965	-5.578	-5.881
CH ₃ O (² A')	-5.613	-7.371	-10.746	-10.876	-8.922	-7.689	-7.803
CH ₃ CH ₂ O (² A'')	-8.731	-11.191	-15.918	-15.059	-11.956	-10.346	-10.638
CH ₃ S (² A')	-3.660	-6.737	-5.552	-7.009	-5.548	-4.925	-5.043
C ₂ H ₅ (² A')	-8.630	-8.437	-11.686	-8.989	-7.019	-6.628	-7.192
(CH ₃) ₂ CH (² A')	-11.700	-11.662	-16.422	-12.664	-9.581	-8.837	-9.589
(CH ₃) ₂ CH (² A')	-11.700	-11.662	-16.422	-12.664	-9.581	-8.837	-9.589
NO ₂	-1.476	-13.635	-11.987	-14.997	-11.989	-9.855	-9.083
Methyl allene (C ₄ H ₆)	-7.918	-7.192	-14.782	-10.023	-5.709	-4.410	-5.143
Isoprene (C ₅ H ₈)	-7.995	-7.242	-16.749	-10.749	-5.339	-3.710	-4.634
Cyclopentane (C ₅ H ₁₀)	-8.610	-10.140	-18.251	-12.754	-7.052	-5.323	-6.318
n-Pentane (C ₅ H ₁₂)	-8.313	-8.787	-17.092	-11.642	-5.966	-4.240	-5.209
Neo pentane (C ₅ H ₁₂)	-6.580	-8.076	-16.122	-10.787	-5.128	-3.394	-4.351
1,4 Cyclohexadiene (C ₆ H ₈)	-10.464	-9.852	-20.989	-13.911	-7.327	-5.359	-6.498
Cyclohexane (C ₆ H ₁₂)	-10.149	-11.918	-21.759	-15.174	-8.329	-6.243	-7.432
n-Hexane (C ₆ H ₁₄)	-9.910	-11.386	-20.938	-14.458	-7.642	-5.552	-6.723
3-Methyl pentane (C ₆ H ₁₄)	-8.396	-10.350	-19.805	-13.332	-6.513	-4.436	-5.607
Toluene (C ₆ H ₅ CH ₃)	-17.063	-15.427	-29.048	-20.665	-13.085	-10.780	-12.027
n-Heptane (C ₇ H ₁₆)	-12.561	-13.046	-24.679	-17.051	-9.087	-6.669	-8.033
Cyclooctatetraene (C ₈ H ₈)	-14.862	-12.823	-28.540	-18.725	-10.109	-7.538	-9.015
n-Octane (C ₈ H ₁₈)	-13.714	-15.599	-28.310	-19.684	-10.566	-7.775	-9.342
Naphthalene (C ₁₀ H ₈)	-28.555	-25.406	-45.599	-33.373	-22.624	-19.318	-21.090
Azulene (C ₁₀ H ₈)	-28.941	-26.524	-46.202	-34.044	-23.255	-19.963	-21.741
Acetic acid methyl ester (CH ₃ COOCH ₃)	-3.630	-11.300	-18.500	-18.124	-12.484	-8.931	-9.434
t-Butanol ((CH ₃) ₃ COH)	-3.363	-7.715	-15.843	-12.782	-7.177	-4.522	-5.219
Aniline (C ₆ H ₅ NH ₂)	-17.441	-20.168	-28.714	-21.139	-13.912	-12.146	-12.472
Phenol (C ₆ H ₅ OH)	-13.913	-15.660	-28.982	-23.054	-15.506	-12.278	-13.285
Divinyl ether (C ₄ H ₆ O)	-6.711	-9.115	-18.479	-14.763	-9.381	-6.880	-7.601
Tetrahydrofuran (C ₄ H ₈ O)	-6.331	-9.966	-18.595	-15.193	-9.616	-6.984	-7.725
Cyclopentanone (C ₅ H ₈ O)	-8.899	-13.800	-23.266	-19.266	-12.462	-9.447	-10.372
Benzoquinone(C ₆ H ₄ O ₂)	-10.968	-15.618	-30.342	-25.830	-17.068	-12.508	-13.509
Pyrimidine (C ₄ H ₄ N ₂)	-15.751	-21.928	-24.133	-18.160	-12.418	-11.610	-10.809

Continued on next page

TABLE VII: Deviation of SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Dimethyl sulphone (C ₂ H ₆ O ₂ S)	3.462	-12.938	-9.818	-17.114	-11.673	-7.872	-7.966
Chlorobenzene (C ₆ H ₅ Cl)	-20.608	-14.758	-29.158	-22.071	-15.324	-12.823	-14.051
Butanedinitrile (NC-CH ₂ -CH ₂ -CN)	-1.686	-7.627	-10.457	-4.332	1.390	2.179	2.912
Pyrazine (C ₄ H ₄ N ₂)	-11.835	-17.633	-20.044	-14.107	-8.373	-7.554	-6.750
Acetyl acetylene (CH ₃ -C(=O)-CCH)	-1.078	-3.609	-12.737	-9.222	-3.798	-1.262	-2.011
Crotonaldehyde (CH ₃ -CH=CH-CHO)	-7.116	-10.095	-18.902	-15.618	-10.124	-7.515	-8.209
Acetic anhydride (CH ₃ -C(=O)-O-C(=O)-CH ₃)	-5.758	-17.770	-27.541	-27.625	-19.664	-14.520	-15.218
2,5-Dihydrothiophene (C ₄ H ₆ S)	-6.531	-11.083	-13.870	-13.387	-8.290	-6.445	-7.042
Isobutane nitrile ((CH ₃) ₂ CH-CN)	-2.601	-5.850	-10.605	-5.349	-0.240	0.847	0.834
Methyl ethyl ketone (CH ₃ -CO-CH ₂ -CH ₃)	-4.851	-9.791	-17.530	-14.594	-8.966	-6.308	-7.018
Isobutanal ((CH ₃) ₂ CH-CHO)	-3.492	-7.430	-15.447	-12.400	-6.780	-4.116	-4.842
1,4-Dioxane (C ₄ H ₈ O ₂)	-7.722	-13.501	-24.367	-21.747	-15.138	-11.295	-11.991
Tetrahydrothiophene (C ₄ H ₈ S)	-5.412	-11.002	-13.130	-12.857	-7.645	-5.739	-6.356
t-Butyl chloride ((CH ₃) ₃ C-Cl)	-10.952	-7.866	-16.965	-13.150	-8.259	-6.272	-7.248
n-Butyl chloride (CH ₃ -CH ₂ -CH ₂ -CH ₂ -Cl)	-10.998	-7.263	-16.371	-12.296	-7.447	-5.500	-6.465
Tetrahydropyrrole (C ₄ H ₈ NH)	-7.881	-12.830	-16.487	-11.368	-6.080	-4.961	-5.034
Nitro-s-butane (CH ₃ -CH ₂ -CH(CH ₃)-NO ₂)	-6.746	-19.152	-24.952	-23.295	-15.614	-11.838	-11.744
Diethyl ether (CH ₃ -CH ₂ -O-CH ₂ -CH ₃)	-6.373	-9.716	-18.134	-14.838	-9.289	-6.655	-7.384
Dimethyl acetal (CH ₃ -CH(OCH ₃) ₂)	-5.465	-12.007	-22.313	-19.966	-13.303	-9.452	-10.155
t-Butanethiol ((CH ₃) ₃ C-SH)	-4.121	-10.102	-11.961	-12.030	-6.781	-4.825	-5.410
Diethyl disulfide (CH ₃ -CH ₂ -S-S-CH ₂ -CH ₃)	-4.171	-13.478	-11.127	-15.218	-9.381	-6.937	-7.351
t-Butylamine ((CH ₃) ₃ C-NH ₂)	-5.991	-10.653	-14.043	-9.150	-3.907	-2.772	-2.793
Tetramethylsilane (Si(CH ₃) ₄)	-1.480	-3.070	-10.369	-5.434	-1.203	-0.081	-1.012
2-Methyl thiophene (C ₅ H ₆ S)	-9.650	-14.844	-19.194	-17.923	-11.735	-9.439	-10.173
N-methyl pyrrole (cyc-C ₄ H ₄ N-CH ₃)	-12.581	-17.459	-22.724	-16.768	-10.541	-9.095	-9.263
Tetrahydropyran (C ₅ H ₁₀ O)	-8.532	-12.726	-22.798	-18.362	-11.596	-8.631	-9.575
Diethyl ketone (CH ₃ -CH ₂ -CO-CH ₂ -CH ₃)	-8.520	-12.430	-22.133	-18.044	-11.274	-8.252	-9.161
Isopropyl acetate (CH ₃ -C(=O)-O-CH(CH ₃) ₂)	-7.322	-15.759	-25.982	-23.582	-15.663	-11.403	-12.295
Tetrahydrothiopyran (C ₅ H ₁₀ S)	-7.886	-13.641	-17.255	-15.960	-9.613	-7.345	-8.160
Piperidine (cyc-C ₅ H ₁₀ NH)	-9.140	-14.608	-19.990	-13.752	-7.347	-5.873	-6.137
t-Butyl methyl ether ((CH ₃) ₃ C-O-CH ₃)	-7.749	-10.985	-21.070	-16.686	-9.973	-7.015	-7.924
1,3-Difluorobenzene (C ₆ H ₄ F ₂)	-26.828	-18.559	-31.748	-26.671	-19.341	-16.268	-18.599
1,4-Difluorobenzene (C ₆ H ₄ F ₂)	-29.212	-20.326	-33.327	-28.317	-20.932	-17.859	-20.202
Fluorobenzene (C ₆ H ₅ F)	-21.106	-15.943	-28.641	-22.411	-15.520	-13.002	-14.701
Di-isopropyl ether ((CH ₃) ₂ CH-O-CH(CH ₃) ₂)	-9.489	-13.343	-25.130	-19.624	-11.750	-8.439	-9.542
PF ₅	-8.813	11.946	10.607	0.199	2.992	6.248	3.395
SF ₆	-22.845	-5.833	-1.148	-15.502	-12.047	-7.569	-10.783
P ₄	9.473	-6.849	12.287	-6.669	-6.415	-4.838	-2.888
SO ₃	11.684	-8.353	-3.105	-13.902	-9.538	-5.168	-4.853
SCl ₂	-4.313	-1.222	-1.536	-6.203	-5.104	-3.556	-3.575
POCl ₃	-6.586	-2.209	-4.906	-14.150	-11.245	-7.602	-7.849
PCl ₅	-23.160	-5.207	-12.969	-21.000	-18.722	-15.170	-15.684
Cl ₂ O ₂ S	-0.053	-8.993	-7.302	-17.982	-13.944	-9.699	-9.699
PCl ₃	-9.514	0.586	-2.869	-10.274	-8.500	-5.956	-6.296
Cl ₂ S ₂	-9.472	-10.981	-6.376	-16.255	-14.098	-11.853	-11.807
SiCl ₂ singlet	-4.249	5.245	-1.715	-4.714	-3.319	-1.749	-2.522
CF ₃ Cl	-18.823	-6.193	-11.021	-14.235	-11.093	-8.493	-10.708
Hexafluoro ethane (C ₂ F ₆)	-30.675	-11.656	-17.535	-22.828	-17.455	-13.298	-17.477
CF ₃	-19.471	-9.337	-12.250	-14.651	-12.340	-10.553	-12.775
C ₆ H ₅ (phenyl radical)	-21.391	-19.806	-31.271	-23.846	-17.750	-16.048	-17.300
Bicyclo[1.1.0]butane (C ₄ H ₆)	-6.294	-8.637	-14.492	-10.004	-5.592	-4.284	-5.068
(CH ₃) ₃ C (t-butyl radical)	-12.936	-13.437	-19.718	-14.970	-10.755	-9.663	-10.598
Trans-butane(C ₄ H ₁₀)	-6.084	-7.066	-13.471	-9.148	-4.621	-3.230	-4.006
1,3 Cyclohexadiene (C ₆ H ₈)	-11.086	-10.597	-21.739	-14.606	-8.023	-6.049	-7.186
ME	-5.612	-6.418	-10.524	-9.736	-6.209	-4.533	-5.036
MAE	7.164	7.723	11.629	10.332	7.128	5.681	6.121

TABLE VIII: Deviation of rSCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
LiH	2.022	2.093	2.081	2.144	2.139	2.125	2.130
BeH	-11.231	-11.181	-10.947	-11.020	-11.013	-10.998	-11.006
CH	1.521	1.462	1.516	1.491	1.387	1.406	1.426
CH ₂ (³ B ₁)	-8.621	-8.971	-8.715	-8.716	-8.799	-8.790	-8.769
CH ₂ (¹ A ₁)	3.671	3.600	3.783	3.736	3.596	3.610	3.644
CH ₃	-7.135	-7.528	-7.236	-7.324	-7.429	-7.409	-7.387
Methane (CH ₄)	-3.790	-3.669	-3.563	-3.537	-3.680	-3.668	-3.636
NH	-1.620	-1.740	-1.830	-2.047	-1.987	-1.930	-1.959
NH ₂	-4.130	-4.009	-4.056	-4.354	-4.335	-4.257	-4.290
Ammonia (NH ₃)	-0.659	-0.105	-0.321	-0.542	-0.550	-0.472	-0.501
OH	-3.766	-3.568	-3.749	-3.810	-3.839	-3.856	-3.867
Water (H ₂ O)	-0.889	-0.755	-1.162	-1.224	-1.257	-1.275	-1.288
Hydrogen fluoride (HF)	-0.290	-0.392	-0.261	-0.123	-0.133	-0.136	-0.132
SiH ₂ (¹ A ₁)	1.263	1.067	1.120	1.130	1.159	1.133	1.135
SiH ₂ (³ B ₁)	-8.897	-8.627	-8.715	-8.645	-8.607	-8.616	-8.621
SiH ₃	-6.450	-6.395	-6.402	-6.369	-6.332	-6.353	-6.355
Silane (SiH ₄)	-2.241	-2.303	-2.227	-2.227	-2.194	-2.228	-2.223
PH ₂	-5.469	-5.024	-5.023	-4.833	-4.853	-4.820	-4.815
PH ₃	-2.224	-1.645	-1.640	-1.399	-1.408	-1.361	-1.366
Hydrogen sulfide (H ₂ S)	-3.163	-2.412	-2.397	-2.393	-2.411	-2.359	-2.354
Hydrogen chloride (HCl)	-1.187	-1.934	-1.824	-1.733	-1.702	-1.700	-1.690
Li ₂	4.834	5.124	5.105	5.225	5.213	5.187	5.197
LiF	-0.271	-0.676	-0.696	-0.400	-0.415	-0.433	-0.424
Acetylene (C ₂ H ₂)	-3.163	-3.109	-2.704	-2.683	-2.959	-2.941	-2.877
Ethylene (H ₂ C=CH ₂)	-6.540	-6.457	-6.125	-6.106	-6.391	-6.370	-6.304
Ethane (H ₃ C-CH ₃)	-8.068	-7.842	-7.620	-7.571	-7.861	-7.835	-7.770
CN	-0.681	-0.256	-0.246	-0.431	-0.545	-0.465	-0.462
Hydrogen cyanide (HCN)	-0.594	-0.012	0.043	-0.159	-0.313	-0.228	-0.223
CO	0.028	0.200	-0.088	-0.153	-0.337	-0.357	-0.329
HCO	-8.358	-8.390	-8.620	-8.689	-8.842	-8.851	-8.835
Formaldehyde (H ₂ C=O)	-5.322	-5.200	-5.401	-5.476	-5.656	-5.665	-5.645
Methanol (CH ₃ -OH)	-6.038	-5.954	-6.188	-6.256	-6.431	-6.436	-6.417
N ₂	5.226	6.345	6.076	5.648	5.601	5.752	5.701
Hydrazine (H ₂ N-NH ₂)	-2.190	-1.189	-1.536	-2.013	-2.025	-1.869	-1.926
NO	-2.141	-1.812	-2.373	-2.655	-2.686	-2.629	-2.671
O ₂	-11.013	-11.904	-12.027	-12.095	-12.152	-12.185	-12.201
Hydrogen peroxide (HO-OH)	-3.602	-3.445	-4.062	-4.215	-4.285	-4.326	-4.353
F ₂	-1.197	-1.189	-0.926	-0.756	-0.785	-0.798	-0.794
Carbon dioxide (CO ₂)	-12.833	-12.419	-13.130	-13.261	-13.484	-13.510	-13.500
Na ₂	1.805	1.974	1.831	1.857	1.877	1.881	1.880
Si ₂	-1.773	-1.865	-1.689	-1.546	-1.482	-1.529	-1.534
P ₂	0.196	1.279	1.272	1.803	1.776	1.865	1.857
S ₂	-11.349	-10.626	-10.636	-10.411	-10.420	-10.364	-10.368
Cl ₂	-0.923	-2.313	-2.172	-2.036	-2.000	-1.986	-1.968
NaCl	-1.194	-1.930	-1.935	-1.819	-1.763	-1.765	-1.755
Silicon monoxide (SiO)	2.559	2.821	2.351	2.316	2.306	2.255	2.250
CS	-0.835	-0.151	0.074	0.037	-0.136	-0.077	-0.032
SO	-9.436	-9.338	-9.466	-9.398	-9.456	-9.428	-9.447
ClO	-6.582	-7.029	-7.264	-7.224	-7.272	-7.271	-7.279
Chlorine monofluoride (FCl)	-1.578	-2.203	-2.051	-1.887	-1.888	-1.885	-1.874
Si ₂ H ₆	-6.677	-6.762	-6.634	-6.631	-6.562	-6.631	-6.621
Methyl chloride (CH ₃ Cl)	-5.687	-6.447	-6.174	-6.101	-6.213	-6.196	-6.155
Methanethiol (H ₃ CSH)	-7.288	-6.495	-6.354	-6.334	-6.495	-6.430	-6.393
Hypochlorous acid (HOCl)	-3.075	-3.440	-3.732	-3.740	-3.762	-3.775	-3.780
Sulfur dioxide (SO ₂)	-6.976	-5.723	-6.709	-6.791	-6.883	-6.873	-6.893
BF ₃	-10.500	-10.946	-10.976	-10.597	-10.559	-10.550	-10.550
BCl ₃	-14.721	-16.929	-16.720	-16.546	-16.353	-16.344	-16.325

Continued on next page

TABLE VIII: Deviation of rSCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
AlF ₃	-3.285	-3.647	-3.414	-2.866	-2.915	-2.923	-2.908
AlCl ₃	-12.659	-13.924	-13.562	-13.284	-13.172	-13.174	-13.145
Carbon tetrafluoride (CF ₄)	-17.244	-18.019	-17.471	-16.942	-17.134	-17.124	-17.076
Carbon tetrachloride (CCl ₄)	-11.675	-14.407	-13.967	-13.628	-13.679	-13.644	-13.574
Carbon oxide sulfide (COS)	-15.481	-14.465	-14.712	-14.777	-14.982	-14.939	-14.910
Carbon bisulfide (CS ₂)	-17.106	-15.485	-15.234	-15.245	-15.436	-15.320	-15.272
Carbonic difluoride (COF ₂)	-10.905	-11.225	-11.331	-11.126	-11.324	-11.334	-11.307
Silicon tetrafluoride (SiF ₄)	1.167	-0.273	0.256	0.918	0.903	0.861	0.886
Silicon tetrachloride (SiCl ₄)	-9.054	-12.368	-11.748	-11.375	-11.200	-11.240	-11.189
Dinitrogen monoxide (N ₂ O)	-10.782	-9.266	-10.016	-10.546	-10.610	-10.472	-10.542
Nitrogen chloride oxide (ClNO)	-10.875	-10.906	-11.375	-11.584	-11.608	-11.551	-11.579
Nitrogen trifluoride (NF ₃)	-16.002	-16.367	-15.985	-15.866	-15.908	-15.846	-15.862
PF ₃	-1.802	-2.279	-1.927	-1.136	-1.175	-1.144	-1.137
O ₃	-8.912	-8.660	-9.767	-9.933	-10.034	-10.099	-10.140
F ₂ O	-9.366	-9.538	-9.517	-9.372	-9.433	-9.466	-9.477
Chlorine trifluoride (ClF ₃)	-18.643	-19.849	-19.485	-18.961	-18.998	-19.000	-18.979
Tetrafluoro Ethene (F ₂ C=CF ₂)	-26.871	-27.518	-26.892	-26.375	-26.691	-26.677	-26.600
Tetrachloro Ethene (C ₂ Cl ₄)	-20.242	-22.545	-21.909	-21.559	-21.741	-21.698	-21.597
Acetonitrile, trifluoro- (CF ₃ CN)	-17.256	-17.272	-16.780	-16.576	-16.911	-16.815	-16.766
Propyne (C ₃ H ₄)	-9.747	-9.542	-9.044	-8.997	-9.417	-9.385	-9.289
Allene (C ₃ H ₄)	-13.913	-13.740	-13.248	-13.199	-13.625	-13.597	-13.497
Cyclopropene (C ₃ H ₄)	-10.753	-10.454	-10.066	-10.013	-10.440	-10.406	-10.309
Propylene (C ₃ H ₆)	-11.797	-11.487	-11.072	-11.023	-11.451	-11.418	-11.319
Cyclopropane (C ₃ H ₆)	-13.767	-13.255	-12.984	-12.904	-13.337	-13.299	-13.202
Propane (C ₃ H ₈)	-12.736	-12.023	-11.743	-11.668	-12.103	-12.064	-11.967
Trans-1,3-butadiene (C ₄ H ₆)	-16.003	-15.725	-15.089	-15.044	-15.614	-15.573	-15.441
Dimethylacetylene (C ₄ H ₆)	-14.981	-14.627	-14.038	-13.964	-14.528	-14.483	-14.355
Methylenecyclopropane (C ₄ H ₆)	-20.864	-20.245	-19.766	-19.683	-20.255	-20.209	-20.079
Bicyclo[1.1.0]butane (C ₄ H ₆)	-17.810	-17.187	-16.814	-16.694	-17.274	-17.225	-17.094
Cyclobutene (C ₄ H ₆)	-14.724	-14.138	-13.627	-13.550	-14.127	-14.081	-13.950
Cyclobutane (C ₄ H ₈)	-17.245	-16.056	-15.676	-15.584	-16.163	-16.112	-15.981
Isobutene (C ₄ H ₈)	-16.732	-15.917	-15.426	-15.348	-15.922	-15.875	-15.744
Trans-butane(C ₄ H ₁₀)	-16.609	-16.062	-15.595	-15.506	-16.089	-16.036	-15.906
Isobutane (C ₄ H ₁₀)	-16.032	-15.633	-15.259	-15.168	-15.749	-15.698	-15.567
Spiropentane (C ₅ H ₈)	-24.963	-23.950	-23.499	-23.363	-24.084	-24.021	-23.860
Benzene (C ₆ H ₆)	-28.756	-28.141	-27.165	-27.098	-27.954	-27.894	-27.695
Difluoromethane (CH ₂ F ₂)	-9.045	-9.594	-9.252	-8.993	-9.156	-9.146	-9.107
Trifluoromethane(CHF ₃)	-12.908	-13.602	-13.128	-12.715	-12.894	-12.885	-12.841
CH ₂ Cl ₂	-7.862	-9.201	-8.848	-8.687	-8.774	-8.752	-8.701
CHCl ₃	-9.718	-11.841	-11.432	-11.180	-11.246	-11.218	-11.158
Methylamine (H ₃ C-NH ₂)	-4.874	-4.212	-4.298	-4.511	-4.663	-4.571	-4.568
Acetonitrile (CH ₃ -CN)	-7.132	-6.401	-6.246	-6.424	-6.721	-6.623	-6.587
Nitromethane (CH ₃ -NO ₂)	-19.367	-18.181	-19.020	-19.382	-19.608	-19.555	-19.578
Methyl nitrite (CH ₃ -O-N=O)	-15.933	-14.996	-15.816	-16.183	-16.411	-16.360	-16.382
Methyl silane (CH ₃ SiH ₃)	-5.976	-5.872	-5.700	-5.667	-5.779	-5.800	-5.763
Formic acid (HCOOH)	-11.162	-10.849	-11.562	-11.691	-11.903	-11.930	-11.921
Methyl formate (HCOOCH ₃)	-16.823	-16.449	-16.966	-17.112	-17.465	-17.478	-17.438
Acetamide (CH ₃ CONH ₂)	-15.989	-14.852	-15.201	-15.474	-15.802	-15.719	-15.694
Aziridine (C ₂ H ₄ NH)	-11.222	-10.192	-10.177	-10.368	-10.665	-10.561	-10.526
Cyanogen (NCCN)	-5.812	-4.594	-4.511	-4.914	-5.221	-5.047	-5.040
Dimethylamine ((CH ₃) ₂ NH)	-9.569	-8.692	-8.666	-8.867	-9.163	-9.058	-9.023
Trans ethylamine (CH ₃ CH ₂ NH ₂)	-10.573	-9.692	-9.640	-9.831	-10.130	-10.025	-9.989
Ketene (CH ₂ CO)	-15.129	-14.819	-14.956	-14.982	-15.303	-15.304	-15.248
Oxirane (C ₂ H ₄ O)	-12.333	-12.040	-12.155	-12.214	-12.537	-12.531	-12.479
Acetaldehyde (CH ₃ CHO)	-11.384	-11.044	-11.157	-11.199	-11.523	-11.520	-11.466
Glyoxal (HCOCOH)	-13.284	-12.895	-13.268	-13.414	-13.775	-13.796	-13.754
Ethanol (CH ₃ CH ₂ OH)	-10.520	-10.273	-10.405	-10.448	-10.769	-10.761	-10.709

Continued on next page

TABLE VIII: Deviation of rSCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Dimethylether (CH ₃ OCH ₃)	-11.131	-10.844	-10.922	-10.993	-11.313	-11.304	-11.253
Thiirane (C ₂ H ₄ S)	-13.579	-12.611	-12.328	-12.300	-12.606	-12.529	-12.460
Dimethyl sulfoxide ((CH ₃) ₂ SO)	-16.753	-15.423	-15.639	-15.632	-15.978	-15.921	-15.864
Ethanethiol (C ₂ H ₅ SH)	-11.359	-10.382	-10.130	-10.087	-10.395	-10.317	-10.248
Dimethyl sulfide (CH ₃ SCH ₃)	-12.084	-11.056	-10.823	-10.776	-11.083	-11.007	-10.938
Vinyl fluoride (CH ₂ =CHF)	-11.726	-11.854	-11.439	-11.288	-11.578	-11.559	-11.491
Ethyl chloride (C ₂ H ₅ Cl)	-10.470	-11.008	-10.618	-10.521	-10.778	-10.749	-10.675
Vinyl chloride (CH ₂ =CHCl)	-13.766	-14.324	-13.884	-13.785	-14.039	-14.013	-13.939
Acrylonitrile (CH ₂ =CHCN)	-9.091	-8.316	-7.958	-8.131	-8.571	-8.464	-8.394
Acetone (CH ₃ COCH ₃)	-17.049	-16.195	-16.228	-16.240	-16.707	-16.693	-16.606
Acetic acid (CH ₃ COOH)	-16.412	-15.730	-16.341	-16.445	-16.801	-16.816	-16.775
Acetyl fluoride (CH ₃ COF)	-15.727	-15.683	-15.733	-15.629	-15.962	-15.960	-15.903
CH ₃ COCl (acetyl chloride)	-16.653	-17.134	-17.154	-17.116	-17.407	-17.401	-17.338
CH ₃ CH ₂ CH ₂ Cl (propyl chloride)	-15.105	-15.269	-14.816	-14.687	-15.089	-15.046	-14.940
Isopropanol (CH ₃) ₂ CHOH)	-15.058	-14.729	-14.769	-14.781	-15.247	-15.226	-15.141
Methyl ethyl ether (C ₂ H ₅ OCH ₃)	-16.103	-15.605	-15.579	-15.621	-16.085	-16.064	-15.980
Trimethylamine ((CH ₃) ₃ N)	-14.470	-13.636	-13.511	-13.683	-14.125	-14.008	-13.940
Furan (C ₄ H ₄ O)	-22.556	-22.095	-21.971	-21.958	-22.562	-22.541	-22.421
C ₄ H ₄ S (thiophene)	-23.212	-21.991	-21.398	-21.312	-21.905	-21.813	-21.675
Pyrrole (C ₄ H ₅ N)	-22.491	-21.347	-21.039	-21.185	-21.762	-21.641	-21.539
Pyridine (C ₅ H ₅ N)	-26.992	-25.883	-25.244	-25.418	-26.143	-26.015	-25.877
H ₂	2.073	2.089	2.089	2.089	2.089	2.089	2.089
HS	-3.432	-3.012	-3.031	-3.052	-3.059	-3.030	-3.027
CCH	-5.561	-5.691	-5.292	-5.234	-5.465	-5.452	-5.391
C ₂ H ₃ (² A')	-11.708	-11.854	-11.457	-11.475	-11.720	-11.702	-11.642
CH ₃ CO (² A')	-14.346	-14.408	-14.463	-14.545	-14.836	-14.831	-14.783
H ₂ COH (² A)	-10.165	-10.346	-10.563	-10.654	-10.798	-10.801	-10.788
CH ₃ O (² A')	-10.978	-10.744	-10.821	-10.875	-11.048	-11.053	-11.032
CH ₃ CH ₂ O (² A'')	-16.855	-16.402	-16.378	-16.413	-16.731	-16.723	-16.669
CH ₃ S (² A')	-9.509	-8.966	-8.870	-8.862	-9.012	-8.969	-8.935
C ₂ H ₅ (² A')	-12.531	-12.722	-12.339	-12.398	-12.649	-12.616	-12.561
(CH ₃) ₂ CH (² A')	-18.293	-18.015	-17.558	-17.591	-17.985	-17.941	-17.853
(CH ₃) ₂ CH (² A')	-18.293	-18.015	-17.558	-17.591	-17.985	-17.941	-17.853
NO ₂	-16.417	-15.713	-16.732	-17.074	-17.142	-17.104	-17.159
Methyl allene (C ₄ H ₆)	-18.364	-17.828	-17.248	-17.170	-17.742	-17.701	-17.569
Isoprene (C ₅ H ₈)	-20.349	-19.822	-19.083	-19.006	-19.722	-19.667	-19.502
Cyclopentane (C ₅ H ₁₀)	-20.613	-19.945	-19.474	-19.367	-20.097	-20.032	-19.869
n-Pentane (C ₅ H ₁₂)	-21.281	-19.772	-19.359	-19.221	-19.946	-19.882	-19.719
Neo pentane (C ₅ H ₁₂)	-19.657	-19.037	-18.557	-18.458	-19.186	-19.121	-18.958
1,4 Cyclohexadiene (C ₆ H ₈)	-23.854	-23.033	-22.211	-22.093	-22.955	-22.887	-22.691
Cyclohexane (C ₆ H ₁₂)	-24.008	-23.320	-22.766	-22.623	-23.502	-23.424	-23.229
n-Hexane (C ₆ H ₁₄)	-25.021	-24.096	-23.380	-23.247	-24.120	-24.039	-23.844
3-Methyl pentane (C ₆ H ₁₄)	-23.702	-23.008	-22.384	-22.233	-23.107	-23.028	-22.833
Toluene (C ₆ H ₅ CH ₃)	-33.196	-32.453	-31.362	-31.261	-32.263	-32.190	-31.958
n-Heptane (C ₇ H ₁₆)	-30.132	-27.812	-27.268	-27.067	-28.083	-27.993	-27.764
Cyclooctatetraene (C ₈ H ₈)	-31.862	-31.018	-29.809	-29.691	-30.836	-30.753	-30.489
n-Octane (C ₈ H ₁₈)	-33.505	-32.002	-31.075	-30.896	-32.052	-31.944	-31.683
Naphthalene (C ₁₀ H ₈)	-50.069	-49.141	-47.554	-47.427	-48.856	-48.756	-48.423
Azulene (C ₁₀ H ₈)	-50.698	-49.588	-48.016	-47.894	-49.324	-49.223	-48.890
Acetic acid methyl ester (CH ₃ COOCH ₃)	-20.915	-20.199	-20.627	-20.743	-21.241	-21.242	-21.169
t-Butanol ((CH ₃) ₃ COH)	-18.995	-18.537	-18.478	-18.471	-19.084	-19.051	-18.933
Aniline (C ₆ H ₅ NH ₂)	-33.205	-31.824	-31.097	-31.242	-32.105	-31.965	-31.796
Phenol (C ₆ H ₅ OH)	-32.896	-32.138	-31.593	-31.578	-32.466	-32.423	-32.238
Divinyl ether (C ₄ H ₆ O)	-20.906	-20.353	-20.139	-20.147	-20.747	-20.722	-20.605
Tetrahydrofuran (C ₄ H ₈ O)	-18.880	-18.682	-18.576	-18.594	-19.202	-19.168	-19.052
Cyclopentanone (C ₅ H ₈ O)	-25.814	-25.273	-25.117	-25.079	-25.837	-25.795	-25.643
Benzoquinone(C ₆ H ₄ O ₂)	-32.751	-32.015	-31.729	-31.815	-32.747	-32.728	-32.552

Continued on next page

TABLE VIII: Deviation of rSCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Pyrimidine (C ₄ H ₄ N ₂)	-26.306	-24.740	-24.406	-24.833	-25.428	-25.231	-25.154
Dimethyl sulphone (C ₂ H ₆ O ₂ S)	-20.350	-18.469	-19.303	-19.348	-19.731	-19.689	-19.644
Chlorobenzene (C ₆ H ₅ Cl)	-31.944	-32.134	-31.048	-30.900	-31.724	-31.660	-31.452
Butanedinitrile (NC-CH ₂ -CH ₂ -CN)	-13.037	-11.649	-11.361	-11.709	-12.310	-12.112	-12.039
Pyrazine (C ₄ H ₄ N ₂)	-22.326	-20.740	-20.446	-20.862	-21.455	-21.260	-21.183
Acetyl acetylene (CH ₃ -C(=O)-CCH)	-15.583	-15.227	-14.979	-14.984	-15.588	-15.567	-15.449
Crotonaldehyde (CH ₃ -CH=CH-CHO)	-21.911	-21.341	-21.152	-21.173	-21.782	-21.759	-21.639
Acetic anhydride (CH ₃ -C(=O)-O-C(=O)-CH ₃)	-31.140	-30.387	-31.156	-31.307	-31.986	-31.995	-31.902
2,5-Dihydrothiophene (C ₄ H ₆ S)	-20.349	-19.277	-18.723	-18.647	-19.241	-19.143	-19.007
Isobutane nitrile ((CH ₃) ₂ CH-CN)	-13.533	-12.347	-12.006	-12.143	-12.733	-12.609	-12.507
Methyl ethyl ketone (CH ₃ -CO-CH ₂ -CH ₃)	-20.426	-20.287	-20.124	-20.130	-20.741	-20.714	-20.595
Isobutanal ((CH ₃) ₂ CH-CHO)	-18.482	-17.814	-17.711	-17.713	-18.326	-18.298	-18.179
1,4-Dioxane (C ₄ H ₈ O ₂)	-23.212	-22.781	-22.951	-23.071	-23.712	-23.695	-23.592
Tetrahydrothiophene (C ₄ H ₈ S)	-19.297	-18.338	-17.863	-17.777	-18.377	-18.273	-18.138
t-Butyl chloride ((CH ₃) ₃ C-Cl)	-18.912	-19.435	-18.859	-18.723	-19.270	-19.216	-19.076
n-Butyl chloride (CH ₃ -CH ₂ -CH ₂ -CH ₂ -Cl)	-18.629	-18.672	-18.095	-17.954	-18.503	-18.447	-18.307
Tetrahydropyrrole (C ₄ H ₈ NH)	-18.370	-17.146	-16.944	-17.094	-17.684	-17.554	-17.452
Nitro-s-butane (CH ₃ -CH ₂ -CH(CH ₃)-NO ₂)	-31.060	-29.648	-30.134	-30.431	-31.096	-31.003	-30.929
Diethyl ether (CH ₃ -CH ₂ -O-CH ₂ -CH ₃)	-20.096	-19.400	-19.312	-19.326	-19.937	-19.902	-19.786
Dimethyl acetal (CH ₃ -CH(OCH ₃) ₂)	-23.108	-22.514	-22.733	-22.857	-23.501	-23.484	-23.381
t-Butanethiol ((CH ₃) ₃ C-SH)	-19.432	-18.308	-17.872	-17.785	-18.382	-18.279	-18.143
Diethyl disulfide (CH ₃ -CH ₂ -S-S-CH ₂ -CH ₃)	-23.884	-21.625	-21.167	-21.060	-21.672	-21.517	-21.379
t-Butylamine ((CH ₃) ₃ C-NH ₂)	-18.208	-16.548	-16.333	-16.458	-17.043	-16.913	-16.811
Tetramethylsilane (Si(CH ₃) ₄)	-16.916	-16.247	-15.808	-15.681	-16.228	-16.211	-16.075
2-Methyl thiophene (C ₅ H ₆ S)	-27.541	-26.351	-25.675	-25.573	-26.312	-26.206	-26.035
N-methyl pyrrole (cyc-C ₄ H ₄ N-CH ₃)	-26.404	-25.192	-24.757	-24.890	-25.610	-25.476	-25.342
Tetrahydropyran (C ₅ H ₁₀ O)	-23.513	-23.021	-22.762	-22.759	-23.520	-23.473	-23.324
Diethyl ketone (CH ₃ -CH ₂ -CO-CH ₂ -CH ₃)	-26.146	-24.879	-24.746	-24.703	-25.463	-25.421	-25.269
Isopropyl acetate (CH ₃ -C(=O)-O-CH(CH ₃) ₂)	-29.430	-28.536	-28.778	-28.847	-29.634	-29.609	-29.471
Tetrahydrothiopyran (C ₅ H ₁₀ S)	-23.969	-22.597	-22.004	-21.895	-22.641	-22.524	-22.357
Piperidine (cyc-C ₅ H ₁₀ NH)	-21.658	-20.625	-20.342	-20.454	-21.191	-21.048	-20.914
t-Butyl methyl ether ((CH ₃) ₃ C-O-CH ₃)	-23.739	-22.760	-22.538	-22.545	-23.294	-23.249	-23.099
1,3-Difluorobenzene (C ₆ H ₄ F ₂)	-36.912	-36.823	-35.593	-35.266	-36.134	-36.078	-35.874
1,4-Difluorobenzene (C ₆ H ₄ F ₂)	-38.528	-38.331	-37.187	-36.850	-37.720	-37.662	-37.459
Fluorobenzene (C ₆ H ₅ F)	-33.023	-32.668	-31.569	-31.369	-32.231	-32.173	-31.972
Di-isopropyl ether ((CH ₃) ₂ CH-O-CH(CH ₃) ₂)	-28.348	-27.149	-26.834	-26.787	-27.690	-27.630	-27.448
PF ₅	-6.765	-7.508	-7.003	-5.968	-6.022	-5.995	-5.982
SF ₆	-21.971	-21.943	-21.152	-20.266	-20.341	-20.309	-20.290
P ₄	-11.276	-9.097	-9.038	-7.958	-8.007	-7.822	-7.841
SO ₃	-13.926	-12.339	-13.784	-13.950	-14.084	-14.089	-14.124
SCl ₂	-6.119	-7.243	-7.009	-6.812	-6.770	-6.711	-6.691
POCl ₃	-12.113	-12.890	-12.978	-12.528	-12.466	-12.444	-12.434
PCL ₅	-17.777	-20.396	-19.761	-19.095	-18.927	-18.880	-18.842
Cl ₂ O ₂ S	-16.285	-16.641	-17.388	-17.306	-17.339	-17.314	-17.319
PCL ₃	-8.891	-10.159	-9.709	-9.204	-9.098	-9.058	-9.034
Cl ₂ S ₂	-16.892	-17.321	-16.986	-16.797	-16.762	-16.656	-16.629
SiCl ₂ singlet	-3.637	-5.580	-5.206	-5.022	-4.914	-4.950	-4.922
CF ₃ Cl	-17.509	-18.881	-18.345	-17.862	-18.015	-18.001	-17.947
Hexafluoro ethane (C ₂ F ₆)	-29.546	-30.830	-29.904	-29.091	-29.450	-29.434	-29.343
CF ₃	-19.080	-19.779	-19.389	-18.996	-19.127	-19.123	-19.085
C ₆ H ₅ (phenyl radical)	-34.220	-33.962	-32.917	-32.881	-33.699	-33.644	-33.450
Bicyclo[1.1.0]butane (C ₄ H ₆)	-17.810	-17.187	-16.814	-16.694	-17.274	-17.225	-17.094
(CH ₃) ₃ C (t-butyl radical)	-22.148	-22.037	-21.444	-21.432	-21.973	-21.917	-21.796
Trans-butane(C ₄ H ₁₀)	-16.609	-16.062	-15.595	-15.506	-16.089	-16.036	-15.906
1,3 Cyclohexadiene (C ₆ H ₈)	-24.454	-23.906	-23.081	-22.964	-23.827	-23.759	-23.561
ME	-14.299	-14.011	-13.826	-13.774	-14.121	-14.080	-14.010
MAE	14.531	14.258	14.069	14.025	14.368	14.328	14.258

TABLE IX: Deviation of r++SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
LiH	2.407	2.428	2.386	2.480	2.511	2.504	2.498
BeH	-10.768	-10.712	-10.485	-10.553	-10.547	-10.532	-10.540
CH	2.197	2.136	2.188	2.160	2.058	2.079	2.098
CH ₂ (³ B ₁)	-7.859	-8.212	-7.957	-7.960	-8.042	-8.031	-8.011
CH ₂ (¹ A ₁)	4.728	4.662	4.840	4.788	4.650	4.667	4.699
CH ₃	-5.860	-6.250	-5.963	-6.055	-6.158	-6.136	-6.115
Methane (CH ₄)	-2.264	-2.142	-2.042	-2.021	-2.163	-2.148	-2.117
NH	-0.853	-0.991	-1.081	-1.297	-1.235	-1.180	-1.208
NH ₂	-2.986	-2.879	-2.929	-3.229	-3.206	-3.130	-3.162
Ammonia (NH ₃)	0.751	1.287	1.068	0.844	0.839	0.917	0.888
OH	-3.456	-3.257	-3.435	-3.495	-3.524	-3.541	-3.552
Water (H ₂ O)	-0.350	-0.213	-0.616	-0.677	-0.710	-0.729	-0.741
Hydrogen fluoride (HF)	-0.088	-0.193	-0.063	0.076	0.066	0.062	0.067
SiH ₂ (¹ A ₁)	2.271	2.063	2.113	2.122	2.150	2.124	2.126
SiH ₂ (³ B ₁)	-8.257	-7.991	-8.078	-8.010	-7.973	-7.982	-7.987
SiH ₃	-5.357	-5.303	-5.309	-5.278	-5.241	-5.263	-5.264
Silane (SiH ₄)	-0.862	-0.936	-0.859	-0.860	-0.829	-0.862	-0.857
PH ₂	-4.600	-4.156	-4.151	-3.964	-3.983	-3.951	-3.946
PH ₃	-1.050	-0.479	-0.468	-0.230	-0.238	-0.193	-0.198
Hydrogen sulfide (H ₂ S)	-2.510	-1.758	-1.742	-1.739	-1.755	-1.704	-1.699
Hydrogen chloride (HCl)	-0.940	-1.684	-1.574	-1.483	-1.453	-1.451	-1.440
Li ₂	5.590	5.778	5.698	5.876	5.936	5.922	5.910
LiF	0.407	-0.044	-0.093	0.233	0.255	0.242	0.241
Acetylene (C ₂ H ₂)	-1.455	-1.399	-1.004	-0.994	-1.268	-1.244	-1.183
Ethylene (H ₂ C=CH ₂)	-4.677	-4.590	-4.269	-4.260	-4.542	-4.516	-4.453
Ethane (H ₃ C-CH ₃)	-6.106	-5.883	-5.671	-5.632	-5.919	-5.888	-5.826
CN	0.391	0.804	0.808	0.615	0.505	0.587	0.589
Hydrogen cyanide (HCN)	0.763	1.333	1.380	1.169	1.019	1.107	1.110
CO	0.181	0.357	0.067	-0.002	-0.186	-0.202	-0.176
HCO	-7.888	-7.921	-8.152	-8.224	-8.376	-8.382	-8.367
Formaldehyde (H ₂ C=O)	-4.648	-4.524	-4.727	-4.806	-4.985	-4.991	-4.972
Methanol (CH ₃ -OH)	-5.083	-4.998	-5.235	-5.306	-5.480	-5.482	-5.464
N ₂	6.234	7.327	7.051	6.615	6.576	6.726	6.674
Hydrazine (H ₂ N-NH ₂)	-0.521	0.456	0.102	-0.381	-0.386	-0.231	-0.289
NO	-1.881	-1.564	-2.124	-2.408	-2.436	-2.379	-2.421
O ₂	-11.403	-12.289	-12.406	-12.473	-12.530	-12.564	-12.579
Hydrogen peroxide (HO-OH)	-3.630	-3.467	-4.078	-4.229	-4.299	-4.340	-4.367
F ₂	-1.754	-1.752	-1.491	-1.321	-1.349	-1.362	-1.358
Carbon dioxide (CO ₂)	-12.959	-12.538	-13.247	-13.381	-13.604	-13.627	-13.618
Na ₂	2.905	2.733	2.743	2.750	2.753	2.756	2.754
Si ₂	-0.772	-0.876	-0.697	-0.558	-0.496	-0.543	-0.548
P ₂	0.979	2.047	2.052	2.576	2.552	2.639	2.630
S ₂	-11.481	-10.754	-10.767	-10.540	-10.547	-10.490	-10.496
Cl ₂	-1.633	-3.017	-2.878	-2.743	-2.708	-2.694	-2.676
NaCl	-0.494	-1.386	-1.319	-1.213	-1.165	-1.167	-1.158
Silicon monoxide (SiO)	2.798	3.055	2.592	2.555	2.544	2.494	2.488
CS	-0.461	0.227	0.447	0.404	0.233	0.296	0.338
SO	-9.778	-9.681	-9.809	-9.739	-9.796	-9.768	-9.788
ClO	-7.165	-7.607	-7.839	-7.798	-7.846	-7.846	-7.853
Chlorine monofluoride (FCl)	-2.299	-2.917	-2.765	-2.602	-2.603	-2.599	-2.588
Si ₂ H ₆	-4.643	-4.739	-4.605	-4.606	-4.540	-4.609	-4.598
Methyl chloride (CH ₃ Cl)	-5.041	-5.796	-5.529	-5.462	-5.573	-5.552	-5.513
Methanethiol (H ₃ CSH)	-6.228	-5.429	-5.292	-5.278	-5.437	-5.369	-5.334
Hypochlorous acid (HOCl)	-3.481	-3.838	-4.127	-4.134	-4.156	-4.169	-4.174
Sulfur dioxide (SO ₂)	-7.803	-6.545	-7.524	-7.604	-7.696	-7.685	-7.706
BF ₃	-11.197	-11.656	-11.687	-11.308	-11.268	-11.263	-11.262
BCl ₃	-15.681	-17.898	-17.688	-17.515	-17.320	-17.314	-17.293

Continued on next page

TABLE IX: Deviation of r++SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
AlF ₃	-3.805	-4.169	-3.939	-3.392	-3.440	-3.449	-3.434
AlCl ₃	-13.161	-14.421	-14.064	-13.786	-13.674	-13.675	-13.646
Carbon tetrafluoride (CF ₄)	-18.851	-19.633	-19.092	-18.568	-18.758	-18.747	-18.701
Carbon tetrachloride (CCl ₄)	-13.999	-16.726	-16.291	-15.959	-16.009	-15.970	-15.902
Carbon oxide sulfide (COS)	-15.533	-14.514	-14.761	-14.831	-15.034	-14.988	-14.960
Carbon bisulfide (CS ₂)	-17.056	-15.432	-15.185	-15.202	-15.391	-15.270	-15.226
Carbonic difluoride (COF ₂)	-11.785	-12.108	-12.214	-12.014	-12.210	-12.217	-12.192
Silicon tetrafluoride (SiF ₄)	-0.463	-1.920	-1.388	-0.728	-0.745	-0.787	-0.763
Silicon tetrachloride (SiCl ₄)	-11.104	-14.421	-13.801	-13.431	-13.258	-13.297	-13.245
Dinitrogen monoxide (N ₂ O)	-10.286	-8.796	-9.550	-10.086	-10.142	-10.006	-10.075
Nitrogen chloride oxide (ClNO)	-11.110	-11.148	-11.617	-11.830	-11.850	-11.793	-11.822
Nitrogen trifluoride (NF ₃)	-16.995	-17.378	-17.000	-16.885	-16.923	-16.862	-16.878
PF ₃	-3.179	-3.671	-3.312	-2.524	-2.561	-2.533	-2.526
O ₃	-9.781	-9.518	-10.615	-10.779	-10.880	-10.945	-10.986
F ₂ O	-10.388	-10.567	-10.543	-10.397	-10.458	-10.492	-10.502
Chlorine trifluoride (ClF ₃)	-20.572	-21.768	-21.407	-20.883	-20.920	-20.922	-20.902
Tetrafluoro Ethene (F ₂ C=CF ₂)	-28.348	-29.001	-28.381	-27.875	-28.187	-28.169	-28.095
Tetrachloro Ethene (C ₂ Cl ₄)	-22.399	-24.677	-24.059	-23.719	-23.900	-23.850	-23.753
Acetonitrile, trifluoro- (CF ₃ CN)	-17.811	-17.845	-17.365	-17.176	-17.504	-17.404	-17.359
Propyne (C ₃ H ₄)	-7.622	-7.414	-6.932	-6.899	-7.316	-7.276	-7.185
Allene (C ₃ H ₄)	-11.750	-11.575	-11.098	-11.064	-11.487	-11.451	-11.356
Cyclopropene (C ₃ H ₄)	-8.820	-8.520	-8.148	-8.110	-8.534	-8.491	-8.399
Propylene (C ₃ H ₆)	-9.475	-9.159	-8.759	-8.725	-9.150	-9.108	-9.015
Cyclopropane (C ₃ H ₆)	-11.668	-11.151	-10.897	-10.832	-11.261	-11.215	-11.123
Propane (C ₃ H ₈)	-10.289	-9.570	-9.306	-9.247	-9.678	-9.631	-9.538
Trans-1,3-butadiene (C ₄ H ₆)	-13.290	-13.008	-12.393	-12.368	-12.933	-12.881	-12.755
Dimethylacetylene (C ₄ H ₆)	-12.467	-12.109	-11.541	-11.487	-12.046	-11.990	-11.868
Methylenecyclopropane (C ₄ H ₆)	-18.476	-17.857	-17.399	-17.336	-17.904	-17.847	-17.723
Bicyclo[1.1.0]butane (C ₄ H ₆)	-15.592	-14.970	-14.620	-14.520	-15.096	-15.035	-14.911
Cyclobutene (C ₄ H ₆)	-12.514	-11.931	-11.440	-11.384	-11.957	-11.899	-11.774
Cyclobutane (C ₄ H ₈)	-14.908	-13.716	-13.357	-13.287	-13.860	-13.798	-13.674
Isobutene (C ₄ H ₈)	-13.927	-13.113	-12.641	-12.584	-13.153	-13.095	-12.971
Trans-butane(C ₄ H ₁₀)	-13.664	-13.115	-12.670	-12.601	-13.179	-13.115	-12.991
Isobutane (C ₄ H ₁₀)	-13.063	-12.663	-12.311	-12.240	-12.817	-12.754	-12.630
Spiropentane (C ₅ H ₈)	-22.358	-21.339	-20.915	-20.805	-21.520	-21.443	-21.289
Benzene (C ₆ H ₆)	-26.044	-25.411	-24.469	-24.432	-25.281	-25.204	-25.015
Difluoromethane (CH ₂ F ₂)	-9.161	-9.702	-9.367	-9.113	-9.275	-9.262	-9.225
Trifluoromethane(CHF ₃)	-13.748	-14.450	-13.982	-13.574	-13.751	-13.740	-13.698
CH ₂ Cl ₂	-8.111	-9.446	-9.100	-8.945	-9.030	-9.005	-8.956
CHCl ₃	-10.953	-13.071	-12.669	-12.423	-12.488	-12.457	-12.398
Methylamine (H ₃ C-NH ₂)	-3.048	-2.408	-2.502	-2.724	-2.871	-2.777	-2.775
Acetonitrile (CH ₃ -CN)	-5.338	-4.620	-4.478	-4.670	-4.962	-4.858	-4.825
Nitromethane (CH ₃ -NO ₂)	-18.864	-17.696	-18.536	-18.903	-19.124	-19.069	-19.094
Methyl nitrite (CH ₃ -O-N=O)	-15.518	-14.599	-15.420	-15.793	-16.017	-15.964	-15.986
Methyl silane (CH ₃ SiH ₃)	-4.059	-3.961	-3.791	-3.765	-3.877	-3.895	-3.859
Formic acid (HCOOH)	-10.948	-10.634	-11.345	-11.477	-11.688	-11.712	-11.704
Methyl formate (HCOOCH ₃)	-16.187	-15.806	-16.326	-16.481	-16.831	-16.838	-16.801
Acetamide (CH ₃ CONH ₂)	-14.369	-13.247	-13.606	-13.892	-14.214	-14.126	-14.105
Aziridine (C ₂ H ₄ NH)	-9.348	-8.329	-8.329	-8.533	-8.824	-8.716	-8.684
Cyanogen (NCCN)	-4.182	-2.986	-2.921	-3.341	-3.639	-3.460	-3.457
Dimethylamine ((CH ₃) ₂ NH)	-7.252	-6.390	-6.377	-6.592	-6.881	-6.772	-6.740
Trans ethylamine (CH ₃ CH ₂ NH ₂)	-8.248	-7.383	-7.345	-7.550	-7.841	-7.731	-7.699
Ketene (CH ₂ CO)	-14.092	-13.782	-13.926	-13.961	-14.280	-14.275	-14.222
Oxirane (C ₂ H ₄ O)	-11.319	-11.021	-11.144	-11.212	-11.533	-11.521	-11.471
Acetaldehyde (CH ₃ CHO)	-10.168	-9.822	-9.942	-9.994	-10.315	-10.307	-10.256
Glyoxal (HCOCOH)	-12.844	-12.442	-12.821	-12.975	-13.333	-13.349	-13.309
Ethanol (CH ₃ CH ₂ OH)	-9.055	-8.805	-8.945	-8.997	-9.316	-9.302	-9.253

Continued on next page

TABLE IX: Deviation of r++SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Dimethylether (CH ₃ OCH ₃)	-9.661	-9.374	-9.459	-9.539	-9.856	-9.842	-9.794
Thiirane (C ₂ H ₄ S)	-12.469	-11.492	-11.221	-11.204	-11.506	-11.423	-11.358
Dimethyl sulfoxide ((CH ₃) ₂ SO)	-15.753	-14.415	-14.638	-14.641	-14.984	-14.920	-14.867
Ethanethiol (C ₂ H ₅ SH)	-9.822	-8.848	-8.604	-8.573	-8.877	-8.794	-8.728
Dimethyl sulfide (CH ₃ SCH ₃)	-10.579	-9.544	-9.323	-9.287	-9.590	-9.508	-9.443
Vinyl fluoride (CH ₂ =CHF)	-10.691	-10.821	-10.416	-10.274	-10.562	-10.538	-10.473
Ethyl chloride (C ₂ H ₅ Cl)	-9.359	-9.894	-9.514	-9.428	-9.682	-9.647	-9.576
Vinyl chloride (CH ₂ =CHCl)	-12.803	-13.357	-12.929	-12.840	-13.092	-13.060	-12.989
Acrylonitrile (CH ₂ =CHCN)	-6.961	-6.194	-5.855	-6.047	-6.480	-6.365	-6.300
Acetone (CH ₃ COCH ₃)	-15.327	-14.471	-14.516	-14.543	-15.007	-14.984	-14.902
Acetic acid (CH ₃ COOH)	-15.723	-15.030	-15.646	-15.758	-16.112	-16.121	-16.082
Acetyl fluoride (CH ₃ COF)	-15.326	-15.276	-15.333	-15.239	-15.569	-15.561	-15.508
CH ₃ COCl (acetyl chloride)	-16.324	-16.800	-16.825	-16.798	-17.086	-17.075	-17.015
CH ₃ CH ₂ CH ₂ Cl (propyl chloride)	-13.508	-13.669	-13.232	-13.119	-13.517	-13.466	-13.364
Isopropanol (CH ₃) ₂ CHOH)	-13.080	-12.745	-12.798	-12.825	-13.286	-13.257	-13.177
Methyl ethyl ether (C ₂ H ₅ OCH ₃)	-14.131	-13.631	-13.616	-13.672	-14.132	-14.102	-14.023
Trimethylamine ((CH ₃) ₃ N)	-11.592	-10.778	-10.672	-10.863	-11.297	-11.173	-11.109
Furan (C ₄ H ₄ O)	-21.071	-20.603	-20.498	-20.504	-21.104	-21.071	-20.957
C ₄ H ₄ S (thiophene)	-21.697	-20.468	-19.897	-19.832	-20.419	-20.315	-20.184
Pyrrole (C ₄ H ₅ N)	-20.059	-18.921	-18.638	-18.808	-19.377	-19.245	-19.149
Pyridine (C ₅ H ₅ N)	-24.573	-23.464	-22.857	-23.059	-23.775	-23.633	-23.503
H ₂	2.073	2.089	2.089	2.089	2.089	2.089	2.089
HS	-3.066	-2.643	-2.663	-2.684	-2.689	-2.660	-2.658
CCH	-4.227	-4.357	-3.965	-3.916	-4.146	-4.128	-4.069
C ₂ H ₃ (² A')	-10.086	-10.228	-9.839	-9.866	-10.109	-10.085	-10.028
CH ₃ CO (² A')	-13.279	-13.338	-13.398	-13.489	-13.778	-13.767	-13.722
H ₂ COH (² A)	-9.367	-9.551	-9.767	-9.862	-10.004	-10.005	-9.993
CH ₃ O (² A')	-10.187	-9.948	-10.027	-10.086	-10.257	-10.259	-10.240
CH ₃ CH ₂ O (² A'')	-15.533	-15.082	-15.067	-15.111	-15.427	-15.413	-15.362
CH ₃ S (² A')	-8.667	-8.121	-8.032	-8.029	-8.177	-8.131	-8.099
C ₂ H ₅ (² A')	-10.720	-10.913	-10.540	-10.607	-10.855	-10.817	-10.765
(CH ₃) ₂ CH (² A')	-15.963	-15.685	-15.243	-15.291	-15.680	-15.628	-15.546
(CH ₃) ₂ CH (² A')	-15.963	-15.685	-15.243	-15.291	-15.680	-15.628	-15.546
NO ₂	-16.569	-15.876	-16.891	-17.234	-17.298	-17.261	-17.316
Methyl allene (C ₄ H ₆)	-15.774	-15.228	-14.668	-14.611	-15.179	-15.127	-15.001
Isoprene (C ₅ H ₈)	-17.207	-16.677	-15.965	-15.915	-16.625	-16.556	-16.399
Cyclopentane (C ₅ H ₁₀)	-17.917	-17.237	-16.793	-16.712	-17.435	-17.356	-17.201
n-Pentane (C ₅ H ₁₂)	-17.848	-16.335	-15.947	-15.836	-16.555	-16.476	-16.321
Neo pentane (C ₅ H ₁₂)	-16.180	-15.562	-15.110	-15.036	-15.757	-15.679	-15.523
1,4 Cyclohexadiene (C ₆ H ₈)	-21.149	-20.316	-19.526	-19.439	-20.294	-20.209	-20.022
Cyclohexane (C ₆ H ₁₂)	-20.944	-20.243	-19.722	-19.611	-20.482	-20.388	-20.201
n-Hexane (C ₆ H ₁₄)	-21.099	-20.168	-19.483	-19.381	-20.246	-20.148	-19.963
3-Methyl pentane (C ₆ H ₁₄)	-19.781	-19.079	-18.486	-18.366	-19.233	-19.137	-18.951
Toluene (C ₆ H ₅ CH ₃)	-30.042	-29.290	-28.239	-28.172	-29.167	-29.074	-28.853
n-Heptane (C ₇ H ₁₆)	-25.715	-23.392	-22.884	-22.720	-23.727	-23.617	-23.400
Cyclooctatetraene (C ₈ H ₈)	-28.642	-27.776	-26.610	-26.532	-27.668	-27.563	-27.311
n-Octane (C ₈ H ₁₈)	-28.609	-27.093	-26.207	-26.068	-27.214	-27.084	-26.836
Naphthalene (C ₁₀ H ₈)	-46.605	-45.655	-44.124	-44.048	-45.465	-45.337	-45.020
Azulene (C ₁₀ H ₈)	-47.155	-46.029	-44.514	-44.443	-45.862	-45.732	-45.415
Acetic acid methyl ester (CH ₃ COOCH ₃)	-19.798	-19.079	-19.516	-19.645	-20.140	-20.132	-20.064
t-Butanol ((CH ₃) ₃ COH)	-16.521	-16.059	-16.020	-16.032	-16.640	-16.596	-16.484
Aniline (C ₆ H ₅ NH ₂)	-30.187	-28.810	-28.120	-28.298	-29.151	-28.994	-28.834
Phenol (C ₆ H ₅ OH)	-30.749	-29.985	-29.469	-29.483	-30.364	-30.305	-30.129
Divinyl ether (C ₄ H ₆ O)	-18.705	-18.144	-17.950	-17.978	-18.573	-18.537	-18.426
Tetrahydrofuran (C ₄ H ₈ O)	-17.186	-16.967	-16.876	-16.916	-17.519	-17.473	-17.363
Cyclopentanone (C ₅ H ₈ O)	-23.906	-23.347	-23.213	-23.201	-23.952	-23.897	-23.752
Benzoquinone(C ₆ H ₄ O ₂)	-31.519	-30.772	-30.514	-30.628	-31.554	-31.517	-31.350

Continued on next page

TABLE IX: Deviation of r++SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Pyrimidine (C ₄ H ₄ N ₂)	-24.177	-22.632	-22.326	-22.779	-23.363	-23.156	-23.086
Dimethyl sulphone (C ₂ H ₆ O ₂ S)	-19.945	-18.052	-18.889	-18.944	-19.323	-19.275	-19.233
Chlorobenzene (C ₆ H ₅ Cl)	-30.220	-30.393	-29.341	-29.223	-30.041	-29.960	-29.762
Butanedinitrile (NC-CH ₂ -CH ₂ -CN)	-10.498	-9.125	-8.865	-9.241	-9.830	-9.621	-9.555
Pyrazine (C ₄ H ₄ N ₂)	-20.185	-18.615	-18.351	-18.794	-19.375	-19.170	-19.099
Acetyl acetylene (CH ₃ -C(=O)-CCH)	-13.778	-13.404	-13.176	-13.200	-13.800	-13.768	-13.656
Crotonaldehyde (CH ₃ -CH=CH-CHO)	-19.893	-19.311	-19.140	-19.180	-19.785	-19.750	-19.636
Acetic anhydride (CH ₃ -C(=O)-O-C(=O)-CH ₃)	-30.369	-29.603	-30.380	-30.549	-31.223	-31.221	-31.134
2,5-Dihydrothiophene (C ₄ H ₆ S)	-18.813	-17.732	-17.200	-17.145	-17.733	-17.623	-17.495
Isobutane nitrile ((CH ₃) ₂ CH-CN)	-10.808	-9.632	-9.316	-9.478	-10.059	-9.924	-9.829
Methyl ethyl ketone (CH ₃ -CO-CH ₂ -CH ₃)	-18.291	-18.139	-17.994	-18.020	-18.626	-18.587	-18.474
Isobutanal ((CH ₃) ₂ CH-CHO)	-16.310	-15.636	-15.551	-15.573	-16.181	-16.142	-16.028
1,4-Dioxane (C ₄ H ₈ O ₂)	-22.035	-21.605	-21.792	-21.931	-22.566	-22.538	-22.440
Tetrahydrothiophene (C ₄ H ₈ S)	-17.579	-16.613	-16.160	-16.096	-16.690	-16.574	-16.446
t-Butyl chloride ((CH ₃) ₃ C-Cl)	-16.867	-17.386	-16.834	-16.718	-17.260	-17.194	-17.060
n-Butyl chloride (CH ₃ -CH ₂ -CH ₂ -CH ₂ -Cl)	-16.550	-16.583	-16.029	-15.910	-16.454	-16.386	-16.253
Tetrahydropyrrole (C ₄ H ₈ NH)	-15.778	-14.568	-14.392	-14.566	-15.148	-15.006	-14.911
Nitro-s-butane (CH ₃ -CH ₂ -CH(CH ₃)-NO ₂)	-29.216	-27.807	-28.311	-28.631	-29.287	-29.184	-29.115
Diethyl ether (CH ₃ -CH ₂ -O-CH ₂ -CH ₃)	-17.632	-16.921	-16.852	-16.886	-17.492	-17.446	-17.336
Dimethyl acetal (CH ₃ -CH(OCH ₃) ₂)	-21.228	-20.631	-20.865	-21.008	-21.647	-21.619	-21.521
t-Butanethiol ((CH ₃) ₃ C-SH)	-16.945	-15.811	-15.399	-15.332	-15.922	-15.808	-15.679
Diethyl disulfide (CH ₃ -CH ₂ -S-S-CH ₂ -CH ₃)	-21.923	-19.647	-19.208	-19.122	-19.727	-19.560	-19.429
t-Butylamine ((CH ₃) ₃ C-NH ₂)	-14.887	-13.222	-13.033	-13.182	-13.758	-13.617	-13.522
Tetramethylsilane (Si(CH ₃) ₄)	-13.256	-12.603	-12.182	-12.077	-12.620	-12.592	-12.462
2-Methyl thiophene (C ₅ H ₆ S)	-25.636	-24.443	-23.794	-23.719	-24.450	-24.330	-24.168
N-methyl pyrrole (cyc-C ₄ H ₄ N-CH ₃)	-23.521	-22.323	-21.919	-22.081	-22.791	-22.644	-22.518
Tetrahydropyran (C ₅ H ₁₀ O)	-21.386	-20.890	-20.656	-20.679	-21.433	-21.372	-21.230
Diethyl ketone (CH ₃ -CH ₂ -CO-CH ₂ -CH ₃)	-23.584	-22.308	-22.198	-22.180	-22.934	-22.878	-22.734
Isopropyl acetate (CH ₃ -C(=O)-O-CH(CH ₃) ₂)	-27.359	-26.446	-26.707	-26.800	-27.581	-27.543	-27.412
Tetrahydrothiopyran (C ₅ H ₁₀ S)	-21.864	-20.484	-19.919	-19.837	-20.576	-20.445	-20.286
Piperidine (cyc-C ₅ H ₁₀ NH)	-18.710	-17.685	-17.432	-17.574	-18.301	-18.145	-18.019
t-Butyl methyl ether ((CH ₃) ₃ C-O-CH ₃)	-20.821	-19.840	-19.642	-19.674	-20.417	-20.357	-20.216
1,3-Difluorobenzene (C ₆ H ₄ F ₂)	-35.899	-35.808	-34.612	-34.315	-35.176	-35.104	-34.909
1,4-Difluorobenzene (C ₆ H ₄ F ₂)	-37.526	-37.314	-36.204	-35.897	-36.759	-36.686	-36.491
Fluorobenzene (C ₆ H ₅ F)	-31.159	-30.795	-29.730	-29.560	-30.415	-30.340	-30.148
Di-isopropyl ether ((CH ₃) ₂ CH-O-CH(CH ₃) ₂)	-24.876	-23.664	-23.378	-23.362	-24.257	-24.181	-24.007
PF ₅	-9.379	-10.134	-9.624	-8.594	-8.647	-8.623	-8.610
SF ₆	-25.684	-25.637	-24.851	-23.966	-24.041	-24.011	-23.993
P ₄	-10.909	-8.759	-8.674	-7.608	-7.651	-7.471	-7.492
SO ₃	-15.346	-13.749	-15.186	-15.350	-15.483	-15.488	-15.522
SCl ₂	-7.287	-8.409	-8.174	-7.979	-7.936	-7.876	-7.857
POCl ₃	-14.325	-15.091	-15.175	-14.729	-14.666	-14.645	-14.634
PCL ₅	-21.155	-23.761	-23.122	-22.463	-22.294	-22.248	-22.210
Cl ₂ O ₂ S	-18.727	-19.052	-19.796	-19.715	-19.747	-19.721	-19.726
PCL ₃	-10.382	-11.643	-11.192	-10.693	-10.586	-10.546	-10.522
Cl ₂ S ₂	-18.474	-18.891	-18.559	-18.374	-18.336	-18.230	-18.203
SiCl ₂ singlet	-4.005	-5.949	-5.575	-5.392	-5.286	-5.321	-5.293
CF ₃ Cl	-19.274	-20.646	-20.117	-19.640	-19.792	-19.774	-19.723
Hexafluoro ethane (C ₂ F ₆)	-32.258	-33.547	-32.634	-31.831	-32.187	-32.166	-32.080
CF ₃	-20.093	-20.797	-20.412	-20.022	-20.153	-20.146	-20.110
C ₆ H ₅ (phenyl radical)	-31.721	-31.451	-30.436	-30.429	-31.241	-31.170	-30.984
Bicyclo[1.1.0]butane (C ₄ H ₆)	-15.592	-14.970	-14.620	-14.520	-15.096	-15.035	-14.911
(CH ₃) ₃ C (t-butyl radical)	-19.292	-19.184	-18.611	-18.618	-19.155	-19.087	-18.973
Trans-butane(C ₄ H ₁₀)	-13.664	-13.115	-12.670	-12.601	-13.179	-13.115	-12.991
1,3 Cyclohexadiene (C ₆ H ₈)	-21.677	-21.115	-20.322	-20.236	-21.092	-21.007	-20.819
ME	-13.182	-12.896	-12.722	-12.683	-13.025	-12.977	-12.912
MAE	13.487	13.237	13.053	13.012	13.349	13.305	13.239

TABLE X: Deviation of r²SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
LiH	2.455	2.525	2.484	2.553	2.586	2.583	2.575
BeH	-10.418	-10.465	-10.225	-10.267	-10.247	-10.233	-10.242
CH	2.617	2.454	2.557	2.508	2.417	2.440	2.455
CH ₂ (³ B ₁)	-6.222	-6.642	-6.398	-6.398	-6.469	-6.455	-6.439
CH ₂ (¹ A ₁)	5.934	5.708	5.938	5.884	5.748	5.769	5.797
CH ₃	-3.602	-4.222	-3.879	-3.970	-4.069	-4.042	-4.025
Methane (CH ₄)	0.930	0.535	0.881	0.849	0.707	0.723	0.751
NH	-0.347	-0.728	-0.764	-0.953	-0.901	-0.853	-0.881
NH ₂	-1.531	-1.933	-1.805	-2.092	-2.090	-2.019	-2.051
Ammonia (NH ₃)	3.638	3.297	3.479	3.203	3.174	3.244	3.215
OH	-2.582	-2.388	-2.407	-2.543	-2.561	-2.583	-2.593
Water (H ₂ O)	1.811	2.105	2.018	1.792	1.758	1.737	1.725
Hydrogen fluoride (HF)	1.773	1.793	1.767	1.848	1.848	1.838	1.842
SiH ₂ (¹ A ₁)	2.613	2.510	2.611	2.564	2.597	2.573	2.574
SiH ₂ (³ B ₁)	-7.694	-7.426	-7.511	-7.461	-7.422	-7.426	-7.434
SiH ₃	-4.661	-4.577	-4.556	-4.553	-4.513	-4.528	-4.532
Silane (SiH ₄)	0.017	-0.035	0.082	0.053	0.087	0.060	0.062
PH ₂	-3.849	-3.382	-3.677	-3.454	-3.506	-3.471	-3.467
PH ₃	0.207	0.872	0.545	0.761	0.732	0.778	0.772
Hydrogen sulfide (H ₂ S)	-0.578	-0.269	-0.693	-0.378	-0.494	-0.401	-0.406
Hydrogen chloride (HCl)	-0.364	-0.455	-0.394	-0.421	-0.502	-0.464	-0.462
Li ₂	5.642	5.908	5.840	5.970	6.033	6.028	6.013
LiF	2.359	2.194	1.980	2.189	2.228	2.210	2.208
Acetylene (C ₂ H ₂)	2.694	2.322	2.918	2.875	2.621	2.646	2.701
Ethylene (H ₂ C=CH ₂)	0.502	-0.097	0.510	0.464	0.196	0.227	0.282
Ethane (H ₃ C-CH ₃)	0.133	-0.693	0.037	-0.046	-0.331	-0.298	-0.243
CN	3.139	2.495	3.001	2.728	2.605	2.675	2.674
Hydrogen cyanide (HCN)	4.155	3.827	4.298	4.027	3.860	3.946	3.944
CO	2.678	2.708	2.927	2.648	2.476	2.458	2.482
HCO	-4.878	-4.927	-4.753	-5.013	-5.158	-5.167	-5.154
Formaldehyde (H ₂ C=O)	-1.017	-0.962	-0.765	-1.023	-1.196	-1.204	-1.188
Methanol (CH ₃ -OH)	-0.286	-0.406	-0.171	-0.442	-0.614	-0.617	-0.602
N ₂	8.758	8.531	8.813	8.336	8.265	8.406	8.351
Hydrazine (H ₂ N-NH ₂)	4.624	4.073	4.417	3.846	3.799	3.941	3.883
NO	-0.034	0.070	-0.016	-0.462	-0.518	-0.465	-0.507
O ₂	-9.598	-9.716	-9.795	-10.006	-10.055	-10.099	-10.112
Hydrogen peroxide (HO-OH)	-0.413	0.195	-0.043	-0.436	-0.497	-0.548	-0.572
F ₂	0.465	0.228	0.271	0.484	0.456	0.436	0.440
Carbon dioxide (CO ₂)	-8.180	-7.765	-7.609	-8.131	-8.342	-8.371	-8.364
Na ₂	3.115	2.930	2.938	2.944	2.947	2.950	2.948
Si ₂	-0.369	-0.224	0.106	0.141	0.218	0.171	0.164
P ₂	2.182	3.536	2.905	3.268	3.224	3.305	3.297
S ₂	-9.481	-9.097	-9.612	-9.201	-9.282	-9.181	-9.200
Cl ₂	-0.944	-1.299	-1.041	-1.152	-1.257	-1.195	-1.187
NaCl	0.072	-0.074	-0.098	-0.130	-0.212	-0.171	-0.173
Silicon monoxide (SiO)	4.492	4.879	4.916	4.646	4.631	4.585	4.578
CS	2.141	2.243	2.091	2.321	2.072	2.161	2.196
SO	-7.749	-7.440	-7.626	-7.635	-7.722	-7.686	-7.707
ClO	-6.162	-5.928	-5.873	-6.090	-6.198	-6.166	-6.183
Chlorine monofluoride (FCl)	-0.668	-0.943	-0.783	-0.753	-0.785	-0.771	-0.764
Si ₂ H ₆	-2.754	-2.763	-2.537	-2.615	-2.538	-2.598	-2.592
Methyl chloride (CH ₃ Cl)	-1.782	-2.225	-1.835	-1.923	-2.135	-2.075	-2.047
Methanethiol (H ₃ CSH)	-1.391	-1.469	-1.541	-1.288	-1.538	-1.429	-1.407
Hypochlorous acid (HOCl)	-1.499	-1.198	-1.168	-1.432	-1.508	-1.504	-1.512
Sulfur dioxide (SO ₂)	-3.119	-2.012	-2.525	-2.854	-3.003	-2.974	-2.999
BF ₃	-4.514	-4.431	-4.872	-4.769	-4.687	-4.711	-4.708
BCl ₃	-12.435	-12.660	-12.654	-12.843	-13.026	-12.912	-12.915

Continued on next page

TABLE X: Deviation of r^2 SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
AlF ₃	1.845	2.048	1.787	2.061	2.093	2.047	2.062
AlCl ₃	-11.011	-10.175	-10.142	-10.186	-10.453	-10.333	-10.331
Carbon tetrafluoride (CF ₄)	-9.966	-10.598	-10.463	-10.241	-10.383	-10.395	-10.351
Carbon tetrachloride (CCl ₄)	-9.515	-10.331	-9.643	-9.901	-10.344	-10.166	-10.133
Carbon oxide sulfide (COS)	-10.578	-10.206	-10.439	-10.359	-10.665	-10.579	-10.563
Carbon bisulfide (CS ₂)	-11.811	-11.458	-12.059	-11.417	-11.802	-11.611	-11.583
Carbonic difluoride (COF ₂)	-4.974	-5.166	-5.067	-5.218	-5.385	-5.408	-5.385
Silicon tetrafluoride (SiF ₄)	7.092	6.459	6.457	6.671	6.755	6.672	6.698
Silicon tetrachloride (SiCl ₄)	-7.978	-8.472	-8.158	-8.287	-8.618	-8.500	-8.481
Dinitrogen monoxide (N ₂ O)	-5.466	-5.469	-5.292	-6.007	-6.108	-5.987	-6.055
Nitrogen chloride oxide (ClNO)	-8.059	-8.004	-7.877	-8.417	-8.575	-8.484	-8.521
Nitrogen trifluoride (NF ₃)	-11.109	-11.895	-11.684	-11.722	-11.751	-11.711	-11.727
PF ₃	2.528	2.831	2.514	2.982	2.969	2.976	2.982
O ₃	-6.384	-5.387	-5.721	-6.347	-6.459	-6.533	-6.572
F ₂ O	-6.925	-6.840	-7.013	-6.973	-7.022	-7.071	-7.080
Chlorine trifluoride (ClF ₃)	-15.604	-15.938	-15.685	-15.556	-15.613	-15.600	-15.592
Tetrafluoro Ethene (F ₂ C=CF ₂)	-17.332	-18.071	-17.792	-17.584	-17.836	-17.841	-17.773
Tetrachloro Ethene (C ₂ Cl ₄)	-15.540	-16.140	-15.151	-15.455	-16.030	-15.831	-15.776
Acetonitrile, trifluoro- (CF ₃ CN)	-7.114	-8.077	-7.405	-7.522	-7.832	-7.750	-7.712
Propyne (C ₃ H ₄)	-0.243	-1.019	-0.069	-0.152	-0.547	-0.505	-0.423
Allene (C ₃ H ₄)	-4.256	-5.099	-4.186	-4.245	-4.641	-4.599	-4.516
Cyclopropene (C ₃ H ₄)	-1.295	-2.251	-1.301	-1.397	-1.804	-1.756	-1.675
Propylene (C ₃ H ₆)	-1.100	-2.044	-1.074	-1.161	-1.572	-1.526	-1.443
Cyclopropane (C ₃ H ₆)	-3.005	-4.101	-3.048	-3.170	-3.596	-3.550	-3.467
Propane (C ₃ H ₈)	-0.916	-1.822	-0.769	-0.894	-1.322	-1.274	-1.190
Trans-1,3-butadiene (C ₄ H ₆)	-2.837	-3.977	-2.761	-2.855	-3.391	-3.329	-3.219
Dimethylacetylene (C ₄ H ₆)	-1.878	-3.064	-1.758	-1.883	-2.419	-2.360	-2.251
Methylenecyclopropane (C ₄ H ₆)	-7.655	-8.869	-7.554	-7.682	-8.234	-8.173	-8.062
Bicyclo[1.1.0]butane (C ₄ H ₆)	-4.610	-6.161	-4.724	-4.879	-5.454	-5.395	-5.282
Cyclobutene (C ₄ H ₆)	-1.505	-2.767	-1.401	-1.554	-2.109	-2.048	-1.936
Cyclobutane (C ₄ H ₈)	-2.978	-3.943	-2.503	-2.696	-3.264	-3.201	-3.088
Isobutene (C ₄ H ₈)	-2.332	-3.357	-2.031	-2.164	-2.718	-2.657	-2.545
Trans-butane(C ₄ H ₁₀)	-1.217	-2.799	-1.289	-1.475	-2.049	-1.983	-1.871
Isobutane (C ₄ H ₁₀)	-0.575	-2.312	-0.894	-1.075	-1.648	-1.584	-1.472
Spiropentane (C ₅ H ₈)	-8.169	-9.864	-8.099	-8.301	-9.013	-8.936	-8.796
Benzene (C ₆ H ₆)	-10.385	-11.955	-10.110	-10.242	-11.058	-10.963	-10.798
Difluoromethane (CH ₂ F ₂)	-3.466	-4.033	-3.868	-3.770	-3.903	-3.903	-3.869
Trifluoromethane(CHF ₃)	-6.480	-7.101	-6.913	-6.749	-6.885	-6.894	-6.854
CH ₂ Cl ₂	-4.538	-4.932	-4.474	-4.613	-4.900	-4.799	-4.771
CHCl ₃	-6.962	-7.600	-7.035	-7.233	-7.599	-7.459	-7.429
Methylamine (H ₃ C-NH ₂)	2.688	2.003	2.538	2.206	2.039	2.127	2.125
Acetonitrile (CH ₃ -CN)	1.275	0.554	1.378	1.067	0.759	0.862	0.888
Nitromethane (CH ₃ -NO ₂)	-10.901	-10.682	-10.352	-11.123	-11.367	-11.324	-11.350
Methyl nitrite (CH ₃ -O-N=O)	-8.080	-7.915	-7.616	-8.396	-8.638	-8.591	-8.617
Methyl silane (CH ₃ SiH ₃)	0.117	-0.335	0.164	0.093	-0.016	-0.029	0.001
Formic acid (HCOOH)	-5.160	-4.901	-4.765	-5.280	-5.485	-5.512	-5.506
Methyl formate (HCOOCH ₃)	-7.722	-7.706	-7.256	-7.816	-8.157	-8.164	-8.133
Acetamide (CH ₃ CONH ₂)	-4.447	-4.906	-4.142	-4.723	-5.068	-4.989	-4.972
Aziridine (C ₂ H ₄ NH)	-1.449	-2.254	-1.381	-1.755	-2.061	-1.958	-1.934
Cyanogen (NCCN)	2.658	2.010	2.950	2.395	2.062	2.239	2.234
Dimethylamine ((CH ₃) ₂ NH)	1.448	0.498	1.381	0.997	0.690	0.795	0.820
Trans ethylamine (CH ₃ CH ₂ NH ₂)	0.619	-0.396	0.549	0.168	-0.143	-0.040	-0.014
Ketene (CH ₂ CO)	-7.915	-8.169	-7.631	-7.914	-8.218	-8.214	-8.168
Oxirane (C ₂ H ₄ O)	-4.507	-4.889	-4.332	-4.642	-4.954	-4.945	-4.901
Acetaldehyde (CH ₃ CHO)	-3.291	-3.583	-3.024	-3.315	-3.632	-3.624	-3.579
Glyoxal (HCOCOH)	-5.533	-5.294	-4.863	-5.374	-5.719	-5.739	-5.704
Ethanol (CH ₃ CH ₂ OH)	-1.135	-1.631	-1.024	-1.338	-1.656	-1.642	-1.599

Continued on next page

TABLE X: Deviation of r²SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Dimethylether (CH ₃ OCH ₃)	-2.107	-2.428	-1.858	-2.177	-2.490	-2.476	-2.433
Thiirane (C ₂ H ₄ S)	-5.363	-5.705	-5.469	-5.253	-5.644	-5.526	-5.473
Dimethyl sulfoxide ((CH ₃) ₂ SO)	-5.964	-5.917	-5.636	-5.758	-6.160	-6.072	-6.031
Ethanethiol (C ₂ H ₅ SH)	-1.897	-2.328	-2.030	-1.820	-2.215	-2.091	-2.041
Dimethyl sulfide (CH ₃ SCH ₃)	-2.726	-3.069	-2.798	-2.603	-2.991	-2.869	-2.820
Vinyl fluoride (CH ₂ =CHF)	-4.086	-4.722	-4.195	-4.173	-4.433	-4.413	-4.355
Ethyl chloride (C ₂ H ₅ Cl)	-2.990	-3.731	-2.978	-3.108	-3.465	-3.390	-3.334
Vinyl chloride (CH ₂ =CHCl)	-7.275	-7.842	-7.155	-7.259	-7.601	-7.528	-7.474
Acrylonitrile (CH ₂ =CHCN)	1.690	0.821	1.894	1.579	1.144	1.261	1.313
Acetone (CH ₃ COCH ₃)	-5.211	-5.563	-4.656	-4.985	-5.444	-5.421	-5.348
Acetic acid (CH ₃ COOH)	-6.737	-6.633	-6.155	-6.703	-7.054	-7.064	-7.031
Acetyl fluoride (CH ₃ COF)	-6.927	-7.363	-6.854	-7.092	-7.402	-7.404	-7.355
CH ₃ COCl (acetyl chloride)	-9.045	-9.490	-8.846	-9.207	-9.602	-9.553	-9.508
CH ₃ CH ₂ CH ₂ Cl (propyl chloride)	-4.021	-4.952	-3.869	-4.035	-4.534	-4.443	-4.359
Isopropanol (CH ₃) ₂ CHOH)	-1.980	-2.946	-1.978	-2.332	-2.792	-2.764	-2.692
Methyl ethyl ether (C ₂ H ₅ OCH ₃)	-3.444	-4.095	-3.153	-3.511	-3.967	-3.936	-3.866
Trimethylamine ((CH ₃) ₃ N)	0.129	-1.352	-0.131	-0.551	-1.000	-0.880	-0.826
Furan (C ₄ H ₄ O)	-9.116	-10.053	-8.913	-9.250	-9.824	-9.781	-9.683
C ₄ H ₄ S (thiophene)	-9.462	-10.394	-9.482	-9.295	-9.953	-9.800	-9.695
Pyrrrole (C ₄ H ₅ N)	-6.809	-8.251	-6.834	-7.205	-7.782	-7.645	-7.565
Pyridine (C ₅ H ₅ N)	-9.667	-11.128	-9.447	-9.821	-10.531	-10.379	-10.270
H ₂	2.073	2.089	2.088	2.088	2.088	2.088	2.088
HS	-2.202	-2.059	-2.287	-2.103	-2.174	-2.112	-2.118
CCH	-0.726	-1.295	-0.734	-0.715	-0.929	-0.910	-0.859
C ₂ H ₃ (² A')	-5.657	-6.310	-5.731	-5.791	-6.016	-5.990	-5.940
CH ₃ CO (² A')	-7.132	-7.683	-7.120	-7.443	-7.723	-7.713	-7.674
H ₂ COH (² A)	-5.246	-5.483	-5.298	-5.579	-5.718	-5.717	-5.709
CH ₃ O (² A')	-6.360	-6.558	-6.248	-6.441	-6.601	-6.607	-6.589
CH ₃ CH ₂ O (² A'')	-8.548	-9.090	-8.408	-8.650	-8.954	-8.944	-8.898
CH ₃ S (² A')	-4.760	-4.973	-4.868	-4.732	-4.940	-4.861	-4.841
C ₂ H ₅ (² A')	-5.296	-6.265	-5.562	-5.693	-5.937	-5.895	-5.849
(CH ₃) ₂ CH (² A')	-7.348	-8.410	-7.361	-7.537	-7.921	-7.865	-7.792
(CH ₃) ₂ CH (² A)	-7.348	-8.410	-7.361	-7.537	-7.921	-7.865	-7.792
NO ₂	-12.349	-12.138	-12.246	-12.926	-13.008	-12.983	-13.035
Methyl allene (C ₄ H ₆)	-5.145	-6.179	-4.907	-5.012	-5.549	-5.491	-5.381
Isoprene (C ₅ H ₈)	-3.487	-4.960	-3.364	-3.499	-4.181	-4.103	-3.964
Cyclopentane (C ₅ H ₁₀)	-2.800	-4.827	-3.008	-3.267	-3.982	-3.902	-3.762
n-Pentane (C ₅ H ₁₂)	-2.243	-3.457	-1.734	-1.937	-2.652	-2.572	-2.432
Neo pentane (C ₅ H ₁₂)	-0.506	-2.573	-0.777	-1.023	-1.740	-1.660	-1.519
1,4 Cyclohexadiene (C ₆ H ₈)	-4.599	-6.373	-4.368	-4.551	-5.379	-5.288	-5.121
Cyclohexane (C ₆ H ₁₂)	-2.548	-5.093	-2.931	-3.219	-4.083	-3.987	-3.820
n-Hexane (C ₆ H ₁₄)	-2.441	-4.723	-2.428	-2.711	-3.570	-3.471	-3.303
3-Methyl pentane (C ₆ H ₁₄)	-0.979	-3.512	-1.308	-1.576	-2.435	-2.338	-2.170
Toluene (C ₆ H ₅ CH ₃)	-11.179	-13.201	-10.958	-11.130	-12.092	-11.980	-11.787
n-Heptane (C ₇ H ₁₆)	-3.879	-5.386	-2.998	-3.278	-4.279	-4.167	-3.971
Cyclooctatetraene (C ₈ H ₈)	-7.673	-9.833	-7.355	-7.550	-8.633	-8.510	-8.289
n-Octane (C ₈ H ₁₈)	-3.736	-6.518	-3.477	-3.855	-4.994	-4.861	-4.637
Naphthalene (C ₁₀ H ₈)	-20.460	-23.262	-20.181	-20.396	-21.756	-21.597	-21.322
Azulene (C ₁₀ H ₈)	-21.123	-23.781	-20.692	-20.918	-22.287	-22.126	-21.849
Acetic acid methyl ester (CH ₃ COOCH ₃)	-8.120	-8.312	-7.529	-8.119	-8.605	-8.597	-8.538
t-Butanol ((CH ₃) ₃ COH)	-2.230	-3.620	-2.284	-2.689	-3.295	-3.251	-3.151
Aniline (C ₆ H ₅ NH ₂)	-11.566	-13.375	-11.314	-11.734	-12.580	-12.414	-12.279
Phenol (C ₆ H ₅ OH)	-13.077	-14.422	-12.634	-13.007	-13.855	-13.780	-13.627
Divinyl ether (C ₄ H ₆ O)	-6.486	-7.212	-6.049	-6.395	-6.959	-6.918	-6.821
Tetrahydrofuran (C ₄ H ₈ O)	-3.869	-5.347	-4.046	-4.465	-5.059	-5.014	-4.916
Cyclopentanone (C ₅ H ₈ O)	-8.037	-9.788	-8.105	-8.548	-9.291	-9.237	-9.107

Continued on next page

TABLE X: Deviation of r²SCAN atomization energies (kcal/mol) obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Benzoquinone(C ₆ H ₄ O ₂)	-13.851	-14.713	-13.036	-13.621	-14.509	-14.467	-14.321
Pyrimidine (C ₄ H ₄ N ₂)	-10.036	-11.422	-9.872	-10.498	-11.106	-10.898	-10.844
Dimethyl sulphone (C ₂ H ₆ O ₂ S)	-8.084	-7.408	-7.276	-7.678	-8.116	-8.044	-8.014
Chlorobenzene (C ₆ H ₅ Cl)	-14.208	-15.922	-13.991	-14.178	-15.067	-14.930	-14.765
Butanedinitrile (NC-CH ₂ -CH ₂ -CN)	2.574	1.052	2.705	2.073	1.452	1.660	1.711
Pyrazine (C ₄ H ₄ N ₂)	-6.029	-7.400	-5.891	-6.510	-7.114	-6.905	-6.852
Acetyl acetylene (CH ₃ -C(=O)-CCH)	-2.696	-3.439	-2.290	-2.624	-3.195	-3.163	-3.063
Crotonaldehyde (CH ₃ -CH=CH-CHO)	-7.788	-8.545	-7.392	-7.732	-8.317	-8.279	-8.179
Acetic anhydride (CH ₃ -C(=O)-O-C(=O)-CH ₃)	-14.696	-15.158	-14.111	-14.983	-15.645	-15.647	-15.570
2,5-Dihydrothiophene (C ₄ H ₆ S)	-6.090	-7.173	-6.244	-6.102	-6.762	-6.610	-6.504
Isobutane nitrile ((CH ₃) ₂ CH-CN)	1.962	0.615	2.155	1.742	1.148	1.282	1.364
Methyl ethyl ketone (CH ₃ -CO-CH ₂ -CH ₃)	-5.074	-6.641	-5.271	-5.668	-6.266	-6.229	-6.127
Isobutanal ((CH ₃) ₂ CH-CHO)	-3.245	-4.273	-2.964	-3.362	-3.961	-3.922	-3.821
1,4-Dioxane (C ₄ H ₈ O ₂)	-7.196	-8.035	-6.874	-7.504	-8.130	-8.101	-8.017
Tetrahydrothiophene (C ₄ H ₈ S)	-3.945	-5.433	-4.392	-4.293	-4.973	-4.818	-4.711
t-Butyl chloride ((CH ₃) ₃ C-Cl)	-4.183	-5.962	-4.518	-4.737	-5.385	-5.278	-5.165
n-Butyl chloride (CH ₃ -CH ₂ -CH ₂ -CH ₂ -Cl)	-3.952	-5.306	-3.829	-4.058	-4.704	-4.595	-4.483
Tetrahydropyrrole (C ₄ H ₈ NH)	-1.280	-3.023	-1.391	-1.885	-2.481	-2.345	-2.263
Nitro-s-butane (CH ₃ -CH ₂ -CH(CH ₃)-NO ₂)	-11.849	-12.992	-11.527	-12.443	-13.119	-13.026	-12.968
Diethyl ether (CH ₃ -CH ₂ -O-CH ₂ -CH ₃)	-3.788	-4.797	-3.529	-3.928	-4.530	-4.483	-4.384
Dimethyl acetal (CH ₃ -CH(OCH ₃) ₂)	-5.875	-6.614	-5.453	-6.101	-6.731	-6.702	-6.617
t-Butanethiol ((CH ₃) ₃ C-SH)	-2.678	-4.050	-3.039	-2.919	-3.602	-3.445	-3.338
Diethyl disulfide (CH ₃ -CH ₂ -S-S-CH ₂ -CH ₃)	-6.148	-6.702	-6.128	-5.734	-6.513	-6.272	-6.170
t-Butylamine ((CH ₃) ₃ C-NH ₂)	0.374	-0.976	0.669	0.214	-0.381	-0.247	-0.164
Tetramethylsilane (Si(CH ₃) ₄)	0.885	-0.722	0.895	0.695	0.153	0.184	0.300
2-Methyl thiophene (C ₅ H ₆ S)	-10.258	-11.769	-10.509	-10.371	-11.176	-11.005	-10.872
N-methyl pyrrole (cyc-C ₄ H ₄ N-CH ₃)	-7.316	-9.162	-7.395	-7.814	-8.528	-8.375	-8.268
Tetrahydropyran (C ₅ H ₁₀ O)	-4.763	-6.521	-4.791	-5.256	-6.004	-5.942	-5.815
Diethyl ketone (CH ₃ -CH ₂ -CO-CH ₂ -CH ₃)	-7.202	-8.228	-6.622	-7.036	-7.782	-7.727	-7.598
Isopropyl acetate (CH ₃ -C(=O)-O-CH(CH ₃) ₂)	-9.366	-10.451	-8.948	-9.629	-10.400	-10.361	-10.245
Tetrahydrothiopyran (C ₅ H ₁₀ S)	-4.933	-6.579	-5.134	-5.090	-5.913	-5.741	-5.607
Piperidine (cyc-C ₅ H ₁₀ NH)	-0.947	-3.389	-1.416	-1.936	-2.677	-2.526	-2.416
t-Butyl methyl ether ((CH ₃) ₃ C-O-CH ₃)	-3.699	-4.961	-3.301	-3.764	-4.501	-4.439	-4.312
1,3-Difluorobenzene (C ₆ H ₄ F ₂)	-17.361	-19.136	-17.371	-17.362	-18.164	-18.088	-17.918
1,4-Difluorobenzene (C ₆ H ₄ F ₂)	-18.967	-20.647	-18.954	-18.941	-19.744	-19.668	-19.498
Fluorobenzene (C ₆ H ₅ F)	-14.056	-15.731	-13.924	-13.986	-14.795	-14.709	-14.542
Di-isopropyl ether ((CH ₃) ₂ CH-O-CH(CH ₃) ₂)	-4.677	-6.266	-4.236	-4.715	-5.605	-5.527	-5.372
PF ₅	0.578	0.810	0.377	0.933	0.942	0.927	0.941
SF ₆	-13.028	-12.488	-13.026	-12.359	-12.435	-12.406	-12.398
P ₄	-7.667	-4.968	-6.398	-5.867	-5.861	-5.724	-5.743
SO ₃	-8.855	-7.223	-7.836	-8.447	-8.637	-8.624	-8.662
SCL ₂	-5.017	-5.106	-5.263	-5.151	-5.383	-5.225	-5.227
POCl ₃	-9.920	-8.498	-8.616	-8.756	-9.082	-8.947	-8.960
PCL ₅	-17.057	-16.190	-16.097	-16.110	-16.532	-16.299	-16.301
Cl ₂ O ₂ S	-12.396	-11.570	-11.967	-12.389	-12.698	-12.578	-12.605
PCL ₃	-7.992	-7.003	-7.068	-6.984	-7.260	-7.107	-7.108
Cl ₂ S ₂	-14.280	-14.108	-14.687	-14.245	-14.608	-14.355	-14.362
SiCl ₂ singlet	-2.573	-2.951	-2.719	-2.827	-2.969	-2.921	-2.913
CF ₃ Cl	-11.537	-12.298	-12.034	-11.925	-12.145	-12.110	-12.069
Hexafluoro ethane (C ₂ F ₆)	-17.625	-18.855	-18.478	-18.147	-18.428	-18.443	-18.362
CF ₃	-13.405	-14.006	-13.918	-13.760	-13.851	-13.862	-13.828
C ₆ H ₅ (phenyl radical)	-16.804	-18.612	-16.759	-16.912	-17.687	-17.597	-17.436
Bicyclo[1.1.0]butane (C ₄ H ₆)	-4.610	-6.161	-4.724	-4.879	-5.454	-5.395	-5.282
(CH ₃) ₃ C (t-butyl radical)	-7.491	-9.261	-7.806	-8.006	-8.539	-8.468	-8.365
Trans-butane(C ₄ H ₁₀)	-1.217	-2.799	-1.289	-1.475	-2.049	-1.983	-1.871
1,3 Cyclohexadiene (C ₆ H ₈)	-5.141	-7.187	-5.185	-5.370	-6.196	-6.103	-5.937
ME	-4.700	-5.329	-4.612	-4.784	-5.158	-5.097	-5.042

Continued on next page

TABLE XI: Deviation of r^4 SCAN atomization energies (kcal/mol) for the G3 test set obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
MAE	5.578	6.119	5.500	5.637	5.964	5.916	5.866

TABLE XI: Deviation of r^4 SCAN atomization energies (kcal/mol) for the G3 test set obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
LiH	2.403	2.537	2.562	2.503	2.547	2.553	2.583
BeH	-10.549	-10.555	-10.337	-10.462	-10.363	-10.166	-10.358
CH	2.203	2.564	2.348	2.400	2.449	2.293	2.375
CH ₂ (³ B ₁)	-6.302	-6.343	-6.300	-6.009	-6.037	-6.256	-6.188
CH ₂ (¹ A ₁)	5.376	5.926	5.755	5.800	5.758	5.603	5.724
CH ₃	-4.008	-4.250	-4.020	-4.022	-3.956	-4.130	-4.078
Methane (CH ₄)	0.527	0.574	0.493	0.530	0.544	0.419	0.530
NH	0.413	-1.697	-0.425	-0.814	-0.948	-0.957	-0.987
NH ₂	-0.208	-3.232	-1.197	-2.168	-2.274	-2.187	-2.177
Ammonia (NH ₃)	5.643	1.965	4.299	3.459	3.189	3.213	3.275
OH	-3.108	-1.348	-2.753	-3.108	-2.774	-2.486	-2.464
Water (H ₂ O)	0.774	3.718	1.761	1.687	2.057	2.299	2.156
Hydrogen fluoride (HF)	1.838	3.626	1.241	2.129	2.302	2.382	2.339
SiH ₂ (¹ A ₁)	2.480	2.309	2.310	2.435	2.516	2.423	2.381
SiH ₂ (³ B ₁)	-7.987	-7.620	-7.765	-7.745	-7.632	-7.562	-7.561
SiH ₃	-5.159	-4.897	-5.018	-5.025	-4.916	-4.911	-4.907
Silane (SiH ₄)	-1.005	-0.512	-0.538	-0.699	-0.600	-0.663	-0.628
PH ₂	-3.804	-3.587	-3.660	-3.353	-3.674	-3.722	-3.883
PH ₃	-0.546	0.460	0.079	0.184	-0.104	-0.178	-0.263
Hydrogen sulfide (H ₂ S)	-0.674	0.624	0.193	-0.408	-0.587	-0.871	-0.704
Hydrogen chloride (HCl)	-1.011	-0.238	0.164	-0.034	-0.621	-0.079	-0.306
Li ₂	5.635	5.940	6.013	5.874	5.973	5.990	6.043
LiF	3.181	5.286	-0.150	0.776	3.519	2.769	2.859
Acetylene (C ₂ H ₂)	1.580	3.505	2.963	3.177	3.084	2.694	2.934
Ethylene (H ₂ C=CH ₂)	-0.858	0.048	-0.273	-0.138	-0.182	-0.521	-0.292
Ethane (H ₃ C-CH ₃)	-1.651	-1.513	-1.633	-1.560	-1.526	-1.793	-1.591
CN	4.927	1.250	4.166	3.231	3.141	2.876	2.970
Hydrogen cyanide (HCN)	5.007	3.315	4.851	4.172	4.125	3.965	4.048
CO	1.409	4.084	2.391	1.531	2.058	2.295	2.369
HCO	-6.905	-3.818	-6.041	-6.308	-5.923	-5.694	-5.647
Formaldehyde (H ₂ C=O)	-3.305	-0.472	-2.188	-2.508	-2.195	-2.039	-2.004
Methanol (CH ₃ -OH)	-3.000	-0.147	-2.093	-2.145	-1.892	-1.817	-1.783
N ₂	11.372	6.766	10.081	8.383	8.332	8.358	8.367
Hydrazine (H ₂ N-NH ₂)	6.769	-0.071	4.991	3.164	2.844	2.760	2.732
NO	-0.733	0.020	-0.499	-1.511	-1.182	-0.750	-0.894
O ₂	-11.568	-5.991	-11.715	-11.600	-10.861	-9.748	-10.019
Hydrogen peroxide (HO-OH)	-2.141	2.135	-1.535	-2.266	-1.713	-0.889	-0.979
F ₂	2.756	-1.003	-1.266	-0.877	-0.848	0.765	0.182
Carbon dioxide (CO ₂)	-10.238	-5.578	-8.099	-10.142	-8.564	-7.625	-8.077
Na ₂	3.050	2.982	2.833	2.954	2.927	2.911	3.042
Si ₂	-0.804	-0.042	0.035	0.258	0.584	0.257	0.226
P ₂	2.377	3.814	3.465	3.956	3.445	3.228	3.021
S ₂	-9.776	-7.810	-8.122	-8.995	-9.070	-9.929	-9.806
Cl ₂	-2.564	-2.003	-1.349	-0.880	-1.802	-1.410	-1.190
NaCl	1.262	1.715	1.070	-0.467	-1.626	0.638	-0.366
Silicon monoxide (SiO)	3.348	5.887	4.101	2.940	3.628	4.074	4.050
CS	2.069	4.296	3.834	2.768	2.315	1.658	2.010
SO	-10.227	-6.309	-7.913	-8.224	-8.240	-8.562	-8.340
ClO	-9.146	-6.094	-7.708	-6.577	-7.600	-6.874	-6.836

Continued on next page

TABLE XI: Deviation of r^4 SCAN atomization energies (kcal/mol) for the G3 test set obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Chlorine monofluoride (FCl)	-2.133	-0.867	-2.022	-0.533	-2.072	-1.265	-1.217
Si ₂ H ₆	-4.870	-3.789	-3.887	-4.255	-4.026	-4.176	-4.129
Methyl chloride (CH ₃ Cl)	-3.931	-3.043	-2.709	-2.659	-3.206	-2.909	-3.051
Methanethiol (H ₃ CSH)	-2.728	-1.267	-1.776	-2.248	-2.436	-2.888	-2.648
Hypochlorous acid (HOCl)	-3.283	-0.558	-2.053	-2.015	-2.393	-1.928	-1.717
Sulfur dioxide (SO ₂)	-6.736	-0.538	-3.226	-5.877	-4.313	-4.066	-4.245
BF ₃	-4.734	2.825	-9.572	-6.631	-2.933	-4.081	-3.669
BCl ₃	-13.824	-10.851	-10.409	-13.325	-15.941	-12.319	-13.679
AlF ₃	2.488	10.060	-3.966	-1.206	4.914	3.152	3.544
AlCl ₃	-9.983	-6.581	-7.013	-11.209	-13.865	-8.404	-10.906
Carbon tetrafluoride (CF ₄)	-12.313	-5.587	-14.437	-11.383	-10.974	-10.562	-10.753
Carbon tetrachloride (CCl ₄)	-15.091	-12.969	-11.341	-10.856	-12.583	-11.367	-12.126
Carbon oxide sulfide (COS)	-11.338	-7.031	-9.040	-11.162	-10.848	-10.790	-10.485
Carbon bisulfide (CS ₂)	-10.955	-7.146	-8.246	-10.899	-11.681	-12.598	-11.696
Carbonic difluoride (COF ₂)	-7.769	-2.216	-8.592	-7.336	-6.249	-5.990	-6.190
Silicon tetrafluoride (SiF ₄)	6.340	16.511	-0.151	3.277	9.878	7.246	8.119
Silicon tetrachloride (SiCl ₄)	-8.552	-5.121	-4.674	-9.530	-12.135	-6.942	-9.426
Dinitrogen monoxide (N ₂ O)	-0.428	-7.868	-3.061	-6.646	-5.897	-5.237	-5.572
Nitrogen chloride oxide (ClNO)	-8.300	-7.707	-6.996	-9.317	-9.718	-8.109	-8.900
Nitrogen trifluoride (NF ₃)	-11.331	-12.118	-14.163	-13.265	-14.346	-12.343	-13.096
PF ₃	1.360	8.568	-2.548	0.329	4.147	2.261	2.757
O ₃	-10.916	-0.457	-8.987	-8.634	-7.846	-6.495	-6.822
F ₂ O	-7.670	-5.262	-9.843	-9.103	-9.463	-7.257	-7.914
Chlorine trifluoride (ClF ₃)	-18.920	-13.889	-20.090	-16.468	-18.038	-16.036	-16.629
Tetrafluoro Ethene (F ₂ C=CF ₂)	-21.787	-14.927	-23.621	-20.219	-19.925	-19.733	-19.834
Tetrachloro Ethene (C ₂ Cl ₄)	-21.924	-18.464	-17.426	-16.679	-18.622	-17.362	-18.009
Acetonitrile, trifluoro- (CF ₃ CN)	-9.086	-5.386	-10.601	-8.882	-8.681	-8.577	-8.605
Propyne (C ₃ H ₄)	-2.202	-0.355	-0.888	-0.725	-0.771	-1.296	-0.937
Allene (C ₃ H ₄)	-6.114	-4.331	-4.971	-4.708	-4.727	-5.246	-4.913
Cyclopropene (C ₃ H ₄)	-1.938	-1.222	-1.546	-1.201	-1.301	-1.813	-1.484
Propylene (C ₃ H ₆)	-3.650	-2.548	-2.945	-2.814	-2.822	-3.292	-2.959
Cyclopropane (C ₃ H ₆)	-4.693	-4.626	-4.757	-4.394	-4.493	-4.981	-4.678
Propane (C ₃ H ₈)	-3.694	-3.531	-3.562	-3.510	-3.447	-3.858	-3.561
Trans-1,3-butadiene (C ₄ H ₆)	-6.315	-4.256	-4.829	-4.581	-4.670	-5.339	-4.884
Dimethylacetylene (C ₄ H ₆)	-4.749	-2.975	-3.482	-3.376	-3.373	-4.028	-3.555
Methylenecyclopropane (C ₄ H ₆)	-10.010	-9.019	-9.507	-9.146	-9.170	-9.811	-9.392
Bicyclo[1.1.0]butane (C ₄ H ₆)	-6.268	-6.914	-6.812	-6.160	-6.353	-7.014	-6.620
Cyclobutene (C ₄ H ₆)	-5.367	-3.901	-4.214	-3.991	-4.005	-4.632	-4.243
Cyclobutane (C ₄ H ₈)	-7.550	-6.172	-6.364	-6.200	-6.152	-6.722	-6.366
Isobutene (C ₄ H ₈)	-6.061	-4.553	-5.003	-4.858	-4.823	-5.424	-4.991
Trans-butane(C ₄ H ₁₀)	-5.521	-5.212	-5.300	-5.205	-5.124	-5.652	-5.265
Isobutane (C ₄ H ₁₀)	-4.658	-4.771	-4.834	-4.729	-4.636	-5.165	-4.776
Spiropentane (C ₅ H ₈)	-11.215	-11.056	-11.129	-10.559	-10.672	-11.470	-11.014
Benzene (C ₆ H ₆)	-16.232	-12.570	-13.282	-13.130	-13.254	-14.295	-13.575
Difluoromethane (CH ₂ F ₂)	-6.003	-2.108	-7.176	-5.308	-4.914	-4.824	-4.868
Trifluoromethane(CHF ₃)	-9.279	-3.547	-10.738	-8.120	-7.679	-7.613	-7.659
CH ₂ Cl ₂	-8.095	-6.214	-5.766	-5.461	-6.508	-5.744	-6.180
CHCl ₃	-11.537	-9.405	-8.476	-8.202	-9.596	-8.546	-9.165
Methylamine (H ₃ C-NH ₂)	2.978	-0.486	1.790	1.053	0.720	0.602	0.779
Acetonitrile (CH ₃ -CN)	1.422	-0.542	1.118	0.429	0.365	0.088	0.275
Nitromethane (CH ₃ -NO ₂)	-12.283	-10.601	-11.833	-14.487	-13.811	-12.995	-12.825
Methyl nitrite (CH ₃ -O-N=O)	-11.627	-8.063	-10.185	-11.635	-11.071	-10.564	-10.591
Methyl silane (CH ₃ SiH ₃)	-1.326	-0.811	-0.778	-0.859	-0.748	-0.940	-0.815
Formic acid (HCOOH)	-9.335	-3.588	-7.358	-7.869	-6.899	-6.488	-6.742
Methyl formate (HCOOCH ₃)	-13.812	-7.654	-11.404	-11.860	-11.000	-10.735	-10.856
Acetamide (CH ₃ CONH ₂)	-6.225	-7.823	-5.673	-7.542	-7.228	-7.266	-7.226
Aziridine (C ₂ H ₄ NH)	-1.413	-3.971	-2.265	-2.812	-3.054	-3.314	-3.095

Continued on next page

TABLE XI: Deviation of r^4 SCAN atomization energies (kcal/mol) for the G3 test set obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Cyanogen (NCCN)	4.302	0.890	4.069	2.612	2.535	2.228	2.400
Dimethylamine ((CH ₃) ₂ NH)	0.058	-3.171	-0.814	-1.531	-1.915	-2.176	-1.880
Trans ethylamine (CH ₃ CH ₂ NH ₂)	-0.265	-3.598	-1.263	-1.968	-2.305	-2.548	-2.266
Ketene (CH ₂ CO)	-9.971	-7.020	-8.820	-8.929	-8.383	-8.325	-8.388
Oxirane (C ₂ H ₄ O)	-7.301	-4.366	-6.163	-6.714	-6.427	-6.397	-6.206
Acetaldehyde (CH ₃ CHO)	-6.560	-3.546	-5.391	-5.667	-5.305	-5.278	-5.150
Glyoxal (HCOCHO)	-10.073	-4.571	-7.850	-8.639	-8.000	-7.704	-7.616
Ethanol (CH ₃ CH ₂ OH)	-4.934	-1.971	-3.948	-3.963	-3.719	-3.745	-3.611
Dimethylether (CH ₃ OCH ₃)	-6.776	-3.475	-5.517	-5.430	-5.273	-5.360	-5.179
Thiirane (C ₂ H ₄ S)	-6.647	-4.784	-5.597	-6.104	-6.274	-6.972	-6.703
Dimethyl sulfoxide ((CH ₃) ₂ SO)	-9.613	-6.655	-7.086	-7.150	-8.857	-9.131	-8.797
Ethaneithiol (C ₂ H ₅ SH)	-4.422	-2.806	-3.388	-3.871	-4.042	-4.634	-4.292
Dimethyl sulfide (CH ₃ SCH ₃)	-5.264	-3.506	-4.092	-4.447	-4.644	-5.279	-4.945
Vinyl fluoride (CH ₂ =CHF)	-6.596	-4.107	-6.902	-5.717	-5.555	-5.806	-5.646
Ethyl chloride (C ₂ H ₅ Cl)	-6.343	-5.161	-4.864	-4.868	-5.489	-5.242	-5.305
Vinyl chloride (CH ₂ =CHCl)	-10.192	-8.173	-8.156	-8.095	-8.709	-8.526	-8.577
Acrylonitrile (CH ₂ =CHCN)	0.895	-0.058	1.308	0.765	0.680	0.202	0.483
Acetone (CH ₃ COCH ₃)	-9.385	-5.975	-7.928	-8.200	-7.801	-7.903	-7.666
Acetic acid (CH ₃ COOH)	-11.907	-5.748	-9.536	-10.055	-9.123	-8.827	-8.964
Acetyl fluoride (CH ₃ COF)	-10.685	-5.960	-10.669	-9.989	-8.969	-9.034	-9.001
CH ₃ COCl (acetyl chloride)	-13.098	-9.341	-10.753	-11.623	-11.819	-10.855	-11.286
CH ₃ CH ₂ CH ₂ Cl (propyl chloride)	-8.385	-7.279	-6.926	-6.898	-7.456	-7.379	-7.352
Isopropanol (CH ₃) ₂ CHOH)	-6.693	-3.897	-5.930	-5.874	-5.564	-5.730	-5.508
Methyl ethyl ether (C ₂ H ₅ OCH ₃)	-9.108	-5.826	-7.816	-7.712	-7.525	-7.736	-7.460
Trimethylamine ((CH ₃) ₃ N)	-2.742	-6.089	-3.733	-4.387	-4.837	-5.245	-4.823
Furan (C ₄ H ₄ O)	-15.396	-9.859	-13.034	-12.652	-12.394	-12.793	-12.445
C ₄ H ₄ S (thiophene)	-12.961	-9.272	-10.783	-11.164	-11.260	-12.362	-11.675
Pyrrole (C ₄ H ₅ N)	-8.965	-11.410	-8.516	-9.319	-9.597	-10.455	-10.044
Pyridine (C ₅ H ₅ N)	-13.768	-14.409	-12.450	-13.016	-13.466	-14.296	-13.697
H ₂	2.073	2.089	2.088	2.088	2.089	2.088	2.088
HS	-2.145	-1.489	-1.600	-1.846	-2.116	-2.288	-2.256
CCH	-0.786	-0.624	-0.743	-0.382	-0.340	-0.627	-0.451
C ₂ H ₃ (² A')	-6.569	-6.078	-6.255	-6.089	-6.055	-6.378	-6.173
CH ₃ CO (² A')	-9.965	-7.098	-9.156	-9.552	-9.176	-9.065	-8.891
H ₂ COH (² A)	-7.775	-4.186	-7.003	-6.812	-6.473	-6.287	-6.293
CH ₃ O (² A')	-8.503	-6.745	-8.091	-8.389	-8.128	-7.981	-7.915
CH ₃ CH ₂ O (² A'')	-11.616	-10.026	-11.258	-11.608	-11.320	-11.257	-11.074
CH ₃ S (² A')	-5.812	-4.943	-5.119	-5.315	-5.619	-5.980	-5.828
C ₂ H ₅ (² A')	-6.774	-6.917	-6.696	-6.665	-6.623	-6.940	-6.778
(CH ₃) ₂ CH (² A')	-9.818	-9.656	-9.489	-9.452	-9.405	-9.845	-9.570
(CH ₃) ₂ CH (² A')	-9.818	-9.656	-9.489	-9.452	-9.405	-9.845	-9.570
NO ₂	-11.911	-11.371	-12.099	-15.134	-14.080	-13.025	-13.142
Methyl allene (C ₄ H ₆)	-8.375	-6.270	-6.804	-6.555	-6.550	-7.214	-6.775
Isoprene (C ₅ H ₈)	-8.068	-5.904	-6.499	-6.264	-6.322	-7.119	-6.559
Cyclopentane (C ₅ H ₁₀)	-8.471	-8.416	-8.454	-8.309	-8.219	-8.930	-8.480
n-Pentane (C ₅ H ₁₂)	-7.305	-6.840	-6.827	-6.752	-6.639	-7.328	-6.844
Neo pentane (C ₅ H ₁₂)	-5.772	-5.687	-5.733	-5.623	-5.505	-6.162	-5.675
1,4 Cyclohexadiene (C ₆ H ₈)	-11.118	-8.718	-9.352	-9.113	-9.102	-10.042	-9.400
Cyclohexane (C ₆ H ₁₂)	-9.919	-10.066	-9.967	-9.827	-9.664	-10.480	-9.930
n-Hexane (C ₆ H ₁₄)	-9.279	-8.733	-8.774	-8.655	-8.520	-9.314	-8.745
3-Methyl pentane (C ₆ H ₁₄)	-7.697	-7.532	-7.448	-7.378	-7.252	-8.059	-7.490
Toluene (C ₆ H ₅ CH ₃)	-18.297	-14.495	-15.186	-15.059	-15.160	-16.323	-15.500
n-Heptane (C ₇ H ₁₆)	-11.228	-10.448	-10.391	-10.293	-10.129	-11.096	-10.425
Cyclooctatetraene (C ₈ H ₈)	-16.058	-11.721	-12.774	-12.440	-12.578	-13.884	-12.960
n-Octane (C ₈ H ₁₈)	-13.067	-12.191	-12.121	-12.003	-11.818	-12.877	-12.125
Naphthalene (C ₁₀ H ₈)	-31.084	-24.971	-26.025	-25.805	-26.069	-27.714	-26.546

Continued on next page

TABLE XI: Deviation of r^4 SCAN atomization energies (kcal/mol) for the G3 test set obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Azulene (C ₁₀ H ₈)	-30.897	-25.352	-26.715	-26.361	-26.510	-28.229	-27.035
Acetic acid methyl ester (CH ₃ COOCH ₃)	-15.028	-8.756	-12.585	-12.953	-12.127	-11.990	-11.987
t-Butanol ((CH ₃) ₃ COH)	-8.044	-5.091	-7.173	-7.122	-6.776	-7.056	-6.723
Aniline (C ₆ H ₅ NH ₂)	-16.557	-16.687	-14.564	-15.392	-15.772	-16.855	-16.062
Phenol (C ₆ H ₅ OH)	-21.248	-14.448	-17.512	-17.241	-16.934	-17.777	-17.218
Divinyl ether (C ₄ H ₆ O)	-12.913	-7.347	-10.536	-9.963	-9.609	-10.104	-9.808
Tetrahydrofuran (C ₄ H ₈ O)	-11.146	-8.223	-10.034	-10.084	-9.858	-10.183	-9.816
Cyclopentanone (C ₅ H ₈ O)	-15.110	-12.354	-14.100	-14.325	-13.857	-14.243	-13.880
Benzoquinone(C ₆ H ₄ O ₂)	-23.303	-15.175	-19.349	-19.652	-19.064	-19.398	-18.855
Pyrimidine (C ₄ H ₄ N ₂)	-12.511	-17.282	-12.771	-14.039	-14.753	-15.401	-14.907
Dimethyl sulphone (C ₂ H ₆ O ₂ S)	-13.003	-8.113	-9.534	-14.271	-11.996	-11.693	-11.502
Chlorobenzene (C ₆ H ₅ Cl)	-21.487	-16.982	-17.474	-17.328	-18.002	-18.525	-18.080
Butanedinitrile (NC-CH ₂ -CH ₂ -CN)	1.818	-1.965	1.393	0.099	-0.003	-0.602	-0.265
Pyrazine (C ₄ H ₄ N ₂)	-8.471	-13.346	-8.370	-9.924	-10.674	-11.332	-10.862
Acetyl acetylene (CH ₃ -C(=O)-CCH)	-7.285	-2.607	-5.054	-5.095	-4.763	-5.106	-4.737
Crotonaldehyde (CH ₃ -CH=CH-CHO)	-13.228	-9.050	-11.173	-11.355	-10.998	-11.329	-10.978
Acetic anhydride (CH ₃ -C(=O)-O-C(=O)-CH ₃)	-24.157	-14.799	-20.346	-21.290	-19.889	-19.582	-19.644
2,5-Dihydrothiophene (C ₄ H ₆ S)	-10.983	-8.338	-9.229	-9.488	-9.689	-10.628	-10.078
Isobutane nitrile ((CH ₃) ₂ CH-CN)	-0.324	-2.061	-0.457	-1.087	-1.119	-1.687	-1.311
Methyl ethyl ketone (CH ₃ -CO-CH ₂ -CH ₃)	-10.626	-8.053	-9.784	-9.999	-9.559	-9.835	-9.504
Isobutanal ((CH ₃) ₂ CH-CHO)	-9.068	-5.965	-7.674	-7.917	-7.500	-7.776	-7.470
1,4-Dioxane (C ₄ H ₈ O ₂)	-17.482	-10.945	-14.893	-14.831	-14.584	-14.621	-14.288
Tetrahydrothiophene (C ₄ H ₈ S)	-9.134	-7.754	-8.439	-8.764	-8.875	-9.759	-9.245
t-Butyl chloride ((CH ₃) ₃ C-Cl)	-9.305	-8.494	-8.444	-8.549	-9.146	-9.016	-8.920
n-Butyl chloride (CH ₃ -CH ₂ -CH ₂ -CH ₂ -Cl)	-9.380	-8.401	-7.991	-8.039	-8.614	-8.617	-8.510
Tetrahydropyrrole (C ₄ H ₈ NH)	-5.379	-8.419	-6.175	-6.735	-7.125	-7.675	-7.204
Nitro-s-butane (CH ₃ -CH ₂ -CH(CH ₃)-NO ₂)	-16.628	-15.141	-16.383	-18.967	-18.181	-17.789	-17.334
Diethyl ether (CH ₃ -CH ₂ -O-CH ₂ -CH ₃)	-10.302	-7.048	-9.008	-9.043	-8.876	-9.187	-8.816
Dimethyl acetal (CH ₃ -CH(OCH ₃) ₂)	-15.031	-8.548	-12.543	-12.510	-12.026	-12.178	-11.924
t-Butanethiol ((CH ₃) ₃ C-SH)	-7.341	-5.945	-6.489	-6.966	-7.062	-7.928	-7.365
Diethyl disulfide (CH ₃ -CH ₂ -S-S-CH ₂ -CH ₃)	-11.451	-8.213	-9.286	-10.259	-10.505	-11.744	-11.075
t-Butylamine ((CH ₃) ₃ C-NH ₂)	-2.645	-5.403	-3.133	-3.809	-4.098	-4.592	-4.101
Tetramethylsilane (Si(CH ₃) ₄)	-1.525	-1.132	-0.957	-0.818	-0.651	-1.287	-0.882
2-Methyl thiophene (C ₅ H ₆ S)	-15.147	-11.358	-12.903	-13.340	-13.403	-14.630	-13.829
N-methyl pyrrole (cyc-C ₄ H ₄ N-CH ₃)	-10.990	-13.341	-10.546	-11.276	-11.592	-12.607	-12.050
Tetrahydropyran (C ₅ H ₁₀ O)	-13.338	-10.200	-12.286	-12.223	-11.913	-12.385	-11.955
Diethyl ketone (CH ₃ -CH ₂ -CO-CH ₂ -CH ₃)	-13.680	-10.271	-12.257	-12.518	-12.073	-12.459	-12.043
Isopropyl acetate (CH ₃ -C(=O)-O-CH(CH ₃) ₂)	-18.388	-12.109	-15.882	-16.348	-15.446	-15.549	-15.335
Tetrahydrothiopyran (C ₅ H ₁₀ S)	-11.784	-10.067	-10.547	-10.896	-11.014	-12.034	-11.419
Piperidine (cyc-C ₅ H ₁₀ NH)	-6.441	-10.050	-7.619	-8.268	-8.561	-9.216	-8.650
t-Butyl methyl ether ((CH ₃) ₃ C-O-CH ₃)	-11.524	-7.842	-9.831	-9.792	-9.508	-9.941	-9.451
1,3-Difluorobenzene (C ₆ H ₄ F ₂)	-24.873	-18.248	-24.181	-21.976	-21.698	-22.476	-21.941
1,4-Difluorobenzene (C ₆ H ₄ F ₂)	-26.668	-20.065	-25.879	-23.559	-23.332	-24.076	-23.546
Fluorobenzene (C ₆ H ₅ F)	-20.756	-15.546	-18.946	-17.744	-17.632	-18.566	-17.932
Di-isopropyl ether ((CH ₃) ₂ CH-O-CH(CH ₃) ₂)	-13.279	-9.579	-11.726	-11.674	-11.404	-11.971	-11.401
PF ₅	-2.048	10.604	-5.561	-1.551	2.992	0.895	1.528
SF ₆	-20.116	-3.159	-17.193	-14.617	-12.643	-12.940	-13.038
P ₄	-5.908	-2.635	-3.277	-3.147	-4.240	-5.083	-5.626
SO ₃	-11.975	-5.286	-7.458	-13.123	-10.330	-9.549	-9.682
SCl ₂	-7.998	-5.916	-5.130	-5.764	-6.853	-6.096	-6.619
POCl ₃	-13.193	-7.551	-7.428	-11.337	-12.167	-9.082	-10.632
PCl ₅	-22.522	-17.015	-15.014	-17.722	-20.656	-17.044	-19.084
Cl ₂ O ₂ S	-17.248	-12.148	-12.427	-17.813	-16.306	-14.471	-15.362
PCl ₃	-9.704	-6.705	-5.965	-8.201	-10.379	-7.212	-9.042
Cl ₂ S ₂	-17.105	-13.944	-13.691	-15.524	-16.450	-15.741	-16.358
SiCl ₂ singlet	-2.265	-1.010	-1.528	-4.145	-5.774	-2.385	-4.071
CF ₃ Cl	-14.974	-9.368	-15.540	-13.283	-13.494	-12.746	-13.150

Continued on next page

TABLE XI: Deviation of r^4 SCAN atomization energies (kcal/mol) for the G3 test set obtained with increasingly dense numerical grids. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	1	2	3	4	5	6	7
Hexafluoro ethane (C_2F_6)	-22.890	-12.265	-25.936	-21.000	-20.361	-20.022	-20.200
CF_3	-15.403	-10.249	-17.423	-14.911	-14.270	-14.116	-14.224
C_6H_5 (phenyl radical)	-21.894	-18.951	-19.817	-19.454	-19.428	-20.456	-19.766
Bicyclo[1.1.0]butane (C_4H_6)	-6.268	-6.914	-6.812	-6.160	-6.353	-7.014	-6.620
$(CH_3)_3C$ (t-butyl radical)	-10.850	-11.132	-10.862	-10.814	-10.758	-11.339	-10.952
Trans-butane(C_4H_{10})	-5.521	-5.212	-5.300	-5.205	-5.124	-5.652	-5.265
1,3 Cyclohexadiene (C_6H_8)	-11.876	-9.572	-10.045	-9.775	-9.781	-10.728	-10.091
ME	-7.725	-5.631	-6.915	-7.104	-6.998	-7.054	-6.939
MAE	8.607	6.749	7.653	7.765	7.841	7.805	7.716

Appendix G: Weakly Bound Complexes

TABLE XII: Error in dissociation energies (kcal/mol) for the S22 test set. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
0	-0.032	-0.259	-0.048	-0.236	-0.245
1	0.383	0.051	0.392	0.135	0.142
2	2.157	1.332	2.087	1.248	1.480
3	0.478	-0.181	0.511	-0.181	-0.131
4	0.025	-0.573	0.105	-0.640	-0.674
5	0.031	-0.434	0.207	-0.547	-0.563
6	-0.502	-1.045	-0.343	-1.108	-1.080
7	-0.176	-0.298	-0.196	-0.238	-0.233
8	-0.450	-0.618	-0.451	-0.614	-0.599
9	-0.498	-0.862	-0.516	-0.660	-0.630
10	-1.375	-2.431	-1.530	-1.824	-1.697
11	-1.370	-2.379	-1.525	-1.868	-1.737
12	-1.366	-2.605	-1.522	-2.000	-1.885
13	-2.335	-3.678	-2.540	-2.962	-2.803
14	-2.553	-4.153	-2.812	-3.500	-3.297
15	-0.076	-0.276	-0.089	-0.191	-0.179
16	0.075	-0.421	0.061	-0.108	-0.049
17	-0.235	-0.689	-0.257	-0.406	-0.360
18	-0.348	-0.797	-0.351	-0.580	-0.507
19	-1.012	-1.494	-1.035	-1.262	-1.206
20	-1.344	-1.950	-1.314	-1.619	-1.563
21	-1.017	-1.599	-1.022	-1.454	-1.416
ME	-0.524	-1.153	-0.554	-0.937	-0.874
MAE	0.798	1.273	0.846	1.057	1.015

TABLE XIII: Error in reaction barrier heights (kcal/mol) for the BH76 test set. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
0	-8.791	-8.273	-8.096	-7.781	-8.427
1	-18.390	-21.305	-21.950	-20.121	-20.565
2	-13.497	-13.512	-13.451	-13.044	-13.569
3	-13.497	-13.512	-13.451	-13.044	-13.569
4	-8.797	-9.233	-9.052	-8.601	-8.918
5	-8.797	-9.233	-9.052	-8.601	-8.918
6	-10.468	-11.605	-11.294	-11.038	-10.433

Continued on next page

TABLE XIII: Error in reaction barrier heights (kcal/mol) for the BH76 test set. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
7	-10.804	-10.046	-10.548	-10.149	-11.007
8	-13.704	-12.920	-12.713	-12.261	-12.871
9	-16.482	-14.953	-15.504	-15.065	-16.433
10	-12.131	-11.556	-11.777	-11.402	-11.465
11	-15.415	-15.440	-15.763	-15.448	-15.031
12	-7.986	-6.873	-7.332	-7.131	-6.897
13	-7.986	-6.873	-7.332	-7.131	-6.897
14	-5.314	-4.739	-4.867	-4.886	-4.815
15	-5.314	-4.739	-4.867	-4.886	-4.815
16	-8.027	-6.176	-6.608	-6.558	-7.048
17	-8.027	-6.176	-6.608	-6.558	-7.048
18	-6.536	-5.304	-5.403	-5.543	-5.977
19	-6.536	-5.304	-5.403	-5.543	-5.977
20	-9.200	-7.443	-7.774	-7.723	-8.191
21	-5.157	-4.601	-5.210	-4.936	-4.557
22	-4.379	-3.428	-3.444	-3.603	-3.976
23	-4.722	-4.626	-4.901	-4.813	-4.551
24	-7.495	-6.323	-6.857	-6.592	-6.559
25	-8.043	-6.594	-6.931	-6.913	-6.942
26	-5.832	-5.135	-5.402	-5.373	-5.525
27	-3.076	-1.944	-1.759	-2.358	-1.935
28	-10.529	-10.642	-10.509	-10.129	-10.550
29	-0.959	-1.323	-1.339	-1.411	-1.087
30	-6.871	-6.904	-6.734	-6.456	-6.675
31	1.453	1.574	1.426	1.126	1.301
32	-6.209	-5.053	-4.981	-4.880	-5.185
33	1.272	1.077	1.207	1.095	1.162
34	-6.322	-4.408	-4.877	-4.491	-5.078
35	-2.213	-1.624	-1.244	-1.774	-0.915
36	-1.909	-1.600	-1.838	-1.931	-2.238
37	-0.981	-1.217	-1.265	-1.416	-1.469
38	-7.107	-5.951	-5.821	-5.895	-6.641
39	-8.749	-9.552	-9.175	-8.213	-8.855
40	-7.180	-8.073	-7.730	-6.681	-7.342
41	-10.266	-8.153	-8.042	-8.579	-9.469
42	-4.921	-5.684	-5.324	-4.523	-4.827
43	-8.426	-7.503	-7.390	-7.419	-7.504
44	-8.464	-8.399	-8.782	-7.819	-8.601
45	-8.046	-6.660	-7.029	-6.822	-8.051
46	-7.262	-7.834	-7.482	-7.119	-7.120
47	-7.262	-7.834	-7.482	-7.119	-7.120
48	-10.639	-10.641	-11.085	-10.071	-11.052
49	-9.564	-9.015	-9.426	-8.716	-9.803
50	-4.920	-4.845	-5.246	-4.667	-5.144
51	-10.066	-10.265	-10.666	-9.880	-10.036
52	-8.343	-8.513	-8.876	-7.882	-8.504
53	-7.101	-5.577	-6.025	-5.898	-7.215
54	-9.789	-10.369	-10.092	-9.174	-9.382
55	-10.996	-8.393	-8.313	-9.257	-9.883
56	-12.313	-11.802	-12.111	-11.340	-11.701
57	-4.879	-4.797	-5.173	-4.606	-5.249
58	-6.416	-6.726	-6.614	-6.368	-6.243
59	-4.142	-4.791	-4.378	-3.612	-4.134
60	-7.468	-7.637	-7.554	-7.482	-7.749
61	-11.397	-12.823	-12.426	-11.321	-11.524
62	-6.302	-6.444	-6.313	-6.066	-6.141
63	-6.170	-7.599	-7.186	-6.139	-6.414

Continued on next page

TABLE XIII: Error in reaction barrier heights (kcal/mol) for the BH76 test set. Mean error (ME) and Mean Absolute Error (MAE) are given at end of table.

	SCAN	rSCAN	r++SCAN	r ² SCAN	r ⁴ SCAN
64	-13.951	-14.410	-14.775	-13.839	-14.466
65	-11.664	-12.825	-13.257	-12.318	-12.905
66	-3.552	-3.700	-4.064	-3.398	-3.902
67	-10.413	-9.764	-9.998	-9.317	-9.674
68	-1.562	-1.598	-2.065	-1.442	-2.078
69	-9.247	-8.860	-9.097	-8.347	-8.589
70	-5.726	-5.726	-6.070	-5.123	-5.709
71	-5.558	-4.416	-4.878	-4.494	-5.670
72	-6.974	-6.714	-7.068	-6.169	-6.872
73	-7.532	-6.501	-6.873	-6.427	-7.471
74	-4.794	-5.678	-5.849	-5.286	-5.268
75	-4.794	-5.678	-5.849	-5.286	-5.268
ME	-7.653	-7.365	-7.488	-7.125	-7.463
MAE	7.724	7.434	7.556	7.182	7.527