Fragmentation fraction f_{Ω_b} and the $\Omega_b \rightarrow \Omega J/\psi$ decay in the light-front formalism

Yu-Kuo Hsiao
1,* and Chong-Chung ${\rm Lih}^{2,\,\dagger}$

 ¹School of Physics and Information Engineering, Shanxi Normal University, Taiyuan, 030031
 ²Department of Optometry, Central Taiwan University of Science and Technology, Taichung 40601

(Dated: April 14, 2022)

Abstract

One has measured $f_{\Omega_b}\mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi)$ at the level of 10^{-6} , where the fragmentation faction f_{Ω_b} is to evaluate the *b*-quark to Ω_b^- production rate. Using the $\Omega_b \to \Omega$ transition form factors calculated in the light-front quark model, we predict $\mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi) = (5.3^{+3.3+3.8}_{-2.1-2.7}) \times 10^{-4}$. In particular, we extract $f_{\Omega_b} = (0.54^{+0.34+0.39+0.21}_{-0.22-0.28-0.15}) \times 10^{-2}$, demonstrating that the *b* to Ω_b productions are much more difficult than the *b* to $\Lambda_b(\Xi_b)$ ones. Since f_{Ω_b} has not been determined experimentally, f_{Ω_b} added to theoretical branching fractions can be compared to future measurements of the Ω_b decays.

^{*} yukuohsiao@gmail.com

[†] cclih@phys.nthu.edu.tw

I. INTRODUCTION

The anti-triplet *b*-baryons $(\Lambda_b, \Xi_b^0, \Xi_b^-)$ and Ω_b^- all decay weakly [1], where Ω_b belongs to the sextet *b*-baryon states. Interestingly, only Ω_b is allowed to have a direct transition to \mathbf{B}^* in the weak interaction, where \mathbf{B}^* stands for a spin-3/2 decuplet baryon. This is due to the fact that Ω_b and \mathbf{B}^* both have totally symmetric quark orderings. By contrast, the anti-triplet baryon \mathbf{B}_b consisting of $(q_1q_2 - q_2q_1)b$ mismatches \mathbf{B}^* with $(q_1q_2 + q_2q_1)q_3$ in the \mathbf{B}_b to \mathbf{B}^* transition. Clearly, the Ω_b decay into \mathbf{B}^* worths an investigation.

One has barely measured the Ω_b decays. Moreover, the fragmentation fraction $f_{\mathbf{B}_b(\Omega_b)}$ that evaluates the *b*-quark to $\mathbf{B}_b(\Omega_b)$ production rate has not been determined yet. Consequently, the charmful Ω_b decay channel $\Omega_b^- \to \Omega^- J/\Psi$ can only be partially measured. In addition to $\Lambda_b \to \Lambda J/\psi$ and $\Xi_b^- \to \Xi^- J/\psi$, the partial branching fractions are given by [1]

$$f_{\Omega_b} \mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi) = (2.9^{+1.1}_{-0.8}) \times 10^{-6} ,$$

$$f_{\Lambda_b} \mathcal{B}(\Lambda_b \to \Lambda J/\psi) = (5.8 \pm 0.8) \times 10^{-5} ,$$

$$f_{\Xi_b} \mathcal{B}(\Xi_b^- \to \Xi^- J/\psi) = (1.02^{+0.26}_{-0.21}) \times 10^{-5} ,$$
(1)

where $f_{\Xi_b} = f_{\Xi_b^{-(0)}}$. Some theoretical attempts have been given to extract $f_{\mathbf{B}_b(\Omega_b)}$ [2–4]. Using the calculations of $\mathcal{B}(\Lambda_b \to \Lambda J/\psi)$ and $\mathcal{B}(\Xi_b^- \to \Xi^- J/\psi)$ [2, 3], one extracts f_{Λ_b} and f_{Ξ_b} as some certain numbers. Without a careful study of $\Omega_b^- \to \Omega^- J/\Psi$ [2, 3], it is roughly estimated that $f_{\Omega_b} < 0.108$. Therefore, it can be an important task to explore the charmful $\Omega_b^- \to \Omega^- J/\Psi$ decay.

See Fig. 1, $\Omega_b^- \to \Omega^- J/\Psi$ is depicted to proceed through the $\Omega_b^- \to \Omega^-$ transition, while J/Ψ is produced from the internal W-boson emission. To calculate the branching fraction, the information of the $\Omega_b \to \Omega$ transition is required. On the other hand, the light-front quark model has provided its calculation on the $\Omega_c \to \Omega$ transition form factors, such that one interprets the relative branching fractions of $\Omega_c^0 \to \Omega^- \rho^+$ and $\Omega_c^0 \to \Omega^- \ell^+ \bar{\nu}_\ell$ to that of $\Omega^- \pi^+$ [5]. Therefore, we propose to calculate the $\Omega_b^- \to \Omega^-$ transition form factors in the light-front formalism, as applied to the Ω_c decays as well as the other heavy hadron decays [6–24]. We will be able to predict $\mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi)$, and extract f_{Ω_b} . Besides, we will compare the branching fractions of $\Omega_b^- \to \Omega^- J/\Psi$, $\Lambda_b \to \Lambda J/\psi$ and $\Xi_b^- \to \Xi^- J/\psi$, and their fragmentation fractions.




FIG. 1. Feynman diagram for $\Omega_b^- \to \Omega^- J/\Psi$.

II. FORMALISM

According to Fig. 1, the amplitude of $\Omega_b^- \to \Omega^- J/\Psi$ combines the matrix elements of the $\Omega_b^- \to \Omega^-$ transition and J/Ψ production, written as [2, 3]

$$\mathcal{M}(\Omega_b^- \to \Omega^- J/\Psi) = \frac{G_F}{\sqrt{2}} V_{cb} V_{cs}^* a_2 \left\langle J/\psi | \bar{c} \gamma^\mu (1 - \gamma_5) c | 0 \right\rangle \left\langle \Omega^- | \bar{s} \gamma_\mu (1 - \gamma_5) b | \Omega_b^- \right\rangle, \tag{2}$$

where G_F is the Fermi constant, and $V_{cb(s)}^{(*)}$ the Cabibbo-Kobayashi-Maskawa (CKM) matrix element. The factorization derives that $a_2 = c_2^{eff} + c_1^{eff}/N_c$, where $c_{1,2}^{eff}$ are the effective Wilson coefficients, and N_c the color number [25, 26]. For the J/Ψ production, the matrix elements read [27]

$$\langle J/\psi | \bar{c} \gamma^{\mu} (1 - \gamma_5) c | 0 \rangle = m_{J/\psi} f_{J/\psi} \varepsilon^*_{\mu} , \qquad (3)$$

where $m_{J/\psi}$, $f_{J/\psi}$ and ε^*_{μ} are the mass, decay constant and polarization four-vector, respectively. The matrix elements of the $\Omega^-_b(bss) \to \Omega^-(sss)$ transition are parameterized as [21, 28]

$$\begin{split} \langle T^{\mu} \rangle &\equiv \langle \Omega(sss) | \bar{s} \gamma^{\mu} (1 - \gamma_5) b | \Omega_b(bss) \rangle \\ &= \bar{u}_{\alpha} \left[\frac{P^{\alpha}}{M} \left(\gamma^{\mu} F_1^V + \frac{P^{\mu}}{M} F_2^V + \frac{P'^{\mu}}{M'} F_3^V \right) + g^{\alpha \mu} F_4^V \right] \gamma_5 u \\ &- \bar{u}_{\alpha} \left[\frac{P^{\alpha}}{M} \left(\gamma^{\mu} F_1^A + \frac{P^{\mu}}{M} F_2^A + \frac{P'^{\mu}}{M'} F_3^A \right) + g^{\alpha \mu} F_4^A \right] u \,, \end{split}$$
(4)

where $M^{(\prime)}$ and $P^{(\prime)}$ represent the mass and momentum of $\Omega_b(\Omega)$, respectively, and $F_i^{V,A}$ (i = 1, 2, ..., 4) are the form factors. By substituting the matrix elements of Eqs. (3, 4) for those of Eq. (2), we derive the amplitude in the helicity basis [28],

$$\mathcal{M} = c_W \sum_{\lambda_\Omega, \lambda_J} (H^V_{\lambda_\Omega \lambda_J} - H^A_{\lambda_\Omega \lambda_J}), \qquad (5)$$

where $c_W \equiv (G_F/\sqrt{2})V_{cb}V_{cs}^* a_2 m_{J/\psi} f_{J/\psi}$, and $\lambda_{\Omega} = (\pm 3/2, \pm 1/2)$ and $\lambda_J = (0, \pm 1)$ denote the helicity states of Ω and J/Ψ , respectively. Due to the helicity conservation, $\lambda_{\Omega_b} = \lambda_{\Omega} - \lambda_J$ should be respected, where $\lambda_{\Omega_b} = \pm 1/2$. Subsequently, we obtain [28]

$$\begin{aligned} H_{\frac{1}{2}0}^{V(A)} &= \sqrt{\frac{2}{3} \frac{Q_{\pm}^2}{q^2}} \left[F_1^{V(A)} \left(\frac{Q_{\pm}^2 M_{\pm}}{2MM'} \right) \mp \left(F_2^{V(A)} + F_3^{V(A)} \frac{M}{M'} \right) \left(\frac{|\vec{P}'|^2}{M'} \right) \mp F_4^{V(A)} \bar{M}'_{-} \right] , \\ H_{\frac{1}{2}1}^{V(A)} &= -\sqrt{\frac{Q_{\pm}^2}{3}} \left[F_1^{V(A)} \left(\frac{Q_{\pm}^2}{MM'} \right) - F_4^{V(A)} \right] , \\ H_{\frac{3}{2}1}^{V(A)} &= \pm \sqrt{Q_{\pm}^2} F_4^{V(A)} , \end{aligned}$$
(6)

and $H_{-\lambda_{\Omega}-\lambda_{J}}^{V(A)} = \mp H_{\lambda_{\Omega}\lambda_{f}}^{V(A)}$, with $M_{\pm} = M \pm M'$, $Q_{\pm}^{2} = M_{\pm}^{2} - q^{2}$, $\bar{M}_{\pm}^{(\prime)} = (M_{+}M_{-} \pm q^{2})/(2M^{(\prime)})$ and $|\vec{P}'| = \sqrt{Q_{+}^{2}Q_{-}^{2}}/(2M)$.

In the light-front quark model, we can calculate the form factors. To start with, we consider the baryon as a bound state that consists of three quarks q_1 , q_2 and q_3 , where $q_{2,3}$ are combined as a diquark, denoted by $q_{[2,3]}$. Explicitly, the baryon bound state can be written as [9]

$$|\mathbf{B}(P, S, S_z)\rangle = \int \{d^3 p_1\} \{d^3 p_2\} \\ \times 2(2\pi)^3 \delta^3 (\tilde{P} - \tilde{p}_1 - \tilde{p}_2) \sum_{\lambda_1, \lambda_2} \Psi^{SS_z}(\tilde{p}_1, \tilde{p}_2, \lambda_1, \lambda_2) |q_1(p_1, \lambda_1) q_{[2,3]}(p_2, \lambda_2)\rangle,$$
(7)

where p_i and λ_i stand for the momentum and helicity state, respectively, and $\Psi^{SS_z}(\tilde{p}_1, \tilde{p}_2, \lambda_1, \lambda_2)$ is the momentum-space wave function. In the light-front frame, one defines $P = (P^-, P^+, P_\perp)$ with $P^{\pm} = P^0 \pm P^3$ and $P_\perp = (P^1, P^2)$, and $p_i = (p_i^-, p_i^+, p_{i\perp})$ with $p_i^{\pm} = p_i^0 \pm p_i^3$ and $p_{i\perp} = (p_i^1, p_i^2)$, together with $\tilde{P} = (P^+, P_\perp)$ and $\tilde{p}_i = (p_i^+, p_{i\perp})$, which result in $P^+P^- = M^2 + P_\perp^2$ and $p_i^+p_i^- = m_i^2 + p_{i\perp}^2$ with $(m_1, m_2) = (m_{q_1}, m_{q_2} + m_{q_3})$. Moreover, P and p_i are related as $P^+ = p_1^+ + p_2^+$ and $P_\perp = p_{1\perp} + p_{2\perp}$, where

$$p_1^+ = (1 - x)P^+, \ p_2^+ = xP^+,$$

$$p_{1\perp} = (1 - x)P_\perp - k_\perp, \ p_{2\perp} = xP_\perp + k_\perp,$$
(8)

with k_{\perp} from $\vec{k} = (k_{\perp}, k_z)$ the relative momentum. By means of $e_i \equiv \sqrt{m_i^2 + \vec{k}^2}$ the energy of the (di)quark and $M_0 \equiv e_1 + e_2$, the above parameters can be rewritten as

$$(x, 1-x) = (e_2 - k_z, e_1 + k_z)/(e_1 + e_2), \ k_z = \frac{xM_0}{2} - \frac{m_2^2 + k_\perp^2}{2xM_0}.$$
(9)

In addition, we obtain $M_0^2 = (m_1^2 + k_\perp^2)/(1 - x) + (m_2^2 + k_\perp^2)/x$. We also get $(\bar{P}_\mu \gamma^\mu - M_0)u(\bar{P}, S_z) = 0$ with $\bar{P} \equiv p_1 + p_2$, where $p_{1,2}$ describe the internal motions of the internal

quarks. Under the Melosh transformation [7], we derive Ψ^{SS_z} as [19–22]

$$\Psi^{SS_z}(\tilde{p}_1, \tilde{p}_2, \lambda_1, \lambda_2) = \sqrt{\frac{C}{2(p_1 \cdot \bar{P} + m_1 M_0)}} \,\bar{u}(p_1, \lambda_1) \Gamma u(\bar{P}, S_z) \phi(x, k_\perp) \,, \tag{10}$$

where $\Gamma = \Gamma_S(\Gamma_A^{(\alpha)})$ represents the vertex function for the scalar (axial-vector) quantity of the diquark, given by [19–22]

$$\Gamma_S = 1 ,$$

$$\Gamma_A = -\frac{1}{\sqrt{3}} \gamma_5 \not\in^*(p_2, \lambda_2) , \ \Gamma_A^{\alpha} = \epsilon^{*\alpha}(p_2, \lambda_2) .$$
(11)

Moreover, the parameter C for $(\Gamma_{S(A)}, \Gamma_A^{\alpha})$ is given by

$$C = \left(\frac{3(m_1M_0 + p_1 \cdot \bar{P})}{3m_1M_0 + p_1 \cdot \bar{P} + 2(p_1 \cdot p_2)(p_2 \cdot \bar{P})/m_2^2}, \frac{3m_2^2M_0^2}{2m_2^2M_0^2 + (p_2 \cdot \bar{P})^2}\right).$$
 (12)

In Eq. (10), $\phi(x, k_{\perp})$ is the wave function that illustrates the momentum distribution of the constituent quark-diquark states. Here, we present $\phi(x, k_{\perp})$ in the Gaussian form [5, 6, 19–23]:

$$\phi(x,k_{\perp}) = 4\left(\frac{\pi}{\beta^2}\right)^{3/4} \sqrt{\frac{e_1 e_2}{x(1-x)M_0}} \exp\left(\frac{-\vec{k}^2}{2\beta^2}\right) , \qquad (13)$$

with $\beta \equiv \beta_{b[ss]}(\beta_{s[ss]})$ to shape the momentum distribution of the *b*-[*ss*] (*s*-[*ss*]) system in the Ω_b (Ω) bound state.

Using the bound states of $|\Omega_b(P, S, S_z)\rangle$ and $|\Omega(P, S', S'_z)\rangle$ in Eq. (7) and the above identities, we derive the matrix elements of the $\Omega_b \to \Omega$ transition in the light-front frame, given by [21]

$$\langle \bar{T}^{\mu} \rangle \equiv \langle \Omega(P', S' = 3/2, S'_z) | \bar{s} \gamma^{\mu} (1 - \gamma_5) b | \Omega_b(P, S = 1/2, S_z) \rangle$$

$$= \int \{ d^3 p_2 \} \hat{C}^{-1/2} \phi'(x', k'_\perp) \phi(x, k_\perp)$$

$$\times \sum_{\lambda_2} \bar{u}_\alpha(\bar{P}', S'_z) \left[\bar{\Gamma}_A'^\alpha(p'_1 + m'_1) \gamma^\mu (1 - \gamma_5) (p'_1 + m_1) \Gamma_A \right] u(\bar{P}, S_z) ,$$

$$(14)$$

where $m_1^{(\prime)} = m_{b(s)}, \ \bar{\Gamma} = \gamma^0 \Gamma^{\dagger} \gamma^0$ and $\hat{C} = 4p_1^+ p_1^{\prime +} (p_1 \cdot \bar{P} + m_1 M_0) (p_1^{\prime} \cdot \bar{P}^{\prime} + m_1^{\prime} M_0^{\prime}).$

To determine $F_i^{V,A}$, the identities $J_{(5)}^{\mu} \equiv \bar{u}\Gamma^{\mu\beta}(\gamma_5)u_{\beta}$ and $\bar{J}_{(5)}^{\mu} \equiv \bar{u}\bar{\Gamma}^{\mu\beta}(\gamma_5)u_{\beta}$ can be useful, where $\Gamma^{\mu\beta} = (\gamma^{\mu}P^{\beta}, P'^{\mu}P^{\beta}, P^{\mu}P^{\beta}, g^{\mu\beta})$ and $\bar{\Gamma}^{\mu\beta} = (\gamma^{\mu}\bar{P}^{\beta}, \bar{P}'^{\mu}\bar{P}^{\beta}, \bar{P}^{\mu}\bar{P}^{\beta}, g^{\mu\beta})$. We can hence perform the following calculations [5, 21],

$$J_{5} \cdot \langle T \rangle = Tr \left\{ u_{\beta} \bar{u}_{\alpha} \left[\frac{P^{\alpha}}{M} \left(\gamma^{\mu} F_{1}^{V} + \frac{P^{\mu}}{M} F_{2}^{V} + \frac{P'^{\mu}}{M'} F_{3}^{V} \right) + g^{\alpha \mu} F_{4}^{V} \right] \gamma_{5} \bar{u} \Gamma_{\mu}^{\beta} \gamma_{5} \right\},$$

$$\bar{J}_{5} \cdot \langle \bar{T} \rangle = \int \{ d^{3} p_{2} \} \hat{C}^{-1/2} \phi'(x', k'_{\perp}) \phi(x, k_{\perp})$$

$$\times \sum_{\lambda_{2}} Tr \left\{ u_{\beta} \bar{u}_{\alpha} \left[\bar{\Gamma}_{A}'^{\alpha} (p'_{1} + m'_{1}) \gamma^{\mu} (p'_{1} + m_{1}) \Gamma_{A} \right] u \bar{\Gamma}_{\mu}^{\beta} \gamma_{5} \right\}.$$
(15)

By connecting $J_5 \cdot \langle T \rangle$ to $\bar{J}_5 \cdot \langle \bar{T} \rangle$, that is, $J_5 \cdot \langle T \rangle = \bar{J}_5 \cdot \langle \bar{T} \rangle$, F_i^V in $J_5 \cdot \langle T \rangle$ can be extracted with $\bar{J}_5 \cdot \langle \bar{T} \rangle$ in the light-front quark model, as the other extractions of the $\mathbf{B}_{b(c)} \to \mathbf{B}^{(*)}$ transition form factors [5, 6, 19–23]. Similarly, $J \cdot \langle T \rangle = \bar{J} \cdot \langle \bar{T} \rangle$ enables us to get F_i^A . We will present our results in the next section.

III. NUMERICAL ANALYSIS

For the numerical analysis, the CKM matrix elements and the mass (decay constant) of the J/Ψ meson state are given by [1]

$$(V_{cb}, V_{cs}) = (A\lambda^2, 1 - \lambda^2/2),$$

 $(m_{J/\Psi}, f_{J/\Psi}) = (3.097, 0.418) \text{ GeV},$ (16)

with $\lambda = 0.2265$ and A = 0.790 in the Wolfenstein parameterization. The effective Wilson coefficients $(c_1^{eff}, c_2^{eff}) = (1.168, -0.365)$ come from Refs. [25, 26]. In the generalized version of the factorization approach, N_c is taken as a floating number, in order that the nonfactorizable effects from QCD corrections can be estimated. By adopting $N_c = 2.15 \pm 0.17$ in [2, 3], we obtain $a_2 = 0.18^{+0.05}_{-0.04}$, which has been used to interpret $\mathcal{B}(\Lambda_b \to \Lambda J/\psi)$ and $\mathcal{B}(\Xi_b^- \to \Xi^- J/\psi)$.

In terms of $J_5 \cdot \langle T \rangle = \overline{J}_5 \cdot \langle \overline{T} \rangle$ and $J \cdot \langle T \rangle = \overline{J} \cdot \langle \overline{T} \rangle$ and the theoretical inputs in Eqs. (13, 14, 15), given by [23]

$$(m_b, \beta_{b[ss]}) = (5.00 \pm 0.20, 0.78 \pm 0.04) \text{ GeV},$$

 $(m_s, \beta_{s[ss]}) = (0.38, 0.48) \text{ GeV},$ (17)

we derive $F_i^{V(A)}$ as the functions of q^2 , depicted in Fig. 2. It is common that one parameterizes the form factors in the dipole expressions [16, 17, 24], which reproduce the momentum dependences derived in the quark model. Subsequently, the form factors can have simple forms to be used in the weak decays. In our case, we present [5, 6]

$$F(q^2) = \frac{F(0)}{1 - a\left(q^2/m_F^2\right) + b\left(q^4/m_F^4\right)},$$
(18)

with m_F , a, b and F(0) at $q^2 = 0$ given in Table I, in order to describe the momentum behaviors of $F_i^{V,A}$ in Fig. 2.




FIG. 2. $F_i^{V(A)}$ versus q^2 (i = 1, 2, 3, 4).

Thus, we calculate the branching fraction and fragmentation fraction as

$$\mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi) = (5.3^{+3.3+3.8}_{-2.1-2.7}) \times 10^{-4},$$

$$f_{\Omega_b} = (0.54^{+0.34+0.39+0.21}_{-0.22-0.28-0.15}) \times 10^{-2},$$
 (19)

where f_{Ω_b} is extracted with $\mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi)$ and the data in Eq. (1). Moreover, the first and second uncertainties come from a_2 and $F_i^{V,A}$, respectively, and the third one for f_{Ω_b} is from the measurement.

IV. DISCUSSIONS AND CONCLUSIONS

Because of the insufficient information on the $\Omega_b \to \mathbf{B}^*$ transition, the Ω_b decays have not been richly explored. In the light-front quark model, we calculate the $\Omega_b \to \Omega$ transition form factors. We can hence predict $\mathcal{B}(\Omega_b \to \Omega J/\Psi) = (5.3^{+3.3+3.8}_{-2.1-2.7}) \times 10^{-4}$, which is compatible with those of the anti-triplet *b*-baryon decays $\mathcal{B}(\Lambda_b \to \Lambda J/\Psi) = (3.3 \pm 2.0) \times 10^{-4}$ and

TABLE I. The $\Omega_b \to \Omega$ transition form factors with (F(0), a, b) in Eq. (18), where $m_F = 6.05$ GeV is from m_{Ω_b} . The uncertainties come from m_b and $\beta_{b[ss]}$ in Eq. (17).

	F(0)	a	b		F(0)	a	b
F_1^V	$0.371\substack{+0.045\\-0.042}$	-2.22	2.37	F_1^A	$0.329\substack{+0.121\\-0.110}$	-1.93	2.73
F_2^V	$-0.104\substack{+0.022\\-0.025}$	-3.19	4.69	F_2^A	$-0.081\substack{+0.022\\-0.020}$	-3.31	4.36
F_3^V	$0.040\substack{+0.042\\-0.035}$	4.11	11.38	F_3^A	$-0.064^{+0.130}_{-0.140}$	-3.16	0.77
F_4^V	$0.692\substack{+0.054\\-0.051}$	-2.05	1.91	F_4^A	$-0.416\substack{+0.092\\-0.082}$	-1.89	0.99

 $\mathcal{B}(\Xi_b^- \to \Xi^- J/\Psi) = (5.1 \pm 3.2) \times 10^{-4} [2, 3].$ On the other hand, $\mathcal{B}(\Omega_b \to \Omega J/\Psi) = 8.1 \times 10^{-4}$ is given by the authors of Ref. [28]. In addition, the total decay width $\Gamma(\Omega_b \to \Omega J/\Psi) = 3.15a_2^2 \times 10^{10} \text{ s}^{-1}$ [29] leads to $\mathcal{B}(\Omega_b \to \Omega J/\Psi) = 16.7 \times 10^{-4}$, where we have used $a_2 = 0.18$ for the demonstration.

In the helicity basis, the branching fraction is given by

$$\mathcal{B} \propto \left(|\mathcal{H}_V|^2 + |\mathcal{H}_A|^2 \right),\tag{20}$$

where $|\mathcal{H}_{V(A)}|^2 \equiv |H_{\frac{3}{2}1}^{V(A)}|^2 + |H_{\frac{1}{2}1}^{V(A)}|^2 + |H_{\frac{1}{2}0}^{V(A)}|^2$. It is found that $(|\mathcal{H}_V|^2, |\mathcal{H}_A|^2)$ give (19,81)% of \mathcal{B} ; besides, $(|H_{\frac{3}{2}1}^A|^2, |H_{\frac{1}{2}1}^A|^2, |H_{\frac{1}{2}0}^A|^2)/|\mathcal{H}_A|^2 = (54.0, 22.4, 23.6)\%$, such that F_4^A gives the main contribution to $\mathcal{B}(\Omega_b \to \Omega J/\Psi)$.

In Eq. (19), $f_{\Omega_b} = 0.54 \times 10^{-2}$ agrees with the previous upper limit of 0.108 [2]. By comparing our extraction to $f_{\Lambda_b} = 0.175 \pm 0.106$ and $f_{\Xi_b} = 0.019 \pm 0.013$ [2], it demonstrates that the *b* to Ω_b productions are much more difficult than the *b* to \mathbf{B}_b ones. Since the fragmentation fraction has not been determined experimentally, the branching fractions of the Ω_b decays should be partially measured with the factor f_{Ω_b} . Therefore, our extraction for f_{Ω_b} can be useful. With f_{Ω_b} of Eq. (19) added to the branching fractions, one can compare his theoretical results to future measurements of the Ω_b decays.

In summary, we have investigated the charmful Ω_b decay channel $\Omega_b^- \to \Omega^- J/\Psi$. In the light-front quark model, we have studied the $\Omega_b \to \Omega$ transition form factors (F_i^V, F_i^A) (i = 1, 2, ..., 4). We have hence predicted $\mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi) = (5.3^{+3.3+3.8}_{-2.1-2.7}) \times 10^{-4}$, which is compatible with those of the $\Lambda_b \to \Lambda J/\Psi$ and $\Xi_b^- \to \Xi^- J/\Psi$ decays. In addition, F_4^A has been found to give the main contribution. Particularly, we have extracted $f_{\Omega_b} =$ $(0.54^{+0.34+0.39+0.21}_{-0.22-0.28-0.15}) \times 10^{-2}$ from the partial observation $f_{\Omega_b} \mathcal{B}(\Omega_b^- \to \Omega^- J/\Psi) = (2.9^{+1.1}_{-0.8}) \times 10^{-6}$. Since f_{Ω_b} has not been determined experimentally, by adding f_{Ω_b} to the branching fractions, one is allowed to compare his calculations to future observations of the Ω_b decays.

ACKNOWLEDGMENTS

YKH was supported in part by National Science Foundation of China (Grants No. 11675030 and No. 12175128). CCL was supported in part by CTUST (Grant No. CTU109-P-108).

- [1] P.A. Zyla *et al.* [Particle Data Group], PTEP **2020**, 083C01 (2020).
- [2] Y.K. Hsiao, P.Y. Lin, L.W. Luo and C.Q. Geng, Phys. Lett. B 751, 127 (2015).
- [3] Y.K. Hsiao, P.Y. Lin, C.C. Lih and C.Q. Geng, Phys. Rev. D 92, 114013 (2015).
- [4] H.Y. Jiang and F.S. Yu, Eur. Phys. J. C 78, 224 (2018).
- [5] Y.K. Hsiao, L. Yang, C.C. Lih and S.Y. Tsai, Eur. Phys. J. C 80, 1066 (2020).
- [6] Z.X. Zhao, Chin. Phys. C 42, 093101 (2018).
- [7] H.J. Melosh, Phys. Rev. D 9, 1095 (1974).
- [8] P.L. Chung, F. Coester and W.N. Polyzou, Phys. Lett. B 205, 545 (1988).
- [9] H.G. Dosch, M. Jamin and B. Stech, Z. Phys. C 42, 167 (1989).
- [10] W. Jaus, Phys. Rev. D 44, 2851 (1991).
- [11] F. Schlumpf, Phys. Rev. D 47, 4114 (1993); Erratum: [Phys. Rev. D 49, 6246 (1994)].
- [12] C.Q. Geng, C.C. Lih and W.M. Zhang, Mod. Phys. Lett. A 15, 2087 (2000).
- [13] C.R. Ji and C. Mitchell, Phys. Rev. D 62, 085020 (2000).
- [14] B.L.G. Bakker and C.R. Ji, Phys. Rev. D 65, 073002 (2002).
- [15] B.L.G. Bakker, H.M. Choi and C.R. Ji, Phys. Rev. D 67, 113007 (2003).
- [16] H.Y. Cheng, C.K. Chua and C.W. Hwang, Phys. Rev. D 69, 074025 (2004).
- [17] H.M. Choi and C.R. Ji, Few Body Syst. 55, 435 (2014).
- [18] C.Q. Geng and C.C. Lih, Eur. Phys. J. C 73, 2505 (2013).
- [19] H.W. Ke, X.H. Yuan, X.Q. Li, Z.T. Wei and Y.X. Zhang, Phys. Rev. D 86, 114005 (2012).
- [20] H.W. Ke, N. Hao and X.Q. Li, J. Phys. G 46, 115003 (2019).
- [21] Z.X. Zhao, Eur. Phys. J. C 78, 756 (2018).
- [22] X.H. Hu, R.H. Li and Z.P. Xing, Eur. Phys. J. C 80, 320 (2020).
- [23] H.W. Ke, N. Hao and X.Q. Li, Eur. Phys. J. C 79, 540 (2019).
- [24] Y.K. Hsiao, S.Y. Tsai, C.C. Lih and E. Rodrigues, JHEP 2004, 035 (2020).
- [25] A. Ali, G. Kramer, and C.D. Lu, Phys. Rev. **D** 58, 094009 (1998).

- [26] Y.K. Hsiao and C.Q. Geng, Phys. Rev. D 91, 116007 (2015).
- [27] D. Bečirević, G. Duplančić, B. Klajn, B. Melić and F. Sanfilippo, Nucl. Phys. B 883, 306 (2014).
- [28] T. Gutsche, M.A. Ivanov, J.G. Korner and V.E. Lyubovitskij, Phys. Rev. D 98, 074011 (2018).
- [29] H.Y. Cheng, Phys. Rev. D 56, 2799 (1997).