

# LIOUVILLE THEOREMS FOR KIRCHHOFF-TYPE HYPOELLIPTIC PARTIAL DIFFERENTIAL EQUATIONS AND SYSTEMS. I. HEISENBERG GROUP

AIDYN KASSYMOV, MICHAEL RUZHANSKY, NIYAZ TOKMAGAMBETOV,  
AND BERIKBOL TOREBEK

ABSTRACT. In this paper, we show the nonexistence results for the Kirchhoff elliptic, parabolic, and hyperbolic type equations on the Heisenberg groups. Also, the pseudo-parabolic and pseudo-hyperbolic equations of the Kirchhoff-type are under consideration. To prove these results we use the test function method. In addition, the analogous results are transferred to the cases of systems. Also, we give some examples of non-local nonlinearities.

## CONTENTS

1. Introduction	2
1.1. Heisenberg groups	2
1.2. Historical background	2
2. Kirchhoff type elliptic equations and systems	4
2.1. Elliptic type equations	4
2.2. Elliptic type systems	8
3. Kirchhoff type parabolic equations and systems	13
3.1. Parabolic type equations	13
3.2. Parabolic type systems	16
3.3. Pseudo-parabolic type equations and systems	20
4. Kirchhoff type hyperbolic equations and systems	23
4.1. Hyperbolic type equations	23
4.2. Hyperbolic Kirchhoff type systems	27
4.3. Pseudo-hyperbolic equations and systems	30
References	35

---

*Date:* October 5, 2021.

*2010 Mathematics Subject Classification.* 35R11, 34B10, 35R03.

*Key words and phrases.* Kohn-Laplacian, blow-up, Kato-type exponent, Fujita-type exponent, Kirchhoff type equation, Heisenberg group, heat equation, wave equation, pseudo-parabolic equation, pseudo-hyperbolic equation.

The authors were supported by the FWO Odysseus 1 grant G.0H94.18N: Analysis and Partial Differential Equations. MR was supported in parts by the EPSRC Grant EP/R003025/2 and by the Methusalem programme of the Ghent University Special Research Fund (BOF) (Grant number 01M01021).

## 1. INTRODUCTION

**1.1. Heisenberg groups.** Let us briefly recall the definition of the Heisenberg group. For  $n \in \mathbb{N}$ ,  $\mathbb{H}^n := (\mathbb{R}^{2n+1}, \cdot)$  is called the Heisenberg group if the group law is given by

$$x \cdot x' = (\xi_1 + \xi_2, \tilde{\xi}_1 + \tilde{\xi}_2, \tau_1 + \tau_2 + 2(\xi_1 \tilde{\xi}_2 - \xi_2 \tilde{\xi}_1)), \quad (1.1)$$

for any  $x = (\xi_1, \tilde{\xi}_1, \tau_1)$  and  $x' = (\xi_2, \tilde{\xi}_2, \tau_2)$ , where  $\xi_1, \xi_2, \tilde{\xi}_1, \tilde{\xi}_2 \in \mathbb{R}^n$  and  $\tau_1, \tau_2 \in \mathbb{R}$ . The family of dilations has the following form

$$\delta_\lambda(x) := (\lambda\xi, \lambda\tilde{\xi}, \lambda^2\tau), \quad \forall \lambda > 0. \quad (1.2)$$

So, the homogeneous dimension of  $\mathbb{H}^n$  is  $Q = 2n + 2$  and the topological dimension is  $2n + 1$ . The Lie algebra  $\mathfrak{h}$  of the left-invariant vector fields on the Heisenberg group  $\mathbb{H}^n$  is spanned by

$$X_i = \frac{\partial}{\partial \xi_i} + 2\tilde{\xi}_i \frac{\partial}{\partial \tau}, \quad Y_i = \frac{\partial}{\partial \tilde{\xi}_i} - 2\xi_i \frac{\partial}{\partial \tau}, \quad x = (\xi, \tilde{\xi}, \tau),$$

for  $i = 1, \dots, n$ , with their non-zero commutator

$$[X_i, Y_i] = -4\partial_\tau.$$

The horizontal gradient on  $\mathbb{H}^n$  is given by

$$\nabla_{\mathbb{H}} := (X_1, \dots, X_n, Y_1, \dots, Y_n).$$

So, the Kohn-Laplacian or sub-Laplacian is given by

$$\Delta_{\mathbb{H}} := \sum_{i=1}^n (X_i^2 + Y_i^2).$$

Finally, we define the (Kaplan) distance on the Heisenberg group by

$$|x|_H^4 = \tau^2 + \sum_{i=1}^n (|\xi|^2 + |\tilde{\xi}|^2)^2,$$

for  $x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n$ . For further details, we refer to the book [9].

Now let us denote by  $S_1^2(\mathbb{H}^n)$  the Sobolev space on the Heisenberg group:

$$S_1^2(\mathbb{H}^n) := \{u : u \in L^2(\mathbb{H}^n), |\nabla_{\mathbb{H}} u| \in L^2(\mathbb{H}^n)\}. \quad (1.3)$$

Similarly, let us denote  $S_2^2(\mathbb{H}^n)$  in the following form:

$$S_2^2(\mathbb{H}^n) := \{u : u \in L^2(\mathbb{H}^n), |\nabla_{\mathbb{H}} u| \in L^2(\mathbb{H}^n), |\Delta_{\mathbb{H}} u| \in L^2(\mathbb{H}^n)\}. \quad (1.4)$$

**1.2. Historical background.** In the celebrated paper [10], Fujita studied the following nonlinear heat equation

$$\begin{cases} u_t(x, t) - \Delta u(x, t) = u^{1+p}(x, t), & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ u(x, 0) = u_0(x) \geq 0, & x \in \mathbb{R}^N. \end{cases} \quad (1.5)$$

He showed that if  $0 < p < \frac{2}{N}$  then a solution of the problem (1.5) blows up in finite time for  $N > 2$  while being globally well-posed for  $p > \frac{2}{N}$ , causing to call this critical exponent “the critical exponent of Fujita” or just “Fujita’s exponent”. One of the further generalisations of the problem (1.5) is considering the fractional Laplacian

$(-\Delta)^s$  instead of the classical one  $(-\Delta)$ . That is, in [15, 30] the authors considered the Cauchy problem

$$\begin{cases} u_t(x, t) + (-\Delta)^s u(x, t) = a(x, t)|u(x, t)|^{1+p}, & (x, t) \in \mathbb{R}^N \times (0, \infty), \\ u(x, 0) = u_0(x) \geq 0, & x \in \mathbb{R}^N, \end{cases}$$

for  $s > 0$ .

Consequently, in [31], Véron and Pohozaev found a critical exponent for the following non-linear diffusion equation with the Kohn-Laplacian on Heisenberg groups

$$\frac{\partial u(x, t)}{\partial t} - \Delta_{\mathbb{H}} u(x, t) = |u(x, t)|^p, \quad (x, t) \in \mathbb{H}^n \times (0, +\infty).$$

Also, the critical exponents for other equations with the Kohn-Laplacian on the Heisenberg groups were derived in [1, 2, 7, 17]. In addition, in [28] the authors found the Fujita exponent on general unimodular Lie groups.

In the paper [20], T. Kato showed the nonexistence of the global solutions for the Cauchy problem for a non-linear wave equation. This work was a pioneering work in this field for the hyperbolic type equations. The corresponding order is called the Kato exponent. As a generalisation to the Heisenberg group  $\mathbb{H}^n$ , Véron and Pohozaev in [31] considered the following problem

$$\frac{\partial^2 u(x, t)}{\partial t^2} - \Delta_{\mathbb{H}} u(x, t) = |u(x, t)|^p, \quad (x, t) \in \mathbb{H}^n \times (0, +\infty),$$

with the Cauchy data

$$u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x),$$

where  $p > 1$ . Also, in the work [1] a fractional analogue was studied, namely, the following non-linear wave equation

$$\frac{\partial^2 u(x, t)}{\partial t^2} + (-\Delta_{\mathbb{H}})^s u(x, t) = |u(x, t)|^p, \quad (x, t) \in \mathbb{H}^n \times (0, +\infty),$$

with the Cauchy data

$$u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x),$$

where  $(-\Delta_{\mathbb{H}})^s$  is the fractional Kohn-Laplacian,  $s \in (0, 1)$ ,  $p > 1$ . The wave equation with nonlinear damping on the Heisenberg and graded groups was studied in [27]. In [13, 14], one found a critical exponents for the semilinear heat equation and wave equation with damping term on the Heisenberg group. Also, comparison principles for heat equations with Rockland operators on general graded groups were considered in [26] and [29].

In [3], under certain assumptions on  $f$  for the semilinear equation

$$\begin{cases} -\Delta u = f(x, u), & x \in \Omega \subset \mathbb{R}^n, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \quad (1.6)$$

Ambrosetti and Rabinowitz proved the existence of solutions by the mountain pass theorem, leading to a number of extensions and generalisations. Indeed, the authors proved the existence results of the weak solutions of (1.6) in the model case  $f(x, u) = a(x)|u|^{q-1}u$ , where  $1 < q < \frac{n+2}{n-2}$ .

By the fibering method the authors of the work [8] proved the existence of the weak solutions to

$$\begin{cases} -\Delta_p u(x) = \lambda|u|^{p-2}u + f(x)|u|^{q-2}u, & x \in \Omega \subset \mathbb{R}^n, \\ u(x) = 0, & x \in \partial\Omega, \end{cases} \quad (1.7)$$

where  $p, \lambda, q \in \mathbb{R}$  and  $1 < p < q < p^* = \frac{np}{n-p}$ .

In the work [6], the author considered and showed that there is at least one weak solution to the following problem on Heisenberg groups

$$\begin{cases} -\Delta_{\mathbb{H}} u + au = |u|^{\frac{Q+2}{Q-2}}, & x \in \Omega \subset \mathbb{H}^n, \\ u(x) = 0, & x \in \partial\Omega. \end{cases} \quad (1.8)$$

Also, see [12].

An important generalisation of the semilinear elliptic type equations is the Kirchhoff type equations, which arise in the description of nonlinear vibrations of an elastic string. Also, the study of Kirchhoff-type problems has been receiving considerable attention in recent years, see [4, 5, 16, 18, 19, 21, 22, 23, 24, 25] and [32]. This interest arises from their contributions to the modeling of many physical and biological phenomena. The main aim of this note is to extend the above results to the Heisenberg group case, or more general cases. Thus, we will consider nonlinear Kirchhoff type problems on the Heisenberg group. Note that there is already a number of results related to the non-existence of solutions to the semilinear equations, mentioning only few of the pioneering works on the Heisenberg group, for instance, [1], [2] and [31]. In the present paper, we will use the test function approach, and new interesting results on the Heisenberg group will be proved. Some results obtained here can be extended to the general stratified Lie groups and graded Lie groups.

In this paper, we also study the Kirchhoff type nonlinear systems of elliptic, parabolic, and hyperbolic type equations. In all of our equations the coefficient which expresses the Kirchhoff type non-linearity will be denoted by  $M$ . In each individual case we will ask different properties on  $M$ . But the following particular cases could be a few examples of the function  $M$ :

- a)  $M = 1$ ,
- b)  $M = \frac{1}{1+a \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx + b \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx}$ , where  $a, b$  are some real constants.
- c)  $M = (1 + c|t|^\gamma)^{-\beta}$ , where  $c, \gamma$  are some real constants and  $\beta > 0$ .
- d)  $M = \exp(- (a \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx + b \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx))$ , where  $a, b$  are real constants.
- e)  $M = \log \left( \frac{a+b \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx + c \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx}{a_1 + b_1 \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx + c_1 \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx} \right)$ ,  $a, b, c, a_1, b_1, c_1$  are some real constants.

## 2. KIRCHHOFF TYPE ELLIPTIC EQUATIONS AND SYSTEMS

In this section we find the critical exponents for the Kirchhoff type elliptic equations on the Heisenberg group.

**2.1. Elliptic type equations.** Let us consider the following Kirchhoff type elliptic equation

$$-M \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u(x)|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u(x)|^2 dx \right) \Delta_{\mathbb{H}} u(x) = |u(x)|^p, \quad x \in \mathbb{H}^n, \quad p > 1, \quad (2.1)$$

where  $u \in C^2(\mathbb{H}^n)$  and  $M : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  is a bounded function such that

$$0 < M(\cdot, \cdot) \leq C_0. \quad (2.2)$$

Let us introduce a notion of the weak solution to the equation (2.1).

**Definition 2.1.** We say that  $u \in S_2^2(\mathbb{H}^n) \cap L^p(\mathbb{H}^n)$  with  $p > 1$  is a weak solution of the Kirchhoff type elliptic equation (2.1) on  $\mathbb{H}^n$ , if the following identity

$$\int_{\mathbb{H}^n} |u|^p \varphi dx = - \int_{\mathbb{H}^n} M \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx$$

holds true, for any test function  $0 \leq \varphi \in C_0^2(\mathbb{H}^n)$ .

Then we have the following result.

**Theorem 2.2.** Assume that the function  $M(\cdot, \cdot)$  satisfies (2.2) and

$$1 < p \leq \frac{Q}{Q-2}, \quad (2.3)$$

with  $Q = 2n + 2$  the homogeneous dimension of  $\mathbb{H}^n$ . Then the problem (2.1) does not have a global non-trivial weak solution.

*Proof.* In the proof we widely use the method of test functions. We start from Definition 2.1. From the identity (2.1) one has

$$\begin{aligned} \int_{\mathbb{H}^n} |u|^p \varphi dx &= - \int_{\mathbb{H}^n} M \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx \\ &\leq \left| \int_{\mathbb{H}^n} M \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx \right| \\ &\leq \int_{\mathbb{H}^n} \left| M \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) \right| |u| |\Delta_{\mathbb{H}} \varphi| dx \\ &\leq C_0 \int_{\mathbb{H}^n} |u| |\Delta_{\mathbb{H}} \varphi| dx. \end{aligned} \quad (2.4)$$

Now, we choose the following test function

$$\varphi_R(x) = \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right), \quad x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n, \quad R > 0, \quad (2.5)$$

with  $\Phi \in C_0^\infty(\mathbb{R}_+)$  satisfying the following structure

$$\Phi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2, \end{cases} \quad (2.6)$$

with  $\text{supp } \Phi = [0, 2]$ . We note that  $\text{supp}(\varphi_R)$  and  $\text{supp}(\Delta_{\mathbb{H}} \varphi_R)$  are subsets of  $\Omega_1$ , where

$$\Omega_1 := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\}.$$

By using  $\varepsilon$ -Young's inequality in (2.4), we obtain

$$\begin{aligned}
\int_{\mathbb{H}^n} |u|^p \varphi_R dx &\leq C_0 \int_{\mathbb{H}^n} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\
&\stackrel{(2.6)}{=} C_0 \int_{\Omega_1} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\
&= C_0 \int_{\Omega_1} |u| \varphi_R^{\frac{1}{p}} \varphi_R^{-\frac{1}{p}} |\Delta_{\mathbb{H}} \varphi_R| dx \\
&= C_0 \left( \frac{\int_{\Omega_1} |u|^p \varphi_R dx}{\varepsilon^p p} + \frac{(p-1) \varepsilon^{\frac{p}{p-1}} \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx}{p} \right) \\
&= \frac{C_0}{p \varepsilon^p} \left( \int_{\Omega_1} |u|^p \varphi_R dx + (p-1) \varepsilon^{p+\frac{p}{p-1}} \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right).
\end{aligned} \tag{2.7}$$

By choosing  $\varepsilon$  such that  $C_0 < p \varepsilon^p$  we get

$$\int_{\mathbb{H}^n} |u|^p \varphi_R dx \leq C \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right). \tag{2.8}$$

Let us denote by

$$\rho = \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4}.$$

Then from [31], one has the following property

$$\begin{aligned}
\Delta_{\mathbb{H}} \varphi_R(x) &= \frac{4(n+4)}{R^4} \left( |\xi|^2 + |\tilde{\xi}|^2 \right) \Phi'(\rho) \\
&\quad + \frac{16}{R^8} \left( (|\xi|^6 + |\tilde{\xi}|^6) + 2\tau(|\xi|^2 - |\tilde{\xi}|^2) \xi \cdot \tilde{\xi} + \tau^2(|\xi|^2 + |\tilde{\xi}|^2) \right) \Phi''(\rho).
\end{aligned} \tag{2.9}$$

Taking into account (2.9), we obtain

$$|\Delta_{\mathbb{H}} \varphi_R| \leq C R^{-2}. \tag{2.10}$$

Let us denote

$$\Theta := \{y = (\bar{\xi}, \hat{\xi}, \tilde{\tau}) \in \mathbb{H}^n : |\bar{\xi}|^4 + |\hat{\xi}|^4 + \tilde{\tau}^2 \leq 2\},$$

and

$$\mu := |\bar{\xi}|^4 + |\hat{\xi}|^4 + \tilde{\tau}^2.$$

By changing variables  $R\bar{\xi} = \xi$ ,  $R\hat{\xi} = \tilde{\xi}$ ,  $R^2\tilde{\tau} = \tau$  in (2.8) and by using (2.10), we get

$$\begin{aligned}
\int_{\mathbb{H}^n} |u|^p \varphi_R dx &\leq C \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right) \\
&\leq C R^{-\frac{2p}{p-1}+Q} \left( \int_{\Theta} (\Phi \circ \mu)^{-\frac{1}{p-1}} dy \right) \\
&\leq C R^{-\frac{2p}{p-1}+Q}.
\end{aligned} \tag{2.11}$$

Let us consider the case  $p < \frac{Q}{Q-2}$ , and we note that  $-\frac{2p}{p-1} + Q < 0$  is equivalent to  $p < \frac{Q}{Q-2}$ . Then letting  $R \rightarrow \infty$  in (2.11), we arrive at

$$\int_{\mathbb{H}^n} |u|^p dx \leq C \lim_{R \rightarrow \infty} R^{-\frac{2p}{p-1}+Q} = 0, \tag{2.12}$$

concluding that  $u = 0$ . This is a contradiction.

Let us consider the case  $p = \frac{Q}{Q-2} > 1$ . In this case, from (2.12), we get

$$\int_{\mathbb{H}^n} |u|^p \varphi_R dx \leq C, \quad (2.13)$$

where  $C > 0$ .

Let us introduce the following domain:

$$\bar{\Omega}_R := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : R^4 \leq |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\}, \quad (2.14)$$

then by combining (2.13) and the Lebesgue dominated convergence theorem, we have

$$\lim_{R \rightarrow \infty} \int_{\bar{\Omega}_R} |u|^p \varphi_R dx = 0. \quad (2.15)$$

Let us denote by

$$\Theta_1 := \{y = (\bar{\xi}, \hat{\xi}, \tilde{\tau}) \in \mathbb{H}^n : 1 \leq |\bar{\xi}|^4 + |\hat{\xi}|^4 + \tilde{\tau}^2 \leq 2\},$$

By using Hölder's inequality in (2.4), we get

$$\begin{aligned} \int_{\mathbb{H}^n} |u|^p \varphi_R dx &\leq C_0 \int_{\Omega_1} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\ &\leq C_0 \int_{\Omega_1 \setminus \bar{\Omega}_R} |u| |\Delta_{\mathbb{H}} \varphi_R| dx + C_0 \int_{\bar{\Omega}_R} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\ &\stackrel{(2.6)}{=} C_0 \int_{\bar{\Omega}_R} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\ &= C_0 \int_{\bar{\Omega}_R} |u| \varphi_R^{\frac{1}{p}} \varphi_R^{-\frac{1}{p}} |\Delta_{\mathbb{H}} \varphi_R| dx \\ &\leq C_0 \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx \right)^{\frac{1}{p}} \left( \int_{\bar{\Omega}_R} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \\ &\stackrel{(2.11)}{=} C_0 R^{-2 + \frac{Q(p-1)}{p}} \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx \right)^{\frac{1}{p}} \left( \int_{\Theta_1} (\Phi \circ \mu)^{-\frac{1}{p-1}} dy \right)^{\frac{p-1}{p}} \\ &\stackrel{p = \frac{Q}{Q-2}}{=} C \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx \right)^{\frac{1}{p}} \left( \int_{\Theta_1} (\Phi \circ \mu)^{-\frac{1}{p-1}} dy \right)^{\frac{p-1}{p}} \\ &\leq C \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx \right)^{\frac{1}{p}}. \end{aligned}$$

Finally, we get

$$\int_{\mathbb{H}^n} |u|^p \varphi_R dx \leq C \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx \right)^{\frac{1}{p}}.$$

By letting  $R \rightarrow \infty$  and using (2.15), we get

$$\int_{\mathbb{H}^n} |u|^p dx = \lim_{R \rightarrow \infty} \int_{\mathbb{H}^n} |u|^p \varphi_R dx \leq C \lim_{R \rightarrow \infty} \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx \right)^{\frac{1}{p}} = 0,$$

that is,  $u = 0$ , ending the proof.  $\square$

**2.2. Elliptic type systems.** In this subsection we consider the following system of Kirchhoff type elliptic equations

$$\begin{cases} -M_1 \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) \Delta_{\mathbb{H}} u = |v|^q, & x \in \mathbb{H}^n, \quad q > 1, \\ -M_2 \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) \Delta_{\mathbb{H}} v = |u|^p, & x \in \mathbb{H}^n, \quad p > 1, \end{cases} \quad (2.16)$$

where  $u, v \in C^2(\mathbb{H}^n)$  and  $M_1, M_2 : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  are bounded functions such that

$$0 < M_1(\cdot, \cdot) \leq C_1, \quad (2.17)$$

and

$$0 < M_2(\cdot, \cdot) \leq C_2. \quad (2.18)$$

Now, we formulate a definition of weak solutions to the system (2.16).

**Definition 2.3.** We say that a pair of functions  $(u, v) \in S_2^2(\mathbb{H}^n) \cap L^p(\mathbb{H}^n) \times S_2^2(\mathbb{H}^n) \cap L^q(\mathbb{H}^n)$  with  $p, q > 1$  is a weak solution of the system of equations (2.16) on the group  $\mathbb{H}^n$ , if the following identities

$$\int_{\mathbb{H}^n} |v|^q \varphi dx = - \int_{\mathbb{H}^n} M_1 \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx, \quad (2.19)$$

and

$$\int_{\mathbb{H}^n} |u|^p \psi dx = - \int_{\mathbb{H}^n} M_2 \left( \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) v \Delta_{\mathbb{H}} \psi dx, \quad (2.20)$$

hold true, for any test functions  $0 \leq \psi, \varphi \in C_0^2(\mathbb{H}^n)$ .

In the following statement, we give a nonexistence result for the system (2.16).

**Theorem 2.4.** *Let  $p, q > 1$ . Suppose  $M_1$  and  $M_2$  satisfy (2.17) and (2.18), respectively. Assume that the following inequality*

$$Q \left( 1 - \frac{1}{pq} \right) \leq 2 \max \left\{ 1 + \frac{1}{q}, 1 + \frac{1}{p} \right\}, \quad (2.21)$$

*holds with  $Q = 2n + 2$  the homogeneous dimension of  $\mathbb{H}^n$ . Then the system of elliptic type equations (2.16) does not have global non-trivial weak solution.*

*Proof.* In analogy with the single equation case, one can show that

$$\int_{\mathbb{H}^n} |v|^q \varphi dx \leq C_0 \int_{\mathbb{H}^n} |u| |\Delta_{\mathbb{H}} \varphi| dx, \quad \int_{\mathbb{H}^n} |u|^p \psi dx \leq C_1 \int_{\mathbb{H}^n} |v| |\Delta_{\mathbb{H}} \psi| dx. \quad (2.22)$$

Now, we choose the following test functions

$$\varphi_R(x) = \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right), \quad x = (\xi, \tilde{\xi}, \tau), \quad R > 0,$$

and

$$\psi_R(x) = \Psi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right), \quad x = (\xi, \tilde{\xi}, \tau), \quad R > 0,$$



where  $\Phi, \Psi \in C_0^\infty(\mathbb{R}_+)$  have the following property

$$\Phi(r), \Psi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2, \end{cases}$$

with  $\text{supp } \Psi = \text{supp } \Phi = [0, 2]$ . We note that  $\text{supp}(\varphi_R)$ ,  $\text{supp}(\psi_R)$ ,  $\text{supp}(\Delta_{\mathbb{H}}\varphi_R)$  and  $\text{supp}(\Delta_{\mathbb{H}}\psi_R)$  are subsets of  $\Omega_1$ , where

$$\Omega_1 := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\}.$$

Then for the first inequality, by using Hölder's inequality, one obtains

$$\begin{aligned} \int_{\mathbb{H}^n} |v|^q \varphi_R dx &\leq C_0 \int_{\mathbb{H}^n} |u| |\Delta_{\mathbb{H}}\varphi_R| dx \\ &= C_0 \int_{\Omega_1} |u| |\Delta_{\mathbb{H}}\varphi_R| dx \\ &= C_0 \int_{\Omega_1} |u| \psi_R^{\frac{1}{p}} \psi_R^{-\frac{1}{p}} |\Delta_{\mathbb{H}}\varphi_R| dx \\ &\leq C_0 \left( \int_{\Omega_1} |u|^p \psi_R dx \right)^{\frac{1}{p}} \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}\varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \\ &= C_0 \left( \int_{\mathbb{H}^n} |u|^p \psi_R dx \right)^{\frac{1}{p}} \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}\varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}}. \end{aligned} \quad (2.23)$$

Similarly, we have

$$\int_{\mathbb{H}^n} |u|^p \psi_R dx \leq C_1 \left( \int_{\mathbb{H}^n} |v|^q \varphi_R dx \right)^{\frac{1}{q}} \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}}\psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{q}}. \quad (2.24)$$

By combining (2.23) and (2.24), one finds

$$\begin{aligned} \int_{\mathbb{H}^n} |v|^q \varphi_R dx &\leq C_0 \left( \int_{\mathbb{H}^n} |u|^p \psi_R dx \right)^{\frac{1}{p}} \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}\varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \\ &\leq C_0 C_1^{\frac{1}{p}} \left( \int_{\mathbb{H}^n} |v|^q \varphi_R dx \right)^{\frac{1}{pq}} \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}}\psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{pq}} \\ &\quad \times \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}\varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}}. \end{aligned} \quad (2.25)$$

Finally, we obtain

$$\begin{aligned} \left( \int_{\mathbb{H}^n} |v|^q \varphi_R dx \right)^{1-\frac{1}{pq}} &\leq C_0 C_1^{\frac{1}{p}} \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}}\psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{pq}} \\ &\quad \times \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}\varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}}. \end{aligned} \quad (2.26)$$

Also, we have

$$\begin{aligned} \left( \int_{\mathbb{H}^n} |u|^p \psi_R dx \right)^{1-\frac{1}{pq}} &\leq C_1 C_0^{\frac{1}{q}} \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}} \psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{q}} \\ &\times \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{pq}}. \end{aligned} \quad (2.27)$$

Taking into account (2.9), one obtains

$$|\Delta_{\mathbb{H}} \psi_R| \leq CR^{-2}, \quad |\Delta_{\mathbb{H}} \varphi_R| \leq CR^{-2}. \quad (2.28)$$

Similarly to equation case, let us denote

$$\Theta := \{y = (\bar{\xi}, \hat{\xi}, \tilde{\tau}) \in \mathbb{H}^n : |\bar{\xi}|^4 + |\hat{\xi}|^4 + \tilde{\tau}^2 \leq 2\},$$

and

$$\mu := |\bar{\xi}|^4 + |\hat{\xi}|^4 + \tilde{\tau}^2.$$

By changing variables  $R\bar{\xi} = \xi$ ,  $R\hat{\xi} = \tilde{\xi}$  and  $R^2\tilde{\tau} = \tau$  in (2.26), we get

$$\begin{aligned} &\left( \int_{\mathbb{H}^n} |v|^q \varphi_R dx \right)^{1-\frac{1}{pq}} \\ &\leq C_0 C_1^{\frac{1}{p}} \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}} \psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{pq}} \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \\ &\leq C_0 C_1^{\frac{1}{p}} R^{-\frac{2}{p} + \frac{Q(q-1)}{pq} + \frac{Q(p-1)}{p} - 2} \left( \int_{\Theta} (\Phi \circ \mu)^{-\frac{1}{q-1}} dy \right)^{\frac{q-1}{pq}} \left( \int_{\Theta} (\Psi \circ \mu)^{-\frac{1}{p-1}} dy \right)^{\frac{p-1}{p}} \quad (2.29) \\ &= C_0 C_1^{\frac{1}{p}} R^{l_1} \left( \int_{\Theta} (\Phi \circ \mu)^{-\frac{1}{q-1}} dy \right)^{\frac{q-1}{pq}} \left( \int_{\Theta} (\Psi \circ \mu)^{-\frac{1}{p-1}} dy \right)^{\frac{p-1}{p}} \\ &\leq CR^{l_1}, \end{aligned}$$

where  $l_1 = -\frac{2}{p} + \frac{Q(q-1)}{pq} + \frac{Q(p-1)}{p} - 2$ .

Also, we have

$$\begin{aligned} &\left( \int_{\mathbb{H}^n} |u|^p \psi_R dx \right)^{1-\frac{1}{pq}} \\ &\leq C_1 C_0^{\frac{1}{q}} \left( \int_{\Omega_1} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}} \psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{q}} \left( \int_{\Omega_1} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{pq}} \\ &= C_1 C_0^{\frac{1}{q}} R^{l_2} \left( \int_{\Theta} (\Phi \circ \mu)^{-\frac{1}{q-1}} dy \right)^{\frac{q-1}{q}} \left( \int_{\Theta} (\Psi \circ \mu)^{-\frac{1}{p-1}} dy \right)^{\frac{p-1}{pq}} \\ &\leq CR^{l_2}, \end{aligned}$$

where  $l_2 = -\frac{2}{q} + \frac{Q(p-1)}{pq} + \frac{Q(q-1)}{q} - 2$ .

Simple calculations show that

$$\begin{aligned}
 l_1 &= -\frac{2}{p} + \frac{Q(q-1)}{pq} + \frac{Q(p-1)}{p} - 2 \\
 &= -\frac{2}{p} - 2 + Q \left( \frac{1}{p} - \frac{1}{pq} + 1 - \frac{1}{p} \right) \\
 &= -\frac{2}{p} - 2 + Q \left( 1 - \frac{1}{pq} \right) < 0,
 \end{aligned} \tag{2.30}$$

provided that

$$Q < \frac{2 + \frac{2}{p}}{1 - \frac{1}{pq}}.$$

Then letting  $R \rightarrow \infty$ , we obtain

$$\left( \int_{\mathbb{H}^n} |v|^q \varphi_R dx \right)^{1 - \frac{1}{pq}} \leq 0,$$

showing that  $\int_{\mathbb{H}^n} |v|^q dx = 0$ .

Similar calculations works in the case  $l_2 < 0$ , i.e.,

$$Q \left( 1 - \frac{1}{pq} \right) < 2 \left( 1 + \frac{1}{q} \right).$$

Finally, letting  $R \rightarrow \infty$ , we get

$$\int_{\mathbb{H}^n} |u|^p dx = 0.$$

Let us consider the case  $l_1 = 0$  (or  $l_2 = 0$ ). By denoting

$$\bar{\Omega}_R := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : R^4 \leq |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\},$$

we have

$$\lim_{R \rightarrow \infty} \int_{\bar{\Omega}_R} |v|^q \varphi_R dx = 0, \tag{2.31}$$

and

$$\lim_{R \rightarrow \infty} \int_{\bar{\Omega}_R} |u|^p \psi_R dx = 0. \tag{2.32}$$

Let us denote by

$$\Theta_1 := \{y = (\bar{\xi}, \hat{\xi}, \tilde{\tau}) \in \mathbb{H}^n : 1 \leq |\bar{\xi}|^4 + |\hat{\xi}|^4 + \tilde{\tau}^2 \leq 2\}.$$

From (2.25), (2.27) and changing variables  $R\bar{\xi} = \xi$ ,  $R\hat{\xi} = \tilde{\xi}$  and  $R^2\tilde{\tau} = \tau$ , we establish

$$\begin{aligned}
\int_{\mathbb{H}^n} |v|^q \varphi_R dx &\leq C_0 \int_{\mathbb{H}^n} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\
&= C_0 \int_{\bar{\Omega}_R} |u| |\Delta_{\mathbb{H}} \varphi_R| dx \\
&\leq C_0 \left( \int_{\bar{\Omega}_R} |v|^q \varphi_R dx \right)^{\frac{1}{pq}} \left( \int_{\bar{\Omega}_R} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}} \psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{pq}} \\
&\quad \times \left( \int_{\bar{\Omega}_R} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \\
&\leq C_0 R^{l_1} \left( \int_{\bar{\Omega}_R} |v|^q \varphi_R dx \right)^{\frac{1}{pq}} \left( \int_{\Theta_1} [\Phi \circ \rho]^{-\frac{1}{q-1}} dy \right)^{\frac{q-1}{pq}} \\
&\quad \times \left( \int_{\Theta_1} [\Psi \circ \rho]^{-\frac{1}{p-1}} dy \right)^{\frac{p-1}{p}} \\
&\stackrel{l_1=0}{\leq} C \left( \int_{\bar{\Omega}_R} |v|^q \varphi_R dx \right)^{\frac{1}{pq}},
\end{aligned} \tag{2.33}$$

and

$$\begin{aligned}
\int_{\mathbb{H}^n} |u|^p \psi_R dx &\leq C \left( \int_{\bar{\Omega}_R} |u|^p \psi_R dx \right)^{\frac{1}{pq}} \left( \int_{\bar{\Omega}_R} \varphi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}} \psi_R|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{q}} \\
&\quad \times \left( \int_{\bar{\Omega}_R} \psi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx \right)^{\frac{p-1}{pq}} \\
&\leq C \left( \int_{\bar{\Omega}_R} |u|^p \psi_R dx \right)^{\frac{1}{pq}}.
\end{aligned} \tag{2.34}$$

Hence, we obtain

$$\int_{\mathbb{H}^n} |v|^q dx = \lim_{R \rightarrow \infty} \int_{\mathbb{H}^n} |v|^q \varphi_R dx \leq C \lim_{R \rightarrow \infty} \left( \int_{\bar{\Omega}_R} |v|^q \varphi_R dx \right)^{\frac{1}{pq}} = 0, \tag{2.35}$$

and

$$\int_{\mathbb{H}^n} |u|^p dx = \lim_{R \rightarrow \infty} \int_{\mathbb{H}^n} |u|^p \psi_R dx \leq C \lim_{R \rightarrow \infty} \left( \int_{\bar{\Omega}_R} |u|^p \psi_R dx \right)^{\frac{1}{pq}} = 0, \tag{2.36}$$

ending the proof of the theorem.  $\square$

**Corollary 2.5.** *Let  $p = q > 1$ ,  $u = v$  and  $M_1 = M_2$ . Then we arrive at the result of Theorem 2.2.*

*Proof.* It is easy to see that

$$Q \left( \frac{p^2 - 1}{p^2} \right) = Q \left( 1 - \frac{1}{p^2} \right) \leq 2 \left( 1 + \frac{1}{p} \right) = 2 \left( \frac{p+1}{p} \right),$$

showing

$$p \leq \frac{Q}{Q-2}.$$

The proof is finished.  $\square$

### 3. KIRCHHOFF TYPE PARABOLIC EQUATIONS AND SYSTEMS

In this section we prove the Liouville type theorems for the Kirchhoff type parabolic equations and systems on the Heisenberg group  $\mathbb{H}^n$ .

**3.1. Parabolic type equations.** Let us consider the following Kirchhoff type heat equation on  $\mathbb{H}^n$ , that is,

$$\begin{cases} u_t - M\left(t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx\right) \Delta_{\mathbb{H}} u = |u|^p, \\ u(x, 0) = u_0(x) \geq 0, \quad x \in \mathbb{H}^n, \end{cases} \quad (3.1)$$

where  $(x, t) \in \mathbb{H}^n \times (0, T)$ ,  $T > 0$ ,  $p > 1$ ,  $M : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  is satisfying the following condition

$$0 < M(t, \cdot, \cdot) \leq C_1 t^\beta, \quad \forall t > 0, \quad 0 \leq \beta < 1. \quad (3.2)$$

Let us formulate the following definition of the weak solution of the equation (3.1).

**Definition 3.1.** We say that  $u$  is a weak solution to (3.1) in  $\Omega = \mathbb{H}^n \times (0, T)$  with initial data  $0 \leq u_0 \in L^1(\mathbb{H}^n)$ , if  $u \in C^1(0, T; S_2^2(\mathbb{H}^n)) \cap L_{loc}^p(\Omega)$  and satisfies

$$\begin{aligned} \int_{\Omega} |u|^p \varphi dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi(x, 0) dx = - \int_{\Omega} u \varphi_t dx dt \\ - \int_{\Omega} M\left(t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx\right) u \Delta_{\mathbb{H}} \varphi dx dt, \end{aligned} \quad (3.3)$$

for any test function  $0 \leq \varphi \in C_{x,t}^{2,1}(\Omega)$ .

Now we give one of the main results of this section.

**Theorem 3.2.** *Suppose that  $M(\cdot, \cdot, \cdot)$  satisfies the condition (3.2). If*

$$1 < p < p_c = 1 + \frac{2-2\beta}{Q+2\beta}, \quad (3.4)$$

*with  $Q = 2n + 2$  the homogeneous dimension of  $\mathbb{H}^n$ , then the problem (3.1) admits no global weak non-trivial solution.*

*Proof.* Let us choose

$$\varphi_R(x, t) = \Phi\left(\frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2}{R^4}\right), \quad x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n, \quad t > 0, \quad R > 0, \quad (3.5)$$

with  $\Phi \in C_0^\infty(\mathbb{R}_+)$  and the following property

$$\Phi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2. \end{cases}$$

We note that  $\text{supp}(\varphi_R)$  and  $\text{supp}(\Delta_{\mathbb{H}}\varphi_R)$  are subsets of  $\Omega_1$ , where

$$\Omega_1 := \{(x, t) = (\xi, \tilde{\xi}, \tau, t) \in \mathbb{H}^n \times \mathbb{R}_+ : |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2 \leq 2R^4\}. \quad (3.6)$$

We seek the test function  $\varphi_R$  with the following properties

$$\int_{\Omega_1} |(\varphi_R(x, t))_t|^{p'} |\varphi_R(x, t)|^{-\frac{p'}{p}} dxdt < \infty,$$

and

$$\int_{\Omega_1} t^{\beta p'} |\Delta_{\mathbb{H}}\varphi_R(x, t)|^{p'} |\varphi_R(x, t)|^{-\frac{p'}{p}} dxdt < \infty.$$

Consider the following integral in the domain  $\Omega$ ,

$$\begin{aligned} & \int_{\Omega} \varphi_R(x, t) |u(x, t)|^p dxdt \leq \int_{\Omega} \varphi_R(x, t) |u(x, t)|^p dxdt + \int_{\mathbb{H}^n} u_0(x) \varphi_R(x, 0) dxdt \\ & = - \int_{\Omega} M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u(x, t) \Delta_{\mathbb{H}} \varphi_R(x, t) dxdt \\ & \quad - \int_{\Omega} u(x, t) (\varphi_R(x, t))_t dxdt \\ & \leq C_1 \int_{\Omega} t^{\beta} |u(x, t)| |\Delta_{\mathbb{H}} \varphi_R(x, t)| dxdt + \int_{\Omega} |u(x, t)| |(\varphi_R(x, t))_t| dxdt. \end{aligned}$$

Then we have

$$\int_{\Omega} \varphi_R(x, t) |u(x, t)|^p dxdt \leq C_1 \int_{\Omega} t^{\beta} |u(x, t)| |\Delta_{\mathbb{H}} \varphi_R(x, t)| dxdt + \int_{\Omega} |u(x, t)| |(\varphi_R(x, t))_t| dxdt.$$

By using  $\varepsilon$ -Young's inequality in the first integral of the right hand side, one obtains

$$\begin{aligned} & \int_{\Omega} t^{\beta} |u(x, t)| |\Delta_{\mathbb{H}} \varphi_R(x, t)| dxdt \\ & = \int_{\Omega_1} (\varphi_R(x, t))^{\frac{1}{p}} |u(x, t)| t^{\beta} |\Delta_{\mathbb{H}} \varphi(x, t)| (\varphi_R(x, t))^{-\frac{1}{p}} dxdt \quad (3.7) \\ & \leq \varepsilon \int_{\Omega} \varphi_R(x, t) |u(x, t)|^p dxdt + C(\varepsilon) \int_{\Omega_1} |t|^{\beta p'} |\Delta_{\mathbb{H}} \varphi_R(x, t)|^{p'} (\varphi_R(x, t))^{-\frac{p'}{p}} dxdt, \end{aligned}$$

and

$$\begin{aligned} \int_{\Omega} |u(x, t)| |(\varphi_R(x, t))_t| dxdt & \leq \varepsilon_1 \int_{\Omega} |u(x, t)|^p \varphi_R(x, t) dxdt \\ & \quad + C(\varepsilon_1) \int_{\Omega_1} |(\varphi_R(x, t))_t|^{p'} |\varphi_R(x, t)|^{-\frac{p'}{p}} dxdt. \quad (3.8) \end{aligned}$$

Then from these facts and choosing  $C_1\varepsilon + \varepsilon_1 < 1$ , we have

$$\begin{aligned} & \int_{\Omega} |u(x, t)|^p \varphi_R(x, t) dxdt \\ & \leq C \left( \int_{\Omega_1} (t^{\beta p'} |\Delta_{\mathbb{H}} \varphi_R(x, t)|^{p'} + |(\varphi_R(x, t))_t|^{p'}) (\varphi_R(x, t))^{-\frac{p'}{p}} dxdt \right). \quad (3.9) \end{aligned}$$

We note that from [31] we have

$$|\Delta_{\mathbb{H}} \varphi_R| \leq CR^{-2},$$

and

$$\left| \frac{\partial \varphi_R}{\partial t} \right| \leq CR^{-2}.$$

Let us denote

$$\Theta := \{(y, t_1) = (\bar{\xi}, \widehat{\xi}, \tilde{\tau}, t_1) \in \mathbb{H}^n \times \mathbb{R}_+ : |\bar{\xi}|^4 + |\widehat{\xi}|^4 + \tilde{\tau}^2 + t_1^2 \leq 2\},$$

and

$$\mu := |\bar{\xi}|^4 + |\widehat{\xi}|^4 + \tilde{\tau}^2 + t_1^2.$$

Since the homogeneous dimension of  $\mathbb{H}^n$  is equal to  $Q$ , and by changing variables  $R^2 \tilde{t} = t$ ,  $R\bar{\xi} = \xi$ ,  $R\widehat{\xi} = \tilde{\xi}$  and  $R^2 \tilde{\tau} = \tau$  in  $\Omega_1$ , we calculate

$$\begin{aligned} \int_{\Omega_1} |\varphi_R(x, t)|^{-\frac{1}{p-1}} \left| \frac{\partial \varphi_R(x, t)}{\partial t} \right|^{\frac{p}{p-1}} dx dt &= R^{Q+2-\frac{2p}{p-1}} \int_{\Theta} |(\Phi \circ \mu)|^{-\frac{p}{p-1}} \left| \frac{\partial(\Phi \circ \mu)}{\partial t} \right|^{\frac{p}{p-1}} dy dt_1 \\ &\leq CR^{Q+2-\frac{2p}{p-1}}, \end{aligned} \quad (3.10)$$

and

$$\begin{aligned} \int_{\Omega_1} t^{\beta p'} |\varphi_R(x, t)|^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R(x, t)|^{\frac{p}{p-1}} dx dt \\ = R^{\frac{2\beta p}{p-1} - \frac{2p}{p-1} + Q+2} \int_{\Theta} t_1^{\beta p'} |(\Phi \circ \mu)|^{-\frac{p}{p-1}} |\Delta_{\mathbb{H}}(\Phi \circ \mu)|^{\frac{p}{p-1}} dy dt_1 \\ \leq CR^{\frac{2\beta p}{p-1} - \frac{2p}{p-1} + Q+2}. \end{aligned} \quad (3.11)$$

By using (3.10)–(3.11) in (3.9), one obtains

$$\begin{aligned} \int_{\Omega} |u|^p \varphi_R dx dt &\leq C \left( R^{Q+2-\frac{2p}{p-1}} + R^{Q+2-\frac{2p}{p-1}+\frac{2\beta p}{p-1}} \right) \\ &\leq CR^{Q+2-\frac{2p}{p-1}+\frac{2\beta p}{p-1}}. \end{aligned} \quad (3.12)$$

Here, we choose

$$Q + 2 - \frac{2p}{p-1} + \frac{2\beta p}{p-1} < 0,$$

that is,

$$p < 1 + \frac{2-2\beta}{Q+2\beta}.$$

Letting  $R \rightarrow \infty$  with  $p < 1 + \frac{2-2\beta}{Q+2\beta}$ , one has

$$\int_{\Omega} |u(x, t)|^p dx dt \leq 0, \quad (3.13)$$

arriving at a contradiction.  $\square$

**3.2. Parabolic type systems.** Now, we will show a Liouville type result for the following system of Kirchhoff type parabolic equations in  $\Omega := \mathbb{H}^n \times (0, T)$ ,  $T > 0$

$$\begin{cases} u_t - M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) \Delta_{\mathbb{H}} u = |\omega|^p, \\ \omega_t - M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) \Delta_{\mathbb{H}} \omega = |u|^q, \\ u(x, 0) = u_0(x) \geq 0, \quad x \in \mathbb{H}^n, \\ \omega(x, 0) = \omega_0(x) \geq 0, \quad x \in \mathbb{H}^n, \end{cases} \quad (3.14)$$

where  $p, q > 1$  and  $M_1, M_2 : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  are bounded functions such that

$$0 < M_1(t, \cdot, \cdot, \cdot, \cdot) \leq C_1 t^{\beta_1}, \quad \forall t \in (0, \infty), \quad 0 \leq \beta_1 < 1, \quad (3.15)$$

and

$$0 < M_2(t, \cdot, \cdot, \cdot, \cdot) \leq C_2 t^{\beta_2}, \quad \forall t \in (0, \infty), \quad 0 \leq \beta_2 < 1. \quad (3.16)$$

Let us give a definition of the weak solution to (3.14) as follows.

**Definition 3.3.** We say that the pair  $u \in C^1(0, T; S_2^2(\mathbb{H}^n)) \cap L^p(\Omega)$  and  $\omega \in C^1(0, T; S_2^2(\mathbb{H}^n)) \cap L^q(\Omega)$  with  $p, q > 1$  is a weak solution of the system (3.14) on  $\Omega = \mathbb{H}^n \times (0, T)$  with the Cauchy data  $(u_0, \omega_0) \in L_{loc}^1(\mathbb{H}^n) \times L_{loc}^1(\mathbb{H}^n)$  with  $u_0, \omega_0 \geq 0$  on  $\mathbb{H}^n$ , if the following identities

$$\begin{aligned} & \int_{\Omega} |\omega|^p \varphi dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi(x, 0) dx = - \int_{\Omega} u \varphi_t dx dt \\ & - \int_{\Omega} M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx dt, \end{aligned}$$

and

$$\begin{aligned} & \int_{\Omega} |u|^q \psi dx dt + \int_{\mathbb{H}^n} \omega_0(x) \psi(x, 0) dx = - \int_{\Omega} \omega \psi_t dx dt \\ & - \int_{\Omega} M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) \omega \Delta_{\mathbb{H}} \psi dx dt, \end{aligned}$$

hold for any test functions  $0 \leq \psi, \varphi \in C_{x,t}^{2,1}(\Omega)$ .

**Theorem 3.4.** Let  $p, q > 1$  and  $Q < \max\{A_1, A_2\}$ , where  $Q$  is the homogeneous dimension of  $\mathbb{H}^n$ ,

$$A_1 = \frac{\frac{1}{q} - \beta_1 - \frac{\beta_2}{q} + \frac{1}{pq}}{\frac{1}{2qp'} + \frac{1}{2q'}}, \quad (3.17)$$

and

$$A_2 = \frac{\frac{1}{p} - \beta_2 - \frac{\beta_1}{p} + \frac{1}{pq}}{\frac{1}{2q'p} + \frac{1}{2p'}}, \quad (3.18)$$

where  $\beta_1, \beta_2 \in [0, 1)$ . Then the system (3.14) does not admit a nontrivial weak solution.



*Proof.* Firstly, let us define the test functions

$$\varphi_R = \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2}{R^4} \right), \quad x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n, \quad t > 0, \quad R > 0, \quad (3.19)$$

and

$$\psi_R = \Psi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2}{R^4} \right), \quad x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n, \quad t > 0, \quad R > 0, \quad (3.20)$$

with  $\Psi, \Phi \in C_0^\infty(\mathbb{R}_+)$  and the following property

$$\Psi(r), \Phi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2, \end{cases}$$

with  $\text{supp } \Psi = \text{supp } \Phi = [0, 2]$ . We note that  $\text{supp}(\psi_R)$ ,  $\text{supp}(\varphi_R)$ ,  $\text{supp}(\Delta_{\mathbb{H}}\psi_R)$  and  $\text{supp}(\Delta_{\mathbb{H}}\varphi_R)$  are subsets of  $\Omega_1$ , where

$$\Omega_1 := \{(x, t) = (\xi, \tilde{\xi}, \tau, t) \in \mathbb{H}^n \times \mathbb{R}_+ : |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2 \leq 2R^4\}. \quad (3.21)$$

Assume that there exists a weak solution to (3.14), and let  $M_1$  and  $M_2$  satisfies the conditions (3.15) and (3.16), then

$$\begin{aligned} & \int_{\Omega} |\omega|^p \varphi_R dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi_R(x, 0) dx \\ & \leq \int_{\Omega} |u| |(\varphi_R)_t| dx dt + C_1 \int_{\Omega} t^{\beta_1} |u| |\Delta_{\mathbb{H}} \varphi_R| dx dt \\ & = \int_{\Omega_1} |u| |(\varphi_R)_t| dx dt + C_1 \int_{\Omega_1} t^{\beta_1} |u| |\Delta_{\mathbb{H}} \varphi_R| dx dt, \end{aligned} \quad (3.22)$$

and

$$\begin{aligned} & \int_{\Omega} |u|^q \psi_R dx dt + \int_{\mathbb{H}^n} \omega_0(x) \psi_R(x, 0) dx \\ & \leq \int_{\Omega} |\omega| |(\psi_R)_t| dx dt + C_2 \int_{\Omega} t^{\beta_2} |\omega| |\Delta_{\mathbb{H}} \psi_R| dx dt \\ & = \int_{\Omega_1} |\omega| |(\psi_R)_t| dx dt + C_2 \int_{\Omega_1} t^{\beta_2} |\omega| |\Delta_{\mathbb{H}} \psi_R| dx dt. \end{aligned}$$

By using Hölder's inequality, we have

$$\begin{aligned} \int_{\Omega_1} |u| |(\varphi_R)_t| dx dt &= \int_{\Omega_1} |u| \psi_R^{\frac{1}{q}} |(\varphi_R)_t| \psi_R^{-\frac{1}{q}} dx dt \\ &\leq \left( \int_{\Omega_1} |u|^q \psi_R dx dt \right)^{\frac{1}{q}} \left( \int_{\Omega_1} |(\varphi_R)_t|^{q'} \psi_R^{-\frac{q'}{q}} dx dt \right)^{\frac{1}{q'}} \\ &\leq \left( \int_{\Omega} |u|^q \psi_R dx dt \right)^{\frac{1}{q}} \left( \int_{\Omega_1} |(\varphi_R)_t|^{q'} \psi_R^{-\frac{q'}{q}} dx dt \right)^{\frac{1}{q'}}, \end{aligned}$$

and

$$\int_{\Omega_1} t^{\beta_1} |u| |\Delta_{\mathbb{H}} \varphi_R| dx dt \leq \left( \int_{\Omega_1} |u|^q \psi_R dx dt \right)^{\frac{1}{q}} \left( \int_{\Omega_1} t^{\beta_1 q'} |\Delta_{\mathbb{H}} \varphi_R|^{q'} \psi_R^{-\frac{q'}{q}} dx dt \right)^{\frac{1}{q'}}.$$

By using (3.22) and the previous estimates, we obtain

$$\begin{aligned} \int_{\Omega} |\omega|^p \varphi_R dx dt &\leq \int_{\Omega} |\omega|^p \varphi_R dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi_R(x, 0) dx dt \\ &\leq \int_{\Omega} |u| (\varphi_R)_t dx dt + \int_{\Omega} t^{\beta_1} |u| |\Delta_{\mathbb{H}} \varphi_R| dx dt \\ &\leq \left( \int_{\Omega} |u|^q \psi_R dx dt \right)^{\frac{1}{q}} I_1, \end{aligned} \quad (3.23)$$

where

$$I_1 = \left( \int_{\Omega_1} t^{\beta_1 q'} |\Delta_{\mathbb{H}} \varphi_R|^{q'} \psi_R^{-\frac{q'}{q}} dx dt \right)^{\frac{1}{q'}} + \left( \int_{\Omega_1} |(\varphi_R)_t|^{q'} \psi_R^{-\frac{q'}{q}} dx dt \right)^{\frac{1}{q'}}. \quad (3.24)$$

Analogous calculations show that

$$\int_{\Omega} |u|^q \psi_R dx dt \leq \left( \int_{\Omega} |\omega|^p \varphi_R dx dt \right)^{\frac{1}{p}} I_2, \quad (3.25)$$

where

$$I_2 = \left( \int_{\Omega_1} t^{\beta_2 p'} |\Delta_{\mathbb{H}} \psi_R|^{p'} \varphi_R^{-\frac{p'}{p}} dx dt \right)^{\frac{1}{p'}} + \left( \int_{\Omega_1} |(\psi_R)_t|^{p'} \varphi_R^{-\frac{p'}{p}} dx dt \right)^{\frac{1}{p'}}.$$

By using (3.23) and (3.25), one obtains

$$\int_{\Omega} |\omega|^p \varphi_R dx dt \leq \left( \int_{\Omega} |u|^q \psi_R dx dt \right)^{\frac{1}{q}} I_2 \leq \left( \int_{\Omega} |\omega|^p \varphi_R dx dt \right)^{\frac{1}{pq}} I_2^{\frac{1}{q}} I_1,$$

implying

$$\left( \int_{\Omega} |\omega(x, t)|^p \varphi_R(x, t) dx dt \right)^{1 - \frac{1}{pq}} \leq I_1 I_2^{\frac{1}{q}}.$$

Similarly, we have

$$\left( \int_{\Omega} |u(x, t)|^q \psi_R(x, t) dx dt \right)^{1 - \frac{1}{pq}} \leq I_1^{\frac{1}{p}} I_2.$$

Let us denote

$$\Theta := \{(y, t_1) = (\bar{\xi}, \widehat{\xi}, \tilde{\tau}, t_1) \in \mathbb{H}^n \times \mathbb{R}_+ : 0 \leq |\bar{\xi}|^4 + |\widehat{\xi}|^4 + \tilde{\tau}^2 + t_1^2 \leq 2\},$$

and

$$\mu := |\bar{\xi}|^4 + |\widehat{\xi}|^4 + \tilde{\tau}^2 + t_1^2.$$

By using the homogeneous dimension  $Q$  of  $\mathbb{H}^n$  and by choosing variables  $R^2 t_1 = t$ ,  $R \bar{\xi} = \xi$ ,  $R \widehat{\xi} = \widehat{\xi}$  and  $R^2 \tilde{\tau} = \tau$  in the domain  $\Omega_1$ , one calculates

$$\begin{aligned} \int_{\Omega_1} |(\psi_R)_t|^{p'} \varphi_R^{-\frac{p'}{p}} dx dt &= R^{-2p'+2+Q} \int_{\Theta} |(\Psi \circ \mu)_{t_1}|^{p'} (\Phi \circ \mu)^{-\frac{p'}{p}} dy dt_1 \\ &\leq C R^{-2p'+2+Q}, \end{aligned}$$

similarly, we get

$$\int_{\Omega_1} |(\varphi_R)_t|^{q'} \psi_R^{-\frac{q'}{q}} dxdt \leq CR^{-2q'+2+Q},$$

$$\int_{\Omega_1} t^{\beta_1 q'} |\Delta_{\mathbb{H}} \varphi_R|^{q'} \psi_R^{-\frac{q'}{q}} dxdt \leq CR^{-2q'+2+Q+2\beta_1 q'},$$

and

$$\int_{\Omega_1} t^{\beta_2 p'} |\Delta_{\mathbb{H}} \psi_R|^{p'} \varphi_R^{-\frac{p'}{p}} dxdt \leq CR^{-2p'+2+Q+2\beta_2 p'}.$$

By using the last estimates in (3.24), we obtain

$$I_1 \leq CR^{-2+\frac{2+Q}{q'}+2\beta_1}.$$

Analogously, with  $R^2 t_1 = t$ ,  $R\bar{\xi} = \xi$ ,  $R\hat{\xi} = \tilde{\xi}$  and  $R^2 \tilde{\tau} = \tau$  in the domain  $\Omega_1$ , we get

$$I_2 \leq CR^{-2+\frac{2+Q}{p'}+2\beta_2}. \quad (3.26)$$

By using these facts, one has

$$\left( \int_{\Omega} |\omega(x, t)|^p \varphi_R(x, t) dxdt \right)^{1-\frac{1}{pq}} \leq I_1 I_2^{\frac{1}{q}} \leq CR^{\theta_1 + \frac{\theta_2}{q}}, \quad (3.27)$$

where  $\theta_1 = -2 + \frac{2+Q}{q'} + 2\beta_1$  and  $\theta_2 = -2 + \frac{2+Q}{p'} + 2\beta_2$ .

Now we show

$$\theta_1 + \frac{\theta_2}{q} < 0.$$

By calculating, we have

$$-2 + \frac{2+Q}{q'} + 2\beta_1 - \frac{2}{q} + \frac{2\beta_2}{q} + \frac{2+Q}{p'q} < 0.$$

Thus, we can require

$$Q < \frac{\frac{1}{q} - \beta_1 - \frac{\beta_2}{q} + \frac{1}{pq}}{\frac{1}{2qp'} + \frac{1}{2q'}}. \quad (3.28)$$

Repeating the arguments above, we obtain

$$\left( \int_{\Omega} |u(x, t)|^q \psi_R(x, t) dxdt \right)^{1-\frac{1}{pq}} \leq I_1^{\frac{1}{p}} I_2 \leq CR^{\theta_2 + \frac{\theta_1}{p}}, \quad (3.29)$$

and we can require

$$Q < \frac{\frac{1}{p} - \beta_2 - \frac{\beta_1}{p} + \frac{1}{pq}}{\frac{1}{2q'p} + \frac{1}{2p'}}. \quad (3.30)$$

Finally, letting  $R \rightarrow \infty$  we arrive at a contradiction.  $\square$

**Corollary 3.5.** *Let  $p = q$ ,  $u = \omega$ ,  $\beta_1 = \beta_2$  and  $M_1 = M_2$ . Then we get the results of Theorem 3.2.*

*Proof.* By simple calculations we have

$$Q < \frac{\frac{1}{p} - \beta_1 - \frac{\beta_1}{p} + \frac{1}{p^2}}{\frac{1}{2p'} \left( \frac{1}{p} + 1 \right)} = \frac{2 - 2p\beta_1}{p - 1},$$

which means

$$p < 1 + \frac{2 - 2\beta_1}{Q + 2\beta_1},$$

implying the statement.  $\square$

**3.3. Pseudo-parabolic type equations and systems.** In this subsection we show nonexistence results for pseudo-parabolic equations and systems on the Heisenberg groups. We consider the following Cauchy problem for the Kirchhoff type pseudo-parabolic equation

$$\begin{cases} u_t - M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) \Delta_{\mathbb{H}} u - \Delta_{\mathbb{H}} u_t = |u|^p, \\ u(x, 0) = u_0(x) \geq 0, \quad x \in \mathbb{H}^n, \end{cases} \quad (3.31)$$

where  $(x, t) \in \Omega := \mathbb{H}^n \times (0, T)$ ,  $T > 0$ ,  $p > 1$ ,  $M : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  is a bounded function such that

$$0 < M(\cdot, \cdot, \cdot) \leq C_0. \quad (3.32)$$

We give a definition of the weak solution of the equation (3.31).

**Definition 3.6.** We say that  $u \in C^1(0, T; S_2^2(\mathbb{H}^n)) \cap L^p(\Omega)$  with  $p > 1$  is a weak solution to (3.31) on  $\Omega$  with the initial data  $0 \leq u_0(x) \in L_{loc}^1(\mathbb{H}^n)$ , if  $u \in L_{loc}^p(\Omega)$  and satisfies

$$\begin{aligned} \int_{\Omega} |u|^p \varphi dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi(x, 0) dx &= - \int_{\Omega} u \varphi_t dx dt \\ &- \int_{\Omega} M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx dt \\ &+ \int_{\Omega} u \Delta_{\mathbb{H}} \varphi_t dx dt + \int_{\mathbb{H}^n} u_0(x) \Delta_{\mathbb{H}} \varphi(x, 0) dx, \end{aligned} \quad (3.33)$$

for any test function  $0 \leq \varphi \in C_{x,t}^{2,1}(\Omega)$ .

Let us formulate one of the main results of this section.

**Theorem 3.7.** *Suppose that  $M(\cdot, \cdot, \cdot)$  satisfies the condition (3.32). If the rate  $p$  is from the following interval*

$$1 < p \leq p_c = 1 + \frac{2}{Q}, \quad (3.34)$$

*with the homogeneous dimension  $Q$  of  $\mathbb{H}^n$ , then the problem (3.31) admits no global weak solution.*

*Proof.* Let us give a short proof of Theorem 3.7. Here, we choose the test functions as follows

$$\varphi_R(x, t) = \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2}{R^4} \right), \quad x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n, \quad t > 0, \quad R > 0, \quad (3.35)$$

with  $\Phi \in C_0^\infty(\mathbb{R}_+)$  satisfying

$$\Phi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2. \end{cases}$$

Note that  $\text{supp}(\varphi_R)$  and  $\text{supp}(\Delta_{\mathbb{H}}\varphi_R)$  are subsets of  $\Omega_1$ , where

$$\Omega_1 := \{(x, t) \in \mathbb{H}^n \times \mathbb{R}_+ : |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 + t^2 \leq 2R^4\}. \quad (3.36)$$

By using (3.32) and Young's inequality, from the identity (3.33) we obtain

$$\int_{\Omega} |u|^p \varphi_R dx dt \leq C \left( A_p(\varphi_R) + B_p(\varphi_R) + \int_{\mathbb{H}^n} u_0 |\Delta_{\mathbb{H}}\varphi_R(x, 0)| dx \right), \quad (3.37)$$

where

$$\begin{aligned} A_p(\varphi_R) &= \int_{\Omega_1} |\varphi_R|^{-\frac{1}{p-1}} (|(\varphi_R)_t|^{\frac{p}{p-1}} + |\Delta_{\mathbb{H}}\varphi_R|^{\frac{p}{p-1}}) dx dt, \\ B_p(\varphi_R) &= \int_{\Omega_1} |\varphi_R|^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}(\varphi_R)_t|^{\frac{p}{p-1}} dx dt. \end{aligned} \quad (3.38)$$

Then we have

$$\left| \frac{\partial \varphi_R}{\partial t} \right| \leq CR^{-2},$$

and

$$|\Delta_{\mathbb{H}}(\varphi_R)_t| \leq CR^{-4}.$$

Now, by using property of homogeneous dimension of  $\mathbb{H}^n$  and by choosing variables  $R^2\bar{t} = t$ ,  $R\bar{\xi} = \xi$ ,  $R\tilde{\xi} = \tilde{\xi}$  and  $R^2\bar{\tau} = \tau$  in the domain  $\Omega_1$ , from (3.38) we obtain

$$\begin{aligned} |A_p(\varphi_R)| &\leq CR^{Q+2-\frac{2p}{p-1}}, \\ |B_p(\varphi_R)| &\leq CR^{Q+2-\frac{4p}{p-1}}. \end{aligned} \quad (3.39)$$

Thus, we conclude

$$\begin{aligned} \int_{\Omega} |u|^p \varphi_R dx dt &\leq \int_{\Omega} |u|^p \varphi_R dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi_R(x, 0) dx \\ &\leq C \left( R^{Q+2-\frac{2p}{p-1}} + R^{Q+2-\frac{4p}{p-1}} + \int_{\mathbb{H}^n} u_0 |\Delta_{\mathbb{H}}\varphi_R(x, 0)| dx \right) \\ &\leq C \left( R^{Q+2-\frac{2p}{p-1}} + \int_{\Omega_1} u_0 |\Delta_{\mathbb{H}}\varphi_R(x, 0)| dx \right) \\ &\leq CR^{Q+2-\frac{2p}{p-1}} + CR^{-2}. \end{aligned}$$

Finally, by taking  $p < 1 + \frac{2}{Q}$  and letting  $R \rightarrow \infty$ , one concludes

$$\int_{\Omega} |u|^p dx dt \leq 0,$$

that is,  $u = 0$ . This is a contradiction.

The case  $p = 1 + \frac{2}{Q}$  can be dealt with Hölder's inequality instead of Young's inequality in (3.37).  $\square$

As the last problem of this section, we consider the system of the Kirchhoff type pseudo-parabolic equations

$$\left\{ \begin{array}{l} u_t - M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) \Delta_{\mathbb{H}} u \\ \quad - \Delta_{\mathbb{H}} u_t = |\omega|^q, \quad (x, t) \in \Omega, \\ \omega_t - M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) \Delta_{\mathbb{H}} \omega \\ \quad - \Delta_{\mathbb{H}} \omega_t = |u|^p, \quad (x, t) \in \Omega, \\ u(x, 0) = u_0(x) \geq 0, \quad x \in \mathbb{H}^n, \\ \omega(x, 0) = \omega_0(x) \geq 0, \quad x \in \mathbb{H}^n, \end{array} \right. \quad (3.40)$$

where  $\Omega := \mathbb{H}^n \times (0, T)$ ,  $T > 0$ ,  $p, q > 1$  and  $M_1, M_2 : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  are bounded functions such that

$$0 < M_1(\cdot, \cdot, \cdot, \cdot, \cdot) \leq C_1, \quad (3.41)$$

and

$$0 < M_2(\cdot, \cdot, \cdot, \cdot, \cdot) \leq C_2. \quad (3.42)$$

We formulate a definition of the weak solution to (3.40) in the following form.

**Definition 3.8.** We say that the pair of functions  $(u, \omega) \in C^1(0, T; S_2^2(\mathbb{H}^n)) \cap L^p(\Omega) \times C^1(0, T; S_2^2(\mathbb{H}^n)) \cap L^q(\Omega)$  with  $p, q > 1$  is a weak solution of the Cauchy problem (3.40) on  $\Omega$  with the initial data  $(u(x, 0), \omega(x, 0)) = (u_0, \omega_0) \in L_{loc}^1(\mathbb{H}^n) \times L_{loc}^1(\mathbb{H}^n)$  and  $u_0, \omega_0 \geq 0$ , if

$$\begin{aligned} & \int_{\Omega} |\omega|^q \varphi dx dt + \int_{\mathbb{H}^n} u_0(x) \varphi(x, 0) dx = - \int_{\Omega} u \varphi_t dx dt + \int_{\mathbb{H}^n} u_0(x) \Delta_{\mathbb{H}} \varphi(x, 0) dx \\ & - \int_{\Omega} M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx dt \\ & + \int_{\Omega} u \Delta_{\mathbb{H}} \varphi_t dx dt, \end{aligned}$$

and

$$\begin{aligned} & \int_{\Omega} |u|^p \psi dx dt + \int_{\mathbb{H}^n} \omega_0(x) \psi(x, 0) dx = - \int_{\Omega} \omega \psi_t dx dt + \int_{\mathbb{H}^n} \omega_0(x) \Delta_{\mathbb{H}} \psi(x, 0) dx \\ & - \int_{\Omega} M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} \omega|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} \omega|^2 dx \right) \omega \Delta_{\mathbb{H}} \psi dx dt \\ & + \int_{\Omega} \omega \Delta_{\mathbb{H}} \psi_t dx dt, \end{aligned}$$

for any test functions  $0 \leq \psi, \varphi \in C_{x,t}^{2,1}(\Omega)$ .

**Theorem 3.9.** *Assume that  $p, q > 1$ . Let  $Q$  be a homogeneous dimension of  $\mathbb{H}^n$  such that  $Q \leq \max\{A_1, A_2\}$ , where*

$$A_1 = \frac{\frac{1}{pq} + \frac{1}{q}}{\frac{1}{2qp'} + \frac{1}{2q'}} \quad \text{and} \quad A_2 = \frac{\frac{1}{pq} + \frac{1}{p}}{\frac{1}{2q'p} + \frac{1}{2p'}}. \quad (3.43)$$

Then the system (3.40) does not admit nontrivial weak solution.

*Proof.* Repeating the proofs similar to those of Theorems 3.4 and 3.7, we conclude this theorem.  $\square$

#### 4. KIRCHHOFF TYPE HYPERBOLIC EQUATIONS AND SYSTEMS

In this section we show a nonexistence result for the Kirchhoff type hyperbolic equations and systems on the Heisenberg group.

**4.1. Hyperbolic type equations.** We consider the equation

$$\begin{cases} u_{tt} - M\left(t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx\right) \Delta_{\mathbb{H}} u = |u|^p, & (x, t) \in \Omega, \\ u(x, 0) = u_0(x), & x \in \mathbb{H}^n, \\ u_t(x, 0) = u_1(x), & x \in \mathbb{H}^n, \end{cases} \quad (4.1)$$

where  $\Omega := \mathbb{H}^n \times (0, T)$ ,  $T > 0$ ,  $p > 1$  and  $M : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  is satisfying the following condition

$$0 < M(t, \cdot, \cdot) \leq C_0 t^\beta, \quad \forall t > 0, \quad \beta > -2, \quad \text{and} \quad Q > \frac{2}{2 + \beta}. \quad (4.2)$$

Let us formulate a definition of the weak solution of the Cauchy problem (4.1).

**Definition 4.1.** We say that  $u$  is a weak solution to (4.1) on  $\Omega$  with the initial data  $u_1, u_0 \in L^1_{loc}(\mathbb{H}^n)$ , if  $u \in C^2(0, T; S^2_2(\Omega)) \cap L^p(\Omega)$  and it satisfies

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi dx dt + \int_{\mathbb{H}^n} (u_1(x) \varphi(x, 0) - u_0(x) \varphi_t(x, 0)) dx \\ &= \int_{\Omega} u \varphi_{tt} dx dt - \int_{\Omega} M\left(t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx\right) u \Delta_{\mathbb{H}} \varphi dx dt, \end{aligned} \quad (4.3)$$

for any test function  $0 \leq \varphi \in C^{2,2}_{x,t}(\Omega)$ .

**Theorem 4.2.** *Let  $u_1, u_0 \in L^1(\mathbb{H}^n)$ , and assume that  $\int_{\mathbb{H}^n} u_1 dx \geq 0$ . Moreover, assume that the conditions in (4.2) hold. If*

$$1 < p \leq p_c = 1 + \frac{4}{(2 + \beta)Q - 2}, \quad (4.4)$$

with  $Q = 2n + 2$  the homogeneous dimension of  $\mathbb{H}^n$ , then there exists no nontrivial weak solution to the equation (4.1).

*Proof.* Let us consider the case  $1 < p < p_c$ . By taking into account the property (4.2) and by Definition 4.1, for any test function  $\varphi$ , we have

$$\begin{aligned}
& \int_{\Omega} |u|^p \varphi dx dt + \int_{\mathbb{H}^n} (u_1(x)\varphi(x, 0) - u_0(x)\varphi_t(x, 0)) dx \\
&= \int_{\Omega} u \varphi_{tt} dx dt - \int_{\Omega} M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx dt \\
&\leq \left| \int_{\Omega} u \varphi_{tt} dx dt - \int_{\Omega} M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx dt \right| \\
&\leq \int_{\Omega} |u \varphi_{tt}| dx dt + \int_{\Omega} \left| M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) \right| |u| |\Delta_{\mathbb{H}} \varphi| dx dt \\
&\leq \int_{\Omega} |u \varphi_{tt}| dx dt + C_0 \int_{\Omega} t^{\beta} |u| |\Delta_{\mathbb{H}} \varphi| dx dt.
\end{aligned} \tag{4.5}$$

Let us choose a test function in the following form

$$\varphi_R(x, t) = \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right) \Phi \left( \frac{t^2}{R^{\theta}} \right), \quad (x, t) \in \mathbb{H}^n \times \mathbb{R}_+, \quad \theta > 0, \quad R > 0, \tag{4.6}$$

with  $\Phi \in C_0^{\infty}(\mathbb{R}_+)$  as

$$\Phi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2. \end{cases}$$

By noting that

$$\frac{\partial \varphi_R(x, t)}{\partial t} = \frac{2t}{R^{\theta}} \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right) \Phi'_t \left( \frac{t^2}{R^{\theta}} \right), \tag{4.7}$$

one has

$$\frac{\partial \varphi_R(x, 0)}{\partial t} = 0. \tag{4.8}$$

We note that  $\text{supp}(\varphi_R)$  and  $\text{supp}(\Delta_{\mathbb{H}} \varphi_R)$  are subsets of  $\Omega_1 = \Gamma \cup \tilde{\Gamma}$ , where

$$\Gamma := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\}, \quad \tilde{\Gamma} := \{t : 0 \leq t^2 \leq 2R^{\theta}\}. \tag{4.9}$$

We seek the test function  $\varphi_R$  with the following properties

$$A_p(\varphi_R) := \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |(\varphi_R)_{tt}|^{\frac{p}{p-1}} dx dt < \infty,$$

and

$$B_p(\varphi_R) := \int_{\Omega_1} t^{\frac{\beta p}{p-1}} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx dt < \infty.$$



Then by using Young's inequality, we get

$$\begin{aligned} \int_{\Omega_1} |u| |(\varphi_R)_{tt}| dx dt &= \int_{\Omega_1} u \varphi_R^{\frac{1}{p}} \varphi_R^{-\frac{1}{p}} |(\varphi_R)_{tt}| dx dt \\ &\leq \varepsilon \int_{\Omega_1} |u|^p \varphi_R dx dt + C(\varepsilon) \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |(\varphi_R)_{tt}|^{\frac{p}{p-1}} dx dt, \end{aligned}$$

and

$$\int_{\Omega_1} t^\beta |u| |\Delta_{\mathbb{H}} \varphi_R| dt dx \leq \varepsilon \int_{\Omega_1} |u|^p \varphi_R dt dx + C(\varepsilon) \int_{\Omega_1} t^{\frac{\beta p}{p-1}} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx dt,$$

for some positive constant  $C(\varepsilon)$ . Hence by using the last fact and (4.8), we obtain

$$\int_{\Omega} |u|^p \varphi_R dx dt + C_1 \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \leq C (A_p(\varphi_R) + B_p(\varphi_R)), \quad (4.10)$$

where  $C, C_1 > 0$ .

Then from [31], one obtains

$$\begin{aligned} \Delta_{\mathbb{H}} \varphi_R(x, t) &= \frac{4(n+4)}{R^4} \left( |\xi|^2 + |\tilde{\xi}|^2 \right) \Phi' \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right) \Phi \left( \frac{t^2}{R^2} \right) \\ &+ \frac{16}{R^8} \left( (|\xi|^6 + |\tilde{\xi}|^6) + 2\tau(|\xi|^2 - |\tilde{\xi}|^2)\xi \cdot \tilde{\xi} + \tau^2(|\xi|^2 + |\tilde{\xi}|^2) \right) \\ &\times \Phi'' \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right) \Phi \left( \frac{t^2}{R^2} \right). \end{aligned} \quad (4.11)$$

By choosing variables  $R^{\frac{\theta}{2}} \bar{t} = t$ ,  $R \bar{\xi} = \xi$ ,  $R \hat{\xi} = \tilde{\xi}$  and  $R^2 \bar{\tau} = \tau$  in the domain  $\Omega_1$ ,  $\theta = \frac{2}{1+\frac{\beta}{2}}$  and (4.8), one calculates

$$\begin{aligned} \int_{\Omega} |u|^p \varphi_R dx dt + C_1 \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ \stackrel{(4.10)}{\leq} C \left( R^{-\theta p' + Q + \frac{\theta}{2}} + R^{-2p' + (\frac{\beta\theta}{2})p' + Q + \frac{\theta}{2}} \right) \\ \stackrel{\theta = \frac{2}{1+\frac{\beta}{2}}}{=} 2C R^{-\theta p' + Q + \frac{\theta}{2}}. \end{aligned} \quad (4.12)$$

On the other hand, we have

$$\begin{aligned} \int_{\Omega} |u|^p \varphi_R dx dt + C_1 \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ \geq \liminf_{R \rightarrow \infty} \int_{\Omega} |u|^p \varphi_R dx dt + C_1 \liminf_{R \rightarrow \infty} \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx. \end{aligned}$$

Using the monotone convergence theorem, we obtain

$$\liminf_{R \rightarrow \infty} \int_{\Omega} |u|^p \varphi_R dx dt = \int_{\Omega} |u|^p dx dt.$$

Since  $u_1 \in L^1(\mathbb{H}^n)$ , by the dominated convergence theorem, one has

$$\liminf_{R \rightarrow \infty} \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx = \int_{\mathbb{H}^n} u_1(x) dx.$$

Now, we get

$$\liminf_{R \rightarrow \infty} \left( \int_{\Omega} |u|^p \varphi_R dx dt + C_1 \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \right) \geq \int_{\Omega} |u|^p dx dt + C_1 d,$$

where

$$d = \int_{\mathbb{H}^n} u_1(x) dx \geq 0.$$

Then,

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi_R dx dt + C_1 \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ & \geq \liminf_{R \rightarrow \infty} \left( \int_{\Omega} |u|^p \varphi_R dx dt + C_1 \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \right) \geq \int_{\Omega} |u|^p dx dt + C_1 d. \end{aligned}$$

From (4.12) and  $u_1 \in L^1(\mathbb{H}^n)$ , it can be shown that

$$\int_{\Omega} |u|^p dx dt + C_1 d \leq R^{-\theta p' + Q + \frac{\theta}{2}}. \quad (4.13)$$

By the assumption  $p < 1 + \frac{\theta}{Q - \frac{\theta}{2}} = p_c$ , and where we take  $\theta = \frac{2}{1 + \frac{\theta}{2}} > 0$ . Finally, letting  $R \rightarrow \infty$ , one has

$$\int_{\Omega} |u|^p dx dt \leq \int_{\Omega} |u|^p dx dt + C_1 d \leq 0.$$

That is a contradiction, concluding that

$$\int_{\Omega} |u|^p dx dt \leq 0.$$

Let us consider the critical case  $p = 1 + \frac{\theta}{Q - \frac{\theta}{2}}$ . By using inequality (4.13), we obtain

$$\int_{\Omega} |u|^p dx dt \leq C < \infty, \quad (4.14)$$

and

$$\lim_{R \rightarrow \infty} \int_{\bar{\Omega}_R} |u|^p \varphi_R dx dt = 0, \quad (4.15)$$

where  $\bar{\Omega}_R = \Gamma_R \cup \tilde{\Gamma}_R$ ,

$\Gamma_R := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : R^4 \leq |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\}$ ,  $\tilde{\Gamma}_R := \{t : R^\theta \leq t^2 \leq 2R^\theta\}$ .

Using the Hölder inequality instead of Young's inequality in (4.5), one has

$$\int_{\Omega} |u|^p \varphi_R dx dt \leq \int_{\Omega} |u|^p \varphi_R dx dt + d \leq C \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx dt \right)^{\frac{1}{p}}.$$

Letting  $R \rightarrow \infty$ , we get

$$\int_{\Omega} |u|^p \varphi_R dx dt = 0,$$

completing the proof.  $\square$

**4.2. Hyperbolic Kirchhoff type systems.** Let us consider the Kirchhoff type system of wave equations in  $\Omega = \mathbb{H}^n \times (0, T)$ ,  $T > 0$  with the Cauchy data

$$\begin{cases} u_{tt} - M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) \Delta_{\mathbb{H}} u = |v|^q, \\ v_{tt} - M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) \Delta_{\mathbb{H}} v = |u|^p, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), \quad x \in \mathbb{H}^n, \\ v(x, 0) = v_0(x), v_t(x, 0) = v_1(x), \quad x \in \mathbb{H}^n, \end{cases} \quad (4.16)$$

where  $p, q > 1$  and  $M_1, M_2 : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  are satisfying the following conditions

$$0 < M_i(t, \cdot, \cdot, \cdot, \cdot) \leq C_i t^{\beta_i}, \quad \forall t > 0, \quad \beta_i > -2, \quad \text{and } Q > \frac{4}{2 + \beta_i}, \quad (4.17)$$

for  $i = 1, 2$ .

Now, we formulate a definition of the weak solution to (4.16).

**Definition 4.3.** We say that the pair  $(u, v)$  is a weak solution to the equation (4.16) on  $\Omega$  with the initial data  $(u(x, 0), v(x, 0)) = (u_0, v_0) \in L^1_{loc}(\mathbb{H}^n) \times L^1_{loc}(\mathbb{H}^n)$ , if  $(u, v) \in C^2(0, T; S^2_2(\mathbb{H}^n)) \cap L^p(\Omega) \times C^2(0, T; S^2_2(\mathbb{H}^n)) \cap L^q(\Omega)$  satisfies

$$\begin{aligned} & \int_{\Omega} |v|^q \varphi dx dt + \int_{\mathbb{H}^n} u_1(x) \varphi(x, 0) dx - \int_{\mathbb{H}^n} u_0(x) \varphi_t(x, 0) dx = \int_{\Omega} u \varphi_{tt} dx dt \\ & - \int_{\Omega} M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) u \Delta_{\mathbb{H}} \varphi dx dt, \end{aligned}$$

and

$$\begin{aligned} & \int_{\Omega} |u|^p \psi dx dt + \int_{\mathbb{H}^n} v_1(x) \psi(x, 0) dx - \int_{\mathbb{H}^n} v_0(x) \psi_t(x, 0) dx = \int_{\Omega} v \psi_{tt} dx dt \\ & - \int_{\Omega} M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) v \Delta_{\mathbb{H}} \psi dx dt, \end{aligned}$$

for any test function  $0 \leq \varphi, \psi \in C^{2,2}_{x,t}(\Omega)$ .

For simplicity, let us denote

$$\theta_i = \frac{4}{2 + \beta_i}, \quad i = 1, 2, \quad (4.18)$$

and  $\tilde{\theta} = \max\{\theta_1, \theta_2\}$ . We give the following main result for the system of wave equations.

**Theorem 4.4.** *Suppose that the condition (4.17) holds. Assume that the initial data  $(u_0, v_0), (u_1, v_1) \in L^1(\mathbb{H}^n) \times L^1(\mathbb{H}^n)$  satisfy*

$$\int_{\mathbb{H}^n} u_1 dx \geq 0, \quad \text{and} \quad \int_{\mathbb{H}^n} v_1 dx \geq 0. \quad (4.19)$$

If

$$1 < pq \leq 1 + \frac{\theta_2 q + \theta_1}{Q + \frac{\tilde{\theta}}{2} - \theta_1}, \quad \text{or} \quad 1 < pq \leq 1 + \frac{\theta_1 p + \theta_2}{Q + \frac{\tilde{\theta}}{2} - \theta_2},$$

with  $Q = 2n + 2$  the homogeneous dimension of  $\mathbb{H}^n$ , then there exists no nontrivial weak solution of the system (4.16).

*Proof.* By analogy with the single equation case, one obtains

$$\int_{\Omega} |v|^q \varphi dxdt + \int_{\mathbb{H}^n} u_1(x) \varphi(x, 0) dx \leq \int_{\Omega} |u| |\varphi_{tt}| dxdt + C_1 \int_{\Omega} t^{\beta_1} |u| |\Delta_{\mathbb{H}} \varphi| dxdt$$

and

$$\int_{\Omega} |u|^p \psi dxdt + \int_{\mathbb{H}^n} v_1(x) \psi(x, 0) dx \leq \int_{\Omega} |v| |\psi_{tt}| dxdt + C_2 \int_{\Omega} t^{\beta_2} |v| |\Delta_{\mathbb{H}} \psi| dxdt.$$

By taking  $\varphi = \varphi_R$  and  $\psi = \psi_R$  as in (4.6) and using the Hölder inequality, we calculate

$$\begin{aligned} \int_{\Omega} |v|^q \varphi_R dxdt + \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ \leq C(A_p(\varphi_R)^{\frac{p-1}{p}} + B_p(\varphi_R)^{\frac{p-1}{p}}) \left( \int_{\Omega} |u|^p \psi_R dxdt \right)^{\frac{1}{p}}, \end{aligned}$$

and

$$\begin{aligned} \int_{\Omega} |u|^p \psi_R dxdt + \int_{\mathbb{H}^n} v_1(x) \psi_R(x, 0) dx \\ \leq C(A_q(\psi_R)^{\frac{q-1}{q}} + B_q(\psi_R)^{\frac{q-1}{q}}) \left( \int_{\Omega} |v|^q \varphi_R dxdt \right)^{\frac{1}{q}}, \end{aligned}$$

where  $C > 0$ ,

$$A_p(\varphi_R) = \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |(\varphi_R)_{tt}|^{\frac{p}{p-1}} dxdt < \infty,$$

$$B_p(\varphi_R) = \int_{\Omega_1} t^{\frac{\beta_1 p}{p-1}} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dxdt < \infty,$$

$$A_q(\psi_R) = \int_{\Omega_1} \psi_R^{-\frac{1}{q-1}} |(\psi_R)_{tt}|^{\frac{q}{q-1}} dxdt < \infty,$$

and

$$B_q(\psi_R) = \int_{\Omega_1} t^{\frac{\beta_2 q}{q-1}} \psi_R^{-\frac{1}{q-1}} |\Delta_{\mathbb{H}} \psi_R|^{\frac{q}{q-1}} dxdt < \infty.$$

By using the dominated convergence theorem with  $u_1, v_1 \in L^1_{loc}(\mathbb{H}^n)$ , and conditions  $\int_{\mathbb{H}^n} u_1 dx \geq 0$  and  $\int_{\mathbb{H}^n} v_1 dx \geq 0$ , we obtain

$$\begin{aligned} \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx &\geq \liminf_{R \rightarrow \infty} \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx = \int_{\mathbb{H}^n} u_1 dx \geq 0, \\ \int_{\mathbb{H}^n} v_1(x) \psi_R(x, 0) dx &\geq \liminf_{R \rightarrow \infty} \int_{\mathbb{H}^n} v_1(x) \psi_R(x, 0) dx = \int_{\mathbb{H}^n} v_1 dx \geq 0. \end{aligned}$$

Thus, we get

$$\int_{\Omega} |v|^q \varphi_R dx dt \leq C(A_p(\varphi_R)^{\frac{p-1}{p}} + B_p(\varphi_R)^{\frac{p-1}{p}}) \left( \int_{\Omega} |u|^p \psi_R dx dt \right)^{\frac{1}{p}},$$

and

$$\int_{\Omega} |u|^p \psi_R dx dt \leq C(A_q(\psi_R)^{\frac{q-1}{q}} + B_q(\psi_R)^{\frac{q-1}{q}}) \left( \int_{\Omega} |v|^q \varphi_R dx dt \right)^{\frac{1}{q}}.$$

By choosing variables  $R^{\frac{\theta_i}{2}} \bar{t} = t$ ,  $R\bar{\xi} = \xi$ ,  $R\widehat{\xi} = \tilde{\xi}$  and  $R^2\tilde{\tau} = \tau$  in the domain  $\Omega_1$  (in (4.9)), for  $i = 1, 2$ , we get

$$\int_{\Omega} |v|^q \varphi_R dx dt \leq CR^{\frac{(Q+\frac{\theta_1}{2})(p-1)}{p} - \theta_1} \left( \int_{\Omega} |u|^p \psi_R dx dt \right)^{\frac{1}{p}},$$

and

$$\int_{\Omega} |u|^p \psi_R dx dt \leq CR^{\frac{(Q+\frac{\theta_2}{2})(q-1)}{q} - \theta_2} \left( \int_{\Omega} |v|^q \varphi_R dx dt \right)^{\frac{1}{q}}.$$

Combining the last two facts, one concludes

$$\left( \int_{\Omega} |u|^p \psi_R dx dt \right)^{1 - \frac{1}{pq}} \leq CR^{\alpha_1}, \quad (4.20)$$

and

$$\left( \int_{\Omega} |v|^q \varphi_R dx dt \right)^{1 - \frac{1}{pq}} \leq CR^{\alpha_2}, \quad (4.21)$$

where

$$\alpha_1 = \frac{\left(Q + \frac{\tilde{\theta}}{2}\right)(pq - 1) - \theta_1 pq - \theta_2 q}{pq} \quad \text{and} \quad \alpha_2 = \frac{\left(Q + \frac{\tilde{\theta}}{2}\right)(pq - 1) - \theta_2 pq - \theta_1 p}{pq},$$

with  $\tilde{\theta} = \max\{\theta_1, \theta_2\}$ .

**Case 1:**  $1 < pq < 1 + \frac{\theta_2 q + \theta_1}{Q + \frac{\tilde{\theta}}{2} - \theta_1}$ , (or  $1 < pq < 1 + \frac{\theta_1 p + \theta_2}{Q + \frac{\tilde{\theta}}{2} - \theta_2}$ ). By letting  $R \rightarrow \infty$  in (4.20) with  $1 < pq < 1 + \frac{\theta_2 q + \theta_1}{Q + \frac{\tilde{\theta}}{2} - \theta_1}$ , we obtain

$$\int_{\Omega} |u|^p dx dt = 0,$$

which is a contradiction. Similarly, from (4.21) and  $1 < pq < \frac{\theta_1 p + \theta_2}{Q + \frac{\tilde{\theta}}{2} - \theta_2}$ , we obtain

$$\int_{\Omega} |v|^q dx dt = 0.$$

**Case 2:**  $pq = 1 + \max\left\{\frac{\theta_2 q + \theta_1}{Q + \frac{\tilde{\theta}}{2} - \theta_1}, \frac{\theta_1 p + \theta_2}{Q + \frac{\tilde{\theta}}{2} - \theta_2}\right\}$ . This case can be treated in the same way as in the proof of Theorem 4.2.  $\square$

**Corollary 4.5.** *If  $p = q > 1$ ,  $u = v$  and  $M_1 = M_2$  in Theorem 4.4, we obtain the result for a single equation given by Theorem 4.2.*

*Proof.* From Theorem 4.4, we get

$$p^2 \leq 1 + \frac{\frac{4}{2+\beta}(p+1)}{Q - \frac{2}{2+\beta}},$$

which means

$$p^2 - 1 \leq \frac{4(p+1)}{(2+\beta)Q - 2}.$$

Thus, by dividing both sides by  $p+1$ , we obtain

$$p - 1 \leq \frac{4}{(2+\beta)Q - 2}, \quad (4.22)$$

finishing the proof.  $\square$

**4.3. Pseudo-hyperbolic equations and systems.** In this subsection we show a nonexistence result for the Kirchhoff type pseudo-hyperbolic equation on Heisenberg groups in the following form:

$$\begin{cases} u_{tt} - M\left(t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx\right) (\Delta_{\mathbb{H}} u + \Delta_{\mathbb{H}} u_{tt}) = |u|^p, & (x, t) \in \Omega, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), & x \in \mathbb{H}^n, \end{cases} \quad (4.23)$$

where  $p > 1$ ,  $\Omega = \mathbb{H}^n \times (0, T)$  and  $M : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  is a bounded function such that

$$0 < M(\cdot, \cdot, \cdot) \leq C_0. \quad (4.24)$$

Let us formulate a definition of the weak solution of the equation (4.23).

**Definition 4.6.** We say that  $u$  is a weak solution to (4.23) on  $\mathbb{H}^n \times (0, T)$  with  $u_0, u_1 \in L^1_{loc}(\mathbb{H}^n)$ , if  $u \in C^2(0, T; S^2_2(\mathbb{H}^n)) \cap L^p(\Omega)$  and satisfies

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi dx dt + \int_{\mathbb{H}^n} u_1(x) (\varphi(x, 0) - \Delta_{\mathbb{H}} \varphi(x, 0)) dx \\ & - \int_{\mathbb{H}^n} u_0(x) (\varphi_t(x, 0) - \Delta_{\mathbb{H}} \varphi_t(x, 0)) dx = \int_{\Omega} u \varphi_{tt} dx dt \\ & - \int_{\Omega} M\left(t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx\right) u (\Delta_{\mathbb{H}} \varphi + \Delta_{\mathbb{H}} \varphi_{tt}) dx dt, \end{aligned} \quad (4.25)$$

for any test function  $0 \leq \varphi \in C^{2,2}_{x,t}(\Omega)$ .

**Theorem 4.7.** Assume that  $u_1, u_0 \in L^1(\mathbb{H}^n)$ , and

$$\int_{\mathbb{H}^n} u_1 dx \geq 0. \quad (4.26)$$

Suppose that the condition (4.24) is true. If

$$1 < p \leq p_c = \frac{Q+1}{Q-1}, \quad (4.27)$$

with  $Q$  the homogeneous dimension of  $\mathbb{H}^n$ , then there exists no nontrivial weak solution to (4.23).

*Proof.* Consider the case  $1 < p < p_c$ . Let us choose a test function in the following form

$$\varphi_R(x, t) = \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right) \Phi \left( \frac{t^2}{R^2} \right), \quad (x, t) \in \mathbb{H}^n \times \mathbb{R}_+, \quad R > 0, \quad (4.28)$$

with  $\Phi \in C_0^\infty(\mathbb{R}_+)$  as

$$\Phi(r) = \begin{cases} 1, & \text{if } 0 \leq r \leq 1, \\ \searrow, & \text{if } 1 < r \leq 2, \\ 0, & \text{if } r > 2. \end{cases}$$

By noting that

$$\frac{\partial \varphi_R(x, t)}{\partial t} = \frac{2t}{R^2} \Phi \left( \frac{|\xi|^4 + |\tilde{\xi}|^4 + \tau^2}{R^4} \right) \Phi'_t \left( \frac{t^2}{R^2} \right), \quad (4.29)$$

one has

$$\frac{\partial \varphi_R(x, 0)}{\partial t} = 0. \quad (4.30)$$

By Definition 4.6, we obtain

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi_R dx dt + \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \stackrel{(4.30)}{=} \int_{\Omega} |u|^p \varphi_R dx dt \\ & + \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx - \int_{\mathbb{H}^n} u_0(x) ((\varphi_R)_t(x, 0) - \Delta_{\mathbb{H}}(\varphi_R)_t(x, 0)) dx \\ & = \int_{\Omega} u(\varphi_R)_{tt} dx dt - \int_{\Omega} M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u(\Delta_{\mathbb{H}} \varphi + \Delta_{\mathbb{H}} \varphi_{tt}) dx dt \\ & + \int_{\mathbb{H}^n} u_1(x) \Delta_{\mathbb{H}} \varphi_R(x, 0) dx \\ & \leq \left| \int_{\Omega} u(\varphi_R)_{tt} dx dt - \int_{\Omega} M \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx \right) u(\Delta_{\mathbb{H}} \varphi + \Delta_{\mathbb{H}} \varphi_{tt}) dx dt \right. \\ & \left. + \int_{\mathbb{H}^n} u_1(x) \Delta_{\mathbb{H}} \varphi_R(x, 0) dx \right| \\ & \leq \int_{\Omega} |u(\varphi_R)_{tt}| dx dt + C_0 \int_{\Omega} |u \Delta_{\mathbb{H}}(\varphi_R)_{tt}| dx dt + C_0 \int_{\Omega} |u \Delta_{\mathbb{H}} \varphi_R| dx dt \\ & + \int_{\mathbb{H}^n} |u_1(x) \Delta_{\mathbb{H}} \varphi_R(x, 0)| dx. \end{aligned} \quad (4.31)$$

Then by using the Young inequality, we get

$$\begin{aligned} \int_{\Omega_1} |u| |(\varphi_R)_{tt}| dx dt &= \int_{\Omega_1} |u| \varphi_R^{\frac{1}{p}} \varphi_R^{-\frac{1}{p}} |(\varphi_R)_{tt}| dx dt \\ &\leq \varepsilon \int_{\Omega_1} |u|^p \varphi_R dx dt + C(\varepsilon) \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |(\varphi_R)_{tt}|^{\frac{p}{p-1}} dx dt, \end{aligned} \quad (4.32)$$

$$\int_{\Omega_1} u |\Delta_{\mathbb{H}} \varphi_R| dt dx \leq \varepsilon \int_{\Omega_1} |u|^p \varphi_R dt dx + C(\varepsilon) \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}} \varphi_R|^{\frac{p}{p-1}} dx dt, \quad (4.33)$$

and

$$\int_{\Omega_1} |u| \Delta_{\mathbb{H}}(\varphi_R)_{tt} dt dx \leq \varepsilon \int_{\Omega_1} |u|^p \varphi_R dt dx + C(\varepsilon) \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}(\varphi_R)_{tt}|^{\frac{p}{p-1}} dx dt, \quad (4.34)$$

for some positive constant  $C(\varepsilon)$ . By using the above facts, we obtain

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi_R dx dt + C \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ & \leq C \left( A_p(\varphi_R) + B_p(\varphi_R) + \int_{\mathbb{H}^n} |u_1(x)| |\Delta_{\mathbb{H}} \varphi_R(x, 0)| dx \right), \end{aligned} \quad (4.35)$$

where  $C > 0$ ,

$$A_p(\varphi_R) = \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |(\varphi_R)_{tt}|^{\frac{p}{p-1}} dx dt < \infty, \quad (4.36)$$

and

$$B_p(\varphi_R) = \int_{\Omega_1} \varphi_R^{-\frac{1}{p-1}} |\Delta_{\mathbb{H}}(\varphi_R)_{tt}|^{\frac{p}{p-1}} dx dt < \infty. \quad (4.37)$$

From (4.8), we have

$$\frac{\partial \varphi_R(x, 0)}{\partial t} = 0. \quad (4.38)$$

Also, let us denote  $\bar{\Omega}_R = \Gamma_R \cup \tilde{\Gamma}_R$ ,

$$\Gamma_R := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : R^4 \leq |\xi|^4 + |\tilde{\xi}|^4 + \tau^2 \leq 2R^4\}, \quad \tilde{\Gamma}_R := \{t : R^2 \leq t^2 \leq 2R^2\}.$$

Let us estimate  $A_p(\varphi_R), B_p(\varphi_R)$ . By using the homogeneous dimension of the Heisenberg groups, we calculate

$$|A_p(\varphi_R)| \leq CR^{Q+1-\frac{2p}{p-1}}, \quad (4.39)$$

and

$$|B_p(\varphi_R)| \leq CR^{Q+1-\frac{4p}{p-1}}. \quad (4.40)$$

Choose variables  $R\bar{t} = t$ ,  $R\bar{\xi} = \xi$ ,  $R\hat{\xi} = \tilde{\xi}$  and  $R^2\bar{\tau} = \tau$  in the domain  $\Omega_1$ , and by using (4.39), (4.40) in (4.35), one calculates

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi_R dx dt + C \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ & \leq C \left( R^{Q+1-\frac{2p}{p-1}} + C \int_{\Gamma} |u_1(x)| |\Delta_{\mathbb{H}} \varphi_R(x, 0)| dx \right). \end{aligned} \quad (4.41)$$

On the other hand, we have

$$\begin{aligned} & \int_{\Omega} |u|^p \varphi_R dx dt + C \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ & \geq \liminf_{R \rightarrow \infty} \int_{\Omega} |u|^p \varphi_R dx dt + C \liminf_{R \rightarrow \infty} \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx. \end{aligned}$$

Using the monotone convergence theorem, we obtain

$$\liminf_{R \rightarrow \infty} \int_{\Omega} |u|^p \varphi_R dx dt = \int_{\Omega} |u|^p dx dt.$$



Since  $u_1 \in L^1(\mathbb{G})$ , by the dominated convergence theorem, one has

$$\liminf_{R \rightarrow \infty} \int_{\Gamma} u_1(x) \varphi_R(x, 0) dx = \int_{\mathbb{H}^n} u_1(x) dx.$$

Now, we have

$$\liminf_{R \rightarrow \infty} \left( \int_{\Omega} |u|^p \varphi_R dx dt + C \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \right) \geq \int_{\Omega} |u|^p dx dt + Cd,$$

where

$$d = \int_{\mathbb{H}^n} u_1(x) dx \geq 0.$$

By the definition of the limit, for every  $\alpha > 0$  there exists  $R_0 > 0$  such that

$$\begin{aligned} \int_{\Omega} |u|^p \varphi_R dx dt + C \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \\ \geq \liminf_{R \rightarrow \infty} \left( \int_{\Omega} |u|^p \varphi_R dx dt + \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \right) - \alpha \\ \geq \int_{\Omega} |u|^p dx dt + Cd - \alpha, \end{aligned}$$

for every  $R \geq R_0$ . By choosing  $\alpha = \frac{Cd}{2}$ , we get

$$\int_{\Omega} |u|^p \varphi_R dx dt + \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx \geq \int_{\Omega} |u|^p dx dt + \frac{Cd}{2},$$

for every  $R \geq R_0$ . Then from  $u_1 \in L^1(\mathbb{H}^n)$ , one obtains

$$\begin{aligned} \int_{\Omega} |u|^p dx dt + \frac{Cd}{2} &\leq C \left( R^{Q+1-\frac{2p}{p-1}} + \int_{\Gamma} |u_1(x)| |\Delta_{\mathbb{H}} \varphi_R(x, 0)| dx \right) \\ &\leq C \left( R^{Q+1-\frac{2p}{p-1}} + R^{-2} \int_{\Gamma} |u_1(x)| dx \right) \\ &\leq C \left( R^{Q+1-\frac{2p}{p-1}} + R^{-2} \right). \end{aligned} \tag{4.42}$$

From assumption we have  $p < 1 + \frac{2}{p-1}$ , or, in other words  $Q + 1 - \frac{2p}{p-1} < 0$  and by using this fact with letting  $R \rightarrow \infty$ , we obtain

$$\int_{\Omega} |u|^p dx dt \leq \int_{\Omega} |u|^p dx dt + C \frac{d}{2} \leq 0,$$

that is,

$$\int_{\Omega} |u|^p dx dt = 0.$$

It is a contradiction.

Now, we consider the critical case  $p = 1 + \frac{2}{Q-1}$ . By using (4.42), we have

$$\int_{\Omega} |u|^p dx dt \leq C < \infty. \tag{4.43}$$

Then

$$\lim_{R \rightarrow \infty} \int_{\Omega_R} |u|^p \varphi_R dx dt = 0, \tag{4.44}$$

where  $\bar{\Omega}_R = \Gamma_{1,R} \cup \tilde{\Gamma}_{1,R}$ ,

$$\Gamma_{1,R} := \{x = (\xi, \tilde{\xi}, \tau) \in \mathbb{H}^n : R^4 \leq |x|^4 \leq 2R^4\}, \quad \tilde{\Gamma}_{1,R} := \{t : R^2 \leq t^2 \leq 2R^2\}.$$

Using the Hölder inequality instead of the Young's inequality in (4.31), one obtains

$$\int_{\Omega} |u|^p \varphi_R dx dt \leq \int_{\Omega} |u|^p \varphi_R dx dt + C \frac{d}{2} \leq C \left( \int_{\bar{\Omega}_R} |u|^p \varphi_R dx dt \right)^{\frac{1}{p}}.$$

Letting  $R \rightarrow \infty$  in the above, we get

$$\int_{\Omega} |u|^p \varphi_R dx dt = 0,$$

arriving at a contradiction.  $\square$

Let us consider the Cauchy problem for the system of Kirchhoff type pseudo-hyperbolic equations

$$\begin{cases} u_{tt} - M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) (\Delta_{\mathbb{H}} u + \Delta_{\mathbb{H}} u_{tt}) \\ \quad = |v|^q, \quad (x, t) \in \Omega, \\ v_{tt} - M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) (\Delta_{\mathbb{H}} v + \Delta_{\mathbb{H}} v_{tt}) \\ \quad = |u|^p, \quad (x, t) \in \Omega, \\ u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), \quad x \in \mathbb{H}^n, \\ v(x, 0) = v_0(x), v_t(x, 0) = v_1(x), \quad x \in \mathbb{H}^n, \end{cases} \quad (4.45)$$

where  $\Omega = \mathbb{H}^n \times (0, T)$ ,  $T > 0$ ,  $p, q > 1$ , and  $M_1, M_2 : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$  are bounded functions such that

$$0 < M_1(\cdot, \cdot, \cdot, \cdot, \cdot) \leq C_1, \quad (4.46)$$

and

$$0 < M_2(\cdot, \cdot, \cdot, \cdot, \cdot) \leq C_2. \quad (4.47)$$

Now, we formulate the following definition of the weak solution to the system (4.45).

**Definition 4.8.** We say that the pair  $(u, v)$  is a weak solution to (4.45) on  $\Omega = \mathbb{H}^n \times (0, T)$  with the initial data  $(u(x, 0), v(x, 0)) = (u_0, v_0) \in L^1_{loc}(\mathbb{H}^n) \times L^1_{loc}(\mathbb{H}^n)$ , if  $(u, v) \in C^2(0, T; S^2_2(\mathbb{H}^n)) \cap L^p(\Omega) \times C^2(0, T; S^2_2(\mathbb{H}^n)) \cap L^q(\Omega)$  with  $p, q > 1$  and they satisfy

$$\begin{aligned} & \int_{\Omega} |v|^q \varphi dx dt + \int_{\mathbb{H}^n} u_1(x) (\varphi(x, 0) - \Delta_{\mathbb{H}} \varphi(x, 0)) dx \\ & - \int_{\mathbb{H}^n} u_0(x) (\varphi_t(x, 0) - \Delta_{\mathbb{H}} \varphi_t(x, 0)) dx = \int_{\Omega} u \varphi_{tt} dx dt \\ & - \int_{\Omega} M_1 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) u (\Delta_{\mathbb{H}} \varphi + \Delta_{\mathbb{H}} \varphi_{tt}) dx dt, \end{aligned}$$

and

$$\begin{aligned} & \int_{\Omega} |u|^p \psi dx dt + \int_{\mathbb{H}^n} v_1(x) (\psi(x, 0) - \Delta_{\mathbb{H}} \psi(x, 0)) dx \\ & - \int_{\mathbb{H}^n} v_0(x) (\psi_t(x, 0) - \Delta_{\mathbb{H}} \psi_t(x, 0)) dx = \int_{\Omega} v \psi_{tt} dx dt \\ & - \int_{\Omega} M_2 \left( t, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\nabla_{\mathbb{H}} v|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} u|^2 dx, \int_{\mathbb{H}^n} |\Delta_{\mathbb{H}} v|^2 dx \right) v (\Delta_{\mathbb{H}} \psi + \Delta_{\mathbb{H}} \psi_{tt}) dx dt, \end{aligned}$$

for any test functions  $0 \leq \varphi, \psi \in C_{x,t}^{2,2}(\Omega)$ .

Then we have the following theorem.

**Theorem 4.9.** *Let  $p, q > 1$ ,  $(u_0, v_0), (u_1, v_1) \in L^1(\mathbb{H}^n) \times L^1(\mathbb{H}^n)$  with*

$$\int_{\mathbb{H}^n} u_1 dx \geq 0, \quad \text{and} \quad \int_{\mathbb{H}^n} v_1 dx \geq 0. \quad (4.48)$$

*Assume that the conditions (4.46) and (4.47) hold. If*

$$1 < pq \leq 1 + \frac{2}{Q-1} \max\{p+1; q+1\},$$

*with homogeneous dimension  $Q = 2n + 2$  of  $\mathbb{H}^n$ , then there exists no nontrivial weak solution to (4.45).*

*Proof.* The proof of this theorem is similar to the proof of Theorem 4.7. The only thing that we need to note is that for the large  $R$ , we have

$$\begin{aligned} & \int_{\mathbb{H}^n} u_1(x) \varphi_R(x, 0) dx - \int_{\mathbb{H}^n} |u_1(x)| |\Delta_{\mathbb{H}} \varphi_R(x, 0)| dx \geq 0, \\ & \int_{\mathbb{H}^n} v_1(x) \psi_R(x, 0) dx - \int_{\mathbb{H}^n} |v_1(x)| |\Delta_{\mathbb{H}} \psi_R(x, 0)| dx \geq 0. \end{aligned} \quad (4.49)$$

And, by repeating the proof of Theorem 4.7, we finish our proof.  $\square$

## REFERENCES

- [1] B. Ahmad, A. Alsaedi and M. Kirane. Nonexistence of global solutions of some nonlinear space-nonlocal evolution equations on the Heisenberg groups, *Electronic Journal of Differential Equations*, 2015(227):1–10, 2015.
- [2] B. Ahmad, A. Alsaedi and M. Kirane. Blow-up of solutions to parabolic inequalities in the Heisenberg group, *Electronic Journal of Differential Equations*, 2015(167):1–9, 2015.
- [3] A. Ambrosetti and P. Rabinowitz. Dual methods in critical point theory and applications. *J. Func. Anal.*, 14:349–381, 1973.
- [4] G. Autuori, P. Pucci and M.C. Salvatori. Asymptotic stability for anisotropic Kirchhoff systems. *J. Math. Anal. Appl.*, 352:149–165, 2009.
- [5] G. Autuori and P. Pucci. Asymptotic stability for Kirchhoff systems in variable exponent Sobolev spaces. *Complex Var. Elliptic Equ.*, 56:715–753, 2011.
- [6] G. Citti. Semilinear Dirichlet problem involving critical exponent for the Kohn Laplacian. *Ann. Mat. Pura Appl.*, 169:375–392, 1995.
- [7] L. D’Ambrosio. Critical degenerate inequalities on the Heisenberg group. *Manuscripta Mathematica*, 106(4):519–536, 2001.
- [8] P. Drábek and S.I. Pohozaev. Positive solutions for the  $p$ -Laplacian: application of the fibering method. *Proceedings of the Royal Society of Edinburgh*, 127(4):703–726, 1997.

- [9] V. Fischer and M. Ruzhansky. Quantization on Nilpotent Lie Groups. Progress in Mathematics. Birkhauser, 2016.
- [10] H. Fujita. On the blowing up of solutions of the Cauchy problem for  $u_t = \Delta u + u^{1+\alpha}$ . J. Fac. Sci. Univ. Tokyo Sect. I, 13:109–124, 1966.
- [11] C. Garetto and M. Ruzhansky. Wave equation for sums of squares on compact Lie groups. J. Differential Equations, 258:4324–4347, 2015.
- [12] N. Garofalo and E. Lanconelli. Existence and nonexistence results for semilinear equations on the Heisenberg group. Indiana Univ. Math. J., 41:71–98, 1992.
- [13] V. Georgiev and A. Palmieri. Lifespan estimates for local in time solutions to the semilinear heat equation on the Heisenberg group. Annali di Matematica Pura ed Applicata, doi:10.1007/s10231-020-01023-z, 2020.
- [14] V. Georgiev, A. Palmieri. Critical exponent of Fujita-type for the semilinear damped wave equation on the Heisenberg group with power nonlinearity. Journal of Differential Equations, 269(1):420–448, 2020.
- [15] M. Guedda and M. Kirane. A note on nonexistence of global solutions to a nonlinear integral equation. Bull. Belg. Math. Soc. Simon Stevin, 6:491–497, 1999.
- [16] H. Hashimoto and T. Yamazaki. Hyperbolic-parabolic singular perturbation for quasilinear equations of Kirchhoff type. J. Differ. Equ., 237:491–525, 2007.
- [17] M. Jleli, M. Kirane and B. Samet. Nonexistence results for a class of evolution equations in the Heisenberg group. Fractional Calculus and Applied Analysis, 18(3):717–734, 2015.
- [18] A. Kassymov and D. Suragan. Existence of solutions for p-sub-Laplacians with nonlinear sources on the Heisenberg group. Complex Variables and Elliptic Equations, 66(4):614–625, 2021.
- [19] A. Kassymov and D. Suragan. Multiplicity of positive solutions for a nonlinear equation with a Hardy potential on the Heisenberg group. Bulletin des Sciences Mathématiques, 165:102916, 2020.
- [20] T. Kato. Blow-up of solutions of some nonlinear hyperbolic equations in three space dimensions. Comm. Pure Appl. Math., 37:443–455, 1984.
- [21] T. Matsuyama and M. Ruzhansky. On the Gevrey well-posedness of the Kirchhoff equation. J. Anal. Math., 137:449–468, 2019.
- [22] T. Matsuyama and M. Ruzhansky. Almost global well-posedness of Kirchhoff equation with Gevrey data. C. R. Math. Acad. Sci. Paris, 355(5):522–525, 2017.
- [23] T. Matsuyama and M. Ruzhansky. Global well-posedness of Kirchhoff systems. J. Math. Pures Appl., 100:220–240, 2013.
- [24] X. Mingqi and V.D. Rădulescu and B. Zhang. Fractional Kirchhoff problems with critical Trudinger-Moser nonlinearity. Calc. Var. Partial Differential Equations, 58(2):1–27, 2019
- [25] X. Mingqi, V. D. Rădulescu and B. Zhang. Nonlocal Kirchhoff diffusion problems: local existence and blow-up of solutions. Nonlinearity, 31(7):3228–325, 2018.
- [26] M. Ruzhansky and D. Suragan. A comparison principle for nonlinear heat Rockland operators on graded groups. Bulletin of the London Mathematical Society, 50(5):753–758, 2018.
- [27] M. Ruzhansky and N. Tokmagambetov. Nonlinear damped wave equations for the sub-Laplacian on the Heisenberg group and for Rockland operators on graded Lie groups. J. Differential Equations, 265:5212–5236, 2018.
- [28] M. Ruzhansky and N. Yessirkegenov. Existence and non-existence of global solutions for semilinear heat equations and inequalities on sub-Riemannian manifolds, and Fujita exponent on unimodular Lie groups. arXiv:1812.01933, 2019, to appear in J. Differential Equations.
- [29] M. Ruzhansky and N. Yessirkegenov. A comparison principle for higher order nonlinear hypoelliptic heat operators on graded Lie groups. arXiv:2005.11503, 2020, to appear in Nonlinear Analysis.
- [30] S. Sugitani. On nonexistence of global solutions for some nonlinear integral equations. Osaka J. Math., 12:45–51, 1975.
- [31] L. Véron and S.I. Pohozaev. Nonexistence results of solutions of semilinear differential inequalities on the Heisenberg group. Manuscripta Math., 102:85–99, 2000.

- [32] M. Xiang, B. Zhang, and M. Ferrara. Existence of solutions for Kirchhoff type problem involving the non-local fractional  $p$ -Laplacian. *Journal of Mathematical Analysis and Applications*, 424(2):1021–1041, 2015.

AIDYN KASSYMOV:  
DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, BELGIUM  
AND  
INSTITUTE OF MATHEMATICS AND MATHEMATICAL MODELING  
ALMATY, KAZAKHSTAN  
*E-mail address* kassymov@math.kz AND aidyn.kassymov@ugent.be

MICHAEL RUZHANSKY:  
DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, BELGIUM  
AND  
SCHOOL OF MATHEMATICAL SCIENCES  
QUEEN MARY UNIVERSITY OF LONDON  
UNITED KINGDOM  
*E-mail address* michael.ruzhansky@ugent.be

NIYAZ TOKMAGAMBETOV:  
DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, BELGIUM  
AND  
AL-FARABI KAZAKH NATIONAL UNIVERSITY  
71 AL-FARABI AVENUE  
050040 ALMATY  
KAZAKHSTAN  
*E-mail address* niyaz.tokmagambetov@ugent.be

BERIKBOL TOREBEK:  
DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, BELGIUM  
AND  
INSTITUTE OF MATHEMATICS AND MATHEMATICAL MODELING  
ALMATY, KAZAKHSTAN  
*E-mail address* berikbol.torebek@ugent.be