CONSTANT SCALAR CURVATURE KÄHLER METRICS ON RAMIFIED GALOIS COVERINGS

CLAUDIO AREZZO, ALBERTO DELLA VEDOVA, AND YALONG SHI

ABSTRACT. We give sufficient conditions for the existence of Kähler-Einstein and constant scalar curvature Kähler (cscK) metrics on finite ramified Galois coverings of a cscK manifold in terms of cohomological conditions on the Kähler classes and the branching divisor. This result generalizes previous work on Kähler-Einstein metrics by Li-Sun [21], and extends Chen-Cheng's existence results for cscK metrics in [8].

CONTENTS

1.	Introduction	1
2.	Preliminaries	4
3.	K-energy under Galois coverings	7
4.	Proof of Theorem 1.1	9
5.	Proof of Theorem 1.5	11
6.	Applications and Examples	12
6.1	. Existence of covers	12
6.2	. KE and cscK metrics: the abelian case	14
6.3	. KE and cscK metrics: the non-abelian case	15
References		19

1. INTRODUCTION

In this paper we provide a new geometric construction of Kähler-Einstein and constant scalar curvature Kähler (cscK) metrics on finite ramified Galois coverings of cscK manifolds. In order to state our results let us first recall the notion of a Galois cover.

Let M, N be complex manifolds of dimension n. A holomorphic map $f : M \to N$ is called "finite", if it is proper and has finite fibers. If f is moreover surjective, it is called an "analytic covering". For such an analytic covering f, we can define the Galois group

$$Gal(f) := \{ \gamma \in Aut(M) \mid f \circ \gamma = f \}.$$

f is called a "Galois covering" if any two points of *M* lie on the same fiber of *f* is and if they belong to the same Gal(f)-orbit. In other words, $N \cong M/Gal(f)$.

Now we assume both M and N are smooth compact Kähler manifolds, and N has a constant scalar curvature Kähler metric ("cscK metric" for short) with Kähler form ω_N . If $f: M \to N$ is an unramified (or "étale") Galois covering, then $f^*\omega_N$ is a cscK metric on M. The central question addressed in this paper is whether (or when) cscK metrics still exist in the class $[f^*\omega_N]$ also when f is ramified. Note that in this case while $f^*\omega_N$ is degenerate, thanks to Demailly-Paun's result [12], the class $[f^*\omega_N]$ is still a Kähler class (see Section 2).

When both *M* and *N* are Fano manifolds and ω_N is Kähler-Einstein, this problem is well studied and partial results have been found by Arezzo-Ghigi-Pirola [1], Ghigi-Kollar [15] and Li-Sun [21]. Their methods are all variational, using the (non-trivial) fact that the properness of Ding-functional (resp. log-K-energy) implies the existence of Kähler-Einstein metrics (resp. conical Kähler-Einstein metrics).

A recent breakthrough in Kähler geometry is Chen-Cheng's work [8], showing that the properness of K-energy in a general Kähler class implies the existence of Kähler metrics of constant scalar curvature. In this paper, we shall use a (*G*-invariant version) of Chen-Cheng's theorem (Theorem 4.1) to prove that the above mentioned theorems on Kähler-Einstein metrics generalize to the cscK setting.

To state our main theorem we introduce the following notations: let $f : M \to N$ be a finite Galois covering of degree d with Galois group G. Denote the ramification divisor by $R_f = \sum_j r_j R_j$. It is known that locally on the regular part of R_j , f is of the form $(z_1, \ldots, z_n) \mapsto (z_1, \ldots, z_{n-1}, z_n^{r_j+1})$ under appropriate holomorphic coordinates (Lemma 5 of [15]). So we have $K_M = f^*K_N + R_f$. As in [15], we have $R_f = f^*B_f$ where $B_f = \sum_i (1 - \frac{1}{|\Gamma_i|})B_i$ is the branching divisor. Here $\Gamma_i := \{\gamma \in Gal(f) \mid R_j \subset Fix(\gamma)\}$ is the inertia subgroup of any R_j such that $f(R_j) = B_i$. Write $d_i := |\Gamma_i|$. Note that we do not assume R_f or B_f is simple normal crossing but we only require that both M and N are smooth.

The main result of this paper is then the following theorem:

Theorem 1.1. Let (N, ω_N) be a n-dimensional compact Kähler manifold, and $f : M \to N$ a finite ramified Galois covering, branched along divisor $B_f = \sum_{i=1}^k (1 - \frac{1}{d_i})B_i$ of N with ramification divisor R_f . Assume that the K-energy on N of the class $[\omega_N]$ is bounded from below (For example, if $[\omega_N]$ has a cscK metric). Write $B_{\delta} := \delta \sum_{i=1}^k (1 - \frac{1}{d_i})B_i$ and assume that for some $\delta_0 \in [1, \min\{\frac{d_i}{d_i-1}\})$, the classes $K_N + B_{\delta_0}$ and

$$\frac{n(K_N + B_{\delta_0}) \cdot [\omega_N]^{n-1}}{[\omega_N]^n} [\omega_N] - (n-1)(K_N + B_{\delta_0})$$

are both nef. Then there exists a cscK metric on M in the Kähler class $[f^*\omega_N]$.

The simplest examples of ramified Galois coverings are finite cyclic coverings branching along an irreducible divisor (which is necessarily smooth by Catanese [5, Proposition 1.1]). In this special case, Theorem 1.1 reduces to:

Corollary 1.2. Let (N, ω_N) be a n-dimensional compact cscK manifold, and $f : M \to N$ a finite cyclic cover of order d, branched along an irreducible divisor $B \subset N$ with ramification divisor R_f . If the Kähler class $[\omega_N]$ and [B] satisfy the condition that there is a positive constant $\beta_0 \in (0, \frac{1}{d}]$ such that both $K_N + (1 - \beta_0)B$ and

$$\frac{n(K_N + (1 - \beta_0)B) \cdot [\omega_N]^{n-1}}{[\omega_N]^n} [\omega_N] - (n-1)(K_N + (1 - \beta_0)B)$$

are nef, then there exists a cscK metric on M in the Kähler class $[f^*\omega_N]$.

The idea of the proof to Theorem 1.1 is clearest in the case of simple cyclic cover, i.e. Corollary 1.2. In this case, one should have cscK metrics with cone angle $2\pi\beta$ along *B* for any $\beta_0 \le \beta < 1$. Then the pull back of the conical cscK metric with angle $\frac{2\pi}{d}$ will give rise to a smooth cscK metric in the class $[f^*\omega_N]$. This is indeed true in the conical Kähler-Einstein case as observed in [21]. Even though there are already systematic studies on conic cscK metrics (e.g. [31]), we can in fact avoid such "conical cscK metrics" altogether except the concepts of "log *K*-energy" and "log- α -invariant".

Our strategy is as follows:

cscK on N \implies K-energy bounded from below on N \implies log K-energy proper on N \implies K-energy proper in the space of *G*-invariant potentials on M.

The first step is already well known. The main part of our proof is devoted to step 2 and step 3.

Remark 1.3. When B_f is simple normal crossing, one can use Zheng's theorem on conical cscK metrics [31]. Our method avoids analytical difficulties involving conical metrics. Moreover, our theorem works even if B_f is not simple normal crossing, while a suitable "conical theory" is still missing.

Remark 1.4. In view of the recent work of Z. Sjöström Dyrefelt [24] and K. Zhang [30], our main result can be improved. See, for example Remark 4.3. However, we stick to this weaker version since the conditions here are much easier to check.

When *f* is a cyclic covering branched over a smooth hypersurface *B*, and $[\omega_N]$ is proportional to $c_1(B)$, we can strengthen Corollary 1.2 using Berman's estimate of log- α -invariants [3]:

Theorem 1.5. Let (N, ω_N) be a n-dimensional compact Kähler manifold, and $f : M \to N$ a finite cyclic Galois covering, branched along a smooth hypersurface B of N. Assume ω_N is a cscK metric

on N with $[\omega_N] = \lambda 2\pi c_1(B)$ for some $\lambda > 0$. If both $K_N + B$ and

$$\Big(\frac{nK_N\cdot B^{n-1}}{B^n}+1\Big)B-(n-1)K_N$$

are nef, then there exists a cscK metric on M in the class $[f^*\omega_N]$.

This contains as a special case Theorem 6.1 of Li-Sun [21], which covers the case of N Fano and ω_N Kähler-Einstein. In fact, if $B \in |-\beta K_N|$ with $\beta \ge 1$, both conditions are satisfied, we can apply Theorem 1.5 to get the Kähler-Einstein metric on M.

The structure of the paper is as follows. In section 2, we discuss basic properties of *K*-energy on the space of potentials with finite energies. Most crucial is Lemma 2.2, saying that to check properness of the *K*-energy on \mathcal{E}^1 space, it suffices to check it on the space of smooth potentials. Then in section 3, we compute the K-energy in the space of *G*-invariant potentials and relate it to the log-K-energy downstairs. Theorem 1.1 is proved in section 4, using a log-version of Li-Shi-Yao's criterion for properness of the K-energy (which is independently discovered by K. Zheng in the conic setting), as well as an equivariant version of Chen-Cheng's theorem. This generalized version of Chen-Cheng's theorem is essentially contained in [8]. However, since it is neither stated explicitly, nor implied directly by their theorems, we decide to include a sketch of proof for the convenience of readers. Then in section 5, making use of Berman's estimate of the log- α invariant, we prove Theorem 1.5. In the last Section 6, we provide large families of new examples both of Kähler-Einstein and constant scalar curvature Kähler (cscK) metrics and compare on explicit cases our results with other known existence Theorems, notably Li-Sun [21] and Chen-Cheng's [8] ones.

Aknowledgements: The fist author is very grateful to Fabio Perroni for beautiful explanations of the Catanese-Perroni's theory largely used in the last Section to produce many interesting examples. The third author thanks K. Zheng for helpful discussions on conical metrics. This work began when the second and third authors were both visiting ICTP, they would like to thank ICTP for its hospitality.

2. Preliminaries

Before entering the analytical details, let us observe that the class $[f^*\omega_N]$ is indeed a Kähler class even when f has a non-empty ramification. In fact, when N is projective algebraic, this is standard result, see, for example, Proposition 1.2.13 of [19]. In the general case, we can apply Demailly-Paun's Theorem [12] generalizing the Nakai-Moishezon criterion, stating that hat the Kähler cone is one of the connected components of the set of all real (1, 1)-class α such that $\int_V \alpha^{\dim V} > 0$ for any analytic cycle V. In our case it is immediate to observe that

$$\int_{V} (f^*\omega_N)^p = d \int_{f(V)} \omega_N^p > 0,$$

since f(V) is a *p*-dimensional analytic cycle of *N*. Since $f^*\omega_N \ge 0$, it is in the closure of the Kähler cone and hence in the Kähler cone.

Let us now introduce the spaces of Kähler potentials we shall use: let X be a compact Kähler manifold and ω be a smooth non-negative closed (1, 1) form with $[\omega]$ is a Kähler class. We define

$$\mathcal{H}(X,\omega) := \{ \varphi \in C^{\infty}(X;\mathbb{R}) \mid \omega + \sqrt{-1}\partial \overline{\partial} \varphi > 0 \},\$$

and

$$PSH(X,\omega) := \{ \varphi \in L^1(X;\mathbb{R}) \mid \omega + \sqrt{-1}\partial\overline{\partial}\varphi \ge 0 \}.$$

Here $L^1(X; \mathbb{R})$ is defined with respect to any smooth volume form, and " $\omega + \sqrt{-1}\partial\overline{\partial}\varphi \ge 0$ " is in the sense of currents. Another important space is the space of potentials with finite energy, defined as follows: write $\omega_{\varphi} := \omega + \sqrt{-1}\partial\overline{\partial}\varphi$

$$\mathcal{E}^{1}(X,\omega) := \Big\{ \varphi \in L^{1}(X;\mathbb{R}) \, \Big| \, \int_{X} \omega_{\varphi}^{n} = \int_{X} \omega^{n}, \quad \int_{X} |\varphi| \omega_{\varphi}^{n} < +\infty \Big\},$$

where $\omega_{\varphi}^{n} := \lim_{j \to \infty} \mathbf{1}_{\varphi > -j} \left(\omega + \sqrt{-1} \partial \overline{\partial} \max\{\varphi, -j\} \right)^{n}$ is defined as the pluripolar product.

Obviously, $\mathcal{H}(X,\omega) \subset PSH(X,\omega) \cap C^0(X;\mathbb{R}) \subset \mathcal{E}^1(X,\omega) \subset PSH(X,\omega)$. It is proved by Darvas that $\mathcal{E}^1(X,\omega)$ is the metric completion of $\mathcal{H}(X,\omega)$ with respect to the L^1 -Finsler metric on \mathcal{H} . The \mathcal{E}^1 potentials appear naturally in our problem in view of the following lemma:

Lemma 2.1. Let $f : M \to N$ be a Galois covering between compact Kähler manifolds with Galois group G and branching locus B_{red} . Let ω_N be a Kähler form on N. Then for any G-invariant $\varphi \in \mathcal{H}(M, f^*\omega_N)$, we can find a function $\phi \in C^0(N; \mathbb{R}) \cap C^{\infty}(N \setminus B_{red}; \mathbb{R}) \cap PSH(N, \omega_N)$ such that $\varphi = f^*\phi = \phi \circ f$. In particular, $\phi \in \mathcal{E}^1(N, \omega_N)$.

Proof. The existence of $\phi \in C^0(N; \mathbb{R}) \cap C^{\infty}(N \setminus B_{red}; \mathbb{R})$ such that $\varphi = f^*\phi = \phi \circ f$ is obvious. Also $\omega_N + \sqrt{-1}\partial\overline{\partial}\phi > 0$ on $N \setminus B_{red}$. To show $\phi \in PSH(N, \omega_N)$, just observe that the direct image of the Kähler form $f^*\omega_N + \sqrt{-1}\partial\overline{\partial}\varphi$ is precisely $d(\omega_N + \sqrt{-1}\partial\overline{\partial}\phi)$, which is a Kähler current by Lemma 2.6 of [1].

For a Q-divisor of the form $D = \sum_i (1 - \beta_i) D_i$ on *N*, and let the defining section of D_i be s_i with metric h_i on the bundle $O(D_i)$. We can define the log-K-energy by

$$K_{D,\omega_N}(\varphi) := \int_N \log \frac{\omega_{\varphi}^n \, \omega_{\varphi}^n}{\omega_N^n \, n!} + \sum_i (1 - \beta_i) \int_N \log \|s_i\|_{h_i}^2 \left(\frac{\omega_{\varphi}^n}{n!} - \frac{\omega_N^n}{n!}\right) + \hat{J}_{-Ric(\omega_N) + \sum_i (1 - \beta_i)\Theta_{h_i}}(\varphi). \tag{2.1}$$

Lemma 2.2. To check the properness of log-K-energy on the space of \mathcal{E}^1 potentials, it suffices to check the properness on the space of smooth potentials.

Proof. Suppose we have a uniform estimate of the form

$$K_{D,\omega_N}(\varphi) \ge \epsilon (I-J)(\varphi) - C$$

on $\mathcal{H}(N, \omega_N)$. Now let $\varphi \in \mathcal{E}^1(N, \omega_N)$. We can find a sequence $\varphi_i \in \mathcal{H}(N, \omega_N)$ converging to φ in the \mathcal{E}^1 -topology.

Now it is well-known that $(I - J)(\varphi_i) \rightarrow (I - J)(\varphi)$. We need to show that

$$K_{D,\omega_N}(\varphi) \geq \liminf_{i\to\infty} K_{\beta,\omega_N}(\varphi_i).$$

Then from

$$K_{D,\omega_N}(\varphi_i) \ge \epsilon (I-J)(\varphi_i) - C$$

we will conclude that

$$K_{D,\omega_N}(\varphi) \ge \epsilon (I-J)(\varphi) - C$$

holds for \mathcal{E}^1 potential φ . However, the K-energy is only lower semi-continuous with respect to the d_1 -distance. So we need Berman-Darvas-Lu's result (Lemma 3.1 of [4]): there is a sequence of smooth potentials φ_k such that $d_1(\varphi_k, \varphi) \to 0$ and

$$\int_{N} \log \frac{\omega_{\varphi_{k}}^{n}}{\omega_{N}^{n}} \frac{\omega_{\varphi_{k}}^{n}}{n!} \to \int_{N} \log \frac{\omega_{\varphi}^{n}}{\omega_{N}^{n}} \frac{\omega_{\varphi}^{n}}{n!}, (k \to \infty).$$

Now both I - J and $\hat{J}_{-Ric(\omega_N)+\sum(1-\beta_i)\Theta_{h_i}}$ are continuous on \mathcal{E}^1 with respect to the d_1 distance (see, for example, Lemma 5.23 of [11]), so

$$(I-J)(\varphi_k) \to (I-J)(\varphi), \quad \hat{J}_{-Ric(\omega_N)+\sum(1-\beta_i)\Theta_{h_i}}(\varphi_i) \to \hat{J}_{-Ric(\omega_N)+\sum(1-\beta_i)\Theta_{h_i}}(\varphi), \quad (k \to \infty).$$

Finally we need to cope with the terms $\int_{N} \log \|s_i\|_{h_i}^2 \frac{w_{\varphi}^n}{n!}$. For simplicity, we omit the subscript *i* in the following arguments. We shall use the fact the d_1 convergence implies the weak convergence of Monge-Amperè measures (Corollary 5.8 of [10]). So for the continuous function $\log(\|s\|_h^2 + \delta)$ ($\delta > 0$), we have

$$\int_{N} \log(\|s\|_{h}^{2} + \delta) \frac{\omega_{\varphi_{k}}^{n}}{n!} \to \int_{N} \log(\|s\|_{h}^{2} + \delta) \frac{\omega_{\varphi}^{n}}{n!}, \quad (k \to \infty).$$

So we conclude that

$$K_{D,\omega_N}(\varphi) + \sum_i (1-\beta_i) \int_N \left(\log(||s_i||_{h_i}^2 + \delta) - \log ||s_i||_{h_i}^2 \right) \frac{\omega_{\varphi}^n}{n!}$$

$$= \lim_{k \to \infty} \left(K_{D,\omega_N}(\varphi_k) + \sum_i (1-\beta_i) \int_N \left(\log(||s_i||_{h_i}^2 + \delta) - \log ||s_i||_{h_i}^2 \right) \frac{\omega_{\varphi_k}^n}{n!} \right)$$

$$\geq \liminf_{k \to \infty} K_{\beta,\omega_N}(\varphi_k)$$

$$\geq \liminf_{k \to \infty} \epsilon (I-J)(\varphi_k) - C$$

$$= \epsilon (I-J)(\varphi) - C.$$

Finally, let $\delta \to 0$ and use the monotone convergence theorem, we get $K_{D,\omega_N}(\varphi) \ge \epsilon (I - J)(\varphi) - C$. \Box

3. K-ENERGY UNDER GALOIS COVERINGS

Let $f: M \to N$ be a finite Galois covering of degree d with Galois group G with ramification divisor R_f and branching divisor $B_f = \sum_i \left(1 - \frac{1}{d_i}\right) B_i$. To study the behavior of K-energy under Galois coverings, it is convenient to use the degenerate reference form $f^*\omega_N$ as the reference form in the definition of K-energy. This causes no trouble, since for smooth φ satisfying $\omega_{\varphi} := \omega_0 + \sqrt{-1}\partial\overline{\partial}\varphi > 0$, set $\psi := \varphi + \varphi_0$, then for any $\epsilon > 0$ and sufficiently small, we have $f^*\omega_N + \epsilon \sqrt{-1}\partial\overline{\partial}\varphi_0 = (1 - \epsilon)f^*\omega_N + \epsilon \omega_0 > 0$. Since the K-energy satisfies the cocycle conditions, we get

$$\begin{split} K_{\omega_0}(\varphi) &=: \quad \mathcal{M}(\omega_0, \omega_{\varphi}) = \mathcal{M}(\omega_0, f^* \omega_N + \epsilon \sqrt{-1} \partial \overline{\partial} \varphi_0) - \mathcal{M}(\omega_{\varphi}, f^* \omega_N + \epsilon \sqrt{-1} \partial \overline{\partial} \varphi_0) \\ &= \quad K_{\omega_0}(-(1-\epsilon)\varphi_0) - K_{\omega_{\varphi}}(-\psi + \epsilon \varphi_0). \end{split}$$

Using Chen-Tian's formula, it is easy to see that when $\epsilon \to 0$, both terms have finite limits: the entropy term converges by bounded convergence theorem and the convergence of the \hat{J} term is trivial. Then we can write this as a generalized cocycle condition:

$$K_{\omega_0}(\varphi) = K_{\omega_0}(-\varphi_0) + K_{f^*\omega_N}(\psi),$$

where we define $K_{f^*\omega_N}(\psi)$ as

$$K_{f^*\omega_N}(\psi) := -K_{\omega_{\varphi}}(-\psi) := -\lim_{\epsilon \to 0^+} K_{\omega_{\varphi}}(-\psi + \epsilon \varphi_0).$$

So the properness of K_{ω_0} on $\mathcal{H}(\omega_0)$ is equivalent to the properness of $K_{f^*\omega_N}$ on $\mathcal{H}(f^*\omega_N) := \{\psi \in C^{\infty}(M; \mathbb{R}) \mid f^*\omega_N + \sqrt{-1}\partial\overline{\partial}\psi > 0\}.$

The key observation of our work is the following lemma, generalizing Lemma 2.7 of [1] for the Ding-functional and Proposition 6.2 of [21] for the K-energy.

Lemma 3.1. Let $f: M \to N$ be as above. If $\varphi \in \mathcal{H}(M, f^*\omega_N)$ is *G*-invariant, then we have

$$K^M_{f^*\omega_N}(\varphi) = d \cdot K^N_{B_{f,\omega_N}}(\phi), \tag{3.1}$$

where ϕ is the function on N such that $\varphi = f^* \phi$.

Proof. Let us write $\omega_{t\varphi} := f^* \omega_N + t \sqrt{-1} \partial \overline{\partial} \varphi$ and $\omega_{t\phi} := \omega_N + t \sqrt{-1} \partial \overline{\partial} \phi$, then we have

$$\begin{split} K_{f^*\omega_N}^M(\varphi) &= -\int_0^1 dt \int_M \varphi(s(\omega_{t\varphi}) - \underline{s}^M) \frac{\omega_{t\varphi}^n}{n!} \\ &= \int_0^1 dt \int_M \varphi\Big[\sqrt{-1}\partial\overline{\partial}\log \frac{\omega_{t\varphi}^n}{f^*\omega_N^n} + \sqrt{-1}\partial\overline{\partial}\log f^*\omega_N^n\Big] \wedge \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} + \underline{s}^M \int_0^1 dt \int_M \varphi \frac{\omega_{t\varphi}^n}{n!} \\ &= \int_M \log \frac{\omega_{\varphi}^n}{f^*\omega_N^n} \frac{\omega_{\varphi}^n}{n!} + \int_0^1 dt \int_M \varphi\Big[\frac{\underline{s}^M}{n} \omega_{t\varphi} - f^*Ric(\omega_N)\Big] \wedge \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} \\ &+ \sum_j 2\pi r_j \int_0^1 dt \int_{R_j} \varphi \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} \\ &= d\Big(\int_N \log \frac{\omega_{\varphi}^n}{\omega_N^n} \frac{\omega_{\varphi}^n}{n!} + \int_0^1 dt \int_N \varphi\Big[\frac{\underline{s}^M}{n} \omega_{t\varphi} - Ric(\omega_N)\Big] \wedge \frac{\omega_{t\varphi}^{n-1}}{(n-1)!}\Big) \\ &+ \sum_i \sum_{j:f(R_j)=B_i} 2\pi r_j \int_0^1 dt \int_{R_j} \varphi \frac{\omega_{t\varphi}^{n-1}}{(n-1)!}. \end{split}$$

On one hand,

$$\underline{s}^{M} = \frac{2\pi n c_{1}(M) \cdot [f^{*}\omega_{N}]^{n-1}}{[f^{*}\omega_{N}]^{n}} = \frac{2n\pi f^{*}(c_{1}(N) - [B_{f}]) \cdot [f^{*}\omega_{N}]^{n-1}}{[f^{*}\omega_{N}]^{n}}$$

$$= \frac{2n\pi (c_{1}(N) - [B_{f}]) \cdot [\omega_{N}]^{n-1}}{[\omega_{N}]^{n}} = \underline{s}^{N} - \sum_{i} \left(1 - \frac{1}{|\Gamma_{i}|}\right) \frac{2n\pi [B_{i}] \cdot [\omega_{N}]^{n-1}}{[\omega_{N}]^{n}}$$

$$= \underline{s}^{N} - \sum_{i} \left(1 - \frac{1}{|\Gamma_{i}|}\right) \frac{n[\Theta_{h_{i}}] \cdot [\omega_{N}]^{n-1}}{[\omega_{N}]^{n}}.$$

On the other hand, for any R_j such that $f(R_j) = B_i$, suppose $f|_{R_j} : R_j \to B_i$ has degree $d_j \ge 1$. Since f is Galois, both d_j and $r_j = |\Gamma(R_j)| - 1$ depends only on B_i , not on R_j , so we can just write d_i and $r_i = |\Gamma_i| - 1$. Then we have

$$\sum_{i} \sum_{j:f(R_j)=B_i} 2\pi r_j \int_0^1 dt \int_{R_j} \varphi \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} = \sum_{i} 2\pi (|\Gamma_i| - 1) d_i k_i \int_0^1 dt \int_{B_i} \varphi \frac{\omega_{t\varphi}^{n-1}}{(n-1)!},$$

where k_i is the number of different R_j 's mapped onto B_i .

Now a generic point $q \in N$ should have *d* pre-images. Choose a point near B_i , we will have $d_i(r_i + 1) = d_i |\Gamma_i|$ pre-images near R_j , so we get $d = k_i d_i |\Gamma_i|$. So we get

$$\begin{split} \sum_{i} \sum_{j:f(R_{j})=B_{i}} 2\pi r_{j} \int_{0}^{1} dt \int_{R_{j}} \varphi \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} &= \sum_{i} 2\pi (|\Gamma_{i}|-1) \frac{d}{|\Gamma_{i}|} \int_{0}^{1} dt \int_{B_{i}} \varphi \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} \\ &= d \sum_{i} \left(1 - \frac{1}{|\Gamma_{i}|}\right) \int_{0}^{1} dt \int_{N} \phi \left[\sqrt{-1}\partial\overline{\partial}\log\|s_{i}\|_{h_{i}}^{2} + \Theta_{h_{i}}\right] \wedge \frac{\omega_{t\varphi}^{n-1}}{(n-1)!} \end{split}$$

Also note that

$$\int_{0}^{1} dt \quad \int_{N} \phi \sqrt{-1} \partial \overline{\partial} \log \|s_{i}\|_{h_{i}}^{2} \wedge \frac{\omega_{t\phi}^{n-1}}{(n-1)!} = \int_{0}^{1} dt \int_{N} \log \|s_{i}\|_{h_{i}}^{2} \frac{d}{dt} \frac{\omega_{t\phi}^{n}}{n!}$$
$$= \left(\int_{N} \log \|s_{i}\|_{h_{i}}^{2} \frac{\omega_{t\phi}^{n}}{n!}\right)_{t=0}^{t=1} = \int_{N} \log \|s_{i}\|_{h_{i}}^{2} \left(\frac{\omega_{\phi}^{n}}{n!} - \frac{\omega_{N}^{n}}{n!}\right)$$

From these identities we at once get (3.1).

4. Proof of Theorem 1.1

The following theorem is essentially the same as Chen-Cheng [8]. Since it is not stated explicitly, we include a sketched proof here for the convenience of the readers:

Theorem 4.1. If *K*-energy is proper on the space of *G*-invariant Kähler potentials for a finite subgroup G of Aut(N, J), then there is a cscK metric.

Proof. We follow the same strategy as Tian did in [27], using Chen-Cheng's continuity method. Start with a *G*-invariant Kähler form ω_0 in the given *G*-invariant Kähler class. We use Chen's continuity path proposed in [7]:

$$t(s(\omega_{\varphi}) - \underline{s}) = (1 - t)(tr_{\omega_{\varphi}}\omega_0 - n).$$

In [7], Chen proved that the set of $t \in (0, 1)$ such that the above equation is solvable is open, and the openness at t = 0 is proved later by Hashimoto [18] and Zeng [29]. Their arguments work also for *G*-invariant function spaces when *G* is finite. Then we apply Chen-Cheng's *a priori* estimates in [8] to get the closedness.

Given this theorem, to prove Theorem 1.1, it suffices to prove the properness of $K_{f^*\omega_N}^M$ on the space of *G*-invariant potentials $\mathcal{H}^G(M, f^*\omega_N)$. In turn, by Lemma 3.1 and 2.1, it suffices to prove the properness of K_{B_f,ω_N}^N on the space of \mathcal{E}^1 potentials $\mathcal{E}^1(N, \omega_N)$ with $B_f = \sum_i (1 - \beta_i) B_i := \sum_i (1 - \frac{1}{|\Gamma_i|}) B_i$. Finally by Lemma 2.2, it suffices to prove the properness of log-K-energy on $\mathcal{H}(N, \omega_N)$. The idea, as in [22] is again to use the log- α -invariant:

$$\alpha(\omega_N, D) := \sup\{\alpha > 0 \Big| \int_N e^{-\alpha(\varphi - \sup\varphi)} \frac{\omega_N^n}{\prod_i |s_i|_{h_i}^{2(1-\beta_i)} n!} \le C_\alpha \quad \forall \varphi \in PSH(N, \omega_N) \}.$$

Also one can prove that $\alpha(\omega_N, D) > 0$.

We have the following lemma, which is first observed by K. Zheng (Proposition 4.2 of [31]) when D is simple normal crossing. The proof here is essentially the same. For the convenience of readers, we include a sketched proof here.

Lemma 4.2. Let $D = \sum_{i} (1 - \beta_i) D_i$. Assume we can find a constant $\epsilon \ge 0$ such that

(1) $\alpha(\omega_N, D) > \frac{n\epsilon}{n+1}$;

(2) $2\pi(K_N + D) + \epsilon[\omega_N] > 0;$ (3) $2\pi - (K_N + D)$

$$\Big(\frac{2n\pi(K_N+D)\cdot[\omega_N]^{n-1}}{[\omega_N]^n}+\epsilon\Big)[\omega_N]>2\pi(n-1)(K_N+D),$$

then K_{D,ω_N}^N is proper.

Proof. Choose $\frac{n\epsilon}{n+1} < \alpha < \alpha(\omega_N, D)$. Then by definition, we have

$$C_{\alpha} \geq \int_{N} e^{-\alpha(\varphi - \sup\varphi)} \frac{\omega_{N}^{n}}{\prod_{i} |s_{i}|_{h_{i}}^{2(1-\beta_{i})} n!}$$

=
$$\int_{N} e^{-\alpha(\varphi - \sup\varphi) - \sum_{i} (1-\beta_{i}) \log |s_{i}|_{h_{i}}^{2} - \log \frac{\omega_{\varphi}^{n}}{\omega_{N}^{n}}} \frac{\omega_{\varphi}^{n}}{n!}.$$

By Jensen inequality, we get

$$\int_{N} \log \frac{\omega_{\varphi}^{n}}{\omega_{N}^{n}} \frac{\omega_{\varphi}^{n}}{n!} + \sum_{i} (1 - \beta_{i}) \int_{N} \log |s_{i}|_{h_{i}}^{2} \frac{\omega_{\varphi}^{n}}{n!}$$

$$\geq \alpha \int_{N} (\sup \varphi - \varphi) \frac{\omega_{\varphi}^{n}}{n!} - C$$

$$\geq \alpha \int_{N} \varphi \left(\frac{\omega_{N}^{n}}{n!} - \frac{\omega_{\varphi}^{n}}{n!} \right) - C = \alpha I_{\omega_{N}}(\varphi) - C$$

$$\geq \frac{(n+1)\alpha}{n} (I - J)(\varphi) - C.$$

Using the above inequality, we have

$$K_{D,\omega_N}^N(\phi) \ge \Big(\frac{(n+1)\alpha}{n} - \epsilon\Big)(I-J)(\varphi) + \hat{J}_{-Ric(\omega_N) + \sum_i (1-\beta_i)\Theta_{h_i} + \epsilon\omega_N,\omega_N}(\varphi) - C.$$

Write $\chi_{\epsilon} := -Ric(\omega_N) + \sum_i (1 - \beta_i)\Theta_{h_i} + \epsilon \omega_N$, then condition (2) implies $[\chi_{\epsilon}] > 0$. Now condition (3) implies the solvability of the corresponding *J*-equation, so $\hat{J}_{\chi_{\epsilon},\omega_N}$ is bounded from below. \Box

Remark 4.3. In the above proof, we can write instead

$$K_{D,\omega_N}^N(\phi) \ge \left(\frac{(n+1)\alpha}{n}\right)(I-J)(\varphi) + \hat{J}_{-Ric(\omega_N) + \sum_i (1-\beta_i)\Theta_{h_i},\omega_N}(\varphi) - C$$

Set $\theta := -Ric(\omega_N) + \sum_i (1 - \beta_i) \Theta_{h_i}$, then $[\theta] = 2\pi(K_N + D)$. Using the work of Sjöström Dyrefelt [24], we have

$$\sup\{\delta \in \mathbb{R} | \hat{J}_{\theta,\omega_N} \ge \delta(I-J)_{\omega_N} - C_{\delta}\} \ge \min\left\{\mathcal{T}([\theta], [\omega_N]), \inf_{V} \frac{C_{\theta,\omega_N} \int_{V} \omega_N^p - p \int_{V} \theta \wedge \omega_N^{p-1}}{(n-p) \int_{V} \omega_N^p}\right\}$$

where $\mathcal{T}([\theta], [\omega_N]) := \sup\{\delta \in \mathbb{R} | [\theta] - \delta[\omega_N] \ge 0\}, C_{\theta, \omega_N} := \frac{n \int_N \theta \wedge \omega_N^{n-1}}{\int_N \omega_N^n}$ and the inf is taken with respect all subvarieties of *N* of dimension $p \le n-1$.

Then Lemma 4.2 can be strengthened to: If

$$\frac{n+1}{n}\alpha(\omega_N,D) + \min\left\{\mathcal{T}([\theta],[\omega_N]), \inf_V \frac{C_{\theta,\omega_N} \int_V \omega_N^p - p \int_V \theta \wedge \omega_N^{p-1}}{(n-p) \int_V \omega_N^p}\right\} > 0,$$

where $[\theta] = 2\pi(K_N + D)$, then K_{D,ω_N}^N is proper. In view of K. Zhang's work on the analytic δ -invariant, we can even replace the log- α invariant by certain δ -invariant [31].

Now we can finish the proof of Theorem 1.1:

Proof of Theorem 1.1: By Lemmma 4.2, the assumptions in Theorem 1.1 imply that $K_{B_{\delta_0},\omega_N}$ is proper on the space of smooth Kähler potentials. By the linearity of K_{B_{δ},ω_N} on δ , we get, for $0 < \delta \le \delta_0$,

$$K_{B_{\delta},\omega_N} = \frac{\delta}{\delta_0} K_{B_{\delta_0},\omega_N} + (1 - \frac{\delta}{\delta_0}) K_N$$

Since K_N is bounded from below, K_{B_{δ},ω_N} is proper on the space of smooth Kähler potentials for all $\delta \in (0, \delta_0]$. Take $\delta = 1$, we get the properness of K_{B_f,ω_N} on the space of smooth Kähler potentials. This implies the properness of K_M on the space of smooth *G*-invariant Kähler potentials and hence the existence of cscK metrics in $[f^*\omega_N]$.

5. Proof of Theorem 1.5

Now let $f : M \to N$ be a cyclic Galois covering of order *d* branched over a smooth hypersurface $B \subset N$, where both *M* and *N* are project algebraic manifolds of dimension *n*. Moreover, we assume that *N* has a cscK metric with $[\omega_N] = \lambda 2\pi c_1(B)$ for some $\lambda > 0$. We can make use of the following estimate of Berman [3] for the log- α -invariant:

Lemma 5.1 ([3] Proposition 6.2). Let N be a projective algebraic manifold with $B \subset N$ a smooth hypersurface. Let $0 < \beta < 1$, then we have

$$\alpha(2\pi c_1(B), (1-\beta)B) \ge \min\{\beta, \alpha(2\pi c_1(B)), \alpha(2\pi c_1(B)|_B)\}.$$

From this we get

$$\alpha([\omega_N], (1-\beta)B) \ge \min\{\frac{\beta}{\lambda}, \alpha([\omega_N]), \alpha([\omega_N]|_B)\}.$$
(5.1)

To show the properness of $K_{(1-\beta)B,\omega_N}$, we apply Lemma 4.2: we need to find a number $\epsilon \ge 0$ such that

(1) $\alpha(\omega_N, (1-\beta)B) > \frac{n\epsilon}{n+1};$ (2) $K_N + (1-\beta+\lambda\epsilon)B > 0;$ (3) $\left(\frac{nK_N \cdot B^{n-1}}{B^n} + 1 - \beta + \lambda\epsilon\right)B - (n-1)K_N > 0.$ When $\beta > 0$ is sufficiently small, for condition (1), we only need to choose ϵ such that

$$\frac{n\epsilon}{n+1} < \frac{\beta}{\lambda}.$$

So in case both $K_N + B$ and

$$\Big(\frac{nK_N\cdot B^{n-1}}{B^n}+1\Big)B-(n-1)K_N$$

are nef, we can take ϵ slightly bigger than $\frac{\beta}{\lambda}$ such that both (2) and (3) are true. This proves that $K_{(1-\beta)B,\omega_N}$ is proper for sufficiently small $\beta > 0$. On the other hand, when $\beta = 1$, K_{ω_N} is bounded from below. The remaining arguments are identical to the proof of Theorem 1.1.

6. Applications and Examples

6.1. **Existence of covers.** In order to discuss the range of applicability of our results, the first point to address is certainly the existence problem for Galois covers between smooth complex manifolds.

The simplest and most classical situation is the so called "simple cyclic cover" (see e.g. Lazarsfeld [19] Proposition 4.1.6): let *N* be a smooth Kähler manifold and *L* a holomorphic line bundle over *N*, with total space **L**. Suppose there is a holomorphic section $\sigma \in H^0(N, O(L^{\otimes d}))$ ($d \in \mathbb{N}$ and $d \ge 2$) such that ($\sigma = 0$) is a smooth prime divisor. Then one can define

$$M := \{ (p, v) \in \mathbf{L} \mid v^{\otimes d} = \sigma(p) \}.$$

Then it is easy to check that *M* is a smooth Kähler manifold and the restriction of the projection map of **L** is a cyclic cover of degree *d*, branched along ($\sigma = 0$).

One should note concerning this type of construction that the smoothness of ($\sigma = 0$) suffices to prove smoothness of M, but it is clearly in general not necessary. Moving to the general abelian case this relationship has been well understood. Indeed on one hand, given a branched covering $f : M \to N$ with M and N smooth, and abelian Galois group Gal(f), Catanese ([5, Proposition 1.1]) proved that both the ramification divisor R_f and the branching divisor B_f are simple normal crossing.

Viceversa, given a smooth base N and a simple normal crossing divisor B in N, Bloch-Gieseker-Kawamata proved the existence of a smooth abelian Galois cover with *some* Galois group G. (see e.g. Lazarsfeld [19] Proposition 4.1.12 and Theorem 1.1.1 in Kawamata-Matsuda-Matsuki [16]).

A complete general theory for the existence of Galois covers with preassigned abelian Galois group G has been developed in the seminal paper by Pardini [23].

Concerning the general abelian case, let us observe that thanks to the above mentioned result by Catanese, the inertia group G_j of each component R_j of R_f , i.e. the intersection $G_j = \bigcap_{x \in R_j} G_x$ of all the isotropy groups of points of R_j , is a cyclic group. Also, the isotropy group $G_x = \{g \in G : gx = x\}$ of a point $x \in M$ is the direct product of the inertia groups of the irreducible components of R_f passing through *x*. Furthermore, note that given a component B_i of B_f , all the components of $f^{-1}(B_i)$ have the same inertia group since *G* is abelian.

As a consequence, one can deduce that an abelian Galois covering $f : M \to N$ factorizes as a composition of branched cyclic coverings and an unramified abelian covering (as indeed happens for the above mentioned Kawamata's coverings), a result which seems to be of independent interest and makes the production of further cscK examples straightforward.

Proposition 6.1. Let $f : M \to N$ be an abelian Galois covering, and put $M_0 = M$. There exist cyclic subgroups $G_1, \ldots, G_\ell \subset Gal(f)$ together with Galois G_j -coverings $f_j : M_{j-1} \to M_j$ for all $j \in \{1, \ldots, \ell\}$, and an unramified \hat{G} -covering $\hat{f} : M_\ell \to N$, where $\hat{G} = Gal(f) / \prod_{j=1}^{\ell} G_j$, such that

$$f = \hat{f} \circ f_{\ell} \circ \cdots \circ f_{1}.$$

Proof. Let $K = \prod_{x \in M} G_x \subset G$ be the subgroup generated by all the isotropy groups for the action of *G* on *M*. Since *K* is a finite abelian group, there exists a subset $S \subset K$ such that $K = \bigoplus_{g \in S} \langle g \rangle$, where $\langle g \rangle \simeq \mathbb{Z}_{\operatorname{ord}(g)}$ is the cyclic group generated by *g*, and the order $\operatorname{ord}(g)$ of *g* is a power of a prime.

Pick $g \in S$, and let $\operatorname{ord}(g) = p^m$ for some prime p and integer m. Note that, in general, $Fix(g) = \{x \in M : gx = x\}$ may be empty. On the other hand, since $\langle g \rangle$ is a direct summand of K, it has a subgroup that fixes some point of M. Therefore, there is a maximal m' < m such that $g' = g^{p^{m'}}$ fixes some point of M. As a consequence, Fix(g') is a smooth, non-necessarily connected, codimension one submanifold of M. Moreover, $\langle g' \rangle$ acts freely on $M \setminus Fix(g')$. Thus, for all $x \in Fix(g')$ there is a $\langle g' \rangle$ -invariant neighborhood U with local coordinates z_1, \ldots, z_n such that $Fix(g') \cap U$ is the locus $z_1 = 0$, and the action of $\langle g' \rangle$ on U is generated by $z \mapsto (e^{2\pi i/p^{m-m'}}z_1, z_2, \ldots, z_n)$. Therefore, the quotient $M/\langle g' \rangle$ is smooth and the projection from M is a $\langle g' \rangle$ -covering ramified along Fix(g'). Note that $M/\langle g' \rangle$ is acted on freely by the quotient group $\langle g \rangle/\langle g' \rangle$. Therefore, letting $G_1 = \langle g \rangle$, and $M_1 = M_0/G_1$ yields a cyclic covering $f_1 : M_0 \to M_1$ with Galois group G_1 .

On the other hand, since *G* is abelian, the action of *G* on *M* descends to an action of $H_1 = G/G_1$ on M_1 . Clearly one has $N = M_1/H_1$ and, denoted by $p_1 : M_1 \to N$ the projection on the quotient, one can factorize $f = p_1 \circ f_1$. Note that p_1 is a Galois covering with Galois group H_1 . Now, the subset of H_1 generated by all the isotropy groups for the action of H_1 on M_1 is isomorphic to $\bigoplus_{g \in S_1} \langle g \rangle$ where S_1 is *S* without the generator of G_1 .

Applying the same argument as above to the covering p_1 one eventually gets a cyclic G_2 -covering $f_2: M_1 \to M_2$ and a H_2 -covering $p_2: M_2 \to N$, where $H_2 = G/(G_1 \times G_2)$, that satisfies $p_1 = p_2 \circ f_2$. Hence $f = p_2 \circ f_2 \circ f_1$.

Let $\ell = |S|$. Iterating this argument until $j = \ell$, and letting $\hat{f} = p_{\ell}$ and $\hat{G} = H_{\ell}$, then yields the thesis.

When we turn to Galois non-abelian coverings, the possibility of constructing such coverings is much more restricted than for cyclic and abelian ones, and indeed it has been subject of intensive research by many authors for different type of groups (see e.g. [5], [6], [14], [20]). In the last subsection of this paper we will focus on the example of the Dihedral group, where thanks to the fundamental work of Catanese-Perroni [6], we have a fairly complete understanding. In particular we observe that such covers do exist in abundance even with very singular branch divisors.

6.2. **KE and cscK metrics: the abelian case.** Given the great flexibility in constructing cyclic Galois covers described above, it is also clear that the range of applicability of Theorem 1.5 is extremely large.

Let us first observe that we can recover immediately the known results for Kähler-Einstein manifolds and cyclic covers proved in Theorem 6.1 in [21] (see also Theorem 2.5 in [1]).

Indeed, let us assume that *N* is Fano with Kähler-Einstein metric ω_N and assume $B \in |-\beta K_N|$ (note that this β is not the one in [1]). It is then immediate to check that Theorems 1.5 and 1.1 apply if $\beta \ge 1$, hence providing Kähler-Einstein metrics on the covering space *M*. Note that *M* will be Fano again iff $d < \frac{\beta}{\beta-1}$ (which is the condition appearing in Theorem 2.5 of [1]). This application extends the range of application of Theorem 6.1 in [21] where *B* is supposed to be smooth and irreducible (and similar results in [1]).

Even in this simplest situation it is worth observing how our results extend Corollary 2.7 in Cheng-Chen [8]: let us indeed consider as a simple example cyclic coverings of $N = \mathbb{P}^n$, |G| = d and $B \in |\lambda H|$, where *H* is the hyperplane bundle. In this case, Corollary 1.2 applies.

Our first condition reads:

$$K_N + (1 - \beta_0)B = -(n+1)H + \lambda(1 - \beta_0)H \ge 0$$

As for the second one, having fixed $\omega_N = \mu \omega_{FS}$, i.e. $[\omega_N] = \mu c_1(H)$, it is immediate to check that we get again the same condition as above.

Therefore, we can conclude that

$$\lambda > n+1$$

in order to get a cscK metric on M. (Note that by Theorem 1.5 one gets the better condition $\lambda \ge n + 1$). On the other hand, Cheng-Chen require

$$\lambda \ge (n+1)\frac{d}{d-1}.$$

In particular for all cyclic coverings of \mathbb{P}^n with $d \le n + 2$ we can find an integer λ which falls into the range of our results and not in Cheng-Chen's.

In fact, Chen-Cheng's Corollary 2.7 is simply a combination of their main theorem and previous works (e.g. [22]) on the properness of K-energy. It is easy to check that a direct application of

Chen-Cheng in our setting gives precisely the $\delta = 1$ case of Theorem 1.1. So our theorems are both theoretical and practical improvements of Chen-Cheng's Corollary 2.7 in our setting.

Let us now illustrate one simple example, where the Kaähler class on the base manifold is not taken to be necessarily KE and yet produces via our Theorems also new KE metrics: fix as base space

$$(N, \omega_N) = (\mathbb{P}^1 \times \mathbb{P}^1, \omega),$$

where $\omega \in c_1(L)$, $L = a_1C_1 + a_2C_2$, $a_1, a_2 > 0$, and C_i denotes the homology class of the *i*-th factor \mathbb{P}^1 .

Set $B \in |d(b_1C_1 + b_2C_2)|$ the (smooth) branching divisor. In order for this data to satisfy our setup we need to assume that $b_i > 0$. It is then an elementary computation to check that conditions of Corollary 1.2 are satisfied iff $db_1 > 2$ and $db_2 > 2$. Moreover, whenever (a_1, a_2) is proportional to (b_1, b_2) , the conditions of Theorem 1.5 reduce to $db_1 \ge 2$, $db_2 \ge 2$, thus adding the case d = 2, $b_1 = b_2 = 1$, $a_1 = a_2$ to the ones given by Corollary 1.2.

Let us remark on this example that the scalar curvature of the cscK metric ω_M is also easily computed by

$$s_M = 2 \cdot \frac{L \cdot (-K_M)}{L^2} = (2 - (d - 1)b_1)a_2 + (2 - (d - 1)b_2)a_1,$$

which is always non positive, except for the trivial case d = 1 and the case d = 2, $b_1 = b_2 = 1$, $a_1 = a_2$. Moreover, s_M vanishes if and only if (d = 3 and $b_1 = b_2 = 1$) or $d = b_1 = b_2 = 2$.

Note that this construction has 4 free parameters (plus the group *G*): b_1 , b_2 upon which the geometry of the resulting manifold *M* depends, and a_1 , a_2 which control the Kähler class of ω_M , showing in practice the large flexibility of our construction in this case. One can check that ω_M is KE if and only if (a_1, a_2) and $(2 - (d - 1)b_1, 2 - (d - 1)b_2)$ are linearly dependent. Note that this implies one can get KE metrics on the covering even starting from cscK not KE on the base (e.g. $a_1 = 1, a_2 = 2, b_1 = 2, b_2 = 3, d = 3$).

6.3. **KE and cscK metrics: the non-abelian case.** As a prototype example, in this Section we focus on the case of Galois coverings with structure group D_p , the dihedral group of order 2p, whose existence has been largely studied in the above mentioned works. We assume hereafter that $p \ge 3$ so that D_p is not abelian.

As described by Catanese-Perroni, in the theory of D_p -Galois coverings the parameter B is "a posteriori" determined by the choice of a line bundle L in such a way that there exists sections $a \in H^0(N, O_N(pL))$ and $f \in H^0(N, O_N(2L))$, such that the following conditions are satisfied (see [6], Section 6 and Theorem 6.1):

- (1) the zero-locus of $a^2 f^p$ is smooth in the open subset where $f \neq 0$;
- (2) the divisors $A := \{a = 0\}$ and $F := \{f = 0\}$ intersect each other transversely;
- (3) $A \cap F \neq \emptyset$;

A beautiful result (Theorem 6.1 of [6]) by Catanese-Perroni shows how with these data at hand one can construct a D_p -Galois covering $\pi_L \colon M_L \to N$ ramified along the locus $\{a^2 - f^p = 0\} \subset N$. It is interesting to observe that in this case the ramification divisor has p components all with inertia group \mathbb{Z}_2 , while the branch locus has one component for odd and two for even p respectively.

We want to apply this result in few significant cases, in order to analyze the possibility of applying our Theorem 1.1. From now on G is assumed to be the dihedral group D_p .

(1) Let us start again with $N = \mathbb{P}^n$ and we consider $L = \lambda H$ which implies $B \in |2p\lambda H|$, where *H* is the hyperplane bundle. In this case we also assume for simplicity $[\omega_N] = c_1(H)$.

This implies that in this situation $B_{\delta} \sim \delta p \lambda H$, hence our first condition reads

$$K_N + B_\delta \sim (\delta p \lambda - n - 1) H \ge 0$$
,

while the second

$$\frac{n(K_N+B_{\delta})\cdot[\omega_N]^{n-1}}{[\omega_N]^n}[\omega_N]-(n-1)(K_N+B_{\delta})\geq 0,$$

hence both reduce to

$$\delta p\lambda \ge n+1$$
,

for some $\delta \in [1, 2)$.

As for the cyclic case described above, it is easy to observe that there are plenty of situations where our result applies while Chen-Cheng's doesn't (the simplest being n = 2, p = 3 and $\lambda = 1$, or n = p = 3 and $\lambda = 1$).

Yet, all these examples fall in the category of Kähler-Einstein manifolds, but again, contrary to Li-Sun, we do not have to assume smoothness and irreducibility of the branch locus.

(2) In order to construct csck non KE metrics we can apply Catanese-Perroni's theory taking as base manifold $\mathbb{P}^n \times \mathbb{P}^n$ with $L = H_1 + H_2$, having denoted by H_i the hyperplane bundle on the *i*-th factor. We also choose $\omega_N \in b_1c_1(H_1) + b_2c_1(H_2)$.

The first condition

$$K_N + B_\delta \sim (\delta p - n - 1)H_1 + (\delta p - n - 1)H_2 \ge 0$$

translates into $\delta p \ge n + 1$, while the second

$$\frac{2n(K_N+B_{\delta})\cdot[\omega_N]^{2n-1}}{[\omega_N]^{2n}}[\omega_N] - (2n-1)(K_N+B_{\delta}) \ge 0$$

becomes

$$(n\frac{b_1}{b_2} - (n-1))(\delta p - n - 1)H_1 + (n\frac{b_2}{b_1} - (n-1))(\delta p - n - 1)H_2 \ge 0$$

We then have to require either

$$\delta p = n+1$$
 or $\begin{cases} \delta p > n+1\\ n\frac{b_1}{b_2} - (n-1) \ge 0\\ n\frac{b_2}{b_1} - (n-1) \ge 0 \end{cases}$

for some $\delta \in [1, 2)$. Equivalently, either $1 \le \frac{n+1}{p} < 2$ or

$$\begin{cases} p > \frac{n+1}{2} \\ \frac{n-1}{n}b_2 \le b_1 \le \frac{n}{n-1}b_2 \end{cases}$$

Note that the above system gives large families of new cscK manifolds for any *n* (not KE for $b_1 \neq b_2$) and these were not covered by Chen-Cheng's Theorem for $n \ge 3$.

(3) Let us end this Section by applying our results and Catanese-Perroni's construction for dihedral D_p -Galois covering to the case when $N = Bl_{P_1,P_2,P_3}\mathbb{P}^2$ is the the projective plane blown-up at three non-collinear points P_i . We denote by $\sigma: N \to \mathbb{P}^2$ the blow-up at $\{P_1, P_2, P_3\}$, and

$$L = kH - \sum_{i=1}^{3} a_i E_i,$$

where *H* is the pull-back via σ of any line not passing through the points P_i 's, $E_i = \sigma^{-1}(P_i)$ is the exceptional curve over P_i for all i = 1, 2, 3, and k, a_1, a_2, a_2 are integers.

Since $Bl_{P_1,P_2,P_3}\mathbb{P}^2$ is toric, *L* is very ample if and only if it is ample [9, Theorem 6.1.15]. On the other hand, whenever 2*L* is very ample, a simple application of Bertini's Theorem gives the existence of the seeked holomorphic sections $a \in H^0(N, O_N(pL))$ and $f \in H^0(N, O_N(2L))$ which satisfy the above requirements (the transversality property required follows at once from [17, Lemma V,1.2]). Summarizing, in order to apply [6], the requirement is just that *L* is ample. As a strightforward application of the Nakai-Moishezon Criterion (see e.g. [17]) this is equivalent to

$$a_1, a_2, a_3 > 0$$
, and $k > a_i + a_j$ $\forall i \neq j$

Having now $\pi_L \colon M_L \to N$, branched along $B \in |2pL|$, we have to choose the cscK class $[\omega_N]$. We can make (at least) two very different, but equally interesting, choices:

- $[\omega_N] = -K_N$ and ω_N is the Kähler-Einstein metric found by Siu [26] and Tian-Yau [28].
- $[\omega_N] = F$ where $F = rH \sum_{i=1}^{3} E_i$, with *r* positive and sufficiently large so that *F* contains a cscK metric via Theorem 1.3 in [2].

Clearly, we can treat these cases simultaneously denoting by $[\omega_N] = rH - \sum_{i=1}^{3} E_i$, where r = 3 gives the Siu-Tian-Yau case, while $r \gg 3$ gives the Arezzo-Pacard one.

When is $K_N + B_{\delta}$ nef?

$$K_N + B_\delta = (\delta pk - 3)H - \sum_{i=1}^3 (\delta pa_i - 1)E_i,$$

hence the nef condition translates into $k \ge (a_i + a_j) + \frac{1}{\delta p}$ for all $i \ne j$. Equivalently, thanks to the fact that k and a_i 's are integers, $p \ge 3$, and $1 \le \delta < 2$, nef condition reduces to

$$k \ge a_i + a_j + 1 \qquad \forall i \ne j. \tag{6.1}$$

When is $\frac{2(K_N+B_{\delta})\cdot[\omega_N]}{[\omega_N]^2}[\omega_N] - (K_N + B_{\delta})$ nef? Notice that

$$\frac{2(K_N + B_{\delta}) \cdot [\omega_N]}{[\omega_N]^2} [\omega_N] = 2 \frac{r(\delta pk - 3) - \sum_{i=1}^3 (\delta pa_i - 1)}{r^2 - 3} (rH - \sum_{i=1}^3 E_i).$$

Given the number of parameters involved let us first fix r = 3 hence looking at Siu-Tian-Yau's KE metric on the base. In this case $\frac{2(K_N + B_{\delta}) \cdot [\omega_N]}{[\omega_N]^2} [\omega_N] - (K_N + B_{\delta})$ reduces to

$$[\delta p(2k - \sum_{j=1}^{3} a_j) - 3]H - \sum_{i=1}^{3} [\delta p(k - a_i - \frac{1}{3}\sum_{j=1}^{3} a_j) - 1]E_i$$

hence the nef condition becomes $\delta p(a_i + a_j - \frac{1}{3}\sum_{\ell=1}^3 a_\ell) \ge 1$ for all $i \ne j$, or equivalently $a_i \le -\frac{1}{\delta p} + \frac{2}{3}\sum_{\ell=1}^3 a_\ell$ for all $i \in \{1, 2, 3\}$. Since all a_i 's are integers and $p \ge 3$, this condition is satisfied for some $\delta \in [1, 2)$ whenever

$$a_i \leq \frac{1}{3} \sum_{\ell=1}^{3} (2a_\ell - 1), \quad \forall i \in \{1, 2, 3\}.$$

As a consequence, as soon a_i 's satisfy this condition and k is chosen so that L is ample, M admits a cscK metric in $\pi_L^*(-K_N)$ thanks to Theorem 1.1.

Going back to the adiabatic cscK classes given by $r \gg 3$, let us analyze for simplicity the case $a_i = 1$ for all *i*, yet is clearly not a necessary condition for our Theorem to apply.

In this case, $\frac{2(K_N+B_{\delta})\cdot[\omega_N]}{[\omega_N]^2}[\omega_N] - (K_N + B_{\delta})$ becomes

$$\frac{\delta p(r^2k+3k-6r)-3(r^2-2r+3)}{r^2-3}H - \sum_{i=1}^3 \frac{\delta p(2rk+3-r^2-6)+r^2-6r+3}{r^2-3}E_i$$

and it is nef if and only if the coefficient of H is not smaller than the coefficient of E_i , or equivalently

$$(k+2)r^2 - (4k+6)r + 3k + 6 \ge \frac{5r^2 - 18r + 15}{\delta p}.$$

This inequality is satisfied for some $\delta \in [1, 2)$ whenever $k \ge 4$ and $r \gg 3$, hence providing new large families of csck manifolds by Theorem 1.1.

Finally notice that in all cases discussed above, under assumption of generality of the parameters involved, cscK metric constructed on M are not KE. In order to see this, note that by Catanese-Perroni's construction one has $K_M = \pi_L^*(K_N + pL)$. On the other hand, the cscK metric we construct over M represents the class of π^*F , thus it is KE if and only if there is a real constant λ such that $K_N + pL = \lambda F$. In turns, this is equivalent to

$$(pk-3-\lambda r)H - \sum_{i=1}^{3} (pa_i-1-\lambda)E_i = 0$$
, for some $\lambda \in \mathbf{R}$.

Therefore, in order to make sure that the cscK metric we constructed on M is not KE, the parameters must satisfy:

- $k \neq 3a_i$ for some *i* (Siu-Tian-Yau case, i.e. r = 3),
- $k \neq \frac{3 + (pa_i 1)r}{p}$ for some *i* (Arezzo-Pacard case, i.e. $r \gg 3$).

It is then clear that the cscK metric we constructed on *M* is never KE in the Arezzo-Pacard case up to taking *r* sufficiently large, once *k* and the a_i 's are fixed. On the other hand, that metric is KE in the Siu-Tian-Yau case if and only if $k = 3a_1 = 3a_2 = 3a_3$.

We end up this subsection by noting that for all coverings of $Bl_{P_1,P_2,P_3}\mathbb{P}^2$ discussed here does not apply a result of Chen-Cheng stating that any complex surface with negative first Chern class and no curves with negative self-intersection admits a cscK metric in any Kähler class [8, Corollary 1.7]. Indeed, our examples are surfaces having negative first Chern class since $K_M = \pi_L^*(K_N + pL)$, and

$$K_N + pL = (pk - 3)H - \sum_{i=1}^3 (pa_i - 1)E_i$$

is ample since $k > a_i + a_j + \frac{1}{p}$ for all $i \neq j$. On the other hand, the finiteness of the map π_L implies that pulling back any (-1)-curve of $Bl_{P_1,P_2,P_3}\mathbb{P}^2$ yields a curve with negative self-intersection on M.

References

- [1] Arezzo, C., Ghigi, A., and Pirola, G. Symmetries, quotients and Kähler-Einstein metrics, J. reine angew. Math. 591 (2006), 177-200.
- [2] Arezzo, C., Pacard, F., *Blowing up Kähler manifolds with constant scalar curvature. II.* Ann. of Math. (2) 170 (2009), no. 2, 685–738.
- [3] Berman, R. A thermodynamical formalism for Monge-Ampère equations, Moser-Trudinger inequalities and Kähler-Einstein metrics. Adv. Math. 248 (2013), 1254-1297.
- [4] Berman, R., Darvas, T., and Lu, C. *Convexity of extended K-energy and the large time behavior of the weak Calabi flow*, Geometry and Topology, 21 (2017), 2945-2988.
- [5] Catanese, F. On the moduli spaces of surfaces of general type, J. Differential Geometry, 19 (1984), 483-515.

- [6] Catanese, F., Perroni, F., *Dihedral Galois covers of algebraic varieties and the simple cases*, J. Geom. Phys. 118 (2017), 67-93.
- [7] Chen, X.X. On the existence of constant scalar curvature Kähler metrics: a new perspective, Ann. Math. Qué. 42 (2018), no. 2, 169-189.
- [8] Chen, X.X. and Cheng, J. On the constant scalar curvature Kähler metrics (II)-Existence results, J. Amer. Math. Soc. 34 (2021), no. 4, 937-1009.
- [9] Cox, D. A., Little, J. B., and Schenck, H. K. *Toric varieties*. Graduate Studies in Mathematics, 124. American Mathematical Society, Providence, RI, 2011.
- [10] Darvas, T. The Mabuchi geometry of finite energy classes, Adv. Math. 285 (2015), 182-219.
- [11] Darvas, T. and Rubinstein, Y. *Tian's properness conjectures and Finsler geometry of the space of Kähler metrics*, Journal of the AMS. 30 (2017), 347-387.
- [12] Demailly, J.P. and Paun, M. Numerical characterization of the Kähler cone of a compact Kähler manifold, Ann. of Math. (2) 159 (2004), no. 3, 1247-1274.
- [13] Dervan, R. On K-stability of finite covers, Bull. Lond. Math. Soc. 48 (2016), no. 4, 717-728.
- [14] Fantechi B. and Pardini, R., Automorphisms and moduli spaces of varieties with ample canonical class via deformations of abelian covers. Comm. Algebra 25 (1997), no. 5, 1413–1441.
- [15] Ghigi, A. and Kollár, J. Kähler-Einstein metrics on orbifolds and Einstein metrics on spheres, Comment. Math. Helv. 82 (2007), 877-902.
- [16] Y. Kawamata, K. Matsuda and K. Matsuki, *Introduction to the Minimal Model Problem*, Advanced Studies in Pure Mathematics 10, 1987 Algebraic Geometry, Sendai, 1985
- [17] Hartshorne, R., Algebraic Geometry, Springer-Verlag, Berlin, 1977.
- [18] Hashimoto, Y. Existence of twisted constant scalar curvature Kähler metrics with a large twist, Math. Zeit. 292 (2019), 791-803.
- [19] R. Lazarsfeld. Positivity in Algebraic Geometry, I, Springer-Verlag, Berlin, 2004.
- [20] M. Manetti Iterated double covers and connected components of moduli spaces. Topology 36 (1997), no. 3, 745–764.
- [21] Li, C. and Sun, S. Conical Kähler-Einstein metrics revisited, Comm. Math. Phys. 331 (2014), no. 3, 927-973.
- [22] Li, H., Shi, Y. and Yao, Y. A criterion for the properness of the K-energy in a general Kähler class, Math. Ann. 361 (2015), no. 1-2, 135-156.
- [23] Pardini, R., Abelian covers of algebraic varieties, J. Reine Angew. Math. 417 (1991), 191-213.
- [24] Sjöström Dyrefelt, Z. Optimal lower bounds for Donaldson's J-functional, Advances in Mathematics, 374 (2020), 107271.
- [25] Song, J. and Weinkove, B. On the convergence and singularities of the J-flow with applications to the Mabuchi energy, Comm. Pure Appl. Math. 61 (2008), no. 2, 210-229.
- [26] Siu, Y. T. *The existence of Kähler-Einstein metrics on manifolds with positive anticanonical line bundle and a suitable finite symmetry group.* Ann. of Math. (2) 127 (1988), no. 3, 585–627.
- [27] Tian, G. On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$, Invent. Math. 89 (1987), no. 2, 225–246.
- [28] Tian, G. and Yau, S.-T. *Kähler-Einstein metrics on complex surfaces with* $C_1 > 0$. Comm. Math. Phys. 112 (1987), no. 1, 175–203.
- [29] Zeng, Y. Deformations from a given Kähler metric to a twisted cscK metric, Asian J. Math. 23 (2019), no. 6, 985-1000.

- [30] Zhang, K., *Continuity of delta invariants and twisted Kähler-Einstein metrics*, Advances in Mathematics, 388 (2021), 107888.
- [31] Zheng, K. Existence of constant scalar curvature Kaehler cone metrics, properness and geodesic stability, arXiv:1803.09506.

ICTP and Università di Parma Email address: arezzo@ictp.it

UNIVERSITÀ DI MILANO - BICOCCA Email address: alberto.dellavedova@unimib.it

NJU

Email address: shiyl@nju.edu.cn