# Navier-Stokes equations, the algebraic aspect

Zharinov V.V. \*

#### Abstract

Analysis of the Navier-Stokes equations in the frames of the algebraic approach to systems of partial differential equations (formal theory of differential equations) is presented.

**Keywords:** Navier–Stokes equations, integrability conditions, evolution, constraints, differential algebra, symmetries, cohomologies.

# 1 Preliminaries.

# 1.1 Navier-Stokes equations

The Navier-Stokes equations in physical notation are (see [1] - [6], for example)

$$\mathbf{u}_t + (\mathbf{u} \cdot \nabla)\mathbf{u} = \nu \Delta \mathbf{u} - \nabla p, \tag{1}$$

$$\nabla \cdot \mathbf{u} = 0, \tag{2}$$

where  $\mathbf{u}=(u^1,u^2,u^3)$  is the velocity field, t is the time variable, the dot "·" denotes the scalar product,  $\nabla=(\nabla_1,\nabla_2,\nabla_3)$  is the gradient with respect to the spacial variables  $\mathbf{x}=(x^1,x^2,x^3)$ , the parameter  $\nu>0$  is the viscosity of the flow (do not confuse it with the index  $\nu$ ),  $\Delta=\nabla_1^2+\nabla_2^2+\nabla_3^2$  is the Laplacian, p is the pressure.

Here we study the Navier-Stokes equations from the algebra-geometrical point of view. The system (1)-(2) is not formally integrable, as we need, to

<sup>\*</sup>Steklov Mathematical Institute, e-mail: zharinov@mi-ras.ru

get the equivalent formally integrable system we add (see [7], for example) the trivial differential prolongations

$$\nabla \cdot \mathbf{u}_t = 0, \quad \nabla(\nabla \cdot \mathbf{u}) = 0, \tag{3}$$

and the non-trivial differential prolongation (hidden integrability condition)

$$\Delta p + \nabla ((\mathbf{u} \cdot \nabla)\mathbf{u}) = \Delta p + \nabla \mathbf{u} \cdot \nabla \mathbf{u} = 0 \tag{4}$$

(we have used the equation (2)).

**Remark 1.** The equation (4) is a Poisson equation for the pressure p with the density  $\rho = -\nabla \mathbf{u} \cdot \nabla \mathbf{u}$ , it can be considered as the *inner constraint* for the Navier-Stokes equations.

#### 1.2 Notation

We use rather sophisticated notation:

- $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\} \supset \mathbb{Z}_+ = \{0, 1, 2, \dots\} \supset \mathbb{N} = \{1, 2, 3, \dots\};$
- $M = \overline{1, m} = \{1, 2, \dots, m\}, \quad N = \overline{2, m} = \{2, \dots, m\}, \quad M = \{1\} \cup N;$
- $\mathbb{I} = \mathbb{Z}_{+}^{\mathcal{M}} = \{ \mathbf{i} = (i^{1}, \dots, i^{m}) \mid i^{\mu} \in \mathbb{Z}_{+}, \ \mu \in \mathcal{M} \}, \quad |\mathbf{i}| = i^{1} + \dots + i^{m};$
- $\mathbf{i} + \mathbf{j} = (i^1 + j^1, \dots, i^m + j^m), \quad \mathbf{i} + (\mu) = (i^1, \dots, i^{\mu} + 1, \dots, i^m),$  $(\mu) = 0 + (\mu), \text{ for all } \mathbf{i}, \mathbf{j} \in \mathbb{I}, \ \mu \in \mathcal{M};$
- $\mathbb{J} = \mathbb{Z}_+^{\mathbb{N}} = \{ \mathbf{j} = (j^2, \dots, j^m) \mid j^\alpha \in \mathbb{Z}_+, \ \alpha \in \mathbb{N} \};$
- $\mathbb{I} = \mathbb{Z}_+ \times \mathbb{J} = \{ \mathbf{i} = (i^1, \mathbf{j}) \mid i^1 \in \mathbb{Z}_+, \ \mathbf{j} \in \mathbb{J} \};$
- $\mathbb{I}_0 = \{ i \in \mathbb{I} \mid i^1 = 0 \} = \{ 0 \} \times \mathbb{J}, \quad \mathbb{I}_1 = \{ i \in \mathbb{I} \mid i^1 = 0, 1 \} = \{ 0, 1 \} \times \mathbb{J};$
- $\mathbb{I}'_0 = \mathbb{I} \setminus \mathbb{I}_0 = \{ i \in \mathbb{I} \mid i^1 > 0 \}, \ \mathbb{I}'_1 = \mathbb{I} \setminus \mathbb{I}_1 = \{ i \in \mathbb{I} \mid i^1 > 1 \}, \ \mathbb{I} = \mathbb{I}_0 \cup \mathbb{I}'_0 = \mathbb{I}_1 \cup \mathbb{I}'_1.$

We shall also use the notation:

- $X = \mathbb{R}^{M} = \{x = (x^{1}, \dots, x^{m}) \mid x^{\mu} \in \mathbb{R}, \ \mu \in M\};$
- $$\begin{split} \bullet \ \ \mathbb{R}_{\mathbb{I}}^{\mathcal{M}} &= \{\mathbf{u} = (u_{\mathbf{i}}^{\mu}) \mid u_{\mathbf{i}}^{\mu} \in \mathbb{R}, \ \mu \in \mathcal{M}, \ \mathbf{i} \in \mathbb{I} \}, \\ \mathbb{R}_{\mathbb{I}}^{\mathcal{N}} &= \{\mathbf{u} = (u_{\mathbf{i}}^{\alpha}) \mid u_{\mathbf{i}}^{\alpha} \in \mathbb{R}, \ \alpha \in \mathcal{N}, \ \mathbf{i} \in \mathbb{I} \}, \\ \mathbb{R}_{\mathbb{I}}^{\mathcal{M}} &= \mathbb{R}_{\mathbb{I}}^{1} \times \mathbb{R}_{\mathbb{I}}^{\mathcal{N}} = \mathbb{R}_{\mathbb{I}_{0}}^{1} \times \mathbb{R}_{\mathbb{I}_{0}}^{1} \times \mathbb{R}_{\mathbb{I}}^{\mathcal{N}}; \end{split}$$

- $\mathbb{R}_{\mathbb{I}} = \{ \mathbf{p} = (p_i) \mid p_i \in \mathbb{R}, i \in \mathbb{I} \} = \mathbb{R}_{\mathbb{I}_1} \times \mathbb{R}_{\mathbb{I}'_1};$
- $$\begin{split} \bullet & \ \mathbf{B} = X \times \mathbb{R}_{\mathbb{I}}^{M} \times \mathbb{R}_{\mathbb{I}}, \\ & \mathbf{CE} = X \times \mathbb{R}_{\mathbb{I}_{0}}^{1} \times \mathbb{R}_{\mathbb{I}}^{N} \times \mathbb{R}_{\mathbb{I}}, \\ & \mathbf{CPE} = X \times \mathbb{R}_{\mathbb{I}_{0}}^{1} \times \mathbb{R}_{\mathbb{I}}^{N} \times \mathbb{R}_{\mathbb{I}_{1}}. \end{split}$$

We assume the summation over repeated upper and lower indices in the prescribed limits.

The paper is rather calculative, the calculations are straightforward but tiresome, and we omit them as a rule.

#### 1.3 The base components

The base space for the Navier–Stokes equations in the algebraic approach is the infinite dimensional space  $\mathbf{B} = X \times \mathbb{R}^M_{\mathbb{I}} \times \mathbb{R}_{\mathbb{I}}$ , where X is the set of independent variables (space coordinates),  $\mathbb{R}^M_{\mathbb{I}}$  is the set of differential variables (velocity coordinates and their partial derivatives),  $\mathbb{R}_{\mathbb{I}}$  is the set of differential variables (pressure and its partial derivatives).

The base algebra in the algebraic approach is the unital commutative associative algebra  $\mathcal{A}(\mathbf{B}) = \mathcal{C}^{\infty}_{\mathrm{fin}}(\mathbf{B})$  of all smooth real functions on the base space  $\mathbf{B}$  of a finite order, i.e., depending on a finite number of the variables  $x^{\mu}, u^{\mu}_{\mathbf{i}}, p_{\mathbf{i}}$ , where  $\mu \in \mathcal{M}$ ,  $\mathbf{i} \in \mathbb{I}$ . In more detail, the integer  $r \in \mathbb{Z}_{+}$  is called the  $\mathbf{u}$ -order of a function  $f(x, \mathbf{u}, \mathbf{p}) \in \mathcal{A}(\mathbf{B})$ , we write  $\mathrm{ord}_{\mathbf{u}} f = r$ , if the partial derivative  $\partial_{u^{\mu}_{\mathbf{i}}} f \neq 0$  for some variable  $u^{\mu}_{\mathbf{i}}, |\mathbf{i}| = r$ , while partial derivatives  $\partial_{u^{\mu}_{\mathbf{i}}} f = 0$  for all  $|\mathbf{i}| > r$ . In the same way, the  $\mathbf{p}$ -order is defined.

The base algebra of derivations of the algebra  $\mathcal{A}(\mathbf{B})$  is the Lie algebra

$$\mathfrak{D}(\mathbf{B}) = \mathfrak{D}(\mathcal{A}(\mathbf{B})) = \left\{ \zeta = \zeta^{\mu} \partial_{x^{\mu}} + \zeta_{i}^{\mu} \partial_{u_{i}^{\mu}} + \zeta_{i} \partial_{p_{i}} \mid \zeta^{\mu}, \zeta_{i}^{\mu}, \zeta_{i} \in \mathcal{A}(\mathbf{B}) \right\}.$$

The Lie algebra  $\mathfrak{D}(\mathbf{B})$  splits into the *vertical* and *horizontal* subalgebras,

$$\mathfrak{D}(\mathbf{B}) = \mathfrak{D}_{\mathrm{V}}(\mathbf{B}) \oplus_{\mathcal{A}(\mathbf{B})} \mathfrak{D}_{\mathrm{H}}(\mathbf{B}),$$

where

• 
$$\mathfrak{D}_{V}(\mathbf{B}) = \{ \zeta \in \mathfrak{D}(\mathbf{B}) \mid \zeta |_{\mathcal{C}^{\infty}(X)} = 0 \}$$
 (note,  $\mathcal{C}^{\infty}(X) \subset \mathcal{A}(\mathbf{B})$ )  
=  $\{ \zeta = \zeta_{i}^{\mu} \partial_{u_{i}^{\mu}} + \zeta_{i} \partial_{p_{i}} \mid \zeta_{i}^{\mu}, \zeta_{i} \in \mathcal{A}(\mathbf{B}) \};$ 

• 
$$\mathfrak{D}_{\mathrm{H}}(\mathbf{B}) = \{ \zeta = \zeta^{\mu} D_{\mu} \mid \zeta^{\mu} \in \mathcal{A}(\mathbf{B}) \}, \qquad (D_{\mu}|_{\mathcal{C}^{\infty}(\mathrm{X})} = \partial_{x^{\mu}})$$
  

$$D_{\mu} = \partial_{x^{\mu}} + u^{\lambda}_{\mathrm{i}+(\mu)} \partial_{u^{\lambda}_{\mathrm{i}}} + p_{\mathrm{i}+(\mu)} \partial_{p_{\mathrm{i}}}, \qquad \lambda, \mu \in \mathrm{M};$$

•  $[D_{\mu}, D_{\nu}] = 0$ ,  $[D_{\mu}, \zeta] = (D_{\mu}\zeta_{i}^{\lambda} - \zeta_{i+(\mu)}^{\lambda})\partial_{u_{i}^{\lambda}} + (D_{\mu}\zeta_{i} - \zeta_{i+(\mu)})\partial_{p_{i}}$ , for all  $\lambda, \mu, \nu \in M$ ,  $\zeta = \zeta_{i}^{\lambda}\partial_{u_{i}^{\lambda}} + \zeta_{i}\partial_{p_{i}} \in \mathfrak{D}_{V}(\mathbf{B})$ .

We denote

• 
$$D_{\mathbf{i}} = (D_1)^{i^1} \circ \ldots \circ (D_m)^{i^m}$$
 for all  $\mathbf{i} = (i^1, \ldots, i^m) \in \mathbb{I}$ .

The pair  $(\mathcal{A}(\mathbf{B}), \mathfrak{D}_{\mathbf{H}}(\mathbf{B}))$  is called the *differential algebra* associated with the base space  $\mathbf{B}$ .

The Lie algebra

$$Sym(\mathcal{A}(\mathbf{B}), \mathfrak{D}_{H}(\mathbf{B})) = \{ \zeta = ev_f \in \mathfrak{D}_{V}(\mathbf{B}) \mid [D_{\mu}, ev_f] = 0, \ \mu \in M \}$$

is the Lie algebra of symmetries of the differential algebra  $(\mathcal{A}(\mathbf{B}), \mathfrak{D}_{\mathbf{H}}(\mathbf{B}))$ , where

• 
$$f = (f^{\mu}, f) \in \mathcal{A}^{M}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}), \quad f^{\mu} = \zeta_{0}^{\mu}, f = \zeta_{0}$$

• 
$$\operatorname{ev}_{\mathbf{f}} = D_{\mathbf{i}} f^{\mu} \cdot \partial_{u_{\mathbf{i}}} + D_{\mathbf{i}} f \cdot \partial_{p_{\mathbf{i}}}, \quad D_{\mathbf{i}} f^{\mu} = \zeta_{\mathbf{i}}^{\mu}, \ D_{\mathbf{i}} f = \zeta_{\mathbf{i}}.$$

The  $\mathbb{Z}$ -graded  $\mathcal{A}(\mathbf{B})$ -module  $\Omega_{\mathrm{H}}(\mathbf{B}) = \bigoplus_{q \in \mathbb{Z}} \Omega_{\mathrm{H}}^{q}(\mathbf{B})$  of horizontal differential forms is defined as follows,

$$\Omega_{\mathrm{H}}^{q}(\mathbf{B}) = \begin{cases}
0, & q < 0, q > m; \\
\mathcal{A}(\mathbf{B}), & q = 0; \\
\operatorname{Hom}_{\mathcal{A}(\mathbf{B})}(\wedge^{q}\mathfrak{D}_{\mathrm{H}}(\mathbf{B}); \mathcal{A}(\mathbf{B})), & 1 \leq q \leq m.
\end{cases}$$

Here,

$$\operatorname{Hom}_{\mathcal{A}(\mathbf{B})}(\wedge^{q}\mathfrak{D}_{\mathbf{H}}(\mathbf{B}); \mathcal{A}(\mathbf{B})) = \{ \omega^{q} = \omega_{\mu_{1}...\mu_{q}} \cdot dx^{\mu_{1}} \wedge ... \wedge dx^{\mu_{q}} \mid \omega_{\mu_{1}...\mu_{q}} \in \mathcal{A}(\mathbf{B}), + \text{s-s} \},$$

where the abbreviation "+s-s" states that components  $\omega_{\mu_1...\mu_q}$  are skew-symmetric in indices  $\mu_1, \ldots, \mu_q \in M$ .

The horizontal differential  $d_{\rm H} \in \operatorname{End}_{\mathbb{R}}(\Omega_{\rm H}(\mathbf{B})), d_{\rm H} \circ d_{\rm H} = 0$ , is defined by the rule,

$$d_{\mathrm{H}}^{q} = d_{\mathrm{H}}|_{\Omega_{\mathrm{H}}^{q}(\mathbf{B})} : \Omega_{\mathrm{H}}^{q}(\mathbf{B}) \to \Omega_{\mathrm{H}}^{q+1}(\mathbf{B}),$$
  
$$\omega_{\mu_{1}...\mu_{q}} \cdot dx^{\mu_{1}} \wedge ... \wedge dx^{\mu_{q}} \mapsto D_{[\mu_{0}}\omega_{\mu_{1}...\mu_{q}]} \cdot dx^{\mu_{0}} \wedge ... \wedge dx^{\mu_{q}},$$

the brackets [...] denote the skew-symmetrization in indices  $\mu_0, \ldots, \mu_q \in M$ , so  $d_H^{q+1} \circ d_H^q = 0, q \in \mathbb{Z}$ .

The  $\mathbb{Z}$ -graded linear space  $H(\Omega_{\mathrm{H}}(\mathbf{B}); d_{\mathrm{H}}) = \bigoplus_{q \in \mathbb{Z}} H_{\mathrm{H}}^{q}(\mathbf{B})$  of cohomologies of the differential algebra  $(\mathcal{A}(\mathbf{B}); \mathfrak{D}_{\mathrm{H}}(\mathbf{B}))$  is defined in the usual way,  $H(\Omega_{\mathrm{H}}(\mathbf{B}); d_{\mathrm{H}}) = \operatorname{Ker} d_{\mathrm{H}} / \operatorname{Im} d_{\mathrm{H}}, H_{\mathrm{H}}^{q}(\mathbf{B}) = \operatorname{Ker} d_{\mathrm{H}} / \operatorname{Im} d_{\mathrm{H}}^{q-1}, q \in \mathbb{Z}.$ 

Theorem 1 (The main theorem of the formal calculus of variations, see, for example, [8], and the references therein). The linear spaces

$$H_{H}^{q}(\mathbf{B}) = \begin{cases} 0, & q < 0, q > m; \\ \mathbb{R}, & q = 0; \\ 0, & 1 \le q \le m - 1; \\ \mathcal{H}(\mathbf{B}), & q = m; \end{cases}$$

where (in our setting) the Helmholtz linear space

$$\mathcal{H}(\mathbf{B}) = \{ \chi = (\chi_{\mu}, \chi) \in \mathcal{A}_{M}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}) \mid \chi_{*} = \chi^{*} \},$$

the linear mappings  $\chi_*, \chi^* : \mathcal{A}^M(\mathbf{B}) \times \mathcal{A}(\mathbf{B}) \to \mathcal{A}_M(\mathbf{B}) \times \mathcal{A}(\mathbf{B})$  act by the rules:

$$\mathcal{A}^{\mathrm{M}}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}) \ni \mathbf{f} = (f^{\mu}, f) \mapsto \chi_{*}\mathbf{f} = \mathbf{g} = (g_{\mu}, g) \in \mathcal{A}_{\mathrm{M}}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}),$$

$$g_{\mu} = \partial_{u_{i}^{\nu}}\chi_{\mu} \cdot D_{i}f^{\nu} + \partial_{p_{i}}\chi_{\mu} \cdot D_{i}f, \quad g = \partial_{u_{i}^{\nu}}\chi \cdot D_{i}f^{\nu} + \partial_{p_{i}}\chi \cdot D_{i}f;$$

$$\mathcal{A}^{\mathrm{M}}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}) \ni \mathbf{f} = (f^{\mu}, f) \mapsto \chi^{*}\mathbf{f} = \mathbf{g} = (g_{\mu}, g) \in \mathcal{A}_{\mathrm{M}}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}),$$

$$g_{\mu} = (-D)_{i}(f^{\nu} \cdot \partial_{u_{i}^{\mu}}\chi_{\nu} + f \cdot \partial_{u_{i}^{\mu}}\chi), \quad g = (-D)_{i}(f^{\nu} \cdot \partial_{p_{i}}\chi_{\nu} + f \cdot \partial_{p_{i}}\chi).$$

The isomorphism  $\delta = (\delta_{u^{\mu}}, \delta_p) : H_{H}^{m}(\mathbf{B}) \simeq \mathcal{H}(\mathbf{B})$  of linear spaces is defined as the variational derivatives:

$$\Omega_{\mathrm{H}}^{m}(\mathbf{B}) \ni \omega = L \cdot d^{m}x \mapsto \chi = \delta L = (\delta_{u^{\mu}}L, \delta_{p}L),$$
  
$$\delta_{u^{\mu}}L = (-D)_{i}\partial_{u_{i}^{\mu}}L, \quad \delta_{p}L = (-D)_{i}\partial_{p_{i}}L.$$

Remark 2. Note, cohomologies are defined up to isomorphisms.

# 2 Constraints

# 2.1 The divergence-free space

The continuity equation  $CE = \{\partial_{x^{\mu}} u^{\mu} = 0\}$  (see (2)) has the algebraic counterpart

$$\mathbf{CE} = \big\{ (x, \mathbf{u}, \mathbf{p}) \in \mathbf{B} \ \big| \ \mathrm{CE}_i = u^{\mu}_{i+(\mu)} = 0, \ i \in \mathbb{I} \big\}.$$

The subset  $\mathbf{CE} \subset \mathbf{B}$  is called the *solution manifold* of the equation CE, because a smooth function  $\phi(\mathbf{x}) = (\phi^{\mu}(\mathbf{x})) \in \mathcal{C}^{\infty}(\mathbf{X}; \mathbb{R}^{\mathbf{M}})$  is a solution of the equation CE iff the graph

graph 
$$\mathbf{j}\phi(\mathbf{x}) = \{(\mathbf{x}, \mathbf{u}, \mathbf{p}) \mid u_i^{\mu} = \partial_{x^i}\phi^{\mu}(\mathbf{x}), \ \mu \in \mathcal{M}, \ i \in \mathbb{I}\} \subset \mathbf{CE},$$

where  $\mathbf{j}\phi(\mathbf{x}) = \{\partial_{x^i}\phi^{\mu}(\mathbf{x}) \mid i \in \mathbb{I}, \ \mu \in \mathbf{M}\}\$ is the jet of the function  $\phi(\mathbf{x})$ , the variables  $\mathbf{p} = (p_i) \in \mathbb{R}_{\mathbb{I}}$  are parameters,  $\partial_{x^i} = (\partial_{x^1})^{i^1} \circ \ldots \circ (\partial_{x^m})^{i^m}$ ,  $i \in \mathbb{I}$ .

With the solution manifold CE there is associated the ideal

$$\mathcal{I} = \left\{ f \in \mathcal{A}(\mathbf{B}) \mid f \big|_{\mathbf{CE}} = 0 \right\} = \left\{ f = P(D) u^{\mu}_{(\mu)} \mid P(D) \in \mathcal{A}(\mathbf{B})[D] \right\}$$
$$= \left\{ f = \phi^{\mathbf{i}} \cdot u^{\mu}_{\mathbf{i} + (\mu)} \mid \phi^{\mathbf{i}} \in \mathcal{A}(\mathbf{B}), \ \mathbf{i} \in \mathbb{I}, + \text{fin} \right\}$$

of the algebra  $\mathcal{A}(\mathbf{B})$ , where  $\mathcal{A}(\mathbf{B})[D]$  is the unital associative algebra of all polynomials in the indeterminates  $D_1, \ldots, D_m$  with coefficients in  $\mathcal{A}(\mathbf{B})$ , i.e.,

$$\mathcal{A}(\mathbf{B})[D] = \{ P(D) = \phi^{i} \cdot D_{i} \mid \phi^{i} \in \mathcal{A}(\mathbf{B}), i \in \mathbb{I}, +fin \},$$

where the abbreviation "+fin" states that only a finite number of the coefficients  $\phi^{i} \neq 0$ . The ideal  $\mathcal{I}$  is differential, i.e.,  $D_{\mu}|_{\mathcal{I}} : \mathcal{I} \to \mathcal{I}$ ,  $\mu \in \mathcal{M}$ .

By definition, we assume  $\mathcal{A}(\mathbf{CE}) = \mathcal{A}(\mathbf{B})/\mathcal{I}$ , i.e., we define the algebra  $\mathcal{A}(\mathbf{CE})$  of smooth functions on the space  $\mathbf{CE}$  as the factor-algebra. We denote  $\bar{f} = f|_{\mathbf{CE}} = f + \mathcal{I}$  the restriction of the function  $f \in \mathcal{A}(\mathbf{B})$  to the solution manifold  $\mathbf{CE}$  and its equivalence class in  $\mathcal{A}(\mathbf{CE})$ .

We shall need:

• the subalgebra of the Lie algebra  $\mathfrak{D}(\mathbf{B})$ 

$$\mathfrak{D}_{\mathcal{I}}(\mathbf{B}) = \left\{ \zeta \in \mathfrak{D}(\mathbf{B}) \mid \zeta|_{\mathcal{I}} : \mathcal{I} \to \mathcal{I} \right\} = \left\{ \zeta \in \mathfrak{D}(\mathbf{B}) \mid \zeta|_{\mathcal{I}} \in \mathfrak{D}(\mathcal{I}) \right\}$$
$$= \left\{ \zeta = \zeta^{\mu} \partial_{x^{\mu}} + \zeta_{i}^{\mu} \partial_{u_{i}^{\mu}} + \zeta_{i} \partial_{p_{i}} \mid \zeta_{i+(\mu)}^{\mu} \in \mathcal{I}, \ i \in \mathbb{I} \right\}$$

of all derivations of the algebra  $\mathcal{A}(\mathbf{B})$  leaving the ideal  $\mathcal{I}$  invariant;

• the subalgebra of the Lie algebra  $\mathfrak{D}(\mathbf{B})$  and the ideal of the Lie algebra  $\mathfrak{D}_{\mathcal{I}}(\mathbf{B}) \subset \mathfrak{D}(\mathbf{B})$ 

$$\mathfrak{D}(\mathbf{B}; \mathcal{I}) = \left\{ \zeta \in \mathfrak{D}(\mathbf{B}) \mid \zeta : \mathcal{A}(\mathbf{B}) \to \mathcal{I} \right\}$$
$$= \left\{ \zeta = \zeta^{\mu} \partial_{x^{\mu}} + \zeta_{\mathbf{i}}^{\mu} \partial_{u_{\mathbf{i}}^{\mu}} + \zeta_{\mathbf{i}} \partial_{p_{\mathbf{i}}} \mid \zeta^{\mu}, \zeta_{\mathbf{i}}^{\mu}, \zeta_{\mathbf{i}} \in \mathcal{I} \right\}$$

of all derivations with the image in the ideal  $\mathcal{I}$ ,

• the Lie factor-algebra  $\mathfrak{D}(\mathbf{CE}) = \mathfrak{D}_{\mathcal{I}}(\mathbf{B})/\mathfrak{D}(\mathbf{B};\mathcal{I})$  of derivations of the factor-algebra  $\mathcal{A}(\mathbf{CE})$ .

Here, the mapping  $\mathfrak{D}_{\mathcal{I}}(\mathbf{B}) \to \mathfrak{D}(\mathbf{CE})$  is defined as follows:

$$\mathfrak{D}_{\mathcal{I}}(\mathbf{B}) \ni \zeta \mapsto \bar{\zeta} = \zeta + \mathfrak{D}(\mathbf{B}; \mathcal{I}) : \mathcal{A}(\mathbf{CE}) \to \mathcal{A}(\mathbf{CE}),$$
$$\bar{f} = f + \mathcal{I} \mapsto \bar{\zeta}\bar{f} = \overline{\zeta}f = \zeta f + \mathcal{I}$$

(we denote  $\bar{\zeta} = \zeta + \mathfrak{D}(\mathbf{B}; \mathcal{I})$  the equivalence class of a derivation  $\zeta \in \mathfrak{D}_{\mathcal{I}}(\mathbf{B})$ ). The Lie algebra  $\mathfrak{D}(\mathbf{CE})$  splits into the *vertical* and *horizontal* subalgebras:

- $\bullet \ \mathfrak{D}(\mathbf{CE}) = \mathfrak{D}_V(\mathbf{CE}) \oplus_{\mathcal{A}(\mathbf{CE})} \mathfrak{D}_H(\mathbf{CE}),$
- $\mathfrak{D}_{V}(CE) = \{\bar{\zeta} \in \mathfrak{D}(CE) \mid \zeta \in \mathfrak{D}_{V}(B) \cap \mathfrak{D}_{\mathcal{I}}(B)\},\$
- $\bullet \ \mathfrak{D}_H(\mathbf{CE}) = \{\bar{\zeta} \in \mathfrak{D}(\mathbf{CE}) \mid \zeta \in \mathfrak{D}_H(\mathbf{B})\} \qquad (\mathrm{note}, \, \mathfrak{D}_H(\mathbf{B}) \subset \mathfrak{D}_{\mathcal{I}}(\mathbf{B})).$

The Lie algebra

$$\operatorname{Sym}(\mathcal{A}(\mathbf{C}\mathbf{E}), \mathfrak{D}_{\mathbf{H}}(\mathbf{C}\mathbf{E})) = \left\{ \bar{\zeta} \in \mathfrak{D}_{\mathbf{V}}(\mathbf{C}\mathbf{E}) \mid [\bar{D}_{\mu}, \bar{\zeta}] = \bar{0} = \mathcal{I}, \ \mu \in \mathbf{M} \right\}$$

$$= \left\{ \bar{\zeta} \mid \zeta = D_{\mathbf{i}} f^{\mu} \cdot \partial_{u_{\mathbf{i}}^{\mu}} + D_{\mathbf{i}} f \cdot \partial_{p_{\mathbf{i}}}, \ D_{\mu} f^{\mu} \in \mathcal{I} \right\}$$

$$= \left\{ \bar{\zeta} \mid \zeta = \operatorname{ev}_{\mathbf{f}}, \ \mathbf{f} = (f^{\mu}, f) \in \mathcal{A}^{\mathbf{M}}(\mathbf{B}) \times \mathcal{A}(\mathbf{B}), \ (D_{\mu} f^{\mu}) \right|_{\mathbf{C}\mathbf{E}} = 0 \right\}$$

is the Lie algebra of symmetries of the differential algebra  $(\mathcal{A}(CE), \mathfrak{D}_H(CE))$ .

The  $\mathbb{Z}$ -graded  $\mathcal{A}(\mathbf{CE})$ -module  $\Omega_{\mathrm{H}}(\mathbf{CE}) = \bigoplus_{q \in \mathbb{Z}} \Omega_{\mathrm{H}}^q(\mathbf{CE})$  of horizontal differential forms is defined as follows,

$$\Omega_{\mathrm{H}}^{q}(\mathbf{CE}) = \begin{cases}
0, & q < 0, q > m \\
\mathcal{A}(\mathbf{CE}), & q = 0; \\
\operatorname{Hom}_{\mathcal{A}(\mathbf{CE})}(\wedge^{q}\mathfrak{D}_{\mathrm{H}}(\mathbf{CE}); \mathcal{A}(\mathbf{CE})), & 1 \leq q \leq m.
\end{cases}$$

Here,

$$\begin{aligned} \operatorname{Hom}_{\mathcal{A}(\mathbf{CE})}(\wedge^{q}\mathfrak{D}_{H}(\mathbf{CE}); \mathcal{A}(\mathbf{CE})) \\ &= \big\{ \bar{\omega}^{q} = \bar{\omega}_{\mu_{1}...\mu_{q}} \cdot d\bar{x}^{\mu_{1}} \wedge \ldots \wedge d\bar{x}^{\mu_{q}} \mid \bar{\omega}_{\mu_{1}...\mu_{q}} \in \mathcal{A}(\mathbf{CE}), + \text{ s-s} \big\}, \end{aligned}$$

where  $d\bar{x}^{\mu}(\bar{D}_{\nu}) = \bar{\delta}^{\mu}_{\nu} = \delta^{\mu}_{\nu} + \mathcal{I}, \, \mu, \nu \in \mathcal{M}.$ 

The horizontal differential  $d_{\rm H} \in \operatorname{End}_{\mathbb{R}}(\Omega_{\rm H}(\mathbf{CE})), d_{\rm H} \circ d_{\rm H} = 0$ , is defined by the rule,

$$d_{\mathrm{H}}^{q} = d_{\mathrm{H}}|_{\Omega_{\mathrm{H}}^{q}(\mathbf{C}\mathbf{E})} : \Omega_{\mathrm{H}}^{q}(\mathbf{C}\mathbf{E}) \to \Omega_{\mathrm{H}}^{q+1}(\mathbf{C}\mathbf{E}),$$
  
$$\bar{\omega}_{\mu_{1}...\mu_{q}} \cdot d\bar{x}^{\mu_{1}} \wedge \ldots \wedge d\bar{x}^{\mu_{q}} \mapsto \bar{D}_{[\mu_{0}}\bar{\omega}_{\mu_{1}...\mu_{q}]} \cdot d\bar{x}^{\mu_{0}} \wedge \ldots \wedge d\bar{x}^{\mu_{q}},$$

where  $d_{\mathbf{H}}^{q+1} \circ d_{\mathbf{H}}^{q} = 0, q \in \mathbb{Z}$ .

The  $\mathbb{Z}$ -graded linear space  $H(\Omega_{\mathrm{H}}(\mathbf{CE}); d_{\mathrm{H}}) = \bigoplus_{q \in \mathbb{Z}} H_{\mathrm{H}}^q(\mathbf{CE})$  of cohomologies of the differential algebra  $(\mathcal{A}(\mathbf{CE}); \mathfrak{D}_{\mathrm{H}}(\mathbf{CE}))$  is defined in the usual way,  $H(\Omega_{\mathrm{H}}(\mathbf{CE}); d_{\mathrm{H}}) = \operatorname{Ker} d_{\mathrm{H}}/\operatorname{Im} d_{\mathrm{H}}, \ H_{\mathrm{H}}^q(\mathbf{CE}) = \operatorname{Ker} d_{\mathrm{H}}/\operatorname{Im} d_{\mathrm{H}}^{q-1}, \ q \in \mathbb{Z}.$ 

To move further, in particular to calculate the cohomology spaces, we introduce global coordinates on the solution manifold **CE**. Namely, the linear space  $\mathbb{R}^M_{\mathbb{I}}$  we split as

$$\mathbb{R}^{\mathrm{M}}_{\mathbb{I}} = \mathbb{R}^{1}_{\mathbb{I}} \times \mathbb{R}^{\mathrm{N}}_{\mathbb{I}} = \mathbb{R}^{1}_{\mathbb{I}_{0}} \times \mathbb{R}^{1}_{\mathbb{I}'_{0}} \times \mathbb{R}^{\mathrm{N}}_{\mathbb{I}}$$

(see Subsection 1.2). Then, the global coordinates on **CE** are defined by the isomorphism of the linear spaces

$$\mathbf{CE} = \left\{ (x^{\mu}, u_{\mathbf{i}}^{\mu}, p_{\mathbf{i}}) \in \mathbf{B} \mid u_{\mathbf{i}+(\mu)}^{\mu} = 0 \right\} \to \mathbf{X} \times \mathbb{R}_{\mathbb{I}_{0}}^{1} \times \mathbb{R}_{\mathbb{I}}^{\mathbf{N}} \times \mathbb{R}_{\mathbb{I}},$$
$$(x^{\mu}, u_{\mathbf{i}}^{1}, u_{\mathbf{i}}^{\alpha}, p_{\mathbf{i}}) \mapsto (x^{\mu}, u_{\mathbf{i}_{0}}^{1}, u_{\mathbf{i}}^{\alpha}, p_{\mathbf{i}}),$$

with the inverse mapping

$$X \times \mathbb{R}^{1}_{\mathbb{I}_{0}} \times \mathbb{R}^{N}_{\mathbb{I}} \times \mathbb{R}_{\mathbb{I}} \to \left\{ (x^{\mu}, u_{i}^{\mu}, p_{i}) \in \mathbf{B} \mid u_{i+(\mu)}^{\mu} = 0 \right\} = \mathbf{CE},$$
$$(x^{\mu}, u_{i_{0}}^{1}, u_{i}^{\alpha}, p_{i}) \mapsto (x^{\mu}, u_{i}^{1}, u_{i}^{\alpha}, p_{i}),$$

where  $u^1_{\mathbf{i}'_0} = -u^{\beta}_{\mathbf{i}'_0 - (1) + (\beta)}$ ,  $\mu = (1, \alpha) \in \mathcal{M}$ ,  $\mathbf{i}_0 \in \mathbb{I}_0$ ,  $\alpha, \beta \in \mathcal{N}$ ,  $\mathbf{i}'_0 \in \mathbb{I}'_0$ . In these global coordinates we have:

- $\mathbf{CE} = X \times \mathbb{R}^1_{\mathbb{I}_0} \times \mathbb{R}^N_{\mathbb{I}} \times \mathbb{R}_{\mathbb{I}}, \qquad \mathcal{A}(\mathbf{CE}) = \mathcal{C}^{\infty}_{\mathrm{fin}}(\mathbf{CE});$
- $\mathfrak{D}_{V}(\mathbf{CE}) = \{ \zeta = \zeta_{i_0}^1 \partial_{u_{i_{\alpha}}^1} + \zeta_{i}^{\alpha} \partial_{u_{i}^{\alpha}} + \zeta_{i} \partial_{p_i} \mid \zeta_{i_0}^1, \zeta_{i}^{\alpha}, \zeta_{i} \in \mathcal{A}(\mathbf{CE}) \};$
- $D_{\mu} = \partial_{x^{\mu}} + u_{i_0+(\mu)}^1 \partial_{u_{i_0}^1} + u_{i_+(\mu)}^{\alpha} \partial_{u_{i}^{\alpha}} + p_{i_+(\mu)} \partial_{p_i}, \quad \mu \in \mathcal{M},$ where  $u_{i_0+(1)}^1 = -u_{i_0+(\alpha)}^{\alpha};$
- $\operatorname{Sym}(\mathcal{A}(\mathbf{CE}), \mathfrak{D}_{H}(\mathbf{CE})) = \{ \zeta = \operatorname{ev}_{f} \in \mathfrak{D}_{V}(\mathbf{CE}) \mid [D_{\mu}, \operatorname{ev}_{f}] = 0, \ \mu \in M \},$  where

$$- f = (f^{\mu}, f) \in \mathcal{A}^{M}(\mathbf{CE}) \times \mathcal{A}(\mathbf{CE}), \quad D_{\mu}f^{\mu} = 0,$$
  
$$- \operatorname{ev}_{f} = D_{i_{0}}f^{1} \cdot \partial_{u_{i_{0}}^{1}} + D_{i}f^{\alpha} \cdot \partial_{u_{i}^{\alpha}} + D_{i}f \cdot \partial_{p_{i}};$$

•  $\Omega_{\mathrm{H}}^{q}(\mathbf{CE}) = \{ \omega = \omega_{\mu_{1}...\mu_{q}} \cdot dx^{\mu_{1}} \wedge ... \wedge dx^{\mu_{q}} \mid \omega_{\mu_{1}...\mu_{q}} \in \mathcal{A}(\mathbf{CE}), + \text{s-s} \},$  $1 \leq q \leq m.$  In more detail, here the equation  $D_{\mu}f^{\mu}=0$  takes the form

$$D_{\mu}f^{\mu} = \partial_{x^{\mu}}f^{\mu} + u^{1}_{i_{0}+(\mu)}\partial_{u^{1}_{i_{0}}}f^{\mu} + u^{\alpha}_{i+(\mu)}\partial_{u^{\alpha}_{i}}f^{\mu} + p_{i+(\mu)}\partial_{p_{i}}f^{\mu} = 0.$$
 (5)

Let us suppose the function  $f = (f^{\mu})$  has the **p**-order  $r = \max_{\mu \in \mathcal{M}} \operatorname{ord}_{\mathbf{p}} f^{\mu}$ , then the term  $\partial_{p_{\kappa,i}} f^{\mu} \cdot p_{\kappa,i+(\mu)}$  will be of the **p**-order r+1, while all other terms of the equation (5) will have **p**-orders  $\leq r$ . This leads to the conclusion  $\partial_{p_{\kappa,i}} f^{\mu} = 0$ ,  $|\mathbf{i}| = r$ , hence the **p**-order of the function f is  $\leq r-1$ . By induction we get  $f^{\mu} = f^{\mu}(x, \mathbf{u})$ ,  $\mu \in \mathcal{M}$ , i.e., f does not depend on the variable  $\mathbf{p} \in \mathbb{R}_{\mathbb{I}}$ , and we get the following statement.

**Proposition 1.** In the above setting the equation (5) is reduced to


$$\partial_{x^{\mu}} f^{\mu} + u^{1}_{i_{0} + (\mu)} \partial_{u^{1}_{i_{0}}} f^{\mu} + u^{\alpha}_{i+(\mu)} \partial_{u^{\alpha}_{i}} f^{\mu} = 0, \quad f^{\mu}(x, \mathbf{u}) \in \mathcal{C}^{\infty}_{fin}(X \times \mathbb{R}^{1}_{\mathbb{I}_{0}} \times \mathbb{R}^{N}_{\mathbb{I}}).$$
 (6)

**Remark 3.** If we want to have symmetries that really depend on the variable  $\mathbf{p} \in \mathbb{R}_{\mathbb{I}}$ , we must narrow down to a subvariety of the space **CE**. Exactly such situation arises when studying the Navier-Stokes equations.

To calculate the cohomology spaces  $H_{\mathrm{H}}^q(\mathbf{CE})$  we introduce the auxiliary complex  $\{\Theta^q, d_{\Theta}^q \mid q \in \mathbb{Z}\}$ , where

- $\Theta^q = \{\theta^q = \theta_{\alpha_1...\alpha_q} dx^{\alpha_1} \wedge ... \wedge dx^{\alpha_q} \mid \alpha_1, ..., \alpha_q \in \mathbb{N}, \theta_{\alpha_1...\alpha_q} \in \mathcal{A}(\mathbf{CE})\};$
- $d_{\Theta}^q: \Theta^q \to \Theta^{q+1}, \ \theta_{\alpha_1...\alpha_q} dx^{\alpha_1} \wedge ... \wedge dx^{\alpha_q} \mapsto D_{[\alpha_0} \theta_{\alpha_1...\alpha_q]} \cdot dx^{\alpha_0} \wedge ... \wedge dx^{\alpha_q};$
- $H^q(\Theta) = \operatorname{Ker} d_{\Theta}^q / \operatorname{Im} d_{\Theta}^{q-1}$ .

There is defined the commutative diagram



where

• 
$$\Omega^q = \Omega^q_H(\mathbf{CE}), \quad \Omega^0 = \Theta^0 = \mathcal{A}(\mathbf{CE});$$

• 
$$\Omega^q = dx^1 \wedge \Theta^{q-1} \oplus_{\mathcal{A}(\mathbf{CE})} \Theta^q;$$

• 
$$d_{\mathrm{H}}(dx^1 \wedge \theta^{q-1} + \theta^q) = dx^1 \wedge (D_1\theta^q - d_{\Theta}\theta^{q-1}) + d_{\Theta}\theta^q$$
;

• 
$$D_1\theta^q = D_1(\theta_{\alpha_1...\alpha_q} \cdot dx^{\alpha_1} \wedge ... \wedge dx^{\alpha_q}) = (D_1\theta_{\alpha_1...\alpha_q}) \cdot dx^{\alpha_1} \wedge ... \wedge dx^{\alpha_q};$$

• 
$$\iota^q: \Theta^q \to \Omega^{q+1}, \quad \theta^q \mapsto \omega^{q+1} = (-1)^q dx^1 \wedge \theta^q + 0;$$

• 
$$\pi^q : \Omega^q \to \Theta^q$$
,  $\omega^q = dx^1 \wedge \theta^{q-1} + \theta^q \mapsto \theta^q$ ;

According to the general results of homology theory (see, for example, [9]) this diagram defines the long exact sequence of the cohomology spaces

$$0 \to H^{0}_{\mathrm{H}}(\mathbf{CE}) \to H^{0}(\Theta) \xrightarrow{D^{0}_{1}} H^{0}(\Theta) \to H^{1}_{\mathrm{H}}(\mathbf{CE}) \to H^{1}(\Theta) \xrightarrow{D^{1}_{1}} \dots$$

$$\dots \xrightarrow{D^{m-2}_{1}} H^{m-2}(\Theta) \to H^{m-1}_{\mathrm{H}}(\mathbf{CE}) \to H^{m-1}(\Theta) \xrightarrow{D^{m-1}_{1}} H^{m-1}(\Theta) \to$$

$$\to H^{m}_{\mathrm{H}}(\mathbf{CE}) \to 0,$$

where  $D_1^q: H^q(\Theta) \to H^q(\Theta)$ ,

$$[\theta^q] = \theta^q + \operatorname{Im} d_{\Theta}^{q-1} \mapsto D_1^q[\theta^q] = [D_1 \theta^q] = D_1 \theta^q + \operatorname{Im} d_{\Theta}^{q-1}.$$

By Theorem 1, we have  $H^q(\Theta) = 0$  for  $q \neq 0$  and  $q \neq m-1$ , while

• 
$$H^0(\Theta) = \{\phi(x^1) \in \mathcal{C}^{\infty}(\mathbb{R})\};$$

• 
$$H^{m-1}(\Theta) = \mathcal{H}(\Theta) = \{ \chi \in \mathcal{F}(\Theta) \mid \chi_* = \chi^* \},$$
  
 $\mathcal{F}(\Theta) = \mathcal{A}_1(\mathbf{CE}) \times \mathcal{A}_N^{\mathbb{Z}_+^1}(\mathbf{CE}) \times \mathcal{A}^{\mathbb{Z}_+^1}(\mathbf{CE}),$ 

where the isomorphism  $\delta: H^{m-1}(\Theta) \simeq \mathcal{H}(\Theta)$  is defined as follows:

$$\begin{split} \Theta^{m-1} \ni \theta &= L \cdot dx^2 \wedge \ldots \wedge dx^m \mapsto \delta L = (\delta_{u_0^1} L, \delta_{u_{i_1}^{\alpha}} L, \delta_{p_{i_1}} L), \\ \delta_{u_0^1} L &= (-D)_{\mathbf{j}} \partial_{u_{0,\mathbf{j}}^1} L, \quad \delta_{u_{i_1}^{\alpha}} L = (-D)_{\mathbf{j}} \partial_{u_{i_1,\mathbf{j}}^{\alpha}} L, \quad \delta_{p_{i_1}} L = (-D)_{\mathbf{j}} \partial_{p_{i_1,\mathbf{j}}} L, \end{split}$$

here  $j \in \mathbb{J}$ ,  $\alpha \in \mathbb{N}$ ,  $i^1 \in \mathbb{Z}_+$ , remind,  $i_0 = (0, j)$ ,  $i = (i^1, j)$  (see Subsection 1.2), so  $u_{i_0}^1 = u_{0,j}^1$ ,  $u_i^{\alpha} = u_{i_1,j}^{\alpha}$ ,  $p_i = p_{i_1,j}$ .

The above long exact sequence defines the short exact sequence

$$0 \to H^0_{\mathrm{H}}(\mathbf{CE}) \to \mathrm{Ker}\, D^0_1 \to 0$$
, hence,  $H^0_{\mathrm{H}}(\mathbf{CE}) = \mathbb{R}$ .

In the same way, we have the exact sequence

$$0 \to \operatorname{Im} D_1^0 \to H^0(\Theta) \to H^1_{\mathrm{H}}(\mathbf{CE}) \to 0 \quad (H^1(\Theta) = 0),$$

hence,  $H^1_{\mathrm{H}}(\mathbf{C}\mathbf{E}) = H^0(\Theta)/\operatorname{Im} D^0_1 = 0$ . Further, for  $1 \leq q \leq m-2$  we have  $H^q(\Theta) = 0$  and  $H^{q-1}(\Theta) \to H^q_{\mathrm{H}}(\mathbf{C}\mathbf{E}) \to H^q(\Theta)$ , so  $H^q_{\mathrm{H}}(\mathbf{C}\mathbf{E}) = 0$  for  $2 \leq q \leq m-2$ . Now, for q=m-1 we have

$$\dots H^{m-2}(\Theta) \to H^{m-1}_{\mathrm{H}}(\mathbf{CE}) \to H^{m-1}(\Theta) \xrightarrow{D_1^{m-1}} \dots$$

Here,  $H^{m-2}(\Theta) = 0$ , hence,  $0 \to H_{\rm H}^{m-1}(\mathbf{CE}) \to \operatorname{Ker} D_1^{m-1} \to 0$  is the exact sequence, i.e.,  $H_{\rm H}^{m-1}(\mathbf{CE}) = \operatorname{Ker} D_1^{m-1}$  (note, cohomologies are defined up to isomorphisms).

**Lemma 1.** (See, for example, [10], [11]) There is defined the commutative diagram

$$0 \longrightarrow H^{m-1}(\Theta) \xrightarrow{\delta} \mathcal{H}(\Theta) \longrightarrow 0$$

$$D_1^{m-1} \downarrow \qquad \tilde{D_1} \downarrow$$

$$0 \longrightarrow H^{m-1}(\Theta) \xrightarrow{\delta} \mathcal{H}(\Theta) \longrightarrow 0,$$

in particular,  $\delta \circ D_1^{m-1} = \tilde{D}_1 \circ \delta$ , where  $D_1 = \partial_{x^1} + \operatorname{ev}_f'$ ,  $\tilde{D}_1 = D_1 + f^*$ ,

$$ev'_{f} = D_{j} f_{0}^{1} \cdot \partial_{u_{0,j}^{1}} + D_{j} f_{i}^{\alpha} \cdot \partial_{u_{i_{1,j}}^{\alpha}} + D_{j} f_{i_{1}} \cdot \partial_{p_{i_{1,j}}},$$

 $j \in \mathbb{J}, \ \alpha \in \mathbb{N}, \ i^1 \in \mathbb{Z}_+.$ 

In our setting,  $f = (f_0^1 = -u_{(\alpha)}^{\alpha}, \ f_{i^1}^{\alpha} = u_{i^1+1,0}^{\alpha}, \ f_{i^1} = p_{i^1+1,0})$ . Hence, here  $\tilde{D}_1, f^* \in \operatorname{End}_{\mathbb{R}}(\mathcal{A}_1(\mathbf{CE}) \times \mathcal{A}_N^{\mathbb{Z}_+^1}(\mathbf{CE}) \times \mathcal{A}^{\mathbb{Z}_+^1}(\mathbf{CE}))$ ,

$$\chi = (\chi^0_1, \chi^{i^1}_\alpha, \chi^{i^1}) \mapsto f^*\chi = ((f^*\chi)^0_1, (f^*\chi)^{i^1}_\alpha, (f^*\chi)^{i^1},$$

where

$$(f^*\chi)_1^0 = 0$$
,  $(f^*\chi)_{\alpha}^{i^1} = \delta_0^{i^1} D_{\alpha} \chi_1^0 + \chi_{\alpha}^{i^1-1}$ ,  $(f^*\chi)^{i^1} = \chi^{i^1-1}$ .

The system  $(D_1 + f^*)\chi = 0$  of the equations for defining an unknown function  $\chi \in \mathcal{A}_1(\mathbf{CE}) \times \mathcal{A}_N^{\mathbb{Z}^1_+}(\mathbf{CE}) \times \mathcal{A}^{\mathbb{Z}^1_+}(\mathbf{CE})$  with a finite number of the components  $\chi_1^0, \chi_{\alpha}^{i^1}, \chi^{i^1} \neq 0$ , reduces to

$$D_1 \chi_1^0 = 0$$
,  $D_1 \chi_{\alpha}^{i^1} + \delta_0^{i^1} D_{\alpha} \chi_1^0 + \chi_{\alpha}^{i^1 - 1} = 0$ ,  $D_1 \chi^{i^1} + \chi^{i^1 - 1} = 0$ ,

and has the general solution:  $\chi=(\chi_1^0,0,0), \ \chi_1^0\in\mathbb{R}$ . Now, we may take  $\delta u^1=(1,0,0)$ , hence the linear space  $\operatorname{Ker} D_1^{m-1}$  is one-dimensional with the basis  $[u^1d_1x]$  (remind,  $d_\mu x=(-1)^{\mu-1}dx^1\wedge\ldots dx^\mu\ldots\wedge dx^m$ , the term  $dx^\mu$  is omitted, so  $dx^\nu\wedge d_\mu x=\delta_\mu^\nu\cdot d^m x,\ \mu,\nu\in M$ ), i.e.,  $\operatorname{Ker} D_1^{m-1}=\mathbb{R}\cdot [u^1d_1x]$  (cf., [10], [11]). The condition  $\chi_*=\chi^*$  is trivially fulfilled, hence the following statement is valid.

**Proposition 2.** The cohomology space  $H_{\mathrm{H}}^{m-1}(\mathbf{CE}) = \mathbb{R} \cdot [u^{\mu}d_{\mu}x]$ .

Note, here 
$$H_H^{m-1}(\mathbf{CE}) \ni [u^{\mu}d_{\mu}x] \mapsto [u^1d_1x] \in \operatorname{Ker} D_1^{m-1}$$
.

**Remark 4.** The cohomology  $[u^{\mu}d_{\mu}x]$  is generated by the constraint CE =  $D_{\mu}u^{\mu} = 0$ .

At last, we have 
$$H^{m-1}(\Theta) \xrightarrow{D_1^{m-1}} H^{m-1}(\Theta) \to H^m_{\mathrm{H}}(\mathbf{CE}) \to 0$$
, i.e.,  $0 \to \mathrm{Im} \, D_1^{m-1} \to H^{m-1}(\Theta) \to H^m(\mathbf{CE}) \to 0$ ,

so  $H_{\mathrm{H}}^{m}(\mathbf{CE}) = H^{m-1}(\Theta) / \mathrm{Im} \, D_{1}^{m-1}$  Thus, the following theorem holds.

**Theorem 2.** The linear spaces of the cohomologies of the differential algebra  $(\mathcal{A}(CE); \mathfrak{D}_H(CE))$  are

$$H_{\mathrm{H}}^{q}(\mathbf{CE}) = \begin{cases} 0, & q < 0, 1 \leq q \leq m - 2, q > m; \\ \mathbb{R}, & q = 0; \\ \operatorname{Ker} D_{1}^{m-1} = \mathbb{R} \cdot [u^{\mu} d_{\mu} x], & q = m - 1; \\ H^{m-1}(\Theta) / \operatorname{Im} D_{1}^{m-1}, & q = m; \end{cases}$$

where  $H^{m-1}(\Theta)/\operatorname{Im} D_1^{m-1} = \mathcal{H}(\Theta)/\operatorname{Im} \tilde{D_1}$ .

#### 2.2 An additional constraint.

According to Remark 3, to apply the algebraic analysis to the Navier-Stokes equations, explicitly containing dependence on the pressure p, we need to add an additional constraint in the space CE. Having in mind the specific form of these equations, we choose the constraint defined by the equation

$$PE = \Delta p + \nabla \mathbf{u} \cdot \nabla \mathbf{u} = \delta^{\lambda \mu} \partial_{x^{\lambda}} \partial_{x^{\mu}} p + \partial_{x^{\lambda}} u^{\mu} \cdot \partial_{x^{\mu}} u^{\lambda} = 0$$

(see the integrability condition (4), we took into account the equation (2)). Namely, we set

$$\mathbf{PE} = \left\{ (x, \mathbf{u}, \mathbf{p}) \in \mathbf{B} \mid PE_{i} = \Delta p_{i} + D_{i} (u_{(\mu)}^{\lambda} u_{(\lambda)}^{\mu}) = 0, \ i \in \mathbb{I} \right\}. \tag{7}$$

where

• 
$$\Delta = \delta^{\lambda\mu} D_{\lambda} \circ D_{\mu} = \delta^{\lambda\mu} D_{(\lambda)+(\mu)} = \sum_{\mu} D_{2(\mu)},$$

• 
$$D_{\mathbf{i}}(u_{(\mu)}^{\lambda}u_{(\lambda)}^{\mu}) = \sum_{\mathbf{k}+\mathbf{l}=\mathbf{i}} {\mathbf{i} \choose \mathbf{k}} u_{\mathbf{k}+(\mu)}^{\lambda} u_{\mathbf{l}+(\lambda)}^{\mu}.$$

The subspace  $CPE = CE \cap PE \subset B$  generates the differential ideal

$$\mathcal{J} = \{ f \in \mathcal{A}(\mathbf{B}) \mid f|_{\mathbf{CPE}} = 0 \} = \{ f = g^i \cdot \mathrm{CE}_i + h^i \cdot \mathrm{PE}_i \mid g^i, h^i \in \mathcal{A}(\mathbf{B}) \}.$$

The factor-algebra  $\mathcal{A}(\mathbf{CPE}) = \mathcal{A}(\mathbf{B})/\mathcal{J}$  is the algebra of *smooth functions* on the space  $\mathbf{CPE}$ .

The Lie algebra  $\mathfrak{D}(\mathbf{CPE})$  of derivations of the algebra  $\mathcal{A}(\mathbf{CPE})$ , the Lie algebra  $\mathrm{Sym}(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathrm{H}}(\mathbf{CPE}))$  of the symmetries of the differential algebra  $(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathrm{H}}(\mathbf{CPE}))$ , the  $\mathbb{Z}$ -graded  $\mathcal{A}(\mathbf{CPE})$ -module  $\Omega_{\mathrm{H}}(\mathbf{CPE}) = \bigoplus_{q \in \mathbb{Z}} \Omega_{\mathrm{H}}^q(\mathbf{CPE})$  of horizontal differential forms and the horizontal differential  $d_{\mathrm{H}} \in \mathrm{End}_{\mathbb{R}}(\Omega_{\mathrm{H}}(\mathbf{CPE}))$  on the space  $\mathbf{CPE}$  are defined as above (see Subsection 2.1). Namely, here:

- $\mathfrak{D}_{\mathcal{J}}(\mathbf{B}) = \{ \zeta \in \mathfrak{D}(\mathbf{B}) \mid \zeta|_{\mathcal{J}} : \mathcal{J} \to \mathcal{J} \};$
- $\bullet \ \mathfrak{D}(\mathbf{B};\mathcal{J}) = \{\zeta \in \mathfrak{D}(\mathbf{B}) \mid \zeta : \mathcal{A}(\mathbf{B}) \to \mathcal{J}\};$
- $\mathfrak{D}(\mathbf{CPE}) = \mathfrak{D}_{\mathcal{J}}(\mathbf{B})/\mathfrak{D}(\mathbf{B}; \mathcal{J})$  is the Lie factor-algebra of derivations of the factor-algebra  $\mathcal{A}(\mathbf{CPE})$ ;
- $\bullet \ \mathfrak{D}(\mathbf{CPE}) = \mathfrak{D}_V(\mathbf{CPE}) \oplus_{\mathcal{A}(\mathbf{CPE})} \mathfrak{D}_H(\mathbf{CPE});$

- $\mathfrak{D}_{V}(CPE) = \{\bar{\zeta} \in \mathfrak{D}(CPE) \mid \zeta \in \mathfrak{D}_{V}(B) \cap \mathfrak{D}_{\mathcal{J}}(B)\};$
- $\mathfrak{D}_{\mathrm{H}}(\mathbf{CPE}) = \{ \bar{\zeta} \in \mathfrak{D}(\mathbf{CPE}) \mid \zeta \in \mathfrak{D}_{\mathrm{H}}(\mathbf{B}) \subset \mathfrak{D}_{\mathcal{J}}(\mathbf{B}) \};$
- Sym( $\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathrm{H}}(\mathbf{CPE})$ )= $\{\bar{\zeta} \in \mathfrak{D}_{\mathrm{V}}(\mathbf{CPE}) \mid [\bar{D}_{\mu}, \bar{\zeta}] = \bar{0} = \mathcal{J}, \ \mu \in \mathrm{M}\}.$

To write the  $\mathbb{Z}$ -graded  $\mathcal{A}(\mathbf{CPE})$ -module  $\Omega_H(\mathbf{CPE}) = \bigoplus_{q \in \mathbb{Z}} \Omega_H^q(\mathbf{CPE})$  and its cohomology spaces it is sufficient to replace  $\mathbf{CE}$  with  $\mathbf{CPE}$  in all relevant formulas. In particular,

$$\Omega_{\mathrm{H}}^{q}(\mathbf{CPE}) = \begin{cases}
0, & q < 0, q > m; \\
\mathcal{A}(\mathbf{CPE}), & q = 0; \\
\operatorname{Hom}_{\mathcal{A}(\mathbf{CPE})}(\wedge^{q}\mathfrak{D}_{\mathrm{H}}(\mathbf{CPE}); \mathcal{A}(\mathbf{CPE})), & 1 \leq q \leq m.
\end{cases}$$

The horizontal differential  $d_{\rm H} \in \operatorname{End}_{\mathbb{R}}(\Omega_{\rm H}(\mathbf{CPE}))$ ,  $d_{\rm H} \circ d_{\rm H} = 0$ , is defined by the rule,

$$d_{\mathrm{H}}^{q} = d_{\mathrm{H}}\big|_{\Omega_{\mathrm{H}}^{q}(\mathbf{CPE})} : \Omega_{\mathrm{H}}^{q}(\mathbf{CPE}) \to \Omega_{\mathrm{H}}^{q+1}(\mathbf{CPE}),$$
$$\bar{\omega}_{\mu_{1}\dots\mu_{q}} \cdot d\bar{x}^{\mu_{1}} \wedge \dots \wedge d\bar{x}^{\mu_{q}} \mapsto \bar{D}_{[\mu_{0}}\bar{\omega}_{\mu_{1}\dots\mu_{q}]} \cdot d\bar{x}^{\mu_{0}} \wedge \dots \wedge d\bar{x}^{\mu_{q}},$$

where  $d_{\mathrm{H}}^{q+1} \circ d_{\mathrm{H}}^{q} = 0, q \in \mathbb{Z}$ .

Now, let us define the global coordinates on the subspace  $\mathbf{CPE} \subset \mathbf{B}$ . Namely, we split  $\mathbb{I} = \mathbb{I}_1 \cup \mathbb{I}_1'$  (see Subsection 1.2) and get

- $\mathbb{R}_{\mathbb{I}} = \mathbb{R}_{\mathbb{I}_1} \times \mathbb{R}_{\mathbb{I}'_1};$
- $\bullet \ \mathbf{B} = X \times \mathbb{R}^1_{\mathbb{I}_0} \times \mathbb{R}^1_{\mathbb{I}_0'} \times \mathbb{R}^N_{\mathbb{I}} \times \mathbb{R}_{\mathbb{I}_1} \times \mathbb{R}_{\mathbb{I}_1'};$
- **CPE** = **CE**  $\cap$  **PE** = X ×  $\mathbb{R}^1_{\mathbb{I}_0}$  ×  $\mathbb{R}^N_{\mathbb{I}}$  ×  $\mathbb{R}_{\mathbb{I}_1}$ , i.e., the subspace **CPE** has the global coordinates  $(x, \mathbf{u}, \mathbf{p}) = \{x^{\mu}, u^1_{i_0}, u^{\alpha}_{i_1}, p_{i_1}\}$ , where the indices  $\mu \in \mathbb{M}, \ \alpha \in \mathbb{N}, \ i_0 \in \mathbb{I}_0, \ i_1 \in \mathbb{I}_1, \ i \in \mathbb{I}$ ;
- CE =  $u^{\mu}_{(\mu)} = 0$ , PE =  $\Delta p + u^{\lambda}_{(\mu)} u^{\mu}_{(\lambda)}|_{\mathbf{CE}} = p_{2(1)} + \Phi(\mathbf{u}, \mathbf{p}) = 0$ ;
- $\Phi(\mathbf{u}, \mathbf{p}) = \Delta' p + u^{\lambda}_{(\mu)} u^{\mu}_{(\lambda)} |_{\mathbf{CE}}, \quad \Delta' = \delta^{\alpha\beta} D_{\alpha} \circ D_{\beta};$
- $u_{\mathbf{i}'_0}^1 = -u_{\mathbf{i}'_0 (1) + (\alpha)}^{\alpha}, \ \mathbf{i}'_0 \in \mathbb{I}'_0, \ p_{\mathbf{i}'_1} = -D_{\mathbf{i}'_1 2(1)}\Phi(\mathbf{u}, \mathbf{p}), \ \mathbf{i}'_1 \in \mathbb{I}'_1;$
- $D_i(u_{(\mu)}^{\lambda}u_{(\lambda)}^{\mu}) = \sum_{k+l=i} \binom{i}{k} u_{k+(\mu)}^{\lambda} u_{l+(\lambda)}^{\mu}, i=i_1'-2(1) \in \mathbb{I}.$

In these coordinates, we have:

$$\begin{split} \bullet \ \ \mathcal{A}(\mathbf{CPE}) &= \mathcal{C}^{\infty}_{\mathrm{fin}}(\mathbf{CPE}) \\ &= \{ f \in \mathcal{A}(\mathbf{B}) \mid \partial_{u^{1}_{\mathbf{i}'_{0}}} f = 0, \partial_{p_{\mathbf{i}'_{1}}} f = 0, \mathbf{i}'_{0} \in \mathbb{I}'_{0}, \mathbf{i}'_{1} \in \mathbb{I}'_{1} \} \subset \mathcal{A}(\mathbf{B}); \end{split}$$

• 
$$\mathfrak{D}_{V}(\mathbf{CPE}) = \{ \zeta = \zeta_{i_0}^1 \partial_{u_{i_0}}^1 + \zeta_{i}^{\alpha} \partial_{u_{i}^{\alpha}}^{\alpha} + \zeta_{i_1} \partial_{p_{i_1}} \}, \text{ where } \zeta_{i_0}^1, \zeta_{i}^{\alpha}, \zeta_{i_1} \in \mathcal{A}(\mathbf{CPE}), i_0 \in \mathbb{I}_0, i_1 \in \mathbb{I}_1, i \in \mathbb{I}, \alpha \in \mathbb{N};$$

• 
$$D_{\mu} = \partial_{x^{\mu}} + u_{i_{0}+(\mu)}^{1} \partial_{u_{i_{0}}^{1}} + u_{i+(\mu)}^{\alpha} \partial_{u_{i}^{\alpha}} + p_{i_{1}+(\mu)} \partial_{p_{i_{1}}}, \quad \mu \in \mathcal{M}, \text{ where}$$
  
 $u_{i_{0}+(1)}^{1} = -u_{i_{0}+(\alpha)}^{\alpha}, \quad p_{i_{0}+2(1)} = -D_{i_{0}} \Phi(\mathbf{u}, \mathbf{p}), \quad i_{0} \in \mathbb{I}_{0}, \quad i_{1} = i_{0}, i_{0} + (1);$ 

• Sym(
$$\mathcal{A}(\mathbf{CPE})$$
,  $\mathfrak{D}_{\mathbf{H}}(\mathbf{CPE})$ )
$$= \{ \zeta = \operatorname{ev}_{\mathbf{f}} \in \mathfrak{D}_{\mathbf{V}}(\mathbf{CPE}) \mid [D_{\mu}, \operatorname{ev}_{\mathbf{f}}] = 0, \ \mu \in \mathbf{M} \}, \text{ where}$$

$$- \operatorname{ev}_{\mathbf{f}} = D_{\mathbf{i}_{0}} f^{1} \cdot \partial_{u_{\mathbf{i}_{0}}^{1}} + D_{\mathbf{i}} f^{\alpha} \cdot \partial_{u_{\mathbf{i}}^{\alpha}} + D_{\mathbf{i}_{1}} f \cdot \partial_{p_{\mathbf{i}_{1}}},$$

$$- \operatorname{f} = (f^{\mu}, f) \in \mathcal{A}(\mathbf{CPE})^{\mathbf{M}} \times \mathcal{A}(\mathbf{CPE}),$$

$$- D_{\mu} f^{\mu} = 0, \quad \Delta f + \operatorname{ev}_{\mathbf{f}} \left(u_{(\mu)}^{\lambda} u_{(\lambda)}^{\mu}\right) = 0,$$

$$- \Delta = \delta^{\lambda \mu} D_{(\lambda) + (\mu)}, \quad \operatorname{ev}_{\mathbf{f}} \left(u_{(\mu)}^{\lambda} u_{(\lambda)}^{\mu}\right) = 2u_{(\mu)}^{\lambda} D_{\lambda} f^{\mu} \Big|_{u_{(1)}^{1} = -u_{(2)}^{\alpha}};$$

- $\Omega_{\mathrm{H}}^{q}(\mathbf{CPE}) = \{ \omega = \omega_{\mu_{1}...\mu_{q}} \cdot dx^{\mu_{1}} \wedge ... \wedge dx^{\mu_{q}} \mid \omega_{\mu_{1}...\mu_{q}} \in \mathcal{A}(\mathbf{CPE}), + \text{s-s} \},$  $1 \leq q \leq m;$
- $d_{\mathrm{H}}^{q} = d_{\mathrm{H}}|_{\Omega_{\mathrm{H}}^{q}(\mathbf{CPE})} : \Omega_{\mathrm{H}}^{q}(\mathbf{CPE}) \to \Omega_{\mathrm{H}}^{q+1}(\mathbf{CPE}), \text{ where}$  $\omega = \omega_{\mu_{1}...\mu_{q}} \cdot dx^{\mu_{1}} \wedge \ldots \wedge dx^{\mu_{q}} \mapsto D_{[\mu_{0}}\omega_{\mu_{1}...\mu_{q}]} \cdot dx^{\mu_{0}} \wedge \ldots \wedge dx^{\mu_{q}}.$

Like it was done in the Subsection 2.1, to calculate the cohomology spaces  $H^q_{\mathrm{H}}(\mathbf{CPE})$  we introduce the auxiliary complex  $\{\Theta^q, d^q_{\Theta} \mid q \in \mathbb{Z}\}$  (see page 9). The only changes are

- the set **CE** is replaced with the set **CPE**;
- the basic derivations  $D_1, D_{\alpha}$  are as in the present section;
- the operator  $\tilde{D}_1 = D_1 + f^*$ ,

• 
$$D_1 = \partial_{x^1} + \text{ev}'_f$$
, where  $f = (f_0^1, f_{i_1}^{\alpha}, f_0, f_1)$ ,  
 $i_0 = (0, j), i_1 = (0, j), (1, j), i = (i^1, j), i^1 \in \mathbb{Z}_+, j \in \mathbb{J}$ ;

• 
$$\operatorname{ev}'_{\mathbf{f}} = D_{\mathbf{j}} f_0^1 \cdot \partial_{u_{0,\mathbf{j}}^1} + D_{\mathbf{j}} f_{i^1}^{\alpha} \cdot \partial_{u_{i_{1,\mathbf{j}}}^{\alpha}} + D_{\mathbf{j}} f_0 \cdot \partial_{p_{0,\mathbf{j}}} + D_{\mathbf{j}} f_1 \cdot \partial_{p_{1,\mathbf{j}}};$$

• 
$$f_0^1 = -u_{(\alpha)}^{\alpha}$$
,  $f_{i^1}^{\alpha} = u_{i^1+1,0}^{\alpha}$ ,  $f_0 = p_{1,0}$ ,  $f_1 = -\Phi(\mathbf{u}, \mathbf{p})$ ;

• 
$$\Delta' = \delta^{\alpha\beta} D_{(\alpha)+(\beta)}, \quad u^{\lambda}_{(\mu)} u^{\mu}_{(\lambda)} = u^{\alpha}_{(\alpha)} u^{\beta}_{(\beta)} + 2u^{1}_{(\alpha)} u^{\alpha}_{(1)} + u^{\alpha}_{(\beta)} u^{\beta}_{(\alpha)};$$

- $\chi \mapsto f^*\chi$ ,  $\chi = (\chi_1^0, \chi_\alpha^{i^1}, \chi^0, \chi^1)$ , where the set  $\chi$  is *finite*, i.e., only finitely many of its elements are nonzero,
- $(f^*\chi) = ((f^*\chi)^0_1, (f^*\chi)^{i^1}_\alpha, (f^*\chi)^0, (f^*\chi)^1);$

$$\begin{aligned} \bullet & & (\mathbf{f}^*\chi)_1^0 = 2D_\alpha(u_{(1)}^\alpha\chi^1), \\ & & (\mathbf{f}^*\chi)_\alpha^{i^1} = \delta_0^{i^1}D_\alpha\chi_1^0 + \chi_\alpha^{i^1-1} \\ & & & + 2\big(\delta_0^{i^1}D_\alpha(u_{(\beta)}^\beta\chi^1) - \delta_1^{i^1}u_{(\alpha)}^1\chi^1 + \delta_0^{i^1}D_\beta(u_{(\alpha)}^\beta\chi^1)\big), \\ & & (\mathbf{f}^*\chi)^0 = -\Delta'\chi^1, \quad (\mathbf{f}^*\chi)^1 = \chi^0, \end{aligned}$$

where  $\alpha, \beta \in \mathbb{N}, \ \lambda, \mu \in \mathbb{M}, \ i^1 \in \mathbb{Z}_+, \ j \in \mathbb{J}.$ 

The resulting system  $(D_1 + f^*)\chi = 0$  for unknown function  $\chi$  with a finite number of nonzero components reduces to

$$D_1 \chi_1^0 + (f^* \chi)_1^0 = 0, \quad D_1 \chi_{\alpha}^{i^1} + (f^* \chi)_{\alpha}^{i^1} = 0,$$
  
 $D_1 \chi^0 + (f^* \chi)^0 = 0, \quad D_1 \chi^1 + (f^* \chi)^1 = 0.$ 

The equation

$$D_{1}\chi_{\alpha}^{i^{1}} + (f^{*}\chi)_{\alpha}^{i^{1}} = D_{1}\chi_{\alpha}^{i^{1}} + \delta_{0}^{i^{1}}D_{\alpha}\chi_{1}^{0} + \chi_{\alpha}^{i^{1}-1} + 2(\delta_{0}^{i^{1}}D_{\alpha}(u_{(\beta)}^{\beta}\chi^{1}) - \delta_{1}^{i^{1}}u_{(\alpha)}^{1}\chi^{1} + \delta_{0}^{i^{1}}D_{\beta}(u_{(\alpha)}^{\beta}\chi^{1})) = 0$$

for  $i^1 \geq 2$ , gives  $\chi_{\alpha}^{i^1} = 0$  for  $i^1 \geq 1$ , for  $i^1 = 1$  it gives  $\chi_{\alpha}^0 = 2u_{(\alpha)}^1 \chi^1$ , and for  $i^1 = 0$  it takes the form  $D_1 \chi_{\alpha}^0 + D_{\alpha} \chi_1^0 + 2 \left( D_{\alpha} (u_{(\beta)}^{\beta} \chi^1) + D_{\beta} (u_{(\alpha)}^{\beta} \chi^1) \right) = 0$ . Further, the equation  $D_1 \chi^1 + (f^* \chi)^1 = D_1 \chi^1 + \chi^0 = 0$  gives  $\chi^0 = -D_1 \chi^1$ , the equation  $D_1 \chi^0 + (f^* \chi)^0 = -D_{2(1)} \chi^1 - \Delta' \chi^1 = -\Delta \chi^1 = 0$  gives  $\Delta \chi^1 = 0$ , and the last one equation  $D_1 \chi_1^0 + (f^* \chi)_1^0 = D_1 \chi_1^0 + 2D_{\alpha} (u_{(1)}^{\alpha} \chi^1) = 0$  gives  $D_1 \chi_1^0 + 2D_{\alpha} (u_{(1)}^{\alpha} \chi^1) = 0$ .

After some algebra we get

**Lemma 2.** The system  $(D_1 + f^*)\chi = 0$ ,  $\chi = (\chi_1^0, \chi_\alpha^{i^1}, \chi^0, \chi^1)$ , reduces to

$$\chi_{\alpha}^{0} = 2u_{(\alpha)}^{1}\chi^{1}, \quad \chi_{\alpha}^{i^{1}} = 0, \quad \chi^{0} = -D_{1}\chi^{1}, \quad i^{1} \ge 1, \ \alpha \in \mathbb{N},$$
 (8)

where  $\chi^1, \chi^0_1$  are solutions of the system

$$\Delta \chi^1 = 0, \quad (u^{\mu}_{(1)} D_{\mu} D_{\alpha} - u^{\mu}_{(\alpha)} D_{\mu} D_1) \chi^1 = 0, \quad \alpha \in \mathbb{N},$$
 (9)

$$D_1 \chi_1^0 = -2D_{\alpha}(u_{(1)}^{\alpha} \chi^1), \ D_{\alpha} \chi_1^0 = -2(u_{(\alpha)}^{\mu} D_{\mu} \chi^1 + D_{\alpha}(u_{(\beta)}^{\beta} \chi^1)), \ \alpha \in \mathbb{N}.$$
 (10)

**Remark 5.** The right subsystem of the system (9) is the *non-trivial com*patibility condition of the system (10).

**Remark 6.** The system (9) has the trivial solution  $\chi^1 = 0$ , then the system (10) gives  $\chi_1^0 = \text{const}$ , and we get the solution  $\chi = (1, 0, 0, 0)$  of the full system  $(D_1 + f^*)\chi = 0$  (see the similar result in Subsection 2.1).

**Theorem 3.** The linear spaces of the cohomologies of the differential algebra  $(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_H(\mathbf{CPE}))$  are (cf., Theorem 1 and Theorem 2)

$$H_{\mathrm{H}}^{q}(\mathbf{CPE}) = \begin{cases} 0, & q < 0, 1 \leq q \leq m-2, q > m; \\ \mathbb{R}, & q = 0; \\ \mathrm{Ker}\,D_{1}^{m-1} \simeq \mathcal{S} \cap \mathcal{H}, & q = m-1; \\ H^{m-1}(\Theta) \big/ \operatorname{Im}D_{1}^{m-1}, & q = m; \end{cases}$$

where

- $S = Sol(D_1 + f^*)$  is the linear space of solutions  $\chi = (\chi_1^0, \chi_{\alpha}^{i^1}, \chi^0, \chi^1)$  of the linear system  $D_1\chi + f^*\chi = 0$  (the mapping  $f^*$  is defined above);
- $\mathcal{H} = \{\chi = (\chi_1^0, \chi_{\alpha}^{i^1}, \chi^0, \chi^1) \mid \chi_* = \chi^* \}$  is the Helmholtz space of the differential algebra  $(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathbf{H}}(\mathbf{CPE}));$
- $H_{\mathrm{H}}^{m-1}(\mathbf{CPE}) = \mathcal{S} \cap \mathcal{H}, \quad [J^{\mu} \cdot d_{\mu}x] \mapsto (\delta_{u_0^1}J^1, \delta_{u_{i_1}^{\alpha}}J^1, \delta_{p_0}J^1, \delta_{p_1}J^1)$  (remind, cohomologies are defined up to isomorphisms).

**Remark 7.** In particular,  $H_{\rm H}^{m-1}(\mathbf{CPE}) \ni [u^{\mu}d_{\mu}x]$ , see Remark 6. Moreover, the additional constraint PE =  $\Delta p + u_{(\mu)}^{\lambda} u_{(\lambda)}^{\mu} = 0$  generates the additional cohomology  $[F^{\mu}d_{\mu}x] \in H_{\rm H}^{m-1}(\mathbf{CPE})$ , where  $F^{\mu} = -u^{\lambda}u_{(\lambda)}^{\mu} + \nu\Delta u^{\mu} + p^{(\mu)}$ , since here

$$D_{\mu}F^{\mu} = -\left(\Delta p + u^{\lambda}_{(\mu)}u^{\mu}_{(\lambda)}\right) + \left(-u^{\lambda}D_{\lambda}u^{\mu}_{(\mu)} + \nu\Delta u^{\mu}_{(\mu)}\right)$$
$$= -\text{PE} + \left(\nu\Delta - u^{\lambda}D_{\lambda}\right)\text{CE} = 0$$

due to the constraints PE = 0 and CE = 0. Here,  $p^{(\mu)} = D^{\mu}p = \delta^{\mu\lambda}D_{\lambda}p = D_{\mu}p = p_{(\mu)}$  (the euclidean metrics).

Remark 8. Every symmetry  $\operatorname{ev}_f \in \operatorname{Sym}(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_H(\mathbf{CPE}))$ ,  $f = (f^{\mu}, f)$ , generates (possibly trivial) cohomology  $[f^{\mu}d_{\mu}x] \in H^{m-1}_H(\mathbf{CPE})$ , because in this case  $D_{\mu}f^{\mu} = 0$  by definition.

# 3 Navier-Stokes equations as evolution in the space CPE.

## 3.1 Evolution in the space CPE.

The evolution in the space

$$\mathbf{CPE} = T \times X \times \left(\mathbb{R}^1_{\mathbb{I}_0} \times \mathbb{R}^N_{\mathbb{I}}\right) \times \mathbb{R}_{\mathbb{I}_1} = \left\{t, \mathbf{x} = (x^\mu), \mathbf{u} = (u^1_{i_0}, u^\alpha_i), \mathbf{p} = (p_{i_1})\right\}$$

(M = 2, 3) is governed by an evolution derivation  $D_t = \partial_t + ev_E$ , where

- we added the time variable  $t \in T = \mathbb{R}$ , one may assume that it was present from the start as a parameter;
- $E = (E^{\mu}, E) \in \mathcal{A}^{M}(\mathbf{CPE}) \times \mathcal{A}(\mathbf{CPE});$
- $D_{\mu}E^{\mu} = 0$ ,  $\Delta E + 2(u^{\lambda}_{(\mu)}D_{\lambda}E^{\mu})|_{u^{1}_{(1)} = -u^{\alpha}_{(\alpha)}} = 0$ ,  $\Delta = \delta^{\lambda\mu}D_{(\lambda)+(\mu)}$ ;
- $\operatorname{ev}_{\mathrm{E}} = D_{\mathrm{i}_0} E^1 \cdot \partial_{u_{\mathrm{i}_0}^1} + D_{\mathrm{i}} E^{\alpha} \cdot \partial_{u_{\mathrm{i}}^{\alpha}} + D_{\mathrm{i}_1} E \cdot \partial_{p_{\mathrm{i}_1}} \in \operatorname{Sym}(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathrm{H}}(\mathbf{CPE}))$

(see Subsection 2.2, remind also Proposition 1 and Remark 3).

**Remark 9.** In particular,  $D_t u^{\mu} = F^{\mu}$ ,  $\mu \in M$ , and  $D_t p = F$ , while

$$D_t \text{CE} = D_t \left( D_\mu u^\mu \right) = D_\mu E^\mu, \quad D_t \text{PE} = D_t \left( \Delta p + u^\lambda_{(\mu)} u^\mu_{(\lambda)} \right) = \Delta E + 2 u^\lambda_{(\mu)} D_\lambda E^\mu.$$

There is defined the differential algebra  $(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{E}(\mathbf{CPE}))$ , where

- $\bullet \ \mathfrak{D}(\mathbf{CPE}) = \mathfrak{D}_V(\mathbf{CPE}) \oplus_{\mathcal{A}(\mathbf{CPE})} \mathfrak{D}_E(\mathbf{CPE});$
- $\mathfrak{D}_{V}(\mathbf{CPE}) = \{ \zeta = \zeta_{i_0}^1 \partial_{u_{i_0}}^1 + \zeta_i^{\alpha} \partial_{u_i^{\alpha}} + \zeta_{i_1} \partial_{p_{i_1}} \mid \zeta_{i_0}^1, \zeta_i^{\alpha}, \zeta_{i_1} \in \mathcal{A}(\mathbf{CPE}) \};$
- $\mathfrak{D}_{\mathrm{E}}(\mathbf{CPE})$  has the  $\mathcal{A}(\mathbf{CPE})$ -basis  $\{D_t, D_\mu \mid \mu \in \mathbf{M}\}$ , the basic time derivation  $D_t = \partial_t + \mathrm{ev}_{\mathrm{E}}, [D_t, D_\mu] = 0, \, \mu \in \mathbf{M}$ , so

$$\mathfrak{D}_{\mathrm{E}}(\mathbf{CPE}) = \{ \zeta = \zeta^t D_t + \zeta^{\mu} D_{\mu} \mid \zeta^t, \zeta^{\mu} \in \mathcal{A}(\mathbf{CPE}) \}.$$

The Lie algebra of symmetries here is

$$\mathrm{Sym}(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathrm{E}}(\mathbf{CPE}))$$

$$= \{ \zeta = \text{ev}_f \in \text{Sym}(\mathcal{A}(\mathbf{CPE}, \mathfrak{D}_H(\mathbf{CPE})) \mid [D_t, \text{ev}_f] = 0 \},$$

where the condition  $[D_t, ev_f] = 0$  reduces to the equation  $(D_t - E_*)f = 0$ . In more detail, here

• 
$$E_* : \mathcal{A}(CPE)^M \times \mathcal{A}(CPE) \to \mathcal{A}(CPE)^M \times \mathcal{A}(CPE)$$
,

• 
$$f = (f^{\mu}, f) \mapsto F_* f = ((F_* f)^{\mu}, (F_* f)),$$

• 
$$(E_*f)^{\mu} = \partial_{u_{i_0}^1} E^{\mu} \cdot D_{i_0} f^1 + \partial_{u_i^{\alpha}} E^{\mu} \cdot D_i f^{\alpha} + \partial_{p_{i_1}} E^{\mu} \cdot D_{i_1} f$$
,

$$\bullet \ (\mathbf{E}_*\mathbf{f}) = \partial_{u_{\mathbf{i}_0}^1} E \cdot D_{\mathbf{i}_0} f^1 + \partial_{u_{\mathbf{i}}^\alpha} E \cdot D_{\mathbf{i}} f^\alpha + \partial_{p_{\mathbf{i}_1}} E \cdot D_{\mathbf{i}_1} f.$$

As above, to study the cohomology spaces  $H_{\rm E}^q(\mathbf{CPE}) = \operatorname{Ker} d_{\rm E}^q / \operatorname{Im} d_{\rm E}^{q-1}$ ,  $q \in \mathbb{Z}$ , of the differential algebra  $(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\rm E}(\mathbf{CPE}))$  we shall use the decomposition (see pages 9 and 15):

• 
$$\Omega_{\mathrm{H}}^{q} = \Omega_{\mathrm{H}}^{q}(\mathbf{CPE})$$
  
=  $\{\omega_{\mathrm{H}}^{q} = \omega_{\mu_{1}...\mu_{q}} \cdot dx^{\mu_{1}} \wedge ... \wedge dx^{\mu_{q}} \mid \omega_{\mu_{1}...\mu_{q}} \in \mathcal{A}(\mathbf{CPE}), + \text{s-s}\};$ 

• 
$$\Omega_{\mathrm{E}}^{q} = \Omega_{\mathrm{E}}^{q}(\mathbf{CPE}) = dt \wedge \Omega_{\mathrm{H}}^{q-1} \oplus_{\mathcal{A}(\mathbf{CPE})} \Omega_{\mathrm{H}}^{q}, \quad q \in \mathbb{Z};$$

• 
$$0 \to \Omega_{\mathrm{H}}^{q-1} \xrightarrow{\iota^{q-1}} \Omega_{\mathrm{E}}^{q} \xrightarrow{\pi^{q}} \Omega_{\mathrm{H}}^{q} \to 0,$$
  
 $\omega_{\mathrm{H}}^{q-1} \mapsto \omega_{\mathrm{E}}^{q} = \iota^{q-1} \omega_{\mathrm{H}}^{q-1} = (-1)^{q-1} dt \wedge \omega_{\mathrm{H}}^{q-1},$   
 $\omega_{\mathrm{E}}^{q} = dt \wedge \omega_{\mathrm{H}}^{q-1} + \omega_{\mathrm{H}}^{q} \mapsto \pi^{q} \omega_{\mathrm{E}}^{q} = \omega_{\mathrm{H}}^{q};$ 

• 
$$d_{\mathrm{E}}^{q} = d_{t}^{q} + d_{\mathrm{H}}^{q} : \Omega_{\mathrm{E}}^{q} \to \Omega_{\mathrm{E}}^{q+1}, \quad d_{t} = dt \wedge D_{t}, \quad d_{\mathrm{H}} = dx^{\mu} \wedge D_{\mu},$$
  
 $\omega_{\mathrm{E}}^{q} = dt \wedge \omega_{\mathrm{H}}^{q-1} + \omega_{\mathrm{H}}^{q} \mapsto d_{\mathrm{E}}\omega_{\mathrm{E}}^{q} = dt \wedge (D_{t}\omega_{\mathrm{H}}^{q} - d_{\mathrm{H}}\omega_{\mathrm{H}}^{q-1}) + d_{\mathrm{H}}\omega_{\mathrm{H}}^{q}.$ 

These constructions lead to the commutative diagram with the long exact sequence of the cohomology spaces

$$0 \to H_{\mathrm{E}}^{0}(\mathbf{CPE}) \to H_{\mathrm{H}}^{0}(\mathbf{CPE}) \xrightarrow{D_{t}^{0}}$$

$$\xrightarrow{D_{t}^{0}} H_{\mathrm{H}}^{0}(\mathbf{CPE}) \to H_{\mathrm{E}}^{1}(\mathbf{CPE}) \to H_{\mathrm{H}}^{1}(\mathbf{CPE}) \xrightarrow{D_{t}^{1}} \dots$$

$$\dots \xrightarrow{D_{t}^{m-2}} H_{\mathrm{H}}^{m-2}(\mathbf{CPE}) \to H_{\mathrm{E}}^{m-1}(\mathbf{CPE}) \to H_{\mathrm{H}}^{m-1}(\mathbf{CPE}) \xrightarrow{D_{t}^{m-1}}$$

$$\xrightarrow{D_{t}^{m-1}} H_{\mathrm{H}}^{m-1}(\mathbf{CPE}) \to H_{\mathrm{E}}^{m}(\mathbf{CPE}) \to H_{\mathrm{H}}^{m}(\mathbf{CPE}) \xrightarrow{D_{t}^{m}}$$

$$\xrightarrow{D_{t}^{m}} H_{\mathrm{H}}^{m}(\mathbf{CPE}) \to H_{\mathrm{E}}^{m+1}(\mathbf{CPE}) \to 0,$$

where  $D_t^q: H^q_{\mathrm{H}}(\mathbf{CPE}) \to H^q_{\mathrm{H}}(\mathbf{CPE})$  by the component-wise rule

$$D_t^q[\omega_{\mu_1\dots\mu_q}\cdot dx^{\mu_1}\wedge\ldots\wedge dx^{\mu_q}]=[(D_t\omega_{\mu_1\dots\mu_q})\cdot dx^{\mu_1}\wedge\ldots\wedge dx^{\mu_q}].$$

**Theorem 4.** The linear spaces of cohomologies of the differential algebra  $(\mathcal{A}(\mathbf{CPE}); \mathfrak{D}_{E}(\mathbf{CPE}))$  are

$$H_{\mathrm{E}}^{q}(\mathbf{CPE}) = \begin{cases} 0, & q < 0, 1 \leq q \leq m-2, q > m+1; \\ \mathbb{R}, & q = 0; \\ \mathrm{Ker} \ D_{t}^{m-1}, & q = m-1; \\ H_{\mathrm{H}}^{m}(\mathbf{CPE}) \big/ \mathrm{Im} \ D_{t}^{m}, & q = m+1; \end{cases}$$

while in the case q = m one has  $H_{\rm E}^m(\mathbf{CPE})/\operatorname{Im} H_{\rm H}^{m-1}(\mathbf{CPE}) = \operatorname{Ker} D_t^m$ .

Proof. Indeed, the exact subsequence  $0 \to H_{\rm E}^0(\mathbf{CPE}) \to \mathrm{Ker}\,D_t^0 \to 0$  gives  $H_{\rm E}^0(\mathbf{CPE}) = \mathbb{R}$  (remind, we added the time variable  $t \in T = \mathbb{R}$ , so now  $H_{\rm H}^0(\mathbf{CPE}) = \mathcal{T} = \{\phi(t) \in \mathcal{C}^\infty(\mathbb{R})\}$ ). Further, the exact subsequence

$$H^0_{\mathrm{H}}(\mathbf{CPE}) \xrightarrow{D^0_t} H^0_{\mathrm{H}}(\mathbf{CPE}) \to H^1_{\mathrm{E}}(\mathbf{CPE}) \to H^1_{\mathrm{H}}(\mathbf{CPE}) = 0$$

gives  $H_{\rm E}^1(\mathbf{CPE}) = \mathcal{T}/\operatorname{Im} D_t^0 = 0$ . Now, the exact subsequence

$$H^{q-1}_{\mathrm{H}}(\mathbf{CPE}) \to H^q_{\mathrm{E}}(\mathbf{CPE}) \to H^q_{\mathrm{H}}(\mathbf{CPE})$$

gives  $H_{\rm E}^q(\mathbf{CPE})=0$  for  $2\leq q\leq m-2$  because in this case  $H_{\rm H}^{q-1}(\mathbf{CPE})=H_{\rm H}^q(\mathbf{CPE})=0$ . Then, the exact subsequence

$$0 = H_{\mathrm{H}}^{m-2}(\mathbf{CPE}) \to H_{\mathrm{E}}^{m-1}(\mathbf{CPE}) \to H_{\mathrm{H}}^{m-1}(\mathbf{CPE}) \xrightarrow{D_t^{m-1}} \dots$$

gives  $H_{\rm E}^{m-1}(\mathbf{CPE})=\operatorname{Ker}D_t^{m-1}$ . At last, for q=m,m+1 the statements follow from the exact subsequence  $H_{\rm H}^{m-1}(\mathbf{CPE})\to H_{\rm E}^m(\mathbf{CPE})\to H_{\rm H}^m(\mathbf{CPE})\to H_{\rm E}^{m}\to 0$  and from the exact subsequence  $0\to\operatorname{Im}D_t^m\to H_{\rm H}^m(\mathbf{CPE})\to H_{\rm E}^{m+1}(\mathbf{CPE})\to 0$ .

**Remark 10.** To calculate  $\operatorname{Ker} D_t^{m-1}$  one may use the technics from lemmas 1 and 2. The cases q=m,m+1 in this approach are uninformative. They demand a special study.

# 3.2 Navier-Stokes equations as the evolution with constraints.

We consider the Navier-Stokes system (1)-(4) as the evolution process governed by the equation (1) in the divergence-free space with the inner constraint (4).

The algebraic counterpart of the equation (1) is the symmetry

$$\operatorname{ev}_{\mathrm{E}} = D_{\mathrm{i}_{0}} E^{1} \cdot \partial_{u_{\mathrm{i}_{0}}^{1}} + D_{\mathrm{i}} E^{\alpha} \cdot \partial_{u_{\mathrm{i}}^{\alpha}} + D_{\mathrm{i}_{1}} E \cdot \partial_{p_{\mathrm{i}_{1}}} \in \operatorname{Sym}(\mathcal{A}(\mathbf{CPE}), \mathfrak{D}_{\mathrm{H}}(\mathbf{CPE})),$$

where

- $E = (E^{\mu}, E) \in \mathcal{A}(\mathbf{CPE})^{M} \times \mathcal{A}(\mathbf{CPE});$
- $E^{\mu} = -u^{\lambda} \nabla_{\lambda} u^{\mu} + \nu \Delta u^{\mu} \nabla^{\mu} p = -u^{\lambda} u^{\mu}_{(\lambda)} + \nu \Delta u^{\mu} p_{(\mu)};$
- $u^1_{(1)} = -u^{\alpha}_{(\alpha)}$ ,  $\Delta u^{\mu} = \sum_{\lambda} u^{\mu}_{2(\lambda)}$ ,  $\Delta u^1 = -u^{\alpha}_{(1)+(\alpha)} + \sum_{\alpha} u^1_{2(\alpha)}$ ;
- $\nabla^{\mu} = \delta^{\mu\nu} \nabla_{\nu} = \nabla_{\mu}$  (the euclidean metrics), so  $p^{(\mu)} = p_{(\mu)}$ ;
- E to be defined from the condition  $ev_E \in Sym(\mathcal{A}(CPE), \mathfrak{D}_H(CPE))$ .

One can easily check that here  $D_{\mu}E^{\mu}=0$ . On the other hand the condition

$$\Delta E + \text{ev}_{E}(u_{(\mu)}^{\lambda}u_{(\lambda)}^{\mu}) = \Delta E + 2u_{(\mu)}^{\lambda}D_{\lambda}E^{\mu} = 0$$
 (11)

is the Poisson equation for the component E (cf., Remark 1).

Thus, for the algebraic analysis of the Navier-Stokes equations, one can use the technique outlined in the previous subsection.

#### 3.3 Conclusion.

It can be seen from the above constructions that the Navier-Stokes equations are subject to meaningful analysis within the framework of the algebraic approach to differential equations. The resulting equations for finding algebraic characteristics of Navier-Stokes equations, such as symmetries and cohomologies, are essentially complicated. One may hope to find their partial solutions at least, especially using analytical computational packets (Mathematica, for example).

### References

- [1] L.I. Sedov, A course in continuum mechanics. Volume 1. Basic equations and analytical techniques. Walters—Noordhoff Publishing, Groningen, 1971.
- [2] L.D. Landau and E.M. Lifshitz, Fluid mechanics. Course of theoretical physics. Volume 6. Pergamon Press, Oxford, 1987.
- [3] Tai-Peng Tsai, Lectures on Navier-Stokes equations, AMS, Providence, Rhode Island, 2018.
- [4] Pierre Gilles Lemarié-Rieusset, The Navier-Stokes Problem in the 21st Century, CRC Press, Boca Raton London New York, 2016.
- [5] M.V. Korobkov, K. Pileckas, V.V. Pukhnachev, and R. Russo, The flux problem for the Navier–Stokes equations, Russian Math. Surveys 69:6 (2014), 1065–1122.
- [6] C.L. Fefferman, J.C. Robinson, J.L. Rodrigo (eds), Partial differential equations in fluid mechanics, London Mathematical Society Lecture Note Series: 452, Cambridge University Press, 2018.
- [7] W.M. Seiler, Involution, Springer-Verlag, Berlin, Heidelberg, 2010.
- [8] P.J. Olver, Applications of Lie groups to differential equations, Springer-Verlag, New York, 1986,1993.
- [9] S. Mac Lane, Homology, Springer-Verlag, Berlin, Heidelberg, 1963, 1994.
- [10] V.V. Zharinov, "Conservation laws of evolution systems", Theoret. and Math. Phys., 68:2 (1986), 745–751.
- [11] V.V. Zharinov, "Conservation laws, differential identities, and constraints of partial differential equations", Theoret. and Math. Phys., 185:2 (2015), 1557–1581.