# Some nonexistence results for space-time fractional Schrödinger equations without gauge invariance

Mokhtar Kirane, Ahmad Z. Fino<sup>1</sup>

#### Abstract

In this paper, we consider the Cauchy problem in  $\mathbb{R}^N$ ,  $N \geq 1$ , for semi-linear Schrödinger equations with space-time fractional derivatives. We discuss the nonexistence of global  $L^1$  or  $L^2$  weak solutions in the subcritical and critical cases under some conditions on the initial data and the nonlinear term. Furthermore, the nonexistence of local  $L^1$  or  $L^2$  weak solutions in the supercritical case are studied.

MSC[2020]: 35A01, 26A33

**Keywords:** Schrödinger equations, Fractional derivatives and integrals, test function method, nonexistence of global solution

#### 1 Introduction

In this paper, we consider the problem

$$\begin{cases}
i^{\alpha} {}^{c} D_{0|t}^{\alpha} u - (-\Delta)^{\beta/2} u = \lambda |u|^{p}, & (t, x) \in (0, T) \times \mathbb{R}^{N}, \\
u(x, 0) = \varepsilon u_{0}(x), & x \in \mathbb{R}^{N},
\end{cases}$$
(1)

where u is a complex-valued unknown function of (t,x),  $0 < \alpha < 1$ ,  $0 < \beta < 2$ ,  $N \ge 1$ , p > 1, T > 0,  $\varepsilon > 0$ ,  $\lambda \in \mathbb{C} \setminus \{0\}$ , and  $i^{\alpha}$  is the principal part of  $i^{\alpha}$ , i.e.

$$i^{\alpha} = \cos(\frac{\alpha\pi}{2}) + i\cos(\frac{\alpha\pi}{2}),$$

 $^cD^{\alpha}_{0|t}$  is the Caputo fractional derivative and  $(-\Delta)^{\beta/2}:L^2(\mathbb{R}^N)\to L^2(\mathbb{R}^N)$  is the fractional Laplacian which can be defined by a pointwise representation as given in Definition 6 below.

Different fractional generalizations of the Schrödinger equation appeared in the literature: The classical Schrödinger equations with nongauge power nonlinearity, i.e. (1) with  $\alpha=1$  and  $\beta=2$ , has been studied by Ikeda and Wakasugi [8] and Ikeda and Inui [9, 10], the spatial fractional Schrödinger equation involving fractional order space derivatives, i.e. (1) with  $\alpha=1$  and  $\beta\in(0,2)$ , has been investigated in [4, 5, 14, 15, 16], the fractional temporal Schrödinger equation involving a fractional time derivative, i.e.  $\alpha\in(0,1)$  and  $\beta=2$ , has been studied in [25, 19, 20], the semirelativistic Schrödinger equation with nongauge invariant power nonlinearity, i.e. (1) with  $\alpha=1$  and  $\beta=1/2$ , got interest by Fujiwara [6], Inui [11], Fujiwara and Ozawa [7], and the spatio-temporal fractional Schrödinger equation with both time and space fractional derivatives attracted the attention of [2, 22].

<sup>\*</sup>Department of Mathematics, Faculty of Arts and Science, Khalifa University, P.O. Box: 127788, Abu Dhabi, UAE; mokhtar.kirane@ku.ac.ae

Department of Mathematics, Sultan Qaboos University, FracDiff Research Group (DR/RG/03), P.O. Box 46, Al-Khoud 123, Muscat, Oman; ahmad.fino01@gmail.com; a.fino@squ.edu.om

The expected critical exponent can be determined by the following scaling argument: If u(x,t) is a solution of (1) with initial data  $u_0$ , then

$$v(t,x) = \gamma^{\frac{\beta}{p-1}} u(\gamma^{\beta/\alpha}t, \gamma x),$$

for all  $\gamma > 0$ , is also a solution of (1) with initial data  $v_0(x) = v(0,x) = \gamma^{\frac{\beta}{p-1}} u_0(\gamma x)$ , for all  $x \in \mathbb{R}^N$ . We choose  $p = p_s$  such that we get an invariant  $H^s$ -norm of the initial data:

$$||v_0||_{H^s} = \gamma^{\frac{\beta}{p-1} - \frac{N-2s}{2}} ||u_0||_{H^s} = ||u_0||_{H^s};$$

this happens if and only if

$$p = p_s = 1 + \frac{2\beta}{N - 2s}.$$

Therefore, the case  $p = p_s$  is called  $H^s$ -critical case; the case  $p < p_s$  (resp.  $p > p_s$ ) is called  $H^s$ -subcritical case (resp.  $H^s$ -supercritical case). On the other hand, the Fujita critical exponent for the corresponding heat equation with fractional Laplacian is

$$p_F = 1 + \frac{\beta}{N}.$$

Our main goal is to study the nonexistence of  $L^1$  or  $L^2$  global weak solutions under the condition  $p \leq p_s$  or  $p \leq p_F$  as well as the nonexistence of  $L^1$  or  $L^2$  local weak solutions under the condition that  $p > p_s$  or  $p > p_F$  (see e.g. [10]), using the test function method (see e.g. [24]) or a fractional differential equation approach (i.e. construct a fractional differential equation for a new function and using comparison principle). The local existence for (1) is expected in the  $H^s$ -subcritical case, but this is not our case. We refer the reader to [8, Appendix] by using the Strichartz estimates recently studied by Lee [17].

$$X_T = \{ \varphi \in C([0,\infty), H^{\beta}(\mathbb{R}^N)) \cap C^1([0,\infty), L^2(\mathbb{R}^N)), \text{ such that supp} \varphi \subset Q_T, \varphi \text{ is } \mathbb{R}\text{-valued} \},$$

and

$$Y_T = \{ \varphi \in C([0,\infty), H^{\beta}(\mathbb{R}^N)) \cap C^1([0,\infty), L^{\infty}(\mathbb{R}^N)), \text{ such that supp} \varphi \subset Q_T, \varphi \text{ is } \mathbb{R}\text{-valued} \},$$

where  $Q_T := [0,T] \times \mathbb{R}^N$  and the fractional Sobolev space  $H^{\beta}(\mathbb{R}^N)$  is defined by

$$H^{\beta}(\mathbb{R}^N) = \{ u \in L^2(\mathbb{R}^N); (-\Delta)^{\beta/2} u \in L^2(\mathbb{R}^N) \}.$$

**Definition 1** ( $L^2$ -weak solution). Let  $u_0 \in L^2(\mathbb{R}^N)$  and T > 0. We say that u is an  $L^2$ -weak solution of (1) if

$$u \in L^1((0,T), L^2(\mathbb{R}^N)) \cap L^p((0,T), L^{2p}(\mathbb{R}^N)),$$

and

$$\lambda \int_{Q_T} |u|^p \varphi(t, x) dt dx + \varepsilon i^\alpha \int_{Q_T} u_0(x)^c D_{t|T}^\alpha \varphi(t, x) dt dx$$

$$= i^\alpha \int_{Q_T} u^c D_{t|T}^\alpha \varphi dt dx - \int_{Q_T} u(-\Delta)^{\beta/2} \varphi(t, x) dt dx, \tag{2}$$

holds for all  $\varphi \in X_T$ . We denote the lifespan for the  $L^2$ -weak solution by

$$T_w(\varepsilon) := \sup\{T \in (0, \infty]; \text{ there exists a unique } L^2\text{-weak solution } u \text{ to } (1)\}.$$

Moreover, if T > 0 can be arbitrary chosen, i.e.  $T_w(\varepsilon) = \infty$ , then u is called a global  $L^2$ -weak solution of (1).

**Definition 2** ( $L^1$ -weak solution). Let  $u_0 \in L^1(\mathbb{R}^N)$  and T > 0. We say that u is an  $L^1$ -weak solution of (1) if  $u, |u|^p \in L^1((0,T), L^1(\mathbb{R}^N))$  and

$$\lambda \int_{Q_T} |u|^p \varphi(t, x) dt dx + \varepsilon i^\alpha \int_{Q_T} u_0(x)^c D_{t|T}^\alpha \varphi(t, x) dt dx$$

$$= i^\alpha \int_{Q_T} u^c D_{t|T}^\alpha \varphi dt dx - \int_{Q_T} u(-\Delta)^{\beta/2} \varphi(t, x) dt dx,$$
(3)

holds for all  $\varphi \in Y_T$ . We denote the lifespan for the  $L^1$ -weak solution by

 $\overline{T}_w(\varepsilon) := \sup\{T \in (0, \infty]; \text{ there exists a unique } L^1\text{-weak solution } u \text{ to } (1)\}.$ 

Moreover, if T > 0 can be arbitrary chosen, i.e.  $\overline{T}_w(\varepsilon) = \infty$ , then u is called a global  $L^1$ -weak solution to (1).

#### 2 Preliminaries

**Definition 3** (Absolutely continuous functions). [21, Chapter 1]

A function  $g:[a,b] \to \mathbb{R}$  with  $a,b \in \mathbb{R}$ , is absolutely continuous if and only if there exists a Lebesgue summable function  $\psi \in L^1(a,b)$  such that

$$g(t) = g(a) + \int_a^t \psi(s) ds$$
, for all  $t \in [a, b]$ .

The space of these functions is denoted by AC[a, b].

**Definition 4** (Riemann-Liouville fractional integrals). [21, Chapter 1]

Let  $g \in L^1(0,T)$  with T > 0. The Riemann-Liouville left- and right-sided fractional integrals of order  $\sigma \in (0,1)$  are, respectively, defined by

$$I_{0|t}^{\sigma}g(t) := \frac{1}{\Gamma(\sigma)} \int_{0}^{t} (t-s)^{-(1-\sigma)}g(s) ds, \quad t > 0,$$

and

$$I_{t|T}^{\sigma}g(t) := \frac{1}{\Gamma(\sigma)} \int_t^T (s-t)^{-(1-\sigma)} g(s) \, ds, \quad t < T,$$

where  $\Gamma$  is the Euler gamma function.

**Definition 5** (Caputo fractional derivatives). [21, Chapter 1]

Let  $f \in AC[0,T]$  with T > 0. The Caputo left- and right-sided fractional derivatives of order  $\delta \in (0,1)$  exists almost everywhere on [0,T] and defined, respectively, by

$${}^{c}D_{0|t}^{\delta}f(t) := \frac{d}{dt}I_{0|t}^{1-\delta}[f(t) - f(0)] = I_{0|t}^{1-\delta}[f'(t)], \quad t > 0,$$

and

$$^{c}D_{t|T}^{\delta}f(t) := -\frac{d}{dt}I_{t|T}^{1-\delta}[f(t) - f(T)] = -I_{t|T}^{1-\delta}[f'(t)], \quad t < T.$$

**Lemma 1.** [12, Lemma 2.22, p. 96]

Let  $0 < \delta < 1$  and T > 0. If  $f \in AC[0,T]$  or  $f \in C^1[0,T]$ , then

$$I_{0|t}^{\delta}{}^{c}D_{0|t}^{\delta}f(t) = f(t) - f(0). \tag{4}$$

Given T > 0, let us define the function  $w : [0, T] \to \mathbb{R}$  by the following formula:

$$w(t) = (1 - t/T)^{\eta} \quad \text{for all } 0 \le t \le T, \tag{5}$$

where  $\eta \gg 1$ . Later on, we need the following properties concerning the function w.

Lemma 2. [12, Property 2.16, p.95]

Let T > 0,  $\eta > \alpha - 1$ , and  $0 < \alpha < 1$ . For all  $t \in [0, T]$ , we have

$$^{c}D_{t|T}^{\alpha}w(t) = \frac{\Gamma(\eta+1)}{\Gamma(\eta+1-\alpha)}T^{-\alpha}(1-t/T)^{\eta-\alpha}.$$
(6)

**Lemma 3.** Let T > 0,  $0 < \alpha < 1$ ,  $\eta > \alpha p/(p-1) - 1$ , and p > 1. Then, we have

$$\int_0^T (w(t))^{-\frac{1}{p-1}} |^c D_{t|T}^{\alpha} w(t)|^{\frac{p}{p-1}} dt = C_1 T^{1-\alpha \frac{p}{p-1}}, \tag{7}$$

and

$$\int_{0}^{T} {}^{c}D_{t|T}^{\alpha}w(t) dt = C_{2}T^{1-\alpha}, \tag{8}$$

where

$$C_1 = \frac{1}{\eta + 1 - \alpha \frac{p}{p-1}} \left[ \frac{\Gamma(\eta + 1)}{\Gamma(\eta + 1 - \alpha)} \right]^{\frac{p}{p-1}}, \quad and \quad C_2 = \frac{\Gamma(\eta + 1)}{\Gamma(\eta + 2 - \alpha)}.$$

Proof. Let we start by proving (7). Using Lemma 2, we have

$$\begin{split} \int_0^T (w(t))^{-\frac{1}{p-1}}|^c D_{t|T}^{\alpha} w(t)|^{\frac{p}{p-1}} \, dt &= \left[\frac{\Gamma(\eta+1)}{\Gamma(\eta+1-\alpha)}\right]^{\frac{p}{p-1}} \, T^{-\alpha \frac{p}{p-1}} \int_0^T (w(t))^{-\frac{1}{p-1}} (w(t))^{\frac{p(\eta-\alpha)}{(p-1)\eta}} \, dt \\ &= \left[\frac{\Gamma(\eta+1)}{\Gamma(\eta+1-\alpha)}\right]^{\frac{p}{p-1}} \, T^{-\alpha \frac{p}{p-1}} \int_0^T (1-t/T)^{\eta-\alpha \frac{p}{p-1}} \, dt \\ &= \left[\frac{\Gamma(\eta+1)}{\Gamma(\eta+1-\alpha)}\right]^{\frac{p}{p-1}} \, T^{1-\alpha \frac{p}{p-1}} \int_0^1 (1-s)^{\eta-\alpha \frac{p}{p-1}} \, ds \\ &= C_1 \, T^{1-\alpha \frac{p}{p-1}}. \end{split}$$

Similarly, we get (8).

**Lemma 4.** Let T > 0,  $0 < \alpha < 1$ , p > 1,  $A, B \ge 0$ , and  $v \in C^1([0,T),\mathbb{R})$  satisfying the following fractional differential inequality

$$^{c}D_{0|t}^{\alpha}v(t) \ge B[|v(t)|^{p} - A], \quad t \in [0, T).$$
 (9)

subject to  $v(0) > A^{\frac{1}{p}}$ . Then  $v(t) \ge A^{\frac{1}{p}}$  for all  $t \in [0, T)$ .

Proof. Fixing  $T_1 \in (0,T)$ , we show that  $v(t) \geq A^{\frac{1}{p}}$  for any  $t \in (0,T_1]$ . Then, since  $T_1$  is arbitrary, the claim follows. Let us start by defining  $T^* = \inf\{t > 0; \ v(t) \geq A^{\frac{1}{p}}\}$ . Since v is continuous and  $v(0) > A^{\frac{1}{p}}$ , we have  $T^* > 0$ . We claim  $T^* = T_1$ . Otherwise, we have  $v(t) > A^{\frac{1}{p}}$  for all  $t \in (0,T^*)$  such that  $v(T^*) = A^{\frac{1}{p}}$ ; this implies, in particular, that

$$F(t, v(t)) := B[|v(t)|^p - A] \ge 0, \quad \text{for all } t \in [0, T^*].$$
(10)

On the other hand, since the right hand side of (9) is continuous on  $[0, T_1]$  and  $v \in C^1([0, T_1])$ , applying the Riemann-Liouville fractional integral  $I_{0|T^*}^{\alpha}$  to (9) on  $[0, T_1]$  and using (4), we get

$$A^{\frac{1}{p}} = v(T^*) = v(0) + \frac{1}{\Gamma(\alpha)} \int_0^{T^*} (T^* - s)^{-(1-\alpha)} F(s, v(s)) \, ds \ge v(0) > A^{\frac{1}{p}},$$

where we have used (10); contradiction. This completes the proof.

Using [18, Proposition 4.6] and applying the same argument as in the proof of Lemma 4, one can define the function  $g \in C([0, T_b), \mathbb{R}^+)$  which is the unique solution of

$$\begin{cases}
{}^{c}D_{0|t}^{\alpha}g(t) = B g^{p}(t), & t \in [0, T_{b}), \\
g(0) > 0,
\end{cases}$$
(11)

where  $T_b$  is the maximal time of existence.

#### **Proposition 1** (Fractional differential inequalities).

Let  $T_b > 0$  be the blow-time of the solution of (11), and let  $T > T_b$ ,  $0 < \alpha < 1$ , p > 1, B > 0, and  $f \in C^1([0,T),\mathbb{R})$  be a nonnegative solution of the following fractional differential inequality

$$\begin{cases} {}^{c}D_{0|t}^{\alpha}f(t) \ge B f^{p}(t), & t \in [0,T), \\ f(0) > 0. \end{cases}$$
 (12)

Then f blows up at  $T_b$ , i.e.  $\lim_{t\to T_b^-} f(t) = +\infty$ . Moreover, the following upper and lower bound of  $T_b$  are also given

$$T_L \le T_b \le T_U,\tag{13}$$

where

$$T_U := \left(\frac{\Gamma(1+\alpha)}{B\left(f(0)\right)^{p-1}H(p,\alpha)}\right)^{1/\alpha} \quad and \quad T_L := \left(\frac{\Gamma(1+\alpha)}{B\left(f(0)\right)^{p-1}G(p)}\right)^{1/\alpha},$$

with

$$G(p) = \min\left(2^{p}, \frac{p^{p}}{(p-1)^{p-1}}\right), \qquad H(p,\alpha) = \max(p-1, 2^{-\frac{p\alpha}{p-1}}). \tag{14}$$

*Proof.* Applying [3, Theorem 5.1], we conclude that the solution g of (11) is an increasing function and

$$\lim_{t \to T_h^-} g(t) = +\infty.$$

On the other hand, by taking g(0) = f(0), applying [18, Theorem 4.10] and using (11), (12), we conclude that

$$f(t) \ge g(t) \ge 0$$
,

this implies that

$$\lim_{t \to T_h^-} f(t) = +\infty.$$

Moreover, using [3, Theorem 5.2], we get (13)

**Definition 6.** [13, 23] Let  $s \in (0,1)$  and X be a suitable set of functions defined on  $\mathbb{R}^N$ . The fractional Laplacian  $(-\Delta)^s$  in  $\mathbb{R}^N$  is a non-local operator defined as the following singular integral

$$(-\Delta)^s: v \in X \mapsto (-\Delta)^s v(x) := C_{N,s} \ p.v. \int_{\mathbb{R}^N} \frac{v(x) - v(y)}{|x - y|^{N+2s}} dy,$$

as long as the right-hand side exists, and p.v. stands for Cauchy's principal value,  $C_{N,s}:=\frac{4^s\Gamma(\frac{N}{2}+s)}{\pi^{\frac{N}{2}}\Gamma(-s)}$  is a normalization constant and  $\Gamma$  denotes the Gamma function.

**Lemma 5.** [1, Lemma 2.3] Let  $\langle x \rangle := (1 + |x|^2)^{1/2}$  for all  $x \in \mathbb{R}^N$ . Let  $s \in (0,1)$  and  $\phi : \mathbb{R}^N \to \mathbb{R}$  be a function defined by  $\phi(x) = \langle x \rangle^{-q}$ , where  $n < q \le N + 2s$ . Then,  $\phi \in H^{2s}(\mathbb{R}^N)$  and the following estimate holds:

$$|(-\Delta)^s \phi(x)| \le C_{N,q} \phi(x), \quad \text{for all } x \in \mathbb{R}^N, \qquad C_{N,q} = C(s, N, q) > 0.$$
(15)

**Lemma 6.** [1, Lemma 2.4] Let  $s \in (0,1)$ , and let Let  $\psi$  be a smooth function satisfying  $\partial_x^2 \psi \in L^{\infty}(\mathbb{R}^N)$ . For any R > 0, let  $\psi_R$  be a function defined by

$$\psi_R(x) := \psi(x/R) \quad \text{ for all } x \in \mathbb{R}^N.$$

Then,  $(-\Delta)^s \psi_R$  satisfies the following scaling properties:

$$(-\Delta)^s \psi_R(x) = R^{-2s}((-\Delta)^s \psi)(x/R), \quad \text{for all } x \in \mathbb{R}^N.$$

**Lemma 7.** Let  $s \in (0,1)$ , R > 0 and p > 1. Then, the following estimate holds

$$\int_{\mathbb{R}^{N}} (\phi_{R}(x))^{-\frac{1}{p-1}} \left| (-\Delta)^{s} \phi_{R}(x) \right|^{\frac{p}{p-1}} dx \le C_{3} R^{-\frac{2sp}{p-1} + N},$$

where  $C_3 = (C_{N,q})^{p/(p-1)} A_0 > 0$ ,  $A_0$  is defined below,  $\phi_R(x) := \phi(x/R)$ , and  $\phi$  is given in Lemma 5.

*Proof.* If 0 < s < 1, then using the change of variable  $\tilde{x} = x/R$  and Lemma 6 we have  $(-\Delta)^s \phi_R(x) = R^{-2s}(-\Delta)^s \phi(\tilde{x})$ . Therefore, by Lemma 5 we conclude that

$$\int_{\mathbb{R}^N} (\phi_R(x))^{-\frac{1}{p-1}} \left| (-\Delta)^s \phi_R(x) \right|^{\frac{p}{p-1}} dx \le (C_{N,q})^{\frac{p}{p-1}} R^{-\frac{2sp}{p-1}+N} \int_{\mathbb{R}^N} \phi(\tilde{x}) d\tilde{x} = (C_{N,q})^{\frac{p}{p-1}} A_0 R^{-\frac{2sp}{p-1}+N},$$

where

$$A_0 = \int_{\mathbb{R}^N} \phi(x) \, dx > 0.$$

# 3 Theorem 1. Non-existence of global $L^1$ -weak solution in the case $p \leq p_F$

To state our first result, we set

$$\lambda = \lambda_1 + i\lambda_2, \quad u_0 = g + ih,$$

where  $\lambda_i \in \mathbb{R}$  (i = 0, 1) and g and h are real-valued functions; the real and imaginary parts of  $i^{\alpha}u_0$  can be written, respectively, as

$$G_1(x) = \cos(\frac{\alpha\pi}{2})g(x) - \sin(\frac{\alpha\pi}{2})h(x)$$
, and  $G_2(x) = \cos(\frac{\alpha\pi}{2})h(x) + \sin(\frac{\alpha\pi}{2})g(x)$ .

**Theorem 1** (Non-existence of global  $L^1$ -weak solution in the case  $p \leq p_F$ ). Let  $0 < \alpha < 1, \ 0 < \beta < 2, \ N \geq 1, \ \varepsilon = 1$ .

1. If  $1 , and <math>u_0 \in L^1(\mathbb{R}^N)$  satisfies

$$\lambda_1 \int_{\mathbb{R}^N} G_1(x) \, dx > 0 \quad or \quad \lambda_2 \int_{\mathbb{R}^N} G_2(x) \, dx > 0, \tag{16}$$

then problem (1) admits no global  $L^1$ -weak solution.

2. If  $p = p_F$ , and  $u_0 \in L^2(\mathbb{R}^N)$  satisfies

$$|\lambda_1|^{\frac{2-p}{p-1}}\lambda_1\int_{\mathbb{R}^N}G_1(x)\,dx > C_0\,A_0 \quad or \quad |\lambda_1|^{\frac{2-p}{p-1}}\lambda_2\int_{\mathbb{R}^N}G_2(x)\,dx > C_0\,A_0,$$

where  $A_0 = \int_{\mathbb{R}^N} \langle x \rangle^{-N-\beta} dx$ , and  $C_0$  is defined in (25) below, then problem (1) admits no global  $L^1$ -weak solution.

*Proof.* We argue by contradiction. Suppose that u is a global weak solution to (1), then

$$\lambda \int_{Q_T} |u|^p \varphi(t, x) dt dx + i^{\alpha} \int_{Q_T} u_0(x)^c D_{t|T}^{\alpha} \varphi(t, x) dt dx$$

$$= i^{\alpha} \int_{Q_T} u^c D_{t|T}^{\alpha} \varphi dt dx - \int_{Q_T} u(-\Delta)^{\beta/2} \varphi(t, x) dt dx, \tag{17}$$

for all T > 0 and all  $\varphi \in Y_T$ . In order to get a non-negativity in the left hand side of (17), we consider four cases:

Case I: If  $\lambda_1 > 0$ , then  $\int_{\mathbb{R}^N} G_1 dx > 0$ , therefore by taking the real part (Re) of the both sides of (17), we get:

$$\lambda_1 \int_{Q_T} |u|^p \varphi(t, x) dt dx + \int_{Q_T} G_1(x)^c D_{t|T}^{\alpha} \varphi(t, x) dt dx$$

$$= \int_{Q_T} \operatorname{Re}(i^{\alpha} u)^c D_{t|T}^{\alpha} \varphi dt dx - \int_{Q_T} \operatorname{Re}(u) (-\Delta)^{\beta/2} \varphi(t, x) dt dx.$$

Case II: If  $\lambda_1 < 0$ , then  $\int_{\mathbb{R}^N} G_1 dx < 0$  therefore by taking (-Re) of the both sides of (17) we get:

$$(-\lambda_1) \int_{Q_T} |u|^p \varphi(t,x) dt dx - \int_{Q_T} G_1(x)^c D_{t|T}^{\alpha} \varphi(t,x) dt dx$$
$$= - \int_{Q_T} \operatorname{Re}(i^{\alpha} u)^c D_{t|T}^{\alpha} \varphi dt dx + \int_{Q_T} \operatorname{Re}(u) (-\Delta)^{\beta/2} \varphi(t,x) dt dx.$$

Case III: If  $\lambda_2 > 0$ , then  $\int_{\mathbb{R}^N} G_2 dx > 0$ , therefore by taking the imaginary part (Im) of the both sides of (17), we get:

$$\lambda_2 \int_{Q_T} |u|^p \varphi(t, x) dt dx + \int_{Q_T} G_2(x)^c D_{t|T}^{\alpha} \varphi(t, x) dt dx$$

$$= \int_{Q_T} \operatorname{Im}(i^{\alpha} u)^c D_{t|T}^{\alpha} \varphi dt dx - \int_{Q_T} \operatorname{Im}(u) (-\Delta)^{\beta/2} \varphi(t, x) dt dx.$$

Case IV: If  $\lambda_2 < 0$ , then  $\int_{\mathbb{R}^N} G_2 dx < 0$ , therefore by taking (-Im) of the both sides of (17), we get:

$$(-\lambda_2) \int_{Q_T} |u|^p \varphi(t,x) dt dx - \int_{Q_T} G_2(x)^c D_{t|T}^{\alpha} \varphi(t,x) dt dx$$
  
= 
$$- \int_{Q_T} \operatorname{Im}(i^{\alpha} u)^c D_{t|T}^{\alpha} \varphi dt dx + \int_{Q_T} \operatorname{Im}(u) (-\Delta)^{\beta/2} \varphi(t,x) dt dx.$$

Then we only consider the Case I, since the other cases can be treated in the same way, by assuming  $\lambda_1 > 0$ ,  $u_0 \in L^1(\mathbb{R}^N)$  and

$$\int_{\mathbb{R}^N} G_1(x) \, dx > 0. \tag{18}$$

Thus we have

$$\lambda_{1} \int_{Q_{T}} |u|^{p} \varphi(t,x) dt dx + \int_{Q_{T}} G_{1}(x)^{c} D_{t|T}^{\alpha} \varphi(t,x) dt dx 
\leq \int_{Q_{T}} \left| \cos(\frac{\alpha \pi}{2}) \operatorname{Re} u - \sin(\frac{\alpha \pi}{2}) \operatorname{Im} u \right| \left| {}^{c} D_{t|T}^{\alpha} \varphi(t,x) \right| dt dx + \int_{Q_{T}} |\operatorname{Re}(u)| \left| (-\Delta)^{\beta/2} \varphi(t,x) \right| dt dx 
\leq 2 \int_{Q_{T}} |u| \left| {}^{c} D_{t|T}^{\alpha} \varphi(t,x) \right| dt dx + \int_{Q_{T}} |u| \left| (-\Delta)^{\beta/2} \varphi(t,x) \right| dt dx,$$
(19)

all  $\varphi \in Y_T$ . Using the  $\varepsilon$ -Young inequality

$$ab \le \varepsilon a^p + C_\varepsilon b^{\frac{p}{p-1}}, \quad \text{for all } \varepsilon > 0, \ a, b \ge 0, \qquad C_\varepsilon = \frac{(p-1)(p\varepsilon)^{-\frac{1}{p-1}}}{p},$$
 (20)

we get

$$2\int_{Q_{T}} |u| \left| {}^{c}D_{t|T}^{\alpha} \varphi(t,x) \right| dt dx$$

$$= \int_{Q_{T}} |u| \varphi^{1/p} \varphi^{-1/p} 2 \left| {}^{c}D_{t|T}^{\alpha} \varphi(t,x) \right| dt dx$$

$$\leq \varepsilon \int_{Q_{T}} |u|^{p} \varphi(t,x) dt dx + C_{4} \int_{Q_{T}} \varphi^{-\frac{1}{p-1}} \left| {}^{c}D_{t|T}^{\alpha} \varphi(t,x) \right|^{\frac{p}{p-1}} dt dx, \tag{21}$$

where

$$C_4 = 2^{\frac{p}{p-1}} C_{\varepsilon} = \frac{2^{\frac{p}{p-1}} (p-1)(p\varepsilon)^{-\frac{1}{p-1}}}{p}.$$

Similarly,

$$\int_{Q_T} |u| \left| (-\Delta)^{\beta/2} \varphi(t, x) \right| dt dx$$

$$\leq \varepsilon \int_{Q_T} |u|^p \varphi(t, x) dt dx + C_5 \int_{Q_T} \varphi^{-\frac{1}{p-1}} \left| (-\Delta)^{\beta/2} \varphi(t, x) \right|^{\frac{p}{p-1}} dt dx, \tag{22}$$

where

$$C_5 = C_{\varepsilon} = \frac{(p-1)(p\varepsilon)^{-\frac{1}{p-1}}}{p}.$$

Combining (21)-(22) with (19), we obtain

$$(\lambda_{1} - 2\varepsilon) \int_{Q_{T}} |u|^{p} \varphi(t, x) dt dx + \int_{Q_{T}} G_{1}(x)^{c} D_{t|T}^{\alpha} \varphi(t, x) dt dx$$

$$\leq C_{4} \int_{Q_{T}} \varphi^{-\frac{1}{p-1}} \left| {}^{c} D_{t|T}^{\alpha} \varphi(t, x) \right|^{\frac{p}{p-1}} dt dx + C_{5} \int_{Q_{T}} \varphi^{-\frac{1}{p-1}} \left| (-\Delta)^{\beta/2} \varphi(t, x) \right|^{\frac{p}{p-1}} dt dx$$

which implies, by taking  $\varepsilon \leq \lambda_1/2$ , that

$$\int_{Q_{T}} G_{1}(x)^{c} D_{t|T}^{\alpha} \varphi(t, x) dt dx$$

$$\leq C_{4} \int_{Q_{T}} \varphi^{-\frac{1}{p-1}} \left| {}^{c} D_{t|T}^{\alpha} \varphi(t, x) \right|^{\frac{p}{p-1}} dt dx + C_{5} \int_{Q_{T}} \varphi^{-\frac{1}{p-1}} \left| (-\Delta)^{\beta/2} \varphi(t, x) \right|^{\frac{p}{p-1}} dt dx, \qquad (23)$$

all  $\varphi \in X_T$ . At this stage, we take the test function

$$\varphi(t,x) := \phi_R(x)w(t),$$

with  $\phi_R(x) := \phi(x/R)$ , R > 0, where  $\phi(x)$  and w(t) are defined in Section 2 with  $s = \beta/2$  and  $q = N + \beta$ . Therefore, from (23) we obtain

$$\int_{\mathbb{R}^{N}} G_{1}(x)\phi_{R}(x) dx \int_{0}^{T} {}^{c}D_{t|T}^{\alpha}w(t) dt 
\leq C_{4} \int_{\mathbb{R}^{N}} \phi_{R}(x) dx \int_{0}^{T} (w(t))^{-\frac{1}{p-1}} \left| {}^{c}D_{t|T}^{\alpha}w(t) \right|^{\frac{p}{p-1}} dt 
+ C_{5} \int_{0}^{T} w(t) dt \int_{\mathbb{R}^{N}} (\phi_{R}(x))^{-\frac{1}{p-1}} \left| (-\Delta)^{\beta/2} \phi_{R}(x) \right|^{\frac{p}{p-1}} dx.$$

As

$$\int_{\mathbb{R}^N} \phi_R(x) dx = \int_{\mathbb{R}^N} \phi(\tilde{x}) R^N d\tilde{x} = A_0 R^N, \quad \text{and} \quad \int_0^T w(t) dt = \frac{T}{\eta + 1},$$

so, using Lemma 3 and Lemma 7 with  $s=\beta/2$  and  $\eta>\alpha p/(p-1)-1$ , we obtain

$$C_2 T^{1-\alpha} \int_{\mathbb{R}^N} G_1(x) \phi_R(x) \, dx \le C_6 R^N T^{1-\alpha \frac{p}{p-1}} + C_7 T R^{-\frac{\beta p}{p-1} + N},$$

where

$$C_6 = C_1 C_4 A_0$$
, and  $C_7 = \frac{C_3 C_5}{\eta + 1}$ .

Choosing  $R = T^{\alpha/\beta}$ , we get

$$\int_{\mathbb{D}^{N}} G_{1}(x)\phi(x/T^{\alpha/\beta}) dx \le C_{8} T^{\alpha[\frac{N}{\beta} - \frac{1}{p-1}]}, \tag{24}$$

where

$$C_8 = \frac{1}{C_2} \max\{C_6, C_7\}.$$

By taking, e.g.,  $\varepsilon = \lambda_1/2$ ,  $C_8$  can be written as

$$C_8 = C_0 A_0 \lambda_1^{-\frac{1}{p-1}},$$

where

$$C_0 = \frac{2^{\frac{1}{p-1}}}{p^{\frac{p}{p-1}}C_2} \max \left\{ C_1 \, 2^{\frac{p}{p-1}}, \frac{(C_{N,N+\beta})^{\frac{p}{p-1}}}{\eta+1} \right\}. \tag{25}$$

If  $p < 1 + \frac{\beta}{N}$ , then  $\frac{N}{\beta} - \frac{1}{p-1} < 0$ . As  $G_1 \in L^1(\mathbb{R}^N)$ , letting  $T \to \infty$  and using the dominated convergence theorem we derive

$$\int_{\mathbb{R}^N} G_1(x) \, dx \le 0;$$

a contraction with (18).

If  $p = 1 + \frac{\beta}{N}$ , using again the same argument, we arrive at

$$\int_{\mathbb{R}^N} G_1(x) \, dx \le C_0 \, A_0 \, \lambda_1^{-\frac{1}{p-1}},$$

which is a contradiction.

**Remark 1.** We note that the regularity of  $u_0$  is not so important in Theorem 1, in fact, we can replace  $u_0 \in L^1(\mathbb{R}^N)$  by  $u_0 \in L^2(\mathbb{R}^N)$  and we get a nonexistence of global  $L^2$ -weak solution. In this case, to ensure the existence of the conditions on  $G_1$  and  $G_2$ , we need also to assume that  $G_1$  or  $G_2$  are in  $L^1(\mathbb{R}^N)$ .

### 4 Theorem 2. Non-existence of global $L^2$ -weak solution in $L^2$ -subcritical case for small data

**Theorem 2** (Non-existence for global  $L^2$ -weak solution in  $L^2$ -subcritical case and for small data). Let  $0 < \alpha < 1, \ 0 < \beta < 2, \ N \ge 1, \ \varepsilon > 0$ . Let  $u_0 \in H^s(\mathbb{R}^N)$ ,  $s \ge 0$ , and u be an  $L^2$ -weak solution on  $[0, T_w(\varepsilon))$ . We assume that  $1 and <math>u_0$  satisfies

$$\lambda_1 G_1(x) \quad or \quad \lambda_2 G_2(x) \ge \begin{cases}
|x|^{-k}, & \text{if } |x| > 1, \\
0, & \text{if } |x| \le 1,
\end{cases}$$
(26)

where  $N/2 < k < \frac{\beta}{p-1}$ . Then, u is not global, i.e.  $T_w(\varepsilon) < \infty$ . More precisely, there exists a constant  $\varepsilon_0 > 0$  such that

$$T_w(\varepsilon) \le \begin{cases} B_0 \, \varepsilon^{-\frac{1}{\alpha \kappa_0}}, & \text{if } \varepsilon \in (0, \varepsilon_0), \\ 1, & \text{if } \varepsilon \in [\varepsilon_0, \infty), \end{cases}$$
 (27)

where  $\kappa_0 = \frac{1}{p-1} - \frac{k}{\beta} > 0$  and

$$B_0 = \left( C_0(k+\beta)\omega_N^{-1} \, 2^{\frac{N+\beta}{2}} A_0 \lambda_1^{\frac{p-2}{p-1}} \right)^{\frac{1}{\alpha \kappa_0}},$$

with  $\omega_N$  stands for the (N-1)-dimensional surface measure of the unit sphere.

*Proof.* Repeating the same calculations as in the proof of Theorem 1, by taking here  $\varepsilon \neq 1$ , and assuming only

$$\lambda_1 > 0$$
 and  $G_1(x) \ge \begin{cases} \lambda_1^{-1} |x|^{-k}, & \text{if } |x| > 1, \\ 0, & \text{if } |x| \le 1, \end{cases}$ 

(the other cases can be treated similarly). From (24), we obtain

$$\varepsilon \int_{\mathbb{R}^N} G_1(x)\phi(x/T^{\alpha/\beta}) dx \le C_0 A_0 \lambda_1^{-\frac{1}{p-1}} T^{\alpha\left[\frac{N}{\beta}+1-\frac{p}{p-1}\right]}, \quad \text{for all } 0 < T < T_w(\varepsilon).$$
 (28)

On the other hand,

$$\varepsilon \int_{\mathbb{R}^{N}} G_{1}(x)\phi(x/T^{\alpha/\beta}) dx = \varepsilon T^{\frac{\alpha N}{\beta}} \int_{\mathbb{R}^{N}} G_{1}(yT^{\alpha/\beta})\phi(y) dy$$

$$\geq \lambda_{1}^{-1}\varepsilon T^{\frac{\alpha(N-k)}{\beta}} \int_{|y|>T^{-\frac{\alpha N}{\beta}}} |y|^{-k}\phi(y) dy$$

$$= \lambda_{1}^{-1}\varepsilon T^{\frac{\alpha(N-k)}{\beta}} K(T),$$

where

$$K(T) := \int_{|y| > T^{-\frac{\alpha N}{\beta}}} |y|^{-k} \phi(y) \, dy.$$

Therefore, from (28), we arrive at

$$\varepsilon K(T) \le C_0 A_0 \lambda_1^{\frac{p-2}{p-1}} T^{\alpha \left[\frac{k}{\beta} - \frac{1}{p-1}\right]}, \quad \text{for all } 0 < T < T_w(\varepsilon).$$
(29)

It remains to estimate from below the last inequality.

First, let  $\varepsilon_0 = B_0^{\alpha k_0}$ , then

$$T_w(\varepsilon) \leq 1$$
,

for all  $\varepsilon \geq \varepsilon_0$ . Indeed, suppose on the contrary that there exists  $\varepsilon \geq \varepsilon_0$  such that  $T_w(\varepsilon) > 1$ . Applying (29) with  $\tau \in (1, T_w(\varepsilon))$ , we obtain

$$\varepsilon K(\tau) < C_0 A_0 \lambda_1^{\frac{p-2}{p-1}} \tau^{\alpha \left[\frac{k}{\beta} - \frac{1}{p-1}\right]}, \quad \text{for all } 1 < \tau < T_m(\varepsilon). \tag{30}$$

Using the fact that

$$|y| \leq (1+|y|^2)^{1/2} \leq \sqrt{2}|y|, \qquad \text{for all } |y| > 1,$$

we have

$$\frac{\omega_N}{(k+\beta)2^{\frac{N+\beta}{2}}} = 2^{-\frac{N+\beta}{2}} \int_{|y|>1} |y|^{-k-N-\beta} \, dy \leq K(1) \leq \int_{|y|>1} |y|^{-k-N-\beta} \, dy = \frac{\omega_N}{(k+\beta)}.$$

Whereupon,

$$K(\tau) \ge K(1) \ge \frac{\omega_N}{(k+\beta)2^{\frac{N+\beta}{2}}}, \quad \text{for all } 1 < \tau < T_w(\varepsilon).$$
 (31)

Combining (30) and (31), we obtain

$$\varepsilon \leq (k+\beta)\omega_N^{-1} 2^{\frac{N+\beta}{2}} C_0 A_0 \lambda_1^{\frac{p-2}{p-1}} \tau^{\alpha[\frac{k}{\beta} - \frac{1}{p-1}]},$$

i.e.

$$\tau \le B_0 \, \varepsilon^{-\frac{1}{\alpha \kappa_0}}, \quad \text{for all } 1 < \tau < T_w(\varepsilon).$$

Letting  $\tau \to T_w(\varepsilon)$ , we get

$$T_w(\varepsilon) \le B_0 \, \varepsilon^{-\frac{1}{\alpha \kappa_0}} \le B_0 \, \varepsilon_0^{-\frac{1}{\alpha \kappa_0}} = 1;$$

contradiction. Therefore,  $T_w(\varepsilon) \leq 1$ , for all  $\varepsilon \geq \varepsilon_0$ .

On the other hand, suppose  $\varepsilon < \varepsilon_0$ . If  $T_w(\varepsilon) \leq 1$ , it follows that

$$T_w(\varepsilon) \le 1 = B_0 \, \varepsilon_0^{-\frac{1}{\alpha \kappa_0}} \le B_0 \, \varepsilon^{-\frac{1}{\alpha \kappa_0}}.$$

Hence, it is sufficient to consider  $T_w(\varepsilon) > 1$ . By the above argument, we get again

$$T_w(\varepsilon) \le B_0 \, \varepsilon^{-\frac{1}{\alpha \kappa_0}}$$

This completes the proof.

**Remark 2.** We note that the condition  $k > \frac{N}{2}$  in Theorem 2 is necessary to ensure the existence of at least an  $H^s$ -function  $u_0$  satisfying (32), for all  $s \ge 0$ .

### 5 Theorem 3. Non-existence of global $L^2$ -weak solution for large data

**Theorem 3** (Non-existence of global  $L^2$ -weak solution for p > 1 and large data). Let  $0 < \alpha < 1$ ,  $0 < \beta < 2$ ,  $N \ge 1$ ,  $\varepsilon > 0$ , and p > 1. Let  $u_0 \in H^s(\mathbb{R}^N)$ ,  $s \ge 0$ , and u be an  $L^2$ -weak solution on  $[0, T_w(\varepsilon))$ . We assume that  $u_0$  satisfies

$$\lambda_1 G_1(x) \quad or \quad \lambda_2 G_2(x) \ge \begin{cases} |x|^{-k}, & \text{if } |x| \le 1, \\ 0, & \text{if } |x| > 1, \end{cases}$$

$$(32)$$

where  $k < \min\{\frac{N}{2} - s, \frac{\beta}{p-1}\}$ . Then, there exists a constant  $\varepsilon_1 > 0$  such that for any  $\varepsilon > \varepsilon_1$ , u is not global, i.e.  $T_w(\varepsilon) < \infty$ . More precisely,

$$T_w(\varepsilon) \leq \overline{C} \, \varepsilon^{-\frac{1}{\alpha \kappa_0}},$$

for all  $\varepsilon > \varepsilon_1$ , where  $\kappa_0 = \frac{1}{p-1} - \frac{k}{\beta} > 0$  and

$$\overline{C} = \left( C_0(N-k)\omega_N^{-1} \, 2^{\frac{N+\beta}{2}} A_0 \lambda_1^{\frac{p-2}{p-1}} \right)^{\frac{1}{\alpha \kappa_0}}.$$

*Proof.* Repeating the same calculations as in the proof of Theorem 1, by taking here  $\varepsilon \neq 1$ , and considering only the case

$$\lambda_1 > 0$$
 and  $G_1(x) \ge \begin{cases} \lambda_1^{-1} |x|^{-k}, & \text{if } |x| \le 1, \\ 0, & \text{if } |x| > 1, \end{cases}$ 

as the other cases can be treated similarly. From (24), we obtain

$$\varepsilon \int_{\mathbb{R}^N} G_1(x)\phi(x/T^{\alpha/\beta}) dx \le C_0 A_0 \lambda_1^{-\frac{1}{p-1}} T^{\alpha[\frac{N}{\beta}+1-\frac{p}{p-1}]}, \quad \text{for all } 0 < T < T_w(\varepsilon).$$
 (33)

On the other hand,

$$\varepsilon \int_{\mathbb{R}^{N}} G_{1}(x)\phi(x/T^{\alpha/\beta}) dx = \varepsilon T^{\frac{\alpha N}{\beta}} \int_{\mathbb{R}^{N}} G_{1}(yT^{\alpha/\beta})\phi(y) dy$$

$$\geq \lambda_{1}^{-1}\varepsilon T^{\frac{\alpha(N-k)}{\beta}} \int_{|y| \leq T^{-\frac{\alpha N}{\beta}}} |y|^{-k}\phi(y) dy$$

$$= \lambda_{1}^{-1}\varepsilon T^{\frac{\alpha(N-k)}{\beta}} L(T),$$

where

$$L(T):=\int_{|y|< T^{-\frac{\alpha N}{\beta}}}|y|^{-k}\phi(y)\,dy.$$

Therefore, from (33), we arrive at

$$\varepsilon L(T) \le C_0 A_0 \lambda_1^{\frac{p-2}{p-1}} T^{\alpha[\frac{k}{\beta} - \frac{1}{p-1}]}, \text{ for all } 0 < T < T_w(\varepsilon).$$
 (34)

It remains to estimate from below the last inequality.

We claim that there exists a constant  $\varepsilon_1 > 0$  such that for any  $\varepsilon > \varepsilon_1$ ,

$$T_w(\varepsilon) \le 1.$$
 (35)

Indeed, suppose on the contrary that for all  $\varepsilon_1 > 0$ , there exists  $\varepsilon > \varepsilon_1$  such that  $T_w(\varepsilon) > 1$ . Applying (34) with T = 1, we have

$$\varepsilon L(1) \le C_0 A_0 \lambda_1^{\frac{p-2}{p-1}}. \tag{36}$$

Using the fact that k < N, and

$$\frac{1}{2^{\frac{N+\beta}{2}}} \le \phi(y) \le 1, \quad \text{for all } 0 \le |y| \le 1,$$

it is easy to check that

$$\frac{\omega_N}{(N-k)2^{\frac{N+\beta}{2}}} \le L(1) \le \frac{\omega_N}{(N-k)}.$$
(37)

Combining (36) and (37), we obtain

$$\varepsilon \le (N-k)\omega_N^{-1} \, 2^{\frac{N+\beta}{2}} \, C_0 \, A_0 \, \lambda_1^{\frac{p-2}{p-1}} =: \varepsilon_1;$$

contradiction. Thus the claim is proved.

Therefore, for all  $T < T_w(\varepsilon) \le 1$ , we have

$$L(T) \ge \int_{|y| \le 1} |y|^{-k} \phi(y) \, dy = L(1) \ge \frac{\omega_N}{(N-k)2^{\frac{N+\beta}{2}}},$$

which implies, using again (34),

$$\varepsilon \, \frac{\omega_N}{(N-k)2^{\frac{N+\beta}{2}}} \leq \, C_0 \, A_0 \, \lambda_1^{\frac{p-2}{p-1}} \, T^{\alpha[\frac{k}{\beta} - \frac{1}{p-1}]}, \quad \text{for all } 0 < T < T_w(\varepsilon),$$

i.e.

$$T \leq \overline{C} \, \varepsilon^{-\frac{1}{\alpha \kappa_0}}, \quad \text{for all } 0 < T < T_w(\varepsilon).$$

Since T is arbitrary in  $(0, T_w(\varepsilon))$ , the proof is completed by letting  $T \to T_w(\varepsilon)$ .

Remark 3. In Theorem 3, it is sufficient to just consider the case  $p \le 1 + 2\beta/(N - 2s)$ , because the other case  $p > 1 + 2\beta/(N - 2s)$  is proved below in Section 6, (non local implies non global existence), and in this case we take  $k < \frac{N}{2} - s$  ( $\le \frac{\beta}{p-1}$ ).

**Remark 4.** We note that the condition  $k < \frac{N}{2} - s$  in Theorem 3 is necessary to ensure the existence of at least an  $H^s$ -function  $u_0$  satisfying (32).

## 6 Theorem 4. Nonexistence of local $L^2$ -weak solution in $H^s$ -supercritical case

**Theorem 4** (Non-existence of local  $L^2$ -weak solution in  $H^s$ -supercritical case). Let  $0 < \alpha < 1$ ,  $0 < \beta < 2$ ,  $N \ge 1$ ,  $\varepsilon > 0$ , and  $p > 1 + 2\beta/(N - 2s)$ . Assume  $u_0 \in H^s(\mathbb{R}^N)$ ,  $0 \le s < N/2$ , such that  $u_0$  satisfies (32) with  $\beta/(p-1) < k < N/2 - s$ . Then there is no local  $L^2$ -weak solution of (1).

*Proof.* Suppose that there exists an  $L^2$ -weak solution u on [0,T) for some  $0 < T < T_w(\varepsilon)$ . Repeating the same proof of Theorem 3, we have

$$\varepsilon L(\tau) \le C_0 A_0 \lambda_1^{\frac{p-2}{p-1}} \tau^{\alpha[\frac{k}{\beta} - \frac{1}{p-1}]}, \text{ for all } 0 < \tau < T.$$

For all  $\tau < 1$ , we have

$$L(\tau) \ge \int_{|y| \le 1} |y|^{-k} \phi(y) \, dy = L(1) \ge \frac{\omega_N}{(N-k)2^{\frac{N+\beta}{2}}},$$

whereupon

$$\varepsilon \, \frac{\omega_N}{(N-k)2^{\frac{N+\beta}{2}}} \leq \, C_0 \, A_0 \, \lambda_1^{\frac{p-2}{p-1}} \, \tau^{\alpha[\frac{k}{\beta} - \frac{1}{p-1}]}, \quad \text{for all } 0 < \tau < \min\{1, T\},$$

i.e.

$$\varepsilon \le (N-k)\omega_N^{-1} \, 2^{\frac{N+\beta}{2}} \, C_0 \, A_0 \, \lambda_1^{\frac{p-2}{p-1}} \, \tau^{\alpha[\frac{k}{\beta} - \frac{1}{p-1}]}, \quad \text{for all } 0 < \tau < \min\{1, T\}.$$

As  $\beta/(p-1) < k$ , we have  $k/\beta - 1/(p-1) > 0$ . Therefore, taking  $\tau \to 0^+$ , we obtain  $\varepsilon = 0$ ; contradiction. This completes the proof.

# 7 Theorem 5. Nonexistence of local $L^1$ -weak solution in the case $p > p_F$

**Theorem 5** (Non-existence of local  $L^1$ -weak solution in the supercritical case). Let  $0 < \alpha < 1, \ 0 < \beta < 2, \ N \ge 1, \ \varepsilon > 0$ , and  $p > 1 + \beta/N = p_F$ . Assume  $u_0 \in L^1(\mathbb{R}^N)$  and satisfying (32) with  $\beta/(p-1) < k < N$ . Then there is no local  $L^1$ -weak solution of (1). *Proof.* Suppose that there exists an  $L^1$ -weak solution u on [0,T) for some  $0 < T < \overline{T}_w(\varepsilon)$ . Applying the proof of Theorem 4, step by step. The only difference is the condition k < N instead of k < N/2 - s, which is required to ensure that there exists an  $L^1$ -function  $u_0$  satisfying (32).

# 8 Theorem 6. Nonexistence of global $L^2$ -weak solution: New approach

**Theorem 6** (Nonexistence for global  $L^2$ -weak solution: New approach). Let  $0 < \alpha < 1$ ,  $0 < \beta < 2$ ,  $N \ge 1$ , p > 1, T > 0, and

$$X(T) = C([0,T), L^{2}(\mathbb{R}^{N})) \cap C^{1}([0,T), H^{-\frac{\beta}{2}}(\mathbb{R}^{N})) \cap L^{\infty}((0,T), L^{p}(\mathbb{R}^{N})).$$

Assume  $u_0 \in L^2(\mathbb{R}^N)$  and satisfies

$$M_R(0) > C_{N,p,\beta,\gamma} R^{N - \frac{\beta}{p-1}},$$
 (38)

for some R > 0 and  $\gamma \in \mathbb{C}$  satisfying  $Re(\gamma \lambda) > 0$ , where

$$M_R(0) = Re\left(i^{\alpha}\gamma \int_{\mathbb{R}^N} u(0,x)\phi_R(x) dx\right),$$

with  $\phi_R(x) := \phi(x/R)$ , R > 0,  $(\phi(x) \text{ is defined in Section 2 with } q = N + \beta)$ , and

$$C_{N,p,\beta,\lambda,\gamma}^{p} = 2 C_{1/2} (Re(\gamma\lambda))^{-\frac{p}{p-1}} |\gamma|^{\frac{p^2}{p-1}} A_0^p (C_{N,N+\beta})^{\frac{p}{p-1}}.$$

Then there is no distributional solution  $u \in X(T)$ , with  $T > T_b$ , for (1), where (see (13))

$$T_b \sim \left(\frac{R^{N(p-1)}\Gamma(1+\alpha)}{D_{N,p,\beta,\lambda,\gamma} \left[M_R(0) - C_{N,p,\beta,\gamma}R^{N-\frac{\beta}{p-1}}\right]^{p-1}}\right)^{1/\alpha},\tag{39}$$

and

$$D_{N,p,\beta,\lambda,\gamma} = 2^{-1} \operatorname{Re}(\gamma \lambda) |\gamma|^{-p} A_0^{-(p-1)}.$$

*Proof.* Suppose, on the contrary, that there exists a distributional solution  $u \in X(T)$  with  $T > T_b$ . Let

$$M_R(t) = \operatorname{Re}\left(i^{\alpha}\gamma \int_{\mathbb{R}^N} u(t,x)\phi_R(x) dx\right).$$

By Lemmas 5 and 6, we have

$${}^{c}D_{t|T}^{\alpha}M_{R}(t) = \operatorname{Re}\left(\gamma \int_{\mathbb{R}^{N}} i^{\alpha} {}^{c}D_{t|T}^{\alpha}u(t,x)\phi_{R}(x) dx\right)$$

$$= \operatorname{Re}\left(\gamma \lambda\right) \int_{\mathbb{R}^{N}} |u(t,x)|^{p} \phi_{R}(x) dx + R^{-\beta} \operatorname{Re}\left(\gamma \int_{\mathbb{R}^{N}} u(t,x) \left((-\Delta)^{\beta/2} \phi\right) (x/R) dx\right)$$

$$\geq \operatorname{Re}\left(\gamma \lambda\right) \int_{\mathbb{R}^{N}} |u(t,x)|^{p} \phi(x/R) dx - C_{N,N+\beta} R^{-\beta} |\gamma| \int_{\mathbb{R}^{N}} |u(t,x)| \phi(x/R) dx. \tag{40}$$

In order to get a differential inequality, we start by estimating the second term in the right hand side of

(40). Using 1/2-Young's inequality (20), we obtain

$$C_{N,N+\beta} R^{-\beta} |\gamma| \int_{\mathbb{R}^{N}} |u(t,x)| \phi(x/R) dx$$

$$= \int_{\mathbb{R}^{N}} |u(t,x)| \left[ \operatorname{Re} (\gamma \lambda) \phi(x/R) \right]^{1/p} C_{N,N+\beta} R^{-\beta} |\gamma| \left[ \operatorname{Re} (\gamma \lambda) \right]^{-1/p} \left[ \phi(x/R) \right]^{(p-1)/p} dx$$

$$\leq \frac{1}{2} \operatorname{Re} (\gamma \lambda) \int_{\mathbb{R}^{N}} |u(t,x)|^{p} \phi(x/R) dx$$

$$+ C_{1/2} |\gamma|^{\frac{p}{p-1}} \left( C_{N,N+\beta} \right)^{\frac{p}{p-1}} R^{-\frac{\beta p}{p-1}} \left( \operatorname{Re} (\gamma \lambda) \right)^{-\frac{1}{p-1}} \int_{\mathbb{R}^{N}} \phi(x/R) dx$$

$$= \frac{1}{2} \operatorname{Re} (\gamma \lambda) \int_{\mathbb{R}^{N}} |u(t,x)|^{p} \phi(x/R) dx$$

$$+ C_{1/2} A_{0} |\gamma|^{\frac{p}{p-1}} \left( \operatorname{Re} (\gamma \lambda) \right)^{-\frac{1}{p-1}} \left( C_{N,N+\beta} \right)^{\frac{p}{p-1}} R^{N-\frac{\beta p}{p-1}}, \tag{41}$$

where

$$C_{1/2} = (p-1)p^{-\frac{p}{p-1}}2^{\frac{1}{p-1}}$$
 and  $A_0 = \int_{\mathbb{R}^N} \phi(\tilde{x}) d\tilde{x}$ .

On the other hand, by estimating the first term in the right hand side of (40) by using Hölder's inequality, we get

$$|M_{R}(t)| = \left| \operatorname{Re} \left( i^{\alpha} \gamma \int_{\mathbb{R}^{N}} u(t, x) \phi(x/R) \, dx \right) \right|$$

$$\leq |\gamma| \int_{\mathbb{R}^{N}} |u(t, x)| \phi(x/R) \, dx$$

$$= |\gamma| \int_{\mathbb{R}^{N}} |u(t, x)| (\phi(x/R))^{\frac{1}{p}} (\phi(x/R))^{\frac{p-1}{p}} \, dx$$

$$\leq |\gamma| \left( \int_{\mathbb{R}^{N}} |u(t, x)|^{p} \phi(x/R) \, dx \right)^{\frac{1}{p}} \left( \int_{\mathbb{R}^{N}} \phi(x/R) \, dx \right)^{\frac{p-1}{p}}$$

$$= |\gamma| A_{0}^{\frac{p-1}{p}} R^{\frac{N(p-1)}{p}} \left( \int_{\mathbb{R}^{N}} |u(t, x)|^{p} \phi(x/R) \, dx \right)^{\frac{1}{p}},$$

i.e.

$$\int_{\mathbb{R}^N} |u(t,x)|^p \phi(x/R) \, dx \ge |\gamma|^{-p} A_0^{-(p-1)} R^{-N(p-1)} |M_R(t)|^p. \tag{42}$$

Inserting (41)-(42) into (40), we conclude that

$${}^{c}D_{t|T}^{\alpha}M_{R}(t) \geq 2^{-1}\operatorname{Re}(\gamma\lambda)\int_{\mathbb{R}^{N}}|u(t,x)|^{p}\phi(x/R)\,dx$$

$$-C_{1/2}A_{0}|\gamma|^{\frac{p}{p-1}}\left(\operatorname{Re}(\gamma\lambda)\right)^{-\frac{1}{p-1}}\left(C_{N,N+\beta}\right)^{\frac{p}{p-1}}R^{N-\frac{\beta p}{p-1}}$$

$$\geq 2^{-1}\operatorname{Re}(\gamma\lambda)|\gamma|^{-p}A_{0}^{-(p-1)}R^{-N(p-1)}|M_{R}(t)|^{p}$$

$$-C_{1/2}A_{0}|\gamma|^{\frac{p}{p-1}}\left(\operatorname{Re}(\gamma\lambda)\right)^{-\frac{1}{p-1}}\left(C_{N,N+\beta}\right)^{\frac{p}{p-1}}R^{N-\frac{\beta p}{p-1}}$$

$$= 2^{-1}\operatorname{Re}(\gamma\lambda)|\gamma|^{-p}A_{0}^{-(p-1)}R^{-N(p-1)}\left[|M_{R}(t)|^{p}-C_{N,p,\beta,\lambda,\gamma}^{p}R^{p(N-\frac{\beta}{p-1})}\right],$$

i.e.

$${}^{c}D_{t|T}^{\alpha}M_{R}(t) \ge D_{N,p,\beta,\lambda,\gamma}R^{-N(p-1)} \left[ |M_{R}(t)|^{p} - C_{N,p,\beta,\gamma}^{p}R^{p(N-\frac{\beta}{p-1})} \right]. \tag{43}$$

Applying Lemma 4 and using (38), we conclude that

$$M_R(t) \ge C_{N,p,\beta,\gamma} R^{N - \frac{\beta}{p-1}} > 0, \quad \text{for all } t \in [0,T),$$
 (44)

which implies, by using (43) and the following elementary inequality

$$a^p - b^p \ge (a - b)^p$$
, for all  $a > b \ge 0$ ,  $p > 1$ ,

that

$$^{c}D_{t|T}^{\alpha}M_{R}(t) \ge D_{N,p,\beta,\lambda,\gamma}R^{-N(p-1)}\left[M_{R}(t) - C_{N,p,\beta,\gamma}R^{N-\frac{\beta}{p-1}}\right]^{p}.$$
 (45)

Apply Proposition 1 and the fact that  ${}^{c}D_{t|T}^{\alpha}C=0$ , for any constant C>0, we infer that

$$\lim_{t \to T_h} M_R(t) = +\infty.$$

Since

$$M_R(t) \le ||u(t)||_{L^{\infty}((0,T),L^2(\mathbb{R}^N))} ||\phi(\cdot/R)||_{L^2(\mathbb{R}^N)} < \infty, \text{ for all } t \in [0,T),$$

we get a contradiction, and this completes the proof.

**Remark 5.** Note that, from (14), we have  $H(p,\alpha) = \max(p-1,2^{-\frac{p\alpha}{p-1}}) \ge p-1$ ; this implies that  $T_b$  can be chosen as

$$T_b = \left(\frac{R^{N(p-1)}\Gamma(1+\alpha)}{(p-1)D_{N,p,\beta,\lambda,\gamma} \left[M_R(0) - C_{N,p,\beta,\gamma}R^{N-\frac{\beta}{p-1}}\right]^{p-1}}\right)^{1/\alpha},$$

which is the same blow-up time as in the ordinary differential equation when  $\alpha = 1$ .

Corollary 1 (Theorem 1: New approach).

Let  $0 < \alpha < 1$ ,  $0 < \beta < 2$ ,  $N \ge 1$ ,  $\gamma \in \mathbb{C}$ ,  $\varepsilon = 1$ , p > 1. Assume that  $p < 1 + \beta/N$ , and  $u_0 \in L^1(\mathbb{R}^N) \cap L^2(\mathbb{R}^N)$  satisfies

$$Re(\gamma\lambda) > 0$$
 and  $Re\left(i^{\alpha}\gamma\int_{\mathbb{R}^N}u_0(x)\,dx\right) > 0.$  (46)

Then there is no distributional solution  $u \in X(T)$  to (1) for sufficiently large T > 0.

*Proof.* By (46), using the dominated convergence theorem, we conclude that

$$\lim_{R \to \infty} M_R(0) = \operatorname{Re}\left(i^{\alpha} \gamma \int_{\mathbb{R}^N} u_0(x) \, dx\right) > 0.$$

On the other hand, as  $p < 1 + \beta/N$ ,

$$C_{N,p,\beta,\gamma}R^{N-\frac{\beta}{p-1}}\longrightarrow 0, \quad \text{when } R\to\infty.$$

Therefore, there exists  $R_0 > 0$  such that condition (38) is satisfied. Using Theorem 6, the proof is completed.

**Remark 6.** Note that, by taking  $\gamma = \pm 1, \pm i$  in Corollary 1, condition (46) implies (16), which means that (46) is more general that (16). Therefore, in the subcritical case, Theorem 1 can be seen as a particular case of Corollary 1, but with different regularity.

Corollary 2 (Theorem 2: New approach).

Let  $0 < \alpha < 1$ ,  $0 < \beta < 2$ ,  $N \ge 1$ ,  $\varepsilon > 0$ ,  $\gamma \in \mathbb{C}$ , and p > 1. Assume that  $p < 1 + 2\beta/N$ , and  $u_0 \in H^s(\mathbb{R}^N)$ ,  $s \ge 0$ , satisfies

$$Re(\gamma\lambda) > 0$$
 and  $Re(i^{\alpha}\gamma u_0(x)) \ge \begin{cases} |x|^{-k}, & \text{if } |x| > 1, \\ 0, & \text{if } |x| \le 1, \end{cases}$  (47)

where  $N/2 < k < \frac{\beta}{p-1}$ . Then, there exists a constant  $\varepsilon_2 > 0$  such that for all  $\varepsilon \in (0, \varepsilon_2]$ , there is no distributional solution  $u \in X(T)$  to (1) for sufficiently large  $T > T_b$  with  $T_b$  defined in (39). Moreover  $T_b$  can be estimated as follows

$$T_b \le B_1 \,\varepsilon^{-\frac{1}{\alpha \kappa_1}},\tag{48}$$

for all  $\varepsilon \in (0, \varepsilon_2]$ , where  $\kappa_1 = \frac{1}{p-1} - \frac{\min(N,k)}{\beta} > 0$ ,

$$B_1 = (p-1)^{-1/\alpha} D_{N,p,\beta,\lambda,\gamma}^{-1/\alpha} \Gamma(1+\alpha)^{1/\alpha} 2^{\frac{1}{\alpha\kappa_1}} (C_{N,p,\beta,\gamma})^{\frac{\min(N,k)(p-1)}{\alpha\beta\kappa_1}} I_1^{-\frac{1}{\alpha\kappa_1}}$$

and

$$I_1 := \begin{cases} 2^{-N-\beta-1}\omega_N(N-k)^{-1}R^{N-k}, & \text{if } k < N, \\ \\ 2^{-N-\beta}\omega_N \int_1^2 r^{N-1-k} dr, & \text{if } k \ge N. \end{cases}$$

*Proof.* In order to apply Theorem 6, we need to estimate  $M_R(0)$  from below, for some R>0. Let

$$\varepsilon_{2} = \begin{cases} I_{1}^{-1} C_{N,p,\beta,\gamma} 2^{1 - \frac{\beta \kappa_{1}}{N - k}}, & \text{if } k < N, \\ I_{1}^{-1} C_{N,p,\beta,\gamma} 2^{1 - \beta \kappa_{1}}, & \text{if } k \ge N. \end{cases}$$

Let  $\varepsilon \in (0, \varepsilon_2]$ . We choose  $R = R(\varepsilon)$  such that

$$\begin{cases}
R \ge 2^{1/(N-k)}, & \text{if } k < N, \\
R \ge 2, & \text{if } k \ge N.
\end{cases}$$
(49)

Then, as  $R^{N-k} - 1 \ge R^{N-k}/2$ , when k < N, using (47), we have

$$\begin{split} M_R(0) & \geq & \varepsilon \operatorname{Re} \left( \gamma \, i^{\alpha} \int_{\mathbb{R}^N} u_0(x) \phi(x/R) \, dx \right) \\ & \geq & \varepsilon \int_{|x| \geq 1} |x|^{-k} \phi(x/R) \, dx \\ & \geq & \varepsilon \int_{1 \leq |x| \leq R} |x|^{-k} \phi(x/R) \, dx \\ & \geq & \varepsilon 2^{-N-\beta} \int_{1 \leq |x| \leq R} |x|^{-k} \, dx \\ & = & \varepsilon 2^{-N-\beta} \omega_N \int_1^R r^{N-1-k} \, dr \\ & \geq & \varepsilon 2^{-N-\beta} \omega_N \begin{cases} & (N-k)^{-1} (R^{N-k} - 1), & \text{if } k < N, \\ & \int_1^2 r^{N-1-k} \, dr, & \text{if } k \geq N, \end{cases} \\ & \geq & \varepsilon I_1 \, R^{(N-k)+}, \end{split}$$

with  $(N-k)_{+} = \max(N-k,0)$ . Therefore

$$M_{R}(0) - C_{N,p,\beta,\gamma} R^{N - \frac{\beta}{p-1}} \ge R^{(N-k)+} \left( \varepsilon I_{1} - C_{N,p,\beta,\gamma} R^{-\beta\kappa_{1}} \right) = R^{(N-k)+} \left( \frac{\varepsilon I_{1}}{2} \right) > 0, \tag{50}$$

where R is chosen to ensure the last equality, namely

$$R = \left(\frac{2C_{N,p,\beta,\gamma}}{\varepsilon I_1}\right)^{\frac{1}{\beta \kappa_1}}.$$
 (51)

It is clear, by our choice of  $\varepsilon_2$ , that condition (49) is satisfied. Applying Theorem 6, we conclude that there is no solution  $u \in X(T)$  to (1) for all  $T > T_b$ . Moreover, from (13),(39) and the fact that  $H(p, \alpha) \ge p - 1$ , we obtain

$$T_b \le T_U \le (p-1)^{-1/\alpha} \left( \frac{R^{N(p-1)}\Gamma(1+\alpha)}{D_{N,p,\beta,\lambda,\gamma} \left[ M_R(0) - C_{N,p,\beta,\gamma} R^{N-\frac{\beta}{p-1}} \right]^{p-1}} \right)^{1/\alpha}.$$

Then, using (50) and (51), we conclude that

$$T_b \le B_1 \, \varepsilon^{-\frac{1}{\alpha \kappa_1}}.$$

This complete the proof.

**Remark 7.** Note that,  $\kappa_1 > \kappa_0$ , this means that (27) is better than (48). Moreover, by taking  $\gamma = \pm 1, \pm i$  in Corollary 2, condition (47) implies (26), which means that (47) is more general that (26). Therefore, Theorem 2 can be seen as a particular case of Corollary 2, but with different regularity.

Corollary 3 (Theorem 3: New approach).

Let  $0 < \alpha < 1, \ 0 < \beta < 2, \ N \ge 1, \ \varepsilon > 0, \ \gamma \in \mathbb{C}$ , and p > 1. Assume that  $u_0 \in H^s(\mathbb{R}^N)$ ,  $s \ge 0$ , satisfies

$$Re(\gamma\lambda) > 0$$
 and  $Re(i^{\alpha}\gamma u_0(x)) \ge \begin{cases} |x|^{-k}, & \text{if } |x| \le 1, \\ 0, & \text{if } |x| > 1, \end{cases}$  (52)

where  $k < \min\{\frac{N}{2} - s, \frac{\beta}{p-1}\}$ . Then, there exists a constant  $\varepsilon_3 > 0$  such that for any  $\varepsilon \geq \varepsilon_3$ , there is no distributional solution  $u \in X(T)$  to (1) for sufficiently large  $T > T_b$  with  $T_b$  is defined in (39). Moreover  $T_b$  can be estimated as follows

$$T_b \le B_2 \,\varepsilon^{-\frac{1}{\alpha \kappa_0}},\tag{53}$$

for all  $\varepsilon \geq \varepsilon_3$ , where  $\kappa_0 = \frac{1}{p-1} - \frac{k}{\beta} > 0$ ,

$$B_2 = (p-1)^{-1/\alpha} D_{N,p,\beta,\lambda,\gamma}^{-1/\alpha} \Gamma(1+\alpha)^{1/\alpha} 2^{\frac{1}{\alpha\kappa_0}} (C_{N,p,\beta,\gamma})^{\frac{k(p-1)}{\alpha\beta\kappa_0}} I_2^{-\frac{1}{\alpha\kappa_2}},$$

and

$$I_2 := 2^{-N-\beta} \omega_N (N-k)^{-1}.$$

*Proof.* In order to apply Theorem 6, we need to estimate  $M_R(0)$  from below, for some R>0. Let

$$\varepsilon_3 = 2 I_2^{-1} C_{N,p,\beta,\gamma}$$
.

Let  $\varepsilon \geq \varepsilon_3$ . We choose  $R = R(\varepsilon) \leq 1$ . Then, using (52), we have

$$M_{R}(0) \geq \varepsilon \operatorname{Re} \left( \gamma i^{\alpha} \int_{\mathbb{R}^{N}} u_{0}(x) \phi(x/R) dx \right)$$

$$\geq \varepsilon \int_{|x| \leq 1} |x|^{-k} \phi(x/R) dx$$

$$\geq \varepsilon \int_{|x| \leq R} |x|^{-k} \phi(x/R) dx$$

$$\geq \varepsilon 2^{-N-\beta} \int_{|x| \leq R} |x|^{-k} dx$$

$$= \varepsilon 2^{-N-\beta} \omega_{N} \int_{0}^{R} r^{N-1-k} dr$$

$$= \varepsilon 2^{-N-\beta} \omega_{N} (N-k)^{-1} R^{N-k}$$

$$= \varepsilon I_{2} R^{N-k}.$$

Therefore

$$M_R(0) - C_{N,p,\beta,\gamma} R^{N - \frac{\beta}{p-1}} \ge R^{N-k} \left( \varepsilon I_2 - C_{N,p,\beta,\gamma} R^{-\beta\kappa_0} \right) = R^{N-k} \left( \frac{\varepsilon I_2}{2} \right) > 0, \tag{54}$$

where

$$R = \left(\frac{2C_{N,p,\beta,\gamma}}{\varepsilon I_2}\right)^{\frac{1}{\beta \kappa_0}}.$$
 (55)

It is clear, by our choice of  $\varepsilon_3$ , that  $R \leq 1$ . Applying Theorem 6, we conclude that there is no solution  $u \in X(T)$  of (1) for all  $T > T_b$ . Moreover, from (13),(39) and  $H(p, \alpha) \geq p - 1$ , we obtain

$$T_b \le T_U \le (p-1)^{-1/\alpha} \left( \frac{R^{N(p-1)}\Gamma(1+\alpha)}{D_{N,p,\beta,\lambda,\gamma} \left[ M_R(0) - C_{N,p,\beta,\gamma} R^{N-\frac{\beta}{p-1}} \right]^{p-1}} \right)^{1/\alpha}.$$

Then, using (54) and (55), we conclude that

$$T_b \leq B_2 \, \varepsilon^{-\frac{1}{\alpha \kappa_0}}.$$

This complete the proof.

**Remark 8.** Note that, by taking  $\gamma = \pm 1, \pm i$  in Corollary 3, condition (52) implies (32), which means that (52) is more general that (32). Therefore, Theorem 3 can be seen as a particular case of Corollary 3, but with different regularity.

#### References

- [1] T.A. Dao, M. Reissig, A blow-up result for semi-linear structurally damped  $\sigma$ -evolution equations, preprint on arXiv:1909.01181v1, 2019.
- [2] J. Dong, M. Xu, Space-time fractional Schrödinger equation with time-independent potentials, J. Math. Anal. Appl., 344 (2) (2008), 1005-1017.
- [3] Y. Feng, Lei Li, Jian-Guo Liu, X. Xu, Continuous and discrete one dimensional autonomous fractional ODEs, Discrete & Continuous Dynamical Systems B, 23(8) (2018), 3109-3135.
- [4] A. Z. Fino, I. Dannawi, M. Kirane Blow-up of solutions for semilinear fractional Schrödinger equations, J. Integral Equations & Appl. 30(1) (2018), 67-80.
- [5] A. Z. Fino, I. Dannawi, M. Kirane Erratum to "Blow-up of solutions for semilinear fractional Schrödinger equations", J. Integral Equations & Appl. accepted, 2020.
- [6] K. Fujiwara, A note for the global nonexistence of semirelativistic equations with nongauge invariant power type nonlinearity, Math Meth Appl Sci. 41(13) (2018), 1-12.
- [7] K. Fujiwara, T. Ozawa, Remarks on global solutions to the Cauchy problem for semirelativistic equations with power type nonlinearity, Int. J. Math. Anal. 9 (2015), 2599-2610.
- [8] M. Ikeda, Y. Wakasugi Small data blow-up of L<sup>2</sup>-solution for the nonlinear Schrödinger equation without gauge invariance, Diff. Int. Equ. 26 (2013), 1275-1285.
- [9] M. Ikeda, T. Inui Small data blow-up of  $L^2$  or  $H^1$ -solution for the semilinear Schrödinger equation without gauge invariance, J. Evol. Equ. 15(3) (2015), 1-11.
- [10] M. Ikeda, T. Inui Some non-existence results for the semilinear Schrödinger equation without gauge invariance, J. Math. Anal. Appl. 425(2) (2015), 758-773.

- [11] T. Inui Some nonexistence results for a semirelativistic Schrödinger equation with nongauge power type nonlinearity, Proc. Amer. Math. Soc. 144(7) (2016), 2901-2909.
- [12] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, Theory and Applications of Fractional Differential Equations, 2006.
- [13] M. Kwaśnicki, Ten equivalent definitions of the fractional Laplace operator, Fract. Calc. Appl. Anal., 20 (2017), 7-51.
- [14] N. Laskin, Fractional quantum mechanics, Phys Rev E., 62 (3) (2000), 3135-3145.
- [15] N. Laskin, Fractional quantum mechanics and Levy path integrals, Phys Lett A., 268 (4-6) (2000), 298-305.
- [16] N. Laskin, Fractals and quantum mechanics, Chaos., 10 (4) (2000), 780-790.
- [17] Jin Bong Lee, Strichartz estimates for space-time fractional Schrödinger equations, J. Math. Anal. Appl. 487(2) (2020), 123999.
- [18] Lei Li, Jian-Guo Liu, A generalized definition of Caputo derivatives and its application to fractional ODEs, SIAM J. Math. Anal. 50(3) (2018), 2867-2900.
- [19] M. Naber, Time fractional Schrödinger equation, J. Math. Phys., 45 (8) (2004), 3339-3352.
- [20] B. N. Narahari Achar, B. T. Yale, J. W. Hanneken, Time fractional Schrödinger equation revisited, Adv. Math. Phys., 2013 (2013), Article ID 290216, 11 pages.
- [21] S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional integrals and derivatives*, Theory and Applications, Gordon and Breach Science Publishers, 1987.
- [22] R. K. Saxena, R. Saxena, S. L. Kalla, Solution of space-time fractional Schrödinger equation occurring in quantum mechanics, Fract. Calc. Appl. Anal., 13 (2) (2010), 177-190.
- [23] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, Comm. Pure Appl. Math., 60(1) (2007), 67-112.
- [24] Qi S. Zhang, A blow up result for a nonlinear wave equation with damping: the critical case, C. R. Acad. Sci. Paris, 333(2) (2001), 109-114.
- [25] Q.G. Zhang, H.R. Sun, Y.N. Li, The nonexistence of global solutions for a time fractional nonlinear Schrödinger equation without gauge invariance, Applied Mathematics Letters 64 (2016) 119-124.