HF, DF, TF: Approximating potential curves, calculating rovibrational states

Laura E. Angeles Gantes^{*} and Horacio Olivares-Pilón[†]

Departamento de Física, Universidad Autónoma Metropolitana-Iztapalapa, Apartado Postal 55-534, 09340 México, D.F., Mexico

Abstract

An analytical expression for the potential energy curve for the ground state $X^{1}\Sigma^{+}$ of the hydrogen fluoride molecule (HF) obtained in the framework of the Born-Oppenheimer approximation is proposed. This analytical expression for the potential energy curve is based in the two point Padé approximation, which correctly reproduces the asymptotic behavior at small $R \to 0$ and large $R \to \infty$ internuclear distances, and position and depth of potential well, leading to the accuracy of 4-5 s.d. when compared with experimental data. The rovibrational spectra of the diatomic molecule HF is calculated by solving the Schrödinger equation for nuclear motion using the Lagrange-mesh method. The ground state $X^{1}\Sigma^{+}$ contains 21 vibrational states (ν , 0) and 724 rovibrational states (ν , L) with maximal angular momentum equal to 55. The change of reduced mass in the Schrödinger equation for nuclear motion allows us to obtain the rovibrational spectrum of the ground state potential curve of the deuterium fluoride (DF) and tritium fluoride (TF) molecules, it contains 1377 and 1967 rovibrational states (ν , L) with maximal angular momenta 76 and 92, respectively. Entire rovibrational spectra is presented for the HF molecule and its two isotopologues DF and TF.

arXiv:2110.01991v2 [physics.chem-ph] 27 Oct 2021

^{*}Electronic address: lauraE_AG@gmail.com

[†]Electronic address: horop@xanum.uam.mx

I. INTRODUCTION

In the analysis of molecular systems within the Born-Oppenheimer approximation, the parametric dependence of the total energy on the nuclear positions defines the potential energy surface (PES). In practice, the PES is given at a set of discrete points in the space of nuclear coordinates which are obtained by solving the electronic motion in the field produced by the fixed nuclei (charged centers).

In the simplest case of diatomic molecules, the potential energy surface becomes a potential energy curve (PEC) as a function of the internuclear distance R. Nowadays, the development of numerical techniques for quantum-mechanical calculations, has made it possible to obtain highly accurate values of the PEC, however, as is well known, as the number of electrons increases, the accuracy deteriorates while the computational time increases dramatically. An alternative is to extract the PEC from experimental results using the Rydberg-Klein-Rees (RKR) method [1–3], from where the turning points of the vibrational levels are obtained. In order to construct a continuous representation of the PEC, the standard approach consists of fitting the discrete points for PEC by traditional potentials (*e.g.* Morse, Lennard-Jones, Pöschl-Teller, etc.) or to consider more sophisticated expressions for potentials [4] mainly, near equilibrium and sometimes involving the domain of large distances.

In this work we consider the ground state $X^1\Sigma^+$ of the Hydrogen Fluoride (HF), a diatomic molecule composed of a Hydrogen atom (1e, 1p) and a Fluorine atom (9e, 9p). This molecule, with 10 electrons in total, has been extensively studied both experimentally and theoretically (see for example [5–10] and references therein). At present, the high accuracy achieved experimentally has made it possible to obtain very accurate potential energy curves. Theoretically, it has been a challenge to obtain a complete description from *ab-initio* calculations. The average absolute deviation between experimental and theoretical vibrational energy levels is ~ 80 cm⁻¹ (~ 4 × 10⁻⁴ Hartree) [9].

The goal of this paper, is to present an analytical expression for the ground state $X^1\Sigma^+$ of the potential energy curve V(R) in all domain of internuclear distances R with a correct asymptotic behavior at small $(R \to 0)$ and large $(R \to \infty)$ values of the distance between nuclei. The analitical expression V(R) is given in terms of a two points Padé approximant. The methodology followed here, has been applied successfully to some specific diatomic molecules: H_2^+ , H_2 , HeH and LiH [11, 12]. With the explicit analytical expression of the PEC, the Schrödinger equation for nuclear motion (see below) is solved in Lagrange-mesh method to obtain the rovibrational spectrum for this diatomic molecule.

Making a simple modification in the nuclear Schrödinger equation - it occurs due to the fact that we make studies in the Born-Oppenheimer approximation - the rovibrational spectra $E_{(\nu,L)}$ is also obtained for the isotopological species with Deuterium (D) and Tritium (T) content: Deuterium Fluoride (DF) and Tritium Fluoride (TF) molecules, respectively.

II. THE GROUND STATE $X^{1}\Sigma^{+}$ OF THE HF MOLECULE

Let us start by describing the two asymptotic limits, small $(R \to 0)$ and large $(R \to \infty)$ internuclear distances of the potential energy curve for the ground state $X^1\Sigma^+$ of the heteronuclear diatomic molecule HF. In the united atom limit $(R \to 0)$, we have a Neon atom (Ne) and the dissociation energy \tilde{E} for small internuclear distances is given by

$$\widetilde{E} = \frac{9}{R} - \varepsilon_0 + O(R^2), \qquad (1)$$

where the first term is the repulsive Coulomb interaction between nuclei and ε_0 is the difference between the energy of the system in the united atom limit (Neon atom E_{Ne}) and the sum of the energies for each individual components of the molecule, the Hydrogen (E_H) and Fluorine atom (E_F)

$$\varepsilon_0 = E_{\rm Ne} - \left(E_H + E_F\right),\tag{2}$$

with $E_{\rm H} = -0.5$ a.u. and $E_{\rm F} = -99.733424$ a.u. [13]. On the other hand at large internuclear distances $R \to \infty$, the molecule dissociates into a Hydrogen atom plus a Fluorine atom, and the dissociation energy is given by

$$\widetilde{E} = -\frac{C_6}{R^6} + O\left(\frac{1}{R^8}\right),\tag{3}$$

where $C_6 = 7.766$ [7] is the Van der Waals coefficient.

In order to construct an analytic representation for the potential energy curve interpolating the two asymptotic limits $R \to 0$ (1) and $R \to \infty$ (3) a two point Padé type approximant of the form $E(R) = \frac{1}{R}$ Pade [N/N + 5] is used. Choosing N = 5, the explicit expression for E(R) is given by

$$E(R)_{\{3,2\}} = \frac{1}{R} \operatorname{Pade}\left[5/10\right](R), \qquad (4)$$

$$= \frac{9 + a_1 R + a_2 R^2 + a_3 R^3 + a_4 R^4 - C_6 R^5}{R(1 + \alpha_1 R + \alpha_2 R^2 + b_3 R^3 + b_4 R^4 + b_5 R^5 + b_6 R^6 + b_7 R^7 + b_8 R^8 + \alpha_3 R^9 + R^{10})},$$

where the three constrains α_i (i = 1, 2, 3)

$$\begin{aligned}
\alpha_1 &= \frac{a_1 - \varepsilon_0}{9}, \\
\alpha_2 &= \frac{-a_1 \varepsilon_0 + \varepsilon_0^2 + 9a_2}{81}, \\
\alpha_3 &= -\frac{a_4}{C_6},
\end{aligned}$$
(5)

guarantee the expected behavior in the series expansion: the three terms R^{-1} , R^0 and R^1 for small internuclear distances $R \to 0$ (1) and the two terms R^{-6} and R^{-7} for large internuclear distances $R \to \infty$ (3) are reproduced. The then free parameters have been obtained through a fit of the experimental energy values reported in [8] with the analytic expression (5). Their values result in:

$$a_1 = 2761.87,$$
 $a_2 = -2269.79,$ $a_3 = -193.29,$ $a_4 = 107.967$
 $b_3 = -660.63,$ $b_4 = 1909.71,$ $b_5 = -1658.58,$ $b_6 = 988.485,$
 $b_7 = -379.066,$ $b_8 = 94.677.$

As can be seen in Table I, the the analytic expression (5) gives no less that 4 s.d. in correspondence with the experimental results [8]. The position and the deep of the potential energy curve can be calculated by simply taking the derivative of (5) and setting it equal to zero. As a result, we obtain $R_{eq} = 1.73254$ a.u. and $E_{min} = -0.224913$ a.u. In comparison, the best numerical result is $R_{eq} = 1.73257$ a.u. and $E_{min} = -0.224908$ a.u. [8]. The analytic expression for the potential energy curve (5) as well as the experimental results [8], are depicted in Figure 1.

1. Rovibrational spectra

In the Born-Oppenheimer approximation, the rovibrational spectra $E_{(\nu,L)}$ can be obtained by solving the nuclear Schrödinger equation for the Hamiltonian

$$-\frac{1}{2\mu}P^2 + \frac{L(L+1)}{2\mu R^2} + V(R), \qquad (6)$$

R_{\min} [u.a.]	$R_{\sf max}$ [u.a.] —	E [Hartree]		
		Data [<mark>8</mark>]	$E_{\{3,2\}}(R_{\min})$	$E_{\{3,2\}}(R_{\max})$
1.57634	1.92856	-0.21556	-0.21557	-0.21556
1.48248	2.10342	-0.19751	-0.19752	-0.19751
1.42634	2.24290	-0.18025	-0.18025	-0.18025
1.38529	2.36979	-0.16375	-0.16374	-0.16375
1.35286	2.49078	-0.14799	-0.14798	-0.14799
1.32613	2.60905	-0.13295	-0.13295	-0.13296
1.30352	2.72653	-0.11863	-0.11863	-0.11864
1.28405	2.84461	-0.10501	-0.10502	-0.10502
1.26707	2.96450	-0.09210	-0.09210	-0.09210
1.25212	3.08735	-0.07987	-0.07987	-0.07987
1.23890	3.21446	-0.06834	-0.06835	-0.06834
1.22716	3.34736	-0.05752	-0.05752	-0.05752
1.21673	3.48807	-0.04742	-0.04742	-0.04742
1.20748	3.63937	-0.03807	-0.03807	-0.03808
1.19934	3.80533	-0.02951	-0.02951	-0.02952
1.19225	3.99227	-0.02181	-0.02181	-0.02181
1.18618	4.21070	-0.01502	-0.01502	-0.01502
1.18116	4.47992	-0.00928	-0.00927	-0.00926
1.17725	4.84130	-0.00471	-0.00470	-0.00471
1.17457	5.41923	-0.00153	-0.00153	-0.00155
1.17337	7.29868	-0.00010	-0.00010	-0.00005

TABLE I: Experimental energy of the diatomic molecule HF [8] (column 3) for their corresponding turning points R_{\min} and R_{\max} of the potential energy curve. Columns 4 and 5 result from fit $E_{\{3,2\}}$ (5) at R_{\min} and R_{\max} .

where L is the angular momentum, μ is the reduced mass and the momentum $P = -i\nabla_R$. For the concrete case of the Hydrogen Fluoride molecule, the interaction potential V(R) is given by the analytic expression (5) and the reduced mass $\mu = m_H m_F/(m_H + m_F)$ is calculated from the nuclear masses of the Hydrogen and Fluorine atoms: $m_H = 1836.1527$ a.u. and $m_F = 34622.974$ a.u. [14], respectively.

In order to solve the one dimension differential equation for the nuclear motion (6), the

Lagrange-mesh method is used [15]. It is found that the potential energy curve for the ground state $X^1\Sigma^+$ supports 21 pure vibrational states $(L = 0) E_{(\nu,0)}$ with $\nu = 0, \dots, 20$ in complete agreement with the experimental results presented in [8] (see Table II). The absolute error in the vibrational spectra is $\leq 4 \times 10^{-5}$. For comparison, the experimental data presented in [5] are also shown in Table II. Figure 2 shows the number N of vibrational levels with the same rotational quantum number L. It can be noticed that the number of rotational levels $E_{(0,L)}$ of the ground state $X^1\Sigma^+$ is 56 ($L = 0, \dots, 55$). In total the potential energy curve supports 724 rovibrational states. The data from $L = 0, \dots, 41$ reported in [10] are also depicted in Figure 2. The presence of 27 extra states between L = 9 and L = 40 is due to the fact that in [10] the breakdown of the Born-Oppenheimer approximation is taken into account.

III. THE GROUND STATE $X^1\Sigma^+$ OF THE DF MOLECULE

The formalism developed previously for the HF molecule allows us to extend the study to the isotopologues DF (Deuterium Fluorine) and TF (Tritium Fluorine). Let's start by considering the deuterium-containing molecule DF.

Due to the fact that we are working inside of the Born-Oppenheimer approximation, we




FIG. 1: Potential energy curve for the HF molecule. The solid curve is the fit $E(R)_{\{3,2\}}$ (5), and the points are the experimental results [8]. The minimum $E_{min} = -0.224913$ a.u. is located at $R_{eq} = 1.73254$ a.u.

		E [Hartree]		
ν —	Data [5]	Data [8]	$E_{(\nu,0)}$	
0	-0.21565	-0.21556	-0.21556	
1	-0.19760	-0.19751	-0.19751	
2	-0.18033	-0.18025	-0.18024	
3	-0.16383	-0.16375	-0.16373	
4	-0.14807	-0.14799	-0.14797	
5	-0.13304	-0.13295	-0.13293	
6	-0.11872	-0.11863	-0.11861	
7	-0.10510	-0.10501	-0.10499	
8	-0.09218	-0.09210	-0.09207	
9	-0.07995	-0.07987	-0.07984	
10	-0.06842	-0.06834	-0.06831	
11	-0.05760	-0.05752	-0.05748	
12	-0.04750	-0.04742	-0.04739	
13	-0.03815	-0.03807	-0.03804	
14	-0.02960	-0.02951	-0.02949	
15	-0.02189	-0.02181	-0.02178	
16	-0.01511	-0.01502	-0.01500	
17	-0.00936	-0.00928	-0.00925	
18	-0.00479	-0.00471	-0.00469	
19	-0.00161	-0.00153	-0.00152	
20		-0.00010	-0.00007	

TABLE II: Vibrational energies $E_{(\nu,0)}$ of the ground state $X^1\Sigma^+$ of the diatomic molecule HF.

are not aware of isotopic corrections: the depth of the minimum is the same for all three systems, HF, DF and TF. For this reason, in order to obtain the rovibrational spectrum by solving the Schrödinger equation for the nuclear motion (with Hamiltonian (6)), the analytic expression for the interaction potential is the same as that for the HF molecule (5). The difference appears in the nuclear reduced mass $\mu = m_D m_F/(m_D + m_F)$ (6), where the Deuteron mass is $m_D = 3670.4833$ a.u. [14]. The resulting differential equation was solved using the Lagrange-mesh method [15].

Column 4 of Table III, presents the 29 vibrational states $E_{(\nu,0)}$, $\nu = 0, \ldots, 28$ supported

by the ground state $X^1\Sigma^+$ of the DF molecule. When comparing with the results obtained from experimental data [10], it is found that the absolute error is $\leq 6 \times 10^{-5}$. For comparison purposes, column 2 of Table III display the results given in [16].

The number of vibrational states as a function of the angular number L are depicted in Figure 3. There are 77 pure rotational states $E_{(0,L)}$, with $L = 0, \ldots, 76$. In total, the ground state $X^1\Sigma^+$ of the DF molecule supports 1377 rovibrational states. Corrections to the Born-Oppenheimer approximation taken into account in [10] explain the presence of the 87 additional states between L = 14 and L = 60 shown in Fig.3.

IV. THE GROUND STATE $X^1\Sigma^+$ OF THE TF MOLECULE

Let us now consider the diatomic molecule TF which contains a Tritium atom and a Fluorine atom. The rovibrational spectra can be calculated in the frame of the Lagrange-mesh method [15] by solving the Schrödinger equation for the nuclear motion with Hamiltonian (6)




FIG. 2: Number of rovibrational states for the ground state $X^1\Sigma^+$ of the HF molecule as a function of the angular momentum L (marked by blue and red). The continuous red line indicates the maximum number of rovibrational states obtained by solving the nuclear Schrödinger equation. Results from [10] for L = 0, ..., 41, where corrections to the B-O approximation are taken into account, are also presented (marked by blue).

and potential curve (5). For this molecule, $\mu = m_T m_F / (m_T + m_F)$, where the Triton mass is $m_T = 5496.9221$ a.u.

As is shown in Table IV, in agreement with the experimental results [10], the ground state $X^{1}\Sigma^{+}$ of the TF molecule contains 35 pure vibrational states (L = 0), $E_{(\nu,0)}$ ($\nu = 0, ..., 34$). The absolute error between $E_{(\nu,0)}$ and the experimental results is $\leq 8 \times 10^{-5}$. It is found that there are 93 rotational states ($\nu = 0$), $E_{(0,L)}$ (L = 0, ..., 92) and in total the ground state keeps 1967 rovibrational states.

The number N of vibrational levels as a function of the angular momentum L is depicted in histogram of Fig.4 together with those reported in [10].

V. CONCLUSIONS

Applying the methodology developed in [11, 12], an accurate analytical representation of the potential energy curve for the ground stated $X^1\Sigma^+$ of the Hydrogen Fluorine (HF)




FIG. 3: Rovibrational states in the ground state $X^1\Sigma^+$ of the DF molecule as a function of the angular momentum L. The continuous red line indicates the maximum number of rovibrational states at given L obtained by solving the nuclear Schrödinger equation. Results reported in [10] from L = 0, ..., 69, where corrections to the B-O approximation are taken into account, are also presented (marked by blue).

molecule is presented. The approximation is based on a two-point Páde approximant which satisfies the two criteria: (i) it should reproduce correctly the coefficients in the series expansion for small $(R \to 0)$ and large $(R \to \infty)$ internuclear distances and (ii) two characteristics of the potential well, the equilibrium position R_{eq} and the dissociation energy E_{min} , are reproduced accurately. Finally, the obtained analytical curve is accurate with not less than 4 s.d. compared with experimental results whenever is possible. It is shown that inside of the Born-Oppenheimer approximation the potential energy curve keeps 21 vibrational states ($\nu = 0, \ldots, 20, L = 0$), 56 rotational states ($\nu = 0, L = 0..., 55$) and, in total, 722 rovibrational states (ν, L).

Modification of the reduced mass μ in the Schrödinger equation for nuclear motion (6) allows us to explore the isotopologues of the HF molecule. It is found that the ground state $X^{1}\Sigma^{+}$ of the Deuterium Fluorine (DF)/ Tritium Fluorine molecule (TF) supports 29/35 vibrational states $E_{(\nu,0)}$ and 77/93 rotational states $E_{(0,L)}$, respectively. The number of vibrational states $E_{(\nu,0)}$ calculated for the HF, DF and TF molecules is in agreement with those reported in [8, 10]. On the other hand, the maximum value of the angular momentum for rotational states $E_{(0,L)}$ is $L_{max} = 55,76$ and 92, respectively, (note that in [10] the rovibrational states are presented up to L = 41,69 and 15) for HF, DF and TF diatomic




FIG. 4: Number of vibrational states for the ground state $X^{1}\Sigma^{+}$ of the TF molecule as a function of the angular momentum L. Results reported in [10] are also presented (marked by blue).

molecules. In the Born-Oppenheimer approximation, the total number of rovibrational states supported by the ground state potential curve are: 724, 1377 and 1967, respectively. For HF and DF, the difference with the results of [10] in the number of vibrational states for a given angular momentum L (Figures 2 and 3), when a comparison is possible, is due to taken into account the non-adiabatic terms in the Schrödinger equation for the nuclear motion.

Finally, it is worth mentioning that the same methodology can be applied to study the potential energy curves for both, attractive and repulsive excited states of the HF molecule and its isotopologies DF and TF. Note that with the analytical expressions for the PEC, radiative transitions between different electronic states can be calculated. It will be done elsewhere.

ACKNOWLEDGMENTS

The authors thank A. Turbiner for bringing the problem to their attention through the work of O. Polyansky about the HF dimer, the interest to the work and for the numerous valuable discussions. HOP also thanks the support through the Programa Especial de Apoyo a la Investigación 2019, Universidad Autónoma Metropolitana. LAG is grateful to CONACyT (Mexico) for a graduate scholarship.

- R. Rydberg, Graphische darstellung einiger bandenspektroskopischer ergebnisse, Zeitschrift für Physik, 73, 376–385 (1932)
- [2] O. Klein, Zur berechnung von potentialkurven f
 ür zweiatomige molek
 üle mit hilfe von spektraltermen,

Zeitschrift für Physik, 76, 226–235 (1932)

- [3] A. L. G. Rees, The calculation of potential-energy curves from band-spectroscopic data, Proc. Phys. Soc, 59, 998 (1947)
- [4] I. G. Kaplan, Intermolecular Interactions: Physical picture, computational methods and model potentials,

John Wiley & Sons, Ltd (2006)

- [5] G. D. Lonardo and A. E. Douglas, The Electronic Spectrum of HF. I. The B 1Σ+-X 1Σ+ Band System,
 Can. J. Phys., 51, 434-445 (1973)
- [6] J. A. Coxon and P. G. Hajigeorgiou, Isotopic dependence of Born-Oppenheimer breakdown effects in diatomic hydrides: The B¹Σ⁺ and X¹Σ⁺ states of HF and DF, J. Mol. Spectrosc., 142, 254-278 (1990)
- [7] W. T. Zemke, W. C. Stwalley, J. A. Coxon, and P. G. Hajigeorgiou, Improved potential energy curves and dissociation energies for HF, DF and TF, Chem. Phys. Lett. 177, 412-418 (1991)
- [8] J. A. Coxon and P. G. Hajigeorgiou, The B¹Σ⁺ and X¹Σ⁺ electronic states of hydrogen fluoride: A direct potential fit analysis,
 J. Chem. Phys, **110**, 6261-6270 (2006)
- [9] X. Li and J. Paldus, Energy- versus amplitude-corrected coupled-cluster approaches. III. Accurate computation of spectroscopy data exemplified on the HF molecule, J. Chem. Phys, 117, 19411955 (2002)
- [10] J. A. Coxon and P. G. Hajigeorgiou, Improved direct potential fit analyses for the ground electronic states of the hydrogen halides: HF/DF/TF, HCl/DCl/TCl, HBr/DBr/TBr and HI/DI/TI,
 - J. Quant. Spectrosc. Radiat. Transf. 151, 133-154 (2015)
- [11] H. Olivares Pilón, and A.V. Turbiner,
 H₂⁺, HeH and H₂: Approximating potential curves, calculating rovibrational states,
 Ann. of Phys. 393, 335-357 (2018)
- [12] A. V. Turbiner and H. Olivares-Pilón, Towards the analytic theory of potential energy curves for diatomic molecules. Studying He₂⁺ and LiH dimers as illustration, arXiv:1904.06614v2 (2020)
- [13] C. Hattig, W. Klopper, A. Kohn and D. P. Tew, Explicitly correlated electrons in molecules, Chem. Rev. bf 112, 4–74 (2012)
- [14] A. H. Wapstra, G. Audi and C. Thibault, The Ame2003 atomic mass evaluation:(I). Evaluation of input data, adjustment procedures, Nucl. Phys. A 729, 129–336 (2003)
- [15] D. Baye, The Lagrange-mesh method,

Phys. Rep., **32**, 1–107 (2015)

[16] R. J. Fallon, J. T. Vanderslice, and E. A. Mason, Potential Energy Curves of Hydrogen Fluoride,

J. Chem. Phys, 32, 698-700 (1960)

ν —	Data [16]	Data [10]	$E_{(\nu,0)}$
0	-0.21807	-0.21806	-0.21812
1	-0.20483	-0.20482	-0.20487
2	-0.19200	-0.19199	-0.19204
3	-0.17953	-0.17957	-0.17962
4	-0.16744	-0.16755	-0.16760
5	-0.15574	-0.15593	-0.15597
6	-0.14441	-0.14469	-0.14473
7	-0.13346	-0.13383	-0.13387
8	-0.12289	-0.12335	-0.12339
9	-0.11271	-0.11324	-0.11328
10	-0.10290	-0.10349	-0.10353
11	-0.09347	-0.09411	-0.09415
12	-0.08443	-0.08510	-0.08513
13	-0.07576	-0.07645	-0.07648
14	-0.06747	-0.06817	-0.06819
15	-0.05957	-0.06026	-0.06028
16	-0.05255	-0.05272	-0.05275
17	-0.04540	-0.04558	-0.04560
18	-0.03865	-0.03883	-0.03885
19	-0.03231	-0.03249	-0.03251
20	-0.02641	-0.02659	-0.02661
21	-0.02097	-0.02115	-0.02117
22	-0.01603	-0.01620	-0.01621
23	-0.01162	-0.01178	-0.01179
24		-0.00794	-0.00795
25		-0.00475	-0.00475
26		-0.00228	-0.00228
27		-0.00066	-0.00066
28		-0.00006	-0.00002

TABLE III: Vibrational energies $E_{(\nu,0)}$ of the ground state $X^1\Sigma^+$ of the diatomic molecule DF.

	e]	
ν —	Data [<mark>10</mark>]	$E_{(\nu,0)}$
0	-0.21915	-0.21922
1	-0.20801	-0.20809
2	-0.19717	-0.19724
3	-0.18662	-0.18669
4	-0.17635	-0.17641
5	-0.16635	-0.16642
6	-0.15664	-0.15670
7	-0.14719	-0.14725
8	-0.13801	-0.13807
9	-0.12909	-0.12915
10	-0.12044	-0.12050
11	-0.11204	-0.11210
12	-0.10390	-0.10396
13	-0.09602	-0.09607
14	-0.08839	-0.08844
15	-0.08102	-0.08107
16	-0.07390	-0.07395
17	-0.06704	-0.06709
18	-0.06044	-0.06049
19	-0.05411	-0.05415
20	-0.04804	-0.04808
21	-0.04225	-0.04229
22	0.03674	-0.03678
23	-0.03153	-0.03157
24	-0.02662	-0.02666
25	-0.02204	-0.02207
26	-0.01779	-0.01782
27	-0.01391	-0.01393
28	-0.01041	-0.01042
29	-0.00733	-0.00734
30	-0.00471	-0.00472
31	-0.00260	-0.00261
32	-0.00107	-0.00107
33	-0.00024	-0.00020
34	-0.00002	-0.00002

TABLE IV: Vibrational energies $E_{(\nu,0)}$ of the ground state $X^1\Sigma^+$ of the Tritium Fluorine molecule TF.